1

A Snapshot of the Global Drinking Water Virome: Diversity and Metabolic Potential Vary with Residual Disinfectant Use

6
7

Authors
Bridget Hegarty[1]
Zihan Dai[2]
Lutgarde Raskin[1]
Ameet Pinto[3]
Krista Wigginton[1,+]
Melissa Duhaime[4,*]

15
16
17
18

Corresponding Authors
+ Mailing address: Department of Civil and Environmental Engineering, 1351 Beal Ave., 181 Environmental and Water Resources Engineering Building, Ann Arbor, MI 48109-2125, USA
Phone: +1 (734) 763-9661
Email: kwigg@umich.edu

* Mailing address: Department of Ecology and Evolutionary Biology, 1105 N University Ave., 4068 Biological Sciences Building, Ann Arbor, MI 48109-1085, USA.
Phone: +1 (734) 764-6219;
E-mail: duhaimem@umich.edu

---

[1] Civil and Environmental Engineering; University of Michigan

[2] Key Laboratory of Drinking Water Science and Technology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China
[3] University of Chinese Academy of Sciences, Beijing 100049, China
[3] School of Civil and Environmental Engineering; Georgia Institute of Technology.
[4] Ecology and Evolutionary Biology; University of Michigan

Abstract

Viruses are important drivers of microbial community ecology and evolution, influencing microbial mortality, metabolism, and horizontal gene transfer. However, the effects of viruses remain largely unknown in many environments, including in drinking water systems. Drinking water metagenomic studies have offered a whole community perspective of bacterial impacts on water quality, but have not yet considered the influences of viruses. In this study, we address this gap by mining viral DNA sequences from publicly available drinking water metagenomes from distribution systems in six countries around the world. These datasets provide a snapshot of the taxonomic diversity and metabolic potential of the global drinking water virome; and provide an opportunity to investigate the effects of geography, climate, and drinking water treatment practices on viral diversity. Both environmental conditions and differences in sample processing were found to influence the viral composition. Using free chlorine as the residual disinfectant was associated with clear differences in viral taxonomic diversity and metabolic potential, with significantly fewer viral populations and less even viral community structures than observed in distribution systems without residual disinfectant. Additionally, drinking water viruses carry antibiotic resistance genes (ARGs), as well as genes to survive oxidative stress and nitrogen limitation. Through this study, we have demonstrated that viral communities are diverse across drinking water systems and vary with the use of residual disinfectant. Our findings offer directions for future research to develop a more robust understanding of how virus-bacteria interactions in drinking water distribution systems affect water quality.

Keywords: viruses, virome, metagenomics, microbiome, disinfection, drinking water

1. Introduction

In an effort to prevent the spread of waterborne diseases, previous research on viruses in drinking water has focused on tracking and eliminating human viral pathogens (Dong et al., 2010; Gall et al., 2015; Petrovich et al., 2020; Rao et al., 1981; Ye et al., 2012). For example, researchers have tracked specific human viruses with qPCR to understand their distribution throughout drinking water treatment plants and distribution systems (Albinana-Gimenez et al., 2006; Samhan et al., 2015; Ye et al., 2012). While research on the presence and fate of human viruses is essential for assessing drinking water quality, the impacts of bacterial viruses, commonly referred to as bacteriophages (or phages), have rarely been addressed. Phages influence microbial populations (Rohwer et al., 2009; Suttle, 2007) and biogeochemical cycling (Parmar et al., 2017), through mortality (Fuhrman and Noble, 1995; Guidi et al., 2016) and by facilitating horizontal gene transfer between microorganisms and across environments (Rohwer et al., 2009). Similarly, viruses likely impact both beneficial and detrimental bacteria and other microorganisms in drinking water systems that have both direct (e.g., causing illness) and indirect effects (e.g., nitrification and corrosion) on drinking water quality.

Metagenomic studies have become critical to evaluate the cumulative effects of the drinking water microbiome on water quality (Hull et al., 2019; Liu et al., 2013). Past studies have shown that many factors, ranging from source waters (Assche et al., 2019; Stanish et al., 2016) to treatment practices (Bautista-de los Santos et al., 2016; Dai et al., 2020; Pinto et al., 2012), influence the bacterial, archaeal, and eukaryotic inhabitants of drinking water. For instance, the use of residual disinfectants (e.g., free chlorine, chloramine) has been connected to the selection of opportunistic pathogens (Feazel et al., 2009; Kotlarz et al., 2019; Liu et al., 2019; Vaerewijck

et al., 2005), the prevalence of antimicrobial resistance genes (Jia et al., 2019, 2015; Shi et al.,

2012), and shaping microbial communities (Dai et al., 2020; Hull et al., 2019; Krishna et al.,

2020; Wang et al., 2014). At most, these studies have quantified the fraction of viral sequences in

a cellular metagenome (Brumfield et al., 2020; Gomez-Alvarez et al., 2012; Stamps et al., 2018).

This leaves fundamental information about drinking water viruses, such as their taxonomic or

metabolic diversity and influence on the drinking water microbiome, unknown. To fill similar

knowledge gaps in other environments, cellular metagenomes have been re-examined for DNA

viruses, revealing novel viral diversity and offering insights into viral-host interactions (Paez-

Espino et al., 2016; Roux et al., 2015b). Taking such an approach can provide an untapped

opportunity to illuminate the effects of phages on drinking water quality and safety.

The overall goal of this work was to characterize viral communities in drinking water

distribution systems. To do so, we identified viral sequences from publicly available

metagenomic datasets from drinking water distribution systems from around the world to assess

viral taxonomic diversity and metabolic potential. We further investigated the effect of study

characteristics (e.g., the type of filter used for sample processing and the DNA extraction

method) and environmental conditions (e.g., water quality) on the structuring of the observed

viral communities. This study lays the groundwork for understanding the impact of viruses on

the microbial communities in drinking water distribution systems.

2. Methodology

2.1 Data Collection

Publicly available drinking water metagenomic reads were downloaded from NCBI using the SRA Toolkit v2.9.6 (Leinonen et al., 2011) (Table S1). Sequences deposited on MG-RAST were directly requested from researchers. Only water samples that were collected from drinking water distribution systems and that were sequenced using Illumina short read technology were included in our analysis, excluding samples taken from within treatment plants and distribution system biofilm samples. Based on these criteria, 164 samples were used. Sequencing for some samples was replicated, resulting in a total of 312 sequencing runs. Replicate sequencing sets from the same samples were pooled during subsequent analyses ("sample" Table S1). Table S1 provides information about sample processing, sequencing platform, and other metadata for each sample included in this study. Water quality parameters for the Dai et al (2020) and Sevillano et al (2021) samples are provided in Tables S2 and S3, respectively.

2.2 Sequence Read Processing

Sequence reads were filtered and trimmed using fastp v0.20.1 (Chen et al., 2018). All code is deposited on github and made freely available (https://github.com/DuhaimeLab/global_drinking_water_viromes_Hegarty_et_al_2021). The UniVec Core database (as of February 2021) was used to remove contamination from vectors (NCBI, 2016).

2.3 Assembly

115 The filtered and processed reads from each distribution system sampling location were co-

116 assembled using MetaSPAdes v3.10.1 (Nurk et al., 2017). Assembly quality was assessed with

117 QUAST v5.0.2 (Gurevich et al., 2013).

2.4 Viral Population Identification

To define viral populations, putative viral contigs were identified from contigs greater than 3 kb

using a strategy based on VirSorter v1.0.6 (Roux et al., 2015a), VirSorter2 v2.1 (Guo et al.,

2021), CheckV v0.7.0 (Nayfach et al., 2020), VIBRANT v1.2.1 (Kieft et al., 2020), and

VirFinder v1.1 (Ren et al., 2018) predictions (Figure S2). Using the predictions considered high

quality by the tools' authors, any contig classified (i) as category 1 or 2 by VirSorter, (ii) as high

or medium by VIBRANT, (iii) as complete, high, or medium by CheckV, (iv) with a VirSorter2

score greater than 0.95, (v) with at least two hallmark viral genes from VirSorter2, or (vi) with a

VirFinder score greater than 0.9 were considered viral. Additionally, any contig called viral by at

least two of the following were considered viral: VirSorter categories 3-6, VIBRANT low,

CheckV low, Virfinder score between 0.7 and 0.9, and VirSorter2 score between 0.5 and 0.95. If

a contig had (i) no viral genes and more than one host gene according to CheckV, (ii) more than

3 times as many host genes as viral genes based on CheckV, or (iii) a length greater than 50 kb

and no viral hallmark genes from VirSorter2, it was removed from the viral list. Bacterial genes

were excised from the edges of proviruses based on their identification by CheckV, VirSorter2,

and VIBRANT. Based on the community-established standard for defining viral populations

(Roux et al., 2019), all viral contigs were clustered (stampede-clustergenomes) (Roux and

Bolduc, 2017) if they shared average nucleotide identity (ANI) of 95% across 85% of the contig

length ("viral populations" in the following).

The annotations of the three most ubiquitous VPs (based on frequency of occurrence and relative

abundance) were examined to assess the confidence of the phage annotations of each of these.

These three ubiquitous VPs were blasted against the nucleotide collection (nt) in NCBI to assess

whether they matched any previously sequenced proviruses. The top hits were run through

CheckV to identify any provirus sequences in their genome.

143   2.5 Read Mapping

Filtered and trimmed reads were mapped to all contigs from that sample's assembly using

Bowtie2 (Langmead and Salzberg, 2012) and quantified using SAM Tools v1.11 (Li et al.,

2009). To quantify the reads mapped to the trimmed viral sequences, BLAST v2.9.0 (NCBI,

2019) was used to align the trimmed viral sequences to the contigs and then reads overlapping

this region were quantified using FeatureCounts (Liao et al., 2014) from the Subread package

(Liao et al., 2013).When more than one sequencing run was performed for a given sample, the

total read coverage was summed across sampling replicates.

151   2.6 Alpha and Beta Diversity

Only contigs with reads covering at least 3 kb of their length were included in diversity analyses.

Alpha diversity measures (i.e., richness and Shannon Dissimilarity) were calculated using the

vegan v2.5-7 package (Oksanen et al., 2020) in R v4.0.2 (R Core Team, 2021). All analyses were

performed using RStudio v1.3. 595 (R Studio Team, 2020). To normalize for differences in

sequencing depth, taxa per one hundred thousand viral reads (henceforth, "normalized richness")

and the Shannon Evenness Index (calculated by dividing the Shannon Dissimilarity by the

maximum possible Shannon Dissimilarity for that sample) were used to examine alpha diversity

trends.

Transcripts per kilobase million (TPM) was determined for each viral population and used to

calculate the Bray Curtis distance between samples using the vegandist function. Permutational

Multivariate Analysis of Variance (PERMANOVA) using the adonis function (vegan) was used

to test the effect of processing procedures, geography, and residual disinfectant use on the full

community structure (Bray Curtis dissimilarities ~ continent * Köppen zone * filter pore size for

biomass collection * DNA extraction method; with stratas for 'study'). The effect of 'study' and

residual disinfectant were also tested in isolation (Bray Curtis dissimilarities ~ study; Bray Curtis

dissimilarities ~ residual disinfectant).

To avoid the confounding effects of study design and climate on diversity analyses, the Dai et al.

(2020) samples collected in the temperate oceanic climate zone (Cfb Köppen climate

classification scheme) were used to evaluate the effects of residual disinfectant use on taxonomic

diversity and metabolic potential (Figure S1 inset). This study was chosen because it provided

detailed water quality parameters and was the only study to include samples from distribution

systems operated without a residual disinfectant. The six samples taken from the warm summer

continental zone (Dfb Köppen climate classification) were excluded because they were all taken

from distribution systems that use a residual disinfectant, complicating the isolation of climate's

effect from that of residual disinfectant. Additionally, only samples that used free chlorine (and

not chloramines) were included to facilitate comparison with the results presented by Dai et al.

(2020). This reduced set of samples was used for all subsequent comparisons of water quality

and residual disinfectant. Differences in alpha diversity between distribution systems based on

residual disinfectant use were tested using the Wilcoxon rank sum test.

Additionally, both the Sevillano et al. (2021) and Dai et al (2020) samples were used to

investigate the effects of water quality parameters, including residual disinfectant, on the viral

community. These two studies were chosen as the only two to include both samples with and

without residual disinfectant. Distance-based Redundancy Analysis (db-RDA) was performed

using the capscale function in the vegan package v2.5-7 (Oksanen et al., 2020) in R (Bray Curtis

dissimilarities ~ chlorine + phosphate + temperature + pH + conductivity + ammonia + nitrate +

DO + TOC). PERMANOVA was used to test the effect of these water quality parameters and the

drinking water distribution system on the viral community structures of the Dai et al. (2020)

samples (Bray Curtis dissimilarities ~ chlorine + phosphate + temperature + pH + conductivity +

ammonia + nitrate + dissolved oxygen (DO) + total organic carbon (TOC) + drinking water

distribution system location). Parameters were considered significantly correlated if p-value $\leq$

0.05.

196 2.7 Metabolic Potential Analyses

Viral contig gene metabolic annotations were assigned using DRAM based on the KEGG and

Pfam databases (Shaffer et al., 2020). Read coverages of the open reading frames predicted by

DRAM were calculated using FeatureCounts (Liao et al., 2014) from the Subread package (Liao

et al., 2013). Bray Curtis distances between samples were calculated using the vegdist function

and then db-RDA was performed using the capscale function with the vegan package in R (Bray

Curtis dissimilarities ~ chlorine + phosphate + temperature + pH + conductivity + ammonia +

nitrate + DO + TOC). The water quality parameters were used in a PERMANOVA model to

evaluate the effect of water quality parameters on the abundance of KEGG orthologies (KOs),

which are molecular function classes that represent the metabolic potential of the viromes. The

following equation was used in the adonis function (vegan): Bray Curtis dissimilarities ~

chlorine + phosphate + temperature + pH + conductivity + ammonia + nitrate + DO + TOC +

drinking water distribution system location). Parameters were considered significantly correlated

if p-value ≤ 0.05. KEGG pathway modules were considered more abundant in a condition if the

difference in normalized read counts was statistically significant between the two conditions

(Welch's Two Sample t-test p-value ≤ 0.05).

212   2.8 Taxonomy

The open reading frames (ORFs) from DRAM were clustered with all full-length bacterial and

archaeal virus genomes from NCBI's viral reference dataset ('Viral RefSeq', v.85, January 2018)

using vConTACT2 (Bin Jang et al., 2019). Clusters with at least one taxonomic assignment from

vConTACT2, as well as clusters found in ≥ 10% of samples, were separately visualized with

Cytoscape v3.8.2 (Shannon et al., 2003). The lowest common ancestor was identified for each

vContact2 cluster with a taxonomic assignment.

219   3 Results/Discussion

To produce a snapshot of the virome in drinking water distribution systems, we mined publicly

available drinking water metagenomes (Chao et al., 2013; Dai et al., 2020; Douterelo et al.,

2018; Garner et al., 2018; Jia et al., 2015; Ma et al., 2019, 2017; Potgieter et al., 2020; Sevillano

et al., 2021; Shi et al., 2012; Zhang et al., 2019). Table 1 summarizes the metadata for each

study. The majority of the samples were collected in Europe (31%) and Asia (35%), with the

remainder obtained from North America (23%) and Africa (11%) (Figure S1). In total, 164

samples were collected from 69 distribution systems, with 18 samples from systems with no

residual disinfectant and 102 from systems with a residual disinfectant. 46 samples are presumed

to be from distribution systems using a residual disinfectant based on the typical disinfection

choice of their country of origin (i.e., China, Singapore, South Africa, United Kingdom, United

States), but this could not be confirmed (Table S1).

Table 1 - Summary data for the assemblies organized by study.

Study	Number of Samples	Number of Assemblies	Filter Pore Size	DNA Extraction Method	Continent	Köppen Zone	Residual Disinfectant
Chao et al. (2013)	2	1	0.22 μm filter	Promega Maxwell16 DNA extraction system	Asia	Cwa	chlorine
Dai et al. (2020)	49	14	0.22 μm filter	Promega Maxwell16 DNA extraction system	Europe	Cfb (171), Dfb (24)	chloramines (16), chlorine (107), no residual (72)
Douterelo et al. (2018)	3	3	0.22 μm filter	chloroform and isoamyl alcohol extraction	Europe	Cfb	chlorine
Garner et al. (2018)	2	1	0.22 μm filter	MP Biomedicals FastDNA SPIN kit	North America	B	chlorine
Jia et al. (2015)	8	1	0.45 μm filter	MP Biomedicals FastDNA SPIN kit	Asia	Cfa	chlorine
Ma et al. (2017)	25	25	0.45 μm, followed	MP Biomedicals	Africa (1), Asia (23),	Bsk (1), Bwk (2), Cfa (4),	not reported

Study			Filter	DNA extraction	Location	Climate	Disinfectant
			by 0.2μm filter	FastDNA SPIN Kit for Soil	North America (1)	Csb (1), Cwa (8), Cwb (1), Dfb(1), Dwa (4), Dwb (1) Dwc (1)	
Ma et al. (2019)	20	12	0.45 μm, followed by 0.2μm filter	MP Biomedicals FastDNA SPIN Kit for Soil	Asia	Af (1), Bsf (1), Bsk (1), Bwk (1), Cfa (8), Cwa (5), Dwa (2), Dwb (1)	not reported
Potgieter et al. (2020)	18	1	0.22 μm filter	phenol and chloroform DNA extraction	North America	Cwb	chloramines
Sevillano et al. (2021)	33	9	0.22 μm filter	QIAGEN DNeasy PowerWater Kit	North America	Am	chlorine (27), no residual detected (6)
Shi et al. (2012)	1	1	0.45 μm filter	MP Biomedicals FastDNA SPIN kit	Asia	Cfa	chlorine
Zhang et al. (2019)	3	1	0.45 μm filter	MP Biomedicals FastDNA SPIN Kit for Soil	Asia	Cfa	chlorine

All samples from the same drinking water distribution system were co-assembled (Tables S1 and

S4), resulting in 729,724 contigs larger than 3 kb. We then mined the assembled contigs for viral

sequences, identifying 62,500 viral sequences, or 9% of all the contigs longer than 3 kb. The

proportion of viral contigs is typically not reported in studies that have mined metagenomes for

viral sequences. Our results are in agreement with the upper range (0.3-16%) of the percentage

of viral sequences observed in a marine metagenome study (Delong et al., 2006). Additionally,

this is consistent with our expectations that viral genomic material would represent roughly 10%

of community DNA given that viruses are on average an order of magnitude more abundant than

their microbial hosts in most aquatic habitats, while their genomes tend to be one to two orders

of magnitude smaller. To facilitate comparisons of the viral populations across distribution

systems, the viral contigs were clustered into 48,935 viral populations based on 95% sequence

similarity over 85% of the contig length (Roux et al., 2019). These viral populations allowed us

to characterize the taxonomic diversity and metabolic potential of the drinking water virome, as

well as the geographic and treatment-based factors shaping these communities.

3.1 Viral ecology of drinking water communities

3.1.1 Drinking water viral populations are predominantly novel and heterogeneous between samples and studies

We used vConTACT2 to examine the evolutionary relatedness of the viral populations (VPs). Of

the total 62,500 viral sequences, 23,573 (37%) formed 7,897 clusters that approximate genus-

level relatedness (Bin Jang et al., 2019) (Figure 1A; Figure S3). Only 778 viral clusters (10% of

all viral clusters) were given a definitive taxonomic assignment per the International Committee

on Taxonomy of Viruses (ICTV) conventions by vConTACT2 (Figure S4). The vast majority of

these, 763 out of the 778 viral clusters, were assigned to families in the Caudovirales order of

tailed dsDNA phage. *Podoviridae* (373 VPs), *Siphoviridae* (284), and *Myoviridae* (81), were the

three most prevalent families of viruses identified in these samples, mirroring the three dominant

families in many other freshwater environments (Moon et al., 2020; Saparbaevna et al., 2017;

Skvortsov et al., 2016; Zaouri et al., 2020). The dominance of Caudovirales among the classified

VPs in our study may represent compositional bias towards this viral order, as the vConTACT

reference database is 90% Caudovirales. However, the family classification of our VPs does not

reflect database bias, as vConTACT contains 50% *Siphoviridae*, 23% *Myoviridae*, and 15%

Podoviradae (Bin Jang et al., 2019).

To date, the vast majority of VPs described in environmental viromics studies have been novel,

reflecting that viral databases insufficiently represent global viral diversity, as well as the fact

that the viral genomes reconstructed from these datasets tend to be incomplete and thus difficult

to classify. For instance, in recent studies of the Arctic Ocean (Gregory et al., 2019), honey bees

(Deboutte et al., 2020), and New York City wastewater (Gulino et al., 2020), only 9.8%, 9.5%,

and 4% of their viral populations, respectively, received a taxonomic assignment. Only 1% of the

drinking water viral populations in our study had a taxonomic assignment (Fig. 2A). Since many

viruses are strongly habitat-specific (Paez-Espino et al., 2016; Roux et al., 2012), the likelihood

of sequencing novel viruses from an environment poorly represented in the databases is high. In

addition to the lack of drinking water virome studies to date, the typical source waters (i.e.,

groundwater and surface water) for drinking water production have been less frequently studied

than marine aquatic environments (Kallies et al., 2019; Kothari et al., 2021; Malki et al., 2020;

Moon et al., 2020; Paez-Espino et al., 2016; Roux et al., 2012; Skvortsov et al., 2016).

Figure 1: Overview of the drinking water viral populations (VPs). (A) Status of drinking water VPs after taxonomic clustering by vConTACT2. VPs that were given a taxonomic assignment by vConTACT2 are in black. (B) Network clusters of VPs found in more than 10% of samples and their connections to viral reference genomes from vConTACT2. Drinking water VPs are black diamonds and viral reference genomes are colored by family-level taxonomic assignment. (C) Number of samples where a VP was abundant (i.e., greater than 1% of reads) versus the number of samples in which that viral population was found. Only contigs with a relative abundance of greater than 1% in at least one sample were included in this figure. The three most ubiquitous drinking water viral populations were labeled and further described below. Where multiple points overlap, they have a darker fill.

Of the 48,935 VPs (95% ANI across 85% of the contig) identified across all distribution

systems, 156 viral populations were found in more than 10% of the samples. Eight of these had

an order-level taxonomic assignment: seven belonged to the Caudovirales order (two viral

populations of *Myoviridae*, one *Podoviridae*, two *Siphoviridae*, two unassigned families), and

one VP belonged to the *Microviridae* family of isometric ssDNA phage in the *Petitvirales* order

(Figure 1B). Other studies have similarly found a high degree of heterogeneity with few

ubiquitous viral populations across different viromes collected from similar environments (e.g.,

within ecological zones in marine environments) (Gregory et al., 2019; Paez-Espino et al., 2016;

Roux et al., 2016).

The three most ubiquitous viral populations were VP_905, VP_2791, and VP_1288 (Figure 1C).

VP_2791, the most prevalent, was found in 42% of the samples. VP_905 and VP_2791 were

classified as unknown genera within the *Siphoviridae* family (Caudovirales). VP_1288 did not

have a taxonomic assignment at any level. The top blast hits for each of the most ubiquitous VPs

in the NCBI nt database (Table S5) were each within prophages identified by CheckV. Both

VP_905 and VP_2971 were most similar (based on query coverage) to a sequence in

Blastochloris viridis (Figure 2A, Table S5). *B. viridis* is a purple photosynthetic bacterium

(Tsukatani et al., 2015) that was isolated from a German river and has been found in other

freshwaters, including in the biofilm of an ultrapure supply water plant (Bohus et al., 2010).

VP_905 also matched with a similar query coverage to *Rhodobacter* sp. LPB0142 (Figure 2A)

and *Paracoccus contaminans.* VP_1288 was most similar to *Nitrobacter winogradskyi*, a nitrite

oxidizing bacterium that has been found in soils, as well as freshwater (Mellbye et al., 2015)

(Figure 2B). While most of the ORFs on the three contigs were not assigned a metabolic

annotation (Figure 2; Table S6), the high number of unknown and phage-related genes,

combined with the high sequence similarity and synteny with known prophages, increased the

confidence that each are truly phage contigs. By pairing deeply-sequenced viromes with cellular

metagenomes, subsequent studies will build on this first report of ubiquitous viral populations in

drinking water to establish "who infects whom", thereby leading to a more robust understanding

of how viruses impact drinking water quality.

323
Figure 2: Genome comparisons of the three most globally ubiquitous drinking water viral
populations (VPs) with the known sequences (NCBI nt) that share the greatest percent coverage
(blastn). The longest representative contig was visualized for each VP. (A) Comparison of
VP_905 (55.2 kb) with Rhodobacter sp. LPB0142, comparison of VP 905 and VP_2971 (35.0 kb)
with Blastochloris viridis. (B) Comparison of VP_1288 (46.8 kb) with Nitrobacter winogradskyi.
Genes on the VPs are colored based on their Pfam identity.

3.1.2 Environmental conditions and experimental design influence drinking water

viral communities

To evaluate the possible effects of experimental design and environmental conditions on viral

community structure, we tested the effects of the continent and Köppen climate zone where

samples were collected, the filter pore size for biomass collection, and the DNA extraction

method. In contrast to differences reported between viruses sequenced from 0.1-20 µm and <0.1

µm fractions (Williamson et al., 2012), our meta-analysis found that filter pore size did not

impact viral community structures (PERMANOVA; Table S7). As virus sizes typically range

from 0.02-0.3 µm, the relatively large filter pore size (0.22 µm and 0.45 µm) used across all

studies in our analysis may explain why filter pore size did not affect viral community structure.

The other three parameters tested, as well as the interaction of the DNA extraction method and

Köppen climate zone, did have a statistically significant effect on the viral community's structure

(PERMANOVA; Table S7). Combined, the parameters and their interactions tested explained

slightly over a quarter of the variability in the system (residual PERMANOVA R^2=0.73; Table

S7). The inability to explain the full variability in beta diversity in the PERMANOVA model

could reflect the unbalanced nature of the metadata (e.g., many more samples used the Maxwell

16 DNA extraction system than the other three methods; Figure S4), as well as the importance of

additional factors.

We additionally tested the effect of 'study' to encompass the experimental design and

environmental conditions, including those not explicitly reported in all of the original

metagenomics studies (e.g., water source, distribution system water qualities, library

preparation). 'Study' was found to significantly impact the viral community structure

(PERMANOVA R^2=0.21, p-value=0.0001). In a previous meta-analysis of drinking water

bacterial communities, 'study'-specific effects explained nearly a third of the variability among

samples (Bautista-de los Santos et al., 2016). The collection and processing of samples are

known to have a substantial impact on the viruses identified (Brinkman et al., 2018; Dias et al.,

2020; Hurwitz et al., 2013; Percival and Wyn-Jones, 2013). The low biomass of drinking water

samples may make them particularly prone to the influence of 'study'-specific differences from

contamination during sample processing (Bautista-de los Santos et al., 2016). Alternatively, as

most studies only examined one distribution system or focused on a single region, the strong

'study'-specific effects may be partly due to local factors, such as geography, climate, treatment

regime, and water source (e.g., groundwater, river water). Such parameters have been shown to

influence the viral communities of many other environments (Paez-Espino et al., 2016).

366
Figure 3: Diversity of viral populations cluster based on residual disinfectant use and water quality.
dbRDA ordination of the Bray Curtis dissimilarities of the viral community structure for (A) Dai et al.
(2020) and (B) Sevillano et al (2021) samples. Distribution system is represented by shape and residual
disinfectant use by color. The ordination is overlaid with the gradient of the fit between the measured
environmental parameters and the viral community dissimilarities. Only samples from distribution
systems using chlorine in the Cfb Köppen climate zone from the Dai et al. (2020) study were included in
4A.
374

To investigate the influence of water quality parameters independently from study-specific

effects, we focused on the Dai et al. (2020) samples, collected from the UK and Netherlands, and

the Sevillano et al. (2021) samples, collected from Puerto Rico. Both studies found that chlorine

was a major determinant of the differences in the microbial community composition (Dai et al.,

2020; Sevillano et al., 2021). Dai et al. (2020) additionally found that DO and conductivity were

correlated with the microbial community structure of these samples; Sevillano et al. (2021) found

that DO, temperature, and TOC were each significantly associated with changes in microbial

composition at specific sites. As bacterial community structure is a critical driver of

bacteriophage community structure (Gregory et al., 2019; Jonge et al., 2018), we hypothesized

that these parameters would also be correlated with the viral community relatedness (Bray Curtis

dissimilarities). We found that viral beta diversity (i.e., differences in community structure

between samples) correlated with chlorine, phosphate, and nitrate concentration, DO,

conductivity, water temperature, pH, and TOC for both studies; ammonia concentration was also

significantly correlated for the Dai et al. (2020) samples (Figure 3; Tables S8 and S9). These

trends indicated that water quality parameters do likely shape the viral communities of drinking

water. Distribution system sampling location was significantly correlated with the viral

community structure of both the Dai et al. (2020) study (PERMANOVA R^2=0.33, p-

value=0.001; Table S8) and Sevillano et al. (2021) study (PERMANOVA R^2=0.32, p-

value=0.0001; Table S9). Together with the unexplained variability in both the Dai et al. (2020)

(PERMANOVA R^2=0.19; Table S8) and the Sevillano et al. (2021) samples (PERMANOVA

R^2=0.25; Table S9), these trends indicated that unmeasured factors also likely affected the viral

community structures. Altogether, the Sevillano et al. (2021) and Dai et al. (2020) viral

sequences suggest that water quality parameters are influencing the structure of viral

communities independently of study-specific differences in experimental design.

3.1.3 Systems with residual disinfectant have less diverse and less even viral

communities

To assess the effect of residual disinfectant on the viral communities, we first assessed within-

sample diversity (alpha diversity). The average normalized viral richness was 350 ± 820 and the

average Shannon evenness index was 0.28 ± 0.10. The Shannon dissimilarities for the drinking

water viral populations range from 0.26-7.7, which is consistent with ranges reported for other

freshwater sources (Aguirre De Cárcer et al., 2015; Tseng et al., 2013). Distribution systems that

used a residual disinfectant had less diverse viral communities than distribution systems that did

not use a residual disinfectant. This trend held for systems using both chlorine and chloramine,

as well as for the suspected residual disinfectant samples (Tukey's test $p_{adj} \leq 10^{-6}$; Figure S5).

Further, no difference was found between the chlorine and chloramine samples (Tukey's test p_{adj}

$= 0.92$), nor between the suspected residual disinfectant samples and either the chlorine (Tukey's

test $p_{ad\,j} = 0.25$) or chloramine (Tukey's test $p_{adj} = 0.85$) samples. Together, these trends supported

our prediction that samples from studies with suspected residual disinfectant usage were indeed

likely obtained from systems operated with a residual disinfectant.

415
Bacterial communities in drinking water systems with residual disinfectant have been observed

to have lower richness and evenness (Bautista-de los Santos et al., 2016; Dai et al., 2020). Dai et

al. (2020) proposed that disinfectant residuals exert a stronger selective pressure than the low

phosphate and ammonia availability in distribution systems that do not use a residual

disinfectant. For the Dai et al. (2020) samples, we similarly found that viral communities in

drinking water distribution systems with a residual disinfectant (i.e., free chlorine) were less

diverse and less even than distribution systems without a residual disinfectant (Wilcoxon rank

sum test p-values: normalized richness p-value = $3.0*10^{-6}$ and Shannon evenness index p-value =

0.001; Figure 4). Correlations between bacterial and viral diversity have been observed in a

range of environments (Anderson et al., 2017; Dalcin Martins et al., 2018; Yang et al., 2019).

Whether residual disinfectant selects for certain viral populations in drinking water, thereby

reducing viral community diversity, or the viral population dynamics are following those of their

bacterial hosts as they are affected by residual disinfectant and other changing environmental

parameters, is unknown. As phages impose a selective pressure on bacterial communities, phages

can both disrupt desirable bacterial communities during industrial processes (e.g., yogurt

production), as well as be used to prevent contamination by pathogens (e.g., to prevent milk

spoilage) (Marks and Sharp, 2000). Based on the abundance and diversity of phages in drinking

water, we hypothesize that phages are an important factor influencing the stability of bacterial

communities in drinking water. However, more systematic and deeply sequenced analyses of

paired cellular and viral metagenomes is needed to better resolve virus, host, and infection

dynamics in drinking water distribution systems and their consequences for water quality.

438
439 *Figure 4 - Viral diversity varied based on whether samples originated from distribution systems that use*
440 *or do not use a residual disinfectant. (A) Observed normalized viral richness and (B) Shannon Evenness*
441 *Index of the viral community of each sample. Note that only samples from the Cfb Köppen climate zone*
442 *and samples with chlorine as the residual disinfectant from the Dai et al. (2020) study were included.*
443

Residual disinfectant is also a significant determinant of between-sample bacterial community

structure (Bautista-de los Santos et al., 2016; Dai et al., 2020), e.g., beta diversity. Therefore, we

anticipated it to have a similar effect on viral community structure. The viral communities from

the Dai et al. (2020) study clustered based on residual disinfectant use (PERMANOVA R^2=0.09,

p-value=0.0001; Figure 3A). While the Dai et al. (2020) samples considered were all taken from

the same Köppen climatic zone, the residual disinfectant samples were taken in the summer/fall

and have lower conductivities and the no residual disinfectant samples were taken during the

winter/spring and have higher conductivities (Tables S2, S10). For this reason, the differences in

viral community structure may reflect the additive effect of water quality differences that reflect

treatment (e.g., chlorine, phosphate, DO, ammonia) and other factors that may confound the

effect of treatment, such as season and water source (e.g., temperature and conductivity). Both

season (Pinto et al., 2014; Potgieter et al., 2018; Prest et al., 2016) and water source (Assche et

al., 2019; Roeselers et al., 2015) have been shown to influence drinking water microbial

communities. When season and water source have been examined together, systems that are

groundwater fed have lacked the strong seasonal trends observed in surface water fed systems

(Assche et al., 2019; Roeselers et al., 2015). Therefore, season may have a weaker confounding

effect than source water for the no residual disinfectant samples, as their conductivities suggest

groundwater sources (Douterelo et al., 2018).

Providing an additional opportunity to consider the effect of residual disinfectant, one site from

the Sevillano et al. (2021) study, GUA, had chlorine concentrations below the detection limit or

much lower than the other sites (chlorine concentration < 0.05 mg/L). The viral community

structure of the GUA samples was distinct compared to the other samples (PERMANOVA

R^2=0.18, p-value=0.0002; Figure 3B), including those that had similar conductivities,

temperatures, and other water quality parameters. While this further supports the influence of

residual disinfectant on viral community structure, more studies are necessary to determine the

distinct effects of residual disinfectant use on the viral communities of drinking water

distribution systems separate from season and water source.

472   3.2 The metabolic potential of drinking water viromes

Viral genomes often include genes necessary for replication, as well as host-derived auxiliary

metabolic genes (AMGs) thought to reprogram host cellular metabolism to support infection

processes (Warwick-Dugdale et al., 2019). Characterizing and understanding the influences on

the metabolic potential of drinking water viruses is, therefore, an essential step in understanding

their influence on the cellular members of the drinking water microbiome.

3.2.1 Water quality parameters are correlated with differences in viral community

metabolic potential

The viral community metabolic potential varies with water quality parameters in similar ways to

the viral community structure. Specifically, samples clustered primarily by disinfectant status

(Figure 5A) and all tested water quality parameters correlated with the metabolic profiles of

samples from systems using residual disinfectant, other than DO (Figure 7; PERMANOVA

Table S11). Furthermore, the strong correlation between metabolic profiles and drinking water

distribution system (PERMANOVA R^2=0.30, p-value=0.001) and large degree of unexplained

variability (residual PERMANOVA R^2=0.21) suggested that much of the metabolic variability

was due to unreported water quality and treatment specific differences. Unlike the viral

populations, which were more diverse in distribution systems with no residual disinfectant, there

was more diversity in metabolic potential in the residual disinfectant systems (Wilcoxon rank

sum test p-value = 0.03; Figure S6). This may reflect an evolutionary benefit of carrying extra

genes to survive the oxidative stress of residual disinfectant presence.

493
Figure 5: Metabolic potential of the drinking water virome. (A) dbRDA ordination reveals that the
diversity of metabolic potential clusters based on residual disinfectant use and water quality. The shape
of the points represents the distribution system, and the color represents the residual disinfectant. The
ordination is overlaid with the gradient of the fit between the measured environmental parameters and
the Bray Curtis dissimilarities. (B) KEGG Ontology (KO) Pathway module relative abundance for the
samples from Dai et al (2020). The size of the bubble represents the relative abundance of that pathway
module among the reads in that sample. Only samples from the Cfb Köppen climate zone from the Dai et
al. (2020) study were included in A and B. (C) Most abundant protein families (Pfam). The percent of
genes is the percentage of the number of genes with a specific Pfam annotation divided by the total
number of genes with a Pfam annotation. Only those Pfams with a percentage greater than 0.5% are
shown. Bars are colored based on whether they were phage non-structural, phage structural, or genes of
unknown function. Figures 6A and 6B only includes the samples from distribution systems using chlorine
and from the Cfb Köppen climate zone.
507

The relative abundance of KO metabolic pathway modules also differed between distribution

systems that used a residual disinfectant and those that did not (Figure 5B, Table S12). Viral

contigs from distribution systems that did not use a residual disinfectant were enriched in genes

with nucleotide metabolism annotations. Contrastingly, viral contigs from distribution systems

that used a residual disinfectant were enriched in genes involving the metabolism of cofactors

and vitamins and for xenobiotics biodegradation and metabolism. The higher availability of

genes related to these two pathway modules in residual disinfectant systems may reflect the

evolutionary pressure to carry genes capable of mitigating the effects of reactive chlorine species

(Huang et al., 2021; Kavagutti et al., 2019). However, these trends, particularly the

overabundance of nucleotide metabolism in no residual disinfectant systems, could potentially be

an artifact of the increased diversity of these samples (Figure 4). As more than 85% of the KOs

were found in less than a quarter of the samples (Figure S7), further studies with deeper

sequencing or viral-enriched genomes are necessary to robustly investigate differential

abundances of molecular functions.

3.2.2 Drinking water viruses encode metabolic genes with the potential to augment

microbial metabolic processes

As is common in other environmental viromes (Deboutte et al., 2020; Gregory et al., 2019), most

of the predicted viral genes identified in this study were novel, meaning they do not share

homology with known genes in public databases. As a result, only 25% had a Pfam annotation

and 6% could be assigned a KEGG orthology (KO) molecular function. Of the six most

prevalent Pfams, two were phage structural protein families and four were phage non-structural

proteins (Figure 5C). The most abundant was a Caudovirus prohead serine protease

(PF04586.18), an enzyme involved in capsid maturation. The ubiquity of viral-related

annotations among the most abundant Pfams supports that viral contigs were successfully sorted

from the majority cellular metagenomes.

The top four most abundant KO metabolic modules were nucleotide metabolism (44% of the

reads that mapped to viral genes with KO annotations), cofactors and vitamins metabolism

(18%), amino acid metabolism (11%), and carbohydrate metabolism (10%). Genes related to

nucleotide and amino acid metabolism are commonly abundant in viral genomes, due to their

importance in viral replication (Hurwitz et al., 2015). Carbohydrate metabolism genes carried by

viruses may be involved in reprogramming central carbon metabolism to favor energy

production necessary for viral replication (Howard-Varona et al., 2020). In the oligotrophic

environment of the drinking water microbiome, these genes, along with those encoding for

proteins related to cofactor and vitamin metabolism, might contribute to the survival of the

bacteriophages' hosts or enhance viral replication.

Nitrogen metabolism genes are common among drinking water bacteria (Potgieter et al., 2020) to

survive the oligotrophic conditions of drinking water (Kooij and Wielen, 2013). In other

nitrogen-limited environments, phages have carried more genes related to nitrogen cycling

(Warwick-Dugdale et al., 2019). Similarly, genes related to nitrogen metabolism were identified

in the drinking water virome in this study (Figure S8), including carbonic anhydrase (K01673),

glutamine synthetase (K01915), nitrate reductase/nitrite oxidoreductase (K00370), glutamate

dehydrogenase (K15371), nitric oxide reductase (K04748), nitrate/nitrite transporter (K02575),

and nitrogen fixation protein NifU (K04488). The most ubiquitous nitrogen metabolism related

genes were glutamine synthetase (K01915; 11 assemblies), followed by NifU (K04488; 8

assemblies). In addition to being part of nitrogen metabolism, glutamine synthetase is also

involved in amino acid metabolism. The presence of viral encoded nitrogen-related genes

suggests that in drinking water distribution systems, viruses may impact nutrient metabolism,

either through rewiring host metabolism to support infection or by mediating horizontal gene

transfer of nutrient-stress related genes (Warwick-Dugdale et al., 2019; Zimmerman et al., 2020).

To survive in drinking water systems with a residual disinfectant, microorganisms must also

mitigate the effects of reactive chlorine species (Gray et al., 2013). In this study, glutathione

reductase (K00383) and genes responsible for four of the five steps of coenzyme A biosynthesis

(M00120; Figure S9) were identified in the viruses of distribution systems that use a residual

disinfectant (59 samples in 5 studies). Low molecular weight thiols, including glutathione and

coenzyme A, which maintain the reducing potential of the cytosol (Chesney and Eaton, 1996),

have previously been found in bacterial genomes from drinking water distribution systems

treated with chlorine (Dai et al., 2020; Douterelo et al., 2018; Gomez-Alvarez et al., 2012). In

other environments, genes broadly involved in surviving oxidative stress have been found in

phage genomes (Huang et al., 2021; Kavagutti et al., 2019) and have been shown to be beneficial

to viral fitness (Holmgren, 1989; Russel and Model, 1986). While viral genes are implicated in

combating oxidative stress in distribution systems using a residual disinfectant, additional

research is needed to assess the role of viruses in mitigating the harmful effects of reactive

chlorine species in drinking water, specifically.

Taken together, the presence of genes related to nitrogen metabolism and oxidative stress

mitigation on drinking water viral contigs suggest that phages may contribute to the transfer of

genes that are important for microbial survival in drinking water distribution systems.

579 3.3 Opportunities for Future Work

Metagenomic analyses allow researchers to make broad assessments of microbial communities

and suggest areas for more targeted studies. The analyses conducted here suggest that residual

disinfectant use impacts the diversity and metabolic potential of the drinking water virome.

However, future bench-scale experiments will be necessary to separate the effects of residual

disinfectant from other water quality and sampling differences. Confounding methodological

differences between studies, such as DNA extraction method, can mask or be correlated with the

factors, including climate, geography, and treatment regime, that may shape the microbial

communities of drinking water. While standardizing methodologies may be impossible and

premature, coordination between researchers could facilitate cross-study comparisons (Bautista-

de los Santos et al., 2016; Hull et al., 2019). An excellent first step would be to make raw

sequencing data public and report temperature, water quality characteristics (e.g., chlorine,

phosphate, ammonia, nitrate, TOC concentrations), and biomass proxies (e.g., ATP levels, cell

counts), as well as other metadata (e.g., source water characteristics, details of the treatment,

season, location) (Bautista-de los Santos et al., 2016; Yilmaz et al., 2011). Application of

standardized spike-ins, such as "sequins" (Hardwick et al., 2018), to drinking water microbiomes

would further facilitate between study comparisons, as well as reveal whether the changes in

viral gene relative abundances described in this work mirror the trends in absolute abundance.

For the most robust assessments of the drinking water virome, future studies must specifically

develop sample processing methods that enrich for viruses and are optimized for the sample

matrix (Langenfeld et al., 2021). As the drinking water samples of this meta-analysis were not

collected with the intention of enriching for submicron viral particles, the sequenced viruses may

be biased towards particle-associated viruses, viruses actively infecting a cellular host, and free

viruses trapped by the chosen filter. Future studies designed specifically to query the viral

fraction of drinking water (including both DNA and RNA viruses) and with deeper sequencing

efforts are necessary to determine whether the trends described here are reflective of the full viral

community. To more fully capture both the drinking water viral community and their influences

on bacterial communities, future studies should also explore viruses in drinking water biofilms.

As part of this effort, future work should characterize the viral genes that are on DNA within

viral particles, as compared to extracellular DNA. Altogether, these studies will also determine

the extent to which phage-facilitated metabolic reprogramming or gene transfer contributes to

bacterial survival in drinking water systems. By connecting the drinking water virome's

dynamics with that of the bacterial community, improved treatment processes may become

possible.

4 Conclusions

- Mining drinking water metagenomes for viral sequences reveals differences in the viral

community based on geography and water quality parameters.

- More diverse and even viral communities are found in distribution systems that do not

use a residual disinfectant.

-     Viruses in drinking water carry genes, such as those for nitrogen metabolism and chlorine

stress mitigation, that mirror those identified as critical for the drinking water bacterial

community.

Acknowledgements

We would like to thank the Remaking Water Infrastructure Blue Sky team, particularly Sarah

Potgieter, Christopher Anderson, and Katherine Dowdell, as well as the Duhaime and Wigginton

labs for contributing their technical and intellectual expertise for bioinformatic experimental

design and analyses and manuscript preparation. This work was supported by the Blue Sky

Initiative of the University of Michigan College of Engineering and NSF CAREER funding to

Ameet Pinto [NSF CBET 1749530].

Citations

Aguirre De Cárcer, D., López-Bueno, A., Pearce, D.A., Alcamí, A., 2015. Biodiversity and

distribution of polar freshwater DNA viruses. Sci. Adv.

https://doi.org/10.1126/sciadv.1400127

Albinana-Gimenez, N., Clemente-Casares, P., Bofill-Mas, S., Hundesa, A., Ribas, F., Girones,

R., 2006. Distribution of Human Polyomaviruses, Adenoviruses, and Hepatitis E Virus in

the Environment and in a Drinking-Water Treatment Plant. Environ. Sci. Technol. 40,

7416–7422. https://doi.org/10.1021/es060343i

Anderson, C.L., Sullivan, M.B., Fernando, S.C., 2017. Dietary energy drives the dynamic

response of bovine rumen viral communities. Microbiome 5, 155.

https://doi.org/10.1186/s40168-017-0374-3

Assche, A. Van, Lievens, B., Crauwels, S., Brabanter, J. De, Willems, K.A., 2019.

Characterization of the bacterial community composition in water of drinking water

production and distribution systems in 1–11. https://doi.org/10.1002/mbo3.726

Bautista-de los Santos, Q.M., Schroeder, J.L., Sevillano-Rivera, M.C., Sungthong, R., Ijaz, U.Z.,

Sloan, W., T, Pinto, A.J., 2016. Emerging investigators series: microbial communities in

full-scale drinking water distribution systems – a meta-analysis. Environ. Sci. Water Res 2,

631–644. https://doi.org/10.1039/c6ew00030d

Bin Jang, H., Bolduc, B., Zablocki, O., Kuhn, J.H., Roux, S., Adriaenssens, E.M., Brister, J.R.,

Kropinski, A.M., Krupovic, M., Lavigne, R., Turner, D., Sullivan, M.B., 2019. Taxonomic

assignment of uncultivated prokaryotic virus genomes is enabled by gene-sharing networks.

Nat. Biotechnol. 37, 632–639. https://doi.org/10.1038/s41587-019-0100-8

Bohus, V., Toth, E.M., Szekely, A.J., Makk, J., Baranyi, K., Patek, G., Schunk, J., Marialigeti,

K., 2010. Microbiological investigation of an industrial ultra pure supply water plant using

cultivation-based and cultivation-independent methods. Water Res. 44, 6124–6132.

https://doi.org/10.1016/j.watres.2010.07.006

Brinkman, N.E., Villegas, E.N., Garland, J.L., Keely, S.P., 2018. Reducing inherent biases

introduced during DNA viral metagenome analyses of municipal wastewater. PLoS One 13,

1–23. https://doi.org/10.1371/journal.pone.0195350

Brumfield, K.D., Hasan, N.A., Id, M.B.L., Cotruvo, J.A., Rashed, M., Id, R.R.C., Huq, A., 2020.

A comparative analysis of drinking water employing metagenomics. PLoS One 15, 1–27.

https://doi.org/10.1371/journal.pone.0231210

Chao, Y., Ma, L., Yang, Y., Ju, F., Zhang, X.X., Wu, W.M., Zhang, T., 2013. Metagenomic

analysis reveals significant changes of microbial compositions and protective functions

during drinking water treatment. Sci. Rep. 3, 1–9. https://doi.org/10.1038/srep03550

Chen, S., Zhou, Y., Chen, Y., Gu, J., 2018. fastp : an ultra-fast all-in-one FASTQ preprocessor

884–890. https://doi.org/10.1093/bioinformatics/bty560

Chesney, J.A., Eaton, J.W., 1996. Bacterial Glutathione : a Sacrificial Defense against Chlorine

Compounds. J. Bacteriol. 178, 2131–2135.

Dai, Z., Sevillano-rivera, M.C., Calus, S.T., Santos, Q.M.B.L., Eren, A.M., Wielen, P.W.J.J. Van

Der, Ijaz, U.Z., Pinto, A.J., 2020. Disinfection exhibits systematic impacts on the drinking

water microbiome. Microbiome 1–19. https://doi.org/10.1186/s40168-020-00813-0

Dalcin Martins, P., Danczak, R.E., Roux, S., Frank, J., Borton, M.A., Wolfe, R.A., Burris, M.N.,

Wilkins, M.J., 2018. Viral and metabolic controls on high rates of microbial sulfur and

carbon cycling in wetland ecosystems. Microbiome 6, 1–17.

https://doi.org/10.1186/s40168-018-0522-4

Deboutte, W., Beller, L., Kwe, C., Maes, P., Graaf, D.C. De, 2020. Honey-bee – associated

prokaryotic viral communities reveal wide viral diversity and a profound metabolic coding

potential 117, 10511–10519. https://doi.org/10.1073/pnas.1921859117

Delong, E.F., Preston, C.M., Mincer, T., Rich, V., Hallam, S.J., Frigaard, N., Martinez, A.,

Sullivan, M.B., Edwards, R., Brito, B.R., Chisholm, S.W., Karl, D.M., 2006. Community

Genomics Among Stratified Microbial Assemblages in the Ocean's Interior. Science (80-.).

311, 496–504. https://doi.org/10.1126/science.1120250

Dias, R.S., Abe, A.E., Lima, H.S., Silva, L.C.F., de Paula, S.O., da Silva, C.C., 2020. Viral

concentration methods for diversity studies in soil samples. Appl. Soil Ecol. 155, 103666.

https://doi.org/10.1016/j.apsoil.2020.103666

Dong, Y., Kim, J., Lewis, G.D., 2010. Evaluation of methodology for detection of human

adenoviruses in wastewater , drinking water , stream water and recreational waters. J. Appl.

Microbiol. 108, 800–809. https://doi.org/10.1111/j.1365-2672.2009.04477.x

Douterelo, I., Calero-Preciado, C., Soria-Carrasco, V., Boxall, J.B., 2018. Whole metagenome

sequencing of chlorinated drinking water distribution systems. Environ. Sci. Water Res.

Technol. 4, 2080–2091. https://doi.org/10.1039/c8ew00395e

Feazel, L.M., Baumgartner, L.K., Peterson, K.L., Frank, D.N., Harris, J.K., Pace, N.R., 2009.

Opportunistic pathogens enriched in showerhead biofilms. Proc. Natl. Acad. Sci. 106,

16393–16399. https://doi.org/10.1073/pnas.0908446106

Fuhrman, J.A., Noble, R.T., 1995. Viruses and protists cause similar bacterial mortality in

coastal seawater 40, 1236–1242.

Gall, A.M., Mari, B.J., Lu, Y., Shisler, J.L., 2015. Waterborne Viruses : A Barrier to Safe

Drinking Water. PLOS Pathog. 1–7. https://doi.org/10.1371/journal.ppat.1004867

Garner, E., Chen, C., Xia, K., Bowers, J., Engelthaler, D.M., Mclain, J., Edwards, M.A., Pruden,

A., 2018. Metagenomic Characterization of Antibiotic Resistance Genes in Full- Scale

Reclaimed Water Distribution Systems and Corresponding Potable Systems. Environ. Sci.

Technol. 52, 6113–6125. https://doi.org/10.1021/acs.est.7b05419

Gomez-Alvarez, V., Revetta, R.P., Domingo, J.W.S., 2012. Metagenomic Analyses of Drinking

Water Receiving Different Disinfection Treatments. AEM 78, 6095–6102.

https://doi.org/10.1128/AEM.01018-12

Gray, M.J., Wholey, W.-Y., Jakob, U., 2013. Bacterial Responses to Reactive Chlorine Species.

Annu. Rev. Microbiol. 141–160. https://doi.org/10.1146/annurev-micro-102912-

142520.Bacterial

Gregory, A.C., Zayed, A.A., Conceição-Neto, N., Temperton, B., Bolduc, B., Alberti, A.,

Ardyna, M., Arkhipova, K., Carmichael, M., Cruaud, C., Dimier, C., Domínguez-Huerta,

G., Ferland, J., Kandels, S., Liu, Y., Marec, C., Pesant, S., Picheral, M., Pisarev, S., Poulain,

J., Tremblay, J.É., Vik, D., Acinas, S.G., Babin, M., Bork, P., Boss, E., Bowler, C.,

Cochrane, G., de Vargas, C., Follows, M., Gorsky, G., Grimsley, N., Guidi, L., Hingamp,

P., Iudicone, D., Jaillon, O., Kandels-Lewis, S., Karp-Boss, L., Karsenti, E., Not, F., Ogata,

H., Poulton, N., Raes, J., Sardet, C., Speich, S., Stemmann, L., Sullivan, M.B., Sunagawa,

S., Wincker, P., Culley, A.I., Dutilh, B.E., Roux, S., 2019. Marine DNA Viral Macro- and

Microdiversity from Pole to Pole. Cell 177, 1109-1123.e14.

https://doi.org/10.1016/j.cell.2019.03.040

Guidi, L., Chaffron, S., Bittner, L., Eveillard, D., Larhlimi, A., Roux, S., Darzi, Y., Audic, S.,

Berline, L., Brum, J.R., Coelho, L.P., Cesar, J., Espinoza, I., Malviya, S., Sunagawa, S.,

Dimier, C., Kandels-lewis, S., Picheral, M., Poulain, J., 2016. Plankton networks driving

carbon export in the oligotrophic ocean. Nature. https://doi.org/10.1038/nature16942

Gulino, K., Rahman, J., Badri, M., Morton, J., Bonneau, R., Ghedin, E., 2020. Initial Mapping of

the New York City Wastewater Virome. mSystems 5, 1–18.

https://doi.org/10.1128/msystems.00876-19

Guo, J., Bolduc, B., Zayed, A.A., Varsani, A., Dominguez-huerta, G., Delmont, T.O., Pratama,

A.A., Gazitúa, M.C., Vik, D., Sullivan, M.B., Roux, S., 2021. VirSorter2: a multi-classifier,

expert-guided approach to detect diverse DNA and RNA viruses. Microbiome 9, 9–37.

https://doi.org/10.1186/s40168-020-00990-y

Gurevich, A., Saveliev, V., Vyahhi, N., Tesler, G., 2013. QUAST : quality assessment tool for

genome assemblies. Bioinformatics 29, 1072–1075.

https://doi.org/10.1093/bioinformatics/btt086

Hardwick, S.A., Chen, W.Y., Wong, T., Kanakamedala, B.S., Deveson, I.W., Ongley, S.E.,

Santini, N.S., Marcellin, E., Smith, M.A., Nielsen, L.K., Lovelock, C.E., Neilan, B.A.,

Mercer, T.R., 2018. Synthetic microbe communities provide internal reference standards for

metagenome sequencing and analysis. Nat. Commun. 9, 1–10.

https://doi.org/10.1038/s41467-018-05555-0

Holmgren, A., 1989. Electron Transport to Reductive Enzymes 264, 13963–13966.

Howard-Varona, C., Lindback, M.M., Bastien, G.E., Solonenko, N., Zayed, A.A., Jang, H.,

Andreopoulos, B., Brewer, H.M., Rio, T.G. del, Adkins, J.N., Paul, S., Sullivan, M.B.,

Duhaime, M.B., 2020. Phage-specific metabolic reprogramming of virocells. ISME J. 14,

881–895. https://doi.org/10.1038/s41396-019-0580-z

Huang, X., Jiao, N., Zhang, R., 2021. The genomic content and context of auxiliary metabolic

genes in roseophages. Environ. Microbiol. 23, 3743–3757.

https://doi.org/https://doi.org/10.1111/1462-2920.15412

Hull, N.M., Ling, F., Pinto, A.J., Albertsen, M., Jang, H.G., Hong, P.Y., Konstantinidis, K.T.,

LeChevallier, M., Colwell, R.R., Liu, W.T., 2019. Drinking Water Microbiome Project: Is it

Time? Trends Microbiol. 27, 670–677. https://doi.org/10.1016/j.tim.2019.03.011

Hurwitz, B.L., Brum, J.R., Sullivan, M.B., 2015. Depth-stratified functional and taxonomic niche

specialization in the ' core ' and ' flexible ' Pacific Ocean Virome 472–484.

https://doi.org/10.1038/ismej.2014.143

Hurwitz, B.L., Deng, L., Poulos, B.T., Sullivan, M.B., 2013. Evaluation of methods to

concentrate and purify ocean virus communities through comparative, replicated

metagenomics. Environ. Microbiol. 15, 1428–1440. https://doi.org/10.1111/j.1462-

2920.2012.02836.x

Jia, S., Shi, P., Hu, Q., Li, B., Zhang, T., Zhang, X., 2015. Bacterial Community Shift Drives

Antibiotic Resistance Promotion during Drinking Water Chlorination. Environ. Sci.

Technol. 49, 12271–12279. https://doi.org/10.1021/acs.est.5b03521

Jia, S., Wu, J., Ye, L., Zhao, F., Li, T., Zhang, X., 2019. Metagenomic assembly provides a deep

insight into the antibiotic resistome alteration induced by drinking water chlorination and its

correlations with bacterial host changes. J. Hazard. Mater. 379, 120841.

https://doi.org/10.1016/j.jhazmat.2019.120841

Kallies, R., Hölzer, M., Toscan, R.B., Nunes, U., 2019. Evaluation of Sequencing Library

Preparation Protocols for Viral Metagenomic Analysis from Pristine Aquifer Groundwaters.

Viruses 11. https://doi.org/doi:10.3390/v11060484

Kavagutti, V.S., Andrei, A.-Ş., Mehrshad, M., Salcher, M.M., Ghai, R., 2019. Phage-centric

ecological interactions in aquatic ecosystems revealed through ultra- deep metagenomics.

Microbiome 7, 1–15. https://doi.org/10.1186/s40168-019-0752-0

Kieft, K., Zhou, Z., Anantharaman, K., 2020. VIBRANT: Automated recovery, annotation and

curation of microbial viruses, and evaluation of viral community function from genomic

sequences. Microbiome 8, 1–23. https://doi.org/10.1186/s40168-020-00867-0

Kooij, D. van der, Wielen, P.W.J.J. van der, 2013. Microbial Growth in Drinking Water

Supplies. IWA Publishing.

Kothari, A., Roux, S., Zhang, H., Prieto, A., Soneja, D., Chandonia, J., Spencer, S., 2021.

Ecogenomics of Groundwater Phages Suggests Niche Differentiation Linked to Specific

Environmental Tolerance. mSystems 6, 1–12. https://doi.org/10.1128/mSystems.00537-21

Kotlarz, N., Raskin, L., Zimbric, M., Errickson, J., Lipuma, J.J., Caverly, L.J., 2019.

Retrospective Analysis of Nontuberculous Mycobacterial Infection and Monochloramine

Disinfection of Municipal Drinking Water in Michigan. mSphere 4, 1–8.

https://doi.org/10.1128/mSphere.00160-19

Krishna, K.C.B., Sathasivan, A., Listowski, A., 2020. Influence of treatment processes and

disinfectants on bacterial community compositions and opportunistic pathogens in a full-

scale recycled water distribution system. J. Clean. Prod. 274, 123034.

https://doi.org/10.1016/j.jclepro.2020.123034

Langenfeld, K., Chin, K., Roy, A., Wigginton, K., Duhaime, M.B., 2021. Comparison of

ultrafiltration and iron chloride flocculation in the preparation of aquatic viromes from

contrasting sample types. PeerJ 9:e11111. https://doi.org/10.7717/peerj.11111

Langmead, B., Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2. Nat. Methods 9,

357–9. https://doi.org/10.1038/nmeth.1923

Leinonen, R., Sugawara, H., Shumway, M., 2011. The Sequence Read Archive 39, 2010–2012.

https://doi.org/10.1093/nar/gkq1019

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,

Durbin, R., Subgroup, 1000 Genome Project Data Processing, 2009. The Sequence

Alignment/Map format and SAMtools. Bioinformatics 25, 2078–2079.

https://doi.org/10.1093/bioinformatics/btp352

Liao, Y., Smyth, G.K., Shi, W., 2014. featureCounts : an efficient general purpose program for

assigning sequence reads to genomic features. Bioinformatics 30, 923–930.

https://doi.org/10.1093/bioinformatics/btt656

Liao, Y., Smyth, G.K., Shi, W., 2013. The Subread aligner : fast , accurate and scalable read

mapping by seed-and-vote. Nucleic Acids Res. 41. https://doi.org/10.1093/nar/gkt214

Liu, G., Verberk, J.Q.J.C., Van Dijk, J.C., 2013. Bacteriology of drinking water distribution

systems: An integral and multidimensional review. Appl. Microbiol. Biotechnol. 97, 9265–

9276. https://doi.org/10.1007/s00253-013-5217-y

Liu, L., Xing, X., Hu, C., Wang, H., Lyu, L., 2019. Chemosphere Effect of sequential UV / free

chlorine disinfection on opportunistic pathogens and microbial community structure in

simulated drinking water distribution systems. Chemosphere 219, 971–980.

https://doi.org/10.1016/j.chemosphere.2018.12.067

Ma, L., Li, B., Jiang, X.T., Wang, Y.L., Xia, Y., Li, A.D., Zhang, T., 2017. Catalogue of

antibiotic resistome and host-tracking in drinking water deciphered by a large scale survey.

Microbiome 5, 154. https://doi.org/10.1186/s40168-017-0369-0

Ma, L., Li, B., Zhang, T., 2019. New insights into antibiotic resistome in drinking water and

management perspectives : A metagenomic based study of small-sized microbes. Water

Res. 152, 191–201. https://doi.org/10.1016/j.watres.2018.12.069

Malki, K., Rosario, K., Sawaya, N.A., Székely, A.J., Tisza, M.J., Breitbart, M., 2020.

Prokaryotic and Viral Community Composition of Freshwater Springs in Florida, USA.

mBiom 11, e00436-20. https://doi.org/https://doi.org/10.1128/mBio.00436-20

Marks, T., Sharp, R., 2000. Bacteriophages and biotechnology : a review. J. Chem. Technol.

Biotechnol. 17, 6–17.

Mellbye, B.L., Bottomley, P.J., Sayavedra-soto, L.A., 2015. Nitrite-Oxidizing Bacterium

Nitrobacter winogradskyi Produces N -Acyl-Homoserine Lactone Autoinducers. AEM.

https://doi.org/10.1128/AEM.01103-15

Moon, K., Kim, S., Kang, I., Cho, J., 2020. Viral metagenomes of Lake Soyang , the largest

freshwater lake in South Korea. Sci. Data 1–6. https://doi.org/10.1038/s41597-020-00695-9

Nayfach, S., Camargo, A.P., Schulz, F., Eloe-fadrosh, E., Roux, S., Kyrpides, N.C., 2020.

CheckV assesses the quality and completeness of metagenome-assembled viral genomes.

Nat. Biotechnol. https://doi.org/10.1038/s41587-020-00774-7

NCBI, N.C. for B.I., 2019. BLAST.

NCBI, N.C. for B.I., 2016. UniVec.

Nurk, S., Meleshko, D., Korobeynikov, A., Pevzner, P.A., 2017. metaSPAdes : a new versatile

metagenomic assembler 824–834. https://doi.org/10.1101/gr.213959.116.4

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Wagner,

P.R.M., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner,

H., 2020. vegan: Community Ecology Package.

Paez-Espino, D., Eloe-Fadrosh, E.A., Pavlopoulos, G.A., Thomas, A.D., Huntemann, M.,

Mikhailova, N., Rubin, E., Ivanova, N.N., Kyrpides, N.C., 2016. Uncovering Earth's

virome. Nature 536, 425–430. https://doi.org/10.1038/nature19094

Parmar, K.M., Gaikwad, S.L., Dhakephalkar, P.K., Kothari, R., Singh, R.P., 2017. Intriguing

interaction of bacteriophage-host association: An understanding in the era of omics. Front.

Microbiol. 8. https://doi.org/10.3389/fmicb.2017.00559

Percival, S.L., Wyn-Jones, P., 2013. Methods for the Detection of Waterborne Viruses, Second

Edi. ed, Microbiology of Waterborne Diseases: Microbiological Aspects and Risks: Second

Edition. Elsevier. https://doi.org/10.1016/B978-0-12-415846-7.00022-6

Petrovich, M.L., Zilberman, A., Kaplan, A., Eliraz, G.R., Wang, Y., Langenfeld, K., Duhaime,

M., Wigginton, K., Poretsky, R., Avisar, D., Wells, G.F., 2020. Microbial and Viral

Communities and Their Antibiotic Resistance Genes Throughout a Hospital Wastewater

Treatment System. Front. Microbiol. 11, 1–13. https://doi.org/10.3389/fmicb.2020.00153

Pinto, A., Schroeder, J., Lunn, M., Sloan, W., Raskin, L., 2014. Spatial-Temporal Survey and

Occupancy-Abundance Modeling To Predict Bacterial Community Dynamics in the

Drinking Water Microbiome. MBio 5, e01135-14. https://doi.org/10.1128/mBio.01135-14

Pinto, A.J., Xi, C., Raskin, L., 2012. Bacterial community structure in the drinking water

microbiome is governed by filtration processes. Environ. Sci. Technol. 46, 8851–8859.

https://doi.org/10.1021/es302042t

Potgieter, S., Pinto, A., Sigudu, M., du Preez, H., Ncube, E., Venter, S., 2018. Long-term spatial

and temporal microbial community dynamics in a large-scale drinking water distribution

system with multiple disinfectant regimes. Water Res. 139, 406–419.

https://doi.org/10.1016/j.watres.2018.03.077

Potgieter, S.C., Dai, Z., Venter, S.N., Sigudu, M., Pinto, A.J., 2020. Microbial Nitrogen

Metabolism in Chloraminated Drinking. mSphere2 5. https://doi.org/https://doi

.org/10.1128/mSphere.00274-20. Editor

Prest, E.I., Weissbrodt, D.G., Hammes, F., Loosdrecht, M.C.M. Van, Vrouwenvelder, J.S., 2016.

Long-Term Bacterial Dynamics in a Full-Scale Drinking Water Distribution System. PLoS

One 11, 1–20. https://doi.org/10.1371/journal.pone.0164445

Rao, C., Waghmare, S., Lakhe, S., 1981. Detection of viruses in drinking water by concentration

on magnetic iron oxide. Appl. Environ. Microbiol. 42, 421–426.

Ren, J., Song, K., Deng, C., Ahlgren, N.A., Fuhrman, J.A., Li, Y., Xie, X., Sun, F., 2018.

Identifying viruses from metagenomic data by deep learning. Quant. Biol. 8, 64–77.

https://doi.org/10.1007/s40484-019-0187-4

Roeselers, G., Coolen, J., Wielen, P.W.J.J. Van Der, Jaspers, M.C., Atsma, A., Graaf, B. De,

Schuren, F., 2015. Microbial biogeography of drinking water: patterns in phylogenetic

diversity across space and time. Environ. Microbiol. 17, 2505–2514.

https://doi.org/10.1111/1462-2920.12739

Rohwer, F., Prangishvili, D., Lindell, D., 2009. Roles of viruses in the environment. Environ.

Microbiol. 11, 2771–2774. https://doi.org/10.1111/j.1462-2920.2009.02101.x

Roux, S., Adriaenssens, E.M., Dutilh, B.E., Koonin, E. V, Kropinski, A.M., Krupovic, M., Kuhn,

J.H., Lavigne, R., Brister, J.R., Varsani, A., Amid, C., Aziz, R.K., Bordenstein, S.R., Bork,

P., Breitbart, M., Cochrane, G.R., Daly, R.A., Desnues, C., Duhaime, M.B., Emerson, J.B.,

Enault, F., Fuhrman, J.A., Hingamp, P., Hugenholtz, P., Hurwitz, B.L., Ivanova, N.N.,

Labonté, J.M., Lee, K., Malmstrom, R.R., Martinez-garcia, M., Mizrachi, I.K., Ogata, H.,

Rodriguez-valera, F., Rosario, K., Schriml, L., Schulz, F., Steward, G.F., Sullivan, M.B.,

Sunagawa, S., Suttle, C.A., Temperton, B., Tringe, S.G., Thurber, R.V., Webster, N.S.,

Whiteson, K.L., Wilhelm, S.W., Wommack, K.E., Woyke, T., Wrighton, K.C., Yilmaz, P.,

Yoshida, T., 2019. Minimum Information about an Uncultivated Virus Genome (MIUViG

). Nat. Biotechnol. 37, 29–37. https://doi.org/10.1038/nbt.4306

Roux, S., Bolduc, B., 2017. stampede-clustergenomes.

https://doi.org/https://bitbucket.org/MAVERICLab/stampede-clustergenomes/src/master/

Roux, S., Brum, J.R., Dutilh, B.E., Sunagawa, S., Duhaime, M.B., Loy, A., Poulos, B.T.,

Solonenko, N., Lara, E., Poulain, J., Pesant, S., Kandels-lewis, S., Dimier, C., Picheral, M.,

Searson, S., Cruaud, C., Alberti, A., Duarte, C.M., Gasol, J.M., Vaqué, D., Tara Oceans

Coordinators, Bork, P., Acinas, S.G., Wincker, P., Sullivan, M.B., 2016. Ecogenomics and

potential biogeochemical impacts of globally abundant ocean viruses. Nat. Publ. Gr. 537,

689–693. https://doi.org/10.1038/nature19366

Roux, S., Enault, F., Hurwitz, B.L., Sullivan, M.B., 2015a. VirSorter: Mining viral signal from

microbial genomic data. PeerJ 2015, 1–20. https://doi.org/10.7717/peerj.985

Roux, S., Enault, F., Robin, A., Ravet, V., Personnic, S., Theil, S., Colombet, J., Sime-Ngando,

T., Debroas, D., 2012. Assessing the Diversity and Specificity of Two Freshwater Viral

Communities through Metagenomics. PLoS One 7.

https://doi.org/10.1371/journal.pone.0033641

Roux, S., Hallam, S.J., Woyke, T., Sullivan, M.B., 2015b. Viral dark matter and virus – host

interactions resolved from publicly available microbial genomes. Elife 1–20.

https://doi.org/10.7554/eLife.08490

Russel, M., Model, P., 1986. The Role of Thioredoxin in Filamentous Phage Assembly. J. Biol.

Chem. 261, 14997–15005. https://doi.org/10.1016/S0021-9258(18)66819-X

Samhan, F.A., Kronlein, M.R., Fakher, U., Kronlein, C., Robert, D., Hashsham, S.A., 2015.

Detection and Occurrence of Indicator Organisms and Pathogens 87, 883–900.

https://doi.org/10.2175/106143015X14338845155147

Saparbaevna, M., Aizhan, A., Turmagambetova, S., Gennadievich, P., Andrey, A.,

Bogoyavlenskiy, P., Berezin, V.E., 2017. Comparative study of viromes from freshwater

samples of the Ile-Balkhash region of Kazakhstan captured through metagenomic analysis.

VirusDisease 28, 18–25. https://doi.org/10.1007/s13337-016-0353-5

Sevillano, M., Vosloo, S., Cotto, I., Dai, Z., Jiang, T., Santiago, J.M., Padilla, I.Y., Rosario-

pabon, Z., Velez, C., Cordero, F., Alshawabkeh, A., Gu, A., Pinto, A.J., 2021. Spatial-

temporal targeted and non-targeted surveys to assess microbiological composition of

drinking water in Puerto Rico following Hurricane Maria 13.

https://doi.org/10.1016/j.wroa.2021.100123

Shaffer, M., Borton, M.A., McGivern, B.B., Zayed, A.A., La Rosa, S.L. 0003 3527 8101,

Solden, L.M., Liu, P., Narrowe, A.B., Rodríguez-Ramos, J., Bolduc, B., Gazitúa, M.C.,

Daly, R.A., Smith, G.J., Vik, D.R., Pope, P.B., Sullivan, M.B., Roux, S., Wrighton, K.C.,

2020. DRAM for distilling microbial metabolism to automate the curation of microbiome

function. Nucleic Acids Res. 48, 8883–8900. https://doi.org/10.1093/nar/gkaa621

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N.,

Schwikowski, B., Ideker, T., 2003. Cytoscape : A Software Environment for Integrated

Models of Biomolecular Interaction Networks. Genome Res. 2498–2504.

https://doi.org/10.1101/gr.1239303.metabolite

Shi, P., Jia, S., Zhang, X., Zhang, T., Cheng, S., Li, A., 2012. Metagenomic insights into

chlorination effects on microbial antibiotic resistance in drinking water. Water Res. 47,

111–120. https://doi.org/10.1016/j.watres.2012.09.046

Skvortsov, T., Leeuwe, C. De, Quinn, J.P., Mcgrath, J.W., Christopher, C., Allen, R., Mcelarney,

Y., Watson, C., Arkhipova, K., Lavigne, R., 2016. Metagenomic Characterisation of the

Viral Community of Lough Neagh , the Largest Freshwater Lake in Ireland. PLoS One 11,

e0140361. https://doi.org/10.1371/journal.pone.0150361

Stamps, B.W., Leddy, M.B., Plumlee, M.H., Hasan, N.A., Colwell, R.R., Spear, J.R., 2018.

Characterization of the microbiome at the world’s largest potable water reuse facility. Front.

Microbiol. 9, 1–16. https://doi.org/10.3389/fmicb.2018.02435

Stanish, L.F., Hull, N.M., Robertson, C.E., Harris, J.K., Stevens, J., Spear, J.R., Pace, N.R.,

2016. Factors Influencing Bacterial Diversity and Community Composition in Municipal

Drinking Waters in the Ohio River Basin, USA. PLoS One 11, 1–21.

https://doi.org/10.1371/journal.pone.0157966

Suttle, C.A., 2007. Marine viruses — major players in the global ecosystem. Nat. Rev.

Microbiol. 5. https://doi.org/10.1038/nrmicro1750

Team, R.C., 2021. R: A language and environment for statistical computing.

https://doi.org/https://www.r-project.org/

Team, Rs., 2020. RStudio: Integrated Development for R.

https://doi.org/https://www.rstudio.com/

Tseng, C.H., Chiang, P.W., Shiah, F.K., Chen, Y.L., Liou, J.R., Hsu, T.C., Maheswararajah, S.,

Saeed, I., Halgamuge, S., Tang, S.L., 2013. Microbial and viral metagenomes of a

subtropical freshwater reservoir subject to climatic disturbances. ISME J. 7, 2374–2386.

https://doi.org/10.1038/ismej.2013.118

Tsukatani, Y., Hirose, Y., Harada, J., Misawa, N., Mori, K., Inoue, K., 2015. Complete Genome

Sequence of the Bacteriochlorophyll b -Producing Photosynthetic Bacterium Blastochloris

viridis. Genome Announc. 3, 5–6. https://doi.org/10.1128/genomeA.01006-15.Copyright

Vaerewijck, M.J.M., Huys, G., Carlos, J., Swings, J., 2005. Mycobacteria in drinking water

distribution systems : ecology and significance for human health. FEMS Microbiol. Rev.

29, 911–934. https://doi.org/10.1016/j.femsre.2005.02.001

Wang, H., Proctor, C.R., Edwards, M.A., Pryor, M., Domingo, J.W.S., Ryu, H., Camper, A.K.,

Olson, A., Pruden, A., 2014. Microbial Community Response to Chlorine Conversion in a

Chloraminated Drinking Water Distribution System. ES&T 48, 10624–10633.

https://doi.org/10.1021/es502646d

Warwick-Dugdale, J., Buchholz, H.H., Allen, M.J., Temperton, B., 2019. Host-hijacking and

planktonic piracy : how phages command the microbial high seas. Virol. J. 1, 1–13.

https://doi.org/10.1186/s12985-019-1120-1

Williamson, S.J., Allen, L.Z., Lorenzi, H.A., Fadrosh, D.W., Brami, D., Thiagarajan, M.,

McCrow, J.P., Tovchigrechko, A., Yooseph, S., Venter, J.C., 2012. Metagenomic

Exploration of Viruses throughout the Indian Ocean. PLoS One 7.

https://doi.org/10.1371/journal.pone.0042047

Yang, Y., Gu, X., Te, S.H., Goh, S.G., Mani, K., He, Y., Gin, K.Y.H., 2019. Occurrence and

distribution of viruses and picoplankton in tropical freshwater bodies determined by flow

cytometry. Water Res. 149, 342–350. https://doi.org/10.1016/j.watres.2018.11.022

Ye, X.Y., Xing, M., Zhang, Y.L., Xiao, W.Q., Huang, X.N., Cao, Y.G., Gu, K.D., 2012. Real-

Time PCR Detection of Enteric Viruses in Source Water and Treated Drinking Water in

Wuhan , China. Curr. Microbiol. 65, 244–253. https://doi.org/10.1007/s00284-012-0152-1

Yilmaz, P., Kottmann, R., Field, D., Knight, R., Cole, J.R., Amaral-zettler, L., Gilbert, J.A.,

Karsch-mizrachi, I., Johnston, A., Cochrane, G., Vaughan, R., Hunter, C., Park, J.,

Morrison, N., Rocca-serra, P., Sterk, P., Arumugam, M., Bailey, M., Baumgartner, L.,

Birren, B.W., Blaser, M.J., Bonazzi, V., Booth, T., Bork, P., Bushman, F.D., Buttigieg,

P.L., Chain, P.S.G., Charlson, E., Costello, E.K., Huot-creasy, H., Dawyndt, P., Desantis,

T., Fierer, N., Fuhrman, J.A., Gallery, R.E., Gevers, D., Gibbs, R.A., Gil, I.S., Gonzalez, A.,

Gordon, J.I., Guralnick, R., Hankeln, W., Highlander, S., Hugenholtz, P., Jansson, J., Kau,

A.L., Kelley, S.T., Kennedy, J., Knights, D., Koren, O., Kuczynski, J., Kyrpides, N.,

Larsen, R., Lauber, C.L., Legg, T., Ley, R.E., Lozupone, C.A., Ludwig, W., Lyons, D.,

Maguire, E., Methé, B.A., Meyer, F., Muegge, B., Nakielny, S., Nelson, K.E., Nemergut,

D., Neufeld, J.D., Newbold, L.K., Oliver, A.E., Pace, N.R., Palanisamy, G., Peplies, J.,

Ravel, J., Relman, D.A., Assunta-sansone, S., Schloss, P.D., Schriml, L., Sinha, R., Smith,

M.I., Sodergren, E., Spor, A., Stombaugh, J., Tiedje, J.M., Ward, D. V, Weinstock, G.M.,

Wendel, D., White, O., Whiteley, A., Wilke, A., Wortman, J.R., Yatsunenko, T., Glöckner,

F.O., 2011. Minimum information about a marker gene sequence (MIMARKS) and

minimum information about any (x) sequence (MIxS) specifications. Nat. Biotechnol. 29,

415–420. https://doi.org/10.1038/nbt.1823

Zaouri, N., Jumat, M.R., Cheema, T., Hong, P.Y., 2020. Metagenomics-based evaluation of

groundwater microbial profiles in response to treated wastewater discharge. Environ. Res.

180, 108835. https://doi.org/10.1016/j.envres.2019.108835

Zhang, H., Chang, F., Shi, P., Ye, L., Zhou, Q., Pan, Y., Li, A., 2019. Antibiotic Resistome

Alteration by Different Disinfection Strategies in a Full-Scale Drinking Water Treatment

Plant Deciphered by Metagenomic Assembly. Environ. Sci. Technol. 53, 2141–2150.

https://doi.org/10.1021/acs.est.8b05907

Zimmerman, A.E., Varona, C.H.-, Needham, D.M., John, S.G., Worden, A.Z., Sullivan, M.B.,

Waldbauer, J.R., Coleman, M.L., 2020. Metabolic and biogeochemical consequences of

viral infection in aquatic ecosystems. Nat. Rev. Microbiol. 18.

https://doi.org/10.1038/s41579-019-0270-x

998