Vernalization-triggered expression of the antisense transcript *COOLAIR* is mediated by *CBF* genes

Myeongjune Jeon[1,2][†], Goowon Jeong[1,2][†], Yupeng Yang[3], Xiao Luo[4], Daesong Jeong[1,2], Jinseul Kyung[1,2], Youbong Hyun[1,2], Yuehui He[3,4,*], and Ilha Lee[1,2,*]

[1] School of Biological Sciences, Seoul National University, Seoul 08826, Korea
[2] Research Center for Plant Plasticity, Seoul National University, Seoul 08826, Korea
[3] Shanghai Center for Plant Stress Biology & National Key Laboratory for Plant Molecular Genetics, CAS Center for Excellence in Molecular Plant Sciences, Chinese Academy of Sciences (CAS), Shanghai 201602, China
[4] Peking University Institute of Advanced Agricultural Sciences, Weifang 261325, China

[*] For correspondence: ilhalee@snu.ac.kr, yhhe@pku.edu.cn
[†] These authors contributed equally to this work.

Abstract

To synchronize flowering time with spring, many plants undergo vernalization, a floral-promotion process triggered by exposure to long-term winter cold. In *Arabidopsis thaliana*, this is achieved through cold-mediated epigenetic silencing of the floral repressor, *FLOWERING LOCUS C* (*FLC*). *COOLAIR*, a cold-induced antisense RNA transcribed from the *FLC* locus, has been proposed to facilitate *FLC* silencing. Here, we show that C-repeat (CRT)/dehydration-responsive elements (DREs) at the 3′-end of *FLC* and CRT/DRE-binding factors (CBFs) are required for cold-mediated expression of *COOLAIR*. CBFs bind to CRT/DREs at the 3′-end of *FLC*, both *in vitro* and *in vivo*, and CBF levels increase gradually during vernalization. Cold-induced *COOLAIR* expression is severely impaired in *cbfs* mutants in which all *CBF* genes are knocked-out. Conversely, *CBF*-overexpressing plants show increased *COOLAIR* levels even at warm temperatures. We show that *COOLAIR* is induced by CBFs during early stages of vernalization but COOLAIR levels decrease in later phases as *FLC* chromatin transitions to an inactive state to which CBFs can no longer bind. We also demonstrate that *cbfs* and *FLC*$_{\Delta COOLAIR}$ mutants exhibit a normal vernalization response despite their inability to activate *COOLAIR* expression during cold, revealing that *COOLAIR* is not required for the vernalization process.

Introduction

Appropriate timing of flowering provides evolutionary advantage of plant reproductive success. As sessile organisms, plants have evolved mechanisms through which seasonal cues coordinate the transition to flowering. One of the significant environmental factors affecting flowering time of plants adapted to temperate climates is the temperature changes during the seasons, and plants have evolved complex sensory mechanisms to monitor the surrounding temperature to properly control the timing of flowering and tolerate thermal stress (*Went, 1953; Penfield, 2008; Ding and Yang, 2022*).

Cold acclimation and vernalization are two responses of plants to low temperatures. Cold acclimation is generally initiated by a short period of non-freezing cold exposure and increases the frost tolerance of plants (*Weiser, 1970; Gilmour et al., 1988; Guy, 1990; Thomashow, 1999; Chinnusamy et al., 2007*). The three C-REPEAT(CRT)/DEHYDRATION-RESPONSIVE ELEMENT (DRE) BINDING FACTORs (CBFs) and their encoding genes serve as signaling hubs for cold acclimation in *Arabidopsis thaliana* (*Stockinger et al., 1997; Medina et al., 1999; Thomashow, 2010; Ding et al., 2019*). When exposed to low temperatures, the transcription of *CBFs* is rapidly promoted by a group of cold-signal transducers, including the Ca^{2+}/calmodulin-binding proteins, CALMODULIN-BINDING TRANSCRIPTION ACTIVATORs (CAMTAs) (*Doherty et al., 2009; Kim et al., 2013; Kidokoro et al., 2017*); the clock proteins, CIRCADIAN CLOCK-ASSOCIATED 1 (CCA1) and LATE-ELONGATED HYPOCOTYL (LHY) (*Dong et al., 2011*); and the brassinosteroid-responsive proteins, BRASSINAZOLE-RESISTANT 1 (BZR1) and CESTA (CES) (*Eremina et al., 2016; Li et al., 2017*). In addition, cold also enhances the

activity of CBF proteins by triggering degradation of BASIC TRANSCRIPTION FACTOR 3s (BTF3s) and HISTONE DEACETYLASE 2C (HD2C) which prevent CBFs from activating their targets (*Ding et al., 2018; J. Park et al., 2018*), or by reducing oxidized CBFs, which increases active CBF monomers (*Lee et al., 2021*). Low temperature-induced CBFs, in turn, activate the expression of *co*ld-*r*egulated (*COR*) genes by binding to CRT/DREs in their promoters (*Stockinger et al., 1997; Medina et al., 1999*). Diverse arrays of cryoprotective proteins encoded by *COR* genes allow plants to overcome freezing stress (*Gilmour et al., 1988; Hajela et al., 1990; Thomashow et al., 1997; Shinozaki et al., 2003; Shi et al., 2018; Ding et al., 2019*).

In contrast to cold acclimation, vernalization, a floral-promotion process that occurs during winter, requires an extended cold period (*Napp-Zinn, 1955; Chouard, 1960; Michaels and Amasino, 2000*). This allows plants to synchronize the timing of flowering with favorable spring conditions. Vernalization in *A. thaliana* is mainly achieved by silencing the floral repressor gene, *FLOWERING LOCUS C* (*FLC*) (*Michaels and Amasino, 1999; Sheldon et al., 1999; Sheldon et al., 2000; Michaels and Amasino, 2001*). *FLC* encodes a MADS-box protein that represses the expression of floral activator genes, *SUPPRESSOR OF OVEREXPRESSION OF CO 1* (*SOC1*) and *FLOWERING LOCUS T* (*FT,* encoding florigen), by directly binding to their promoter regions (*Lee et al., 2000; Michaels et al., 2005; Helliwell et al., 2006*). In Arabidopsis winter annuals, such as San Feliu-2, Löv-1, and Sweden (SW) ecotypes, flowering is prevented by the high expression of *FLC* before exposure to winter cold (*Lee et al., 1993; Shindo et al., 2005; S. Park et al., 2018*). This is caused by the strong transcriptional activation of *FLC* by the FRIGIDA (FRI) supercomplex, which recruits general transcription factors and several chromatin modifiers (*Michaels and Amasino, 1999; Sheldon et al., 1999; Johanson et al., 2000; Choi et al., 2011; Li et al., 2018*).

Prior to vernalization, *FLC* chromatin is highly enriched with active histone marks such as histone H3 acetylation and trimethylation of Lys4 or Lys36 at H3 (H3K4me3/H3K36me3) (*Bastow et al., 2004; Yang et al., 2014*). In contrast, prolonged cold exposure results in gradual deacetylation of *FLC* chromatin and concomitant removal of H3K4me3 and H3K36me3 from the *FLC* (*Bastow et al., 2004; Yang et al., 2014; Nishio et al., 2016*). Additionally, VP1/ABI3-LIKE 1 (VAL1) and VAL2 recruit Polycomb Repressive Complex 2 (PRC2) onto *FLC* chromatin, thereby accumulating the repressive histone mark, H3 Lys27 trimethylation (H3K27me3), in the nucleation region around the first exon and intron of *FLC* (*Sung and Amasino, 2004b; Wood et al., 2006; De Lucia et al., 2008; Angel et al., 2011; Nishio et al., 2016; Qüesta et al., 2016; Yuan et al., 2016*). Subsequently, upon returning to warm temperatures, H3K27me3 marks are spread over the entire *FLC* chromatin region by LIKE HETEROCHROMATIN PROTEIN 1 (LHP1), which ensures stable *FLC* suppression and renders plants competent to flower (*Mylne et al., 2006; Sung et al., 2006; Yang et al., 2017*).

Several long-term cold-induced factors have been shown to play crucial roles in the epigenetic silencing of *FLC*. *VERNALIZATION INSENSITIVE 3* (*VIN3*) family genes, which are upregulated by prolonged cold, encode plant homeodomain (PHD) proteins that recognize H3K9me2 enriched in *FLC* chromatin during vernalization (*Sung and Amasino, 2004b; Kim and Sung, 2013, 2017a*). These proteins mediate the recruitment of PRC2 and the subsequent deposition of H3K27me3 at the *FLC* nucleation region (*Sung and Amasino, 2004b; De Lucia et al., 2008; Kim and Sung, 2013*). In addition, vernalization-induced long non-coding RNAs (lncRNAs) are involved in such histone modifications. *COLDAIR* and *COLDWRAP*, the two lncRNAs transcribed from the first intron and promoter region of *FLC*, respectively, are required for H3K27me3 deposition in response to long-

term cold (*Heo and Sung, 2011; Kim and Sung, 2017b*). *COLDAIR* and *COLDWRAP* are also thought to affect the formation of the intragenic chromatin loop at the *FLC*, which may be a part of the *FLC* silencing mechanism (*Kim and Sung, 2017b*). Unlike *COLDAIR* and *COLDWRAP*, which are transcribed in the sense direction of *FLC*, another lncRNA, *COOLAIR,* is an antisense transcript expressed from the 3′-end of *FLC* (*Swiezewski et al., 2009*). The gradual accumulation of *COOLAIR* reaches a maximum within a few weeks of cold exposure, whereas *COLDAIR* and *COLDWRAP* showed peaks at a later phase of vernalization (*Csorba et al., 2014; Kim and Sung, 2017b*). *COOLAIR* was reported to remove active histone marks from *FLC* chromatin (*Liu et al., 2007; Csorba et al., 2014; Fang et al., 2020; Xu, Fang, et al., 2021; Zhu et al., 2021*). Particularly, in summer annuals, phase-separated RNA-processing complexes favor co-transcriptional proximal polyadenylation of *COOLAIR* (*Marquardt et al., 2014; Wang et al., 2014; Fang et al., 2019; Wu et al., 2020; Xu, Wu, et al., 2021*). The *COOLAIR*-processing machinery exhibits transient and dynamic interactions with an H3K4 demethylation complex, leading to *FLC* suppression at warm temperatures (*Liu et al., 2007; Fang et al., 2020; Xu, Fang, et al., 2021*). *COOLAIR* is also likely to be involved in reducing H3K36me3 at the *FLC* during vernalization process (*Csorba et al., 2014*). *COOLAIR* was reported to promote the sequestration of the FRI complex from the *FLC* promoter by condensing it into phase-separated nuclear bodies (*Zhu et al., 2021*). This has been suggested to cause the inactivation of *FLC,* which is probably accompanied by the silencing of *FLC* chromatin through the removal of H3K36me3. However, a previous study has raised the issue that *COOLAIR* appears not to be necessary for vernalization (*Helliwell et al., 2011; Luo et al., 2019*).

Compared with the signaling pathway of cold acclimation, how a long-term cold signal is

transduced to trigger the induction of *VIN3* and lncRNAs is less well understood. It is marginally

known that some chromatin modifiers, NAC WITH TRANSMEMBRANE MOTIF 1-LIKE 8

(NTL8), and CCA1/LHY act as positive regulators of *VIN3* during the vernalization process (*Kim*

et al., 2010; Finnegan et al., 2011; Zhao et al., 2020; Kyung et al., 2022). However, little is known

about the upstream regulators of *COOLAIR* required for cold-induction. Recent reports have

shown that an NAC domain-containing protein, NTL8, and the WRKY transcription factor,

WRKY63, can bind to the promoter of *COOLAIR* and activate its expression (*Zhao et al., 2021;*

Hung et al., 2022). However, whether NTL8 and WRKY63 are necessary for the full extent of

COOLAIR induction during vernalization has not been thoroughly addressed. In this study, we

identified that CRT/DREs at the 3′-end of the *FLC* are required for the long-term cold response of

COOLAIR. Additionally, we show that CBFs, which accumulate during the long-term winter cold,

act as upstream regulators of *COOLAIR* during vernalization.

Results

A CRT/DRE-binding factor, CBF3, directly binds to CRT/DREs at the 3′-end of the *FLC*

The proximal promoter region of *COOLAIR* is highly conserved among *FLC* orthologs from *A. thaliana* relatives (*Castaings et al., 2014*). Thus, we assumed that the cis-element conferring a long-term cold response would exist within that block. A comparison of the region near the transcriptional start site (TSS) of *COOLAIR* revealed the conservation of two CRT/DREs among six *FLC* orthologs from five related species of the Brassicaceae family (*Figure 1A*). CRT/DRE, the core sequence of which is CCGAC, is a regulatory element that imparts cold- or drought-responsive gene expression (*Baker et al., 1994*). *CBF1*, *2*, and *3* encode APETALA2 (AP2) domain-containing proteins that can bind to CRT/DRE and are usually present in the promoters of cold- and drought-responsive genes (*Stockinger et al., 1997; Medina et al., 1999*). Consistent with this, the Arabidopsis cistrome database and the genome-wide chromatin immunoprecipitation sequencing (ChIP-seq) results from a previous study suggested that CBFs bind to the 3′-end sequence of *FLC* containing CRT/DREs (*Figure 1—figure supplement 1*; *O'Malley et al., 2016; Song et al., 2021*).

We performed an electrophoretic mobility shift assay (EMSA) to confirm this binding using probes harboring CRT/DREs (named DRE1 or 2) from the *COOLAIR* promoter. The mobility of these two Cy5-labeled probes was retarded by maltose-binding protein (MBP)-fused CBF3 (*Figure 1B, Figure 1—figure supplement 2*). The band shift was competed out by adding an excess amount of unlabeled DRE1 or 2 oligonucleotides. In contrast, competitors containing mutant forms of DRE1 or 2 (DRE1^m or 2^m, respectively) failed to compete (*Figure 1B*). We also

tested the binding of other CRT/DRE-like sequences in the *FLC* locus. Two (DREa and b) were

present in the *FLC* promoter, while the other two (DREc and d), were present in the first and last

exons, respectively (**Figure 1—figure supplement 2**). Only DREc competed with the band shift

caused by the CBF3-DRE1 interaction (**Figure 1B, Figure 1—figure supplement 2**). The presence

or absence of bases that determine the binding affinity between CBFs and CRT/DRE could explain

the differences in CBF3-binding patterns among CRT/DRE-like sequences at the *FLC* locus

(*Maruyama et al., 2004*).

Subsequently, a chromatin immunoprecipitation (ChIP) assay using a *CBF3*-overexpressing

transgenic plant, *pSuper:CBF3-myc* (*Liu et al., 2017*), was conducted to determine whether CBF3

is associated with the *FLC* region containing CRT/DREs *in vivo*. Consistent with the *in vitro* results,

CBF3-myc protein was enriched at the 3′-end of *FLC* and the first exon where DREc was located

(**Figure 1C**). Without vernalization (NV), CBF3-myc was highly enriched at both the 5′- and 3′-

ends of *FLC* (P1 and P3), indicating that CBF3 protein can activate *FLC* under warm conditions,

as previously reported (*Seo et al., 2009*). However, enrichment in the P1 region rapidly disappeared

during the vernalization period. In contrast, enrichment on P3 was maintained until 10 days of

vernalization (10V) and was subsequently reduced. This is coincident with the expression pattern

of *COOLAIR*, which declines during the late phase of vernalization (*Csorba et al., 2014*).

COOLAIR is one of the CBF regulons in Arabidopsis

CBF genes are rapidly and transiently induced upon exposure to cold (*Medina et al., 1999*). CBF

regulons, which are CBF-targeted genes, are correspondingly up- or down-regulated after cold

intervals of a few hours (*Gilmour et al., 1998; Liu et al., 1998; Fowler and Thomashow, 2002*).

As *COOLAIR* contains CBF3-binding sites in its promoter (**Figure 1**), we analyzed whether

COOLAIR shows CBF regulon-like expression. The expression of genes assigned to the CBF

regulon is up- or down-regulated by *CBF* overexpression under warm conditions (*Park et al.,*

2015). Similarly, we found that the total transcript level of *COOLAIR* increased in *CBF3*-

overexpressing transgenic plants (*pSuper:CBF3-myc*) grown at room temperature (22°C)

compared to that in the wild type, Columbia-0 (Col-0) (**Figure 2A**). Such upregulation was mainly

due to the type I.i or type II.i *COOLAIR* isoforms. As the levels of both proximal (type I) and distal

(type II) variants of *COOLAIR* were higher in the *CBF3* overexpressor than in the wild type, it is

likely that *COOLAIR* transcription, instead of the 3′-processing events, is affected by CBF3

(**Figure 2A;** *Liu et al., 2010; Marquardt et al., 2014*). It has been reported that the targets of CBF1

or 2 are not entirely identical to those of CBF3, although *CBF* genes show high sequence similarity

(*Novillo et al., 2007*). Thus, we compared the expression levels of all *COOLAIR* isoforms in

transgenic plants overexpressing *CBF1*, *2*, or *3* (*Seo et al., 2009*). Overall, all CBF-overexpressing

plants showed increased levels of *COOLAIR* (**Figure 2—figure supplement 1**). However, each

CBF-overexpressing plant showed a subtly different effect on *COOLAIR* expression. The *CBF2*

overexpressor showed the highest level of the proximal *COOLAIR* isoform, whereas the *CBF3*

overexpressor showed the highest level of distal *COOLAIR*. In contrast, the *CBF1*-overexpressing

plants showed a slightly smaller increase in proximal *COOLAIR* levels and a negligible increase

in distal *COOLAIR*, compared to those in the wild type, Wassilewskija-2 (Ws-2). Such results

imply that the three CBFs regulate *COOLAIR* transcription subtly differently.

Another characteristic of CBF regulons is that their expression is activated by a day of cold

exposure (*Park et al., 2015*). Thus, we analyzed if *COOLAIR* is also induced by short-term cold

exposure. We treated wild-type (*FRI* Col) plants with 0, 0.5, 1, 1.5, 2, 3, 6, and 24 h of cold (4°C),

then measured the levels of total *COOLAIR*. The results showed that cold treatment for less than

3 h failed to induce *COOLAIR* expression, although *CBF* transcripts reached a peak at 3 h of cold

exposure (**Figures 2B and 3A;** *Gilmour et al., 1998*). However, after 6 h of cold treatment, when

CBF3 protein levels peaked, *COOLAIR* was strongly induced (**Figures 2B and 3D**). Thus, short-

term cold-triggered *COOLAIR* expression dynamics were similar to those of other CBF-targeted

genes; that is, the expression was highly increased after *CBF* transcript levels reached a peak upon

cold exposure (*Gilmour et al., 1998; Fowler and Thomashow, 2002*). To confirm whether *CBFs*

are responsible for the rapid cold response of *COOLAIR*, we examined *COOLAIR* induction after

a day of cold treatment in the wild type and *cbfs-1* mutant, in which all three *CBFs* were knocked

out (**Figure 2C;** *Jia et al., 2016*). All isoforms, as well as total *COOLAIR*, failed to be induced by

short-term cold exposure in the *cbfs* mutant, although a drastic increase was observed in the wild

type. Thus, these results strongly suggest that *CBFs* are required for *COOLAIR* induction, even

under a short period of cold exposure.

CBFs accumulate during vernalization

We subsequently investigated whether *CBFs* are also responsible for vernalization-induced

COOLAIR activation. It has been reported that *COOLAIR* is gradually upregulated as the cold

period persists and peaks after 2–3 weeks of vernalization (*Csorba et al., 2014*). However, most

studies on *CBF* expression have been performed within a few days since the function of *CBFs* has

been analyzed only in the context of short-term cold (*Gilmour et al., 1998; Liu et al., 1998; Medina*

et al., 1999; McKhann et al., 2008). Therefore, we investigated the expression patterns of *CBFs*

before and after long-term cold exposure to determine the correlation between the expression of

CBFs and *COOLAIR* during vernalization. As previously shown, the levels of all three *CBFs*

peaked within 3 h of cold exposure and then decreased rapidly (***Figure 3A***). However, *CBF* levels

increased again as the cold period was prolonged (> 20V), suggesting that both short-term and

vernalizing cold treatments upregulated *CBF* expression (***Figure 3B***).

Because *CBFs* exhibit rhythmic oscillations (*Dong et al., 2011*), we also investigated whether

the rhythmic expression of *CBFs* is affected by vernalization. Wild-type (Col-0) plants were

collected every 4 h under a short-day cycle (SD, 8 h light/16 h dark) under both NV and 40V

conditions. As shown in ***Figure 3C***, although the rhythmic pattern of each *CBF* was variable, the

overall transcript levels of all three *CBFs* were much higher at 40V than under NV. Consistent with

the transcript level, the level of CBF3 protein in vernalized plants was higher than that in NV

plants during all circadian cycles (***Figure 3—figure supplement 1***).

To verify whether the increased transcription of *CBF3* led to protein accumulation, we

measured the level of CBF3 at each time-point of cold treatment using the *pCBF3:CBF3-myc* plant

(*Jia et al., 2016*). Following the rapid and transient induction of *CBF3* under short-term cold

conditions (***Figure 3A;*** *Medina et al., 1999*), CBF3-myc protein levels peaked within 6 h of cold

exposure and then decreased (***Figure 3D***). Thus, the peak of the *CBF3* transcript and that of the

CBF3 protein showed a gap of few hours. Notably, the peak of the CBF3 protein correlated with

the time when *COOLAIR* was rapidly induced (***Figures 2B and 3D***). During vernalization, the

level of CBF3-myc protein gradually declined until 10V, but then increased again (***Figure 3E***).

Thus, the level at 40V is similar to that at 1V. When the plants were transferred to room temperature

for 4 d after 40V (40VT4), the protein level rapidly decreased. These results suggest that CBFs are

responsible for the progressive upregulation of *COOLAIR*, at least during the early phase of

vernalization (*Csorba et al., 2014*).

***CBFs* are involved in vernalization-induced *COOLAIR* expression**

To clarify whether *CBFs* are responsible for long-term cold-mediated *COOLAIR* induction, we

quantified *COOLAIR* levels in wild type and *cbfs* mutant during vernalization. As shown in

Figures 2B and 2C, the *COOLAIR* level was considerably increased by a day of low-temperature

treatment in the wild type but quickly declined afterward (***Figure 4A***). The *COOLAIR* level

increased again after 4V and reached a secondary peak at approximately 10V to 20V. It then

decreased during the remaining vernalization period, as similar to previous results (*Csorba et al.,*

2014). The expression dynamics of *COOLAIR* were similar to those of two well-known CBF

targets, *COR15A* and *RD29A*, although their levels remained high until the end of the vernalization

period (***Figure 4—figure supplement 1***). In contrast to the wild type plants, the *cbfs* mutant

showed severely reduced *COOLAIR* expression, as well as *COR15A* and *RD29A* levels, during all

the period of vernalization (***Figure 4A, Figure 4—figure supplement 1***). Analysis of RNA

sequencing (RNA-seq) data from previous studies also revealed that the levels of total *COOLAIR,*

and proximal and distal *COOLAIR* variants were reduced in the *cbfs* mutant in Col-0 or SW

ecotype background when exposed to either short-term (3 and 24 h) or long-term (14V) cold, which

supports our results (***Figure 4—figure supplement 2;*** *Zhao et al., 2016; S. Park et al., 2018*). All

of these data support that *CBFs* are necessary to fully induce *COOLAIR* in the early phase of

vernalization.

Consistent with the reduced *COOLAIR* levels in *cbfs* mutants, the *CBF3* overexpressor,

pSuper:CBF3-myc, showed a much higher *COOLAIR* level than the wild type (Col-0), throughout

the vernalization period (***Figure 4B***). It is noteworthy that *COOLAIR* expression in *pSuper:CBF3-*

myc was further upregulated by cold, especially during the early vernalization phase (10V), and

then suppressed during the late phase. This result implies that both the transcriptional and post-

transcriptional regulations of CBFs are involved in the long-term cold response of *COOLAIR*.

A recent study showed that the first seasonal frost ($< 0°C$) during the winter season strongly

induces *COOLAIR* expression (*Zhao et al., 2021*). Therefore, we tested whether *CBFs* are also

required for the strong induction of *COOLAIR* triggered by freezing temperatures. Wild-type and

cbfs plants were subjected to NV, 10V, and 20V, with or without an additional 8 h of −1°C freezing

treatment. Irrespective of the pre-treatment with non-freezing cold, the freezing temperature

increased the levels of both *COOLAIR* and *CBF* in the wild type (***Figure 4C, Figure 4—figure***

supplement 3). However, the *cbfs* mutants showed a much smaller increase in the *COOLAIR* level

than the wild type when exposed to 8 h of sub-zero cold after 10V and 20V (***Figure 4C***). In

particular, *COOLAIR* levels were not elevated by freezing treatment in NV *cbfs*. Thus, *CBFs* seem

responsible for both the gradual increase of *COOLAIR* during vernalization and the strong

COOLAIR induction triggered by freezing temperatures.

CRT/DREs at the 3′-end of *FLC* are necessary for CBFs-mediated induction of *COOLAIR*

during vernalization.

Since CBF3 could bind to the CRT/DREs in the first exon (DREc) and the 3′-end of *FLC* (DRE1

and 2) (***Figures 1B and 1C***), we investigated whether CRT/DRE is responsible for the CBF-

mediated long-term cold response of *COOLAIR*. To address this, we performed an *A. thaliana*

protoplast transfection assay using the *35S:CBF3-HA* effector construct and the

pCOOLAIR$_{DRE}$:LUC reporter construct, in which the 1-kb wild-type *COOLAIR* promoter was

fused to the luciferase reporter gene. The luciferase activity assay showed that CBF3-HA protein

activated the transcription of the *COOLAIR* promoter (***Figure 5A***). In contrast, CBF3-HA failed

to increase luciferase activity when the mutant version of the *COOLAIR* promoter with mutations

in the DRE1 and 2 sequences (*pCOOLAIR$_{DRE}{}^{m}$:LUC*) was used as a reporter. Additionally, co-

transfection of the *35S:CBF3-HA* effector and *pFLC:LUC* reporter construct, which harbors the

1-kb sequence with the promoter and first exon of *FLC*, did not show increased luciferase activity.

These results strongly indicate that CBF3 can activate *COOLAIR* transcription through DRE1 and

2 located in the *COOLAIR* promoter, but not *via* the first exon of *FLC*.

To further determine whether the 3′-end sequence of *FLC* is required for cold-induction of

COOLAIR expression, we measured *COOLAIR* expression before and after vernalization in the

FLC$_{ΔCOOLAIR}$ mutant (*FLC$_{ΔCOOLAIR-1}$*), that lacks a 324-bp portion of the *COOLAIR* promoter region

(*Luo et al., 2019*). Reverse transcription PCR (RT-PCR) analysis showed that neither proximal nor

distal *COOLAIR* variants were detected in this mutant, and the levels did not increase after 14-d

cold exposure (***Figure 5B***). Using the CRISPR-Cas9 system, we generated additional *FLC$_{ΔCOOLAIR}$*

mutant lines, *FLC$_{ΔCOOLAIR-3,}$* and *FLC$_{ΔCOOLAIR-4}$* (***Figure 5B; Figure 5—figure supplement 1***).

FLC$_{ΔCOOLAIR-3}$ has a 685-bp deletion in the *COOLAIR* promoter region where both CRT/DREs are

located. As expected, this mutant failed to express *COOLAIR* even after 14V, which is similar to

the *FLC$_{ΔCOOLAIR-1}$* mutant. In contrast, the *FLC$_{ΔCOOLAIR-4}$* mutant, which bears a 301-bp deletion in

the distal *COOLAIR* promoter region (upstream of CRT/DREs) and a 13-bp sequence replacement

downstream of CRT/DREs, showed a normal vernalization-induced increase in all *COOLAIR*

variants, similar to the wild type. This strongly suggests that CRT/DREs located at the *COOLAIR*

promoter are critical for CBF-mediated *COOLAIR* induction during vernalization. Notably, the

low-abundance convergent antisense transcripts (*CAS*; *Zhao et al., 2021*) were similarly reduced

by 14V in all genotypes, i.e., the wild type and *FLC$_{\Delta COOLAIR}$* mutants (***Figure 5B***), suggesting their

regulation is independent of *COOLAIR*.

Although the common *COOLAIR* TSS was eliminated in *FLC$_{\Delta COOLAIR-1}$* (***Figures 5B***), the

proximal *COOLAIR* isoform (type I.i) was still slightly detected by quantitative PCR (qPCR),

probably due to alternative TSSs (***Figure 5C***). However, the expression of the proximal *COOLAIR*

variant in the *FLC$_{\Delta COOLAIR}$* mutant was not induced throughout the vernalization period, while the

wild type showed type I.i *COOLAIR* peaks at 1V and 20V (***Figure 5C***). Furthermore, *CBF3*

overexpression did not significantly increase type I.i *COOLAIR* levels in the *FLC$_{\Delta COOLAIR-1}$* mutant,

while it caused a strong increase in proximal *COOLAIR* in the wild-type background (Col-0)

(***Figure 5—figure supplement 2***). Taken together, our results strongly suggest that the *COOLAIR*

promoter region containing the two CRT/DREs is necessary for the long-term cold response of

COOLAIR.

CBFs-mediated *COOLAIR* induction during vernalization is not required for *FLC* silencing

Finally, we investigated the role of *COOLAIR* induced by CBFs in the vernalization-triggered *FLC*

silencing process. *COOLAIR* was reported to facilitate the removal of H3K36me3 from *FLC*

chromatin during vernalization, whereas the other lncRNAs, *COLDAIR* and *COLDWRAP*, are

required for H3K27me3 deposition (*Csorba et al., 2014; Berry and Dean, 2015; Kim and Sung,*

2017b; Zhu et al., 2021). Therefore, we compared the enrichment of H3K4me3, H3K36me3, and

H3K27me3 on *FLC* chromatin in non-vernalized or vernalized wild-type and *cbfs* plants. As

reported (*Bastow et al., 2004; Sung and Amasino, 2004b; Yang et al., 2014*), the enrichments of

H3K4me3 and H3K36me3 were reduced, but that of H3K27me3 was increased in *FLC* chromatin

at 40V in the wild type (***Figures 6A to 6C***). However, although *COOLAIR* levels were significantly

reduced in the *cbfs* mutant (***Figure 4A***), vernalization-mediated reductions in H3K4me3 and

H3K36me3 levels and an increase in H3K27me3 levels in the *cbfs* mutant were comparable to

those in the wild type (***Figures 6A to 6C***). This result suggests that CBF-mediated *COOLAIR*

induction is not required for vernalization-induced epigenetic changes on *FLC* chromatin.

Consistent with the epigenetic changes, the *FLC* level in the *cbfs* mutant was gradually

suppressed during the vernalization process, and the suppression was maintained after 40VT4

similar to that observed in the wild type (***Figure 6D***). Such result is consistent with the RNA-seq

data previously reported using *cbfs* mutants in an SW ecotype (*S. Park et al., 2018*). RNA-seq data

analysis showed that the *cbfs* mutant in the SW background exhibited reduced *FLC* levels after

14V, similar to the wild type (***Figure 4—figure supplement 2***).

Because our results indicate that *COOLAIR* induction is not required for *FLC* silencing and

the function of *COOLAIR* in vernalization-induced *FLC* silencing is still debated (*Helliwell et al.,*

2011; Csorba et al., 2014; Zhao et al., 2021; Zhu et al., 2021), we analyzed whether the loss-of-

function mutations *FLC$_{\Delta COOLAIR\text{-}1}$*, and *FLC$_{\Delta COOLAIR\text{-}3}$* result in any defect in the vernalization

response. In contrast to the transgenic lines used in previous reports, *FLC$_{\Delta COOLAIR\text{-}1}$* and

FLC$_{\Delta COOLAIR\text{-}3}$, which have small deletions in the *COOLAIR* promoter region, are unlikely to cause

any unexpected effects on *FLC* chromatin. *FLC$_{\Delta COOLAIR\text{-}4}$*, which has a deletion outside of the

CRT/DRE region, exhibited a gradual decrease and eventual silencing of *FLC* by long-term cold,

similar to the wild type (**Figure 6E**). Interestingly, both $FLC_{\Delta COOLAIR-1}$ and $FLC_{\Delta COOLAIR-3}$ mutants

exhibited similar decreases and silencing of *FLC* by vernalization (**Figures 5B, 6D, and 6E**).

These results show that *FLC* is normally silenced by vernalization regardless of *COOLAIR*

induction by cold.

Consistent with this, the *cbfs* mutant, compared to the wild type, did not show any difference

in the reduction of *FLC* levels upon exposure to a freezing cold for 8 h after NV, 10V, and 20V

(**Figure 6F**). However, the same freezing cold treatment caused the failure of strong induction of

COOLAIR in the *cbfs* (**Figure 4C**). These results also support that CBF-mediated *COOLAIR*

induction is not required for *FLC* silencing by vernalization.

As expected from the *FLC* levels, vernalization-mediated promotion of flowering was

comparable in the wild type and *cbfs* (**Figures 7A and 7B**), indicating that *cbfs* exhibits a normal

vernalization response. Consistently, the flowering time of $FLC_{\Delta COOLAIR-1}$ and $FLC_{\Delta COOLAIR-3}$ was

accelerated by vernalization, similar to that of the wild type (**Figures 7A, 7C, and 7D**). These

results show that *FLC* silencing occurs regardless of whether *COOLAIR* is induced during a cold

exposure that results in vernalization. In summary, our data strongly support that CBFs,

upregulated by long-term cold exposure, induce *COOLAIR* expression (**Figure 8**); however, cold-

mediated *COOLAIR* induction is not required for *FLC* silencing during vernalization.

Discussion

Winter is challenging for sessile plants: cold weather makes reproduction difficult, and frost threatens survival. Cold acclimation and vernalization, the two types of low-temperature memories, allow plants to overcome this harsh season by making them more viable during winter or flowering in the subsequent warm spring (*Thomashow, 1999; Michaels and Amasino, 2000*). Although it is well understood how plants recognize and remember short-term cold exposure to increase frost hardiness, there is still much to know about the long-term cold sensing mechanism that establishes the ability to flower.

FLC orthologs from a range of *A. thaliana* relatives express the vernalization-induced antisense lncRNA, *COOLAIR* (*Castaings et al., 2014*). As the proximal sequence blocks within the *COOLAIR* promoters are conserved, they are likely targets of cold sensor modules. This study showed that CBF proteins act as cold sensors binding to the CRT/DREs in the conserved promoter region, thereby activating the transcription of *COOLAIR*. Based on the data presented here, we propose a working model (***Figure 8***) in which a prolonged low-temperature environment gradually increases the levels of CBF proteins, which activate *COOLAIR* transcription in the early phase of vernalization by binding to CRT/DREs at the 3′-end of *FLC*. However, as cold period extended, CBF proteins are excluded from the *COOLAIR* promoter, probably because silencing of *FLC* chromatin occurs regardless of CBFs level. This may account for a decrease in *COOLAIR* levels during the later phase of vernalization. Consistent with this, a previous report showed a continuous increase in the *COOLAIR* level throughout the vernalization period in the *vin3* mutant, which has a defect in the maintenance of *FLC* silencing (*Swiezewski et al., 2009*). Thus, VIN3-dependent

Polycomb silencing is at least partially involved in the release of CBF proteins from the *COOLAIR* promoter.

Although CBFs are upregulated by cold to trigger *COOLAIR* expression, CBF-mediated *COOLAIR* induction during vernalization is not solely due to increased CBF levels because the *CBF3* overexpression lines still showed *COOLAIR* induction by vernalization (***Figure 4B***). This may indicate that additional factors participate in the regulation of CBFs function. As aforementioned, it has previously been reported that low temperatures cause functional activation of CBFs through the degradation of BTF3s and HD2C, which maintains an epigenetically inactive state of downstream targets at warm temperatures (*Ding et al., 2018; J. Park et al., 2018*). Moreover, cold triggers the monomerization of CBF proteins, thereby promoting their functions (*Lee et al., 2021*). Similarly, such an additional regulatory mechanism of CBF function may be enhanced by long-term winter cold to fine-tun *COOLAIR* expression.

In addition, other thermosensors are likely to facilitate *COOLAIR* induction by vernalization. Recently, it was suggested that NTL8, an NAC domain-containing transcription factor, acts as an upstream regulator of *COOLAIR* (*Zhao et al., 2021*). NTL8 also binds to the *COOLAIR* promoter region, which is not far from CBF-binding sites (*O'Malley et al., 2016; Zhao et al., 2021*). *COOLAIR* is highly expressed in both dominant mutant alleles, *ntl8-Ds*, and *NTL8*-overexpressing transgenic plants at warm temperatures. Thus, it has been proposed that slow accumulation of the NTL8 protein, due to reduced dilution during slower growth at low temperatures, causes *COOLAIR* induction during vernalization. In addition, a WRKY transcription factor, WRKY63, has been suggested as another inducer of *COOLAIR* transcription during vernalization (*Hung et al., 2022*). WRKY63 is enriched in the *COOLAIR* promoter region, and *COOLAIR* expression is

reduced in the *wrky63* mutant, suggesting that WRKY63 functions as a *COOLAIR* activator.

Unlike NTL8, both transcript and protein levels of *WRKY63* were upregulated by vernalization,

which may lead to an increase in *COOLAIR* expression. Thus, it would be worthwhile to elucidate

whether NTL8, WRKY63, and CBFs synergistically activate *COOLAIR* transcription during

vernalization. The reason why vernalization-triggered *COOLAIR* induction requires more than one

thermosensor is unclear. Presumably, multiple thermosensory modules ensure precise and robust

COOLAIR expression during winter, when the temperature fluctuates.

Transcription of *COOLAIR* has been proposed to govern the *FLC* chromatin environment in

both warm and cold temperatures (*Csorba et al., 2014; Fang et al., 2020; Xu, Wu, et al., 2021*).

Previous studies have inferred that *COOLAIR* is required to remove H3K36me3 from *FLC*

chromatin, especially under low-temperature conditions. This argument is supported by the

delayed H3K36me3 removal in vernalized *FLC-TEX* transgenic plants, where the *COOLAIR*

promoter is substituted by the *rbcs3B* terminator sequence (*Csorba et al., 2014; Marquardt et al.,*

2014; Wang et al., 2014). In addition, *COOLAIR* is reported to promote condensation of FRI upon

cold exposure (*Zhu et al., 2021*). Thus, FRI is sequestered from the *FLC* promoter as a nuclear

condensate, which causes the transcriptional repression of *FLC*. Therefore, it has been suggested

that vernalization-induced *COOLAIR* prevents the activation of *FLC* by the FRI complex and

causes epigenetic silencing of *FLC* chromatin through a decrease of H3K36me3 during the

vernalization process.

However, there is some controversy as to whether *COOLAIR* is required for vernalization and

the associated *FLC* silencing. In contrast to the *FLC-TEX* lines, some T-DNA insertion lines which

lack *COOLAIR* induction by long-term cold exposure show a normal vernalization response

(*Helliwell et al., 2011*). Likewise, the results from this study show that *COOLAIR* transcription is

not required for the epigenetic silencing of *FLC*. The *cbfs* mutant, which exhibits severely reduced

COOLAIR induction, shows relatively normal vernalization responses such that flowering time is

fairly well accelerated and active histone marks, H3K36me3 and H3K4me3, are reduced in *FLC*

chromatin while the repressive histone mark, H3K27me3, is increased in *FLC* chromatin similar

to the wild type (**Figures 6 and 7**). Consistently, $FLC_{\Delta COOLAIR-1}$, and $FLC_{\Delta COOLAIR-3}$ mutants, which

have almost undetectable levels of *COOLAIR*, exhibit both *FLC* suppression and flowering

acceleration by vernalization similar to that seen in the wild type (**Figures 6 and 7**). Moreover, the

freezing treatment mimicking the first frost did not affect *FLC* suppression in the *cbfs*, although

the mutant showed defects in CBF-mediated *COOLAIR* induction (**Figures 4C and 6F**). It is

noteworthy that our finding that vernalization and the associated *FLC* silencing do not require

COOLAIR expression were obtained using *cbfs* and $FLC_{\Delta COOLAIR}$ mutant lines rather than the

transgenic lines used in previous studies. Thus, we can exclude the possibility that transgenes

somehow provide cryptic promoters or cause unexpected epigenetic effects.

COOLAIR is highly conserved among Arabidopsis relatives (*Castaings et al., 2014*); thus, our

finding that *COOLAIR* is not required for vernalization-induced *FLC* silencing is surprising. There

are several caveats in the studies of *COOLAIR* function. First, different experimental schemes may

affect the vernalization response differently. A recent study of plants grown in condition with

fluctuating temperatures designed to mimic natural environments may result in a role for

COOLAIR in vernalization-mediated *FLC* repression (*Zhao et al., 2021*), and perhaps fluctuating

temperatures that mimic field conditions may have different effects on the vernalization response

of *cbfs* or $FLC_{\Delta COOLAIR}$ mutants as well. However, the experiments in most studies that claimed a

role for *COOLAIR* in vernalization were performed in laboratory conditions with controlled

temperatures as in this study. If indeed *COOLAIR* does have a role in vernalization in specific

environmental conditions, the lack of a role for *COOLAIR* in vernalization in other conditions such

as those in this study may be due to functional redundancy between *COOLAIR* and other *FLC*

repressors such that other repressors, such as *COLDAIR*, obscure the effect of *COOLAIR* on *FLC*

silencing, as has been proposed before (*Castaings et al., 2014*). This issue should be addressed in

future studies.

Although short-term and long-term cold memories elicit different developmental responses,

both are triggered by the same physical environment. Therefore, it is not surprising that a common

signaling network governs both processes. *CBF* genes have been highlighted for their roles in cold

acclimation (*Gilmour et al., 1998; Liu et al., 1998; Jia et al., 2016; Zhao et al., 2016*). CBFs

rapidly accumulate upon short-term cold exposure, thereby activating the transcription of cold-

responsive genes, such as *COR15A* and *RD29A* (*Stockinger et al., 1997; Liu et al., 1998*).

Interestingly, CBFs are also highly accumulated during vernalization and mediate long-term cold-

induced *COOLAIR* expression. Hence, our results indicate that the cold signaling pathways that

establish the short-term and long-term cold responses are not sharply distinguishable, as suggested

previously (*Sung and Amasino, 2004a; Seo et al., 2009*).

Materials and Methods

Plant materials and treatments

Arabidopsis thaliana Columbia-0 (Col-0), Wassilewskija-2 (Ws-2), and Col:*FRI*^{Sf2} (*FRI* Col) were used as the wild types in this study. "Wild type" in this paper mainly refers to *FRI* Col unless otherwise specified. The *FRI* Col (*Lee et al., 1994*), *cbfs-1*, *pCBF3:CBF3-myc* (*Jia et al., 2016*), *pSuper:CBF3-myc* (*Liu et al., 2017*), *FLC$_{ΔCOOLAIR-1}$* (*Luo et al., 2019*), *35S:CBF1*, *35S:CBF2*, and *35S:CBF3* (*Gilmour et al., 2004*) have been previously described. *FRI cbfs-1* was generated by crossing the *FRI* Col with *cbfs-1* stated above. *FRI FLC$_{ΔCOOLAIR-1}$* was generated by crossing *FRI* Col with *FLC$_{ΔCOOLAIR-1}$* in the Col-0 background, as described above. *pSuper:CBF3-myc FLC$_{ΔCOOLAIR-1}$* was generated by crossing *pSuper:CBF3-myc* with *FLC$_{ΔCOOLAIR-1}$*. For all experiments using plant materials, seeds were sown on half-strength Murashige & Skoog (MS) medium (Duchefa, Haarlem, Netherlands) containing 1% sucrose and 1% plant agar (Duchefa) and stratified at 4°C for 3 days. Plants were germinated and grown at 22°C under a short-day (SD) cycle (8 h light/16 h dark) with cool white fluorescent lights (100 μmol m^{-2} s^{-1}). For the vernalization treatment, the plants were grown at 22°C for the prescribed period and then transferred to a 4°C growth chamber under an SD cycle. To adjust the developmental stage, the plants vernalized for 40 days (40V), 30 days (30V), 20 days (20V), 10 days (10V), 4 days (4V), and 1 day (1V) were transferred to 4°C after being grown at 22°C for 10, 12, 14, 15, and 16 days, respectively. Non-vernalized (NV) plants were grown at 22°C for 16 days. For 0. 0.5, 1, 1.5, 2, 3, 6, and 24 h of cold treatment, the plants were grown at 22°C for 16 days before the 4°C treatment. For the freezing treatment, NV, 10V, and 20V plants were transferred to −1°C under dark condition

497   for 8 h (F). To measure flowering time of vernalized plants, plants were transferred to 22°C under

498   a long-day (LD) cycle (16 h light/8 h dark) directly after the vernalization treatment.

Generation of $FLC_{\Delta COOLAIR}$ lines by CRISPR-Cas9

The construction of $FLC_{\Delta COOLAIR-3}$ and $FLC_{\Delta COOLAIR-4}$ are as previously described (*Luo et al., 2019*).

Briefly, two pairs of sgRNAs were designed for the deletion of *COOLAIR* promoter regions (5′-

CTTCACAGTGAAGAAGCCTA-3′ and 5′-AAATGCACTCTTACGTAACG-3′ for $FLC_{\Delta COOLAIR-}$

$_3$; 5′-TTATCCTAAACGCGTATGGT-3′ and 5′-CGTAGTTCCGTCATCCATGA-3′ for

$FLC_{\Delta COOLAIR-4}$), and sgRNA-Cas9 cassettes were introduced into Arabidopsis by floral dipping.

Homozygous deletion mutants were screened in T_2 generation, and *sgRNA-Cas9* free mutants were

further isolated in T_3 generation.

Plasmid construction

To generate the *35S:CBF3-HA* construct, the entire coding sequence of CBF3 was amplified using

PCR and cloned into the *HBT-HA-NOS* plasmid (*Yoo et al., 2007*). For $pCOOLAIR_{DRE}:LUC$ and

$pCOOLAIR_{DRE}{}^{m}:LUC$ construction, the 1-kb sequence containing the *COOLAIR* promoter, the 3′-

untranslated region (UTR), and the last exon of *FLC* was amplified by PCR and cloned into the

LUC-NOS plasmid (*Hwang and Sheen, 2001*). To generate $pCOOLAIR_{DRE}{}^{m}:LUC$, two CRT/DREs

in the *COOLAIR* promoter, 5′-AGGTCGGT-3′ and 5′-ACCGACAT-3′, were replaced with 5′-

CGAGGTGT-3′ and 5′-TGAACCCA-3′, respectively. For *pFLC:LUC* construction, the 1-kb

sequence containing the promoter, 5′-UTR, and the first exon of *FLC* was amplified by PCR and

was cloned into the *LUC-NOS* plasmid.

Electrophoretic mobility shift assay (EMSA)

Maltose binding protein (MBP) and MBP-CBF3 recombinant fusion proteins were induced by 500

μM isopropyl β-D-thiogalactoside (IPTG) in the *E. coli* BL21 (DE3) strain. Cell extracts were

isolated with a buffer containing 20 mM Tris-HCl (pH 7.4), 1 mM EDTA, 200 mM NaCl, 10%

glycerol, 1 mM DTT, and 1 mM PMSF. Proteins were purified from cell extracts using MBPTrap™

HP column (GE Healthcare, Chicago, IL, USA) with ÄKTA FPLC system (Amersham Biosciences,

Amersham, UK). The Cy5-labeled probes and unlabeled competitors were generated by annealing

25 bp-length oligonucleotides. EMSA was performed as previously described with a few

modifications (*Seo et al., 2012*). For each EMSA reaction, 5 μM of protein and 100 nM of Cy5-

labeled probe were incubated at room temperature in a binding buffer containing 10 mM Tris-HCl

(pH 7.5), 1 mM EDTA, 50 mM NaCl, 5% glycerol, and 5 mM DTT. For the competition assay, the

reaction mixtures were incubated in the presence of 100-fold molar excess of each competitor. The

reaction mixtures were electrophoresed at 100 V after incubation. The Cy5 signals were detected

using WSE-6200H LuminoGraph II (ATTO, Amherst, NY, USA).

Chromatin immunoprecipitation (ChIP) assay

Approximately 4 g of seedlings grown on MS plates was collected and cross-linked using 1% (v/v)

formaldehyde. Nuclei were isolated from seedlings using a buffer containing 20 mM PIPES-KOH

(pH 7.6), 1 M hexylene glycol, 10 mM $MgCl_2$, 0.1 mM EGTA, 15 mM NaCl, 60 mM KCl, 0.5%

Triton-X, 5 mM β-mercaptoethanol, and 1× cOmplete™ EDTA-free Protease Inhibitor Cocktail

(Roche, Basel, Switzerland) as described previously with a few modifications (*Shu et al., 2014*).

Isolated nuclei were lysed with a buffer containing 50 mM Tris-HCl (pH 7.4), 150 mM NaCl, 1%

(v/v) Triton X-100, and 1% SDS (w/v), and were subsequently sonicated using a Digital Sonifier®

(Branson, Danbury, CT, USA). Sheared chromatin was diluted with a buffer containing 50 mM

Tris-HCl (pH 7.4), 150 mM NaCl, 1% (v/v) Triton X-100, and 1 mM EDTA. Immunoprecipitation

was performed by incubating sheared chromatin with Protein G Sepharose™ 4 Fast Flow beads

(GE Healthcare) and antibodies. Anti-myc (MBL, Woburn, MA, USA, #M192-3) and normal

mouse IgG_1 (Santa Cruz Biotechnology, Santa Cruz, CA, USA, #sc-3877) were used to detect

CBF3 protein enrichment at the *FLC* locus, and anti-H3K36me3 (Abcam, Cambridge, UK,

#Ab9050), H3K4me3 (Millipore, Bedford, MA, USA, #07-473), and anti-H3K27me3 (Millipore,

#07-449) were used to detect histone modifications on the *FLC* chromatin. DNA was extracted

with phenol:chloroform:isoamyl alcohol (25:24:1, v/v) or Chelex® 100 Resin according to the

manufacturer's instruction.

Gene expression analysis

Total RNA was isolated from ~100 mg of seedlings grown on MS agar plate using TRI Reagent®

(Sigma-Aldrich, St Louis, MO, USA) or TS™ Plant Mini Kit (Taeshin Bio Science, Gyeonggi-do,

Korea). cDNA was generated using 4 μg of total RNA, 5 units of recombinant DNase I (TaKaRa,

Kyoto, Japan), 200 units of RevertAid™ reverse transcriptase (Thermo Scientific, Waltham, MA,

USA), and buffer containing 0.25 mM dNTP and 0.1 μg oligo(dT)$_{18}$. Quantitative PCR analysis

was performed using iQ™ SYBR® Green Supermix (Bio-Rad, Hercules, CA, USA) and a CFX96

Real-Time PCR system (Bio-Rad).

To examine *COOLAIR* expression in the *COOLAIR* promoter deletions lines, Total RNAs

were extracted using the RNeasy® Plus Mini Kit (Qiagen, Hilden, Germany) according to the

manufacturers' instructions, followed by the digestion of residual genomic DNA by the gDNA

wiper (Vazyme, Nanjing, China). 1.0-μg RNA from each sample was taken for cDNA synthesis

using the HiScript® III 1st Strand cDNA Synthesis Kit (Vazyme) with the transcript-specific

primers (5′-TGGTTGTTATTTGGTGGTGTGAA-3′ for *COOLAIR* class I and 5′-

GCCCGACGAAGAAAAAGTAG-3′ for class II). Semi-quantitative PCR amplifications was

performed, followed by agarose gel separation of PCR products.

Immunoblotting

Total protein was isolated from ~100 mg of seedlings using a buffer containing 50 mM Tris-HCl

(pH 7.5), 100 mM NaCl, 10 mM MgCl$_2$, 1 mM EDTA, 10% glycerol, 1 mM PMSF, 1 mM DTT

and 1× cOmplete™ EDTA-free Protease Inhibitor Cocktail (Roche). Fifty micrograms of total

protein were loaded onto SDS-PAGE gels and separated by electrophoresis. The proteins were

transferred to PVDF membranes (Amersham Biosciences) and probed with an anti-myc (MBL,

#M192-3; 1:10000 dilution) antibody overnight at 4°C. The samples were then probed with

horseradish peroxidase (HRP)-conjugated anti-mouse IgG (Cell Signaling, Danvers, MA, USA,

#7076; 1:10000 dilution) antibody at room temperature. The signals were detected with

ImageQuant™ LAS 4000 (GE Healthcare) using WesternBright™ Sirius ECL solution (Advansta,

San Jose, CA, USA).

RNA sequencing (RNA-seq) data analysis

We retrieved two sets of RNA-seq data from the NCBI SRA-numbered SRP056607 and

SRP320943. SRP056607 contains raw reads from NV, 1V, and 14V wild-type and *cbfs* plants in a

Sweden-ecotype (SW) background. SRP3209943 contains reads from 0, 3, and 24 h cold-treated

wild-type and *cbfs* plants in a Col-0 background. The reads were aligned to the Arabidopsis TAIR

10 reference genome and annotated in ARAPORT 11 using STAR version 2.7.10a. Isoform

estimation was performed using Salmon version 1.6.0.

Luciferase assay using Arabidopsis protoplast

Protoplast isolation and transfection were performed as previously described with some

modifications (*Yoo et al., 2007*). Protoplasts were isolated from leaves of SD-grown wild-type

plant (Col-0) using a buffer containing 150 mg Cellulase Onozuka™ R-10 (Yakult, Tokyo, Japan),

50 mg Maceroenzyme™ R-10 (Yakult), 20 mM KCl, 20 mM MES-KOH (pH 5.6), 0.4 M D-

mannitol, 10 mM $CaCl_2$, 5 mM β-mercaptoethanol, and 0.1 g bovine serum albumin. For protoplast

transfection, 200 µg of plasmid DNA and isolated protoplasts were incubated in a buffer containing

0.1 M D-mannitol, 50 mM $CaCl_2$, and 20% (w/v) PEG. Luciferase activity in the protoplasts was

measured using the Luciferase Assay System (Promega, Madison, WI, USA) and MicroLumat

Plus LB96V microplate luminometer (Berthold Technologies, Bad Wildbad, Germany).

Supplementary data

Figure 1—figure supplement 1. CBF proteins directly bind to the sequence at the 3′-end of *FLC* in DNA affinity purification sequencing (DAP-seq).

Figure 1—figure supplement 2. CBF3 binds to DRE1 and DRE2 on the *COOLAIR* promoter and DREc on the first exon of *FLC.*

Figure 2—figure supplement 1. Overexpression of *CBF1*, *2*, or *3* causes an increase in *COOLAIR* level, but the effects on *COOLAIR* variants are subtly different.

Figure 3—figure supplement 1. Vernalization increases CBF3 protein levels with a minor effect on the daily rhythm.

Figure 4—figure supplement 1. CBF regulons show strongly reduced expression in *cbfs* mutants similar to *COOLAIR*.

Figure 4—figure supplement 2. Cold-triggered *COOLAIR* expression is impaired in *cbfs* mutants in Col-0 and Sweden-ecotype background.

Figure 4—figure supplement 3. Freezing temperature increases the levels of *CBFs*.

Figure 5—figure supplement 1. Genomic sequences near the deleted regions in two *COOLAIR* promoter deletion lines, $FLC_{\Delta COOLAIR\text{-}3}$ and $FLC_{\Delta COOLAIR\text{-}4}$.

Figure 5—figure supplement 2. Proximal *COOLAIR* level is not significantly increased by *CBF3* overexpression in the $FLC_{\Delta COOLAIR}$ mutant lacking CRT/DREs.

Figure 1—source data 1. Uncropped labelled gel image and the original image file for the EMSA result.

Figure 1—figure supplement 2—source data 1. Uncropped labelled gel images and the original image files for the EMSA results.

Figure 3—source data 1. Uncropped labelled blot images and the original image files for the

immunoblots.

Figure 3—figure supplement 1—source data 1. Uncropped labelled blot images and the original

image files for the immunoblots.

Figure 5—source data 1. Uncropped labelled gel images and the original image files for the RT-

PCR results.

Supplementary File 1. Oligonucleotide sequences used in this study.

Acknowledgements

We would like to thank Dr. Shuhua Yang (China Agricultural University, China) for providing the

cbfs-1, *pCBF3:CBF3-myc*, and *pSuper:CBF3-myc* seeds. This work was supported by the

Cooperative Research Program for Agriculture Science and Technology Development (No.

PJ01315201 and PJ01315401), Rural Development Administration, Republic of Korea. This work

was also supported by National Research Foundation of Korea (NRF) grants funded by the Korean

government (MSIT) (No. 2019R1A2C2004313 and NRF-2021R1A5A1032428). Y. He's

laboratory was funded by the grants from the National Natural Science Foundation of China (No.

31830049 and 31721001). M. Jeon and D. Jeong were supported by Brain Korea 21 Plus Project.

We would like to thank Editage (http://www.editage.co.kr) for English language editing.

References

Angel A, Song J, Dean C, Howard M (2011) **A Polycomb-based switch underlying quantitative epigenetic memory**. *Nature* **476:** 105-108. https://doi.org/10.1038/nature10241

Baker SS, Wilhelm KS, Thomashow MF (1994) **The 5'-region of *Arabidopsis thaliana cor15a* has *cis*-acting elements that confer cold-, drought- and ABA-regulated gene expression**. *Plant Mol Biol* **24:** 701-713. https://doi.org/10.1007/BF00029852

Bastow R, Mylne JS, Lister C, Lippman Z, Martienssen RA, Dean C (2004) **Vernalization requires epigenetic silencing of *FLC* by histone methylation**. *Nature* **427:** 164-167. https://doi.org/10.1038/nature02269

Berry S, Dean C (2015) **Environmental perception and epigenetic memory: Mechanistic insight through *FLC***. *Plant J* **83:** 133-148. https://doi.org/10.1111/tpj.12869

Castaings L, Bergonzi S, Albani MC, Kemi U, Savolainen O, Coupland G (2014) **Evolutionary conservation of cold-induced antisense RNAs of *FLOWERING LOCUS C* in *Arabidopsis thaliana* perennial relatives**. *Nat Commun* **5:** 4457. https://doi.org/10.1038/ncomms5457

Chinnusamy V, Zhu J, Zhu JK (2007) **Cold stress regulation of gene expression in plants**. *Trends Plant Sci* **12:** 444-451. https://doi.org/10.1016/j.tplants.2007.07.002

Choi K, Kim J, Hwang HJ, Kim S, Park C, Kim SY, Lee I (2011) **The FRIGIDA complex activates transcription of *FLC*, a strong flowering repressor in *Arabidopsis*, by recruiting chromatin modification factors**. *Plant Cell* **23:** 289-303. https://doi.org/10.1105/tpc.110.075911

Chouard P (1960) **Vernalization and its relations to dormancy**. *Annu Rev Plant Phys* **11:** 191-238. https://doi.org/10.1146/annurev.pp.11.060160.001203

Csorba T, Questa JI, Sun Q, Dean C (2014) **Antisense *COOLAIR* mediates the coordinated switching of chromatin states at *FLC* during vernalization**. *Proc Natl Acad Sci U S A* **111:** 16160-16165. https://doi.org/10.1073/pnas.1419030111

De Lucia F, Crevillen P, Jones AM, Greb T, Dean C (2008) **A PHD-Polycomb Repressive Complex 2 triggers the epigenetic silencing of *FLC* during vernalization**. *Proc Natl*

Acad Sci U S A **105:** 16831-16836. https://doi.org/10.1073/pnas.0808687105

Ding Y, Jia Y, Shi Y, Zhang X, Song C, Gong Z, Yang S (2018) **OST1-mediated BTF3L phosphorylation positively regulates CBFs during plant cold responses**. *EMBO J* **37**. https://doi.org/10.15252/embj.201798228

Ding Y, Shi Y, Yang S (2019) **Advances and challenges in uncovering cold tolerance regulatory mechanisms in plants**. *New Phytol* **222:** 1690-1704. https://doi.org/10.1111/nph.15696

Ding Y, Yang S (2022) **Surviving and thriving: How plants perceive and respond to temperature stress**. *Dev Cell* **57:** 947-958. https://doi.org/10.1016/j.devcel.2022.03.010

Doherty CJ, Van Buskirk HA, Myers SJ, Thomashow MF (2009) **Roles for *Arabidopsis* CAMTA transcription factors in cold-regulated gene expression and freezing tolerance**. *Plant Cell* **21:** 972-984. https://doi.org/10.1105/tpc.108.063958

Dong MA, Farré EM, Thomashow MF (2011) **CIRCADIAN CLOCK-ASSOCIATED 1 and LATE ELONGATED HYPOCOTYL regulate expression of the C-REPEAT BINDING FACTOR (CBF) pathway in *Arabidopsis***. *Proc Natl Acad Sci U S A* **108:** 7241-7246. https://doi.org/10.1073/pnas.1103741108

Eremina M, Unterholzner SJ, Rathnayake AI, Castellanos M, Khan M, Kugler KG, May ST, Mayer KF, Rozhon W, Poppenberger B (2016) **Brassinosteroids participate in the control of basal and acquired freezing tolerance of plants**. *Proc Natl Acad Sci U S A* **113:** E5982-E5991. https://doi.org/10.1073/pnas.1611477113

Fang X, Wang L, Ishikawa R, Li Y, Fiedler M, Liu F, Calder G, Rowan B, Weigel D, Li P, Dean C (2019) ***Arabidopsis* FLL2 promotes liquid-liquid phase separation of polyadenylation complexes**. *Nature* **569:** 265-269. https://doi.org/10.1038/s41586-019-1165-8

Fang X, Wu Z, Raitskin O, Webb K, Voigt P, Lu T, Howard M, Dean C (2020) **The 3' processing of antisense RNAs physically links to chromatin-based transcriptional control**. *Proc Natl Acad Sci U S A* **117:** 15316-15321. https://doi.org/10.1073/pnas.2007268117

Finnegan EJ, Bond DM, Buzas DM, Goodrich J, Helliwell CA, Tamada Y, Yun JY, Amasino RM, Dennis ES (2011) **Polycomb proteins regulate the quantitative induction of *VERNALIZATION INSENSITIVE 3* in response to low temperatures**. *Plant J* **65:** 382-391. https://doi.org/10.1111/j.1365-313X.2010.04428.x

Fowler S, Thomashow MF (2002) **Arabidopsis transcriptome profiling indicates that multiple regulatory pathways are activated during cold acclimation in addition to the CBF cold response pathway**. *Plant Cell* **14:** 1675-1690. https://doi.org/10.1105/tpc.003483

Gilmour SJ, Fowler SG, Thomashow MF (2004) **Arabidopsis transcriptional activators CBF1, CBF2, and CBF3 have matching functional activities**. *Plant Mol Biol* **54:** 767-781. https://doi.org/10.1023/B:PLAN.0000040902.06881.d4

Gilmour SJ, Hajela RK, Thomashow MF (1988) **Cold acclimation in *Arabidopsis thaliana***. *Plant Physiol* **87:** 745-750. https://doi.org/10.1104/pp.87.3.745

Gilmour SJ, Zarka DG, Stockinger EJ, Salazar MP, Houghton JM, Thomashow MF (1998) **Low temperature regulation of the *Arabidopsis* CBF family of AP2 transcriptional activators as an early step in cold-induced *COR* gene expression**. *Plant J* **16:** 433-442. https://doi.org/10.1046/j.1365-313x.1998.00310.x

Guy CL (1990) **Cold acclimation and freezing stress tolerance: Role of protein metabolism**. *Annu Rev Plant Physiol Plant Mol Biol* **41:** 187-223. https://doi.org/10.1146/annurev.pp.41.060190.001155

Hajela RK, Horvath DP, Gilmour SJ, Thomashow MF (1990) **Molecular cloning and expression of *cor* (*c*old-regulated) genes in *Arabidopsis thaliana***. *Plant Physiol* **93:** 1246-1252. https://doi.org/10.1104/pp.93.3.1246

Helliwell CA, Robertson M, Finnegan EJ, Buzas DM, Dennis ES (2011) **Vernalization-repression of Arabidopsis *FLC* requires promoter sequences but not antisense transcripts**. *PLoS One* **6:** e21513. https://doi.org/10.1371/journal.pone.0021513

Helliwell CA, Wood CC, Robertson M, James Peacock W, Dennis ES (2006) **The Arabidopsis FLC protein interacts directly *in vivo* with *SOC1* and *FT* chromatin and is part of a high-molecular-weight protein complex**. *Plant J* **46:** 183-192. https://doi.org/10.1111/j.1365-313X.2006.02686.x

Heo JB, Sung S (2011) **Vernalization-mediated epigenetic silencing by a long intronic noncoding RNA**. *Science* **331:** 76-79. https://doi.org/10.1126/science.1197349

Hung FY, Shih YH, Lin PY, Feng YR, Li C, Wu K (2022) **WRKY63 transcriptional activation of *COOLAIR* and *COLDAIR* regulates vernalization-induced flowering**. *Plant Physiol*.

https://doi.org/10.1093/plphys/kiac295

Hwang I, Sheen J (2001) **Two-component circuitry in *Arabidopsis* cytokinin signal transduction**. *Nature* **413:** 383-389. https://doi.org/10.1038/35096500

Jia Y, Ding Y, Shi Y, Zhang X, Gong Z, Yang S (2016) **The *cbfs* triple mutants reveal the essential functions of *CBFs* in cold acclimation and allow the definition of CBF regulons in *Arabidopsis***. *New Phytol* **212:** 345-353. https://doi.org/10.1111/nph.14088

Johanson U, West J, Lister C, Michaels S, Amasino R, Dean C (2000) **Molecular analysis of *FRIGIDA*, a major determinant of natural variation in *Arabidopsis* flowering time**. *Science* **290:** 344-347. https://doi.org/10.1126/science.290.5490.344

Kidokoro S, Yoneda K, Takasaki H, Takahashi F, Shinozaki K, Yamaguchi-Shinozaki K (2017) **Different cold-signaling pathways function in the responses to rapid and gradual decreases in temperature**. *Plant Cell* **29:** 760-774. https://doi.org/10.1105/tpc.16.00669

Kim DH, Sung S (2013) **Coordination of the vernalization response through a *VIN3* and *FLC* gene family regulatory network in *Arabidopsis***. *Plant Cell* **25:** 454-469. https://doi.org/10.1105/tpc.112.104760

Kim DH, Sung S (2017a) **The binding specificity of the PHD-finger domain of VIN3 moderates vernalization response**. *Plant Physiol* **173:** 1258-1268. https://doi.org/10.1104/pp.16.01320

Kim DH, Sung S (2017b) **Vernalization-triggered intragenic chromatin loop formation by long noncoding RNAs**. *Dev Cell* **40:** 302-312 e304. https://doi.org/10.1016/j.devcel.2016.12.021

Kim DH, Zografos BR, Sung S (2010) **Mechanisms underlying vernalization-mediated *VIN3* induction in Arabidopsis**. *Plant Signal Behav* **5:** 1457-1459. https://doi.org/10.4161/psb.5.11.13465

Kim Y, Park S, Gilmour SJ, Thomashow MF (2013) **Roles of CAMTA transcription factors and salicylic acid in configuring the low-temperature transcriptome and freezing tolerance of Arabidopsis**. *Plant J* **75:** 364-376. https://doi.org/10.1111/tpj.12205

Kyung J, Jeon M, Jeong G, Shin Y, Seo E, Yu J, Kim H, Park CM, Hwang D, Lee I (2022) **The two clock proteins CCA1 and LHY activate *VIN3* transcription during vernalization**

through the vernalization-responsive cis-element. *Plant Cell* **34:** 1020-1037. https://doi.org/10.1093/plcell/koab304

Lee ES, Park JH, Wi SD, Kang CH, Chi YH, Chae HB, Paeng SK, Ji MG, Kim WY, Kim MG, Yun DJ, Stacey G, Lee SY (2021) **Redox-dependent structural switch and CBF activation confer freezing tolerance in plants**. *Nat Plants* **7:** 914-922. https://doi.org/10.1038/s41477-021-00944-8

Lee H, Suh SS, Park E, Cho E, Ahn JH, Kim SG, Lee JS, Kwon YM, Lee I (2000) **The AGAMOUS-LIKE 20 MADS domain protein integrates floral inductive pathways in *Arabidopsis***. *Genes Dev* **14:** 2366-2376. https://doi.org/10.1101/gad.813600

Lee I, Bleecker A, Amasino R (1993) **Analysis of naturally occurring late flowering in *Arabidopsis thaliana***. *Mol Gen Genet* **237:** 171-176. https://doi.org/10.1007/BF00282798

Lee I, Michaels SD, Masshardt AS, Amasino RM (1994) **The late-flowering phenotype of *FRIGIDA* and mutations in *LUMINIDEPENDENS* is suppressed in the Landsberg *erecta* strain of *Arabidopsis***. *Plant J* **6:** 903-909. https://doi.org/10.1046/j.1365-313X.1994.6060903.x

Li H, Ye K, Shi Y, Cheng J, Zhang X, Yang S (2017) **BZR1 positively regulates freezing tolerance via CBF-dependent and CBF-independent pathways in *Arabidopsis***. *Mol Plant* **10:** 545-559. https://doi.org/10.1016/j.molp.2017.01.004

Li Z, Jiang D, He Y (2018) **FRIGIDA establishes a local chromosomal environment for *FLOWERING LOCUS C* mRNA production**. *Nat Plants* **4:** 836-846. https://doi.org/10.1038/s41477-018-0250-6

Liu F, Marquardt S, Lister C, Swiezewski S, Dean C (2010) **Targeted 3' processing of antisense transcripts triggers *Arabidopsis FLC* chromatin silencing**. *Science* **327:** 94-97. https://doi.org/10.1126/science.1180278

Liu F, Quesada V, Crevillén P, Bäurle I, Swiezewski S, Dean C (2007) **The *Arabidopsis* RNA-binding protein FCA requires a lysine-specific demethylase 1 homolog to downregulate *FLC***. *Mol Cell* **28:** 398-407. https://doi.org/10.1016/j.molcel.2007.10.018

Liu Q, Kasuga M, Sakuma Y, Abe H, Miura S, Yamaguchi-Shinozaki K, Shinozaki K (1998) **Two transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding**

domain separate two cellular signal transduction pathways in drought- and low-
temperature-responsive gene expression, respectively, in Arabidopsis. *Plant Cell* **10:**
1391-1406. https://doi.org/10.1105/tpc.10.8.1391

Liu Z, Jia Y, Ding Y, Shi Y, Li Z, Guo Y, Gong Z, Yang S (2017) **Plasma membrane CRPK1-**
mediated phosphorylation of 14-3-3 proteins induces their nuclear import to fine-tune
CBF signaling during cold response. *Mol Cell* **66:** 117-128 e115.
https://doi.org/10.1016/j.molcel.2017.02.016

Luo X, Chen T, Zeng X, He D, He Y (2019) **Feedback regulation of *FLC* by *FLOWERING***
LOCUS T* (*FT*) and *FD* through a 5' *FLC* promoter region in *Arabidopsis. *Mol Plant*
12: 285-288. https://doi.org/10.1016/j.molp.2019.01.013

Marquardt S, Raitskin O, Wu Z, Liu F, Sun Q, Dean C (2014) **Functional consequences of**
splicing of the antisense transcript *COOLAIR* on *FLC* transcription. *Mol Cell* **54:** 156-
165. https://doi.org/10.1016/j.molcel.2014.03.026

Maruyama K, Sakuma Y, Kasuga M, Ito Y, Seki M, Goda H, Shimada Y, Yoshida S, Shinozaki K,
Yamaguchi-Shinozaki K (2004) **Identification of cold-inducible downstream genes of**
the *Arabidopsis* DREB1A/CBF3 transcriptional factor using two microarray systems.
Plant J **38:** 982-993. https://doi.org/10.1111/j.1365-313X.2004.02100.x

McKhann HI, Gery C, Bérard A, Lévêque S, Zuther E, Hincha DK, De Mita S, Brunel D, Téoulé
E (2008) **Natural variation in CBF gene sequence, gene expression and freezing**
tolerance in the Versailles core collection of Arabidopsis thaliana. *BMC Plant Biol* **8:**
105. https://doi.org/10.1186/1471-2229-8-105

Medina J, Bargues M, Terol J, Pérez-Alonso M, Salinas J (1999) **The Arabidopsis *CBF* gene**
family is composed of three genes encoding AP2 domain-containing proteins whose
expression Is regulated by low temperature but not by abscisic acid or dehydration.
Plant Physiol **119:** 463-470. https://doi.org/10.1104/pp.119.2.463

Michaels SD, Amasino RM (1999) ***FLOWERING LOCUS C* encodes a novel MADS domain**
protein that acts as a repressor of flowering. *Plant Cell* **11:** 949-956.
https://doi.org/10.1105/tpc.11.5.949

Michaels SD, Amasino RM (2000) **Memories of winter: Vernalization and the competence to**

flower. *Plant Cell Environ* **23:** 1145-1153. https://doi.org/10.1046/j.1365-3040.2000.00643.x

Michaels SD, Amasino RM (2001) **Loss of *FLOWERING LOCUS C* activity eliminates the late-flowering phenotype of *FRIGIDA* and autonomous pathway mutations but not responsiveness to vernalization**. *Plant Cell* **13:** 935-941. https://doi.org/10.1105/tpc.13.4.935

Michaels SD, Himelblau E, Kim SY, Schomburg FM, Amasino RM (2005) **Integration of flowering signals in winter-annual Arabidopsis**. *Plant Physiol* **137:** 149-156. https://doi.org/10.1104/pp.104.052811

Mylne JS, Barrett L, Tessadori F, Mesnage S, Johnson L, Bernatavichute YV, Jacobsen SE, Fransz P, Dean C (2006) **LHP1, the *Arabidopsis* homologue of HETEROCHROMATIN PROTEIN1, is required for epigenetic silencing of *FLC***. *Proc Natl Acad Sci U S A* **103:** 5012-5017. https://doi.org/10.1073/pnas.0507427103

Napp-Zinn K (1955) **Genetische grundlagen des Kältebedürfnisses bei *Arabidopsis thaliana* (L.) Heynh**. *Naturwissenschaften* **42:** 650-650. https://doi.org/10.1007/Bf00622395

Nishio H, Buzas DM, Nagano AJ, Suzuki Y, Sugano S, Ito M, Morinaga S, Kudoh H (2016) **From the laboratory to the field: assaying histone methylation at *FLOWERING LOCUS C* in naturally growing *Arabidopsis halleri***. *Genes Genet Syst* **91:** 15-26. https://doi.org/10.1266/ggs.15-00071

Novillo F, Medina J, Salinas J (2007) ***Arabidopsis* CBF1 and CBF3 have a different function than CBF2 in cold acclimation and define different gene classes in the CBF regulon**. *Proc Natl Acad Sci U S A* **104:** 21002-21007. https://doi.org/10.1073/pnas.0705639105

O'Malley RC, Huang SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M, Gallavotti A, Ecker JR (2016) **Cistrome and epicistrome features shape the regulatory DNA landscape**. *Cell* **165:** 1280-1292. https://doi.org/10.1016/j.cell.2016.04.038

Park J, Lim CJ, Shen M, Park HJ, Cha JY, Iniesto E, Rubio V, Mengiste T, Zhu JK, Bressan RA, Lee SY, Lee BH, Jin JB, Pardo JM, Kim WY, Yun DJ (2018) **Epigenetic switch from repressive to permissive chromatin in response to cold stress**. *Proc Natl Acad Sci U S A* **115:** E5400-E5409. https://doi.org/10.1073/pnas.1721241115

Park S, Gilmour SJ, Grumet R, Thomashow MF (2018) **CBF-dependent and CBF-independent**

regulatory pathways contribute to the differences in freezing tolerance and cold-

regulated gene expression of two Arabidopsis ecotypes locally adapted to sites in

Sweden and Italy. *PLoS One* **13:** e0207723.

https://doi.org/10.1371/journal.pone.0207723

Park S, Lee CM, Doherty CJ, Gilmour SJ, Kim Y, Thomashow MF (2015) **Regulation of the**

Arabidopsis CBF regulon by a complex low-temperature regulatory network. *Plant J*

82: 193-207. https://doi.org/10.1111/tpj.12796

Penfield S (2008) **Temperature perception and signal transduction in plants**. *New Phytol* **179:**

615-628. https://doi.org/10.1111/j.1469-8137.2008.02478.x

Qüesta JI, Song J, Geraldo N, An H, Dean C (2016) *Arabidopsis* **transcriptional repressor VAL1**

triggers Polycomb silencing at *FLC* **during vernalization**. *Science* **353:** 485-488.

https://doi.org/10.1126/science.aaf7354

Seo E, Lee H, Jeon J, Park H, Kim J, Noh YS, Lee I (2009) **Crosstalk between cold response and**

flowering in *Arabidopsis* **is mediated through the flowering-time gene** *SOC1* **and its**

upstream negative regulator *FLC*. *Plant Cell* **21:** 3185-3197.

https://doi.org/10.1105/tpc.108.063883

Seo PJ, Park MJ, Lim MH, Kim SG, Lee M, Baldwin IT, Park CM (2012) **A self-regulatory circuit**

of CIRCADIAN CLOCK-ASSOCIATED1 underlies the circadian clock regulation of

temperature responses in *Arabidopsis*. *Plant Cell* **24:** 2427-2442.

https://doi.org/10.1105/tpc.112.098723

Sheldon CC, Burn JE, Perez PP, Metzger J, Edwards JA, Peacock WJ, Dennis ES (1999) **The** *FLF*

MADS box gene: A repressor of flowering in Arabidopsis regulated by vernalization

and methylation. *Plant Cell* **11:** 445-458. https://doi.org/10.1105/tpc.11.3.445

Sheldon CC, Rouse DT, Finnegan EJ, Peacock WJ, Dennis ES (2000) **The molecular basis of**

vernalization: The central role of *FLOWERING LOCUS C* **(***FLC***)**. *Proc Natl Acad Sci*

U S A **97:** 3753-3758. https://doi.org/10.1073/pnas.97.7.3753

Shi Y, Ding Y, Yang S (2018) **Molecular regulation of CBF signaling in cold acclimation**. *Trends*

Plant Sci **23:** 623-637. https://doi.org/10.1016/j.tplants.2018.04.002

Shindo C, Aranzana MJ, Lister C, Baxter C, Nicholls C, Nordborg M, Dean C (2005) **Role of**
***FRIGIDA* and *FLOWERING LOCUS C* in determining variation in flowering time of**
Arabidopsis. *Plant Physiol* **138:** 1163-1173. https://doi.org/10.1104/pp.105.061309

Shinozaki K, Yamaguchi-Shinozaki K, Seki M (2003) **Regulatory network of gene expression**
in the drought and cold stress responses. *Curr Opin Plant Biol* **6:** 410-417.
https://doi.org/10.1016/s1369-5266(03)00092-x

Shu H, Nakamura M, Siretskiy A, Borghi L, Moraes I, Wildhaber T, Gruissem W, Hennig L (2014)
***Arabidopsis* replacement histone variant H3.3 occupies promoters of regulated genes**.
Genome Biol **15:** R62. https://doi.org/10.1186/gb-2014-15-4-r62

Song Y, Zhang X, Li M, Yang H, Fu D, Lv J, Ding Y, Gong Z, Shi Y, Yang S (2021) **The direct**
targets of CBFs: In cold stress response and beyond. *J Integr Plant Biol* **63:** 1874-1887.
https://doi.org/10.1111/jipb.13161

Stockinger EJ, Gilmour SJ, Thomashow MF (1997) ***Arabidopsis thaliana CBF1* encodes an AP2**
domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-
acting DNA regulatory element that stimulates transcription in response to low
temperature and water deficit. *Proc Natl Acad Sci U S A* **94:** 1035-1040.
https://doi.org/10.1073/pnas.94.3.1035

Sung S, Amasino RM (2004a) **Vernalization and epigenetics: how plants remember winter**.
Curr Opin Plant Biol **7:** 4-10. https://doi.org/10.1016/j.pbi.2003.11.010

Sung S, Amasino RM (2004b) **Vernalization in *Arabidopsis thaliana* is mediated by the PHD**
finger protein VIN3. *Nature* **427:** 159-164. https://doi.org/10.1038/nature02195

Sung S, He Y, Eshoo TW, Tamada Y, Jonson L, Nakahigashi K, Goto K, Jacobsen SE, Amasino
RM (2006) **Epigenetic maintenance of the vernalized state in Arabidopsis thaliana**
requires LIKE HETEROCHROMATIN PROTEIN 1. *Nat Genet* **38:** 706-710.
https://doi.org/10.1038/ng1795

Swiezewski S, Liu F, Magusin A, Dean C (2009) **Cold-induced silencing by long antisense**
transcripts of an *Arabidopsis* Polycomb target. *Nature* **462:** 799-802.
https://doi.org/10.1038/nature08618

Thomashow MF (1999) **PLANT COLD ACCLIMATION: Freezing tolerance genes and**

regulatory mechanisms. *Annu Rev Plant Physiol Plant Mol Biol* **50:** 571-599. https://doi.org/10.1146/annurev.arplant.50.1.571

Thomashow MF (2010) **Molecular basis of plant cold acclimation: Insights gained from studying the CBF cold response pathway**. *Plant Physiol* **154:** 571-577. https://doi.org/10.1104/pp.110.161794

Thomashow MF, Stockinger EJ, Jaglo-Ottosen KR, Gilmour SJ, Zarka DG (1997) **Function and regulation of *Arabidopsis thaliana COR* (<u>co</u>ld-<u>r</u>egulated) genes**. *Acta Physiol Plant* **19:** 497-504. https://doi.org/10.1007/s11738-997-0046-1

Wang ZW, Wu Z, Raitskin O, Sun Q, Dean C (2014) **Antisense-mediated *FLC* transcriptional repression requires the P-TEFb transcription elongation factor**. *Proc Natl Acad Sci U S A* **111:** 7468-7473. https://doi.org/10.1073/pnas.1406635111

Weiser CJ (1970) **Cold resistance and injury in woody plants: Knowledge of hardy plant adaptations to freezing stress may help us to reduce winter damage**. *Science* **169:** 1269-1278. https://doi.org/10.1126/science.169.3952.1269

Went FW (1953) **The effect of temperature on plant growth**. *Annu Rev Plant Phys* **4:** 347-362. https://doi.org/10.1146/annurev.pp.04.060153.002023

Wood CC, Robertson M, Tanner G, Peacock WJ, Dennis ES, Helliwell CA (2006) **The *Arabidopsis thaliana* vernalization response requires a polycomb-like protein complex that also includes VERNALIZATION INSENSITIVE 3**. *Proc Natl Acad Sci U S A* **103:** 14631-14636. https://doi.org/10.1073/pnas.0606385103

Wu Z, Fang X, Zhu D, Dean C (2020) **Autonomous pathway: *FLOWERING LOCUS C* repression through an antisense-mediated chromatin-silencing mechanism**. *Plant Physiol* **182:** 27-37. https://doi.org/10.1104/pp.19.01009

Xu C, Fang X, Lu T, Dean C (2021) **Antagonistic cotranscriptional regulation through ARGONAUTE1 and the THO/TREX complex orchestrates *FLC* transcriptional output**. *Proc Natl Acad Sci U S A* **118**. https://doi.org/10.1073/pnas.2113757118

Xu C, Wu Z, Duan HC, Fang X, Jia G, Dean C (2021) **R-loop resolution promotes co-transcriptional chromatin silencing**. *Nat Commun* **12:** 1790. https://doi.org/10.1038/s41467-021-22083-6

Yang H, Berry S, Olsson TSG, Hartley M, Howard M, Dean C (2017) **Distinct phases of**
Polycomb silencing to hold epigenetic memory of cold in *Arabidopsis*. *Science* **357:**
1142-1145. https://doi.org/10.1126/science.aan1121

Yang H, Howard M, Dean C (2014) **Antagonistic roles for H3K36me3 and H3K27me3 in the**
cold-induced epigenetic switch at *Arabidopsis FLC*. *Curr Biol* **24:** 1793-1797.
https://doi.org/10.1016/j.cub.2014.06.047

Yoo SD, Cho YH, Sheen J (2007) *Arabidopsis* **mesophyll protoplasts: a versatile cell system for**
transient gene expression analysis. *Nat Protoc* **2:** 1565-1572.
https://doi.org/10.1038/nprot.2007.199

Yuan W, Luo X, Li Z, Yang W, Wang Y, Liu R, Du J, He Y (2016) **A** *cis* **cold memory element**
and a *trans* **epigenome reader mediate Polycomb silencing of** *FLC* **by vernalization in**
Arabidopsis. *Nat Genet* **48:** 1527-1534. https://doi.org/10.1038/ng.3712

Zhao C, Zhang Z, Xie S, Si T, Li Y, Zhu JK (2016) **Mutational evidence for the critical role of**
CBF transcription factors in cold acclimation in Arabidopsis. *Plant Physiol* **171:** 2744-
2759. https://doi.org/10.1104/pp.16.00533

Zhao Y, Antoniou-Kourounioti RL, Calder G, Dean C, Howard M (2020) **Temperature-**
dependent growth contributes to long-term cold sensing. *Nature* **583:** 825-829.
https://doi.org/10.1038/s41586-020-2485-4

Zhao Y, Zhu P, Hepworth J, Bloomer R, Antoniou-Kourounioti RL, Doughty J, Heckmann A, Xu
C, Yang H, Dean C (2021) **Natural temperature fluctuations promote** *COOLAIR*
regulation of *FLC*. *Genes Dev* **35:** 888-898. https://doi.org/10.1101/gad.348362.121

Zhu P, Lister C, Dean C (2021) **Cold-induced** *Arabidopsis* **FRIGIDA nuclear condensates for**
FLC **repression**. *Nature* **599:** 657-661. https://doi.org/10.1038/s41586-021-04062-5

Figures

Figure 1. CBF3 directly binds to the CRT/DREs at the 3′-end of *FLC*.

(A) Comparison of sequences around the 3′-end of the *FLC* orthologs from Arabidopsis relatives. The upper graphic presents the gene structure of *AtFLC*. The black boxes, black lines, and white boxes indicate exons, introns, and UTRs, respectively. The blue line presents the region used for sequence comparison among six orthologous genes from five plant species. In the sequence alignment below, the grey line indicates the 3′-UTR of *FLC* orthologs, the blue arrow indicates the

transcriptional start site of *AtCOOLAIR*, and the red lines indicate CRT/DREs. At, *Arabidopsis*

thaliana; Cr, *Capsella rubella*; Al, *Arabidopsis lyrata*; Aa, *Arabis alpina*; Th, *Thellungiella*

halophila.

(B) EMSA using one of the CRT/DREs located at the *COOLAIR* promoter, DRE1, as a probe. In

the upper graphic showing *AtFLC* gene structure, CRT/DRE-like sequences are marked as blue

arrows and labelled as a, b, c, and d for CRT/DRE-like sequences and as 1 and 2 for CRT/DRE

sequences. For the competition assay, these CRT/DRE-like sequences and mutant forms of DRE1

and DRE2 were used as competitors. A 100-fold molar excess of unlabelled competitors was added.

No protein (−) or maltose binding protein (MBP) were used as controls. FP, free probe.

(C) ChIP assay result showing the enrichment of CBF3-myc protein on the *AtFLC* locus. Samples

of NV, 10V, 20V, and 40V plants of *pSuper:CBF3-myc* were collected at zeitgeber time (ZT) 4 in

an SD cycle. The CBF3-chromatin complex was immunoprecipitated (IP) with anti-myc

antibodies (blue bars), and mouse IgG (black bars) was used as a control. Positions of qPCR

amplicons used for ChIP-qPCR analysis are illustrated as P1, P2, and P3 in the upper graphic. The

blue arrows in the graphic denote the position of CRT/DREs and CRT/DRE-like sequences located

on the *FLC* gene. ChIP-qPCR results have been represented as mean ± SEM of the three biological

replicates in the lower panel. Open circles and squares represent each data-point. Relative

enrichments of the IP/5% input were normalized to that of *pTUB2*. The blue shading indicates the

cold period. Asterisks indicate a significant difference between IgG and anti-myc ChIP-qPCR

results at each vernalization time-point (*, $P < 0.05$; **, $P < 0.01$; unpaired Student's *t*-test).

Figure 1—figure supplement 1. CBF proteins directly bind to the sequence at the 3′-end of *FLC*

in DNA affinity purification sequencing (DAP-seq).

Figure 1—figure supplement 2. CBF3 binds to DRE1 and DRE2 on the *COOLAIR* promoter and

DREc on the first exon of *FLC*.

Figure 1—source data 1. Uncropped labelled gel image and the original image file for the EMSA

result.

Figure 1—figure supplement 2—source data 1. Uncropped labelled gel images and the original

image files for the EMSA results.

Figure 2. *COOLAIR* shows a similar expression pattern with CBF regulons.

(A) Expression levels of total *COOLAIR* and *COOLAIR* isoforms in the wild-type (Col-0) and

CBF3-overexpressing plants (*pSuper:CBF3-myc*). The gene structures of *FLC* and *COOLAIR*

variants are illustrated in the upper panel. The thin black arrows indicate the transcription start

sites of *FLC* and *COOLAIR*. The thick black arrows indicate the exons of each *COOLAIR* isoform.

The structures of proximal (I.i) and distal (II.i and II.ii) *COOLAIR* isoforms are shown. The primer

sets used for total, proximal (I.i), and distal (II.i, II.ii) *COOLAIR*s are marked in ***Figure 2—figure***

supplement 1. Relative transcript levels of total *COOLAIR* and *COOLAIR* variants were

normalized to that of *UBC* and have been represented as mean ± SEM of three biological replicates.

Dots and squares represent each data-point. Asterisks indicate a significant difference as compared

to the wild type (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; unpaired Student's t-test).

(B) Expression dynamics of total *COOLAIR* after short-term cold (0, 0.5, 1, 1.5, 2, 3, 6, and 24 h)

treatment. Wild types (*FRI* Col) were subjected to 4°C cold and harvested at each time-point.

Relative transcript levels of total *COOLAIR* to *UBC* were normalized to that of non-cold treated

wild type. The values have been represented as mean ± SEM of three biological replicates. The

blue shading indicates periods under cold treatment. Significant differences have been marked

using different letters (a, b; $P < 0.05$; one-way ANOVA followed by Tukey's post-hoc test).

(C) Transcript levels of total *COOLAIR* and *COOLAIR* isoforms in wild type and *cbfs-1* mutant

before (−Cold) and after (+Cold) a day of 4°C cold treatment. Relative levels of total *COOLAIR*

and *COOLAIR* variants were normalized to that of *PP2A*. Values have been represented as

mean ± SEM of three biological replicates. Dots and squares indicate each data-point. Significant

differences have been marked using different letters (a, b; $P < 0.05$; two-way ANOVA followed by

Tukey's post-hoc test).

1018

1019 **Figure 2—figure supplement 1.** Overexpression of *CBF1*, *2*, or *3* causes an increase in *COOLAIR*

1020 level, but the effects on *COOLAIR* variants are subtly different.

1021

Figure 3. Protein levels of CBFs increase during the vernalization process.

(A) Transcript levels of *CBF1*, *2*, and *3* under short-term cold exposure. Wild-type plants were subjected to 0, 1, 3, 6, 12, and 24 h of 4°C cold treatment. Relative levels of *CBF1*, *2*, and *3* were normalized to that of *UBC*. Values have been represented as mean ± SEM of three biological replicates. The blue shadings indicate cold periods. Significant differences have been marked using different letters (a–c; $P < 0.05$; one-way ANOVA followed by Tukey's post-hoc test).

(B) Transcript levels of *CBF1*, *2*, and *3* after 20V and 40V. Wild-type plants were treated with 4°C

vernalization under an SD cycle and collected at ZT4. Relative levels of *CBF1*, *2*, and *3* were

normalized to that of *UBC*. Values have been represented as mean ± SEM of three biological

replicates. The blue shadings denote periods under cold. Significant differences have been marked

using different letters (a–c; $P < 0.05$; one-way ANOVA followed by Tukey's post-hoc test).

(C) Daily rhythms of *CBF1*, *2*, and *3* transcript levels in NV or 40V plants. Col-0 plants grown

under an SD cycle were collected every 4 h between ZT0 and ZT24. Relative transcript levels of

CBF1, *2*, and *3* were normalized to that of *PP2A*. Values have been represented as mean ± SEM of

three biological replicates. The white and black bars represent light and dark periods, respectively.

Asterisks indicate a significant difference between NV and 40V (***, $P < 0.001$; two-way

ANOVA).

(D) Dynamics of CBF3 protein level under short-term cold exposure. The *pCBF3:CBF3-myc*

transgenic plants were subjected to 0, 1, 3, 6, 12, and 24 h of 4°C cold treatment. CBF3 proteins

were detected using anti-myc antibodies. Rubisco was considered the loading control. Numbers

below each band indicate relative signal intensity compared to 6 h. The mean values of two

biological replicates are presented.

(E) Increase of CBF3 protein level during the vernalization process. The *pCBF3:CBF3-myc*

transgenic plants, subjected to 4°C vernalization, were collected at ZT4 of the indicated time-point.

CBF3 proteins were detected using anti-myc antibodies. Rubisco was considered the loading

control. Numbers below each band indicate relative signal intensity compared to 1V. The mean

values of three biological replicates are presented.

1049

Figure 3—figure supplement 1. Vernalization increases CBF3 protein levels with a minor effect

on the daily rhythm.

Figure 3—source data 1. Uncropped labelled blot images and the original image files for the

immunoblots.

Figure 3—figure supplement 1—source data 1. Uncropped labelled blot images and the original

image files for the immunoblots.

1056

Figure 4. *CBF*s are involved in vernalization-induced *COOLAIR* expression.

(A) Expression dynamics of total *COOLAIR* in the wild-type, *cbfs-1*, and *FLC*_{ΔCOOLAIR-1} plants during vernalization. Relative transcript levels of total *COOLAIR* were normalized to that of *PP2A*. Values have been represented as mean ± SEM of three biological replicates. Asterisks indicate a significant difference compared to the wild-type (***, $P < 0.001$; two-way ANOVA followed by Tukey's post-hoc test).

(B) Transcript level of total *COOLAIR* in wild-type (Col-0) and *CBF3*-overexpressing transgenic plant (*CBF3ox; pSuper:CBF3-myc*) during vernalization. Relative levels of total *COOLAIR* were normalized to that of *PP2A*. Values have been represented as mean ± SEM of three biological replicates. Asterisks indicate a significant difference between the wild-type and *CBF3* overexpressor (***, $P < 0.001$; two-way ANOVA). Blue shading denotes cold periods in (A) and

(B).

(C) Effect of first frost-mimicking treatment (8 h of freezing [< 0°C]) on the level of total

COOLAIR in NV, 10V, and 20V wild type and *cbfs-1*. The upper panel shows a schematic of the

experimental procedure. The non-frost treated (−F) wild type and *cbfs-1* were collected at ZT22

after an 8 h of dark treatment at 22°C (NV) or 4°C (10V and 20V). For the first frost treatment,

wild type and *cbfs-1* mutant were treated with an additional 8 h of −1°C (+F) under dark, and then

the whole seedlings were collected at ZT22 for analysis. All the plants were grown under an SD

cycle. The grey shadings denote dark periods. Total *COOLAIR* levels have been represented as

mean ± SEM of three biological replicates in the lower panel. Dots and squares indicate each data-

point. Relative levels of total *COOLAIR* were normalized to that of *PP2A*. Asterisks indicate a

significant difference (*, $P < 0.05$; ***, $P < 0.001$; two-way ANOVA followed by Tukey's post-

hoc test). ns, not significant ($P \geq 0.05$).

1080

1081   **Figure 4—figure supplement 1.** CBF regulons show strongly reduced expression in *cbfs* mutants

1082   similar to *COOLAIR*.

1083   **Figure 4—figure supplement 2.** Cold-triggered *COOLAIR* expression is impaired in *cbfs* mutants

1084   in Col-0 and Sweden-ecotype background.

1085   **Figure 4—figure supplement 3.** Freezing temperature increases the levels of *CBFs*.

Figure 5. *COOLAIR* promoter region containing CRT/DREs is necessary for the *COOLAIR*

induction by vernalization.

(A) Arabidopsis protoplast transfection assay showing that CBF3 activates *COOLAIR* promoter

with wild-type CRT/DRE (DRE) but fails to activate the one with mutant CRT/DRE (DRE^m). A

schematic of the reporter constructs used for the luciferase assay is presented in the upper panel.

pFLC:LUC contains 1 kb of the promoter, the 5′-UTR, and the first exon of *FLC*. The blue line in

the *pFLC:LUC* graphic indicates the location of DREc. Wild-type and mutant forms of *pCOOLAIR:LUC* include 1 kb of the *COOLAIR* promoter with the 3′-UTR and the last exon of *FLC*. The blue and red lines mark the positions of the wild-type (DRE) and mutant (DRE^{m}) forms of DRE1 and DRE2, respectively. Each reporter construct was co-transfected into Arabidopsis protoplast together with the *35S:CBF3-HA* effector construct. In parallel, *35S-NOS* plasmid was transfected as a control. The result is shown below. Relative luciferase activities were normalized to that of the *35S-NOS* control. Values have been represented as mean $\pm$ SEM of three biological replicates. Dots and squares represent each data-point.

(B) Transcript levels of proximal (I.i) and distal (II.i and II.ii) *COOLAIR* isoforms in wild-type, *FLC$_{\Delta COOLAIR-1}$*, *FLC$_{\Delta COOLAIR-3}$*, and *FLC$_{\Delta COOLAIR-4}$* plants before and after 14 days of vernalization. *FLC$_{\Delta COOLAIR-1}$* and *FLC$_{\Delta COOLAIR-3}$* have a 324- and 685-bp deletion of the *COOLAIR* promoter region, respectively, where DRE1 and 2 are located. *FLC$_{\Delta COOLAIR-4}$* has a deletion in the 301-bp *COOLAIR* promoter region outside of DRE1 and -2 location. The positions of the deleted region are marked in red, orange, and green lines with reversed triangle in the upper graphic. Blue arrows denote CRT/DRE-like sequences. Black and grey bars denote exons, thin lines denote introns, and white bars denote UTRs of *FLC* and its neighboring gene (AT5G10130). Results of RT-PCR analysis are shown below. *UBC* was used as a quantitative control.

(C) Transcript levels of proximal (I.i) *COOLAIR* isoform in wild-type, *cbfs-1*, and *FLC$_{\Delta COOLAIR-1}$* plants during vernalization. The position of the deleted region is marked in red lines with a reversed triangle in the upper graphic. Black bars denote exons, thin lines denote introns, and white bars denote UTRs of *FLC*. The thin black arrow below the gene structure indicates the transcriptional start site of *COOLAIR*. The thick black arrow below denotes exons of the type I.i *COOLAIR* variant.

The position of the amplicon used for the qPCR analysis is marked with blue arrows. The result of

qPCR analysis is presented in the lower panel. Relative transcript levels were normalized to that

of *PP2A*. Values have been represented as mean $\pm$ SEM of three biological replicates. The blue

shading indicates periods under cold treatment. Asterisks indicate a significant difference, as

compared to NV (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; two-way ANOVA followed by Tukey's

post-hoc test).

1121

Figure 5—figure supplement 1. Genomic sequences near the deleted regions in two *COOLAIR*

promoter deletion lines, *FLC$_{\Delta COOLAIR-3}$* and *FLC$_{\Delta COOLAIR-4}$*.

Figure 5—figure supplement 2. Proximal *COOLAIR* level is not significantly increased by *CBF3*

overexpression in the *FLC$_{\Delta COOLAIR}$* mutant lacking CRT/DREs.

Figure 5—source data 1. Uncropped labelled gel images and the original image files for the RT-

PCR results.

1128

Figure 6. Vernalization-triggered epigenetic silencing of *FLC* is not affected by CBFs-mediated *COOLAIR* induction.

(A–C) Enrichments of H3K36me3 (A), H3K4me3 (B), and H3K27me3 (C) on the *FLC* locus in NV or 40V wild-type and *cbfs-1* plants. The whole seedlings were collected at ZT4 under an SD cycle. Modified histones were immunoprecipitated with anti-H3K36me3, anti-H3K4me3, or anti-H3K27me3 antibodies. H3K36me3, H3K4me3, and H3K27me3 enrichments of the IP/5% input

were normalized to those of *ACT7* or *FUS3*. Relative enrichments have been represented as

mean ± SEM of three biological replicates. One-way ANOVA and Tukey's post-hoc test were

performed on the ChIP results obtained using two primer pairs corresponding to the *FLC*

nucleation region (*Yang et al., 2017*). Asterisks indicate a significant difference (*, $P < 0.05$). ns,

not significant ($P \geq 0.05$).

(D) Transcript levels of *FLC* mRNA in wild-type, *cbfs-1* and *FLC$_{\Delta COOLAIR-1}$* plants during and after

vernalization. Blue shading denotes cold periods. Relative levels were normalized to that of *PP2A*.

Values have been represented as mean ± SEM of three biological replicates. ns, not significant

between wild type and mutants, as assessed using two-way ANOVA followed by Tukey's post-hoc

test ($P \geq 0.05$).

(E) Transcript levels of *FLC* mRNA in wild type, *FLC$_{\Delta COOLAIR-1}$*, *FLC$_{\Delta COOLAIR-3}$*, and *FLC$_{\Delta COOLAIR-}$*

$_4$ during and after vernalization. Blue shading denotes cold periods. Relative levels were

normalized to that of *PP2A*, and then normalized to NV of each genotype. Values have been

represented as mean ± SEM of three biological replicates. ns, not significant between wild type

and mutants, as assessed using two-way ANOVA followed by Tukey's post-hoc test ($P \geq 0.05$).

(F) Effect of the first frost-mimicking treatment (8 h freezing) on the *FLC* mRNA level in wild

type and *cbfs-1* after vernalization. NV, 10V, 20V, NV+F, 10V+F, and 20V+F plants were subjected

to the treatments described in ***Figure 4C***. Relative levels were normalized to that of *PP2A*. Values

have been represented as mean ± SEM of three biological replicates. Dots and triangles represent

each data-point. Asterisks indicate a significant difference (*, $P < 0.05$; **, $P < 0.01$; two-way

ANOVA followed by Tukey's post-hoc test). ns, not significant ($P \geq 0.05$).

1156

Figure 7. *CBF*-mediated *COOLAIR* induction under long-term cold is not absolutely necessary for vernalization response.

(A) Photographs of NV and 40V wild type, *cbfs-1*, and *FLC$_{ΔCOOLAIR-1}$*. NV plants were grown at 22°C under LD and 40V plants were grown at 22°C under LD after being treated for 40V under SD. The photos were taken when the NV plants started to bolt and the 40V plants showed the first open flower.

(B) Flowering times of NV and vernalized (10V–40V) wild type and *cbfs-1* plants. Flowering time was measured in terms of the number of rosette leaves produced when bolting. Plants were grown under LD after vernalization treatment. Bars and error bars represent the mean ± standard deviation (SD) of three biological replicates. Each dot and square indicate individual flowering time. Significant differences have been marked using different letters (a–e; two-way ANOVA followed

by Tukey's post-hoc test; $P < 0.05$).

(C) Flowering times of non-vernalized (NV) and vernalized (10V–40V) wild-type and

$FLC_{\Delta COOLAIR}$ plants. Flowering time was measured in terms of the number of rosette leaves

produced when bolting. Plants were grown under LD after vernalization. Bars and error bars

indicate the mean ± SD. Each dot and square represent the individual flowering time. Significant

differences have been indicated using different letters (a–e; two-way ANOVA followed by Tukey's

post-hoc test; $P < 0.05$).

(D) Flowering times of non-vernalized (NV) and vernalized (14V–42V) wild-type, $FLC_{\Delta COOLAIR\text{-}}$

$_1$, $FLC_{\Delta COOLAIR\text{-}3}$, and $FLC_{\Delta COOLAIR\text{-}4}$ plants. Flowering time was measured in terms of the number

of rosette leaves produced when bolting. Plants were grown under LD after vernalization. Bars

and error bars indicate the mean ± SD. Each dot, square, triangle, and polygon represent the

individual flowering time. Significant differences have been indicated using different letters (a–g;

two-way ANOVA followed by Tukey's post-hoc test; $P < 0.05$).

1181

Figure 8. Schematic model describing the mechanism of CBFs-mediated *COOLAIR* expression during the early phase of vernalization.

During the early phase of vernalization, increased CBF upregulates *COOLAIR* expression by binding to CRT/DREs at the 3′-end of *FLC*. In the late phase of vernalization, owing to the silencing of *FLC* chromatin, CBF proteins are released from the CRT/DREs in the *COOLAIR* promoter, which leads to a reduction in *COOLAIR* levels.

1188

Figure 1—figure supplement 1. CBF proteins directly bind to the sequence at the 3′-end of

FLC **in DNA affinity purification sequencing (DAP-seq).**

DAP-seq results showed enrichment of CBF proteins at the *AtFLC* locus obtained from the Plant

Cistrome Database (http://neomorph.salk.edu/dev/pages/shhuang/dap_web/pages/index.php;

O'Malley et al., 2016). The top bars represent the transcribed regions of each *COOLAIR* variant.

The bars below represent the structures of the alternatively spliced forms of the *FLC* transcript.

The green, grey, and red bars indicate the exons, introns, and UTRs of *FLC*, respectively. The

peaks below represent the positions of CBF enrichment. CBF-amp; amplification of DAP-seq

results.

1198

Figure 1—figure supplement 2. CBF3 binds to DRE1 and DRE2 on the *COOLAIR* promoter

and DREc on the first exon of *FLC*.

(A) Schematic showing the positions of six CRT/DRE-like sequences distributed on the *AtFLC*

locus. CRT/DRE sequences have been labelled as DRE1, 2, and CRT/DRE-like sequences have

been labelled as DREa, b, c, and d. The black boxes, black lines, and white boxes represent exons,

introns, and UTRs of *AtFLC*, respectively. The red bases indicate the core sequence (CCGAC) of

CRT/DRE. The blue-colored bases represent bases known to determine the binding affinity of

CBF3 to CRT/DRE (*Maruyama et al., 2004*).

1207   **(B–C)** EMSA using DRE1 (B) or DRE2 (C) as probe. For competition assay, DRE1, DRE2, and

1208   DREc were used as competitors. Molar excess (1-, 10-, and 100-fold) of unlabelled competitors

1209   was added. FP, free probe.

1210

Figure 2—figure supplement 1. Overexpression of *CBF1*, *2*, or *3* causes an increase in *COOLAIR* level, but the effects on *COOLAIR* variants are subtly different.

(A) Positions of qPCR amplicons used for detecting proximal (I.i) and distal (II.i and II.ii) *COOLAIR* variants. The black boxes denote exons, black lines denote introns, and white boxes denote UTRs of *FLC*. The thin black arrow below the gene structure indicates the transcription start site of *COOLAIR*. The thick black arrows below denote exons of type I.i, type II.i, and type II.ii *COOLAIR* variants. The positions of qPCR amplicons used for the qPCR analysis have been marked using blue arrow heads.

(B–D) Transcript levels of *COOLAIR* variants in the wild type (Ws-2) and *CBF1*-, *2*-, and *3*-overexpressing transgenic plants (*35S:CBF1* [*CBF1ox*], *35S:CBF2* [*CBF2ox*], and *35S:CBF3* [*CBF3ox*], respectively; *Seo et al., 2009*). Relative levels of type I.i (B), type II.i (C), and type II.ii

1222   (D) *COOLAIR* were normalized to that of *UBC*. Values have been represented as mean $\pm$ SEM of

1223   three biological replicates. Dots, squares, triangles, and polygons represent each data-point.

1224   Asterisks indicate a significant difference, as compared to the wild type (**, $P < 0.01$; ***,

1225 $P < 0.001$; unpaired Student's *t*-test). ns, not significant ($P \geq 0.05$).

1226

Figure 3—figure supplement 1. Vernalization increases CBF3 protein levels with a minor effect on the daily rhythm.

Daily rhythm of CBF3 protein level in NV or 20V plants. The *pCBF3:CBF3-myc* transgenic plants, subjected to 4°C vernalization, were collected every 4 h between ZT0 and ZT24. CBF3 protein was detected using anti-myc antibodies. Rubisco was used as loading control. The numbers below each band indicate the relative signal intensity compared to 20V ZT4. The mean values of three biological replicates are presented.

1234

Figure 4—figure supplement 1. CBF regulons show strongly reduced expression in *cbfs* mutants similar to *COOLAIR*.

(A–B) Expression dynamics of *COR15A* (A) and *RD29A* (B) in the wild type, *cbfs-1*, and *FLC$_{\Delta COOLAIR-1}$* plants during vernalization. Relative transcript levels of *COR15A* and *RD29A* were normalized to that of *PP2A*. Values have been represented as mean ± SEM of three biological replicates. Asterisks indicate a significant difference compared to the wild type (***, $P < 0.001$; two-way ANOVA followed by Tukey's post-hoc test). ns, not significant ($P \geq 0.05$). Blue shading denotes cold periods in (A) and (B).

1243

Figure 4—figure supplement 2. Cold-triggered *COOLAIR* expression is impaired in *cbfs*

mutants in Col-0 and Sweden-ecotype background.

(A–D) Transcript levels of total (A), proximal (B), distal (C) *COOLAIR* and *COR15A* (D) in the

wild type (Col-0) and *cbfs* mutant upon 0, 3, 24 h of cold exposure. The data were obtained from

the analysis using RNA-seq data reported (*Zhao et al., 2016*). The mean values of transcript per

million (TPM) of each gene obtained from three experimental replicates are shown. Error bars

denote SEM.

1251 **(E–H)** Transcript levels of total (E), proximal (F), distal (G) *COOLAIR* and *FLC* (H) in NV, 1V,

1252 and 14V wild type (SW) and *cbfs* mutant. The data were obtained from the analysis using RNA-

1253 seq data reported (*S. Park et al., 2018*). SW, an Arabidopsis ecotype collected from Sweden. The

1254 mean values of the TPM of each gene obtained from two experimental replicates are shown. Open

1255 circles and squares represent each data-point.

1256

Figure 4—figure supplement 3. Freezing temperature increases the levels of *CBFs*.

(A–C) Effect of the first frost-mimicking treatment (8 h freezing) on the levels of *CBF1* (A), *2* (B), and *3* (C) in wild type after vernalization. NV, 10V, 20V, NV+F, 10V+F, and 20V+F plants were subjected to the treatments described in ***Figure 4C***. Relative levels were normalized to that of *PP2A*. Values have been represented as mean ± SEM of three biological replicates. Dots represent each data-point. Asterisks indicate a significant difference (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; unpaired Student's *t*-test).

1264

1265 **Figure 5—figure supplement 1. Genomic sequences near the deleted regions in two**

1266 ***COOLAIR* promoter deletion lines, *FLC$_{\Delta COOLAIR-3}$* and *FLC$_{\Delta COOLAIR-4}$*.**

1267 **(A–B)** Genomic sequences around the deleted regions in newly generated *COOLAIR* promoter

1268 deletion lines, *FLC$_{\Delta COOLAIR-3}$* (A) and *FLC$_{\Delta COOLAIR-4}$* (B). Deleted or replaced sequences are marked

1269 in red and cyan, respectively.

1270

Figure 5—figure supplement 2. Proximal *COOLAIR* level is not significantly increased by

CBF3 overexpression in the *FLC_{ΔCOOLAIR}* mutant lacking CRT/DREs.

(A–B) Expression levels of proximal *COOLAIR* variant I.i (A) and *CBF*3 (B) in non-vernalized

wild type (Col-0), *pSuper:CBF3-myc* (*CBF3ox*), *FLC_{ΔCOOLAIR-1}*, and *pSuper:CBF3-myc*

FLC_{ΔCOOLAIR-1} (*CBF3ox FLC_{ΔCOOLAIR-1}*). The whole seedlings were collected at 16 days after

germination under an SD cycle. Relative levels were normalized to that of *PP2A*. Values have been

represented as mean ± SD of four technical replicates from two biological replicates. Open circles

and squares represent each data-point. Significant differences have been marked using different

letters (a–c; $P < 0.05$; one-way ANOVA followed by Tukey's post-hoc test).

71