Vernalization-triggered expression of the antisense transcript *COOLAIR* is mediated by *CBF* genes

Myeongjune Jeon[1†], Goowon Jeong[1†], Youbong Hyun[1,2], and Ilha Lee[1,2*]

[1]School of Biological Sciences, Seoul National University, Seoul 08826, Korea

[2]Research Center for Plant Plasticity, Seoul National University, Seoul 08826, Korea

[*]For correspondence: ilhalee@snu.ac.kr

[†]These authors contributed equally to this work

Abstract

Many plants undergo vernalization, a long-term winter-triggered acceleration of flowering, to align their flowering time with spring. In *Arabidopsis thaliana*, this is achieved by silencing a floral repressor, *FLOWERING LOCUS C* (*FLC*). *COOLAIR*, an antisense noncoding RNA expressed at the *FLC* locus, is induced during the early phase of vernalization, preceding *FLC* suppression. However, the mechanism by which long-term cold induces *COOLAIR* is not well understood. Here, we showed that C-repeat (CRT)/dehydration-responsive elements (DREs) at the 3′-end of *FLC* and CRT/DRE-binding factors (CBFs) are required for vernalization-induced *COOLAIR* activation. The CBFs bind to CRT/DREs at the 3′-end of *FLC*, both *in vitro* and *in vivo*, and the CBFs levels increased gradually during vernalization. Additionally, vernalization-induced *COOLAIR* expression was highly suppressed in the *cbfs* mutant, in which all *CBF*s were knocked-out. Contrastingly, *CBF*-overexpressing plants showed *COOLAIR* upregulation, even at warm temperatures. We propose that *COOLAIR* is induced by CBFs in the early phase of vernalization but is downregulated as CBFs are evicted from closed *FLC* chromatin during the late vernalization phase. We also demonstrated that *cbfs* and *FLC*$_{\Delta COOLAIR}$ mutants have a normal vernalization response, although they show defects in vernalization-induced *COOLAIR* activation, indicating that *COOLAIR* is not necessary for this process.

Introduction

Appropriate timing of flowering ensures plant reproductive success, which in turn provides evolutionary advantages. As sessile organisms, plants have evolved mechanisms by which reproductive growth is coordinated with seasonal cues. In many flowering plants, exposure to long-term winter cold is required for transition to the reproductive phase (*Chouard, 1960*). Vernalization, an acquisition of flowering competence in response to prolonged winter cold, allows plants to synchronize the timing of flowering with spring (*Chouard, 1960; Michaels and Amasino, 2000*).

Vernalization in *Arabidopsis thaliana* is mainly achieved by silencing a floral repressor gene, *FLOWERING LOCUS C* (*FLC*) (*Michaels and Amasino, 1999; Sheldon et al., 1999; Sheldon et al., 2000; Michaels and Amasino, 2001*). *FLC* encodes a MADS-box protein that directly associates with the promoter regions of floral activator genes, *SUPPRESSOR OF CO1* (*SOC1*) and *FLOWERING LOCUS T* (*FT*), which encodes florigen, and represses their expression (*Lee et al., 2000; Michaels et al., 2005; Helliwell et al., 2006*). *FLC* is highly expressed in winter annuals of Arabidopsis, such as San Feliu-2 and Lov-1, because of the strong transcriptional activation by the FRIGIDA (FRI) supercomplex which recruits general transcription factors, as well as several chromatin modifiers (*Michaels and Amasino, 1999; Sheldon et al., 1999; Johanson et al., 2000; Choi et al., 2011; Li et al., 2018*). Therefore, winter annuals show a late-flowering phenotype, but their flowering is dramatically accelerated by vernalization-induced *FLC* silencing (*Napp-Zinn, 1955; Lee et al., 1993; Clarke and Dean, 1994; Koornneef et al., 1994; Lee et al., 1994*). In contrast to winter annuals, rapid-cycling accessions, such as Columbia (Col) and Landsberg *erecta* (L*er*), exhibit early flowering due to mutations in *FRI* and/or *FLC* (*Johanson et al., 2000; Gazzani et al., 2003; Michaels et al., 2003*). However, the autonomous pathway (AP) mutants in Col and L*er* background also show late flowering with an increased level of *FLC*; vernalization causes the epigenetic suppression of *FLC*, thus accelerating their flowering to a degree similar to that seen in winter annuals (*Koornneef et al., 1998; Michaels and Amasino, 2001*).

In Arabidopsis winter annuals, flowering is prevented by strong expression of *FLC* during fall. Consistently, *FLC* chromatin is highly enriched with active histone marks such as histone

H3 acetylation and trimethylation of Lys4 or Lys36 at H3 (H3K4me3/H3K36me3) (*Bastow et al., 2004; Yang et al., 2014*). During winter, *FLC* chromatin is gradually deacetylated, and the H3K4me3 and H3K36me3 on *FLC* are removed concomitantly (*Bastow et al., 2004; Yang et al., 2014*). Additionally, the repressive histone marks, H3 Lys9 dimethylation (H3K9me2) and H3 Lys27 trimethylation (H3K27me3), are accumulated by the polycomb repressive complex 2 (PRC2) in the nucleation region around the first exon and intron of *FLC* (*Sung and Amasino, 2004; Wood et al., 2006; De Lucia et al., 2008; Angel et al., 2011*). Subsequently in the following spring, H3K27me3 is spread to the entire *FLC* chromatin region by LIKE HETEROCHROMATIN PROTEIN 1, which ensures stable *FLC* suppression and renders plants competent to flower (*Mylne et al., 2006; Yang et al., 2017*).

Several long-term cold-induced factors have been shown to play a crucial role in the epigenetic silencing of *FLC*. *VERNALIZATION INSENSITIVE 3* (*VIN3*) family genes, which are upregulated by prolonged cold, encode PHD (Plant Homeodomain)-finger domain proteins which recognize H3K9me2 enriched in the *FLC* chromatin during the vernalization process (*Sung and Amasino, 2004; Kim and Sung, 2013*, *2017a*). These proteins mediate the recruitment of PRC2 and subsequent deposition of H3K27me3 at the nucleation region of *FLC* (*Sung and Amasino, 2004; De Lucia et al., 2008; Kim and Sung, 2013*). In addition, vernalization-induced long noncoding RNAs (lncRNAs) are involved in such histone modifications. *COLDAIR* and *COLDWRAP*, the two lncRNAs transcribed from the first intron and promoter region of *FLC*, respectively, are necessary for H3K27me3 deposition in response to long-term cold (*Heo and Sung, 2011; Kim and Sung, 2017b*). Additionally, *COLDAIR* and *COLDWRAP* are thought to affect the formation of the intragenic chromatin loop at *FLC*, which may be part of the *FLC* silencing mechanism (*Kim and Sung, 2017b*).

Unlike *COLDAIR* and *COLDWRAP*, which are transcribed in the sense direction of *FLC*, *COOLAIR* is an antisense transcript expressed from the 3′-end of *FLC* (*Swiezewski et al., 2009*). The gradual accumulation of *COOLAIR* reaches a maximum within a few weeks of winter cold, whereas the other lncRNAs, *COLDAIR* and *COLDWRAP*, show peaks at a later phase of vernalization (*Csorba et al., 2014; Kim and Sung, 2017b*). Since the *COOLAIR* promoter is sufficient to confer cold-induced silencing of a reporter gene when added to the 3′-region of

the gene (*Swiezewski et al., 2009*), extensive efforts have been made to elucidate how

COOLAIR antagonizes *FLC* expression. One possibility is that *COOLAIR* facilitates changes

in the chromatin environment of *FLC*. *COOLAIR* has been reported to be involved in removing

active histone marks, H3K4 and H3K36 methylations from *FLC* chromatin, thus leading to

FLC suppression (*Helliwell et al., 2011; Csorba et al., 2014; Berry and Dean, 2015*). Co-

transcriptional events governed by *COOLAIR*-targeted RNA-processing factors, including

some factors involved in the autonomous pathway for flowering, seem to be critical for

switching the *FLC* chromatin state (*Wu et al., 2020*). Furthermore, *COOLAIR* is also likely

necessary for H3K27me3 deposition via interaction with FLOWERING CONTROL LOCUS

A (FCA) protein (*Tian et al., 2019*).

Compared to the functional mechanism of *COOLAIR* in *FLC* silencing, how long-term

cold triggers *COOLAIR* induction is less well-understood. Previous studies have inferred that

the regulatory region conferring long-term cold response of *COOLAIR* is well-conserved

among relatives of Arabidopsis (*Swiezewski et al., 2009; Castaings et al., 2014; Csorba et al.,*

2014; Berry and Dean, 2015). However, little is known about their upstream regulators.

Although a recent report has shown that an NAC domain-containing protein, NTM1-LIKE 8

(NTL8), can bind to the *COOLAIR* promoter and activate its expression, it has not been

thoroughly addressed whether NTL8 is necessary for vernalization-induced *COOLAIR*

activation (*Zhao et al., 2021*). In the present study, we identified that CRT/DREs at the 3′-end

of *FLC* are required for the long-term cold response of *COOLAIR*. Additionally, we showed

that CRT/DRE-binding factors (CBFs), which accumulate through the long-term winter cold,

act as upstream regulators of *COOLAIR* during vernalization.

Results

A CRT/DRE-binding factor, CBF3, directly binds to CRT/DREs at the 3′-end of *FLC*

Since it has already been shown that the proximal promoter region of *COOLAIR* is highly conserved among the *FLC* orthologs from *A. thaliana* relatives (*Castaings et al., 2014*), we assumed that the *cis*-element conferring long-term cold response would exist within that region. A comparison of the region near the transcriptional start site of *COOLAIR* revealed that two CRT/DREs are well-conserved among six *FLC* orthologs from five related species in the Brassicaceae family (*Figure 1A*). CRT/DRE, whose core sequence is CCGAC, is a regulatory element that imparts cold- or drought-responsive gene expression (*Baker et al., 1994*). Three *CBF* family genes, *CBF1*, *2*, and *3*, have been known to encode AP2 (APETALA2) domain-containing proteins that can bind to CRT/DRE, and are usually present in the promoters of cold- and drought-regulated genes (*Stockinger et al., 1997; Medina et al., 1999*). Consistent with this, the Arabidopsis cistrome database suggests that the CBFs bind to the 3′-end sequence of *FLC* containing CRT/DREs (*Figure 1—figure supplement 1; O'Malley et al., 2016*).

To confirm this binding, we performed electrophoretic mobility shift assay (EMSA) using probes harboring the CRT/DREs (DRE1 or -2) from the *COOLAIR* promoter. The mobility of these two Cy5-labelled probes was retarded by maltose-binding protein (MBP)-fused CBF3 (*Figure 1B, Figure 1-figure supplement 2*). The band shift was competed out by adding an excess amount of non-labelled DRE1 or 2 oligonucleotides (*Figure 1B, Figure 1—figure supplement 2*). By contrast, the competitors containing the mutant forms of DRE1 or 2 (DRE1^m or 2^m respectively) failed to compete (*Figure 1B*). We also tested the binding of other CRT/DRE-like sequences distributed in the *FLC* locus. Two of these (DREa and b) were present in the *FLC* promoter, while the other two, DREc and d, were present in the first and last exons, respectively (*Figure 1—figure supplement 2*). Among these, only DREc competed with the band shift caused by the CBF3-DRE1 interaction (*Figure 1B*).

Subsequently, a chromatin immunoprecipitation (ChIP) assay using a *CBF3*-overexpressing transgenic plant, *pSuper::CBF3-myc* (*Liu et al., 2017*), was performed to determine whether CBF3 is associated with the region of *FLC* containing CRT/DREs *in vivo*. Consistent with the *in vitro* results, the CBF3-myc protein was enriched at the 3′-end of *FLC*

as well as the first exon of *FLC*, where DREc was located (*Figure 1C*). Without vernalization

(NV), CBF3-myc was highly enriched at both the 5′- and 3′-ends of *FLC* (P1 and P3), indicating

that CBF3 protein can activate *FLC* in warm conditions, as previously reported (*Seo et al.,*

2009). However, the enrichment on the P1 region rapidly disappeared during the vernalization

period. By contrast, the enrichment on P3 was maintained until 10 days of vernalization (10V)

and was subsequently reduced (*Figure 1C*). This is coincident with the expression pattern of

COOLAIR, which declines in the late phase of vernalization (*Csorba et al., 2014*).

COOLAIR is one of the CBF regulons in Arabidopsis

CBF genes are rapidly and transiently induced upon exposure to cold (*Medina et al., 1999*).

CBF regulons, which are CBF-targeted genes, are correspondingly up- or down-regulated after

a few hours-long cold interval (*Gilmour et al., 1998; Liu et al., 1998; Fowler and Thomashow,*

2002). As *COOLAIR* bears CBF3-binding sites in its promoter (*Figure 1*), it was expected that

COOLAIR would show CBF regulon-like expression.

The expression of genes assigned to the CBF regulon is up- or downregulated by *CBF*

overexpression under warm condition (*Park et al., 2015*). Likewise, we found that the total

transcript level of *COOLAIR* increased in *CBF3*-overexpressing transgenic plants

(*pSuper::CBF3-myc*) grown at room temperature (22°C), as compared to that in the wild-type

(Col-0) (*Figure 2A*). Such upregulation was mainly due to type I or type IIi *COOLAIR* isoforms.

As the levels of both proximal (type I) and distal (type II) isoforms of *COOLAIR* were higher

in the *CBF3* overexpressor than in the wild-type, it is likely that *COOLAIR* transcription,

instead of the 3′-processing event, is affected by CBF3 (*Figure 2A; Liu et al., 2010; Marquardt*

et al., 2014). It has been reported that the targets of CBF1 or -2 are not entirely identical to

those of CBF3, although *CBF* genes show high sequence similarity (*Novillo et al., 2007*). Thus,

we compared the expression levels of all *COOLAIR* isoforms in transgenic plants

overexpressing *CBF1, -2,* and *-3* (*Seo et al., 2009*). Overall, all *CBF*-overexpressing plants

showed increased levels of *COOLAIR* (*Figure 2—figure supplement 1*). However, each *CBF*-

overexpressing plant showed a subtly different effect on *COOLAIR* expression. The *CBF2*

overexpressor showed the highest level of the proximal *COOLAIR* isoform, whereas the *CBF3*

overexpressor showed the highest level of distal *COOLAIR*. By contrast, the *CBF1*-overexpressing plants showed a slightly smaller increase of proximal *COOLAIR* levels and a negligible increase in distal *COOLAIR*, as compared to those in the wild-type, Ws-2. Such results imply that the three CBFs regulate *COOLAIR* transcription subtly differently.

Another characteristic of CBF regulons is that they are induced by a day of cold exposure (*Park et al., 2015*). Thus, we checked if *COOLAIR* is also induced by short-term cold. Wild-type (*FRI* Col) plants were treated with 0, 0.5, 1, 1.5, 2, 3, 6, and 24 h of cold (4°C), then the levels of total *COOLAIR* were measured. The results showed that cold treatment less than 3 h failed to induce *COOLAIR* expression, although the *CBF*s expression reached a peak at 3 h of cold treatment (*Figures 2B and 3A; Gilmour et al., 1998*). Following 6 h of cold treatment, *COOLAIR* was conspicuously induced (*Figure 2B*). Thus, short-term cold induced *COOLAIR* expression similar to those of other CBF-targeted genes, *i.e.* the expression is highly increased after *CBF* levels reach a peak upon cold exposure (*Fowler and Thomashow, 2002*). To confirm whether *CBF*s are responsible for the rapid cold response of *COOLAIR*, we checked *COOLAIR* induction in the wild-type and *cbfs-1* mutant, in which all three *CBF*s are knocked-out (*Jia et al., 2016*), after a day of cold treatment (*Figure 2C*). All isoforms, as well as total *COOLAIR*, failed to be induced by short-term cold exposure in the *cbfs* mutant, although a drastic increase was observed in the wild-type. Thus, the results strongly suggested that *CBF*s are required for *COOLAIR* induction, even under a short period of cold exposure.

CBFs accumulate during vernalization

We subsequently questioned whether *CBF*s are also responsible for the induction of *COOLAIR* during vernalization. *COOLAIR* is gradually upregulated as the cold period persists and peaks after 2-3 weeks of vernalization (*Csorba et al., 2014*). However, most studies on *CBF* expression have been performed within a few days, since the function of *CBF*s has been analyzed only in the context of short-term cold (*Gilmour et al., 1998; Liu et al., 1998; Medina et al., 1999; McKhann et al., 2008*). Therefore, we investigated the expression patterns of *CBF*s before and after long-term cold exposure, to find a correlation between the expression of *CBF*s and *COOLAIR* during vernalization. As previously shown, the levels of all three *CBF*s reached

their peaks within 3 h of cold exposure, thereafter decreased rapidly (*Figure 3A*). However, the expression of *CBF*s increased again as the cold period was prolonged (>20V), suggesting that both short-term and vernalizing cold treatments upregulated *CBF* expression (*Figure 3B*).

Since *CBF*s are known to be regulated by circadian rhythms (*Dong et al., 2011*), we also investigated whether the circadian rhythm affects vernalization-induced *CBF* expression. For this, we collected wild-type (Col-0) plants every 4 h under a short-day cycle (SD, 8 h light/16 h dark), in both the NV and 40V conditions. As shown in *Figure 3C*, although the rhythmic pattern of each *CBF* was variable, the overall transcript levels of all three *CBF*s were much higher at 40V than at NV.

To verify whether the increased transcription of *CBF3* led to protein accumulation, we also measured the level of CBF3 at each time-point of vernalization using the *pCBF3::CBF3-myc* plant (*Jia et al., 2016*). In accordance with the rapid and transient induction of *CBF3* under short-term cold (*Medina et al., 1999*), CBF3 was highly expressed after a day of cold exposure (1V), but decreased at 10V (*Figure 3D*). However, as the period of cold exposure was prolonged, the CBF3 level was re-increased such that the level at 40V was similar to that at 1V. The increased protein level after 40V rapidly decreased if the plants were transferred to room temperature for 4 d (40VT4). Such results suggested that CBFs are responsible for the progressive upregulation of *COOLAIR*, at least during the early phase of vernalization (*Csorba et al., 2014*).

CBFs are involved in vernalization-induced COOLAIR expression

To clarify whether *CBF*s are responsible for long-term cold-mediated *COOLAIR* induction, we quantified *COOLAIR* levels in the wild-type and *cbfs* mutant during the vernalization process. Although the level of total *COOLAIR* gradually increased until 20V, and then decreased afterward in the wild-type as previously reported (*Csorba et al., 2014*), the level in *cbfs* mutant was severely reduced across the entire vernalization period (*Figure 4A*). It suggests that *CBF*s are required for full induction of *COOLAIR* in the early phase of vernalization.

Consistent with the suppressed *COOLAIR* expression in *cbfs* mutants, the *CBF3*

overexpressor, *pSuper::CBF3-myc,* showed a much higher *COOLAIR* level than the wild-type (Col-0), throughout the vernalization period (*Figure 4B*). It is noteworthy that *COOLAIR* expression in *pSuper::CBF3-myc* was further upregulated by cold, especially during the early vernalization phase (10V) (*Figure 4B*). This result implies that both transcriptional and post-transcriptional regulation of CBFs are involved in the long-term cold response of *COOLAIR*.

A recent study has shown that the first seasonal frost (<0°C) during the winter season strongly induces *COOLAIR* expression (*Zhao et al., 2021*). Therefore, we tested whether *CBF*s are also required for the strong induction of *COOLAIR* triggered by freezing temperatures. Wild-type and *cbfs* mutants were subjected to 20V, with or without an additional 8 h of -1°C freezing treatment (8hF). In both the genotypes, the *COOLAIR* levels were increased upon 8hF treatment, but *cbfs* mutants showed a much smaller increase compared to the wild-type (*Figure 4C*). Thus, *CBFs* seem to be responsible for both the gradual increase of *COOLAIR* during vernalization and the strong *COOLAIR* induction triggered by freezing temperatures.

CRT/DREs at the 3′-end of *FLC* are necessary for CBFs-mediated induction of *COOLAIR* during vernalization.

Since CBF3 was able to bind to CRT/DREs in the first exon (DREc) and the 3′-end of *FLC* (DRE1 and 2) (*Figure 1B-C*), we questioned whether CRT/DRE is responsible for the CBF-mediated long-term cold response of *COOLAIR*. To address this, we performed an *A. thaliana* protoplast transfection assay using the *35S::CBF3-HA* effector construct and *pCOOLAIR_{DRE}::LUC* reporter construct, in which the 1 kb wild-type *COOLAIR* promoter was fused to the luciferase reporter gene. The luciferase activity assay suggested that the CBF3-HA protein activates transcription of the *COOLAIR* promoter (*Figure 5A*). By contrast, CBF3-HA failed to increase luciferase activity when the mutant version of the *COOLAIR* promoter with mutant DRE1 and 2 (*pCOOLAIR_{DRE}^{m}::LUC*) was used as a reporter (*Figure 5A*). Additionally, co-transfection of the *35S::CBF3-HA* effector and *pFLC::LUC* reporter construct, which harbors the 1 kb sequence including the promoter and first exon of *FLC*, did not show increased luciferase activity (*Figure 5A*). These results strongly indicated that CBF3 can activate *COOLAIR* transcription through DRE1 and 2 located in the promoter, but not via the first exon

of *FLC*.

To further determine whether the 3′-end region of *FLC* is required for the vernalization

response of *COOLAIR*, we measured *COOLAIR* expression in the wild-type and *FLC~ΔCOOLAIR~*

mutant, which lacks a 324 bp portion of the *COOLAIR* promoter region (*Luo et al., 2019*),

before and after vernalization. Although the common *COOLAIR* transcriptional start site was

eliminated in *FLC~ΔCOOLAIR~*, the proximal *COOLAIR* isoform (type I) was still detected,

probably due to alternative transcription start sites (*Figure 5B; Zhao et al., 2021*). As expected,

the expression of the proximal *COOLAIR* variant in the *FLC~ΔCOOLAIR~* mutant was not induced

at 20V, while the wild-type showed increased type I *COOLAIR* level at 20V (*Figure 5B*). Thus,

the *COOLAIR* promoter region containing two CRT/DREs is necessary for the long-term cold

response of *COOLAIR*.

CBFs-mediated *COOLAIR* induction by vernalization is not necessary for *FLC* silencing

Lastly, we investigated the role of *COOLAIR* induced by CBFs in the vernalization-triggered

FLC silencing process. *COOLAIR* is reported to facilitate the removal of H3K4me3 and

H3K36me3 from *FLC* chromatin during vernalization, whereas the other lncRNAs, *COLDAIR*

and *COLDWRAP*, are required for H3K27me3 deposition (*Csorba et al., 2014; Berry and Dean,*

2015; Kim and Sung, 2017b). Therefore, we compared the enrichment of H3K4me3,

H3K36me3, and H3K27me3 on *FLC* chromatin in non-vernalized or vernalized wild-type and

cbfs mutants. As reported (*Bastow et al., 2004; Sung and Amasino, 2004; Yang et al., 2014*),

there was a reduction in the enrichment of H3K4me3 and H3K36me3, whereas that of

H3K27me3 increased considerably in the nucleation region of *FLC* by 40V in the wild-type

(*Figure 6A-D*). Interestingly, although *COOLAIR* levels were significantly reduced in the *cbfs*

mutant (*Figure 4A*), vernalization-caused reductions in H3K4me3 and H3K36me3 levels and

increase in H3K27me3 levels were comparable to those in the wild-type (*Figure 6A-D*). This

result indicates that CBF-mediated *COOLAIR* induction is not necessary for vernalization-

induced epigenetic changes in *FLC* chromatin.

Consistent with the epigenetic changes in *FLC* chromatin in the *cbfs* mutant, the *FLC* level

was gradually suppressed during the vernalization process, and the suppression was maintained

after 40VT4 (*Figure 7A*), similar to that seen in the wild-type. Accordingly, vernalization-

triggered acceleration of flowering was comparable in the wild-type and *cbfs* (*Figure 7B-C*),

indicating that *cbfs* show a normal vernalization response. It has recently been proposed that

the strong induction of *COOLAIR* by the first seasonal frost is correlated with stronger *FLC*

repression (*Zhao et al., 2021*). However, hyper-repression of *FLC* upon exposure to freezing

temperature for 8 h after 20V (20V+8hF) was not observed under our growth conditions, in

both the wild-type and *cbfs* (*Figure 7D*), although both genotypes showed stronger *COOLAIR*

induction upon 8hF treatment (*Figure 4C*).

Since our results suggested that *COOLAIR* induction is not required for *FLC* silencing

achieved by means of vernalization, we checked if the loss-of-function mutant, $FLC_{\Delta COOLAIR}$,

displays any defect in the vernalization response. The $FLC_{\Delta COOLAIR}$ mutant produced virtually

undetectable levels of total *COOLAIR* even after 20V (*Figure 5B*). Additionally, although the

proximal *COOLAIR* variant was barely detected in $FLC_{\Delta COOLAIR}$ as previously reported (*Zhao*

et al., 2021), the level of the proximal form did not increase by 20V (*Figure 5B*). Therefore,

the results strongly suggested that $FLC_{\Delta COOLAIR}$ fails to show vernalization-induced *COOLAIR*

expression. We then compared *FLC* levels between the wild-type and $FLC_{\Delta COOLAIR}$ after

vernalization (*Figure 7E*). Similar to the wild-type, the *FLC* level gradually decreased and was

silenced by long-term cold in $FLC_{\Delta COOLAIR}$. Consistently, the flowering time of $FLC_{\Delta COOLAIR}$

was normally accelerated by vernalization (*Figure 7F*). Taken together, it is likely that CBFs,

upregulated by long-term cold, induce *COOLAIR* expression (*Figure 8*). However, such

induced *COOLAIR* does not seem to be absolutely necessary for *FLC* silencing during

vernalization.

Discussion

Plants recognize and remember a period of cold during the winter season to determine the proper timing of flowering. Several long-term cold-induced factors are known to facilitate such "memory of winter", which establishes flowering competence (*Michaels and Amasino, 2000; Sung and Amasino, 2004; Heo and Sung, 2011; Csorba et al., 2014; Kim and Sung, 2017b*). However, the mechanism by which long-term cold activates such regulators is not well understood. In the present study, we showed that CBF proteins act as transcriptional activators of *COOLAIR*, a long-term cold-induced lncRNA, during vernalization. CBF3 directly binds to the CRT/DRE elements located in the *COOLAIR* promoter region, which is evolutionarily conserved among *FLC* orthologs from Arabis family plants. CBFs are required for the full induction of *COOLAIR* during vernalization (*Figures 2C and 4A*), but their overexpression upregulates *COOLAIR* even at warm temperatures (*Figures 2A and 4B*). Consistently, our results show a gradual accumulation of CBFs during vernalization (*Figure 3*). Thus we propose that the increase in CBF levels induces *COOLAIR* expression during vernalization (*Figure 8*).

However, CBF-mediated *COOLAIR* induction during vernalization cannot be fully explained by increased CBF levels, as the *CBF3* overexpressor still shows *COOLAIR* induction by vernalization (*Figure 4B*). This suggests the involvement of additional factors that regulate CBF function. Previously, it has been reported that low temperatures cause functional activation of CBFs through degradation of HD2C and BTF3s, which prevent CBFs from activating downstream targets at warm temperatures (*Ding et al., 2018; Park et al., 2018*). Moreover, cold triggers the monomerization of CBF proteins, thereby activating their functions (*Lee et al., 2021*). Similarly, such an additional regulatory mechanism of CBFs, which fine-tunes vernalization-induced *COOLAIR* activation, may be enhanced by long-term winter cold. Additionally, *cbfs* mutant showed weak *COOLAIR* induction at 20V (*Figure 4A*). Thus, there are likely other thermosensors that facilitate *COOLAIR* induction by vernalization. Recently, it has been suggested that NTL8, an NAC domain-containing transcription factor, acts as an upstream regulator of *COOLAIR* (*Zhao et al., 2021*). NTL8 also binds to the *COOLAIR* promoter region, which is not far from the CBF-binding sites (*O'Malley et al., 2016; Zhao et al., 2021*). Furthermore, *COOLAIR* has been shown to be highly expressed in both dominant

mutant alleles, *ntl8-D*s, and *NTL8*-overexpressing transgenic plants at warm temperatures.

Thus, it has been proposed that slow accumulation of NTL8 protein, due to reduced dilution

resulting from slower growth at low temperatures, causes *COOLAIR* induction during

vernalization. It would be worthwhile to elucidate whether CBFs and NTL8 synergistically

activate *COOLAIR* transcription during vernalization. The reason as to why vernalization-

triggered *COOLAIR* induction requires more than one thermosensor is not apparent.

Presumably, multiple thermosensory modules ensure precise and robust *COOLAIR* expression

during the winter season when the temperature fluctuates.

The increasing trend of *COOLAIR* levels under long-term cold is not maintained across

the entire vernalization period (*Swiezewski et al., 2009; Csorba et al., 2014*). After reaching a

maximum at the early phase of vernalization, the *COOLAIR* level gradually reduced, although

the accumulation of CBFs continued as vernalization proceeded (*Figures 3 and 4A*). This is

likely due to the falling off of CBF proteins from the *COOLAIR* promoter during the late phase

of vernalization (*Figure 8*). The molecular mechanisms underlying this phenomenon are yet to

be explored. One possibility is that epigenetic silencing of *FLC*, which occurs after cold-

induced transcriptional silencing (*Antoniou-Kourounioti et al., 2018; Hepworth et al., 2018*),

suppresses both *COOLAIR* and *FLC*. A previous report has shown that there is a continuous

increase in the *COOLAIR* level throughout the vernalization period in the *vin3* mutant

(*Swiezewski et al., 2009*). Thus, VIN3-dependent polycomb silencing appears to be at least

partially involved in this mechanism.

Based on previous reports and the results shown here, CBFs are believed to facilitate both

sense and antisense transcription of *FLC,* depending on the duration of cold exposure (*Seo et

al., 2009*). Under intermittent cold, such as a brief cold treatment during daytime, CBFs activate

FLC expression, thus delaying floral transition (*Seo et al., 2009*). By contrast, under conditions

of long-term winter cold, CBFs favor *COOLAIR* activation over *FLC* transcription (*Figure 4*).

Considering that CBF3 protein is enriched at both the 5′- and 3′-ends of *FLC* under warm

conditions (*Figure 1C*), CBFs are likely to directly activate both sense and antisense

transcription of *FLC*. However, CBF3 enrichment at the 5′-end of *FLC* was quickly reduced to

the basal level by prolonged cold (*Figure 1C*), suggesting that CBF3 loses its control over *FLC*

transcription in the sense direction. By contrast, CBF3 protein binds to the *COOLAIR* promoter

during the early phase of vernalization, which leads to *COOLAIR* induction (*Figure 1C*). Thus,

we propose that different types of cold durations determine the direction of *FLC* transcription

by controlling the binding of CBF to the *FLC* locus. This may allow plants to flower in

favorable environments.

Extensive analyses of genes involved in AP have revealed how processing of proximal

COOLAIR is associated with switching *FLC* chromatin state inactive (*Whittaker and Dean,*

2017; Wu et al., 2020). Under warm temperatures, a group of AP proteins involved in RNA 3′-

end processing, such as FCA, FY, FPA, CLEAVAGE STIMULATION FACTOR 64 (CstF64),

and CstF77, promote proximal polyadenylation of *COOLAIR* (*Hornyik et al., 2010; Liu et al.,*

2010). This process is linked to H3K4 demethylation at the *FLC* locus through the interaction

of FCA and FLOWERING LOCUS D (FLD), an AP protein that acts as an H3K4 demethylase,

thus suppressing *FLC* (*Liu et al., 2010*). In contrast with such a role of *COOLAIR* for *FLC*

repression at warm temperatures, there is some controversy regarding the function of

COOLAIR in vernalization-induced epigenetic silencing of *FLC* (*Swiezewski et al., 2009;*

Helliwell et al., 2011; Li et al., 2018; Luo et al., 2019; Luo and He, 2020; Zhao et al., 2021).

The results shown here favor the proposal that *COOLAIR* expression is not required for

vernalization response. The *cbfs* mutant, which exhibited severely reduced *COOLAIR*

induction, showed almost normal vernalization responses such that flowering time is fairly well

accelerated and active histone marks, H3K36me3 and H3K4me3, are reduced, but inactive

histone mark, H3K27me3, is increased on the *FLC* chromatin by vernalization as seen in the

wild-type (*Figures 6 and 7*). Consistently, the *FLC*$_{\Delta COOLAIR}$ mutant, which had almost

undetectable levels of total *COOLAIR* and failed to induce proximal *COOLAIR* by

vernalization, showed both *FLC* suppression and acceleration of flowering by vernalization

similar to that seen in the wild-type (*Figure 7*). Moreover, the freezing temperature treatment

mimicking the first frost did not affect *FLC* suppression, although it caused CBF-mediated

COOLAIR induction, as reported previously (*Figures 4C and 7D; Zhao et al., 2021*). It is

noteworthy that our hypothesis, favoring against *FLC* suppression mediated by vernalization-

induced *COOLAIR,* has been formulated using mutant lines, *cbfs* and *FLC*$_{\Delta COOLAIR}$, instead of

transgenic lines. Thus, we could exclude the possibility that transgenes somehow affect the

vernalization response.

This discrepancy can be explained in several ways. First, different experimental schemes

may affect the vernalization response differently. The *bona fide* evidence favoring the

hypothesis that vernalization-induced *COOLAIR* acts on *FLC* suppression has been obtained

in natural environments with fluctuating temperatures (*Zhao et al., 2021*). However, most of

the experiments in previous studies were performed in constant environments. Thus, it is

possible that fluctuating temperatures have different effects on the vernalization response.

Consistent with such idea, in our experiment using the first frost-mimicking cold treatment,

FLC suppression was not observed, although *COOLAIR* was strongly activated, as reported

(*Zhao et al., 2021*). The difference between the two experimental schemes is fluctuating versus

constant cold conditions. The other reason may be due to the small level of CAS (convergent

antisense transcript of *COOLAIR*) produced at the 5′-end of *FLC* in transgenic lines defective

of *COOLAIR* promoter or even in the *FLC$_{\Delta COOLAIR}$* mutant (*Zhao et al., 2021*), which may

somehow affect the epigenetic silencing of *FLC*. Otherwise, the effect of *COOLAIR* on *FLC*

silencing may be obscured due to the functional redundancy between *COOLAIR* and other

lncRNA, such as *COLDAIR* in Arabidopsis, as suggested previously (*Castaings et al., 2014*).

Although cold stress and vernalization elicit different developmental responses, both are

triggered by the same physical environment. Therefore, it is logical that a common signaling

network governs both the processes. *CBF* genes have been highlighted for their role in cold

stress responses (*Gilmour et al., 1998; Liu et al., 1998; Jia et al., 2016; Zhao et al., 2016*).

CBFs rapidly accumulate upon short-term cold exposure, thereby activating the transcription

of cold-responsive genes, such as *COR15A* and *RD29A* (*Stockinger et al., 1997; Liu et al.,*

1998). Interestingly, CBFs are also highly accumulated during the vernalization process and

mediate long-term cold-induced *COOLAIR* expression. Hence, our results support that cold

signaling and vernalization pathways are not sharply distinguishable, as suggested previously

(*Seo et al., 2009*).

Materials and Methods

Plant materials and treatments

Arabidopsis thaliana Columbia-0 (Col-0), Wassilewskija-2 (Ws-2), and Col:*FRI*[SF2] (*FRI* Col) were used as wild types in this study. "Wild type" in this paper mainly refers to *FRI* Col unless otherwise specified. *FRI* Col (*Lee et al., 1994*), *cbfs-1*, *pCBF3::CBF3-myc* (*Jia et al., 2016*), *pSuper::CBF3-myc* (*Liu et al., 2017*), *35S::CBF1*, *35S::CBF2*, and *35S::CBF3* (*Gilmour et al., 2004*) have previously been described. *FRI cbfs-1* was generated by crossing *FRI* Col with *cbfs-1* stated above. $FLC_{\Delta COOLAIR}$ was generated by crossing *FRI* Col with $FLC_{\Delta COOLAIR}$ in Col-0 background described previously (*Luo et al., 2019*). For all experiments using plant materials, seeds were sown on half-strength Murashige & Skoog (MS) media (Duchefa) containing 1% sucrose and 1% plant agar (Duchefa) and stratified at 4°C for 3 days. Plants were germinated and grown at 22ºC under a short day (SD) cycle (8 h light/16 h dark) with cool white fluorescent lights (100 μmol m^{-2} s^{-1}). For the vernalization treatment, plants were grown at 22ºC for the prescribed period and then transferred to a 4ºC growth chamber under an SD cycle. To adjust the developmental stage, the plants to be vernalized for 40 days (40V), 30 days (30V), 20 days (20V), 10 days (10V), and 1 day (1V) were transferred to 4ºC after being grown at 22ºC for 10 days, 12 days, 14 days, 15 days, and 16 days respectively. Non-vernalized (NV) plants were grown at 22ºC for 16 days. For 0. 0.5, 1, 1.5, 2, 3, 6, and 24 hours of cold treatment, plants were grown at 22ºC for 16 days before the 4ºC treatment. For the freezing treatment, 20V plants were transferred to –1ºC, dark condition for 8 hours (8hF). To measure flowering time of vernalized plants, plants were transferred to 22ºC under a long day (LD) cycle (16 h light/8 h dark) directly after the vernalization treatment.

Plasmid construction

To generate the *35S::CBF3-HA* construct, the entire coding sequence of *CBF3* was amplified by PCR and was cloned into the *HBT-HA-NOS* plasmid (*Yoo et al., 2007*). For $pCOOLAIR_{DRE}::LUC$ and $pCOOLAIR_{DRE}{}^{m}::LUC$ construction, the 1 kb sequence containing the *COOLAIR* promoter, the 3' UTR and last exon of *FLC* was amplified by PCR and was

443   cloned into *LUC-NOS* plasmid (*Hwang and Sheen, 2001*). To generate *pCOOLAIR$_{DRE}^{m}$::LUC,*

444   two CRT/DREs in the *COOLAIR* promoter, 5'-AGGTCGGT-3' and 5'-ACCGACAT-3', were

445   replaced with 5'-CGAGGTGT-3' and 5'-TGAACCCA-3', respectively. For *pFLC::LUC*

446   construction, the 1 kb sequence containing the promoter, 5' UTR, and first exon of *FLC* was

447   amplified by PCR and was cloned into the *LUC-NOS* plasmid.

Electrophoretic mobility shift assay (EMSA)

Maltose binding protein (MBP) and MBP-CBF3 recombinant fusion protein were induced by

500 μM of isopropyl β-D-thiogalactoside (IPTG) in *E. coli* BL21 (DE3) strain. Cell extracts

were isolated with buffer containing 20 mM Tris-HCl (pH 7.4), 1 mM EDTA, 200 mM NaCl,

10% Glycerol, 1 mM DTT, and 1 mM PMSF. Proteins were purified from cell extracts using

MBPTrap™ HP column (GE Healthcare) with ÄKTA FPLC system (Amersham Biosciences).

The Cy5-labelled probes and unlabelled competitors were generated by annealing 25 bp-length

oligonucleotides. EMSA was performed as previously described with few modifications (*Seo*

et al., 2012). 5 μM of proteins and 100 nM of Cy5-labelled probe were incubated together at

room temperature in a binding buffer containing 10 mM Tris-HCl (pH 7.5), 1 mM EDTA, 50

mM NaCl, 5% glycerol, and 5 mM DTT for each EMSA reaction. For the competition assay,

the reaction mixtures were incubated in the presence of 100-fold molar excess of each

competitor. The reaction mixtures were electrophoresed at 100 V after incubation. The Cy5

signals were detected using WSE-6200H LuminoGraph II (ATTO).

Chromatin immunoprecipitation (ChIP) assay

Approximately 4 g of seedlings grown on MS plate were collected and cross-linked using 1%

(v/v) formaldehyde. Nuclei were isolated from seedlings using buffer containing 20 mM

PIPES-KOH pH 7.6, 1 M hexylene glycol, 10 mM MgCl2, 0.1 mM EGTA, 15 mM NaCl, 60

mM KCl, 0.5% Triton-X, 5 mM β-mercaptoethanol, and 1X cOmplete™ EDTA-free Protease

Inhibitor Cocktail (Roche) as described previously with few modifications (*Shu et al., 2014*).

Isolated nuclei were lysed with buffer containing 50 mM Tris-HCl (pH 7.4), 150 mM NaCl, 1%

(v/v) Triton X-100, and 1% SDS (w/v), and were subsequently sonicated using Digital

Sonifier® (Branson). Sheared chromatin was diluted with buffer containing 50 mM Tris-HCl

(pH 7.4), 150 mM NaCl, 1% (v/v) Triton X-100, and 1 mM EDTA. Immunoprecipitation was

undertaken by incubating sheared chromatin together with Protein G Sepharose™ 4 Fast Flow

beads (GE Healthcare) and antibodies. Anti-myc (MBL, #M192-3) and normal mouse IgG$_1$

(Santa Cruz, #sc-3877) were used to detect CBF3 protein enrichment on *FLC* locus, and anti-

H3K36me3 (Abcam, #Ab9050), H3K4me3 (Millipore, #07-473), and anti-H3K27me3

(Millipore, #07-449) were used to detect histone modifications deposited on *FLC* chromatin.

DNA was extracted with phenol:chloroform:isoamyl alcohol (25:24:1, v/v) or with Chelex®

100 Resin according to the manufacturer's instruction.

Gene expression analysis

Total RNA was isolated from ~100 mg of seedlings grown on MS agar plate using TRI

Reagent® (Sigma-Aldrich) or TS™ Plant Mini Kit (Taeshin Bio Science, Gyeonggi-do,

Republic of Korea). cDNA was generated using 4 μg of total RNA, 5 units of Recombinant

DNase I (TaKaRa), 200 units of RevertAid™ Reverse Transcriptase (Thermo Scientific), and

buffer containing 0.25 mM dNTP and 0.1 μg oligo(dT)$_{18}$. Quantitative PCR analysis was

performed using iQ™ SYBR® Green Supermix (Bio-Rad) and CFX96 Real-Time PCR system

(Bio-Rad).

Immunoblotting

Total protein was isolated from ~100 mg of seedlings using buffer containing 50 mM Tris-HCl

(pH 7.5), 100 mM NaCl, 10 mM MgCl$_2$, 1 mM EDTA, 10% glycerol, 1 mM PMSF, 1 mM DTT

and the 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail (Roche). 50 μg of total protein

was loaded onto SDS-PAGE gels and separated by electrophoresis. The proteins were

transferred to PVDF membranes (Amersham Biosciences) and probed with anti-myc (MBL,

#M192-3; 1:10000 dilution) antibody overnight at 4°C. Then, the samples were probed with

horseradish peroxidase (HRP)-conjugated anti-mouse IgG (Cell Signaling, #7076; 1:10000

dilution) antibody at room temperature. The signals were detected by ImageQuant™ LAS 4000

(GE Healthcare) using WesternBright™ Sirius ECL solution (Advansta).

Luciferase assay using Arabidopsis protoplast

Protoplast isolation and transfection were undertaken as previously described with few

modifications (*Yoo et al., 2007*). Protoplast was isolated from leaves of SD-grown wild-type

plant (Col-0) using a buffer containing 150 mg Cellulase Onozuka™ R-10 (Yakult), 50 mg

Maceroenzyme™ R-10 (Yakult), 20 mM KCl, 20 mM MES-KOH (pH 5.6), 0.4 M D-mannitol,

10 mM $CaCl_2$, 5 mM β-mercaptoethanol, 0.1 g bovine serum albumin. For protoplast

transfection, 200 μg of plasmid DNA and isolated protoplast were incubated together in buffer

containing 0.1 M D-mannitol, 50 mM $CaCl_2$, and 20% (w/v) PEG. Luciferase activity in

protoplast was measured using Luciferase Assay System (Promega) and MicroLumat Plus

LB96V microplate luminometer (Berthold Technologies).

Acknowledgements

We would like to thank Dr. Shuhua Yang (China Agricultural University, China) for providing

cbfs-1, *pSuper::CBF3-myc* and *pCBF3::CBF3-myc* seeds, and Dr. Yuehui He (Peking

University, China) for providing *FLC$_{\Delta COOLAIR}$* seeds. This work was carried out with the support

of Cooperative Research Program for Agriculture Science and Technology Development (No.

PJ01315201 and PJ01315401), Rural Development Administration, Republic of Korea, and of

the National Research Foundation of Korea (NRF) grant funded by the Korea government

(MSIT) (No. 2019R1A2C2004313). This work was also supported by NRF grant funded by

the Ministry of Science, Republic of Korea (NRF-2021R1A5A1032428). M. Jeon was

supported by Brain Korea 21 Plus Project of the NRF grant.

Competing interests

The authors declare that no competing interests exist.

Supplementary Files

Supplementary File 1. Oligonucleotide sequences used in this study.

References

Angel, A, Song, J, Dean, C, Howard, M. 2011. A Polycomb-based switch underlying quantitative epigenetic memory. *Nature* **476:** 105-108. DOI: https://doi.org/10.1038/nature10241

Antoniou-Kourounioti, RL, Hepworth, J, Heckmann, A, Duncan, S, Questa, J, Rosa, S, Sall, T, Holm, S, Dean, C, Howard, M. 2018. Temperature Sensing Is Distributed throughout the Regulatory Network that Controls FLC Epigenetic Silencing in Vernalization. *Cell Syst* **7:** 643-655 e649. DOI: https://doi.org/10.1016/j.cels.2018.10.011

Baker, SS, Wilhelm, KS, Thomashow, MF. 1994. The 5'-region of Arabidopsis thaliana cor15a has cis-acting elements that confer cold-, drought- and ABA-regulated gene expression. *Plant Mol Biol* **24:** 701-713. DOI: https://doi.org/10.1007/BF00029852

Bastow, R, Mylne, JS, Lister, C, Lippman, Z, Martienssen, RA, Dean, C. 2004. Vernalization requires epigenetic silencing of FLC by histone methylation. *Nature* **427:** 164-167. DOI: https://doi.org/10.1038/nature02269

Berry, S, Dean, C. 2015. Environmental perception and epigenetic memory: mechanistic insight through FLC. *Plant J* **83:** 133-148. DOI: https://doi.org/10.1111/tpj.12869

Castaings, L, Bergonzi, S, Albani, MC, Kemi, U, Savolainen, O, Coupland, G. 2014. Evolutionary conservation of cold-induced antisense RNAs of FLOWERING LOCUS C in Arabidopsis thaliana perennial relatives. *Nat Commun* **5:** 4457. DOI: https://doi.org/10.1038/ncomms5457

Choi, K, Kim, J, Hwang, HJ, Kim, S, Park, C, Kim, SY, Lee, I. 2011. The FRIGIDA complex activates transcription of FLC, a strong flowering repressor in Arabidopsis, by recruiting chromatin modification factors. *Plant Cell* **23:** 289-303. DOI: https://doi.org/10.1105/tpc.110.075911

Chouard, P. 1960. Vernalization and Its Relations to Dormancy. *Annual Review of Plant Physiology and Plant Molecular Biology* **11:** 191-238. DOI: https://doi.org/DOI 10.1146/annurev.pp.11.060160.001203

Clarke, JH, Dean, C. 1994. Mapping FRI, a locus controlling flowering time and vernalization response in Arabidopsis thaliana. *Mol Gen Genet* **242:** 81-89. DOI: https://doi.org/10.1007/BF00277351

Csorba, T, Questa, JI, Sun, Q, Dean, C. 2014. Antisense COOLAIR mediates the coordinated switching of chromatin states at FLC during vernalization. *Proc Natl Acad Sci U S A* **111:** 16160-16165. DOI: https://doi.org/10.1073/pnas.1419030111

De Lucia, F, Crevillen, P, Jones, AM, Greb, T, Dean, C. 2008. A PHD-polycomb repressive complex 2 triggers the epigenetic silencing of FLC during vernalization. *Proc Natl Acad Sci U S A* **105:** 16831-16836. DOI: https://doi.org/10.1073/pnas.0808687105

Ding, Y, Jia, Y, Shi, Y, Zhang, X, Song, C, Gong, Z, Yang, S. 2018. OST1-mediated BTF3L

phosphorylation positively regulates CBFs during plant cold responses. *EMBO J* **37**. DOI: https://doi.org/10.15252/embj.201798228

Dong, MA, Farre, EM, Thomashow, MF. 2011. Circadian clock-associated 1 and late elongated hypocotyl regulate expression of the C-repeat binding factor (CBF) pathway in Arabidopsis. *Proc Natl Acad Sci U S A* **108:** 7241-7246. DOI: https://doi.org/10.1073/pnas.1103741108

Fowler, S, Thomashow, MF. 2002. Arabidopsis transcriptome profiling indicates that multiple regulatory pathways are activated during cold acclimation in addition to the CBF cold response pathway. *Plant Cell* **14:** 1675-1690. DOI: https://doi.org/10.1105/tpc.003483

Gazzani, S, Gendall, AR, Lister, C, Dean, C. 2003. Analysis of the molecular basis of flowering time variation in Arabidopsis accessions. *Plant Physiol* **132:** 1107-1114. DOI: https://doi.org/10.1104/pp.103.021212

Gilmour, SJ, Fowler, SG, Thomashow, MF. 2004. Arabidopsis transcriptional activators CBF1, CBF2, and CBF3 have matching functional activities. *Plant Mol Biol* **54:** 767-781. DOI: https://doi.org/10.1023/B:PLAN.0000040902.06881.d4

Gilmour, SJ, Zarka, DG, Stockinger, EJ, Salazar, MP, Houghton, JM, Thomashow, MF. 1998. Low temperature regulation of the Arabidopsis CBF family of AP2 transcriptional activators as an early step in cold-induced COR gene expression. *Plant J* **16:** 433-442. DOI: https://doi.org/10.1046/j.1365-313x.1998.00310.x

Helliwell, CA, Robertson, M, Finnegan, EJ, Buzas, DM, Dennis, ES. 2011. Vernalization-repression of Arabidopsis FLC requires promoter sequences but not antisense transcripts. *PLoS One* **6:** e21513. DOI: https://doi.org/10.1371/journal.pone.0021513

Helliwell, CA, Wood, CC, Robertson, M, James Peacock, W, Dennis, ES. 2006. The Arabidopsis FLC protein interacts directly in vivo with SOC1 and FT chromatin and is part of a high-molecular-weight protein complex. *Plant J* **46:** 183-192. DOI: https://doi.org/10.1111/j.1365-313X.2006.02686.x

Heo, JB, Sung, S. 2011. Vernalization-mediated epigenetic silencing by a long intronic noncoding RNA. *Science* **331:** 76-79. DOI: https://doi.org/10.1126/science.1197349

Hepworth, J, Antoniou-Kourounioti, RL, Bloomer, RH, Selga, C, Berggren, K, Cox, D, Collier Harris, BR, Irwin, JA, Holm, S, Sall, T, Howard, M, Dean, C. 2018. Absence of warmth permits epigenetic memory of winter in Arabidopsis. *Nat Commun* **9:** 639. DOI: https://doi.org/10.1038/s41467-018-03065-7

Hornyik, C, Duc, C, Rataj, K, Terzi, LC, Simpson, GG. 2010. Alternative polyadenylation of antisense RNAs and flowering time control. *Biochem Soc Trans* **38:** 1077-1081. DOI: https://doi.org/10.1042/BST0381077

Hwang, I, Sheen, J. 2001. Two-component circuitry in Arabidopsis cytokinin signal

transduction. *Nature* **413:** 383-389. DOI: https://doi.org/10.1038/35096500

Jia, Y, Ding, Y, Shi, Y, Zhang, X, Gong, Z, Yang, S. 2016. The cbfs triple mutants reveal the
essential functions of CBFs in cold acclimation and allow the definition of CBF
regulons in Arabidopsis. *New Phytol* **212:** 345-353. DOI:
https://doi.org/10.1111/nph.14088

Johanson, U, West, J, Lister, C, Michaels, S, Amasino, R, Dean, C. 2000. Molecular analysis
of FRIGIDA, a major determinant of natural variation in Arabidopsis flowering time.
Science **290:** 344-347. DOI: https://doi.org/10.1126/science.290.5490.344

Kim, DH, Sung, S. 2013. Coordination of the vernalization response through a VIN3 and FLC
gene family regulatory network in Arabidopsis. *Plant Cell* **25:** 454-469. DOI:
https://doi.org/10.1105/tpc.112.104760

Kim, DH, Sung, S. 2017a. The Binding Specificity of the PHD-Finger Domain of VIN3
Moderates Vernalization Response. *Plant Physiol* **173:** 1258-1268. DOI:
https://doi.org/10.1104/pp.16.01320

Kim, DH, Sung, S. 2017b. Vernalization-Triggered Intragenic Chromatin Loop Formation by
Long Noncoding RNAs. *Dev Cell* **40:** 302-312 e304. DOI:
https://doi.org/10.1016/j.devcel.2016.12.021

Koornneef, M, Alonso-Blanco, C, Blankestijn-de Vries, H, Hanhart, CJ, Peeters, AJ. 1998.
Genetic interactions among late-flowering mutants of Arabidopsis. *Genetics* **148:**
885-892.

Koornneef, M, Blankestijndevries, H, Hanhart, C, Soppe, W, Peeters, T. 1994. The
Phenotype of Some Late-Flowering Mutants Is Enhanced by a Locus on Chromosome-
5 That Is Not Effective in the Landsberg Erecta Wild-Type. *Plant Journal* **6:** 911-
919. DOI: https://doi.org/DOI 10.1046/j.1365-313X.1994.6060911.x

Lee, ES, Park, JH, Wi, SD, Kang, CH, Chi, YH, Chae, HB, Paeng, SK, Ji, MG, Kim, WY,
Kim, MG, Yun, DJ, Stacey, G, Lee, SY. 2021. Redox-dependent structural switch and
CBF activation confer freezing tolerance in plants. *Nat Plants* **7:** 914-922. DOI:
https://doi.org/10.1038/s41477-021-00944-8

Lee, H, Suh, SS, Park, E, Cho, E, Ahn, JH, Kim, SG, Lee, JS, Kwon, YM, Lee, I. 2000.
The AGAMOUS-LIKE 20 MADS domain protein integrates floral inductive pathways
in Arabidopsis. *Genes Dev* **14:** 2366-2376. DOI:
https://doi.org/10.1101/gad.813600

Lee, I, Bleecker, A, Amasino, R. 1993. Analysis of naturally occurring late flowering in
Arabidopsis thaliana. *Mol Gen Genet* **237:** 171-176. DOI:
https://doi.org/10.1007/BF00282798

Lee, I, Michaels, SD, Masshardt, AS, Amasino, RM. 1994. The Late-Flowering Phenotype
of Frigida and Mutations in Luminidependens Is Suppressed in the Landsberg Erecta
Strain of Arabidopsis. *Plant Journal* **6:** 903-909. DOI: https://doi.org/DOI
10.1046/j.1365-313X.1994.6060903.x

Li, Z, Jiang, D, He, Y. 2018. FRIGIDA establishes a local chromosomal environment for FLOWERING LOCUS C mRNA production. *Nat Plants* **4:** 836-846. DOI: https://doi.org/10.1038/s41477-018-0250-6

Liu, F, Marquardt, S, Lister, C, Swiezewski, S, Dean, C. 2010. Targeted 3' processing of antisense transcripts triggers Arabidopsis FLC chromatin silencing. *Science* **327:** 94-97. DOI: https://doi.org/10.1126/science.1180278

Liu, Q, Kasuga, M, Sakuma, Y, Abe, H, Miura, S, Yamaguchi-Shinozaki, K, Shinozaki, K. 1998. Two transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought- and low-temperature-responsive gene expression, respectively, in Arabidopsis. *Plant Cell* **10:** 1391-1406. DOI: https://doi.org/10.1105/tpc.10.8.1391

Liu, Z, Jia, Y, Ding, Y, Shi, Y, Li, Z, Guo, Y, Gong, Z, Yang, S. 2017. Plasma Membrane CRPK1-Mediated Phosphorylation of 14-3-3 Proteins Induces Their Nuclear Import to Fine-Tune CBF Signaling during Cold Response. *Mol Cell* **66:** 117-128 e115. DOI: https://doi.org/10.1016/j.molcel.2017.02.016

Luo, X, Chen, T, Zeng, X, He, D, He, Y. 2019. Feedback Regulation of FLC by FLOWERING LOCUS T (FT) and FD through a 5' FLC Promoter Region in Arabidopsis. *Mol Plant* **12:** 285-288. DOI: https://doi.org/10.1016/j.molp.2019.01.013

Luo, X, He, Y. 2020. Experiencing winter for spring flowering: A molecular epigenetic perspective on vernalization. *J Integr Plant Biol* **62:** 104-117. DOI: https://doi.org/10.1111/jipb.12896

Marquardt, S, Raitskin, O, Wu, Z, Liu, F, Sun, Q, Dean, C. 2014. Functional consequences of splicing of the antisense transcript COOLAIR on FLC transcription. *Mol Cell* **54:** 156-165. DOI: https://doi.org/10.1016/j.molcel.2014.03.026

Maruyama, K, Sakuma, Y, Kasuga, M, Ito, Y, Seki, M, Goda, H, Shimada, Y, Yoshida, S, Shinozaki, K, Yamaguchi-Shinozaki, K. 2004. Identification of cold-inducible downstream genes of the Arabidopsis DREB1A/CBF3 transcriptional factor using two microarray systems. *Plant J* **38:** 982-993. DOI: https://doi.org/10.1111/j.1365-313X.2004.02100.x

McKhann, HI, Gery, C, Berard, A, Leveque, S, Zuther, E, Hincha, DK, De Mita, S, Brunel, D, Teoule, E. 2008. Natural variation in CBF gene sequence, gene expression and freezing tolerance in the Versailles core collection of Arabidopsis thaliana. *BMC Plant Biol* **8:** 105. DOI: https://doi.org/10.1186/1471-2229-8-105

Medina, J, Bargues, M, Terol, J, Perez-Alonso, M, Salinas, J. 1999. The Arabidopsis CBF gene family is composed of three genes encoding AP2 domain-containing proteins whose expression Is regulated by low temperature but not by abscisic acid or dehydration. *Plant Physiol* **119:** 463-470. DOI: https://doi.org/10.1104/pp.119.2.463

Michaels, SD, Amasino, RM. 1999. FLOWERING LOCUS C encodes a novel MADS

domain protein that acts as a repressor of flowering. *Plant Cell* **11:** 949-956. DOI: https://doi.org/10.1105/tpc.11.5.949

Michaels, SD, Amasino, RM. 2000. Memories of winter: vernalization and the competence to flower. *Plant Cell and Environment* **23:** 1145-1153. DOI: https://doi.org/DOI 10.1046/j.1365-3040.2000.00643.x

Michaels, SD, Amasino, RM. 2001. Loss of FLOWERING LOCUS C activity eliminates the late-flowering phenotype of FRIGIDA and autonomous pathway mutations but not responsiveness to vernalization. *Plant Cell* **13:** 935-941. DOI: https://doi.org/10.1105/tpc.13.4.935

Michaels, SD, He, Y, Scortecci, KC, Amasino, RM. 2003. Attenuation of FLOWERING LOCUS C activity as a mechanism for the evolution of summer-annual flowering behavior in Arabidopsis. *Proc Natl Acad Sci U S A* **100:** 10102-10107. DOI: https://doi.org/10.1073/pnas.1531467100

Michaels, SD, Himelblau, E, Kim, SY, Schomburg, FM, Amasino, RM. 2005. Integration of flowering signals in winter-annual Arabidopsis. *Plant Physiol* **137:** 149-156. DOI: https://doi.org/10.1104/pp.104.052811

Mylne, JS, Barrett, L, Tessadori, F, Mesnage, S, Johnson, L, Bernatavichute, YV, Jacobsen, SE, Fransz, P, Dean, C. 2006. LHP1, the Arabidopsis homologue of HETEROCHROMATIN PROTEIN1, is required for epigenetic silencing of FLC. *Proc Natl Acad Sci U S A* **103:** 5012-5017. DOI: https://doi.org/10.1073/pnas.0507427103

Napp-Zinn, K. 1955. Genetische Grundlagen des Kältebedürfnisses beiArabidopsis thaliana (L.) HEYNH. *Die Naturwissenschaften* **42:** 650. DOI: https://doi.org/10.1007/BF00622395

Novillo, F, Medina, J, Salinas, J. 2007. Arabidopsis CBF1 and CBF3 have a different function than CBF2 in cold acclimation and define different gene classes in the CBF regulon. *Proc Natl Acad Sci U S A* **104:** 21002-21007. DOI: https://doi.org/10.1073/pnas.0705639105

O'Malley, RC, Huang, SC, Song, L, Lewsey, MG, Bartlett, A, Nery, JR, Galli, M, Gallavotti, A, Ecker, JR. 2016. Cistrome and Epicistrome Features Shape the Regulatory DNA Landscape. *Cell* **166:** 1598. DOI: https://doi.org/10.1016/j.cell.2016.08.063

Park, J, Lim, CJ, Shen, M, Park, HJ, Cha, JY, Iniesto, E, Rubio, V, Mengiste, T, Zhu, JK, Bressan, RA, Lee, SY, Lee, BH, Jin, JB, Pardo, JM, Kim, WY, Yun, DJ. 2018. Epigenetic switch from repressive to permissive chromatin in response to cold stress. *Proc Natl Acad Sci U S A* **115:** E5400-E5409. DOI: https://doi.org/10.1073/pnas.1721241115

Park, S, Lee, CM, Doherty, CJ, Gilmour, SJ, Kim, Y, Thomashow, MF. 2015. Regulation of the Arabidopsis CBF regulon by a complex low-temperature regulatory network.

Plant J **82:** 193-207. DOI: https://doi.org/10.1111/tpj.12796

Seo, E, Lee, H, Jeon, J, Park, H, Kim, J, Noh, YS, Lee, I. 2009. Crosstalk between cold response and flowering in Arabidopsis is mediated through the flowering-time gene SOC1 and its upstream negative regulator FLC. *Plant Cell* **21:** 3185-3197. DOI: https://doi.org/10.1105/tpc.108.063883

Seo, PJ, Park, MJ, Lim, MH, Kim, SG, Lee, M, Baldwin, IT, Park, CM. 2012. A self-regulatory circuit of CIRCADIAN CLOCK-ASSOCIATED1 underlies the circadian clock regulation of temperature responses in Arabidopsis. *Plant Cell* **24:** 2427-2442. DOI: https://doi.org/10.1105/tpc.112.098723

Sheldon, CC, Burn, JE, Perez, PP, Metzger, J, Edwards, JA, Peacock, WJ, Dennis, ES. 1999. The FLF MADS box gene: a repressor of flowering in Arabidopsis regulated by vernalization and methylation. *Plant Cell* **11:** 445-458. DOI: https://doi.org/10.1105/tpc.11.3.445

Sheldon, CC, Rouse, DT, Finnegan, EJ, Peacock, WJ, Dennis, ES. 2000. The molecular basis of vernalization: the central role of FLOWERING LOCUS C (FLC). *Proc Natl Acad Sci U S A* **97:** 3753-3758. DOI: https://doi.org/10.1073/pnas.060023597

Shu, H, Nakamura, M, Siretskiy, A, Borghi, L, Moraes, I, Wildhaber, T, Gruissem, W, Hennig, L. 2014. Arabidopsis replacement histone variant H3.3 occupies promoters of regulated genes. *Genome Biol* **15:** R62. DOI: https://doi.org/10.1186/gb-2014-15-4-r62

Stockinger, EJ, Gilmour, SJ, Thomashow, MF. 1997. Arabidopsis thaliana CBF1 encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. *Proc Natl Acad Sci U S A* **94:** 1035-1040. DOI: https://doi.org/10.1073/pnas.94.3.1035

Sung, S, Amasino, RM. 2004. Vernalization in Arabidopsis thaliana is mediated by the PHD finger protein VIN3. *Nature* **427:** 159-164. DOI: https://doi.org/10.1038/nature02195

Swiezewski, S, Liu, F, Magusin, A, Dean, C. 2009. Cold-induced silencing by long antisense transcripts of an Arabidopsis Polycomb target. *Nature* **462:** 799-802. DOI: https://doi.org/10.1038/nature08618

Tian, Y, Zheng, H, Zhang, F, Wang, S, Ji, X, Xu, C, He, Y, Ding, Y. 2019. PRC2 recruitment and H3K27me3 deposition at FLC require FCA binding of COOLAIR. *Sci Adv* **5:** eaau7246. DOI: https://doi.org/10.1126/sciadv.aau7246

Whittaker, C, Dean, C. 2017. The FLC Locus: A Platform for Discoveries in Epigenetics and Adaptation. *Annu Rev Cell Dev Biol* **33:** 555-575. DOI: https://doi.org/10.1146/annurev-cellbio-100616-060546

Wood, CC, Robertson, M, Tanner, G, Peacock, WJ, Dennis, ES, Helliwell, CA. 2006. The Arabidopsis thaliana vernalization response requires a polycomb-like protein complex

that also includes VERNALIZATION INSENSITIVE 3. *Proc Natl Acad Sci U S A* **103:** 14631-14636. DOI: https://doi.org/10.1073/pnas.0606385103

Wu, Z, Fang, X, Zhu, D, Dean, C. 2020. Autonomous Pathway: FLOWERING LOCUS C Repression through an Antisense-Mediated Chromatin-Silencing Mechanism. *Plant Physiol* **182:** 27-37. DOI: https://doi.org/10.1104/pp.19.01009

Yang, H, Berry, S, Olsson, TSG, Hartley, M, Howard, M, Dean, C. 2017. Distinct phases of Polycomb silencing to hold epigenetic memory of cold in Arabidopsis. *Science* **357:** 1142-1145. DOI: https://doi.org/10.1126/science.aan1121

Yang, H, Howard, M, Dean, C. 2014. Antagonistic roles for H3K36me3 and H3K27me3 in the cold-induced epigenetic switch at Arabidopsis FLC. *Curr Biol* **24:** 1793-1797. DOI: https://doi.org/10.1016/j.cub.2014.06.047

Yoo, SD, Cho, YH, Sheen, J. 2007. Arabidopsis mesophyll protoplasts: a versatile cell system for transient gene expression analysis. *Nat Protoc* **2:** 1565-1572. DOI: https://doi.org/10.1038/nprot.2007.199

Zhao, C, Zhang, Z, Xie, S, Si, T, Li, Y, Zhu, JK. 2016. Mutational Evidence for the Critical Role of CBF Transcription Factors in Cold Acclimation in Arabidopsis. *Plant Physiol* **171:** 2744-2759. DOI: https://doi.org/10.1104/pp.16.00533

Zhao, Y, Zhu, P, Hepworth, J, Bloomer, R, Antoniou-Kourounioti, RL, Doughty, J, Heckmann, A, Xu, C, Yang, H, Dean, C. 2021. Natural temperature fluctuations promote COOLAIR regulation of FLC. *Genes Dev* **35:** 888-898. DOI: https://doi.org/10.1101/gad.348362.121

Figure legends

Figure 1. CBF3 directly binds to the CRT/DREs at the 3′-end of *FLC*.

(A) Comparison of sequences around the 3′-end of the *FLC* orthologs from Arabidopsis relatives. The upper graphic presents the gene structure of *AtFLC*. The black bars, thick lines, thicker lines indicate exons, introns, and UTRs, respectively. The blue-coloured box presents the region used for sequence comparison among 6 orthologous genes from 5 plant species. In the sequence alignment below, the grey box indicates the 3′-UTR of *FLC* orthologs, the blue arrow indicates the transcriptional start site of *AtCOOLAIR*, and the red dotted boxes indicate CRT/DREs. At, *Arabidopsis thaliana*; Cr, *Capsella rubella*; Al, *Arabidopsis lyrata*; Aa, *Arabis*

alpina; Th, *Thellungiella halophila*.

(B) EMSA using one of the CRT/DREs located at the *COOLAIR* promoter, DRE1, as a probe.

In the upper graphic showing *AtFLC* gene structure, CRT/DRE-like sequences are marked as

blue arrows and labelled as a, b, c, d for CRT/DRE-like sequences and as 1, 2 for CRT/DRE

sequences. For the competition assay, these CRT/DRE-like sequences and mutant forms of

DRE1, DRE2 were used as competitors. A 100-fold molar excess of unlabelled competitors

was added. FP, free probe.

(C) ChIP assay result showing the enrichment of CBF3-myc protein on the *AtFLC* locus.

Samples of NV, 10V, 20V, and 40V plants of *pSuper::CBF3-myc* were collected at zeitgeber

time (ZT) 4 in an SD cycle. The CBF3-chromatin complex was immunoprecipitated (IP) with

anti-myc antibodies (blue bars), and mouse IgG (black bars) was used as a control. Positions

of qPCR amplicons used for ChIP-qPCR analysis are illustrated as P1, P2, and P3 in the upper

graphic. The red arrow in the graphic denotes the position of two CRT/DREs located on the

COOLAIR promoter. ChIP-qPCR results have been represented as mean ± SEM of the 3

biological replicates in the lower panel. Dots represent each data-point. Relative enrichments

of the IP/5% input were normalized to that of *pTUB2*. The blue shading indicates a cold period.

Asterisks indicate a significant difference between IgG and anti-myc ChIP-qPCR results at

each vernalization time-point (*, $P<0.05$; **, $P<0.01$; unpaired Student's *t*-test). ns, not

significant.

Figure 1—figure supplement 2. CBF3 binds to DRE1 and DRE2 on the *COOLAIR* promoter and DREc on the first exon of *FLC*.

(A) Schematic showing the positions of six CRT/DRE-like sequences distributed on the *AtFLC* locus. CRT/DRE sequences have been labelled as DRE1, -2, and CRT/DRE-like sequences have been labelled as DREa, -b, -c, and -d. The black bars, white lines, and light blue lines represent exons, introns, and UTRs of *AtFLC*, respectively. The red-coloured bases indicate the core sequence (CCGAC) of CRT/DRE. The blue-coloured bases represent bases known to

832   determine the binding affinity of CBF3 to CRT/DRE (*Maruyama et al., 2004*).

833   (B-C) EMSA using DRE1 (B) or DRE2 (C) as probe. For competition assay, DRE1, DRE2,

834   and DREc were used as competitors. Molar excess (1-, 10-, and 100-fold) of unlabelled

835   competitors was added. FP, free probe.

Figure 2. *COOLAIR* is one of the CBF regulons.

(A) Expression levels of total *COOLAIR* and *COOLAIR* isoforms in the wild-type (Col-0) and *CBF3*-overexpressing plants (*pSuper::CBF3-myc*). The gene structures of *FLC* and *COOLAIR* variants are illustrated in the upper panel. The thin black arrow indicates the transcriptional start site of *COOLAIR*. The thick black arrows indicate exons of each *COOLAIR* isoform. The structures of proximal (I) and distal (IIi and IIii) *COOLAIR* isoforms are shown. The primer sets used for total, proximal (I), and distal (IIi, IIii) COOLAIRs are marked in Figure 2-supplement 1. Relative transcript levels of total *COOLAIR* and *COOLAIR* variants have been represented as mean ± SEM of 3 biological replicates. Relative expression levels of *UBC* were

normalized to that of Col-0. Dots represent each data-point. Asterisks indicate a significant

difference, as compared to the wild-type (*, $P<0.05$; **, $P<0.01$; ***, $P<0.001$; unpaired

Student's *t*-test).

(B) Expression dynamics of total *COOLAIR* after short-term cold (0, 0.5, 1, 1.5, 2, 3, 6, and 24

h) treatment. Wild-types (*FRI* Col) were subjected to 4°C cold and harvested at each time-point.

Relative transcript levels of total *COOLAIR* to *UBC* were normalized to that of non-cold-

treated wild-type. The values have been represented as mean ± SEM of 3 biological replicates.

Dots represent each data-point. The blue shading indicates periods under cold treatment.

Significant differences have been marked using different letters (a, b; $P<0.05$; one-way

ANOVA followed by Tukey's post-hoc test).

(C) Transcript levels of total *COOLAIR* and *COOLAIR* isoforms in wild-type and *cbfs-1* mutant

before (–Cold) and after (+Cold) a day of 4°C cold treatment. Relative levels of total *COOLAIR*

and *COOLAIR* variants were normalized to that of *PP2A*. Values have been represented as

mean ± SEM of 3 biological replicates. Dots indicate each data-point. Significant differences

have been marked using different letters (a, b; $P<0.05$; two-way ANOVA followed by Tukey's

post-hoc test).

Figure 2—figure supplement 1. Overexpression of *CBF1*, *-2*, or *-3* causes an increase in *COOLAIR* level, but the effects on *COOLAIR* variants are subtly different.

(A) Positions of qPCR amplicons used for detecting proximal (I) and distal (IIi and IIii) *COOLAIR* variants. The black bars denote exons, thin lines denote introns, and thick lines denote UTRs of *FLC*. The thin black arrow below the gene structure indicates the transcriptional start site of *COOLAIR*. The thick black arrows below denote exons of the type I, type IIi, and type IIii *COOLAIR* variants. The positions of qPCR amplicons used for the qPCR analysis have been marked using blue arrows.

(B-D) Transcript levels of *COOLAIR* variants in the wild-type (Ws-2) and *CBF1*-, *-2*-, and *-3*-overexpressing transgenic plants (*CBF1ox*, *CBF2ox*, and *CBF3ox*, respectively*; Seo et al., 2009*). Relative levels of type I (B), type IIi (C), and type IIii (D) *COOLAIR* were normalized

874   to that of *UBC*. Values have been represented as mean ± SEM of 3 biological replicates. Dots

875   represent each data-point. Asterisks indicate a significant difference, as compared to the wild-

876   type (**, $P<0.01$; ***, $P<0.001$; unpaired Student's *t*-test). ns, not significant.

Figure 3. Protein levels of CBFs increase during the vernalization process.

(A) Transcript levels of *CBF1*, *-2*, and *-3* under short-term cold exposure. Wild-type plants were subjected to 0, 1, 3, 6, 12, and 24 h of 4°C cold treatment. Relative levels of *CBF1*, *-2*, and *-3* were normalized to that of *UBC*. Values have been represented as mean ± SEM of 3 biological replicates. Dots represent each data-point. The blue shadings indicate periods under cold period. Significant differences have been marked using different letters (a-c; *P*<0.05; one-

way ANOVA followed by Tukey's post-hoc test).

(B) Transcript levels of *CBF1*, *-2*, and *-3* after 20V and 40V. Wild-type plants were treated with

4°C vernalization under an SD cycle and collected at ZT4. Relative levels of *CBF1*, *-2*, and *-3*

were normalized to that of *UBC*. Values have been represented as mean ± SEM of 3 biological

replicates. Dots indicate each data-point. The blue shadings denote periods under cold period.

Significant differences have been marked using different letters (a-c; *P*<0.05; one-way ANOVA

followed by Tukey's post-hoc test).

(C) Daily rhythms of *CBF1*, *-2*, and *-3* transcript levels in NV or 40V plants. Col-0 plants

grown under an SD cycle were collected every 4 h between ZT0 and ZT24. Relative transcript

levels of *CBF1*, *-2*, and *-3* were normalized to that of *PP2A*. Values have been represented as

mean ± SEM of 3 biological replicates. The grey shadings represent dark periods. Asterisks

indicate a significant difference between NV and 40V at each time-point (*, *P*<0.05; **, *P*<0.01;

***, *P*<0.001; two-way ANOVA followed by Tukey's post-hoc test).

(D) Increase of CBF3 protein level during the vernalization process. The *pCBF3::CBF3-myc*

transgenic plants, subjected to 4°C vernalization, were collected at ZT4 of the indicated time-

point. CBF3 proteins were detected using anti-myc antibodies. Rubisco was considered the

loading control. Numbers below each band indicate relative signal intensity compared to 1V.

The mean values of 3 biological replicates are presented.

Figure 4. *CBF*s are involved in vernalization-induced *COOLAIR* expression.

(A) Expression dynamics of total *COOLAIR* in wild type and *cbfs-1* mutant during vernalization. Relative transcript levels of total *COOLAIR* were normalized to that of *PP2A*. Values have been represented as mean ± SEM of 3 biological replicates. Dots represent each data-point. Asterisks indicate a significant difference between wild-type and *cbfs-1* mutant at each time-point (**, $P<0.01$; ***, $P<0.001$; two-way ANOVA followed by Tukey's post-hoc test).

(B) Transcript level of total *COOLAIR* in wild-type (Col-0) and CBF3-overexpressing transgenic plant (*CBF3ox; pSuper::CBF3-myc*) during vernalization. Relative levels of total *COOLAIR* were normalized to that of *PP2A*. Values have been represented as mean ± SEM of 3 biological replicates. Dots represent each data-point. Asterisks indicate a significant difference between the wild-type and *CBF3* overexpressor (*, $P<0.05$; **, $P<0.01$; ***,

P<0.001; two-way ANOVA followed by Tukey's post-hoc test). Blue shading denotes cold

periods in (A) and (B).

(C) Effect of first frost-mimicking treatment [8 h of freezing (<0°C)] on the level of total

COOLAIR in 20V wild-type and *cbfs-1*. The upper panel shows a schematic of the experimental

procedure. For NV, wild-type and *cbfs-1* were collected at ZT4 after growing at 22°C for 16 d.

For the first frost treatment, 20V wild-type and *cbfs-1* mutant were treated with an additional

8 h of 4°C (20V, blue line) or 8 h of -1°C (20V+8hF, light blue line) under dark, and then the

whole seedlings were collected at ZT22 for analysis. All the plants were grown under an SD

cycle. The grey shadings denote dark periods. Total *COOLAIR* levels in wild-type and *cbfs-1*

treated with NV, 20V, and 20V+8hF have been represented as mean ± SEM of 3 biological

replicates in the lower panel. Dots indicate each data-point. Relative levels of total *COOLAIR*

were normalized to that of *PP2A*. Asterisks indicate a significant difference (***, *P*<0.001;

two-way ANOVA followed by Tukey's post-hoc test).

Figure 5. *COOLAIR* promoter containing CRT/DREs is necessary for CBFs-mediated vernalization induction.

(A) Luciferase assay showing that CBF3 activates *COOLAIR* promoter with wild-type CRT/DRE (DRE) but fails to activate the one with mutant CRT/DRE (DRE^m). A schematic of the reporter constructs used for the luciferase assay is presented in the upper panel. *pFLC::LUC* harbors 1 kb of the promoter, the 5′-UTR, and the first exon of *FLC*. The blue arrow in the *pFLC::LUC* graphic indicates the location of DREc. Wild-type (WT) and mutant (Mut) forms of *pCOOLAIR::LUC* include 1 kb of the *COOLAIR* promoter with the 3′-UTR and last exon of *FLC*. The blue and red arrows mark the positions of the wild-type (DRE) and mutant (DRE^m) forms of DRE1 and DRE2, respectively. Each reporter construct was co-transfected into *A. thaliana* protoplast together with the *35S::CBF3-HA* effector construct. In parallel, *35S-NOS* control plasmid was transfected as a control. The result is shown below. Relative luciferase activities were normalized to that of the *35S-NOS* control. Values have been represented as mean ± SEM of 3 biological replicates. Dots represent each data-point.

(B) Transcript levels of total *COOLAIR* and proximal (I) *COOLAIR* isoform in wild-type and

FLC$_{\Delta COOLAIR}$ mutant before (NV) and after vernalization (20V). *FLC$_{\Delta COOLAIR}$* has a 324 bp

deletion of the *COOLAIR* promoter region containing DRE1 and -2 (*Luo et al., 2019*). The

position of the deleted region is marked in red lines with reversed triangle in the upper graphic.

Black bars denote exons, thin lines denote introns, and thick lines denote UTRs of *FLC*. The

thin black arrow below the gene structure indicates the transcriptional start site of *COOLAIR*.

The thick black arrow below denotes exons of the type I *COOLAIR* variant. The positions of

amplicons used for the qPCR analysis are marked as blue arrows. The qPCR analysis result is

presented in the lower panel. Relative transcript levels were normalized to that of *PP2A*. Values

have been represented as mean ± SEM of 3 biological replicates. Dots represent each data-

point. Asterisks indicate a significant difference, as compared to NV (**, $P<0.01$; ***, $P<0.001$;

unpaired Student's *t*-test). ns, not significant.

Figure 6. Histone modifications on *FLC* chromatin are not affected by *cbfs* mutation.

(A) Schematic representation of *FLC* gene structure showing the positions of qPCR amplicons used for ChIP analysis. The red arrow presents the position of two CRT/DREs in the *COOLAIR* promoter. qPCR amplicons are marked as blue bars labelled P1, P2, P3, and P4.

(B-D) Enrichments of H3K36me3 (B), H3K4me3 (C), and H3K27me3 (D) on the *FLC* locus in NV or 40V wild-type and *cbfs-1* plants. The whole seedlings were collected at ZT4 under an SD cycle. Modified histones were immunoprecipitated with anti-H3K36me3, anti-H3K4me3, or anti-H3K27me3 antibodies. H3K36me3, H3K4me3, and H3K27me3 enrichments of the IP/5% input at P1, P2, P3, and P4 were normalized to those of *ACT7* and *FUS3*. Relative enrichments have been represented as mean ± SEM of 3 biological replicates. Dots represent each data-point. ns, not significant between wild-type and *cbfs-1*, as assessed using two-way ANOVA followed by Tukey's post-hoc test (P>0.05).

Figure 7. *CBF*-mediated *COOLAIR* induction under long-term cold is not absolutely necessary for *FLC* silencing.

(A) Transcript levels of *FLC* mRNA in wild-type and *cbfs-1* during vernalization. Blue shading denotes cold periods. Relative levels were normalized to that of *PP2A*. Values have been represented as mean ± SEM of 3 biological replicates. Dots represent each data-point. ns, not significant between wild-type and *cbfs-1*, as assessed using two-way ANOVA followed by Tukey's post-hoc test (*P*>0.05).

(B) Photographs of NV and 40V wild-type and *cbfs-1*. NV plants were grown at 22°C under

LD and 40V plants were grown at 22°C under LD after being treated for 40V under SD. The

photos were taken when the NV plants started to bolt and the 40V plants showed the first open

flower.

(C) Flowering times of NV and vernalized (10V-40V) wild-type and *cbfs-1* plants. Flowering

time was measured in terms of the number of rosette leaves produced when bolting. Plants

were grown under LD after vernalization treatment. Bars and error bars represent the mean ±

standard deviation (SD) of 3 biological replicates. Each dot indicates individual flowering time.

Significant differences have been marked using different letters (a-e; two-way ANOVA

followed by Tukey's post-hoc test; $P<0.05$).

(D) Effect of the first frost-mimicking treatment (8 h freezing) on *FLC* mRNA level in wild-

type and *FLC$_{\Delta COOLAIR}$* after 20V. NV, 20V, and 20V+8hF plants were subjected to the treatments

described in *Figure 4C*. Relative levels were normalized to that of *PP2A*. Values have been

represented as mean ± SEM of 3 biological replicates. Dots represent each data-point. Asterisks

indicate a significant difference (***$P<0.001$; two-way ANOVA followed by Tukey's post-hoc

test). ns, not significant.

(E) Transcript levels of *FLC* mRNA in wild-type and *cbfs-1* during vernalization. Blue shading

denotes cold periods. Bars and error bars represent mean ± SEM of 3 biological replicates. Dots

represent each data-point. ns, not significant between wild-type and *FLC$_{\Delta COOLAIR}$*, as assessed

using two-way ANOVA followed by Tukey's post-hoc test ($P>0.05$).

(F) Flowering times of non-vernalized (NV) and vernalized (10V-40V) wild-type and

FLC$_{\Delta COOLAIR}$. Flowering time was measured in terms of the number of rosette leaves produced

when bolting. Plants were grown under LD after vernalization. Bars and error bars indicate

mean ± SD. Each dot represents individual flowering time. Significant differences have been

indicated using different letters (a-e; two-way ANOVA followed by Tukey's post-hoc test;

$P<0.05$).

Figure 8. Schematic model describing the mechanism of CBFs-mediated *COOLAIR* expression during the early phase of vernalization.

During the early phase of vernalization, increased CBF upregulates *COOLAIR* expression by binding to the CRT/DREs at the 3′-end of *FLC*. In the late phase of vernalization, CBF proteins fall off from CRT/DREs in the *COOLAIR* promoter, which leads to a reduction in *COOLAIR* levels.