Title: Self-regulation of the brain's Beta rhythm using a brain-computer interface improves

inhibitory control

Author names: Nadja Enz[1], Jemima Schmidt[1], Kate Nolan[1], Matthew Mitchell[1], Sandra

Alvarez Gomez[1], Miryam Alkayyali[1], Pierce Cambay[1], Magdalena Gippert[1], Robert Whelan*[1,2],

Kathy L. Ruddy*[1]

*Equal contribution

Affiliations: [1]School of Psychology and Institute of Neuroscience, Trinity College Dublin,

Dublin, Ireland, [2]Global Brain Health Institute, Trinity College Dublin, Dublin, Ireland

Corresponding Authors:

Kathy L. Ruddy (ruddykl@tcd.ie), Lloyd Institute, Trinity College Institute of Neuroscience and

School of Psychology, Trinity College Dublin, College Green, Dublin 2, Ireland

Robert Whelan (robert.whelan@tcd.ie), Lloyd Institute, Trinity College Institute of

Neuroscience and School of Psychology, Trinity College Dublin, College Green, Dublin 2,

Ireland

Acknowledgements: The authors would like to thank Emma Wall, Aisling Martin and David

M. Cole for assistance during data collection. Nadja Enz is supported by Irish Research

Council postgraduate scholarship GOIPG/2018/537. Robert Whelan was supported by

Science Foundation Ireland (16/ERCD/3797); European Foundation for Alcohol Research

(ERAB); Brain & Behavior Research Foundation (23599); Health Research Board HRAPOR-

2015-1075. Kathy L. Ruddy would like to acknowledge funding from the Irish Research

Council GOIPD/2017/798 and Health Research Board, Ireland HRB-EIA-2019-003.

Competing interests: The authors declare no competing financial or non-financial interests.

Abstract

Neural oscillations, or brain rhythms, fluctuate in a manner reflecting ongoing behavior. Whether these fluctuations are instrumental or epiphenomenal to the behavior remains elusive. Attempts to experimentally manipulate neural oscillations exogenously using non-invasive brain stimulation have shown some promise, but difficulty with tailoring stimulation parameters to individuals has hindered progress in this field. We demonstrate here using electroencephalography (EEG) neurofeedback in a brain-computer interface that human participants (n=44) learned over multiple sessions across a 6-day period to self-regulate their Beta rhythm (13-20 Hz) over the right inferior frontal cortex (rIFC). This Beta modulation had observable consequences on cognitive behavior: changes in an individual's inhibitory control performance before and after training were predicted by the magnitude of their training-related change in Beta over rIFC. This was not the case for a control group (n=38) who underwent training of their Alpha rhythm (8-12 Hz). The present data support the view that the right frontal Beta rhythm is a key component of the brain's inhibitory control system. Most importantly, we introduce causality to this relationship for the first time, as experimental modulation of rIFC Beta activity had a measurable impact upon behavior.

Introduction

The synchronous firing of large populations of neurons across distributed brain networks produces rhythmic electric field fluctuations large enough to be detected at the scalp. The role – either causal or epiphenomenological – of the observed neural oscillations ('brain rhythms') for human behavior has been a topic of intense debate for decades. Traditionally, researchers have recorded neural oscillations from the scalp while participants perform cognitive tasks, thus investigating the correlation between brain signals and behavior. However, experimental manipulation is necessary in order to specify a causal role for neural oscillations (Herrmann et al., 2016; Vosskuhl et al., 2018).

Exogenous modulation of neural oscillations has previously been achieved using non-invasive brain stimulation techniques like transcranial alternating current stimulation (tACS, see Vosskuhl et al., 2018 for a review), oscillatory transcranial direct current stimulation (o-tDCS; e.g., Marshall et al., 2006), and repetitive transcranial magnetic stimulation (rTMS, e.g., Chung et al., 2015; Thut & Miniussi, 2009) (for reviews, see Dayan et al., 2013; Thut et al., 2011). These methods have generated mixed results with regards to effectiveness of neural modulation (for reviews, see Demirtas-Tatlidede et al., 2013; Enriquez-Geppert et al., 2013; Polanía et al., 2018; Thut et al., 2011) and impact upon cognitive behavior (Bestmann et al., 2015) with a key issue being heterogenous responses to the same stimulation across individuals (Adeyemo et al., 2012; Bergmann & Hartwigsen, 2020; Kasten et al., 2019). Individuals exhibit subtle idiosyncratic features of brain rhythms even within the commonly described bandwidths (Benwell et al., 2019; Haegens et al., 2014), however, methods like rTMS, tACS or o-tDCS typically target specific frequencies.

In addition to neuromodulation methods, brain-computer interface (BCI)-based neurofeedback can be used to endogenously train volitional modulation of brain signals. This approach enables participants to endogenously modulate brain rhythms which are intrinsic to the individual brain (Ros et al., 2010). Neurofeedback has been frequently tested and used as a therapeutic tool and some studies have shown improvements in disorders like attention-

deficit/hyperactivity disorder (ADHD; Jean Arthur Micoulaud-Franchi et al., 2014; Sonuga-Barke et al., 2013). However, there is a lot of heterogeneity in terms of research designs in neurofeedback protocols and it is difficult to draw conclusions regarding the effectiveness of neurofeedback to modify behavior (Omejc et al., 2019; Simon et al., 2021; Sitaram et al., 2017). As well as targeting individually tailored neural oscillation frequencies, it is important to target brain regions that are directly instrumental to the target behavior. For example, electroencephalography (EEG)-neurofeedback from brain signals recorded over sensorimotor areas has shown some evidence of motor-skill improvement in healthy participants as well as clinical motor symptoms in ADHD or stroke patients (Jeunet et al., 2019). In addition, Hsueh et al. (2016) showed that neurofeedback of the frontoparietal Alpha rhythm improved working memory.

Inhibitory control is a core component of healthy executive function, and deficiencies with this aspect of cognition manifest in disorders such as ADHD (e.g., Lijffijt et al., 2005) or addiction (e.g., Luijten et al., 2011). Inhibitory control is believed to rely on fast and flexible control of the brain's Beta rhythm (13-20 Hz; Enz et al., 2021; Jana et al., 2020; Schaum et al., 2021; Swann et al., 2009; Wagner et al., 2017; Wessel, 2020), primarily in a pathway connecting right inferior frontal cortex (rIFC) and basal ganglia (Aron et al., 2014; Wessel & Aron, 2017). The Stop Signal Task (SST) measures this cognitive process (Logan & Cowan, 1984) by requiring the participant to cancel an already initiated motor response following an infrequent Stop cue. The Stop Signal Reaction Time (SSRT) is an estimation of the covert latency of the action cancellation process (Verbruggen et al., 2019). Despite the important role of the Beta rhythm in inhibitory control and the large body of evidence of involvement of rIFC, neurofeedback training of the Beta rhythm in the past has predominantly targeted sensorimotor areas (e.g., Boulay et al., 2011; Vernon et al., 2003; Witte et al., 2013) and no study to date has targeted solely the Beta rhythm over the rIFC.

In order to test if selective self-regulation of specific brain rhythms could modulate specific cognitive processes, we designed a protocol whereby 82 participants learned over a 6-day period to either upregulate or downregulate the amplitude of their Beta or Alpha rhythm

using direct neurofeedback in a BCI. We measured two distinct aspects of cognitive function; speed of proactive response inhibition (conditional SST [cSST]) and working memory (2-Back Task). In a double-blinded mixed design with between-subject (neurofeedback types) and repeated measures (pre-post neurofeedback cognitive measures) factors, we tested the theory that causal manipulation of brain rhythms following neurofeedback training would have an observable impact upon behavior. Specifically, we hypothesized that learning to modulate the Beta rhythm over rIFC would impact positively upon speed of response inhibition, but not upon working memory. By contrast, we predicted that the control group undertaking Alpha neurofeedback would demonstrate no improvement in response inhibition.

    Results

    Spectral power in the Beta band over rIFC was modulated by Beta neurofeedback

    training

We first tested whether Beta and Alpha band spectral power was successfully modulated over rIFC by 6 days of neurofeedback in the trained directions (UP or DOWN), when quantified offline using optimal artefact rejection procedures. Linear mixed effects models were performed on EEG data recorded during BCI performance comparing spectral power at resting baseline on the first Day (Rest S1 Pre) to that at the end of the final (6th) Day (Block 6.4), within the two Beta subgroups. Beta power was significantly modulated from resting baseline on Day 1 to the final block of neurofeedback on Day 6, in a manner that differed depending on trained direction (**Figure 1A**). This was revealed by a Direction*Timepoint interaction ($F[1,36.77]=4.61$, $p=0.039$, $d=0.71$, $n=44$). N.B.: All following post-hoc tests are Bonferroni corrected at $0.05/2=0.025$ and all means are shown as estimated marginal means (EMM) ± standard error. Beta power modestly increased for the UP group (Pre 62.0±3.06 %, Post 63.2±3.24 %; post-hoc test: $t[37.1]=0.41$, $p=0.68$) and significantly decreased for the DOWN group (Pre 59.5±3.68 %, Post 50.9±3.91 %; post-hoc test: $t[37.2]=-2.44$, $p=0.02$). The same pattern was evident when comparing data averaged within the calibration block performed immediately before training on Day 1 (Cal S1) to performance in the final block on Day 6 ($F[1,35.49]=10.24$, $p=0.003$, $d=1.07$, $n=41$; **Figure 1B**). Beta power modestly increased for the UP group (Pre 61.6±2.71 %, Post 63.0±2.82 %; post-hoc test: $t[36.6]=0.56$, $p=0.58$) and significantly decreased for the DOWN group (Pre 64.1±3.52 %, Post 52.8±3.52 %; post-hoc test: $t[34.2]=-3.69$, $p=0.001$). The calibration block data was used to establish the midline of the on-screen display in the neurofeedback game, which participants were required to keep an avatar above (UP) or below (DOWN). In this block, participants rested for two minutes, then conducted a left-hand finger tapping movement for two minutes in order to establish the full range of raw values associated with synchronization and desynchronization of the individual participant's Beta rhythm. **Figure 1A-B** show the time course of Beta power for S1

and S6 compared to resting baseline (**Figure 1A**) and calibration baseline (**Figure 1B**). See Supplementary Results 1 for results of acute within session modulations for Day 1 and Day 6.

Figure 1: Neurofeedback training performance of Beta groups. Performance of first and last session blocks is shown for Beta UP and Beta DOWN groups separately. Time course of relative power is corrected to the respective baseline. The boxplots show the medians and quartiles of the data, the whiskers extend to the rest of the distribution, except for points that are determined to be outliers. The swarm plots show individual datapoints and the line plots connect the means of each block. **A)** Relative Beta power is shown relative to the resting baseline before the first training block on Day 1. **B)** Relative Beta power is shown relative to the calibration baseline before the first training block on Day 1. **C)** Relative Alpha power is shown relative to the resting baseline before the first training block on Day 1. **D)** Relative Alpha power is shown relative to the calibration baseline before the first training block on Day 1.

Spectral power in the Beta band over rIFC was not significantly modulated during Alpha BCI training (all Direction*Timepoint interactions p>0.08; **Figure 2C-D**). However, neither was power in the Alpha band (all Direction*Timepoint interactions p>0.30; **Figure 2A-B**), suggesting that training Alpha over rIFC was not achieved using this protocol.

Figure 2: Neurofeedback training performance of Alpha groups. Performance of first and last session blocks is shown for Alpha UP and Alpha DOWN groups separately. Time course of relative power is corrected to the respective baseline. The boxplots show the medians and quartiles of the data, the whiskers extend to the rest of the distribution, except for points that are determined to be outliers. The swarm plots show individual datapoints and the line plots connect the means of each block. **A)** Relative Alpha power is shown relative to the resting baseline before the first training block on Day 1. **B)** Relative Alpha power is shown relative to the calibration baseline before the first training block on Day 1. **C)** Relative Beta power is shown relative to the resting baseline before the first training block on Day 1. **D)** Relative Beta power is shown relative to the calibration baseline before the first training block on Day 1.

Cross frequency effects and modulation of neural oscillations beyond rIFC

We first investigated whether power in the Alpha band was modulated during neurofeedback targeting the Beta rhythm (**Figure 1C-D**). When comparing baseline Alpha power on Day 1 to Alpha power during neurofeedback attempting to regulate Beta, significant modulation was detected at the end of Day 1 ($F[1,36.71]=4.71$, $p=0.037$, $d=-0.72$, $n=44$), and end of Day 6 ($F[1,34.65]=4.15$, $p=0.049$, $n=44$). For end of Day 1, Alpha power modestly decreased for the UP group (Pre 16.6±1.46 %, Post 15.1±1.48 %; post-hoc test: $t[37.5]=-1.83$, $p=0.075$) and modestly increased for the DOWN group (Pre 18.9±1.76 %, Post 20.4±1.81 %; post-hoc test: $t[38.0]=1.317$, $p=0.20$). For end of Day 6, Alpha power modestly decreased for the UP group (Pre 16.6±1.43 %, Post 15.8±1.50 %; post-hoc test: $t[36.6]=-0.72$, $p=0.47$) and significantly increased for the DOWN group (Pre 18.9±1.72 %, Post 21.9±1.81 %; post-hoc test: $t[36.7]=2.04$, $p=0.048$). It is notable however that although this demonstrates that Alpha was modulated during training based upon neurofeedback of the Beta rhythm, the direction of change was opposite (Alpha decreased in the Beta UP training and vice versa). Also, the absolute effect sizes for Alpha modulation during Beta training range from 0.37-0.72, whereas Beta modulation absolute effect sizes were substantially larger (0.67-1.07). On the final Day of training, Alpha power was not modulated acutely (i.e., within session) during Beta training when comparing power during the final neurofeedback block to the resting baseline on the same day ($F[1,34.85]=3.82$, $p=0.06$, $d=-0.66$, $n=41$), nor to the calibration block ($F[1,35.88]=1.21$, $p=0.28$, $d=-0.37$, $n=40$). Thus, the effects of Beta training were largely selective to the Beta rhythm.

To investigate effects spanning beyond the trained cluster of electrodes over rIFC, we tested whether Beta Power at three other scalp sites was modulated during neurofeedback of Beta signals recorded from rIFC. We chose clusters of four electrodes over right and left motor cortex and occipital cortex for comparison and performed mixed effects models testing for Direction*Timepoint interactions, as before. No significant interactions in any of the three regions were detected when comparing resting baseline Beta power to Beta power during the final neurofeedback block on Day 6 (all $p>0.22$). However, when comparing Beta power from

the initial calibration block on Day 1 to power during the final block on Day 6, significant

Direction*Timepoint interactions were revealed for both right (F[1,35.79]=6.77, p=0.013, d=0.87, n=41) and left (F[1,35.13]=5.64, p=0.023, d=0.80, n=41) motor regions, but no modulation was evident in the occipital region (F[1,35.03]=0.29, p=0.59, d=–0.18, n=41). For the right motor region, Beta power modestly increased for the UP group (Pre 56.7±2.36 %, Post 8.4±2.47 %; post-hoc test: t[36.9]=0.69, p=0.50) and significantly decreased for the DOWN group (Pre 59.4±3.06 %, Post 51.4±3.06 %; post-hoc test: t[34.2]=–2.81, p=0.008). For the left motor region, Beta power modestly increased for the UP group (Pre 57.6±2.30 %, Post 59.4±2.38 %; post-hoc test: t[36.1]=0.96, p=0.35) and significantly decreased for the DOWN group (Pre 57.4±3.00%, Post 52.1±3.00 %; post-hoc test: t[34.1]=–2.29, p=0.028). Topoplots for post-training minus pre-training are shown for each training group for both Beta power (**Figure 3**) and Alpha power (**Figure S1**).

Figure 3: Change in Beta power pre- to post-neurofeedback training. Topoplots show relative Beta power for the last block of neurofeedback training (Post B6.4) minus resting EEG before first block of neurofeedback training (Pre Rest) as well as for the last block of neurofeedback training (Post B6.4) minus calibration block (Pre Calibration). Topoplots are shown separately for each group (Beta UP, Beta DOWN, Alpha UP, Alpha DOWN).

Change in inhibitory control behavior modulated by neurofeedback training

Behavioral data of the cSST are displayed in **Table S1**. To assess whether neurofeedback training had any effect upon inhibitory control behavior in the cSST (i.e., upon SSRT, the speed of inhibitory control), we performed a mixed effects model with three fixed effects; Rhythm (Alpha or Beta), Direction (UP or DOWN) and Timepoint (pre or post training). There was no 3-way Rhythm*Direction*Timepoint interaction ($F_{[1,54.88]}=2.32$, p=0.13, d=0.41, n=71). There was a fixed effect of Timepoint ($F_{[1,54.88]}=4.35$, p=0.042, d=0.28, n=71), revealing that SSRTs generally improved over time regardless of BCI training type (EMMs averaged over levels of Rhythm and Direction: Pre 166±7.86 ms, Post 147±8.54 ms). **Figure 4** shows the mean pre and post SSRTs for each group.

Figure 4: SSRTs of each training group. The SSRTs are shown for each training group for pre (Day 1) and post (Day 6) neurofeedback training. The boxplots show the median and quartiles of the data, the whiskers extend to the rest of the distribution, except for points that are determined to be outliers. The swarm plots show individual datapoints and the line plots connect the means of each block. Abbreviations: SSRT: Stop signal reaction time; cSST: conditional stop signal task.

To investigate whether each individual's pre-post change in SSRT could be predicted

by the extent to which their Beta rhythm was modulated, we performed Robust Regression

analyses with change in SSRT as outcome variable and change in Beta rhythm (resting

baseline from Day 1 to final block on Day 6) as predictor. The extent of change in the Beta

rhythm as a result of training significantly predicted improvement in SSRT from Pre-Post

(slope=1.46, df=23, F=6.48, p=0.018; **Figure 5A**). Specifically, participants who

desynchronized the Beta rhythm over rIFC most, achieved faster SSRTs from pre-post

training. The same relationship was evident when tested at the right motor electrode cluster

(slope=1.32, df=23, F=5.20, p=0.032, n=25; **Figure 5B**), and left motor cluster (slope=1.58,

df=23, F=4.73, p=0.040, n=25; **Figure 5C**), but not at the occipital cluster (slope=1.15, df=23,

F=0.03, p=0.86, n=25; **Figure 5D**). Training related change in Alpha power for those training

Alpha rhythms was not predictive of behavioral change in SSRT (all p>0.42; **Figure 5E-H**).

We additionally tested whether neurofeedback training impacted other aspects of

cognitive function, including working memory (2-Back Task), processing speed (Go RT; Go

reaction times in the cSST) and performance variability (ICV; intra-coefficient of variation in

the cSST). No 3-way interactions emerged between Rhythm*Direction*Timepoint (all p>0.41),

but for Go RT there was a significant 2-way Rhythm*Timepoint interaction (F[1,75.34]=4.82,

p=0.031, d=0.47, n=82). Post-hoc tests are indicating a general improvement in speed

predominantly in the Beta group, revealing that Go RT significantly decreased for the Beta

group (Pre 484±7.07 ms, Post 463±7.31 ms; post-hoc test: t[5.40]=3.44, p=0.001) but did not

significantly decrease for the Alpha group (Pre 480±7.65 ms, Post 478±7.79 ms; post-hoc test:

t[74.4]=0.24, p=0.82).

Figure 5: Association between change in SSRT and change in relative power. Each plot shows the linear regression fit line of the data in dark grey as well as the confidence interval in light grey. Please note the difference in scale across all plots. The associations are shown for an average of four electrodes in four different brain regions (rIFC, right motor, left motor, occipital). **A-D)** show the association between change in SSRT and change in relative Beta power in the Beta groups (UP and DOWN). **E-H)** show the association between change in SSRT and change in relative Alpha power in the Alpha groups (UP and DOWN). Abbreviations: SSRT: stop signal reaction time; rIFC: right inferior frontal cortex.

Resting spectral Beta power was not altered by neurofeedback training

Comparing resting EEG data from before training (Rest S1 Pre), after Day 1 of training (Rest S1 Post) and Post training on Day 6 (Rest S6 Post), mixed effects models revealed no significant Direction*Timepoint interactions for the Beta group (F[2,79.42]=0.34, p=0.71, n=44), suggesting that training related modulations of Beta power were only evident when engaged in the task but did not lead to a lasting change in the background (resting) tonic Beta level.

Modulation of Beta burst characteristics by training the tonic Beta Rhythm

We investigated whether training to modulate tonic Beta Power over rIFC has consequences for Beta burst characteristics. Beta burst rate was not altered during or after training at any of the timepoints tested (all p>0.277). Burst volume was significantly modulated at the end of the first Day of training when comparing burst volume in the last block of Day 1 to that detected in the calibration block on the same Day (Direction*Timepoint interaction: F[1,37.48]=6.14, p=0.02, d=0.81, n=40). Burst volume significantly increased for the UP group (Pre 5962±1744 a.u., Post 9769±1771 a.u.; post-hoc test: t[36.6]=2.20, p=0.03) and modestly decreased for the DOWN group (Pre 8560±2311 a.u., Post 5336±2251 a.u.; post-hoc test: t[37.0]=−1.43, p=0.16).

Brain activity while performing the cSST was not modified following training

There were no significant differences in neural activity (Beta power, burst rate, burst volume, timing of first burst) recorded during cSST performance at the start of the first day of training compared to the end of Day 6 of training (all p>0.40).

Discussion

We have demonstrated here that using neurofeedback in a BCI it is possible to train human participants to self-regulate their Beta rhythm over the rIFC, and that this has observable consequences upon inhibitory control behavior. Participants trained over a 6-day period to upregulate or downregulate the amplitude of their Beta rhythm over rIFC, resulting in the predicted directional changes to Beta power. Concomitant changes at other (untrained) scalp regions and other frequency bands were of lower magnitude, indicating good specificity of the neurofeedback protocol for modulating the trained rhythm, direction and region. The extent to which each individual's SSRT changed pre-post training was predicted by the magnitude of their training-related change in Beta over rIFC. This was not the case for the group undergoing Alpha training, nor were changes in the Alpha rhythm predictive of behavioral change. The present data support the view that the right frontal Beta rhythm is a critical component of the brain's inhibitory control system. Most importantly, we introduce causality to this relationship for the first time, as experimental modulation of rIFC Beta activity had a notable impact upon measured behavior.

The direction of neurofeedback training, upregulating or downregulating the amplitude of the Beta rhythm, had a notable impact upon the observed behavioral change. Specifically, participants exhibiting more training-related desynchronization (downregulation) of Beta, exhibited the largest decreases (improvements) in SSRT. We included sub-groups training with neurofeedback in opposite directions in order to establish whether direction was a relevant factor, as existing literature on this topic suggests both high and low Beta to positively impact upon behavior depending on task and region of interest. Timing and magnitude of high amplitude so called 'Beta bursts' critically impact upon subsequent motor performance (Little et al., 2019) and whether attempts to inhibit a response are successful or not (Enz et al., 2021; Wessel, 2020). Although participants in the Beta UP group of the current study learned to increase the amplitude of their Beta rhythm over rIFC, this gradual, tonic background increase in Beta during the distinct 3-minute neurofeedback blocks had no impact upon subsequent

inhibitory control behavior. Further analyses revealed that this style of upregulation of tonic Beta power had no impact on the rate or volume of Beta bursts, adding weight to the emerging view that Beta bursts are a phenomena distinct from the ongoing background or 'tonic' oscillation at the same frequency (Bonaiuto et al., 2021; Little et al., 2019). In order to modify Beta bursts using the BCI, it may be necessary to provide feedback specifically tailored to detect and influence bursting in real-time, rather than simply of generalized upregulation of tonic Beta (e.g., He, 2020).

In the protocol used in the current study, neurofeedback encouraging downregulation of Beta oscillations revealed the most behaviorally relevant result. It is likely that downregulation is simply easier to achieve, as it is known that engaging a brain region in a mental process (such as motor imagery), tends to lead Beta (and Alpha) to desynchronize in the region (Jensen & Mazaheri, 2010). Over the 6-day training period, participants learned to tailor their mental imagery strategies to optimally engage rIFC in order to achieve tangible real-time control over the movement of the avatar on screen. Consistent with previous literature, our results suggest that it may be the ability to flexibly engage (and remove) Beta oscillations that predict behavioral performance, rather than the tonic level per se. Indeed, this is further supported by the fact that we detected no lasting changes to the Beta rhythm when measured at rest following BCI performance. Training related modulation of the Beta rhythm was only manifest during the neurofeedback task, and while the extent of the modulation did predict subsequent behavioral improvement, it did not alter EEG signals measured at rest or during cSST task performance.

While the neurofeedback training we employed was effective for both up and down-regulating the Beta rhythm over rIFC, Alpha modulation at this scalp location was not achieved, detracting somewhat from the ability to make direct comparisons between behavior following Beta vs Alpha modulation. The lack of Alpha modulation over rIFC may be due to the fact that in this location, Beta is the predominant resonating frequency for neural processing and has been repeatedly implicated in the functioning of this region (Schaum et al., 2021; Sundby et al., 2021; Swann et al., 2009; Swann et al., 2012; Wagner et al., 2017).

Further, for all participants (even those in the Alpha group) we performed the same functional localizer to detect the precise cluster of electrodes corresponding to the right frontal scalp location showing most substantial Beta synchronization during the cSST. This cluster of electrodes, selected for exhibiting strong Beta activity during inhibitory control, was used to drive the BCI neurofeedback for both Alpha and Beta groups. Optimizing the BCI for Beta using this method may further explain why Alpha modulation was not achieved at the rIFC site.

Previous studies using implanted electrodes have reported that positive effects upon motor behavior could be achieved in macaques (Khanna & Carmena, 2017) and humans with Parkinson's disease (Bichsel et al., 2021) using BCI to train self-regulation of the brain's Beta rhythm. Here we build upon and extend these initial findings by making the advance to non-invasive scalp recorded EEG signals in humans, demonstrating that volitional modulation of Beta oscillations was achieved within 6 days of training, and that the extent of the changes predicted subsequent improvements in cognitive performance. The BCI neurofeedback protocol demonstrated good spatial and temporal specificity, modulating primarily the targeted region, rhythm and direction, and cognitive behavioral effects were restricted to inhibitory control as hypothesized due to targeting the 13-20 Hz rhythm at rIFC. Using non-invasive BCI technology, volitional and causal self-regulation was achieved without the need for exogenous stimulation, paving the way for easier real-world application of neuromodulation to alter brain rhythms for improving human performance and treatment of pathologies.

Materials and Methods

Participants

82 healthy adult human volunteers (age: 24.27 ± 7.74 years [mean ± SD]; 44 female; 69 right handed) participated in the study. Participants were reimbursed with either €5/hour and a completion bonus of €60 or with ECTS points. Inclusion criteria were: aged over 18, no history of traumatic brain injury and not currently experiencing any psychiatric disorder (self-reported). All participants provided written informed consent prior to participation. The experimental procedures were approved by the School of Psychology ethics committee of Trinity College Dublin and conducted in accordance with the Declaration of Helsinki.

Study design

Participants were allocated at random into four groups: *'Beta UP'* (n=26), *'Beta DOWN'* (n=18), *'Alpha UP'* (n=23) and *'Alpha DOWN'* (n=15). 'Beta' groups were trained to either increase ('UP') or decrease ('DOWN') their Beta (13-20 Hz) rhythm over the rIFC whereas 'Alpha' groups were trained to either increase ('UP') or decrease ('DOWN') their Alpha (8-12 Hz) rhythm over the same region. Each participant was trained over six sessions (*'S1-6'*). Where possible, sessions were scheduled for a similar time of the day.

The study design is illustrated in **Figure 6**. Each session consisted of one calibration (*'Cal S1-6'*) and four neurofeedback training blocks (*'B1-6.1-4'*) and each block lasted three minutes. Resting EEG was recorded in each session before (*'Rest S1-6 Pre'*) and after (*'Rest S1-6 Post'*) the four neurofeedback blocks. During S1 and S6, participants additionally performed two behavioral tasks, the cSST (experimental task) and the 2-Back Task (control task). The cSST was performed twice in S1, once before (*'cSST S1 Pre'*) and once after (*'cSST S1 Post'*) neurofeedback training whereas the 2-Back Task was only performed once (*'2-Back S1 Pre'*). In S6, both tasks were performed once after neurofeedback training (*'cSST S6 Post', '2-Back S6 Post')*. For all sessions and tasks, the participants were seated comfortably in a chair in front of a computer screen in a soundproof, darkened room.

Figure 6: Study design. Behavioral tasks, baseline EEG measurements and neurofeedback training blocks are shown in sequential order for each of the six sessions. Each session was performed on a separate day. Abbreviations: cSST: conditional stop signal task.

EEG recording

During S1 and S6, 128-channel EEG data in the 10-5 system format were recorded using a 128-channel BioSemi headcap connected to an ActiveTwo Biosemi system. During S2-S5, EEG was recorded from four active Ag/AgCl electrodes over the individual rIFC locus (see *'Functional rIFC localization'*) with the same hardware. A reference electrode recorded data from Cz. For all sessions, three additional electrodes recorded the electrooculogram (right outer canthi for horizontal eye movements and ~2 cm below the left eye for vertical eye movements) as well as the right masseter muscle to detect facial movements.

Resting EEG

To record resting EEG, participants were instructed to fixate upon a black cross for 3 minutes with their eyes open and in a relaxed and stable body position.

Conditional Stop Signal Task (cSST)

The cSST was performed using Presentation® (Version 18.0, Neurobehavioral Systems, Inc., Berkeley, CA, www.neurobs.com) and EEG was recorded during the task. Each trial lasted 1000 ms and was preceded by a fixation cross (1000 ms duration). During Go trials, participants were presented with black arrows pointing either to the right or left (Go signal; 750 ms duration) and they were instructed to respond with their right or left index finger, respectively, as fast as possible via an Xbox 360 game controller. In one of four Go trials, the Go signal was followed by a black arrow pointing upwards (Stop signal; 250 ms duration) after a varying stop-signal delay (SSD). The participants were instructed to inhibit their button press on these Stop trials, but only if the Go signal was pointing in the critical direction. If a Stop signal appeared after a Go signal pointing in the non-critical direction, the participants were instructed to ignore the Stop signal and respond with a button press. The task was divided into four blocks; for the first two blocks the critical direction was right and for the last two blocks the critical direction was left. Each block consisted of 24 trials (18 Go trials and 6 Stop trials). The number of right and left pointing Go signals was equal in each block and presented in a randomized manner. The SSD was adjusted by a tracking algorithm, aiming to achieve a task difficulty resulting in 50% successful and 50% failed Stop trials. After a successful critical Stop trial (ignoring non-critical Stop trials), the SSD was increased, making the task harder and after a failed critical Stop trial (ignoring non-critical Stop trials), the SSD was decreased, making the task easier. The initial SSD was 250 ms and was subsequently adjusted using a double-limit algorithm (see Richards et al., 1999). The SSD could vary between 50 ms and 450 ms. Following a Stop trial, the subsequent SSD value was chosen randomly between the current SSD and a pair of limits (higher or lower, as appropriate). These limits were designed to converge on the SSD that produced a 50% success rate and to be robust to fluctuations on individual trials. If a participant responded to the Go signal before Stop signal presentation, then the SSD was decreased for subsequent trials. All participants completed one block of fifteen practice trials where they received feedback before the real task.

2-Back Task

The 2-Back Task was programmed in Presentation® and no EEG was recorded. Participants were serially presented with randomized letters (500 ms duration each) with an interstimulus interval of 1000 ms. Participants were instructed to press the keyboard button number 1 if the current letter matched the letter presented 2 letters ago. Each block contained 10 targets (i.e., matches) with the target frequency balanced across the block. The task consisted of two blocks and each block contained 50 letters.

Functional rIFC localization

After completion of the *cSST S1 Pre*, the raw EEG data recorded during the task was subjected to an immediate analysis in order to localize the rIFC region on the participant's topography. EEG data from the cSST were epoched into 2500 ms epochs with respect to the Go/Stop signal for Go trials and Stop trials, respectively. A short version of the Fully Automated Statistical Thresholding for EEG artefact Rejection plug-in (FASTER; Nolan et al., 2010; https://sourceforge.net/projects/faster/) was used to identify the most significant artefacts (e.g., eyeblinks and idiosyncratic muscle movements). EEG data were filtered at 13-20 Hz (Beta rhythm) and the amplitude was squared to obtain power estimates. Preprocessed EEG data were epoched into 100 ms to 300 ms after the Stop signal for successful and failed Stop trials. Additionally, a baseline (resting EEG data) was extracted from 1800 ms to 2000 ms after the Stop signal. These data were averaged over all trials for each channel and illustrated using topoplots. Four topoplots were shown; successful Stop trials, failed Stop trials, successful minus failed Stop trials and successful Stop trials minus baseline. The topoplots were then visually inspected and four right frontal electrodes showing the largest right frontal power increase in the Beta range were identified as the individual participant's rIFC. This four-electrode cluster was used subsequently to provide signals for the neurofeedback.

Calibration and neurofeedback blocks (B1-4)

The OpenViBE Acquisition Server (OpenViBE, Renard et al., 2010, www.openvibe.inria.fr) received the EEG stream, and data were processed in real time using a custom OpenViBE Designer script. Data were processed using a 100-ms sliding window. First, the four individual rIFC channels as well as the reference channel (Cz) were selected. The selected data was then spatially (averaged over four rIFC channels) and temporally (13-20 Hz for 'Beta' groups, 8-12 Hz for 'Alpha' groups) filtered. The data were then epoched into 100 ms windows and a power estimate of each 100 ms epoch was calculated by squaring the amplitude. Using LabStreamingLayer (https://github.com/sccn/labstreaminglayer), the resulting power estimates were then exported to MATLAB (R2017b, Mathworks, USA). Using a custom MATLAB script, the power estimates were visualized using the Psychophysics Toolbox (PsychToolbox; Brainard, 1997; http://psychtoolbox.org).

Calibration

During the calibration block, the participants were instructed to first rest and fixate upon a red cross for 2 minutes and when the fixation cross turned green after 2 minutes, they were instructed to open and close their left hand for another 2 minutes. EEG was measured during the whole duration of the calibration block. This calibration procedure was performed in order to measure the participant's full power range of the respective frequency band (Beta or Alpha). The median power of the entire block was calculated.

Neurofeedback training (B1-4)

During the four neurofeedback blocks, an avatar (bird for UP groups, fish for DOWN groups) was visualized on the screen and represented the participant's real-time power estimate output from OpenViBE. The avatar moved horizontally from left to right of the screen and either upwards or downwards depending on whether the power estimates increased or decreased, respectively. The screen was separated horizontally by the median power from the calibration block into sky (above median) and sea (below median) (**Figure S2**). The top of the screen was equal to the maximum power of the calibration block and the bottom of the screen was

equal to the minimum power of the calibration block. The UP groups were instructed to keep the bird in the sky (i.e., increase the power estimates) whereas the DOWN groups were instructed to keep the fish in the sea (i.e., decrease the power estimates). If the avatar deviated into the wrong environment (i.e., sky or sea), the environment turned red to give negative feedback and immediately turned back to normal when they returned into the desired zone. When participants were frequently reaching the top (UP groups) or bottom (DOWN groups) of the screen, the minimum and maximum limits were expanded by the X*standard deviation of the power estimates of the calibration block. The game thus had 4 difficulty levels (X=0-4) and the level was increased if the participant was able to stay in the correct area (sky or sea) for more than 95% in a block. The participants were instructed to develop a mental strategy that does not involve movements, clenching teeth or tensing muscles. Participants were instructed to not close their eyes and to fixate upon the screen at all times.

Data offline processing

EEG offline preprocessing

EEG data were digitized with a sampling rate of 512 Hz. EEG data preprocessing was carried out using the EEGLAB toolbox (Delorme and Makeig, 2004; http://sccn.ucsd.edu/eeglab) in conjunction with FASTER. The data were initially bandpass filtered between 1 Hz and 95 Hz, notch filtered at 50 Hz and average referenced across all scalp electrodes. Resting data and data from the calibration and neurofeedback blocks were subsequently epoched into windows of 1000 ms. Data from the cSST were epoched from 500 ms prior to Go/Stop signal onset to 2000 ms after Go/Stop signal onset for Go trials and Stop trials, respectively. FASTER identified and removed artefactual (i.e., non-neural) independent components, removed epochs containing large artefacts (e.g., muscle twitches) and interpolated channels with poor signal quality. The remaining EEG data were then visually inspected by trained raters to ensure good quality and that any remaining noisy data were removed. Specifically, trained raters identified any remaining artefacts in independent components (e.g., eyeblinks) and

epochs containing idiosyncratic muscle/movement or transient electrode artifacts, and interpolated any channels that were noisy throughout all epochs of a participant. After preprocessing, EEG data were transformed using the current source density method (CSD; https://psychophysiology.cpmc.columbia.edu/software/CSDtoolbox/index.html; Kayser and Tenke, 2006) which is a reference-free montage to attenuate the effect of volume conduction in scalp EEG.

Time-frequency transformation

For all epochs, 2-dimensional representations of each electrode's time-frequency were estimated using a complex Morlet wavelet (range of logarithmically spaced 4-10 cycles for 39 linearly spaced frequencies across 1-40 Hz). The squared magnitude of the convolved data was calculated to obtain power estimates. The power estimates were subsequently transformed to relative power. Power values of each given band from 1-28 Hz (Delta = 1-4 Hz, Theta = 5-7 Hz, Alpha = 8-12 Hz, Beta = 13-28 Hz) were expressed as a percentage of the total power within the spectrum (per channel and per given epoch). Beta bursts were extracted from non-relative time-frequency power estimates.

Beta burst detection

Beta burst detection was performed according to the method described in Enz et al. (2021). For each time-frequency power matrix, local maxima were detected using the MATLAB function *imregionalmax*. Beta bursts were then defined as local maxima that exceeded a defined threshold of 2x median power of the entire time-frequency matrix (across all trials per participant). Time-frequency matrices were then divided into ~25.39 ms time bins (also for analysis of relative power). The first and last time bins were removed from all trials due to an edge artefact that can occur when applying the MATLAB function *imregionalmax* (it detects artefactual local maxima on the edges of the time-frequency matrix). Beta burst rate (the sum of the number of supra-threshold bursts) and Beta burst volume (the area under the curve of

 supra-threshold datapoints; see Enz et al., 2021) were then extracted per time bin. We also

512 extracted the timing of the first Beta burst after the Stop/Go signal in the cSST EEG data.

Selection of brain regions

For the statistical analysis, EEG data were averaged over clusters of four electrodes from

different regions of the brain. To interrogate EEG data from the rIFC, data were averaged over

the four individually selected electrodes over the right frontal scalp area that were used during

the neurofeedback training. Further, we also averaged clusters of four electrodes over the left

motor cortex (D19/C3, D20, D12, D11), the right motor cortex (B22/C4, B23, B31, B30) and

the occipital cortex (A23/Oz, A24, A28, A27) to test the specificity or generalization of effects

beyond the trained region.

Statistical analysis

Behavioral analysis

Means and standard deviations were extracted for each participant for the following behavioral

cSST measures: SSRT, intraindividual coefficient of variation (ICV), Go trial reaction time

(RT), failed Stop trial RT, SSD, number of successful Stop trials, number of failed Stop trials,

probability of successful stopping, probability of Go omissions, probability of choice errors.

The SSRT was calculated using the integration method with replacement of Go omissions by

the maximum RT (Verbruggen et al., 2019). All Go trials were included in the Go RT

distribution, including Go trials with choice errors. Premature responses on failed Stop trials

were included when calculating the probability of responding on a Stop trial and mean SSD.

Participants with SSRT < 75 ms were excluded from all analyses. The ICV was calculated by

dividing Go RT standard deviation by the mean Go RT.

For the 2-Back Task, the absolute number of target hits were calculated across both

blocks (i.e., the maximum absolute number of target hits was 20).

Neurofeedback training statistical analysis

For resting EEG and for the calibration and neurofeedback blocks, data were averaged over 37 x 25.39 ms time bins. For the cSST, 3 x 25.39 ms time bins were averaged to create ~75 ms time bins (6 time bins from -75 ms to 375 ms with respect to the Stop/Go signal). R (R Core Team, 2020) was used for all statistical analyses.

We first tested whether the neurofeedback training was effective. For this, we fit a linear mixed-effects model (LMM) using restricted maximum likelihood with relative Beta power over the rIFC as the outcome variable, with fixed effects of *Direction* (UP or DOWN) and *Timepoint* (Pre to Post neurofeedback training) and their two-way interaction, and with a random effect of *Participant*. We also calculated Cohen's D for each effect. The models were fit for the Beta and Alpha groups separately. We then also conducted the same analysis with relative Alpha power as outcome variable. We also looked at the same outcome variables from the other three brain regions (left motor cortex, right motor cortex, occipital cortex). This analysis was repeated with Beta burst rate and Beta burst volume as outcome variables. We ran a post-hoc test for significant interactions using the *emmeans* function in R. All post-hoc tests are Bonferroni corrected at 0.05/2=0.025, correcting for the two directions (UP and DOWN).

Next, we looked at the effects of neurofeedback training on inhibitory control behavior. We again fit a LMM using restricted maximum likelihood with SSRT as the outcome variable, with fixed effects of *Direction* (UP or DOWN), *Timepoint* (Pre to Post neurofeedback training), *Rhythm* (Beta or Alpha) and their two- and three-way interactions, and with a random effect of *Participants*. We also calculated Cohen's D for each effect.

We then interrogated the relationship between the magnitude of the Pre-Post change in SSRT and the extent to which relative Beta power was modulated during neurofeedback training. For this we fit a linear model by robust regression using an M-estimator, with change in SSRT being the dependent variable and change in relative Beta power being the independent variable. We repeated this analysis with N-Back score, Go RT and ICV as outcome variables.

Next, we looked at the effects of neurofeedback training on resting EEG data collected

before and after. We fit the same three-way LMM with relative Beta power, Beta burst rate

and Beta burst volume over the rIFC as outcome variables.

Last, we looked at the effect of neurofeedback training on the brain activity while

engaging inhibitory control behavior. Again, the same three-way LMM was fit for the time bins

around the average SSRT with relative Beta power, Beta burst rate, Beta burst volume and

timing of first Beta burst over the rIFC as outcome variables.

Code and data accessibility

571   Custom written scripts and data summary files can be downloaded on the Open Science

572   Framework at [URL to be inserted after acceptance].

References

Adeyemo, B. O., Simis, M., Macea, D. D., & Fregni, F. (2012). Systematic review of parameters of stimulation, clinical trial design characteristics, and motor outcomes in non-invasive brain stimulation in stroke. *Frontiers in Psychiatry*, *3*(November), 1–27. https://doi.org/10.3389/fpsyt.2012.00088

Aron, A. R., Robbins, T. W., & Poldrack, R. A. (2014). Inhibition and the right inferior frontal cortex: One decade on. *Trends in Cognitive Sciences*, *18*(4), 177–185. https://doi.org/10.1016/j.tics.2013.12.003

Benwell, C. S. Y., London, R. E., Tagliabue, C. F., Veniero, D., Gross, J., Keitel, C., & Thut, G. (2019). Frequency and power of human alpha oscillations drift systematically with time-on-task. *NeuroImage*, *192*, 101–114. https://doi.org/10.1016/j.neuroimage.2019.02.067

Bergmann, T. O., & Hartwigsen, G. (2020). Inferring causality from noninvasive brain stimulation in cognitive neuroscience. *Journal of Cognitive Neuroscience*, *33*(2), 195–225. https://doi.org/10.1162/jocn_a_01591

Bestmann, S., de Berker, A. O., & Bonaiuto, J. (2015). Understanding the behavioural consequences of noninvasive brain stimulation. *Trends in Cognitive Sciences*, *19*(1), 13–20. https://doi.org/10.1016/j.tics.2014.10.003

Bichsel, O., Stieglitz, L. H., Oertel, M. F., Baumann, C. R., Gassert, R., & Imbach, L. L. (2021). Deep brain electrical neurofeedback allows Parkinson patients to control pathological oscillations and quicken movements. *Scientific Reports*, *11*, 7973. https://doi.org/10.1038/s41598-021-87031-2

Bonaiuto, J. J., Little, S., Neymotin, S. A., Jones, S. R., Barnes, G. R., & Bestmann, S. (2021). Laminar dynamics of beta bursts in human motor cortex. *BioRxiv*, 431412. Retrieved from https://doi.org/10.1101/2021.02.16.431412

Boulay, C. B., Sarnacki, W. A., Wolpaw, J. R., & McFarland, D. J. (2011). Trained modulation of sensorimotor rhythms can affect reaction time. *Clinical Neurophysiology*,

122(9), 1820–1826. https://doi.org/10.1016/j.clinph.2011.02.016

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*, 433–436.

https://doi.org/10.1163/156856897X00357

Chung, S. W., Rogasch, N. C., Hoy, K. E., & Fitzgerald, P. B. (2015). Measuring brain

stimulation induced changes in cortical properties using TMS-EEG. *Brain Stimulation*,

8(6), 1010–1020. https://doi.org/10.1016/j.brs.2015.07.029

Dayan, E., Censor, N., Buch, E. R., Sandrini, M., & Cohen, L. G. (2013). Noninvasive brain

stimulation: From physiology to network dynamics and back. *Nature Neuroscience*,

16(7), 838–844. https://doi.org/10.1038/nn.3422

Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-

trial EEG dynamics including independent component analysis. *Journal of*

Neuroscience Methods, *134*(1), 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Demirtas-Tatlidede, A., Vahabzadeh-Hagh, A. M., & Pascual-Leone, A. (2013). Can

noninvasive brain stimulation enhance cognition in neuropsychiatric disorders?

Neuropharmacology, *64*, 566–578. https://doi.org/10.1016/j.neuropharm.2012.06.020

Enriquez-Geppert, S., Huster, R. J., & Herrmann, C. S. (2013). Boosting brain functions:

Improving executive functions with behavioral training, neurostimulation, and

neurofeedback. *International Journal of Psychophysiology*, *88*(1), 1–16.

https://doi.org/10.1016/j.ijpsycho.2013.02.001

Enz, N., Ruddy, K. L., Rueda-Delgado, L. M., & Whelan, R. (2021). Volume of β-Bursts, But

Not Their Rate, Predicts Successful Response Inhibition. *The Journal of Neuroscience*,

41(23), 5069–5079. https://doi.org/10.1523/jneurosci.2231-20.2021

Haegens, S., Cousijn, H., Wallis, G., Harrison, P. J., & Nobre, A. C. (2014). Inter- and intra-

individual variability in alpha peak frequency. *NeuroImage*, *92*, 46–55.

https://doi.org/10.1016/j.neuroimage.2014.01.049

He, S. (2020). Neurofeedback-linked suppression of cortical beta bursts speeds up

movement initiation in healthy motor control : A double-blind sham- controlled study.

Journal of Neuroscience. https://doi.org/https://doi.org/10.1523/JNEUROSCI.0208-

20.2020

Herrmann, C. S., Strüber, D., Helfrich, R. F., & Engel, A. K. (2016). EEG oscillations: From

correlation to causality. *International Journal of Psychophysiology*, *103*, 12–21.

https://doi.org/10.1016/j.ijpsycho.2015.02.003

Hsueh, J. J., Chen, T. S., Chen, J. J., & Shaw, F. Z. (2016). Neurofeedback training of EEG

alpha rhythm enhances episodic and working memory. *Human Brain Mapping*, *37*(7),

2662–2675. https://doi.org/10.1002/hbm.23201

Jana, S., Hannah, R., Muralidharan, V., & Aron, A. R. (2020). Temporal cascade of frontal,

motor and muscle processes underlying human action-stopping. *ELife*, *9*, e50371.

https://doi.org/10.7554/elife.50371

Jensen, O., & Mazaheri, A. (2010). Shaping functional architecture by oscillatory alpha

activity: Gating by inhibition. *Frontiers in Human Neuroscience*, *4*(Nov), 1–8.

https://doi.org/10.3389/fnhum.2010.00186

Jeunet, C., Glize, B., McGonigal, A., Batail, J. M., & Micoulaud-Franchi, J. A. (2019). Using

EEG-based brain computer interface and neurofeedback targeting sensorimotor

rhythms to improve motor skills: Theoretical background, applications and prospects.

Neurophysiologie Clinique, *49*(2), 125–136. https://doi.org/10.1016/j.neucli.2018.10.068

Kasten, F. H., Duecker, K., Maack, M. C., Meiser, A., & Herrmann, C. S. (2019). Integrating

electric field modeling and neuroimaging to explain inter-individual variability of tACS

effects. *Nature Communications*, *10*, 5427. https://doi.org/10.1038/s41467-019-13417-6

Kayser, J., & Tenke, C. E. (2006). Principal components analysis of Laplacian waveforms as

a generic method for identifying ERP generator patterns: I. Evaluation with auditory

oddball tasks. *Clinical Neurophysiology*, *117*, 348–368.

https://doi.org/10.1016/j.clinph.2005.08.034

Khanna, P., & Carmena, J. M. (2017). Beta band oscillations in motor cortex reflect neural

population signals that delay movement onset. *ELife*, *6*, e24573.

https://doi.org/10.7554/eLife.24573

Lijffijt, M., Kenemans, J. L., Verbaten, M. N., & Van Engeland, H. (2005). A meta-analytic

review of stopping performance in attention-deficit/ hyperactivity disorder: Deficient

inhibitory motor control? *Journal of Abnormal Psychology*, *114*(2), 216–222.

https://doi.org/10.1037/0021-843X.114.2.216

Little, S., Bonaiuto, J., Barnes, G., & Bestmann, S. (2019). Human motor cortical beta bursts

relate to movement planning and response errors. *PLoS Biology*, *17*(10), e3000479.

https://doi.org/10.1371/journal.pbio.3000479

Logan, G. D., & Cowan, W. B. (1984). On the ability to inhibit thought and action: A theory of

an act of control. *Psychological Review*, *91*(3), 295–327. https://doi.org/10.1037/0033-

295X.91.3.295

Luijten, M., Littel, M., & Franken, I. H. A. (2011). Deficits in inhibitory control in smokers

during a Go/Nogo task: An investigation using event-related brain potentials. *PLoS

ONE*, *6*(4), e18898. https://doi.org/10.1371/journal.pone.0018898

Marshall, L., Helgadóttir, H., Mölle, M., & Born, J. (2006). Boosting slow oscillations during

sleep potentiates memory. *Nature*, *444*, 610–613. https://doi.org/10.1038/nature05278

Micoulaud-Franchi, J. A., Geoffroy, P. A., Fond, G., Lopez, R., Bioulac, S., & Philip, P.

(2014). EEG neurofeedback treatments in children with ADHD: An updated meta-

analysis of randomized controlled trials. *Frontiers in Human Neuroscience*,

8(November), 1–7. https://doi.org/10.3389/fnhum.2014.00906

Nolan, H., Whelan, R., & Reilly, R. B. (2010). FASTER: Fully Automated Statistical

Thresholding for EEG artifact Rejection. *Journal of Neuroscience Methods*, *192*(1),

152–162. https://doi.org/10.1016/j.jneumeth.2010.07.015

Omejc, N., Rojc, B., Battaglini, P. P., & Marusic, U. (2019). Review of the therapeutic

neurofeedback method using electroencephalography: EEG neurofeedback. *Bosnian

Journal of Basic Medical Sciences*, *19*(3), 213–220.

https://doi.org/10.17305/bjbms.2018.3785

Polanía, R., Nitsche, M. A., & Ruff, C. C. (2018). Studying and modifying brain function with

non-invasive brain stimulation. *Nature Neuroscience*, *21*, 174–187.

https://doi.org/10.1038/s41593-017-0054-4

R Core Team. (2020). R: A language and environment for statistical computing. *R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria*.

Renard, Y., Lotte, F., Gibert, G., Congedo, M., Maby, E., Delannoy, V., et al. (2010). OpenViBE: An open-source software platform to design, test, and use brain-computer interfaces in real and virtual environments. *Presence: Teleoperators and Virtual Environments*. https://doi.org/10.1162/pres.19.1.35

Richards, J. B., Zhang, L., Mitchell, S. H., & de Wit, H. (1999). Delay or probability discounting in a model of impulsive behavior: effect of alcohol. *Journal of the Experimental Analysis of Behavior*, *71*(2), 121–143. https://doi.org/10.1901/jeab.1999.71-121

Ros, T., Munneke, M. A. M., Ruge, D., Gruzelier, J. H., & Rothwell, J. C. (2010). Endogenous control of waking brain rhythms induces neuroplasticity in humans. *European Journal of Neuroscience*, *31*(4), 770–778. https://doi.org/10.1111/j.1460-9568.2010.07100.x

Schaum, M., Pinzuti, E., Sebastian, A., Lieb, K., Fries, P., Mobascher, A., et al. (2021). Right inferior frontal gyrus implements motor inhibitory control via beta-band oscillations in humans. *ELife*, *10*, e61679. https://doi.org/10.7554/eLife.61679

Simon, C., Bolton, D. A. E., Kennedy, N. C., Soekadar, S. R., & Ruddy, K. L. (2021). Challenges and opportunities for the future of Brain-Computer Interface in neurorehabilitation. *Frontiers in Neuroscience*, *15*(July). https://doi.org/https://doi.org/10.3389/fnins.2021.699428

Sitaram, R., Ros, T., Stoeckel, L., Haller, S., Scharnowski, F., Lewis-Peacock, J., et al. (2017). Closed-loop brain training: The science of neurofeedback. *Nature Reviews Neuroscience*, *18*(2), 86–100. https://doi.org/10.1038/nrn.2016.164

Sonuga-Barke, E. J. S., Brandeis, D., Cortese, S., Daley, D., Ferrin, M., Holtmann, M., et al. (2013). Nonpharmacological Interventions for ADHD: Systematic Review and Meta-Analyses of Randomized Controlled Trials of Dietary and Psychological Treatments.

American Journal of Psychiatry, *170*, 275–289.

https://doi.org/https://doi.org/10.1176/appi.ajp.2012.12070991

Sundby, K. K., Jana, S., & Aron, A. R. (2021). Double-blind disruption of right inferior frontal cortex with TMS reduces right frontal beta power for action stopping. *Journal of Neurophysiology*, *125*(1), 140–153. https://doi.org/10.1152/JN.00459.2020

Swann, N., Tandon, N., Canolty, R., Ellmore, T. M., McEvoy, L. K., Dreyer, S., et al. (2009). Intracranial EEG Reveals a Time- and Frequency-Specific Role for the Right Inferior Frontal Gyrus and Primary Motor Cortex in Stopping Initiated Responses. *Journal of Neuroscience*, *29*(40), 12675–12685. https://doi.org/10.1523/JNEUROSCI.3359-09.2009

Swann, N. C., Cai, W., Conner, C. R., Pieters, T. A., Claffey, M. P., George, J. S., et al. (2012). Roles for the pre-supplementary motor area and the right inferior frontal gyrus in stopping action: Electrophysiological responses and functional and structural connectivity. *NeuroImage*, *59*(3), 2860–2870. https://doi.org/10.1016/j.neuroimage.2011.09.049

Thut, G., & Miniussi, C. (2009). New insights into rhythmic brain activity from TMS-EEG studies. *Trends in Cognitive Sciences*, *13*(4), 182–189. https://doi.org/10.1016/j.tics.2009.01.004

Thut, G., Schyns, P. G., & Gross, J. (2011). Entrainment of perceptually relevant brain oscillations by non-invasive rhythmic stimulation of the human brain. *Frontiers in Psychology*, *2*(July), 1–10. https://doi.org/10.3389/fpsyg.2011.00170

Verbruggen, F., Aron, A., Band, G., Beste, C., Bissett, P., Brockett, A. T., et al. (2019). Capturing the ability to inhibit actions and impulsive behaviors: A consensus guide to the stop-signal task. *ELife*, *8*, e46323. https://doi.org/10.31219/OSF.IO/8MZDU

Vernon, D., Egner, T., Cooper, N., Compton, T., Neilands, C., Sheri, A., & Gruzelier, J. (2003). The effect of training distinct neurofeedback protocols on aspects of cognitive performance. *International Journal of Psychophysiology*, *47*(1), 75–85. https://doi.org/10.1016/S0167-8760(02)00091-0

Vosskuhl, J., Strüber, D., & Herrmann, C. S. (2018). Non-invasive Brain Stimulation: A

　　　Paradigm Shift in Understanding Brain Oscillations. *Frontiers in Human Neuroscience*,

　　　12(May), 1–19. https://doi.org/10.3389/fnhum.2018.00211

Wagner, J., Wessel, J. R., Ghahremani, A., & Aron, A. R. (2017). Establishing a Right

　　　Frontal Beta Signature for Stopping Action in Scalp EEG: Implications for Testing

　　　Inhibitory Control in Other Task Contexts. *Journal of Cognitive Neuroscience*, *30*(1),

　　　107–118. https://doi.org/10.1162/jocn

Wessel, J. R. (2020). β-Bursts Reveal the Trial-to-Trial Dynamics of Movement Initiation and

　　　Cancellation. *The Journal of Neuroscience*, *40*(2), 411–423.

　　　https://doi.org/10.1523/JNEUROSCI.1887-19.2019

Wessel, J. R., & Aron, A. R. (2017). On the Globality of Motor Suppression: Unexpected

　　　Events and Their Influence on Behavior and Cognition. *Neuron*, *93*(2), 259–280.

　　　https://doi.org/10.1016/j.neuron.2016.12.013

Witte, M., Kober, S. E., Ninaus, M., Neuper, C., & Wood, G. (2013). Control beliefs can

　　　predict the ability to up-regulate sensorimotor rhythm during neurofeedback training.

　　　Frontiers in Human Neuroscience, *7*(July), 1–8.

　　　https://doi.org/10.3389/fnhum.2013.00478