1   **Fluorescence exclusion: a rapid, accurate and powerful method for measuring yeast cell volume**
2

3   Daniel García-Ruano[1], Akanksha Jain[1], Joseph C. Ryan[1], Vasanthakrishnan Radhakrishnan
4   Balasubramaniam[1], Larisa Venkova[2], Matthieu Piel[2] and Damien Coudreuse[1,*]
5
6
7   [1] Institute of Genetics and Development of Rennes, UMR 6290, CNRS – University of Rennes 1,
8   France
9
10   [2] Institut Curie and Institut Pierre Gilles de Gennes, PSL Research University, CNRS UMR 144,
11   Paris, France.
12
[*] Corresponding author: damien.coudreuse@univ-rennes1.fr

SUMMARY STATEMENT

Fluorescence exclusion provides a unique method to accurately measure the volume of yeast cells at

both the population and single-cell levels.

ABSTRACT

Cells exist in an astonishing range of volumes across and within species. However, our understanding

of cell size control remains limited, due in large part to the challenges associated with accurate

determination of cell volume. Much of our comprehension of size regulation derives from models such

as budding and fission yeast, but even for these morphologically stereotypical cells, assessment of cell

volume has relied on proxies and extrapolations from two-dimensional measurements. Recently, the

fluorescence exclusion method (FXm) was developed to evaluate the size of mammalian cells, but

whether it could be applied to smaller cells remained unknown. Using specifically designed

microfluidic chips and an improved data analysis pipeline, we show here that FXm reliably detects

subtle difference in the volume of fission yeast cells, even for those with altered shapes. Moreover, it

allows for the monitoring of dynamic volume changes at the single-cell level with high time resolution.

Collectively, our work reveals how coupling FXm with yeast genetics will bring new insights into the

complex biology of cell growth.

KEYWORDS

Cell volume, yeast, fluorescence exclusion

RUNNING TITLE

Measuring yeast cell volume

INTRODUCTION

Cell volume is a dynamic and complex trait that is modulated by both genetic and environmental components (Amodeo and Skotheim, 2016; Cook and Tyers, 2007; Lloyd, 2013; Marshall et al., 2012; Mueller, 2015). It has an impact on a host of cellular processes (Pedersen et al., 2001; Zhurinsky et al., 2010), playing important roles in tissue architecture and contributing to the overall size and shape of an organism (Amodeo and Skotheim, 2016; Cook and Tyers, 2007; Lloyd, 2013; Marshall et al., 2012; Mueller, 2015). At the single-cell level, volume and surface area not only define how cells sense external stimuli and interact with their environment but also determine their intracellular chemistry and organization. Strikingly, deregulation of cell size has dramatic consequences and is a hallmark of aging and cancer (Li et al., 2015; Pietras, 2011; Yang et al., 2011).

In proliferating cells, volume is intimately linked to the control of cell cycle progression. The dynamics of cell growth are complex across species, and specific stages of the cell cycle are associated with differential changes in cell size and morphology. For instance, while cells of the fission yeast *Schizosaccharomyces pombe* stop growing and maintain their overall shape during mitosis, mammalian cells undergoing division show a transient phase of rounding and volume alteration (Lancaster et al., 2013; Zlotek-Zlotkiewicz et al., 2015). In addition, size thresholds have been proposed to act as key checkpoints during the fission yeast cell cycle, allowing for the coordination of growth and division (Fantes, 1977; Fantes and Nurse, 1978; Fantes and Nurse, 1977). These mechanisms reduce cell-to-cell heterogeneity, thereby maintaining size homeostasis in a growing population.

Alterations in cell size are also triggered by changes in the environment, such as upon modulation of nutritional conditions, and cell cycle exit is often associated with a decrease in cell volume. For example, yeast cells exposed to nitrogen or glucose starvation enter quiescence with a significantly reduced size (Sun and Gresham, 2021). Interestingly, increases in cell volume were frequently observed during long-term experimental evolution in bacteria and were proposed to contribute to improved proliferation (Grant et al., 2021; Lenski and Travisano, 1994; Mongold and Lenski, 1996). This highlights how the volume of dividing cells may result from the necessary balance between growth advantages and evolutionary trade-offs.

The principles and mechanisms underlying cell size control have been the focus of in-depth investigation. While several pathways have been described to play a role in regulating this critical parameter (Amodeo and Skotheim, 2016; Cook and Tyers, 2007; Lloyd, 2013; Marshall et al., 2012; Mueller, 2015), whether they represent the key mechanisms that ensure size homeostasis remains unclear. Furthermore, how cells measure and control their geometric characteristics is surprisingly poorly understood. As a result, multiple models that are not mutually exclusive have been proposed: cells may divide 1) when they have reached a specific size (sizer), 2) after a specific time following a landmark cell cycle event (timer) and 3) after having produced a fixed amount of mass, at a rate that

depends on birth size (adder). While these possibilities are supported by experimental and theoretical

studies (Fantes and Nurse, 1977; Mueller, 2015; Soifer et al., 2016; Sveiczer et al., 1996; Taheri-Araghi

et al., 2015), general conclusions are difficult to draw. This may be due in part to the diversity of

measurements used to describe cell size, including volume, surface area, cell length or dry mass. Indeed,

the question remains whether one or a combination of these characteristics is more relevant for

understanding size regulation and whether different cell types rely on the same or on distinct parameters.

Thus, while the sizes and shapes of individual cell types are commonly accepted as fundamental

hallmarks of cell identity, no unifying principles for size control have emerged.

Experimentally, the regulation of cell size has been difficult to investigate. In mammalian cell

lines, the challenges of advanced genetic manipulation and the lack of simple methods for accurate

measurement of parameters such as volume and surface area have been an obstacle to studying the

molecular mechanisms of size control. Therefore, unicellular eukaryotes such as the budding and fission

yeasts remain the models of choice for deciphering the bases of this critical cellular feature. In these

organisms, regulation of cell size is intimately linked to the operation of the cell cycle machinery.

Different concepts have thus emerged that may provide robust ways of regulating size, including

titration mechanisms and dependence on geometric parameters (*e.g.* length, surface area) (Amodeo and

Skotheim, 2016; Facchetti et al., 2019). However, how cells monitor their dimensions and how variation

from the target size triggers a corrective response are not fully understood.

In the fission yeast *S. pombe*, the existence of cell size checkpoints was initially suggested at both

S phase and, more importantly, mitosis (Fantes, 1977; Fantes and Nurse, 1978; Fantes and Nurse, 1977).

While a mechanism based on an intracellular gradient of a cell cycle regulator was subsequently

proposed (Martin and Berthelot-Grosjean, 2009; Moseley et al., 2009), recent evidence has challenged

this model (Amodeo and Skotheim, 2016; Pan et al., 2014a; Saunders et al., 2012). Moreover, the

commonly-employed approach of using cell length at division as a proxy for size may not be sufficient

for investigating size regulation. First, cell volume is more critical to cell biology and biochemistry than

length. Second, slight changes in the diameter of these rod-shaped cells are usually ignored, although

they have a stronger impact on cell volume than similar alterations in length: the volume of a cylinder

is linearly dependent on its length but varies with the square of its radius. In addition, morphology

mutants or cells grown in conditions that induce alteration of cellular dimensions and scaling are

complex to study and to compare with reference strains using this method. Finally, size at division

restricts the analysis to only a small fraction of a proliferating population.

To investigate the control of yeast cell volume in a more unbiased manner, different techniques

have been applied. These include volume calculation from geometrical assumptions, 2D imaging data

that allow for 3D reconstruction or the integration of cellular sections extrapolated from the cell

outlines, and complex approaches using advanced micromechanical devices (Baybay et al., 2020; Bryan

et al., 2010; Bryan et al., 2014; Facchetti et al., 2019; Model, 2018; Pan et al., 2014b; Zegman et al.,

2015). However, these strategies are either low-throughput, difficult to establish or rely on various
assumptions and complex processing of 2D measurements. For instance, given the impact of cell
diameter on cell volume, accurate evaluation of cell size based on automated recognition of cell outlines
requires this detection to be highly precise (Baybay et al., 2020; Zegman et al., 2015). Therefore, while
these methods represent important steps in our capacity to address the regulation of cell volume, a more
direct and simple methodology that does not introduce the biases of 2D data analysis or multiple steps
of image segmentation and reconstruction is necessary.

Recently, a fluorescence exclusion method (FXm) was developed for evaluating the volume of
morphologically diverse mammalian cells (Cadart et al., 2017). FXm relies on microfluidic devices in
which cells are imaged in the presence of a fluorescently-labelled dextran that does not cross the plasma
membrane. Thus, using chambers of known height, the local reduction in fluorescence due to the
presence of a cell allows for the determination of the volume of this cell with high accuracy (Fig. 1A).
This technique is simple and only requires single-plane imaging with low magnification objectives. It
is unaffected by morphological alterations and is compatible with the monitoring of high numbers of
cells. Although perfectly adapted to large mammalian cells, whether this new approach could be used
for smaller cells such as yeast remained unclear, as this would require microdevices with significantly
lower internal chambers. For example, HeLa cells are around 20-40 μm in height with a volume of
~1500 μm^3 at birth and ~3000 μm^3 at division (Cadart et al., 2018; Cadart et al., 2019). In contrast,
wild-type fission yeast cells divide at a length of ~14 μm, with a diameter of ~4-4.5 μm. Depending on
the growth conditions and exact diameter used for volume calculations, this led to values for size at
division between ~120 and 200 μm^3 (Navarro and Nurse, 2012; Nurse, 1975).

Here we use the fission yeast *Schizosaccharomyces pombe* as a model system and demonstrate that
the fluorescence exclusion method is a unique tool for measuring the volume of small cells. We engineer
specific microdevices for yeast cells, assess the reproducibility and sensitivity of the system, and
provide an improved way to perform large-scale analyses. Furthermore, we demonstrate the
compatibility of the approach with multi-color imaging for evaluating single-cell volume in specific
subpopulations. We then show how time-lapse experiments using these devices is a powerful strategy
to couple yeast genetics with volume measurements for deciphering growth dynamics and size
modulation. Interestingly, our results suggest that the use of FXm to re-evaluate previous models will
be essential for understanding how cell volume is regulated in fission yeast, with implications for the
general principles that govern cell size control in complex eukaryotes.

RESULTS

Microfluidic devices for determining yeast cell volume by FXm

FXm was developed to measure the volume of mammalian cells despite their complex morphologies (Cadart et al., 2017; Cadart et al., 2018). This method is simple and potentially compatible with any cell type, making it an ideal approach for investigating the regulation of cell volume in genetically amenable models such as yeast. However, to ensure an optimal dynamic range for accurate volume measurements, the height of the microfluidic chamber must be of the same order of magnitude as that of the cells. Indeed, the exclusion of fluorescent molecules in a microsystem several times higher than the cell would only marginally impact the total fluorescence intensity in the region of interest, affecting the reliability of the results. Microdevices employed to determine mammalian cell volume are therefore not adapted to small fission yeast cells, which require chambers of ~4-7 μm in height, depending on the strain studied. Interestingly, evaluation of FXm suggests that such reduced heights represent an advantage for the accuracy of these measurements (Model, 2020).

This design constraint led us to build specific and optimized devices for our yeast model. In particular, we determined a density of normalization pillars that is high enough to prevent the collapse of the chambers without significantly reducing the throughput of the method. Indeed, cells in close vicinity of these pillars are excluded from the FXm analyses due to the bias introduced by the presence of neighboring structures (Cadart et al., 2017). For this, we tested chambers with inter-pillar distances of 120, 180 and 270 μm (Fig. 1B). We found a distance of 180 μm between pillars coupled with an optimized cell concentration and preparation procedure to be the most reliable combination for the measurement of high numbers of individual cells in a single frame while limiting cell aggregation (Fig. 1C, see Materials and Methods).

Measuring the volume of individual fission yeast cells using FXm

The necessity of using microchips of reduced height may represent a challenge for precisely measuring the volume of individual yeast cells. Indeed, in these conditions, the difference between the minimum (pillar) and maximum (no cell) fluorescence intensities (Fig. 1A), as well as the extent of fluorescent molecule displacement in the presence of a cell, are inherently low. This may render the technique less sensitive and more susceptible to noise, with consequences for its accuracy and reproducibility.

To evaluate whether FXm provides high quality results for yeast, we measured the size of wild-type fission yeast cells, comparing both the median volumes and volume distributions from independent experiments (see Materials and Methods). To this end, we used a 5.53 μm high chamber design and analyzed five replicates performed in separate devices built from the same master mold. Remarkably,

we obtained highly reproducible data, with a median volume for the pooled datasets of 104.8 ± 24.5
µm³ (standard error for the 5 replicates, s.e.m.: 1.7 µm³), and comparable population profiles (Fig. 2A).
This value for wild-type cells is consistent with previous results established by complex image
processing (Baybay et al., 2020), validating FXm for cell size monitoring in yeast. In contrast to size at
division, which only focuses on a small subpopulation, our method describes the volumes of the cells
over the entire population, from birth to division. It therefore integrates the changes in size that occur
during the division cycle. In addition, our experiments suggest that several repeats may be needed when
evaluating median volume differences of less than 10 %, as such alterations are in the range of the
observed experimental variability (Fig. 2A). When considering a geometric cell model consisting of a
cylinder with half spheres at both ends and a constant diameter of 4 µm, an increase of 10 % in volume
at division is the result of an increase in cell length at division from 14 to ~15 µm. Conversely, a similar
change in cell volume at constant length is brought about by an increase in diameter of ~0.2 µm. In both
cases, these differences are at the limit or below the resolution of the strategies routinely used for
measuring cell length or width in *S. pombe*. This demonstrates the high sensitivity of FXm for yeast
cell volume measurement, even when applying a conservative threshold of 10 % change.

Next, as molds deteriorate when heavily used and given that standard microfabrication procedures
make it difficult to generate highly reproducible chamber heights, we set out to evaluate how changes
in the dimensions of the devices may impact volume determination. We therefore produced microchips
of various heights and measured the size of wild-type fission yeast cells by FXm. Again, we observed
remarkably similar median volumes and volume distributions between measurements (Fig. 2B – median
volume of the pooled datasets: 106.8 ± 24.4 µm³; s.e.m. for the 5 replicates: 1.9 µm³). This shows that
data obtained with different devices can be reliably compared.

Finally, we further validated the approach by assessing the volumes of known fission yeast size
mutants as well as wild-type diploid cells. In particular, we measured the size of well-described
temperature-sensitive cell cycle mutants: *wee1-50^{ts}* (small cells) and *cdc25-22^{ts}* (large cells) (Fantes,
1979; Fantes and Nurse, 1978; Nurse, 1975). As anticipated, *wee1-50^{ts}* and *cdc25-22^{ts}* cells grown at
the permissive temperature of 25 °C showed significantly smaller and larger volumes than wild type,
respectively (Fig. 2C, Fig. S1A). Note that the apparent broader distribution in *cdc25-22^{ts}* results from
increased overall size, as the volume profiles of these strains are similar when the data are normalized
to their respective median sizes (Fig. S1C). In the case of wild-type diploid cells, we found that the
median population volume is 1.84-fold higher than for haploid cells (Fig. 2D, Fig. S1B).

Altogether, these experiments demonstrate that FXm with specifically adapted microfluidic chips
is compatible with the analysis of yeast cell volume and produces highly reproducible results.
Furthermore, in contrast to measurements of cell length and width, FXm does not rely on 2D parameters
that are difficult to accurately determine. In particular, it will allow for identifying changes in volume

that result from limited alterations in cell diameter. Finally, our data also set a first conservative
threshold of 10 % for the estimation of differences in cell volume between populations.

Improving FXm image analysis

FXm image analysis consists of three steps (Cadart et al., 2017): image normalization, cell
selection and volume calculation. However, as FXm requires a low magnification objective, the size of
yeast cells and the high number of cells in a frame make it tedious to manually select individual cells
using the code established for mammalian cells. Interestingly, the normalization procedure generates a
mask that separates pillars and cells from the background (Cadart et al., 2017). Since the majority of
objects extracted at this step corresponds to valid individual cells, we took advantage of this mask and
developed a Python script to assist the user in cell selection (see Supplementary information). Thus,
instead of manually delineating the multiple regions of interest (Fig. 3A), this tool offers two distinct
strategies.

First, a manual cell selection mode can be chosen. Cells pre-selected by the normalization mask
are magnified and displayed (Fig. 3A), allowing the user to rapidly exclude incorrect objects such as
dust particles or cell aggregates. This approach makes the analysis of a high number of cells easier and
faster. Second, an automated and non-discriminatory selection can be applied. To exclude aberrant
measurements (*e.g.* cell aggregates, small dead cells), thresholds are then automatically set to remove
outliers based on the standard deviation (s.d.) of the population data. This is more appropriate than the
use of absolute cut-offs, which may need to be adjusted from strain to strain. After testing different s.d.
thresholds (data not shown), we found that excluding values outside of 2 s.d. (1 IQR for a normal
distribution) for the first and third quartiles was optimal for obtaining results similar to those established
with the manual method, correctly selecting between 87 and 92 % of the objects in our experiments
(Fig. 3B). Finally, our script also provides the option of separating the selected cells in different groups.
This is particularly useful when determining the volumes of distinct subpopulations, for instance based
on additional fluorescence markers (see below).

Collectively, the improvements brought by our analysis code facilitate FXm and provide more
options to the user for complex studies of cell populations. Interestingly, while manual selection is the
most accurate approach, we show that automated analysis with outlier exclusion allows a rapid
assessment of high numbers of cells while remaining sufficiently reliable for screening the volumes of
many different strains.

Small variations in cell dimensions and population heterogeneity

Our initial volume measurements in wild-type fission yeast cells and the evaluation of FXm experimental variability suggest that changes in volume ≥10 % can be reliably detected (Fig. 2). To further assess the sensitivity of FXm, we first determined the volume of cells showing a marginal difference in length at division compared to wild type. To this end, we took advantage of a fission yeast strain whose proliferation relies on a minimal cell cycle control network (MCN). In this background, cell proliferation is solely dependent on the oscillation of a single cyclin-dependent kinase (CDK) activity between two thresholds (high for mitosis and low for S phase) (Coudreuse and Nurse, 2010). This is achieved through the expression of a fusion protein between cyclin B/Cdc13 and Cdk1/Cdc2, in the absence of all other cell cycle cyclins (Coudreuse and Nurse, 2010). Interestingly, while behaving similarly to wild type (Coudreuse and Nurse, 2010), MCN cells show a minor increase of ~7 % in their length at division in our experimental conditions (15 μm for wild type and 16.1 μm for MCN cells, Fig. 4A). This is predicted to contribute to a similar increase in volume according to the geometric model of *S. pombe* cells (see above). Unexpectedly, FXm measurements showed a median volume difference of 17 % between these strains (Fig. 4A, Fig. S2; compare WT and MCN). This suggests that the change in size between wild type and MCN cells is not solely the result of an increase in cell length. Importantly, it demonstrates that FXm is a promising approach for investigating volume regulation in yeast, providing novel and more in-depth information on cell size and geometry.

Next, we tested whether FXm can reliably reveal volume changes below our initial threshold of 10 %. First, building on the results in Fig. 2A and 2C, we showed that a difference in volume of less than 10 % between wild-type cells grown at 32 *vs*. 25 ºC could be robustly established by simply increasing the number of experimental replicates (Fig. 4B and Fig. S3A). We then investigated whether even smaller alterations in volume at the population level could be detected as cells progressively change size over time. As a proof-of-concept, we used MCN cells, as they harbor a mutation in the Cdc2 moiety of the fusion module that makes them sensitive to dose-dependent and reversible inhibition of CDK activity by non-hydrolysable ATP analogues such as 3-MBPP1 (Bishop et al., 2000; Coudreuse and Nurse, 2010). This allowed us to block cells in G2 using the inhibitor, let them synchronously re-enter the cell cycle upon inhibitor wash-off (Fig. S3B, C) and monitor volume changes throughout the following G2 with high time resolution (Fig. 4C). Remarkably, even considering the experimental variability observed at certain time points, very small increases in volume (<5%) and changes in growth rates could be measured, demonstrating the high sensitivity of FXm in this context.

Another critical parameter in our understanding of cell size homeostasis is the size distribution in the population. This reflects the cell-to-cell heterogeneity in cell cycle progression and the strength of the size control operating in these cells. To date, most studies in fission yeast have focused on variability in cell length at division. However, not only does this approach exclude the majority of cells, but our results above suggest that evaluating cell volume distribution may lead to different conclusions. To

establish whether FXm is sufficiently sensitive for monitoring alteration of cell size profiles, we took
advantage of MCN cells in which the target residues of the conserved Wee1/Cdc25 feedback loop on
Cdc2 are mutated (T14A Y15F, MCN-AF) (Coudreuse and Nurse, 2010). Loss of this mitotic switch
results in an increase in cell-to-cell variability in length at division (Coudreuse and Nurse, 2010). In the
non-supplemented medium used in our experiments, we also found that these cells are on average
slightly smaller than wild type (Fig. 4A). As anticipated, we observed a broader distribution of cell
volume in MCN-AF compared to MCN and wild-type cells (Fig. 4A, D). Interestingly, the relative
changes in median volume *vs.* average length at division between MCN-AF, MCN and wild type
provide insights into the dimensions that are altered in these strains. (Fig. 4A, Table S1): while MCN-
AF and MCN appear to differ only in length (same ratios for volume *vs.* size at division), both strains
are likely to show an increase in cell diameter compared to wild type (differing ratios).

Altogether, these results demonstrate that FXm is easier and more sensitive than previously used
methods, enabling the monitoring of small alterations in cell volume and changes in size homeostasis
in populations of yeast cells. Coupling these data with more traditional cell length measurements will
shed light on how changes in volume are mediated and whether modulations of cellular geometry occur.

Measuring the size of cells with altered geometries and morphologies

FXm allows for measuring small changes in size that are difficult to accurately evaluate from cell
length and diameter measurements. The use of length as a proxy for fission yeast size is even more
problematic when studying mutants or conditions where dimension scaling or cell shape is altered.
Furthermore, strategies using cell outline detection to extrapolate cell volume may not be sufficiently
reliable, given the influence of minor changes in diameter on cell volume.

In this context, we determined the volumes of various strains showing defects in morphology and
shape. First, we assessed the size of strains lacking Rga2, Rga4 or Rga6, which are members of the
RhoGAP family. These factors are involved in polarized growth, and their loss impacts cell geometry
(Das et al., 2007; Revilla-Guarinos et al., 2016; Soto et al., 2010; Villar-Tajadura et al., 2008). When
ranking these strains according to their length at division, we found that Δ*rga2* cells are the longest,
followed by Δ*rga6* and Δ*rga4* (Fig. 5A). Strikingly, FXm, which is unaffected by changes in either
morphology or geometry, led to the opposite conclusion, with Δ*rga4* and Δ*rga2* cells having the largest
and smallest volumes, respectively (Fig. 5B, Fig. S4). This shows that cell length measurement is not
reliable as a proxy for cell size when studying such mutants. Our data also demonstrate the advantages
of FXm for establishing size differences between strains and determining the influence of diverse
pathways on cell volume regulation.

Next, we used FXm to evaluate the size of cells for which length can be difficult to measure or
inherently inaccurate due to morphological alterations. To this end, we used bent cells *(Δknk1, Δtea1)*

as well as cells with significant shape defects and irregular diameters (Δ*mal3*) (Beinhauer et al., 1997;

Mata and Nurse, 1997; Scheffler et al., 2014) (Fig. 5C). As anticipated, while cell length at division

could not be determined without excluding a significant fraction of the cells, FXm allowed us to

establish the profiles of cell volume in these populations and compare them with that of wild-type cells.

This again highlights the power of this method for studying size control mechanisms independently of

cell morphology (Fig. 5D, Fig. S5). Interestingly, in cells with morphological defects, both the median

volume and size distribution may change (Figs. S4, S5), potentially reflecting alterations of their cell

cycle organization and associated growth patterns. This opens the door to investigating how cell shape

may delineate the complex interplay between cell growth and the division cycle at the single-cell level.

Finally, we evaluated FXm data for significantly smaller cells that have stopped dividing. Upon

glucose exhaustion, fission yeast cells exit the cell cycle and enter a non-proliferative quiescent state,

which is associated with a dramatic reduction in cell size. Given the shape of the cells in these conditions

(Fig. 5E), determining their length and width to extrapolate their volume is difficult and unreliable. In

contrast, FXm established high-quality profiles of cell volume in these populations (Fig. 5F, Fig. S6).

This will allow for investigating the impact of cell size on cell physiology and aging in this critical

cellular state.

Collectively, this set of experiments demonstrates the versatility of FXm, which provides an

unprecedented ability to measure the size of any strain, irrespective of cell morphology and

physiological status. Our results also suggest that previous conclusions on size control in fission yeast

may need to be re-evaluated using FXm.

Measuring the volume of subpopulations

All experiments presented so far focus on the determination of cell volume in entire cell

populations. However, extracting the size of cells in specific subpopulations is critical for understanding

how different biological processes may interact with the mechanisms underlying cell size control and

how cell growth and proliferation are coupled. Thus, associating FXm with the use of other intracellular

fluorescent markers would represent a powerful method to take advantage of yeast genetics and explore

the complexity of size regulation in different conditions and throughout the cell cycle. Similarly, this

could allow for discriminating between distinct strains expressing specific markers in a complex

population, opening the door to exploring non-cell autonomous processes that may feed into volume

regulation.

To evaluate these approaches, we set out to determine cell volume at mitosis, using the non-histone

chromatin-associated protein Nhp6 coupled to the red fluorescent marker mCherry. First, we validated

that the presence of Nhp6::mCherry in the nucleus did not induce any significant signal in the FITC

channel under FXm conditions (Fig. S7A). This also showed that the slightly reduced volume of these

cells compared to wild type (Fig. 6A, Fig. S7B) results from an alteration of Nhp6 function due to the

tag rather than an experimental bias from mCherry fluorescence. Using our improved FXm code, we

segmented the population of binucleated cells (~11 % of the total number of cells) and determined their

median volume (126.3 ± 18.2 μm^3, compared to 93.5 ± 21.3 μm^3 for mononucleated cells, Fig. 6B). The

partial overlap between mono- and binucleated cells is consistent with the interruption of growth during

mitosis: cell size does not change throughout nuclear division, and growth only resumes in G1. Note

that the volume of binucleated cells measured in our assay is consistent with previous calculations of

cell volume at division (Nurse, 1975) but significantly differs from those presented in a more recent

study (Navarro and Nurse, 2012). This further demonstrates that fission yeast cell size cannot be reliably

and reproducibly determined using cell length and diameter. Altogether, our experiments validate the

possibility of combining FXm with additional markers in yeast for more complex investigations of cell

size control and dynamics.

Dynamic changes in fission yeast cell volume at the population and single-cell levels

Next, we tested the use of FXm for assessing dynamic changes in yeast cell size. First, we

ascertained whether coupling time course experiments with FXm allows for following induced

alterations in cell volume over time using asynchronous populations. A wide range of growth conditions

(*e.g.* heat stress (Vjestica et al., 2013), osmotic stress (Millar et al., 1995), change in nutrient availability

(Fantes and Nurse, 1977; Petersen, 2009)) as well as experimental modulation of specific cellular

pathways can trigger cell size changes. Understanding the dynamics of such changes may be key to

determining the implication of these processes in cell size homeostasis. To evaluate FXm in this context,

we took advantage of the analog-sensitivity of MCN cells. Indeed, reducing Cdc2/CDK activity in

fission yeast using this method leads to an increase in cell size (Chen et al., 2016; Coudreuse and Nurse,

2010). We therefore measured cell volume in asynchronous MCN cells at 40-minute intervals (~1/4 of

the cell cycle) after treatment with 0.05 µM 3-MBPP1 (Fig. 7). Our data show that MCN cells

progressively increase their volume, up to 1.7-fold after one doubling time. Strikingly, comparison of

the volume *vs*. size at division ratios throughout the experiment (Table S1) suggests complex dynamics

in the respective modulations of length and diameter upon CDK inhibition. This demonstrates that FXm

1) can be used to follow short-term alterations in median cell size and size distribution in asynchronous

yeast populations and 2) provides new perspectives for our understanding of how the geometrical

dimensions of these cells contribute to cell volume.

We then asked if FXm could be applied to analyze individual yeast cells in time-lapse experiments.

First, we determined the experimental noise for single cells in our chips by comparing 20 successive

measurements of 10 individual cells over a period of ~6 seconds (Fig. 8A). Making the assumption that

no volume changes occur in such a short time, this showed that the variability inherent to FXm when

using chips of reduced height remains very limited. Interestingly, it also suggests that when monitoring
single cells, volume alterations as low as ~3 % can be detected. Next, to perform time-lapse experiments
of several hours, we began by using modified chips in which the cell injection inlets were made bigger
(from 2.5 to 4 mm), acting as nutritional reservoirs to maintain growth conditions as stable as possible.
As fission yeast cells are non-adherent, this also required pre-coating of the chip coverslips with lectin.
In this context, we observed a linear increase in volume during each cell cycle and a plateau in size
around the time of cell division, corresponding to the well-described growth arrest at mitosis
(Mitchison, 1957; Mitchison and Nurse, 1985) (Fig. 8B). However, we found that in these experiments,
cells divide at smaller sizes in succeeding cell cycles, suggesting that the built-in reservoirs are
insufficient to ensure a favorable growth environment throughout our assay. We therefore applied a
constant flow of fresh medium containing FITC-Dextran (~3-5 μL.min^{-1}) for the entire duration of the
experiments. In this setup, we did not detect any consistent reduction in cell volume at division after
each cycle (Fig. 8C). Furthermore, in the presence of flow, the growth rate (see Materials and Methods)
was significantly higher than without medium renewal (Fig. 8B, C).

These data demonstrate that FXm is a unique tool for monitoring volume dynamics of small cells
with a high time resolution and for detecting changes in growth rate at the single-cell level. This will
allow for investigating the immediate responses of cells to environmental perturbations and the
behaviors of separate yeast cell lineages. In addition, this will make it possible to relate volume at birth,
growth rate and volume at division for a given cell. Coupled with additional markers, this technique
thus offers an unprecedented entry into the investigation of cell size control and its interplay with
complex biological processes.

DISCUSSION

Regulation of cell volume is critical for eukaryotic organisms, but the mechanisms underlying this process are still unclear. Yeast models have played a pioneering role in deciphering this complex trait and remain at the forefront of this very active research field. However, the difficulty of accurately measuring cell volume in these small unicellular organisms has been an obstacle for our understanding of size control and homeostasis. Here we show that the fluorescence exclusion method (FXm) is ideal for evaluating the volume of small yeast cells in diverse contexts. It provides high quality data on the volume profiles of yeast populations irrespective of cell morphology and physiological state, allows for single-cell analysis of volume changes with high time resolution and is compatible with the use of additional fluorescent markers. Our experiments also demonstrate that FXm gives researchers unique access to in-depth analyses of yeast cell volume dynamics and their coupling with cell cycle progression, integrating the size of cells at all stages of their division cycle. Furthermore, combining FXm with time-lapse imaging represents an unprecedented tool for exploring the changes in growth rate and cellular geometry that occur not only during proliferation or upon major transitions such as quiescence entry and exit, but also when cells respond to acute challenges. Thus, we believe that this technique will become a new standard for investigating cell size control in yeast, with implications for our knowledge of size regulation in complex eukaryotes.

Interestingly, our work led to a number of observations that may question existing models of cell size regulation. First, research on *S. pombe* size control and its interplay with cell cycle progression has mostly been based on the assumption that the diameter of proliferating haploid fission yeast cells is 1) constant from one cell to another and 2) homogenous along their growing axis. Our results clearly indicate that these assumptions can lead to incorrect conclusions when comparing strains, as shown using various mutants (Figs. 4, 5). In this context, FXm has the advantage of being a direct method for volume measurement that does not rely on any pre-conceived idea of cell geometry. Coupling FXm with cell length measurements also offers an additional level of understanding of cell size dynamics. For example, our data suggest that when cells are arrested in their division cycle through CDK inhibition, the geometric changes that occur over time are surprisingly complex, with length and volume alterations showing different kinetics (Table S1, MCN + 3-MBPP1). Similarly, we find that simplifying the architecture of cell cycle control or growing cells in distinct environments has unanticipated impacts on cell geometry, affecting both cell length and diameter (Table S1).

Moreover, focusing on subpopulations, as is the case when using cell length at division as a proxy for cell size, may hamper our comprehension of important processes linking cell growth, morphology and proliferation. For instance, the apparent changes in size distribution at the population level in morphology mutants suggest that the interplay between growth rate and cell cycle phases may be altered in these cells. This again may shed light on potentially novel inputs and mechanisms that modulate cell size dynamics. In this context, the different cell geometries of *rga* mutants (Table S1), as indicated by

our comparisons of volume *vs*. cell length at division ratios, may represent a promising model for

uncovering the key parameters used by the cells to "monitor" their size.

Our work demonstrates that previous models that describe the way cell size is regulated in fission

yeast need to be revisited using FXm. Together with the wealth of knowledge on size control in this

organism, this approach will provide new insights into the processes that underlie the biology of cell

growth. Interestingly, the investigation of biological processes has been enriched by the use of natural

yeast isolates. In fission yeast, these strains are known to be less homogenous in morphology and overall

size (Jeffares et al., 2015). FXm may therefore make it possible to fully exploit this unique resource and

take advantage of advanced population genetics strategies to decipher the regulation of cell size. While

we use fission yeast as a proof-of-concept model, our results also show that FXm can be applied to any

type of cell over a broad range of sizes. Our studies therefore establish the versatility and power of FXm

for investigating cell volume homeostasis and its regulation in small cells and model organisms,

contributing to our general understanding of the modulation of cellular dimensions and scaling in

eukaryotes.

MATERIALS AND METHODS

Fission yeast strains and methods

Standard methods and media were used (Hayles and Nurse, 1992; Moreno et al., 1991). All the strains described in this study are detailed in Table S2. The deletions of *rga2*, *rga4*, *rga6*, *mal3*, *knk1* and *tea1* have already been described (Soto et al., 2010; Villar-Tajadura et al., 2008) (Beinhauer et al., 1997; Mata and Nurse, 1997; Revilla-Guarinos et al., 2016; Scheffler et al., 2014). All experiments were carried out in non-supplemented minimal medium (EMM) at 32 ºC, except otherwise noted. To inhibit CDK activity in analog-sensitive MCN strains, the 3-MBPP1 inhibitor (A602960, Toronto Research Chemicals, Inc.) was dissolved in DMSO at stock concentrations of 10 or 0.4 mM and added to liquid cultures at a final concentration of 1 or 0.05 μM. The percentage of binucleated cells (Fig. S3C) was determined from heat fixed samples on microscope slides (70 ºC for 5 min) stained with a 11:1 solution of Blankophor (1 mg/mL): DAPI (1 μg/mL).

Microfabrication of FXm chips

PDMS microfluidic chips for FXm were prepared following standard microfabrication protocols (McDonald and Whitesides, 2002). In brief, master molds were made by spin-coating SU-8 2005 resin (MicroChem Corp., USA) on silicon wafers using a spincoater (Laurell Technologies, USA) according to manufacturer's instructions. Microstructures were then generated using high-resolution chrome masks (JD phototools, UK) and 365nm UV exposure (UV KUB 3, Kloe, France) followed by PGMEA (Sigma-Aldrich) development. For each mold, the height of the structures was determined as the average of three measurements perpendicular to the long axis of the design (Fig. 1B) using a Veeco Wyko NT9100 optical profilometer (Veeco Instruments Inc., USA). Note that molds showing significant variability between the three height measurements should be either discarded or used considering local rather than global averaged height. To produce chips for FXm, a 10:1 mixture of PDMS (Sylgard 184, Dow Corning, USA) was cast on the SU-8 master mold and allowed to cure at 70 °C for 2 hours. Inlets were then made using 2.5 or 4 mm biopsy punches, and chips were bonded to microscopy-grade coverslips by plasma activation (Harrick Plasma, USA).

Preparation of fission yeast cells for FXm

To perform FXm experiments, exponentially growing fission yeast cells were sampled at an optical density between 0.25 and 0.45 OD_{595}. To limit the formation of cell aggregates, a mild sonication cycle of 5 s at 10 % amplitude (Branson 450 Digital Sonifier, Emerson Electric Co.) was applied. This treatment had no effect on the results we obtained by FXm (data not shown). In order to optimize the number of cells that can be measured on a single FXm image in our chip design, cells were concentrated to ~5.5x10^7 cells/mL in their own conditioned minimal medium (note that to prevent bias in our volume measurements, we did not use autofluorescent rich medium). FITC-Dextran was then added to the

sample (FD10S, Sigma Aldrich) at a final concentration of 1 mg/mL, and cells were loaded into the

chip shortly before image acquisition. For evaluating the volume of quiescent cells, cells were sampled

24 hours after they reached 0.4 OD_{595}.

Microscopy

All microscopy experiments were carried out using an inverted Zeiss Axio Observer (Carl Zeiss

Microscopy LLC) equipped with a Lumencor Spectra X illumination system and an Orca Flash 4.0V2

sCMOS camera (Hamamatsu Photonics). Acquisition was performed using the VisiView software

(Visitron Systems GmbH). A Plan-Apochromat 63X/1.4 NA immersion lens and a Plan-Apochromat

20X/0.8 NA Ph2 lens (Carl Zeiss Microscopy LLC) were used for cell length measurements and FXm,

respectively. For routine volume measurements, the acquisition parameters were 100 ms exposure at 20

% illumination power, and 30 to 40 images were taken across the whole chambers to limit potential bias

due to local changes in chamber height. For the time-lapse experiments in Fig 8, 100 ms exposure at 10

% illumination power was used.

Image analysis

Cell length at division was determined from Blankophor images (1 mg/mL Blankophor solution,

A108741, Ambeed Inc.) using FIJI (National Institutes of Health) and the Pointpicker plug-in. For the

blankophor images in Fig. 2, 4, 5 and 7, the brightness and contrast were adjusted for display purposes.

For routine volume measurements, images were first normalized using a custom Matlab software

(Cadart et al., 2017). Cell selection and volume calculation were subsequently performed using the

normalization mask with a new, specifically-developed Python interface (see Results). For all Figures,

cell selection was done using the manual mode, except for Fig. 3B and 4C where automated cell

identification was used. Note that except for the time-lapses in Fig. 8B and C, we systematically

excluded cells that were undergoing cytokinesis but that had not fully separated into two individual

daughter cells; this is similar to studies of cell length at division. While this category can be integrated

as newly-born pairs of cells using the original Matlab software (Cadart et al., 2017), manually

delineating the division site would introduce measurement errors. Finally, our Python code is not

adapted to tracking the volume of individual cells in long time-lapse experiments (Fig. 8B, C), as slight

cell movements make it difficult to reliably follow single cells over time when using the normalization

mask pre-selection. For these assays, we used the original Matlab software (Cadart et al., 2017).

FXm time-lapse and analysis using fission yeast

As fission yeast cells are non-adherent, FXm time-lapses required coverslips to be coated in order to

prevent cells from moving inside the chamber, which would hamper data acquisition and analysis. To

this end, 10 µl of 1mg/ml filtered lectin was spread in a rectangle of ~10x5 mm at the center of the

coverslip and allowed to dry at 37 ºC. The coverslip was then washed once with filtered ultra-pure water

and dried at 37 ºC. For experiments without medium flow (Fig. 8B), Ø 4 mm inlets were made in the PDMS chip using the appropriate biopsy punch. 5 µl of cells prepared as described above were loaded in one inlet, and both inlets were then filled with 45 µl EMM containing the FITC-Dextran dye. Cells were allowed to flow into the chamber prior to imaging. For time-lapse experiments using medium flow (Fig. 8C), two successive layers of lectin were applied as above and inlets of Ø 0.75 mm were fabricated. Cells were then loaded by depositing a drop of cells prepared as above over one of the inlets, and a mild vacuum was applied at the other inlet using the tip of a 10 mL pipet connected to a vacuum pump. The loaded chip was then connected to a flow control system (pressure generator and flow controller, Elvesys, France), and a flow of medium (~3-5 µL/ min) containing the FITC-Dextran dye was used throughout the experiment. For the analysis of the time-lapse experiments, the two daughter cells after a division were manually delineated during the analysis and only one of them was followed for further investigation (Fig. 8B, C). The growth rate of each individual cell was determined by calculating the slopes of the traces during each cycle. This was obtained by linear regression considering the volume at birth as the first time point (when an invagination is detected) and excluding the mitotic plateau: entry into mitosis was assigned to the last point prior to strong reduction of growth (when two consecutive points on the graph decrease the slope).

ACKNOWLEDGMENTS

We thank Pei-Yun Jenny Wu and Snezhana Oliferenko for critically reading the manuscript. We thank Pascal Hersen and Giacomo Gropplero for the measurements of our chamber heights. We are grateful to Paul Nurse, Snezhana Oliferenko, Pilar Perez and Phong Tran for sharing strains and reagents.

COMPETING INTERESTS

No competing interests declared.

FUNDING

This work was supported by a grant to DC from the Agence Nationale de la Recherche (ANR, project eVOLve, ANR-18-CE13-0009; funding for AJ and DGR) and the Région Bretagne (ARED Pombevol, funding for DGR). JCR was supported by a fellowship from the Ministère de l'Enseignement Supérieur et de la Recherche. VRB was supported by an ERC Starting Grant to DC (SyntheCycle, Grant Agreement no. 310849).

 FIGURE LEGENDS

Figure 1. Microfluidic chips for FXm. A. Schematic of the principles of FXm (Cadart et al., 2017).
Left: cells are injected in a FXm chip (see *B*) in the presence of FITC-Dextran. Pillars are integrated for
ceiling support and image normalization. A low-magnification objective is used to acquire single-plane
images (see *C*). Right: After normalization, the volume of the cell can be determined from the loss of
fluorescence (grey area, the precise internal height of the chip must be determined (Cadart et al., 2017)).
B. Schematic of the chamber used for fission yeast. Left: design and dimensions (schematic not to
scale). Right: pillars are 60 µm in diameter and separated by 180 µm. **C.** Image of fission yeast cells in
an FXm experiment (prior to image normalization).

Figure 2. Measuring fission yeast cell volume by FXm. *A-D*, graphs are histograms with box and
whiskers plots, indicating the min, Q1, Q3 and max, with outliers determined by 1.5 IQR (interquartile
range). White dot indicates the median volume. Median volumes with s.d. are shown. **A.** Measurements
of wild-type fission yeast in five independent experiments (grey) and the total pooled dataset (black).
Chamber height: 5.53 µm; $n \geq 316$ for each replicate, $n=2266$ for the total dataset; values outside of the
plot range (n^*) were excluded: $n^* \leq 1$ for each replicate. **B.** Measurements of wild-type cells from a
single culture using five distinct devices of the indicated heights (grey) and the total pooled dataset
(black). $n \geq 225$ for each replicate, $n=1777$ for the total dataset; $n^*=0$ for each replicate. **C.** Top: wild
type (WT), *wee1-50*^{ts} and *cdc25-22*^{ts} temperature-sensitive strains were grown at the permissive
temperature of 25 °C and their volumes measured by FXm. Pooled datasets ($n \geq 1144$) of three
independent experiments ($n \geq 337$ for each replicate, Fig. S1A) are shown. $n^* \leq 6$ for each replicate.
Bottom: blankophor images with cell length at division (averages with s.d. for pooled datasets of three
independent experiments, $n \geq 100$ for each replicate). Scale bars = 10 µm. At 25 °C, WT cells have a
reduced volume and cell length at division compared to 32 °C (see *A*, *B* and Figs. 4-7). **D.** Top:
comparison of volume distributions in populations of WT haploid and diploid fission yeast cells by
FXm. Cells were grown in supplemented EMM6S at 32 °C. Pooled datasets ($n \geq 1073$) for three
independent experiments ($n \geq 207$ for each replicate, Fig. S1B) are shown. $n^* \leq 1$ for each replicate.
Bottom: blankophor images with cell length at division of the corresponding strains (averages with s.d.
for pooled datasets of three independent experiments, $n \geq 100$ for each replicate). Scale bars = 10 µm. In
EMM6S, WT cells have a volume that is similar to that in EMM (compare with *A*, *B* and Figs. 4-7) but
a reduced cell length at division (compare with Fig. 4A, 5A and 7), suggesting a difference in cell width
between these conditions.

Figure 3. Analysis pipeline for FXm data. A. Top: representative image showing the selection of
individual cells using the original FXm analysis code. Each cell must be identified from the FITC image

(see Fig. 1C; the area shown has been magnified for display purposes) and manually selected. Bottom:

representative image showing the tool that we developed, taking advantage of the normalization mask

for cell selection. All objects are automatically identified and displayed at a higher magnification. Cell

aggregates (bottom left), recently-divided cells (bottom right) or artifacts can be excluded by the user

(red-barred). **B.** Comparison of FXm results using manual (black) *vs.* automated (grey, 2 s.d. filter) cell

selection. Data for manual selection are from Fig. 2C. For automated selection, 1) the same images as

for the manual method were used, pooling the data from all replicates, and 2) a filter was applied that

excludes entries that are 2 s.d. above and below Q3 and Q1 (n^{**}), respectively (unfiltered selected

objects: $n \geq 1390$, excluded objects $n^{**} \leq 200$). The overlap between the two approaches for both correctly

assigned cells and outliers is between 87 and 92 %. Graphs are as in Fig. 2.

Figure 4. FXm allows for the detection of subtle changes in cell volume and size distribution. A.

Top: volume measurement of WT, MCN and MCN-AF cells by FXm. Pooled datasets ($n \geq 913$) of three

independent experiments ($n \geq 172$ for each replicate – Fig. S2) are shown. $n^* \leq 1$ for each replicate.

Graphs are as in Fig. 2. Bottom: blankophor images with cell length at division of the corresponding

strains (averages with s.d. for pooled datasets of three independent experiments, $n \geq 100$ for each

replicate). Scale bars = 10 μm. **B.** Comparison of the median volumes for 5 independent replicates of

WT at 25 and 32 °C. Full datasets are in Fig. S3A. Difference in median cell volume: 8.8 %. **C.** MCN

cells grown in EMM6S at 32 °C were G2-arrested using 1 μM 3-MBPP1 for 160 min and allowed to

synchronously re-enter the cell cycle by washing off the inhibitor (Coudreuse and Nurse, 2010) (Fig.

S3B). Cells were then injected in a FXm chip, and images were acquired at 5 min intervals throughout

the next G2, after cells had undergone a first round of mitosis and cell division (60 min, Fig. S3C). Top:

median population volumes at the indicated time points (black line – automated cell selection) with IQR

(grey area) are shown. Note that the higher volume for MCN cells compared to other Figures results

from the initial G2 block, during which cells grow without dividing. The phases of the cell cycle during

which cell volume was measured (first G2 and mitosis/M after release – Fig. S3C) are indicated. G2

growth rate: 0.59 μm^3.min^{-1} (linear regression from 60 to 110 min, prior to the mitotic plateau. From

125 to 140 min, the automated selection does not separate newly divided cells (17.4%, 25.9% and 25.5%

of the selected objects at 130, 135 and 140 min, respectively, are pairs of cells; the remainder of the

cells have not undergone cytokinesis at these time points). This results in an apparent increased growth

rate. Note the delay in cell cycle progression compared to Fig. S3C, which may be due to the growth

conditions in the absence of medium flow (see Fig. 8B). For each time point, $214 \leq n \leq 278$ (unfiltered

automatically selected objects). Outliers were removed using the 2 s.d. filter (see Fig. 3B; $28 \leq n^{**} \leq 49$,

representing a maximum of 18% of the number of selected objects). Bottom: representative images for

cells in FXm chips at 85 min (G2), 120 min (M) and 135 min (pairs of sister cells represent a significant

subset of the population at this time point, see above). **D.** Volume distribution for the indicated strains

as a percentage of the median volume of the population. Data are as in *A*. *n**≤10.

Figure 5. Measuring the volume of yeast cells with altered morphologies. A, C, E. Blankophor

images with cell length at division of the indicated strains (averages with s.d. for pooled datasets of

three independent experiments, *n*≥100 for each replicate). n.d.: not determined. Scale bars = 10 μm. **B.**

Volume measurement by FXm of the strains in *A*. Pooled datasets (*n*≥832) of three independent

experiments (*n*≥268 for each replicate – Fig. S4) are shown. *n**≤1 for each replicate. **D.** Volume

measurement by FXm of the strains in *C*. Pooled datasets (*n*≥940) of three independent experiments

(*n*≥230 for each replicate – Fig. S5) are shown. *n**=0 for each replicate. Data for WT are as in Fig. 2D

(haploids). **E.** Blankophor images of WT cells in exponential growth and quiescence. For quiescent

cells, samples were taken 24 h after the culture reached OD_{595} 0.4. For proliferating cells, cell length at

division is shown (average with s.d. for pooled dataset of three independent experiments, *n*≥100 for

each replicate). n.d.: not determined. Scale bars = 10 μm. **F.** Volume measurement by FXm of the cells

in *E*. Pooled datasets (*n*≥965) of three independent experiments (*n*≥238 for each replicate – Fig. S6) are

shown. *n**=0 for each replicate. *C, D*: Experiments were carried out in EMM6S due to the presence of

auxotrophies in some of the strains (Table S2). *B, D, F:* Graphs are as in Fig. 2.

Figure 6. FXm is compatible with the use of fluorescent markers to study subpopulations. A. Top:

volume measurement of WT and cells expressing *nhp6::mCherry* by FXm. Pooled datasets (*n*≥1568)

of three independent experiments (*n*≥499 for each replicate – Fig. S7B) are shown. *n**=0 for each

replicate. Median volume values with s.d. are shown. Bottom: blankophor images with cell length at

division of the strains above (averages with s.d. for pooled datasets of three independent experiments,

n≥110 for each replicate). Scale bars = 10 μm. **B.** Volume comparison between mono- (grey) and bi-

nucleated (black) cells in the population of *nhp6::mCherry* cells as in *A*. The percentages and median

volumes for the mono- (*n*=1619) and binucleated (*n*=196) subpopulations are indicated. Binucleated

cells include both septated (dashed line) and non-septated cells. *A, B*: Graphs are as in Fig. 2.

Figure 7. Dynamic changes in cell volume at the population level. Left: FXm volume measurement

of MCN cells after treatment with 0.05 μM 3-MBPP1. Untreated WT cells were used as a control. For

MCN, samples were collected every 40 min after addition of the inhibitor (*n*≥265 at each time point,

n=528 for WT). *n**≤2 for each dataset. Graphs are as in Fig. 2. Right: blankophor images with average

cell length at division and s.d. for the experiment in *A*. *n*≥100 for WT and MCN (0, 120 and 160 min).

For the 40 and 80 min time points, the initial delay in mitosis due to the inhibitor treatment leads to a

reduction in the number of dividing cells, making it difficult to obtain similar numbers for the

calculation of cell length at division (*n*=45 and 22, respectively). Septation indices: WT: 11.5 %, MCN

0 min: 20.4 %, MCN 40 min: 1 %, MCN 80 min: 2 %, MCN 120 min: 15.5 %, MCN 160 min: 26.9 %;

n≥400 for each time point. Scale bars = 10 μm.

Figure 8. Determination and real-time monitoring of single-cell volume by FXm. A. 20 consecutive

FXm images were acquired (see Materials and Methods; 100 ms exposure per image over a total time

of 6 s) and the volumes of 10 individual cells were followed throughout the experiment. The generation

time of WT cells in these conditions is ~140 min. The coefficients of variation are between 0.3 and

3.1%. **B.** FXm time-lapse experiment using WT in EMM at 32 °C without medium flow. **C.** FXm time-

lapse experiment using WT in EMM at 32 °C with a constant flow of medium (3-5 μL.min^{-1}) containing

FITC-Dextran. *B*, *C*: individual traces represent the volume dynamics of single-cell lineages (at each

division, only one of the two daughter cells is further monitored). Volumes were determined every 10

min. For display purposes, the traces are aligned to the first division (see Materials and Methods).

Average growth rates with s.d. are shown for each cycle. Note that the growth rate in *C* is consistent to

that obtained from the bulk analysis of a synchronized population (Fig. 4C).

REFERENCES

Amodeo, A. A. and Skotheim, J. M. (2016). Cell-Size Control. *Cold Spring Harb Perspect Biol* **8**,
a019083.

Baybay, E. K., Esposito, E. and Hauf, S. (2020). Pomegranate: 2D segmentation and 3D
reconstruction for fission yeast and other radially symmetric cells. *Scientific reports* **10**, 16580–
15.

Beinhauer, J. D., Hagan, I. M., Hegemann, J. H. and Fleig, U. (1997). Mal3, the fission yeast
homologue of the human APC-interacting protein EB-1 is required for microtubule integrity and
the maintenance of cell form. *The Journal of cell biology* **139**, 717–728.

Bishop, A. C., Ubersax, J. A., Petsch, D. T., Matheos, D. P., Gray, N. S., Blethrow, J., Shimizu,
E., Tsien, J. Z., Schultz, P. G., Rose, M. D., et al. (2000). A chemical switch for inhibitor-
sensitive alleles of any protein kinase. *Nature* **407**, 395–401.

Bryan, A. K., Goranov, A., Amon, A. and Manalis, S. R. (2010). Measurement of mass, density,
and volume during the cell cycle of yeast. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 999–1004.

Bryan, A. K., Hecht, V. C., Shen, W., Payer, K., Grover, W. H. and Manalis, S. R. (2014).
Measuring single cell mass, volume, and density with dual suspended microchannel resonators.
Lab Chip **14**, 569–576.

Cadart, C., Monnier, S., Grilli, J., Sáez, P. J., Srivastava, N., Attia, R., Terriac, E., Baum, B.,
Cosentino-Lagomarsino, M. and Piel, M. (2018). Size control in mammalian cells involves
modulation of both growth rate and cell cycle duration. *Nat Commun* **9**, 3275–15.

Cadart, C., Venkova, L., Recho, P., Lagomarsino, M. C. and Piel, M. (2019). The physics of cell-
size regulation across timescales. *Nat. Phys.* **15**, 993–1004.

Cadart, C., Zlotek-Zlotkiewicz, E., Venkova, L., Thouvenin, O., Racine, V., Le Berre, M.,
Monnier, S. and Piel, M. (2017). Fluorescence eXclusion Measurement of volume in live cells.
Methods Cell Biol. **139**, 103–120.

Chen, T., Gómez-Escoda, B., Munoz-Garcia, J., Babic, J., Griscom, L., Wu, P.-Y. J. and
Coudreuse, D. (2016). A drug-compatible and temperature-controlled microfluidic device for
live-cell imaging. *Open Biol* **6**, 160156.

Cook, M. and Tyers, M. (2007). Size control goes global. *Curr. Opin. Biotechnol.* **18**, 341–350.

Coudreuse, D. and Nurse, P. (2010). Driving the cell cycle with a minimal CDK control network.
Nature **468**, 1074–1079.

Das, M., Wiley, D. J., Medina, S., Vincent, H. A., Larrea, M., Oriolo, A. and Verde, F. (2007).
Regulation of cell diameter, For3p localization, and cell symmetry by fission yeast Rho-GAP
Rga4p. *Molecular biology of the cell* **18**, 2090–2101.

Facchetti, G., Knapp, B., Flor-Parra, I., Chang, F. and Howard, M. (2019). Reprogramming
Cdr2-Dependent Geometry-Based Cell Size Control in Fission Yeast. *Current biology : CB* **29**,
350–358.e4.

Fantes, P. (1979). Epistatic gene interactions in the control of division in fission yeast. *Nature* **279**,
428–430.

Fantes, P. A. (1977). Control of cell size and cycle time in Schizosaccharomyces pombe. *J. Cell. Sci.*
24, 51–67.

Fantes, P. A. and Nurse, P. (1978). Control of the timing of cell division in fission yeast. Cell size
mutants reveal a second control pathway. *Experimental cell research* **115**, 317–329.

Fantes, P. and Nurse, P. (1977). Control of cell size at division in fission yeast by a growth-
modulated size control over nuclear division. *Experimental cell research* **107**, 377–386.

Grant, N. A., Abdel Magid, A., Franklin, J., Dufour, Y. and Lenski, R. E. (2021). Changes in Cell
Size and Shape during 50,000 Generations of Experimental Evolution with Escherichia coli.
Journal of bacteriology **203**.

Hayles, J. and Nurse, P. (1992). Genetics of the fission yeast Schizosaccharomyces pombe. *Annu.*
Rev. Genet. **26**, 373–402.

Jeffares, D. C., Rallis, C., Rieux, A., Speed, D., Převorovský, M., Mourier, T., Marsellach, F. X.,
Iqbal, Z., Lau, W., Cheng, T. M. K., et al. (2015). The genomic and phenotypic diversity of
Schizosaccharomyces pombe. *Nature genetics* **47**, 235–241.

Lancaster, O. M., Le Berre, M., Dimitracopoulos, A., Bonazzi, D., Zlotek-Zlotkiewicz, E.,
Picone, R., Duke, T., Piel, M. and Baum, B. (2013). Mitotic rounding alters cell geometry to
ensure efficient bipolar spindle formation. *Dev. Cell* **25**, 270–283.

Lenski, R. E. and Travisano, M. (1994). Dynamics of adaptation and diversification: a 10,000-
generation experiment with bacterial populations. *Proc. Natl. Acad. Sci. U.S.A.* **91**, 6808–6814.

Li, Q., Rycaj, K., Chen, X. and Tang, D. G. (2015). Cancer stem cells and cell size: A causal link?
766   *Semin. Cancer Biol.* **35**, 191–199.

767   **Lloyd, A. C.** (2013). The regulation of cell size. *Cell* **154**, 1194–1205.

768   **Marshall, W. F., Young, K. D., Swaffer, M., Wood, E., Nurse, P., Kimura, A., Frankel, J.,**
769   **Wallingford, J., Walbot, V., Qu, X., et al.** (2012). What determines cell size? *BMC Biol.* **10**,
770   101.

771   **Martin, S. G. and Berthelot-Grosjean, M.** (2009). Polar gradients of the DYRK-family kinase
772   Pom1 couple cell length with the cell cycle. *Nature* **459**, 852–856.

773   **Mata, J. and Nurse, P.** (1997). tea1 and the microtubular cytoskeleton are important for generating
774   global spatial order within the fission yeast cell. *Cell* **89**, 939–949.

775   **McDonald, J. C. and Whitesides, G. M.** (2002). Poly(dimethylsiloxane) as a material for fabricating
776   microfluidic devices. *Acc. Chem. Res.* **35**, 491–499.

777   **Millar, J. B., Buck, V. and Wilkinson, M. G.** (1995). Pyp1 and Pyp2 PTPases dephosphorylate an
778   osmosensing MAP kinase controlling cell size at division in fission yeast. *Genes & development*
779   **9**, 2117–2130.

780   **Mitchison, J. M.** (1957). The growth of single cells. I. Schizosaccharomyces pombe. *Experimental*
781   *cell research* **13**, 244–262.

782   **Mitchison, J. M. and Nurse, P.** (1985). Growth in cell length in the fission yeast
783   Schizosaccharomyces pombe. *J. Cell. Sci.* **75**, 357–376.

784   **Model, M.** (2020). Comparison of cell volume measurements by fluorescence and absorption
785   exclusion microscopy. *J Microsc* **280**, 12–18.

786   **Model, M. A.** (2018). Methods for cell volume measurement. *Cytometry A* **93**, 281–296.

787   **Mongold, J. A. and Lenski, R. E.** (1996). Experimental rejection of a nonadaptive explanation for
788   increased cell size in Escherichia coli. *Journal of bacteriology* **178**, 5333–5334.

789   **Moreno, S., Klar, A. and Nurse, P.** (1991). Molecular genetic analysis of fission yeast
790   Schizosaccharomyces pombe. *Meth. Enzymol.* **194**, 795–823.

791   **Moseley, J. B., Mayeux, A., Paoletti, A. and Nurse, P.** (2009). A spatial gradient coordinates cell
792   size and mitotic entry in fission yeast. *Nature* **459**, 857–860.

793   **Mueller, R. L.** (2015). Genome Biology and the Evolution of Cell-Size Diversity. *Cold Spring Harb*
794   *Perspect Biol* **7**, a019125.

795   **Navarro, F. J. and Nurse, P.** (2012). A systematic screen reveals new elements acting at the G2/M
796   cell cycle control. *Genome Biol.* **13**, R36.

797   **Nurse, P.** (1975). Genetic control of cell size at cell division in yeast. *Nature* **256**, 547–551.

798   **Pan, K. Z., Saunders, T. E., Flor-Parra, I., Howard, M. and Chang, F.** (2014a). Cortical
799   regulation of cell size by a sizer cdr2p. *Elife* **3**, e02040.

800   **Pan, K. Z., Saunders, T. E., Flor-Parra, I., Howard, M. and Chang, F.** (2014b). Cortical
801   regulation of cell size by a sizer cdr2p. *Elife* **3**, e02040.

Pedersen, S. F., Hoffmann, E. K. and Mills, J. W. (2001). The cytoskeleton and cell volume
regulation. *Comp. Biochem. Physiol., Part A Mol. Integr. Physiol.* **130**, 385–399.

Petersen, J. (2009). TOR signalling regulates mitotic commitment through stress-activated MAPK
and Polo kinase in response to nutrient stress. *Biochem. Soc. Trans.* **37**, 273–277.

Pietras, A. (2011). Cancer stem cells in tumor heterogeneity. *Adv. Cancer Res.* **112**, 255–281.

Revilla-Guarinos, M. T., Martín-García, R., Villar-Tajadura, M. A., Estravís, M., Coll, P. M.
and Pérez, P. (2016). Rga6 is a Fission Yeast Rho GAP Involved in Cdc42 Regulation of
Polarized Growth. *Molecular biology of the cell*.

Saunders, T. E., Pan, K. Z., Angel, A., Guan, Y., Shah, J. V., Howard, M. and Chang, F. (2012).
Noise reduction in the intracellular pom1p gradient by a dynamic clustering mechanism. *Dev.*
Cell **22**, 558–572.

Scheffler, K., Recouvreux, P., Paoletti, A. and Tran, P. T. (2014). Oscillatory AAA+ ATPase
Knk1 constitutes a novel morphogenetic pathway in fission yeast. *Proc. Natl. Acad. Sci. U.S.A.*
111, 17899–17904.

Soifer, I., Robert, L. and Amir, A. (2016). Single-Cell Analysis of Growth in Budding Yeast and
Bacteria Reveals a Common Size Regulation Strategy. *Current biology : CB* **26**, 356–361.

Soto, T., Villar-Tajadura, M. A., Madrid, M., Vicente, J., Gacto, M., Pérez, P. and Cansado, J.
(2010). Rga4 modulates the activity of the fission yeast cell integrity MAPK pathway by acting
as a Rho2 GTPase-activating protein. *The Journal of biological chemistry* **285**, 11516–11525.

Sun, S. and Gresham, D. (2021). Cellular quiescence in budding yeast. *Yeast* **38**, 12–29.

Sveiczer, A., Novak, B. and Mitchison, J. M. (1996). The size control of fission yeast revisited. *J.*
Cell. Sci. **109 (Pt 12)**, 2947–2957.

Taheri-Araghi, S., Bradde, S., Sauls, J. T., Hill, N. S., Levin, P. A., Paulsson, J., Vergassola, M.
and Jun, S. (2015). Cell-size control and homeostasis in bacteria. *Current biology : CB* **25**, 385–
391.

Villar-Tajadura, M. A., Coll, P. M., Madrid, M., Cansado, J., Santos, B. and Pérez, P. (2008).
Rga2 is a Rho2 GAP that regulates morphogenesis and cell integrity in S. pombe. *Mol.*
Microbiol. **70**, 867–881.

Vjestica, A., Zhang, D., Liu, J. and Oliferenko, S. (2013). Hsp70-Hsp40 chaperone complex
functions in controlling polarized growth by repressing Hsf1-driven heat stress-associated
transcription. *PLOS Genetics* **9**, e1003886.

Yang, J., Dungrawala, H., Hua, H., Manukyan, A., Abraham, L., Lane, W., Mead, H., Wright,
J. and Schneider, B. L. (2011). Cell size and growth rate are major determinants of replicative
lifespan. *Cell cycle* **10**, 144–155.

Zegman, Y., Bonazzi, D. and Minc, N. (2015). Measurement and manipulation of cell size
parameters in fission yeast. *Methods Cell Biol.* **125**, 423–436.

Zhurinsky, J., Leonhard, K., Watt, S., Marguerat, S., Bähler, J. and Nurse, P. (2010). A
coordinated global control over cellular transcription. *Current biology : CB* **20**, 2010–2015.

Zlotek-Zlotkiewicz, E., Monnier, S., Cappello, G., Le Berre, M. and Piel, M. (2015). Optical
volume and mass measurements show that mammalian cells swell during mitosis. *The Journal of*
cell biology **211**, 765–774.

FIGURE 1

A

Microfluidic chip

FITC-Dextran

Pillar   Pillar

20X lens

Fluorescence intensity

Max

Min

Area

B

15 mm

0.5 mm

1 mm

2.6 mm

180 µm

60 µm

C

FIGURE 2

FIGURE 3

A

Cell: 12 (99 µm3)   Cell: 13 (89 µm3)

Cell: 17 (292 µm3)   Cell: 18 (150 µm3)

B

FIGURE 4

FIGURE 5

FIGURE 6

A

WT   nhp6::mCherry

14.9 ± 1.0 µm   14.6 ± 1.1 µm

B

FIGURE 7

FIGURE 8

A

B

C