Retrograde adenosine/A$_{2A}$ receptor signaling facilitates excitatory synaptic transmission and seizures

Kaoutsar Nasrallah[1,6], Coralie Berthoux[1], Yuki Hashimotodani[1,4], Andrés E. Chávez[1,5], Michelle Gulfo[1], Rafael Luján[3], Pablo E. Castillo[1, 2, *]

[1] Dominick P. Purpura Department of Neuroscience, [2] Department of Psychiatry & Behavioral Sciences, Albert Einstein College of Medicine, Bronx, NY 10461, U.S.A.; [3] Instituto de Investigación en Discapacidades Neurológicas (IDINE), Facultad de Medicina, Universidad Castilla-La Mancha, 02008 Albacete, Spain; [4] Graduate School of Brain Science, Doshisha University, Kyoto, Japan.

[5] Present address: Centro Interdisciplinario de Neurociencia de Valparaíso, Facultad de Ciencias, Universidad de Valparaíso, Valparaíso 2340000, Chile.
[6] Present address: Department of Biological Sciences, Fordham University, Bronx, NY 10458, U.S.A.

Running title: Adenosine/A$_{2A}$ receptor retrograde signaling mediates presynaptic LTP

Keywords: Mossy cell, hippocampus, dentate gyrus, retrograde signaling, epilepsy, presynaptic, BDNF, TrkB, PKA, LTP.

* To whom correspondence should be addressed:

Pablo E. Castillo, MD/PhD
Dominick P. Purpura Department of Neuroscience
Albert Einstein College of Medicine
1410 Pelham Parkway South
Kennedy Center, Room 703
Bronx, NY 10461, USA
Email: pablo.castillo@einsteinmed.edu

Title: 92 characters (with spaces)
Total number of words: Summary: 150; Introduction: 607; Results: 3,003; Discussion: 1,770
Methods: 3,045
Figures: 7; Supplementary figures: 7

Highlights

- Postsynaptic firing induces presynaptic LTP at mossy cell to granule cell synapses
- Postsynaptic TrkB activation induces adenosine release from granule cells
- Presynaptic adenosine A_{2A} receptors are necessary and sufficient to induce LTP
- Adenosine/A_{2A}R signaling within the dentate gyrus is pro-convulsant

In Brief:

Nasrallah et al. report a novel retrograde signaling pathway at hippocampal synapses that involves postsynaptic TrkB-dependent release of adenosine and the activation of presynaptic A_{2A} receptors. This pathway mediates presynaptic long-term potentiation at a key hippocampal excitatory synapse and can also promote epileptic seizures.

Summary

Retrograde signaling at the synapse is a fundamental way by which neurons communicate and neuronal circuit function is fine-tuned upon activity. While long-term changes in neurotransmitter release commonly rely on retrograde signaling, the mechanisms remain poorly understood. Here, we identified adenosine/A_{2A} receptor ($A_{2A}R$) as a novel retrograde signaling pathway underlying presynaptic long-term potentiation (LTP) at a hippocampal excitatory circuit critically involved in memory and epilepsy. Transient burst activity of a single dentate granule cell induced LTP of mossy cell synaptic inputs, a BDNF/TrkB-dependent form of plasticity that facilitates seizures. Postsynaptic TrkB activation released adenosine from granule cells, uncovering a non-conventional BDNF/TrkB signaling mechanism. Moreover, presynaptic $A_{2A}Rs$ were necessary and sufficient for LTP. Lastly, seizure induction released adenosine in a TrkB-dependent manner, while removing $A_{2A}Rs$ or TrkB from the dentate gyrus had anti-convulsant effects. By mediating presynaptic LTP, adenosine/$A_{2A}R$ retrograde signaling may modulate dentate gyrus-dependent learning and promote epileptic activity.

INTRODUCTION

Retrograde signaling at the synapse is a well-established mechanism whereby a neuron can strengthen or weaken the synaptic inputs it receives (Fitzsimonds and Poo, 1998; Regehr et al., 2009). Diverse types of messengers, including lipids, gases, peptides and conventional neurotransmitters, can be released from the postsynaptic neuron upon activity and activate presynaptic receptors or other molecular targets, establishing a retrograde signaling system that powerfully increases or decreases neurotransmitter release. Retrograde signaling has been implicated in short-term synaptic plasticity (Regehr *et al.*, 2009), presynaptic long-term plasticity (Castillo, 2012; Monday et al., 2018) and also in presynaptic homeostatic plasticity (Davis and Muller, 2015), all of which rely on changes in neurotransmitter release. Despite the significance of retrograde signaling in fine-tuning neuronal circuit function, important knowledge gaps remain regarding the identity and release mechanism of the retrograde messenger, as well as the presynaptic targets and downstream molecular cascades implicated in regulating neurotransmitter release.

The dentate gyrus (DG), the major input area of the hippocampus, contains two main types of excitatory neurons: dentate granule cells (GCs) and hilar mossy cells (MCs). MCs and GCs form an associative circuit that is proposed to play a key role in DG-dependent cognitive functions (Lisman, 1999; Scharfman, 2016) and epilepsy (Botterill et al., 2019; Bui et al., 2018; Jinde et al., 2013; Nasrallah et al., 2022; Ratzliff et al., 2002; Scharfman and Myers, 2012). Repetitive stimulation of MC axons with physiologically relevant patterns of activity triggers robust presynaptic long-term potentiation at MC-GC synapses (MC-GC LTP) but not at MC to interneurons synapses (Hashimotodani et al., 2017). Recent evidence indicates that MC-GC LTP can be induced *in vivo* by enriched environment exposure (Berthoux et al., 2023) and experimental epileptic activity (Nasrallah *et al.*, 2022). Uncontrolled strengthening of MC-GC transmission promotes seizures and likely contributes to the pro-epileptic role of MCs in early epilepsy stages (Botterill *et al.*, 2019; Nasrallah *et al.*, 2022). Thus, it is crucial to determine the precise mechanism underlying MC-GC LTP. This presynaptic form of plasticity is mechanistically unique. Its induction is NMDA receptor-independent but requires postsynaptic BDNF/TrkB signaling upstream of presynaptic cAMP/PKA signaling (Hashimotodani *et al.*, 2017), strongly suggesting the involvement of a retrograde signaling system whose identity is unknown.

The following criteria must be satisfied to establish retrograde signaling as a mechanism of presynaptic long-term plasticity. First, the retrograde messenger must be synthesized and released from the postsynaptic compartment. Second, interfering with the synthesis

and/or release of this messenger should prevent plasticity. Third, the target for the
retrograde messenger must be present in the presynaptic bouton. Fourth, interfering with
the presynaptic target should also prevent plasticity. Lastly, activation of the presynaptic
target by the retrograde messenger or some analogous molecule should mimic long-term
plasticity –although in some cases this activation alone may be insufficient to induce
plasticity.

In this study, we sought to determine the retrograde signal involved in presynaptic LTP at
MC-GC synapses. To our surprise, we found that most conventional retrograde
messengers were not implicated in this form of plasticity. Using selective pharmacology,
immunoelectron microscopy, and a conditional knockout strategy, we discovered that
activation of presynaptic G_s-coupled adenosine A_{2A} receptors ($A_{2A}Rs$) is necessary and
sufficient to induce MC-GC LTP. In addition, TrkB activation mobilized adenosine from
GCs, uncovering a novel TrkB-dependent signaling mechanism, whereas interfering with
GC adenosine release abolished LTP. Furthermore, adenosine was released *in vivo*
during acutely induced epileptic seizures in a TrkB-dependent manner, while removing
$A_{2A}Rs$ or TrkB from hippocampal excitatory neurons had anti-convulsant effects. Our
findings not only establish adenosine/$A_{2A}R$ as a novel retrograde signaling system that
mediates presynaptic plasticity, but also uncover a synapse-specific mechanism by which
BDNF/TrkB and $A_{2A}Rs$ may promote epileptic activity.

RESULTS

Theta-burst firing of a single GC induces presynaptic LTP at MC-GC synapse

To identify the retrograde signal mediating MC-GC LTP, we first tested whether activation
of a single GC, a manipulation previously used to characterize endocannabinoid
retrograde signaling in long-term plasticity (Younts et al., 2013), could trigger MC-GC LTP.
We found that GC theta-burst firing (TBF: 10 bursts at 5 Hz of 5 AP at 50 Hz, repeated 4
times every 5s, **Figure 1A**) induced LTP selectively at MC-GC synapses but not at
neighboring medial perforant path (MPP)-GC synapses (**Figure 1B**). In addition, the group
II mGluR agonist DCG-IV (1 µM), which selectively abolishes GC-MC (Lysetskiy et al.,
2005) but not MC-GC synaptic transmission (Chiu and Castillo, 2008), did not affect the
magnitude of TBF-induced LTP (**Figure S1A**), indicating that this plasticity does not rely
on GC recruitment of MC firing. We then tested whether TBF-LTP shared the same
mechanistic features as synaptically-induced MC-GC LTP. Like synaptically-induced MC-
GC LTP (Hashimotodani *et al.*, 2017), TBF-LTP is likely expressed presynaptically, as
indicated by a significant reduction in both paired-pulse ratio (PPR) and coeCfficient of

variation (CV) (**Figure 1C**) (see Methods). Moreover, TBF-LTP was abolished in the presence of the selective TrkB antagonist ANA-12 (15 µM) (**Figure S1B**), and in postsynaptic (**Figure 1D** and **Figure S1C**) but not presynaptic *TrkB* conditional knockout (cKO) mice (**Figure S1D**) (see Methods). TBF-LTP was also abolished in postsynaptic *Bdnf* cKO mice (**Figure S1E**), in GCs patch-loaded with botulinum toxin-B (Botox, 0.5 µM; **Figure S1F**), which blocks BDNF release (Shimojo et al., 2015), and in GCs loaded with the calcium chelator BAPTA (20 mM), which blocks postsynaptic calcium rise during TBF (**Figure S1G**). We discarded the possibility that alterations in GC membrane properties or basal transmitter release could underlie the lack of LTP in postsynaptic *Bdnf* cKO (Nasrallah et al., 2022) and postsynaptic *TrkB* cKO (**Figure S1H and S1I**). The PKA inhibitors H89 (10 µM, 40- to 60-min pre-incubation and bath applied) (**Figure S1J**) and membrane-permeable PKI_{14-22} myristoylated (1 µM, 40- to 60-min pre-incubation and bath applied)(**Figure 1E**) also abolished TBF-LTP, whereas loading the membrane-impermeable PKI_{6-22} (2.5 µM) in GCs did not affect this plasticity (**Figure 1E**). TBF-LTP was normally induced in the presence of the NMDAR antagonist D-APV (50 µM) (**Figure S1K**). Remarkably, while postsynaptic *TrkB* or *Bdnf* deletion abolished GC TBF-induced LTP (**Figure 1D,1F** and **Figure S1C, E**), it had no impact on the chemical LTP induced by transient activation of PKA with the adenylyl cyclase activator forskolin (Berthoux *et al.*, 2023; Hashimotodani *et al.*, 2017; Nasrallah *et al.*, 2022), supporting the notion that PKA acts downstream of BDNF/TrkB. To remove *TrkB* from GC specifically, we took advantage of a lentivirus encoding Cre under the control of the C1ql2 promoter, which achieves sparse labeling of GCs only (**Figure S1C**) (Barthet et al., 2018; Monday et al., 2022). In all, these manipulations (summarized in **Figure 1F**) indicated that, like synaptically-induced MC-GC LTP, TBF-LTP requires postsynaptic BDNF/TrkB signaling and presynaptic PKA signaling and is NMDAR-independent (**Figure 1G**). Lastly, synaptically-induced MC-GC LTP and TBF-LTP occluded each other (**Figure S2**), indicating a common mechanism. Thus, GC TBF triggered a presynaptic form of LTP with identical properties to the synaptically induced MC-GC LTP (Hashimotodani *et al.*, 2017), thereby establishing a simple, single-cell manipulation to investigate the identity of the retrograde messenger (**Figure 1G**).

A non-conventional retrograde signal likely mediates presynaptic LTP at MC-GC synapses

Previous work discarded endocannabinoids, glutamate, and GABA as retrograde signals mediating MC-GC LTP (Hashimotodani *et al.*, 2017; Jensen et al., 2021), and our current finding showing that TBF-LTP was normally induced in presynaptic *TrkB* cKO mice (**Figure S1D**), also discards BDNF as retrograde messenger. We therefore tested the role of other conventional retrograde messengers, such as nitric oxide (NO) and lipid-derived

messengers (Regehr *et al.*, 2009). Application of the NO synthase inhibitor L-NAME (100
µM, 50- to 90-min pre-incubation and bath applied) did not impair TBF-LTP (**Figure S3A**),
but as a positive control it significantly reduced LTP at CA3-CA1 synapses (**Figure S3B**),
as previously reported (Williams et al., 1993). Blockade of lipid-derived messengers with
a cocktail of inhibitors consisting of the fatty acid amide hydrolase (FAAH) and
anandamide amidase inhibitor AACOCF3 (10 µM, 50- to 80-min pre-incubation and bath
applied), the lipoxygenases inhibitor baicalein (3 µM, 50- to 80-min pre-incubation and
bath applied), and the lipase inhibitor THL (tetrahydrolipstatin or orlistat, 4 µM in the
recording pipette) did not impair TBF-LTP either (**Figure S3C**). As positive controls, we
found that both AACOCF3 and baicalein significantly reduced long-term depression (LTD)
at CA3-CA1 synapses (**Figure S3D**) (Normandin et al., 1996), and loading the lipase
inhibitor THL (4 µM) in the recording pipette efficiently blocked CA1 inhibitory LTD (iLTD)
(**Figure S3E**) (Chevaleyre and Castillo, 2003). Lastly, as was true for TBF-LTP, blocking
NO signaling and lipid-derived messengers in interleaved control experiments did not alter
synaptically-induced LTP either (**Figure S4**). Altogether, our results strongly suggest the
involvement of a non-conventional retrograde messenger in presynaptic MC-GC LTP.
250
251 **Presynaptic A_{2A}Rs are required for MC-GC LTP**
252
Presynaptic PKA signaling is necessary and sufficient to induce LTP of MC-GC synaptic
transmission (Hashimotodani *et al.*, 2017), raising the possibility that the retrograde
messenger induces LTP by activating a G_s-coupled G protein-coupled receptor (GPCR)
located on MC axon terminals. The G_s-coupled adenosine A_{2A} receptor (A_{2A}R) is a good
candidate as it has been implicated in BDNF-mediated plasticity at other synapses
(Diogenes et al., 2004; Fontinha et al., 2008; Sebastiao and Ribeiro, 2009b), and the
endogenous ligand adenosine can be released from neurons upon activity (King et al.,
2006; Latini and Pedata, 2001; Lovatt et al., 2012; Pons-Bennaceur et al., 2019; Wall and
Dale, 2013). To test whether adenosine mediates MC-GC LTP by activating presynaptic
A_{2A}Rs, we first examined whether A_{2A}R antagonism impaired this plasticity. Two different
A_{2A}R antagonists, SCH 58261 (100 nM) and ZM241385 (50 nM), abolished TBF-LTP
(**Figure 2A**). Moreover, bath application of SCH 58261 did not change basal transmission
(**Figure 2B**) but prevented the induction of LTP by BDNF puffs (8 nM, 2 puffs of 3 s)
delivered in the inner molecular layer (IML) (**Figure 2C**) (Hashimotodani *et al.*, 2017),
strongly suggesting that A_{2A}R activation is required downstream of the BDNF/TrkB
cascade. It is unlikely that postsynaptic A_{2A}Rs could be implicated given that including the
GPCR inhibitor GDPβS (1 mM) in the recording pipette did not affect TBF-LTP, whereas
as a positive control it abolished the change in holding current induced by the selective
$GABA_B$ receptor agonist baclofen (10 µM) (**Figure 2D**). Likewise, synaptically-induced
MC-GC LTP was also abolished by SCH 58261 (**Figure S5A**), but not by including GDPβS

in the recording pipette (**Figure S5B**). To demonstrate the role of presynaptic A_{2A}Rs in
MC-GC LTP, we also employed a conditional knockout strategy combined with
optogenetics which allowed us to selectively activate A_{2A}R-deficient MC axons expressing
the fast opsin ChIEF (**Figure 2E**). We found that synaptic responses elicited by the
activation of these axons did not undergo TBF-LTP (**Figure 2F**). In contrast, deleting
Adora2a from MCs did not significantly affect basal neurotransmitter release, as indicated
by the lack of PPR change (**Figure 2G**). Taken together, these findings indicate that
presynaptic, and not postsynaptic, A_{2A}R activation is necessary for LTP at MC-GC
synapses.

Activation of presynaptic A_{2A}Rs is sufficient to induce MC-GC LTP

We next tested whether A_{2A}R activation was sufficient to induce LTP at MC-GC synapses.
Bath application of the A_{2A}R agonist CGS21680 (50 nM, 15 min) selectively enhanced MC
but not MPP EPSC amplitude (**Figure 3A**). This synapse-specific potentiation was
associated with a significant decrease in both PPR and CV (**Figure 3B**), suggesting a
presynaptic mechanism of expression, and supporting a presynaptic location of A_{2A}Rs. In
addition, we tested whether supramammillary (SuM) inputs express functional A_{2A}Rs
using optogenetic activation of SuM inputs (Tabuchi et al., 2022) (see Methods). Unlike
MC EPSCs, SuM EPSCs were not enhanced by CGS21680 bath application (**Figure 3C**).
We then directly assessed the contribution of presynaptic A_{2A}Rs at MC-GC synapses by
testing CGS21680-induced potentiation in presynaptic A_{2A}Rs cKO mice. CGS21680 failed
to increase MC-GC synaptic transmission in presynaptic *Adora2a* cKO mice (**Figure 3D).**
Adding the selective A_{2A}R antagonist SCH 58261 (100 nM) during the washout of
CGS21680 (50 nM) did not impair the long-lasting potentiation, whereas continuous bath
application of SCH 58261 (100 nM) abolished the CGS21680-induced potentiation
(**Figure 3E**). These results indicate that activation of presynaptic A_{2A}Rs was sufficient to
induce LTP at MC-GC synapses.

Because the A_{2A}R is a G_s-coupled receptor, we next tested whether CGS21680-induced
LTP was PKA-dependent. Bath application of the selective, cell-permeable PKA inhibitor
myristoylated PKI_{14-22} (1 µM) (Harris et al., 1997) abolished CGS21680-induced LTP,
whereas loading the membrane-impermeable PKA inhibitor PKI_{6-22} (2.5 µM) in GCs via
the recording pipette (Hashimotodani *et al.*, 2017; Skeberdis et al., 2006) had no effect
(**Figure 3F**), suggesting that presynaptic but not postsynaptic PKA activity is required for
the A_{2A}R agonist-mediated strengthening of MC-GC synaptic transmission. While deleting
A_{2A}Rs from MCs blocked both GC TBF-LTP (**Figure 2E and 2F**) and CGS21680-LTP
(**Figure 3D**), we found that forskolin induced normal LTP in presynaptic *Adora2a* cKO
mice (**Figure 3G**). Altogether, these findings indicate that presynaptic A_{2A}R activation
enhances glutamate release via presynaptic PKA activation, consistent with previous

results showing that PKA activity is necessary and sufficient for MC-GC LTP (**Figure 1E-**
1G, Figure S1J) (Hashimotodani *et al.*, 2017).

Our results thus far indicate that both BDNF/TrkB and adenosine/$A_{2A}R$ signaling are
involved in MC-GC LTP, and previous work demonstrated that $A_{2A}Rs$ facilitate BDNF
signaling at some synapses (Sebastiao and Ribeiro, 2009b). We therefore sought to
determine potential interactions between these signaling cascades at the MC-GC
synapse. The TrkB antagonist ANA-12 did not affect CGS21680-induced LTP (**Figure**
3H), whereas it blocked TBF-LTP in interleaved experiments (**Figure 3I**; see also **Figure**
S1B). Lastly, we found that pre-application of CGS21680 (50 nM, 15 min) occluded both
BDNF puff-induced and GC TBF-induced LTP (**Figure 3J and 3K**), strongly suggesting
that $A_{2A}R$ agonism, BDNF, and GC TBF strengthen MC-GC synapses via a common
mechanism. These findings, together with our previous results (**Figure 2**), indicate that
presynaptic adenosine/$A_{2A}R$ signaling mediates LTP downstream of BDNF/TrkB signaling
(**Figure 3L**).

Activation of presynaptic A_1Rs dampens MC-GC LTP

So far, our data indicate that the release of endogenous adenosine induces MC-GC LTP
by activating presynaptic $A_{2A}Rs$ (**Figures 2 and 3**). In addition to activating presynaptic
$A_{2A}Rs$, adenosine could also activate presynaptic type 1 adenosine receptors (A_1Rs), $G_{i/o}$-
coupled receptors known to suppress neurotransmitter release. To test the role of A_1Rs in
MC-GC plasticity, we first used the selective A_1R antagonist DPCPX (100 nM). DPCPX
increased MC-GC LTP magnitude (**Figure S6A**) but did not affect basal transmission
(**Figure S6B**), indicating that adenosine acts on A_1Rs to dampen MC-GC LTP induction,
but A_1Rs are not tonically activated by adenosine. In addition, we considered that A_1Rs
could mediate presynaptic LTD (Atwood et al., 2014). However, bath application of the
selective A_1R agonist CCPA (50 nM, 15 min) reduced both MC and MPP-mediated
transmission reversibly as the reduction was washed out with the A_1R antagonist DPCPX
(100 nM) (**Figure S6C**). This transient A_1R-induced depression was associated with a
reversible increase in both PPR and CV (**Figures S6D and S6E**), suggesting a
presynaptic mechanism. Of note, $A_{2A}R$ antagonism blocked TBF-LTP induction, but MC-
GC transmission was not reduced in a long-term manner (**Figure 2A**), indicating that the
net effect of adenosine released during GC TBF is to potentiate MC-GC transmission
through $A_{2A}R$ activation. Individual MC boutons could contain both A_1 and A_{2A} receptors.
To test this possibility, we examined whether responses elicited by minimal stimulation, a
manipulation known to activate one to three MC-GC synapses impinging on GCs
(Hashimotodani *et al.*, 2017) were sensitive to both $A_{2A}R$ and A_1R agonists. We found that
the $A_{2A}R$ agonist CGS21680 (50 nM) potentiated the synaptic responses evoked by
minimal stimulation of MC axons, whereas subsequent application of the A_1R agonist

CCPA (50 nM) significantly reduced these potentiated responses recorded from the same
GC (**Figure S6F and S6G**). These results strongly suggest that functional A_1 and A_{2A}
receptors colocalize in the same MC axon boutons.
351

Anatomical evidence for A_{2A} and A_1 receptors in MC axon terminals

To test whether MC terminals express A_1 and A_{2A} receptors, we performed
immunoelectron microscopy. Immunoparticles for A_1Rs were mainly found presynaptically
in the molecular layer of the dentate gyrus at MC-GC and perforant path (PP)-GC
synapses (**Figures 4A, 4B, 4C and 4D**). Remarkably, using an anti-A_{2A}R antibody
previously validated in A_{2A}R-deficient mice (Quiroz et al., 2009), we found strong A_{2A}R
expression in the presynaptic membrane of asymmetric (presumably glutamatergic)
synapses and very few postsynaptic particles in the IML (**Figures 4E, 4F and 4H**). In
contrast, a quasi-exclusive postsynaptic expression was detected at asymmetric PP-GC
synapses (**Figures 4G, 4I and 4H**). Given the strong projection of MC axons in the IML
(Buckmaster et al., 1996), the presynaptic A_{2A}R labeling most likely arises from MC axon
boutons. While the antibodies were not validated in the DG using immunoelectron
microscopy, these anatomical results are entirely consistent with our functional findings.
Although both A_1 and A_{2A} receptors are expressed at MC axon terminals, only the latter
engages long-lasting synaptic plasticity.

Passive adenosine release from GC is required for presynaptic MC-GC LTP

We next sought to determine the source of adenosine that triggers A_{2A}R-mediated LTP.
Adenosine can be released from neurons upon activity (Latini and Pedata, 2001; Lovatt
et al., 2012; Pons-Bennaceur *et al.*, 2019; Wall and Dale, 2013) via equilibrative
nucleoside transporters (ENTs) (King *et al.*, 2006). If adenosine is the retrograde signal
mediating MC-GC LTP, interfering with adenosine release from GCs, by blocking ENTs,
should impair LTP. As a first approach, we tested whether blocking ENTs can interfere
with LTP induction. Bath application of the ENT inhibitors dipyridamole (20 µM) and
NBMPR (10 µM) abolished LTP (**Figure 5A**) but did not affect basal MC-GC synaptic
transmission (**Figure 5B**). Because ENTs are also implicated in adenosine reuptake, bath
application of the ENT blockers may significantly increase extracellular adenosine levels,
which, by activating A_{2A}Rs may potentiate MC-GC transmission and occlude LTP. Further,
the fact that ENT blockers did not affect basal MC-GC synaptic transmission (**Figure 5B**)
could be due to the simultaneous activation of A_1R, which might mask any potential A_{2A}R-
mediated potentiation. To test this possibility, we bath applied both the ENT blockers and
DPCPX (100 nM) to prevent a potential A_1R-mediated depression of MC-GC synaptic
transmission. We found that co-application of the ENT blockers and DPCPX (100 nM)
increased MC-GC transmission, and this effect was abolished in the presence of the A_{2A}R

antagonist SCH 58261 (100 nM) (**Figure 5C**). These results indicate that endogenous adenosine can activate presynaptic A_{2A}Rs to potentiate MC-GC synaptic transmission.

To directly test whether postsynaptic adenosine release is required for MC-GC LTP, we employed a single-cell approach and selectively blocked ENTs in a single GC. Like previous studies (Lovatt *et al.*, 2012; Pons-Bennaceur *et al.*, 2019), we loaded inosine (100 µM), a competitive blocker of adenosine efflux through ENTs (King *et al.*, 2006), intracellularly *via* the recording pipette. We found that intracellular inosine abolished TBF-LTP (**Figure 5D**), whereas bath application of 100 µM inosine had no effect on MC-GC synaptic transmission (**Figure 5E**). Bath application of DPCPX increased MC EPSC amplitude when ENT inhibitors were continuously bath applied but not when inosine (100 µM) was loaded in a single GC (**Figure 5F**), confirming that intracellular inosine did not increase adenosine tone. Intracellular accumulation of adenosine can result from sequential dephosphorylation of ATP (Arch and Newsholme, 1978). Consistent with this possibility, removing ATP from the internal recording solution abolished LTP (**Figure 5G**). Altogether these findings indicate that adenosine, passively released from GC through ENTs, acts as a retrograde messenger at MC-GC synapses (**Figure 5H**).

Repetitive neuronal activity induces adenosine release via a BDNF/TrkB-dependent mechanism

To visualize adenosine release during MC-GC LTP induction, we utilized the genetically encoded sensor for adenosine GRAB$_{Ado1.0m}$ (Peng et al., 2020; Wu et al., 2023), which was selectively expressed in commissural MC axons (see Methods) and generated a clear fluorescence signal in the contralateral IML (**Figures 6A, 6B and 6D**). The MC burst stimulation protocol that triggers synaptically-induced MC-GC LTP in acute hippocampal slices also triggered a robust, albeit transient increase in the GRAB$_{Ado1.0m}$ signal. As expected (Peng *et al.*, 2020), this enhancement was abolished when the MC BS protocol was delivered in the presence of the A_{2A}R antagonist SCH58261 (**Figures 6B, 6C and 6F**). Consistent with A_{2A}R signaling being engaged downstream of TrkB activation, the burst-induced increase of the GRAB$_{Ado1.0m}$-mediated fluorescence was abolished in the presence of the TrkB antagonist ANA-12 (15 µM), and in postsynaptic *TrkB* cKO and *Bdnf* cKO mice (**Figures 6D-6F**). Altogether these results demonstrate that MC repetitive activity releases adenosine in a postsynaptic BDNF/TrkB-dependent manner.

***In vivo* activation of A_{2A}Rs facilitates seizure occurrence**

We recently found that MC-GC LTP can be triggered by seizure activity induced with kainic acid (KA) intraperitoneal (i.p.) administration and that this LTP further promotes seizures (Nasrallah *et al.*, 2022). Given our new findings that neuronal activity releases adenosine via TrkB to induce A_{2A}R-dependent MC-GC LTP, we hypothesized that interfering with

adenosine/A_{2A}R signaling should prevent seizure-induced LTP and reduce KA-induced
seizures. Moreover, seizure activity should release adenosine in a TrkB-dependent
manner. As expected for a presynaptic form of plasticity, seizure-induced LTP increases
neurotransmitter release as indicated by a decrease in PPR (Nasrallah *et al.*, 2022). While
Adora2a deletion from MC axons did not alter basal PPR (**Figure 2G**), it prevented the
KA-induced decrease in PPR (**Figure 7A and 7B**). These results indicate that presynaptic
A_{2A}Rs mediate seizure-induced LTP at MC-GC synapses. Moreover, *Adora2a* deletion
increased latency to KA-induced convulsive seizures (**Figures 7C-7F**), and *TrkB* deletion
increased latency to convulsive seizures and reduced sum seizure score (**Figures 7G and
7H**). These results strongly suggest that both A2ARs and TrkB are proconvulsant,
consistent with the seizure-induced release of adenosine in a TrkB-dependent manner.

Using *in vivo* two-photon imaging in awake mice, we recently reported that KA-induced
seizures trigger a massive increase in both MC and GC activity (Nasrallah *et al.*, 2022).
To test whether seizure activity triggers a TrkB-dependent release of adenosine *in vivo*,
we expressed GRAB$_{Ado1.0m}$ in commissural MC axons and recorded the fluorescence
signals using fiber photometry in freely moving mice (**Figures 7I and 7J**). We found a
large increase in GRAB$_{Ado1.0m}$ fluorescence following i.p. injection of KA, and this
fluorescence signal was significantly reduced in *TrkB* cKO mice (**Figures 7K and 7L**). Of
note, both control and *TrkB* cKO mice reached stage 3 convulsive seizures, discarding
potential failure of seizure induction. Altogether, these results not only demonstrate the
TrkB-mediated *in vivo* release of adenosine following initial seizures, but also strongly
suggest that adenosine signaling may promote seizures via A_{2A}Rs.

DISCUSSION

In this study, we discovered a novel retrograde signaling mechanism that mediates
presynaptic strengthening within a critical hippocampal circuit (**Figure S7**). Specifically,
repetitive firing of a single postsynaptic neuron was sufficient to trigger presynaptic LTP
at MC-GC but not neighboring MPP synapses. We demonstrated that functional A_{2A}Rs
are expressed at MC axon terminals but not at MPP axon terminals and that A_{2A}R
activation is necessary and sufficient to induce MC-GC LTP. During LTP induction GCs
passively release adenosine through ENTs in a postsynaptic TrkB- and activity-dependent
manner. Thus, our findings establish adenosine/A_{2A}R as a novel retrograde signaling
mechanism. Remarkably, our results uncovered that TrkB can signal through adenosine.
Moreover, we found that adenosine is released *in vivo* during initial seizures in a TrkB-
dependent manner and both *Adora2a* and *Trkb* cKO have anti-convulsant effects.

GC TBF induces presynaptic LTP at MC-GC synapses

The present study revealed that direct activation of a single GC with a physiologically relevant pattern of activity (Danielson et al., 2017; Diamantaki et al., 2016; GoodSmith et al., 2017; Henze et al., 2002; Pernia-Andrade and Jonas, 2014; Senzai and Buzsaki, 2017) was sufficient to induce presynaptic LTP at MC-GC synapses without changing the strength of MPP neighboring inputs. Although the activity of a single GC could have induced LTP by recruiting the recurrent GC-MC-GC circuit, the following two observations are against this possibility. First, GC TBF induced normal LTP even in the continuous presence of the selective group II mGluR agonist DCG IV, which abolishes GC-MC (Lysetskiy *et al.*, 2005) but not MC-GC transmission (Chiu and Castillo, 2008). Second, TBF-LTP was normally induced at optogenetically-activated synaptic inputs arising from contralateral MC axons that are cut off from their cell bodies in hippocampal slices. Importantly, TBF-LTP and synaptically-induced LTP likely share a common mechanism (**Figure S2**). Like synaptically-induced MC-GC LTP, which requires BDNF release from GCs (Berthoux *et al.*, 2023), we now report that TBF-LTP requires postsynaptic calcium rise and BDNF release via SNARE-dependent exocytosis (Leschik et al., 2019; Shimojo *et al.*, 2015). While synaptically-induced LTP is an input-specific phenomenon (Hashimotodani *et al.*, 2017), TBF-induced LTP is not, likely reflecting the dissimilar induction protocols. Presynaptic burst stimulation releases adenosine from the postsynaptic compartment of activated synapses only. In contrast, TBF is expected to release adenosine from multiple synapses due to action potential-driven release of BDNF from GCs and autocrine activation of postsynaptic TrkB (Berthoux *et al.*, 2023).

Adenosine is released from GCs in an activity and TrkB-dependent manner

Using the genetically encoded adenosine sensor $GRAB_{Ado1.0m}$ (Peng *et al.*, 2020; Wu *et al.*, 2023), we detected a phasic increase in extracellular adenosine following both MC-GC LTP induction *ex vivo* and during seizure activity *in vivo*. Importantly, adenosine release required intact BDNF/TrkB signaling (**Figure 6D-6F**, **Figure 7K and 7L**). Previous studies have shown that adenosine can be released from neurons upon activity via ENTs (King *et al.*, 2006; Latini and Pedata, 2001; Lovatt *et al.*, 2012; Pons-Bennaceur *et al.*, 2019; Wall and Dale, 2013; Wu *et al.*, 2023). In support of this mechanism, we found that interfering with ENT-mediated release of adenosine from a single postsynaptic GC abolished MC-GC LTP, or not including ATP in the recording pipette, indicating that the passive release of adenosine from GCs is essential for this form of plasticity. Adenosine arising from ATP extracellular conversion (Ribeiro and Sebastiao, 1987; Zimmermann, 2006) or other cell types such as interneurons and glial cells (Latini and Pedata, 2001; Wall and Dale, 2008) could also strengthen MC-GC synaptic transmission.

Intracellular accumulation of adenosine can be the consequence of sequential
dephosphorylation of ATP or the hydrolysis of S-adenosylho-mocysteine (Arch and
Newsholme, 1978). During MC-GC LTP induction, formation of intracellular adenosine in
GCs likely results from robust postsynaptic TrkB activation and sequential
dephosphorylation of ATP through its kinase activity. In support of this scenario are the
following observations. First, postsynaptic ATP is required for MC-GC LTP (**Figure 5F**).
Second, MC-GC LTP induction releases BDNF that activates postsynaptic TrkB on GCs
(Berthoux *et al.*, 2023; Hashimotodani *et al.*, 2017). Third, activity-induced adenosine
release in the IML is postsynaptic BDNF/TrkB-dependent, as it is abolished in the
presence of the TrkB selective antagonist ANA-12 and by genetically removing *Bdnf* and
TrkB from GCs (**Figure 6D-6F and Figure 7I-L**). Fourth, adenosine/A_{2A}R signaling is
engaged downstream of TrkB activation given that BDNF-induced LTP was abolished by
A_{2A}R antagonism (**Figure 2C**), whereas TrkB antagonism did not affect A_{2A}R agonist-
induced LTP (**Figure 3E**). Lastly, while BDNF activates postsynaptic TrkB (Berthoux *et*
al., 2023; Hashimotodani *et al.*, 2017) and adenosine acts via presynaptic A_{2A}Rs to induce
MC-GC LTP, we cannot discard physical interaction or transactivation, a process whereby
A_{2A}R activation can induce TrkB phosphorylation in the absence of neurotrophins (Lee
and Chao, 2001).

To the best of our knowledge, we provided the first direct evidence that adenosine release
can occur in a TrkB-dependent manner. Such a mechanism could explain, at least in part,
BDNF and adenosine signaling interactions reported at other synapses in the brain.
Previous studies in the CA1 area have shown that A_{2A}Rs can affect synaptic transmission
and plasticity by modulating BDNF-TrkB signaling (Diogenes *et al.*, 2004; Fontinha *et al.*,
2008; Rodrigues et al., 2014). Such modulation has recently been implicated in adult
neurogenesis (Ribeiro et al., 2021). While these studies suggest that A_{2A}Rs act upstream
of BDNF, our findings demonstrate that BDNF/TrkB signaling can also act upstream A_{2A}Rs
by promoting adenosine release. Consistent with our findings at the hippocampal MC-GC
synapse, BDNF-induced facilitation at the neuromuscular junction was abolished in
presence of A_{2A}R antagonists but also by PKA inhibitors, while A_{2A}R agonist-induced
increase in neurotransmitter release is TrkB-independent (Correia-de-Sa et al., 1991;
Pousinha et al., 2006; Sebastiao and Ribeiro, 2009b). In conclusion, our findings support
a model whereby neuronal activity releases adenosine in a TrkB-dependent manner, and
therefore, adenosine release emerges as a non-canonical TrkB signaling mechanism.
Further work is required to determine the generalizability of this model in other brain areas.

Retrograde adenosine/A_{2A}R signaling mediates presynaptic LTP

Retrograde signaling is a common mechanism in presynaptic forms of long-term plasticity
(Castillo, 2012; Monday *et al.*, 2018). While several retrograde signals have been
identified throughout the brain (Regehr *et al.*, 2009), here we uncovered adenosine/A_{2A}R
as a novel form of retrograde signaling. Single cell manipulations revealed that interfering
with adenosine release from a single postsynaptic neuron abolished MC-GC LTP, induced
by GC firing alone. Furthermore, by combining pharmacology, presynaptic *Adora2a*
deletion using cKO mice, and immunoelectron microscopy, we demonstrated that
activation of presynaptic A_{2A}Rs was necessary and sufficient for MC-GC LTP. Despite the
relatively low expression levels of A_{2A}R in the hippocampus (Dixon et al., 1996), previous
work has demonstrated that A_{2A}R signaling can regulate hippocampal synaptic plasticity
(Dias et al., 2013; Duster et al., 2014). To our knowledge, our study not only provides the
first evidence of adenosine/A_{2A}R as a retrograde signaling system but also the first
evidence that this signaling system can mediate presynaptic LTP. As G_s-coupled
receptors, and as supported by our findings, A_{2A}Rs on MC axon terminals likely activate
the cAMP/PKA cascade, and as for other forms of presynaptic LTP (Castillo, 2012), this
activation engages a long-lasting increase of glutamate release whose downstream
mechanism is poorly understood. Whether retrograde adenosine/A_{2A}R signaling can
mediate activity-dependent strengthening at other synapses in the brain, including in the
striatum and globus pallidus, where presynaptic A_{2A}Rs are highly expressed (Kase, 2001),
remains to be investigated.
556
557 **Retrograde adenosine/A_1Rs signaling dampens LTP induction**
558
As previously reported at other synapses (Correia-de-Sa *et al.*, 1991; Pousinha et al.,
2010), our findings indicate that postsynaptically released adenosine simultaneously
activates A_1Rs and A_{2A}Rs localized at MC terminals. A_1Rs and A_{2A}Rs can localize at the
same nerve terminal (Correia-de-Sa et al., 1996) and even form heteromers (Sebastiao
and Ribeiro, 2009a). While A_1R agonism transiently inhibited glutamate release, A_{2A}R
agonism increased glutamate release in a long-lasting manner. Consistent with these
pharmacological observations, the net effect mediated by transient release of endogenous
adenosine during LTP induction was a long-lasting enhancement of glutamate release.
We also found A_1R-mediated dampening of LTP which has previously been observed at
other synapses (Alzheimer et al., 1991; Forghani and Krnjevic, 1995). A possible
explanation for this dampening is that presynaptic A_1Rs, which are $G_{i/o}$-coupled, decrease
the level of cAMP, which is required for MC-GC LTP (Hashimotodani *et al.*, 2017). Type 1
cannabinoid receptor, another $G_{i/o}$-coupled receptor that is highly expressed at MC axon
terminals, also dampens MC-GC LTP (Jensen *et al.*, 2021). Thus, the induction of this
form of plasticity is tightly controlled by two distinct retrograde signals, adenosine and
endocannabinoids, which, by activating presynaptic $G_{i/o}$-coupled receptors, dampen LTP

induction. Both retrograde signals may prevent runaway activity of the MC-GC-MC recurrent excitatory circuit.

Adenosine/A_{2A}R retrograde signaling in the dentate gyrus has physiological and pathological relevance

Growing evidence indicates that MCs are critically involved in hippocampal-dependent learning (Scharfman, 2016). For example, silencing MCs selectively impairs the retrieval of spatial memory (Bui *et al.*, 2018) and novelty-induced contextual memory acquisition (Fredes et al., 2021). MC-GC LTP is a robust form of presynaptic plasticity that can be elicited *in vivo* upon experience (Berthoux *et al.*, 2023) and may contribute significantly to learning and memory by changing information flow in the DG (Hashimotodani *et al.*, 2017). While GCs exhibit extremely sparse and selective firing in a single place field (Diamantaki *et al.*, 2016), we found that MC-GC LTP can be triggered by GC firing alone, suggesting that this plasticity could be implicated in experience-driven refinement of DG circuitry and memory. Previous work indicates that adenosine/A_{2A}R signaling may contribute to hippocampal-dependent memory (Duster *et al.*, 2014; Sebastiao and Ribeiro, 2009a). Our findings showing a key role for adenosine/A_{2A}R signaling in MC-GC LTP could explain, at least in part, memory impairments found in both A_{2A}R antagonist-injected animals (Fontinha et al., 2009) and hippocampal *Adora2a* cKO animals (Wei et al., 2014). Demonstrating the precise role of adenosine/A_{2A}R retrograde signaling at MC-GC synapses in memory requires selective manipulation of A_{2A}Rs at MC axon terminals using tools that are not currently available.

Adenosinergic signaling is critically involved in epilepsy (Beamer et al., 2021). Global A_{2A}R genetic deletion and selective A_{2A}R antagonism attenuate both the seizure progression (El Yacoubi et al., 2008; 2009) and seizure-induced neuronal damage (Jones et al., 1998; Lee et al., 2004; Rosim et al., 2011). In contrast, A_{2A}R activation lowers the seizure threshold (Fukuda et al., 2011) and increased A_{2A}R function promotes epilepsy (Sandau et al., 2016). Although the precise mechanism by which A_{2A}Rs participate in temporal lobe epilepsy remains unclear, our current findings provide a potential explanation for the pro-epileptic role of A_{2A}Rs, consistent with our previous findings that initial seizures trigger MC-GC LTP and that uncontrolled LTP is sufficient to promote subsequent seizures (Nasrallah *et al.*, 2022). The BDNF/TrkB-mediated release of adenosine from GCs and the consequent strengthening of MC-GC synaptic transmission may also explain, at least in part, the pro-epileptic role of BDNF/TrkB signaling (McNamara and Scharfman, 2012). Our current study demonstrating adenosine/A_{2A}R-dependent strengthening of MC-GC synapses reveals a novel mechanism whereby MC activity may contribute to early stages of temporal lobe epilepsy.

614
615
616

Acknowledgments

We thank all Castillo lab members for their invaluable feedback. We thank Dr. Anita Autry (Einstein) for sharing her photometry system and for the technical support provided by her lab members, especially Ilaria Carta and Dr. Giovanni Podda., We thank Subrina Persaud for assisting with confocal image acquisition. We thank Dr. Yulong Li (Peking University) for helpful discussions about the adenosine sensor, Dr. Pascal Kaeser (Harvard University) for sharing an AAV-hSyn-Flex-ChIEF-TdTomato plasmid, and Dr. Lisa Monteggia (Vanderbilt University) for sharing *TrkB*$^{fl/fl}$ and *Bdnf*$^{fl/fl}$ mice, and Drs. Gaël Barthet and Christophe Mulle, Université de Bordeaux, for sharing the C1ql2 constructs. This work was supported by NIH grants R01 MH116673, R01MH125772, and R01 NS 113600 to P.E.C.. K.N. was partially supported by a Postdoctoral Research Fellowship (American Epilepsy Society), the Fondation pour la Recherche Médicale (postdoctoral fellowship for research abroad), and the Fondation Bettencourt Schueller (Prix pour les Jeunes Chercheurs 2016). C.B. was partially supported by a Postdoctoral Research Fellowship (American Epilepsy Society). Y.H. was supported by grants from JSPS KAKENHI (20H03358, 23H04240, and 23K18167). A.E.C. was partially supported by a Ruth L. Kirschstein Award (F32 NS071821), a NARSAD Young Investigator Grant from the Brain & Behavior Research Foundation, and by Chilean Fondecyt regular (#1201848). M.G. was supported by a Ruth L. Kirschstein Award (F31MH122134). R.L. was funded by the Spanish MCIN/AEI/ 10.13039/501100011033 and by "ERDF A way of making Europe" (PID2021-125875OB-I00) and Junta de Comunidades de Castilla-La Mancha (SBPLY/21/180501/000064). Confocal images were obtained at the Einstein Imaging Core (supported by the Rose F. Kennedy Intellectual Disabilities Research Center - shared instrument grant 1S10OD25295 to Konstantin Dobrenis).

Author contributions

K.N. and P.E.C. designed studies and wrote the manuscript. K.N. performed experiments and analyzed all data except for the *in vitro* adenosine measurements that were designed, performed, and analyzed by C.B.. R.L. performed and analyzed immunoelectron microscopy experiments. M.G. tested the role of A1 receptors. Y.H and A.C designed, performed, and analyzed early experiments assessing conventional retrograde signals. All authors read and edited the manuscript.

Declaration of interest

The authors declare no competing interest.

650
651
652

MAIN FIGURE LEGENDS

Figure 1: Theta-burst firing of a single GC induces presynaptic LTP at the MC-GC synapse

(A) *Left,* Diagram illustrating the recording configuration. MC and MPP EPSCs were recorded from the same GC and evoked with stimulation electrodes placed in the inner and middle molecular layer, respectively. *Right*, Current clamp recording showing GC theta-burst firing (GC TBF). LTP induction protocol (GC TBF) was composed of 10 bursts at 5 Hz of 5 action potentials at 50 Hz, repeated 4 times every 5 s.

(B) *Left,* Representative traces before (1) and after (2) GC TBF delivery. *Right,* Time-course plot showing that GC TBF induced LTP at MC-GC but not at MPP-GC synapses.

(C) GC TBF-induced LTP was associated with significant reduction in PPR and CV (n = 13 cells). ** $p < 0.01$, *** $p < 0.001$.

(D) LTP was abolished when TrkB was conditionally knocked out from from GCs (Post *TrkB* cKO, *TrkB$^{fl/fl}$* mice injected in the dorsal blade with AAV$_5$.CaMKII.Cre.GFP). LTP was unaffected in control animals (Control, *TrkB$^{fl/fl}$* mice injected in the dorsal blade with AAV$_5$.CaMKII.eGFP).

(E) LTP was normally induced when loading PKI$_{6-22}$ (2.5 µM) in GCs via the recording pipette but completely blocked when the cell-permeable PKA inhibitor PKI$_{14-22}$ myristoylated (1 µM) was bath applied.

(F) Summary bar graph showing the magnitude of GC TBF-induced LTP in the presence of DGC-IV (1 µM), when TrkB was conditionally knocked out from MCs (Pre *TrkB* cKO), when loading the PKI$_{6-22}$ (2.5 µM) in GCs, and in the presence of D-APV (50 µM). LTP was abolished in the presence of the TrkB antagonist ANA-12 (15 µM), when Botox (0.5 µM) was loaded postsynaptically, in postsynaptic BDNF and *TrkB* cKO mice, and during bath application of the PKA inhibitors H89 (10 µM) or myristoylated PKI$_{14-22}$ (1 µM). Time-course summary plots are shown in Figure S1. ** $p < 0.01$, *** $p < 0.001$.

(G) Scheme illustrating the emerging model for the mechanism underlying GC TBF-LTP. GC TBF triggers postsynaptic BDNF release and subsequent TrkB activation in GCs (1). Presynaptic PKA is then engaged downstream of postsynaptic BDNF/TrkB signaling (2), suggesting the requirement of a retrograde signal. Lastly, presynaptic PKA activation resulted in a long-lasting increase in glutamate release (3).

Numbers in parentheses indicate the number of cells. Data are presented as mean ± SEM.

Figure 2: MC-GC LTP requires activation of presynaptic A$_{2A}$Rs

(A) Bath application of the $A_{2A}R$ selective antagonists SCH 58261 (100 nM) and ZM241385 (50 nM) blocked GC TBF-induced LTP.

(B) SCH 58261 (100 nM) did not change basal EPSC amplitude.

(C) SCH 58261 (100 nM) abolished LTP induced by BDNF puffs (8 nM, 2 puffs of 3 s in the IML), as compared with interleaved controls.

(D) *Left,* loading GDPβS in the recording pipette did not affect TBF-induced LTP. *Right,* interleaved, positive control showing that replacing GTP by 1 mM GDPβS in the internal solution efficiently abolished the $GABA_B$ receptor agonist (baclofen, 10 µM)-induced increase in holding current.

(E) *Left,* A mix of AAV_5.CamKII.Cre-mCherry and AAV_{DJ}.hSyn.Flex.ChIEF.Tdtomato was injected unilaterally into the DG of *Adora2a$^{fl/fl}$* (cKO) or WT (control) mice. *Right,* Infrared/differential interference contrast (IR/DIC, top) and fluorescence (bottom) images show that ChIEF-TdTomato was selectively expressed in putative MC axons of contralateral IML.

(F) Light-evoked MC EPSCs were recorded in contralateral DG. MC-GC LTP was abolished in presynaptic *Adora2a* conditional knockout mice as compared with controls.

(G) Basal PPR was similar in control and presynaptic *Adora2a* cKO animals. n.s. $p > 0.05$.,

Numbers in parentheses represent number of cells. Data are presented as mean ± SEM.

Figure 3: $A_{2A}R$ activation is sufficient to trigger PKA-dependent LTP at MC-GC but not at MPP-GC synapses.

(A) Representative traces *(top)* and time-course summary plot *(bottom)* showing that bath application of the $A_{2A}R$ selective agonist CGS21680 (50 nM) potentiated MC-GC but not MPP-GC synaptic transmission.

(B) CGS21680-induced potentiation at MC-GC synapse was associated with a significant reduction of both PPR and CV. ** $p < 0.01$; n = 9 cells.

(C) AAV1-EF1a-DIO-hChR2(H134R)-eYFP was injected into the SuM of VGluT2-Cre mice. Light-evoked SuM EPSCs and electrically-triggered MC EPSCs were monitored in GCs. CGS21680 (50 nM) potentiated MC-GC but not SuM-GC synaptic transmission.

(D) CGS21680-induced potentiation was abolished in presynaptic *Adora2a* cKO mice. A mix of AAV_5.CamKII.Cre-mCherry and AAV_{DJ}.hSyn.Flex.ChIEF.TdTomato was injected

unilaterally into the DG of *Adora2a*$^{fl/fl}$. Light-evoked MC EPSCs were recorded in
contralateral DG.

(E) CGS21680 induced long-lasting potentiation even when the $A_{2A}R$ selective antagonist
SCH 58261 (100 nM) was included during CGS21680 washout. CGS21680-induced LTP
was completely abolished in continuous presence of SCH 58261.

(F) Loading the selective PKA blocker PKI_{6-22} (2.5 µM) in GCs via the recording pipette
did not impair CGS21680-induced LTP while bath application of the cell-permeable PKA
inhibitor PKI_{14-22} myristoylated (1 µM) completely blocked LTP.

(G) Light-evoked MC EPSCs showing that bath application of the adenylyl cyclase
activator forskolin (50 µM) induced LTP in presynaptic *Adora2a* cKO mice.

(H) Bath application of the TrkB antagonist ANA-12 (15 µM) did not impair CGS21680-
induced LTP.

(I) Positive control in interleaved experiments showing that ANA-12 (15 µM) efficiently
blocked GC TBF-induced LTP.

(J, K) Bath application of CGS21680 (50 nM, 15 min) occluded LTP induced with both
BDNF (8 nM, 2 puffs of 3 s in the IML, J) and GC TBF (K).

(L) Cartoon illustrating how activation of presynaptic $A_{2A}Rs$ induces PKA-dependent long-
lasting increase in glutamate (Glu) release. The presynaptic $A_{2A}R$/PKA pathway is
engaged downstream of postsynaptic BDNF/TrkB signaling during LTP induction.

Numbers in parentheses represent number of cells. Data are presented as mean ± SEM.

741
742
743 **Figure 4: Subcellular localization of A$_1$ and A$_{2A}$ receptors in the molecular layer of**
744 **the dentate gyrus.**
745
(A-C, E-G) Electron micrographs of the molecular layer of the dentate gyrus showing
immunoreactivity for A_1Rs and $A_{2A}Rs$ revealed by pre-embedding immunogold methods.

(A-C) Both at MC-GC **(A, B)** and PP-GC **(C)** putative synapses, immunoparticles for A_1R
were mainly observed on the presynaptic plasma membrane (arrowheads) of axon
terminals (at), with very low frequency in postsynaptic sites (arrows) of spines (s) or
dendritic shafts (Den).

(D) Quantitative analysis of the relative number of immunoparticles found in the
presynaptic membrane for A_1R at MC-GC and MPP-GC putative synapses. From the total

number of immunoparticles detected (n = 490 for MC-GC; n = 419 for MPP-GC synapses, N = 3 mice), 468 (95.5%) and 402 (95.9%) were present in presynaptic sites of MC-GC and MPP-GC putative synapses, respectively.

(E-H) Electron micrographs **(E-G)** and quantitative analysis **(H)** showing that, at putative MC-GC synapses **(E, F** and **H)**, immunoparticles for $A_{2A}R$ were mainly detected presynaptically (91.0%) whereas they were mainly found on the postsynaptic plasma membrane (92.0%) of putative PP-GC synapses **(G, H)**. Total number of immunoparticles: n = 668 for MC-GC; n = 690 for MPP-GC synapses, N = 3 mice.

Scale bars: 200 nm. Data are presented as mean ± SEM.

Figure 5: MC-GC LTP requires passive release of adenosine from GCs, via ENTs.

(A) Representative traces *(left)* and time-course summary plot *(right)* showing that GC TBF failed to induce LTP in presence of the ENT blockers (20 µM of dipyridamole and 10 µM of NBMPR) as compared with interleaved controls.

(B) Bath application of the ENT blockers (20 µM of dipyridamole and 10 µM of NBMPR) did not change basal MC-GC EPSC amplitude.

(C) Co-application of the ENT blockers (20 µM of dipyridamole and 10 µM of NBMPR) and the A_1R antagonist DPCPX (100 nM) increased EPSC amplitude in the control condition but not in presence of the $A_{2A}R$ antagonist SCH 58261 (100 nM).

(D) Intracellular loading of inosine (100 µM) via the patch pipette abolished GC TBF-induced LTP.

(E) Time course summary plot (right) and representative traces (left) showing that bath application of inosine (100 µM) did not affect basal EPSC amplitude.

(F) Bath application of DPCPX (100 nM) increased EPSC amplitude when the ENT blockers (20 µM of dipyridamole and 10 µM of NBMPR) were included in the bath but not when inosine (100 µM) was loaded in the postsynaptic neuron via the patch pipette.

(G) GC TBF failed to induce LTP when ATP was removed from the recording solution as compared to controls. To allow for complete intracellular dialysis, LTP induction protocol was applied 25-35 min after break-in.

(H) Scheme summarizing the findings. Adenosine is passively released from GCs via ENTs. Adenosine then activates presynaptic A_1Rs and $A_{2A}Rs$. $A_{2A}R$ activation induces a long-lasting PKA-dependent increase in glutamate (Glu) release, whereas A_1R activation

786 dampens synaptic transmission and LTP.

787 Numbers in parentheses represent number of cells. Data are represented as mean ±
788 SEM.

Figure 6: Induction of MC-GC LTP triggers a transient TrkB-dependent increase in
extracellular adenosine level

(A) AAV_9.hSyn.GRAB.Ado1.0m ($GRAB_{Ado}$) was injected unilaterally in the DG of WT mice
(*top*). Two-photon image (*bottom*) showing $GRAB_{Ado}$ was selectively expressed in putative
MC axons in contralateral IML.

(B, C) Two-photon images of the IML **(B)** showing $GRAB_{Ado}$ fluorescence intensity
increased during burst electrical stimulation of MC axon terminals (MC BS) in normal
ACSF (control) but not in continuous presence of the A_{2A} receptor antagonist SCH 58261
(100 nM). Time-course summary plot of the average fractional fluorescence changes
($\Delta F/F_0$) with time are shown in **C**.

(D, E) Two-photon images of the IML **(D)** and time-course summary plot **(E)** showing how
MC BS failed to increase $GRAB_{Ado}$ fluorescence intensity in the continuous presence of
the TrkB antagonist ANA-12 (15 µM) and when TrkB (Post *TrkB* cKO) or BDNF (Post *Bdnf*
cKO) was conditionally knocked out from GCs.

(F) Quantification of the averaged responses during burst stimulation of MCs (15-25 s) *
$p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; one-way ANOVA.

Number of slices are shown between parentheses. Data are represented as mean ± SEM.

Figure 7: *In vivo* release of adenosine during acute seizures plays a pro-convulsant
role by activating A_{2A}Rs

(A) Experimental timeline**.** A mix of AAV_5.CamKII.Cre-mCherry and
AAV_{DJ}.hSyn.Flex.ChIEF.TdTomato was injected unilaterally into the DG of *Adora2a$^{fl/fl}$*
(*Adora2a* cKO) or WT (control) mice. Seizures were induced using a single KA i.p. (20
mg/kg) injection and acute hippocampal slices were prepared 25 min after KA injection.
Whole-cell recordings were performed from GC in the contralateral DG and MC-GC light-
evoked synaptic responses were monitored.

(B) KA-induced seizures decreased PPR in control but not in presynaptic *Adora2a* cKO
conditions. ** $p < 0.01$.

(C) AAV$_5$-CaMKII-Cre-mCherry was injected bilaterally into ventral and dorsal DG of WT
(control) and *Adora2a$^{fl/fl}$* (cKO) mice. Mouse behavior was assessed for 120 min following
KA (20 mg/kg i.p.) administration.

(D) Confocal images showing the viral expression in the DG of WT (control) and *Adora2a$^{fl/fl}$*
(*Adora2a* cKO) mice.

(E, F) Deletion of *Adora2a* from DG excitatory neurons (*Adora2a$^{fl/fl}$* mice injected with
AAV$_5$-CaMKII-Cre-mCherry) significantly increased latency to convulsive seizures **(E, F),**
but did not affect seizure severity as compared to control animals**.** Note the shift in seizure
score time course toward the right **(E)**. ** $p < 0.01$, n.s. $p > 0.05$.

(G, H) AAV-CaMKII-eGFP (control) or AAV-CaMKII-Cre-GFP (cKO) was injected
bilaterally into ventral and dorsal DG of *TrkB$^{fl/fl}$* mice. Behavioral seizures were
monitored and scored for 120 min **(G)**. Deletion of *TrkB* from hippocampal excitatory
neurons induced significant increase in latency to convulsive seizures **(E)** and a
decrease in sum score **(H)** and as compared with controls. * $p < 0.05$.

(I) Experimental timeline. AAV$_9$.hSyn.GRAB.Ado1.0m (GRAB$_{Ado}$) was injected unilaterally
in the DG of *TrkB$^{fl/fl}$* mice. In the contralateral DG, both control (AAV-CaMKII-mCherry) or
Cre-expressing AAV (AAV-CaMKII-Cre-mCherry, TrkB cKO) was injected and an optic
fiber was implanted above the contralateral IML. Fiber photometry was performed 3-5
weeks later to assess GRAB$_{Ado}$ fluorescence intensity before and after acute seizure
induction with kainic acid (KA, 30 mg/kg i.p.), in control and *TrkB* cKO mice.

(J) Confocal image showing fiber tract and GRAB$_{Ado}$ expression in the contralateral IML.

(K, L) Time-course of a control and a *TrkB* cKO representative experiments **(K)** and
summary histogram **(L)** showing how KA (30 mg/kg i.p.) administration increased the
average fractional fluorescence ($\Delta F/F_0$) of GRAB$_{Ado}$ in control mice, an effect that was
significantly reduced in *TrkB* cKO animals. ** $p < 0.01$, n.s. $p > 0.05$.

Numbers in parentheses represent number of mice. Data are presented as mean ± SEM.
844

STAR METHODS

Experimental Model and Subject Details

P19-P30 Sprague-Dawley rats and P50-P70 C57BL/6, floxed TrkB (*TrkB$^{fl/fl}$*), floxed BDNF (*Bdnf$^{fl/fl}$*), floxed *Adora2a* (*Adora2a$^{fl/fl}$*), and VGluT2-Cre mice (Jackson labs, Slc17a6$^{tm2(cre)Lowl}$/J, stock #016963) mice were used. Males and females were equally used. All animals were group housed in a standard 12 hr light/12 hr dark cycle and had free access to food and water. Animal handling and use followed a protocol approved by the Institutional Animal Care and Use Committee of Albert Einstein College of Medicine, in accordance with the National Institutes of Health guidelines. Animal care and handling for immunoelectron microscopy studies followed a protocol approved by the Institutional Animal Care and Use Committee of Universidad Castilla-La Mancha in accordance with Spanish (RD 1201/2005) and European Union (86/609/EC) regulations. The supramammillary experiments were approved by the Animal Care and Use Committee of Doshisha University and were performed following the committee guidelines. *TrkB$^{fl/fl}$* and *Bdnf$^{fl/fl}$* mice generated by Dr. Luis Parada were kindly donated by Dr. Lisa Monteggia (University of Texas, Southwestern Medical Center). *Adora2$^{fl/fl}$* were obtained from Jackson Laboratory (B6;129-Adora2atm1Dyj/J, Jax 010687).

Hippocampal slice preparation

Acute transverse hippocampal slices were prepared from Sprague-Dawley rats (400 µm thick) and mice (300-µm thick). Animals were anesthetized with isofluorane and euthanized following institutional regulations. The hippocampi were then removed and cut using a VT1200s microslicer (Leica Microsystems Co.) in a cutting solution. Hippocampal slices from rats were prepared using a cutting solution containing (in mM): 215 sucrose, 2.5 KCl, 26 NaHCO$_3$, 1.6 NaH$_2$PO$_4$, 1 CaCl$_2$, 4 MgCl$_2$, 4 MgSO$_4$ and 20 D-glucose. 30 min post-sectioning, the cutting medium was gradually switched to an extracellular artificial cerebrospinal (ACSF) recording solution containing (in mM): 124 NaCl, 2.5 KCl, 26 NaHCO$_3$, 1 NaH$_2$PO$_4$, 2.5 CaCl$_2$, 1.3 MgSO$_4$ and 10 D-glucose. Slices were incubated for at least 40 min in the ACSF solution before recording. Mouse hippocampal slices were prepared using an NMDG-based cutting solution containing (in mM): 93 N-Methyl-d-glucamin, 2.5 KCl, 1.25 NaH$_2$PO$_4$, 30 NaHCO$_3$, 20 HEPES, 25 D-glucose, 2 Thiourea, 5 Na-Ascorbate, 3 Na-Pyruvate, 0.5 CaCl$_2$, 10 MgCl$_2$. These slices were then transferred to 32°C ACSF for 30 min and then kept at room temperature for at least 1h before recording. All solutions were equilibrated with 95% O$_2$ and 5% CO$_2$ (pH 7.4).

Electrophysiology

Recordings were performed at 28 ± 1°C, otherwise stated, in a submersion-type recording chamber perfused at 2 mL/min with ACSF supplemented with the GABA$_A$ and the GABA$_B$ receptor antagonists, picrotoxin (100 µM) and CGP55845 hydrochloride (3 µM), respectively. Whole-cell patch-clamp recordings using a Multiclamp 700A amplifier (Molecular Devices) were obtained from GCs voltage clamped at −60 mV using patch-type pipette electrodes (~3-4 MΩ) containing a potassium-based internal solution (in mM): 135 KMeSO$_4$, 5 KCl, 1 CaCl$_2$, 5 NaOH, 10 HEPES, 5 MgATP, 0.4 Na$_3$GTP, 5 EGTA and 10 D-glucose, pH 7.2 (288-291 mOsm). For IPSC recordings in CA1, whole-cell voltage

clamp recordings were obtained from CA1 pyramidal neurons voltage clamped at V_h= 10
mV using a cesium-based internal solution containing (in mM): 131 cesium gluconate, 8
NaCl, 1 $CaCl_2$, 10 EGTA, 10 D-glucose and 10 HEPES, pH 7.2 (285-290 mOsm). IPSCs
were evoked using an electrical stimulating pipette placed in CA1 *stratum radiatum*, and
recordings were performed in the continuous presence of ionotropic glutamate receptor
antagonists NBQX (10 µM) and D-APV (50 µM). LTD of inhibitory inputs onto CA1
pyramidal neurons (iLTD) was induced using theta burst-stimulation protocol (TBS, 10
bursts at 5 Hz of 5 pulses at 100 Hz, repeated every 5 s, 4 times). Series resistance (~6-
25 MΩ) was monitored throughout all experiments with a −5 mV, 80 ms voltage step, and
cells that exhibited a significant change in series resistance (> 15%) were excluded from
the analysis. Botox experiments were performed at 32°C. Botox was supplemented with
5 mM dithiothreitol (DTT) in the intracellular solution, and interleaved control experiments
included DTT only.
904
A broken tip (~10–20 µm) stimulating patch-type micropipette filled with ACSF was placed
in the inner molecular layer (IML, < 50 µm from the border of the GC body layer) to activate
MC axons, and in the middle third of the molecular layer to activate MPP inputs. For
minimal stimulation (**Figure S6F and S6G**), stimulating pipettes were made from theta
glass capillaries. To elicit synaptic responses, paired, monopolar square-wave voltage or
current pulses (100–200 µs pulse width, 4-27 V) were delivered through a stimulus isolator
(Digitimer DS2A-MKII). Typically, stimulation intensity was adjusted to obtain synaptic
responses comparable in amplitude across experiments; e.g., 30-70 pA EPSCs (V_h −60
mV). For the optogenetic experiments, EPSCs were evoked using 1-3 ms pulses of blue
(470-nm) light, provided by a collimated LED (Thorlabs, M470L3-C5, 470 nm, 300mW)
and delivered through the microscope objective (40X, 0.8 NA). GC TBF-induced LTP was
typically induced with 10 bursts at 5 Hz of 5 AP at 50 Hz, repeated 4 times every 5s, while
the membrane potential was held at −60 mV in current-clamp mode. Extracellular field
excitatory postsynaptic potentials (fEPSPs) were recorded using patch-type pipettes filled
with 1M NaCl, and placed in *CA1 stratum radiatum*. fEPSPs were evoked with an ACSF-
containing broken tip patch pipette in *CA1 stratum radiatum*. LTP at CA3-CA1 synapses
was triggered using high frequency stimulation protocol (4 HFS: 100 pulses at 100 Hz
repeated 4 times every 10 s) and recordings were performed at 25 ± 1°C. LTD was
induced with low-frequency stimulation protocol (LFS: 900 pulses at 1 Hz).
924
Electrophysiological data were acquired at 5 kHz, filtered at 2.4 kHz, and analyzed using
custom-made software for IgorPro (Wavemetrics Inc.). Paired-pulse ratio (PPR) was
defined as the ratio of the amplitude of the second EPSC (baseline taken 1-2 ms before
the stimulus artifact) to the amplitude of the first EPSC. Coefficient of variation (CV) was
calculated as the standard deviation of EPSC amplitude divided by mean EPSC
amplitude. Both PPR and CV were measured 10 min before and 20–30 min after LTP
induction protocol or CGS21680-induced potentiation. PPR and CV were measured over
minutes 0-6 before, 9-15 min after, and 29-35 min after CCPA application. The magnitude
of LTP/LTD was determined by comparing 10 min baseline responses with responses 20-
30 min (or 30-40 min for **Figure 1B**, 40-50 min for **Figure 2B**, 45-55 min for **Figure 2D**)
after induction protocol. Averaged traces include 20 consecutive individual responses.

Postsynaptic *TrkB* and *Bdnf* conditional KO

Adeno-associated virus AAV_5.CamKII.eGFP (control virus) or AAV_5.CamKII.GFP-CRE (University of Pennsylvania Vector Core) was injected (1 µL at a flow rate of 0.1 µL/min) unilaterally into the dorsal blade of the dentate gyrus (2.06 mm posterior to bregma, 1.5 mm lateral to bregma, 1.8 mm ventral from dura) of *TrkB*$^{fl/fl}$ or *Bdnf*$^{fl/fl}$ mice (4-5 week old). Animals were placed in a stereotaxic frame and anesthetized with isoflurane (up to 5% for induction and 1%–3% for maintenance). Slices for electrophysiology were prepared from injected animals 3–5 weeks after injection. For each animal, the absence of GFP-expressing cells in the hilus of the entire ipsilateral hippocampus was verified. We previously reported that this strategy can target GCs almost exclusively, as indicated by the lack of labeling in the hilus (Berthoux *et al.*, 2023; Hashimotodani *et al.*, 2017; Nasrallah *et al.*, 2022). To conditionally KO *TrkB* from GCs, we also injected a control (ChIEFtom2A-GFP) or a Cre-expressing lentivirus selective for GCs (ChIEFtom2A-Cre, Post TrkB cKO) into the DG (relative to bregma: 1.9 mm posterior, 1.25 mm lateral, 2.3 ventral) of *TrkB*$^{fl/fl}$ mice (see **Figure S1C**). The lentivirus encoding ChIEFtom2A-Cre or ChIEFtom2A-GFP was under the control of the GC-specific C1ql2 promoter (Barthet *et al.*, 2018), which was successfully used in our lab (Monday *et al.*, 2022). In all cases, MC EPSCs were monitored in glowing GCs. C1ql2-ChIEFtom2A-GFP or C1ql2-ChIEFtom2A-Cre plasmids were a gift of Drs. Gaël Barthet and Christophe Mulle, Université de Bordeaux, and lentiviruses were made at the Einstein Genetic Engineering and Gene Therapy Core.

Presynaptic *TrkB* and *Adora2a* conditional KO

A mix (1:2 ratio, 1.2 µL at 0.1 µL/min) of Cre recombinase-containing AAV (AAV_5.CaMKII.Cre-mCherry, UNC Vector) and Cre-dependent ChIEF (AAV_{DJ}.Flx.ChIEF.TdTomato) was injected into the dentate gyrus (relative to bregma: 1.9 mm posterior, 1.25 mm lateral, 2.3 ventral) of adult *TrkB*$^{fl/fl}$, *Adora2a*$^{fl/fl}$ or WT control mice (4-5 week old). We then performed electrophysiology experiments 4–5 weeks post-injection in contralateral hippocampal slices. This allowed us to optically activate Cre-expressing MC axons.

Optogenetic activation of supramammillary inputs

A beveled glass capillary pipette connected to a microsyringe pump (UMP3, WPI) was used for viral injection. 200 nL (50 nL/min) of AAV1-EF1a-DIO-hChR2(H134R)-eYFP (Addgene: 20298-AAV1) was injected into the SuM (relative to bregma, AP: −2.2 mm, ML: ±0.3 mm, DV: −4.85 mm) of VGluT2-Cre mice. The glass capillary remained at the target site for 5 min before the beginning of the injection and was removed 10 min after the infusion. Channelrhodopsin (ChR2)-expressing SuM axons were activated at 0.05 Hz by a pulse of 470 nm blue light (5 ms duration, 10.5 mW/mm^2) delivered through a 40X objective attached to a microscope using an LED (Mightex).

Pharmacology

Reagents were bath applied following dilution into ACSF from stock solutions stored at −20°C prepared in water, DMSO or ethanol, depending on the manufacturer's

recommendation. BDNF puffs (8 nM, 2.5-3 PSI, 3 s puffs repeated twice, 5 s interval) were
applied using a Picospritzer III (Parker) connected to a broken patch pipette. The tip of the
puffer pipette was positioned above the IML while monitoring MC-GC transmission. For
experiments requiring postsynaptic loading PKI_{6-22}, LTP was induced at least 20 min after
establishing the whole-cell configuration. The time to the LTP induction was matched in
interleaved controls.
987
Electron Microscopy
Mice were anesthetized by intraperitoneal injection of ketamine⁄xylazine (ratio 1:1, 0.1
mL/kg) and transcardially perfused with an ice-cold fixative containing 4%
paraformaldehyde, with 0.05% glutaraldehyde and 15% (v/v) saturated picric acid made
up in 0.1 M phosphate buffer (PB, pH 7.4). Brains were then removed and immersed in
the same fixative for 2 hours or overnight at 4°C. Tissue blocks were washed thoroughly
in 0.1 M PB. Coronal sections (60-μm thick) were cut using a vibratome (Leica V1000).
Pre-embedding immunohistochemical analyses were performed as described previously
(Lujan et al., 1996). Free-floating sections were incubated in 10% (v/v) normal goat serum
(NGS) diluted in Tris-buffered saline (TBS). Sections were then incubated in, 3-5 μg/mL
diluted in TBS containing 1% (v/v) normal NGS, anti-$A_{2A}R$ [guinea pig anti-$A_{2A}R$ polyclonal
(AB_2571656; Frontier Institute co., Japan)] or anti-A_1R [rabbit-anti-A_1R antibody (2
mg/ml; Affinity Bioreagents, Labome, USA)] antibodies, followed by incubation in goat
anti-guinea pig IgG coupled to 1.4 nm gold or in goat anti-rabbit IgG coupled to 1.4 nm
gold (Nanoprobes Inc., Stony Brook, NY, USA), respectively. Sections were postfixed in
1% (v/v) glutaraldehyde and washed in double-distilled water, followed by silver
enhancement of the gold particles with an HQ Silver kit (Nanoprobes Inc.). Sections were
then treated with osmium tetraoxide (1% in 0.1 m phosphate buffer), block-stained with
uranyl acetate, dehydrated in a graded series of ethanol, and flat-embedded on glass
slides in Durcupan (Fluka) resin. Regions of interest were cut at 70-90 nm on an
ultramicrotome (Reichert Ultracut E, Leica, Austria) and collected on single-slot pioloform-
coated copper grids. Staining was performed on drops of 1% aqueous uranyl acetate
followed by Reynolds's lead citrate. Ultrastructural analyses were performed in a Jeol-
1010 electron microscope. Quantitative analysis of the relative abundance of $A_{2A}R$ or A_1R,
in the molecular layer of the dentate gyrus, was performed from 60 μm coronal slices as
described (Lujan *et al.*, 1996), in the area for MC-GC synapses and the area of PP-GC
synapses. For each of the three animals, three samples of tissue were obtained (nine total
blocks). Electron microscopic serial ultrathin sections were cut close to the surface of each
block because immunoreactivity decreased with depth. Randomly selected areas were
captured at a final magnification of 45,000X, and measurements covered a total section
area of ~5000 μm^2. Dendritic shafts, dendritic spines and axon terminals were assessed
for the presence of immunoparticles. The percentage of immunoparticles for $A_{2A}Rs$ or for
A_1Rs at postsynaptic and presynaptic sites was calculated. The perimeter of each axon
terminal was measured in reference areas totaling ~2,000 μm^2. All axon terminals
establishing excitatory synapses were counted and assessed from single ultrathin
sections.
1024
Two-photon live imaging of GRAB$_{Ado}$ in acute hippocampal slices

AAV$_9$.hSyn.GRAB.Ado1.0m (WZ Biosciences Inc) was injected (1 µL at 0.1 µL/min)
unilaterally into the hilus of WT mice or TrkB$^{fl/fl}$ mice (3 - 4 week old). Slices were prepared
3 - 4 weeks post-injection and expression of GRAB$_{Ado1.0m}$ was confirmed in MCs of the
ipsilateral hippocampus for each injected mouse. Hippocampal slices from GRAB$_{Ado1.0m}$-
injected mice were prepared using an ice-cold dissection buffer maintained in 5%
CO$_2$/95% O$_2$ and containing (in mM): 25 NaHCO$_3$, 1.25 NaH$_2$PO$_4$, 2.5 KCl, 0.5 CaCl$_2$, 7
MgCl$_2$, 25 D-glucose, 110 choline chloride, 11.6 ascorbic acid, 3.1 pyruvic acid. Slices
were transferred to 32°C ACSF for 30 min and kept at room temperature for at least 45
min before imaging. Slices were then transferred to an imaging chamber under an Ultima
2-photon microscope (Bruker Corp.) equipped with a 60X NA 1.0 water-immersion
objective and InSight DeepSee laser (Spectra-Physics). A 920-nm laser was used to
excite GRAB$_{Ado1.0m}$, and the emission signal was acquired using a 525–570 nm band-pass
filter. The field of view (512 x 512 pixels per frame) was chosen in the IML of contralateral
hippocampal slices where MC commissural axons projected onto GCs and expressed
GRAB$_{Ado1.0m}$. A broken tip stimulating patch-type micropipette filled with ACSF was
positioned in the IML ~150 µm from the imaging region to activate MC axons. The region
of interest (ROI) was magnified to 2X and at least 40 consecutive images (at 0.25 Hz) as
a baseline using PrairieView 5.4 (Bruker Corp.). Following baseline acquisition, MC burst
stimulation was applied and at least an additional 100 images (at 0.25 Hz) were acquired.
To verify the reactivity of the ROI, adenosine (100 µM) was added at the end of the
imaging session. The fluorescence intensity of the ROI was measured and the $\Delta F/F_0$ of
the GRAB$_{Ado1.0m}$ signal was calculated using ImageJ software. All recordings were
performed at 28 ± 1°C in a submersion-type recording chamber perfused at 2 mL/min with
ACSF.
1050
1051 Seizure induction and monitoring
Epileptic seizures were induced acutely in 2-3–month-old mice. Intraperitoneal (i.p.)
injections of 20-30 mg/kg of kainic acid (KA, HelloBio HB0355) prepared in saline solution
the same day were performed. For behavioral seizure scoring, mice were monitored
during the 120 min post-injection, and behavioral seizures were scored, by an
experimenter blind to condition (control vs Adora2a cKO), using a modified Racine scale
as follows: stage 0: normal behavior, stage 1: immobility and rigidity, stage 2: head
bobbing, stage 3: forelimb clonus and rearing, stage 4: continuous rearing and falling,
stage 5: clonic-tonic seizure, stage 6: death. The maximum Racine score was recorded
every 10 minutes and the cumulative seizure score was obtained by summing these
scores across all 12 bins of the 120 min experiment.
1062
1063 Fiber photometry in freely behaving mice
AAV$_9$.hSyn.GRAB.Ado1.0m (1.2 µL at 0.1 µL/min) was unilaterally injected into the left
DG (relative to bregma: 1.9 mm posterior, 1.25 mm lateral, 2.3 ventral) of TrkB$^{fl/fl}$ mice (5-
6 week old). Contralaterally, AAV$_5$.CamKII.eGFP (control) or AAV$_5$.CamKII.GFP-CRE
(TrkB cKO) was injected (0.5 µL at a flow rate of 0.1 µL/min) into the right DG (1.9 mm
posterior, 1.25 mm lateral, 2.1 mm ventral). This allowed us to locally knockout TrkB from
DG excitatory neurons. An optic fiber (200 µm diameter, NA = 0.37, Neurophotometrics)
was then implanted unilaterally into the right DG above the IML (relative to bregma: 1.9

mm posterior, 1.25 mm lateral, 1.8 mm ventral) to record $GRAB_{Ado1.0m}$ fluorescence *in vivo* where *TrkB* was KO as compared to control. 3-5 weeks after surgery, the fluorescence signal was monitored before and after KA i.p. injection (30 mg/kg). To record $GRAB_{Ado1.0m}$ fluorescence, a three-channel multi-fiber photometry system (Neurophotometrics v1 Ltd) was used. 470 nm and 415 nm out-of-phase excitation lights were bandpass filtered and directed via a 20X objective (power: 30 µW). A single patch cord connected to the optic fiber implant was used to deliver light and collect the emitted fluorescence, which was filtered, and projected on a CMOS camera sensor. The open-source software Bonsai was used for data acquisition (40 frames/s rate). The fluorescence intensity profile of each channel was calculated as the mean pixel value of the region of interest. To calculate $\Delta F/F_0$, the 470 nm-evoked signal was normalized by the isosbestic signal (415 nm) to correct for photobleaching and potential artifacts. F_0 corresponds to the average of the last three minutes (baseline) before KA injection. Optic fiber location was verified *post hoc*, to ensure that the fiber tip was in the ML of the DG. Animals with optic tips located outside of the DG or lacking AAV expression were excluded from the analysis. It is worth noting that the recording area is restricted to the vicinity (~200 µm) under the fiber tip (Pisano et al., 2019).

Post-hoc analysis of AAV expression and optic fiber location
Viral expression and optic fiber location were verified *post hoc*, at the end of the experiments (**Figure 7**). Mice were anesthetized with isoflurane (3-5%) and transcardially perfused with 4% paraformaldehyde (PFA) in 0.1 M sodium phosphate buffer (PBS). 50 µm-thick brain coronal sections were prepared using a DSK Microslicer (DTK-1000), stained with DAPI (1:1000) to label cell nuclei, and mounted with Prolong diamond antifade reagent montant (ThermoFisher) onto microscope slides. A Zeiss LSM 880 Airyscan Confocal microscope with Super-Resolution and ZEN (black edition) software and a 25X oil-immersion objective were used for the acquisition of all images in this study.

Quantification and Statistical Analysis
The normality of distributions was assessed using the Shapiro-Wilk test. In normal distributions, Student's unpaired and paired two-tailed t-tests were used to assess between-group and within-group differences, respectively. In unpaired t-tests, we checked for unequal variance. In cases where variance was unequal between normal distributions, we used a p-value that does not assume equal variance (Welch Correction). One-way ANOVA and One-way ANOVA repeated measure (RM) were used when more than two groups were compared. The non-parametric paired sample Wilcoxon signed rank test and Mann-Whitney's U test were used in non-normal distributions. Statistical significance was set to $p < 0.05$ (∗∗∗ indicates $p < 0.001$, ∗∗ indicates $p < 0.01$, and ∗ indicates $p < 0.05$). All values are reported as the mean ± SEM. Statistical results are summarized in **Table S1**. All experiments included at least three animals per condition. Statistical analysis was performed using OriginPro software (OriginLab).

REFERENCES

Alzheimer, C., Rohrenbeck, J., and ten Bruggencate, G. (1991). Adenosine depresses induction of LTP at the mossy fiber-CA3 synapse in vitro. Brain Res *543*, 163-165. 10.1016/0006-8993(91)91061-5.

Arch, J.R., and Newsholme, E.A. (1978). Activities and some properties of 5'-nucleotidase, adenosine kinase and adenosine deaminase in tissues from vertebrates and invertebrates in relation to the control of the concentration and the physiological role of adenosine. Biochem J *174*, 965-977. 10.1042/bj1740965.

Atwood, B.K., Lovinger, D.M., and Mathur, B.N. (2014). Presynaptic long-term depression mediated by Gi/o-coupled receptors. Trends Neurosci *37*, 663-673. 10.1016/j.tins.2014.07.010.

Barthet, G., Jorda-Siquier, T., Rumi-Masante, J., Bernadou, F., Muller, U., and Mulle, C. (2018). Presenilin-mediated cleavage of APP regulates synaptotagmin-7 and presynaptic plasticity. Nat Commun *9*, 4780. 10.1038/s41467-018-06813-x.

Beamer, E., Kuchukulla, M., Boison, D., and Engel, T. (2021). ATP and adenosine-Two players in the control of seizures and epilepsy development. Prog Neurobiol *204*, 102105. 10.1016/j.pneurobio.2021.102105.

Berthoux, C., Nasrallah, K., Milner, T.A., and Castillo, P.E. (2023). BDNF induces its own release to mediate presynaptic plasticity. bioRxiv 10.1101/2021.12.30.474558.

Botterill, J.J., Lu, Y.L., LaFrancois, J.J., Bernstein, H.L., Alcantara-Gonzalez, D., Jain, S., Leary, P., and Scharfman, H.E. (2019). An Excitatory and Epileptogenic Effect of Dentate Gyrus Mossy Cells in a Mouse Model of Epilepsy. Cell Rep *29*, 2875-2889 e2876. 10.1016/j.celrep.2019.10.100.

Buckmaster, P.S., Wenzel, H.J., Kunkel, D.D., and Schwartzkroin, P.A. (1996). Axon arbors and synaptic connections of hippocampal mossy cells in the rat in vivo. J Comp Neurol *366*, 271-292. 10.1002/(sici)1096-9861(19960304)366:2<270::aid-cne7>3.0.co;2-2.

Bui, A.D., Nguyen, T.M., Limouse, C., Kim, H.K., Szabo, G.G., Felong, S., Maroso, M., and Soltesz, I. (2018). Dentate gyrus mossy cells control spontaneous convulsive seizures and spatial memory. Science *359*, 787-790. 10.1126/science.aan4074.

Castillo, P.E. (2012). Presynaptic LTP and LTD of excitatory and inhibitory synapses. Cold Spring Harb Perspect Biol *4*. 10.1101/cshperspect.a005728.

Chevaleyre, V., and Castillo, P.E. (2003). Heterosynaptic LTD of hippocampal GABAergic synapses: a novel role of endocannabinoids in regulating excitability. Neuron *38*, 461-472. 10.1016/s0896-6273(03)00235-6.

Chiu, C.Q., and Castillo, P.E. (2008). Input-specific plasticity at excitatory synapses mediated by endocannabinoids in the dentate gyrus. Neuropharmacology *54*, 68-78. 10.1016/j.neuropharm.2007.06.026.

Correia-de-Sa, P., Sebastiao, A.M., and Ribeiro, J.A. (1991). Inhibitory and excitatory effects of adenosine receptor agonists on evoked transmitter release from phrenic nerve ending of the rat. Br J Pharmacol *103*, 1614-1620. 10.1111/j.1476-5381.1991.tb09836.x.

Correia-de-Sa, P., Timoteo, M.A., and Ribeiro, J.A. (1996). Presynaptic A1 inhibitory/A2A facilitatory adenosine receptor activation balance depends on motor nerve stimulation paradigm at the rat hemidiaphragm. J Neurophysiol *76*, 3910-3919. 10.1152/jn.1996.76.6.3910.

Danielson, N.B., Turi, G.F., Ladow, M., Chavlis, S., Petrantonakis, P.C., Poirazi, P., and Losonczy, A. (2017). In Vivo Imaging of Dentate Gyrus Mossy Cells in Behaving Mice. Neuron *93*, 552-559 e554. 10.1016/j.neuron.2016.12.019.

Davis, G.W., and Muller, M. (2015). Homeostatic control of presynaptic neurotransmitter release. Annu Rev Physiol *77*, 251-270. 10.1146/annurev-physiol-021014-071740.

Diamantaki, M., Frey, M., Berens, P., Preston-Ferrer, P., and Burgalossi, A. (2016). Sparse activity of identified dentate granule cells during spatial exploration. Elife *5*. 10.7554/eLife.20252.

Dias, R.B., Rombo, D.M., Ribeiro, J.A., Henley, J.M., and Sebastiao, A.M. (2013). Adenosine: setting the stage for plasticity. Trends Neurosci *36*, 248-257. 10.1016/j.tins.2012.12.003.

Diogenes, M.J., Fernandes, C.C., Sebastiao, A.M., and Ribeiro, J.A. (2004). Activation of adenosine A2A receptor facilitates brain-derived neurotrophic factor modulation of synaptic transmission in hippocampal slices. J Neurosci *24*, 2905-2913. 10.1523/JNEUROSCI.4454-03.2004.

Dixon, A.K., Gubitz, A.K., Sirinathsinghji, D.J., Richardson, P.J., and Freeman, T.C. (1996). Tissue distribution of adenosine receptor mRNAs in the rat. Br J Pharmacol *118*, 1461-1468. 10.1111/j.1476-5381.1996.tb15561.x.

Duster, R., Prickaerts, J., and Blokland, A. (2014). Purinergic signaling and hippocampal long-term potentiation. Curr Neuropharmacol *12*, 37-43. 10.2174/1570159X113119990045.

El Yacoubi, M., Ledent, C., Parmentier, M., Costentin, J., and Vaugeois, J.M. (2008). Evidence for the involvement of the adenosine A(2A) receptor in the lowered susceptibility to pentylenetetrazol-induced seizures produced in mice by long-term treatment with caffeine. Neuropharmacology *55*, 35-40. 10.1016/j.neuropharm.2008.04.007.

El Yacoubi, M., Ledent, C., Parmentier, M., Costentin, J., and Vaugeois, J.M. (2009). Adenosine A2A receptor deficient mice are partially resistant to limbic seizures. Naunyn Schmiedebergs Arch Pharmacol *380*, 223-232. 10.1007/s00210-009-0426-8.

Fitzsimonds, R.M., and Poo, M.M. (1998). Retrograde signaling in the development and modification of synapses. Physiol Rev *78*, 143-170. 10.1152/physrev.1998.78.1.143.

Fontinha, B.M., Delgado-Garcia, J.M., Madronal, N., Ribeiro, J.A., Sebastiao, A.M., and Gruart, A. (2009). Adenosine A(2A) receptor modulation of hippocampal CA3-CA1 synapse plasticity during associative learning in behaving mice. Neuropsychopharmacology *34*, 1865-1874. 10.1038/npp.2009.8.

Fontinha, B.M., Diogenes, M.J., Ribeiro, J.A., and Sebastiao, A.M. (2008). Enhancement of long-term potentiation by brain-derived neurotrophic factor requires adenosine A2A receptor activation by endogenous adenosine. Neuropharmacology *54*, 924-933. 10.1016/j.neuropharm.2008.01.011.

Forghani, R., and Krnjevic, K. (1995). Adenosine antagonists have differential effects on induction of long-term potentiation in hippocampal slices. Hippocampus *5*, 71-77. 10.1002/hipo.450050109.

Fredes, F., Silva, M.A., Koppensteiner, P., Kobayashi, K., Joesch, M., and Shigemoto, R. (2021). Ventro-dorsal Hippocampal Pathway Gates Novelty-Induced Contextual Memory Formation. Curr Biol *31*, 25-38 e25. 10.1016/j.cub.2020.09.074.

Fukuda, M., Suzuki, Y., Hino, H., Morimoto, T., and Ishii, E. (2011). Activation of central adenosine A(2A) receptors lowers the seizure threshold of hyperthermia-induced seizure in childhood rats. Seizure *20*, 156-159. 10.1016/j.seizure.2010.11.012.

GoodSmith, D., Chen, X., Wang, C., Kim, S.H., Song, H., Burgalossi, A., Christian, K.M., and Knierim, J.J. (2017). Spatial Representations of Granule Cells and Mossy Cells of the Dentate Gyrus. Neuron *93*, 677-690 e675. 10.1016/j.neuron.2016.12.026.

Harris, T.E., Persaud, S.J., and Jones, P.M. (1997). Pseudosubstrate inhibition of cyclic AMP-dependent protein kinase in intact pancreatic islets: effects on cyclic AMP-dependent and glucose-dependent insulin secretion. Biochem Biophys Res Commun *232*, 648-651. 10.1006/bbrc.1997.6344.

Hashimotodani, Y., Nasrallah, K., Jensen, K.R., Chavez, A.E., Carrera, D., and Castillo, P.E. (2017). LTP at Hilar Mossy Cell-Dentate Granule Cell Synapses Modulates Dentate Gyrus Output by Increasing Excitation/Inhibition Balance. Neuron *95*, 928-943 e923. 10.1016/j.neuron.2017.07.028.

Henze, D.A., Wittner, L., and Buzsaki, G. (2002). Single granule cells reliably discharge targets in the hippocampal CA3 network in vivo. Nat Neurosci *5*, 790-795. 10.1038/nn887.

Jensen, K.R., Berthoux, C., Nasrallah, K., and Castillo, P.E. (2021). Multiple cannabinoid signaling cascades powerfully suppress recurrent excitation in the hippocampus. Proc Natl Acad Sci U S A *118*. 10.1073/pnas.2017590118.

Jinde, S., Zsiros, V., and Nakazawa, K. (2013). Hilar mossy cell circuitry controlling dentate granule cell excitability. Front Neural Circuits *7*, 14. 10.3389/fncir.2013.00014.

Jones, P.A., Smith, R.A., and Stone, T.W. (1998). Protection against hippocampal kainate excitotoxicity by intracerebral administration of an adenosine A2A receptor antagonist. Brain Res *800*, 328-335. 10.1016/s0006-8993(98)00540-x.

Kase, H. (2001). New aspects of physiological and pathophysiological functions of adenosine A2A receptor in basal ganglia. Biosci Biotechnol Biochem *65*, 1447-1457. 10.1271/bbb.65.1447.

King, A.E., Ackley, M.A., Cass, C.E., Young, J.D., and Baldwin, S.A. (2006). Nucleoside transporters: from scavengers to novel therapeutic targets. Trends Pharmacol Sci *27*, 416-425. 10.1016/j.tips.2006.06.004.

Latini, S., and Pedata, F. (2001). Adenosine in the central nervous system: release mechanisms and
extracellular concentrations. J Neurochem *79*, 463-484. 10.1046/j.1471-4159.2001.00607.x.
Lee, F.S., and Chao, M.V. (2001). Activation of Trk neurotrophin receptors in the absence of
neurotrophins. Proc Natl Acad Sci U S A *98*, 3555-3560. 10.1073/pnas.061020198.
Lee, H.K., Choi, S.S., Han, K.J., Han, E.J., and Suh, H.W. (2004). Roles of adenosine receptors in the
regulation of kainic acid-induced neurotoxic responses in mice. Brain Res Mol Brain Res *125*, 76-
85. 10.1016/j.molbrainres.2004.03.004.
Leschik, J., Eckenstaler, R., Endres, T., Munsch, T., Edelmann, E., Richter, K., Kobler, O., Fischer, K.D.,
Zuschratter, W., Brigadski, T., et al. (2019). Prominent Postsynaptic and Dendritic Exocytosis of
Endogenous BDNF Vesicles in BDNF-GFP Knock-in Mice. Mol Neurobiol *56*, 6833-6855.
10.1007/s12035-019-1551-0.
Lisman, J.E. (1999). Relating hippocampal circuitry to function: recall of memory sequences by reciprocal
dentate-CA3 interactions. Neuron *22*, 233-242. 10.1016/s0896-6273(00)81085-5.
Lovatt, D., Xu, Q., Liu, W., Takano, T., Smith, N.A., Schnermann, J., Tieu, K., and Nedergaard, M. (2012).
Neuronal adenosine release, and not astrocytic ATP release, mediates feedback inhibition of
excitatory activity. Proc Natl Acad Sci U S A *109*, 6265-6270. 10.1073/pnas.1120997109.
Lujan, R., Nusser, Z., Roberts, J.D., Shigemoto, R., and Somogyi, P. (1996). Perisynaptic location of
metabotropic glutamate receptors mGluR1 and mGluR5 on dendrites and dendritic spines in the
rat hippocampus. Eur J Neurosci *8*, 1488-1500. 10.1111/j.1460-9568.1996.tb01611.x.
Lysetskiy, M., Foldy, C., and Soltesz, I. (2005). Long- and short-term plasticity at mossy fiber synapses on
mossy cells in the rat dentate gyrus. Hippocampus *15*, 691-696. 10.1002/hipo.20096.
McNamara, J.O., and Scharfman, H.E. (2012). Temporal Lobe Epilepsy and the BDNF Receptor, TrkB. In
Jasper's Basic Mechanisms of the Epilepsies, th, J.L. Noebels, M. Avoli, M.A. Rogawski, R.W.
Olsen, and A.V. Delgado-Escueta, eds.
Monday, H.R., Kharod, S.C., Yoon, Y.J., Singer, R.H., and Castillo, P.E. (2022). Presynaptic FMRP and
local protein synthesis support structural and functional plasticity of glutamatergic axon terminals.
Neuron *110*, 2588-2606 e2586. 10.1016/j.neuron.2022.05.024.
Monday, H.R., Younts, T.J., and Castillo, P.E. (2018). Long-Term Plasticity of Neurotransmitter Release:
Emerging Mechanisms and Contributions to Brain Function and Disease. Annu Rev Neurosci *41*,
299-322. 10.1146/annurev-neuro-080317-062155.
Nasrallah, K., Frechou, M.A., Yoon, Y.J., Persaud, S., Goncalves, J.T., and Castillo, P.E. (2022). Seizure-
induced strengthening of a recurrent excitatory circuit in the dentate gyrus is proconvulsant. Proc
Natl Acad Sci U S A *119*, e2201151119. 10.1073/pnas.2201151119.
Normandin, M., Gagne, J., Bernard, J., Elie, R., Miceli, D., Baudry, M., and Massicotte, G. (1996).
Involvement of the 12-lipoxygenase pathway of arachidonic acid metabolism in homosynaptic
long-term depression of the rat hippocampus. Brain Res *730*, 40-46. 10.1016/0006-
8993(96)00428-3.
Peng, W., Wu, Z., Song, K., Zhang, S., Li, Y., and Xu, M. (2020). Regulation of sleep homeostasis
mediator adenosine by basal forebrain glutamatergic neurons. Science *369*.
10.1126/science.abb0556.
Pernia-Andrade, A.J., and Jonas, P. (2014). Theta-gamma-modulated synaptic currents in hippocampal
granule cells in vivo define a mechanism for network oscillations. Neuron *81*, 140-152.
10.1016/j.neuron.2013.09.046.
Pisano, F., Pisanello, M., Lee, S.J., Lee, J., Maglie, E., Balena, A., Sileo, L., Spagnolo, B., Bianco, M.,
Hyun, M., et al. (2019). Depth-resolved fiber photometry with a single tapered optical fiber implant.
Nat Methods *16*, 1185-1192. 10.1038/s41592-019-0581-x.
Pons-Bennaceur, A., Tsintsadze, V., Bui, T.T., Tsintsadze, T., Minlebaev, M., Milh, M., Scavarda, D.,
Giniatullin, R., Giniatullina, R., Shityakov, S., et al. (2019). Diadenosine-Polyphosphate Analogue
AppCH2ppA Suppresses Seizures by Enhancing Adenosine Signaling in the Cortex. Cereb Cortex
29, 3778-3795. 10.1093/cercor/bhy257.
Pousinha, P.A., Correia, A.M., Sebastiao, A.M., and Ribeiro, J.A. (2010). Predominance of adenosine
excitatory over inhibitory effects on transmission at the neuromuscular junction of infant rats. J
Pharmacol Exp Ther *332*, 153-163. 10.1124/jpet.109.157255.

Pousinha, P.A., Diogenes, M.J., Ribeiro, J.A., and Sebastiao, A.M. (2006). Triggering of BDNF facilitatory
1275   action on neuromuscular transmission by adenosine A2A receptors. Neurosci Lett *404*, 143-147.
1276   10.1016/j.neulet.2006.05.036.
1277   Quiroz, C., Lujan, R., Uchigashima, M., Simoes, A.P., Lerner, T.N., Borycz, J., Kachroo, A., Canas, P.M.,
1278   Orru, M., Schwarzschild, M.A., et al. (2009). Key modulatory role of presynaptic adenosine A2A
1279   receptors in cortical neurotransmission to the striatal direct pathway. ScientificWorldJournal *9*,
1280   1321-1344. 10.1100/tsw.2009.143.
1281   Ratzliff, A., Santhakumar, V., Howard, A., and Soltesz, I. (2002). Mossy cells in epilepsy: rigor mortis or
1282   vigor mortis? Trends Neurosci *25*, 140-144. 10.1016/s0166-2236(00)02122-6.
1283   Regehr, W.G., Carey, M.R., and Best, A.R. (2009). Activity-dependent regulation of synapses by
1284   retrograde messengers. Neuron *63*, 154-170. 10.1016/j.neuron.2009.06.021.
1285   Ribeiro, F.F., Ferreira, F., Rodrigues, R.S., Soares, R., Pedro, D.M., Duarte-Samartinho, M., Aroeira, R.I.,
1286   Ferreiro, E., Valero, J., Sola, S., et al. (2021). Regulation of hippocampal postnatal and adult
1287   neurogenesis by adenosine A(2A) receptor: Interaction with brain-derived neurotrophic factor.
1288   Stem Cells *39*, 1362-1381. 10.1002/stem.3421.
1289   Ribeiro, J.A., and Sebastiao, A.M. (1987). On the role, inactivation and origin of endogenous adenosine at
1290   the frog neuromuscular junction. J Physiol *384*, 571-585. 10.1113/jphysiol.1987.sp016470.
1291   Rodrigues, T.M., Jeronimo-Santos, A., Sebastiao, A.M., and Diogenes, M.J. (2014). Adenosine A(2A)
1292   Receptors as novel upstream regulators of BDNF-mediated attenuation of hippocampal Long-
1293   Term Depression (LTD). Neuropharmacology *79*, 389-398. 10.1016/j.neuropharm.2013.12.010.
1294   Rosim, F.E., Persike, D.S., Nehlig, A., Amorim, R.P., de Oliveira, D.M., and Fernandes, M.J. (2011).
1295   Differential neuroprotection by A(1) receptor activation and A(2A) receptor inhibition following
1296   pilocarpine-induced status epilepticus. Epilepsy Behav *22*, 207-213. 10.1016/j.yebeh.2011.07.004.
1297   Sandau, U.S., Colino-Oliveira, M., Jones, A., Saleumvong, B., Coffman, S.Q., Liu, L., Miranda-Lourenco,
1298   C., Palminha, C., Batalha, V.L., Xu, Y., et al. (2016). Adenosine Kinase Deficiency in the Brain
1299   Results in Maladaptive Synaptic Plasticity. J Neurosci *36*, 12117-12128.
1300   10.1523/JNEUROSCI.2146-16.2016.
1301   Scharfman, H.E. (2016). The enigmatic mossy cell of the dentate gyrus. Nat Rev Neurosci *17*, 562-575.
1302   10.1038/nrn.2016.87.
1303   Scharfman, H.E., and Myers, C.E. (2012). Hilar mossy cells of the dentate gyrus: a historical perspective.
1304   Front Neural Circuits *6*, 106. 10.3389/fncir.2012.00106.
1305   Sebastiao, A.M., and Ribeiro, J.A. (2009a). Adenosine receptors and the central nervous system. Handb
1306   Exp Pharmacol, 471-534. 10.1007/978-3-540-89615-9_16.
1307   Sebastiao, A.M., and Ribeiro, J.A. (2009b). Triggering neurotrophic factor actions through adenosine A2A
1308   receptor activation: implications for neuroprotection. Br J Pharmacol *158*, 15-22. 10.1111/j.1476-
1309   5381.2009.00157.x.
1310   Senzai, Y., and Buzsaki, G. (2017). Physiological Properties and Behavioral Correlates of Hippocampal
1311   Granule Cells and Mossy Cells. Neuron *93*, 691-704 e695. 10.1016/j.neuron.2016.12.011.
1312   Shimojo, M., Courchet, J., Pieraut, S., Torabi-Rander, N., Sando, R., 3rd, Polleux, F., and Maximov, A.
1313   (2015). SNAREs Controlling Vesicular Release of BDNF and Development of Callosal Axons. Cell
1314   Rep *11*, 1054-1066. 10.1016/j.celrep.2015.04.032.
1315   Skeberdis, V.A., Chevaleyre, V., Lau, C.G., Goldberg, J.H., Pettit, D.L., Suadicani, S.O., Lin, Y., Bennett,
1316   M.V., Yuste, R., Castillo, P.E., and Zukin, R.S. (2006). Protein kinase A regulates calcium
1317   permeability of NMDA receptors. Nat Neurosci *9*, 501-510. 10.1038/nn1664.
1318   Tabuchi, E., Sakaba, T., and Hashimotodani, Y. (2022). Excitatory selective LTP of supramammillary
1319   glutamatergic/GABAergic cotransmission potentiates dentate granule cell firing. Proc Natl Acad
1320   Sci U S A *119*, e2119636119. 10.1073/pnas.2119636119.
1321   Wall, M., and Dale, N. (2008). Activity-dependent release of adenosine: a critical re-evaluation of
1322   mechanism. Curr Neuropharmacol *6*, 329-337. 10.2174/157015908787386087.
1323   Wall, M.J., and Dale, N. (2013). Neuronal transporter and astrocytic ATP exocytosis underlie activity-
1324   dependent adenosine release in the hippocampus. J Physiol *591*, 3853-3871.
1325   10.1113/jphysiol.2013.253450.

Wei, C.J., Augusto, E., Gomes, C.A., Singer, P., Wang, Y., Boison, D., Cunha, R.A., Yee, B.K., and Chen,
J.F. (2014). Regulation of fear responses by striatal and extrastriatal adenosine A2A receptors in
forebrain. Biol Psychiatry *75*, 855-863. 10.1016/j.biopsych.2013.05.003.
Williams, J.H., Li, Y.G., Nayak, A., Errington, M.L., Murphy, K.P., and Bliss, T.V. (1993). The suppression
of long-term potentiation in rat hippocampus by inhibitors of nitric oxide synthase is temperature
and age dependent. Neuron *11*, 877-884. 10.1016/0896-6273(93)90117-a.
Wu, Z., Cui, Y., Wang, H., Wu, H., Wan, Y., Li, B., Wang, L., Pan, S., Peng, W., Dong, A., et al. (2023).
Neuronal activity-induced, equilibrative nucleoside transporter-dependent, somatodendritic
adenosine release revealed by a GRAB sensor. Proc Natl Acad Sci U S A *120*, e2212387120.
10.1073/pnas.2212387120.
Younts, T.J., Chevaleyre, V., and Castillo, P.E. (2013). CA1 pyramidal cell theta-burst firing triggers
endocannabinoid-mediated long-term depression at both somatic and dendritic inhibitory
synapses. J Neurosci *33*, 13743-13757. 10.1523/JNEUROSCI.0817-13.2013.
Zimmermann, H. (2006). Ectonucleotidases in the nervous system. Novartis Found Symp *276*, 113-128;
discussion 128-130, 233-117, 275-181.
1341

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7