Covid-19 genomic analysis reveals clusters of emerging sublineages within the delta variant

Evans K. Rono[1*, 2a,b,c]

Author details:

[1]*Independent researcher*, 10315 Berlin, Germany: *E-mail: ronoevan@gmail.com

[2]*Former staff as a:*

[a]Research Associate, LabClinic/MNB Health Lab GmbH, 14467 Potsdam, Germany;

[b]Scientific Staff, Vector Biology Unit, Max Plank Institute for Infection Biology, 10117
Berlin, Germany; and,

[c]Lecturer, Pwani University, Department of Biochemistry and Biotechnology, P.O. Box
195-80108 Kilifi, Kenya.

Abstract

The emerging SARS-CoV-2 variants may potentially have enhanced transmissibility and virulence
of the virus, and impacts on performance of diagnostic tools and efficacy of vaccines. Genomic
surveillance provides an opportunity to detect and characterize new mutations early enough for
effective deployment of control strategies. Here, genomic data from Germany and United
Kingdom were examined for genetic diversity by assessing gene mutations and inferring
phylogeny, and simplifying genomic data workflow for faster visualization and interpretation.
Delta variant sublineages were grouped into seven distinct clusters of spike mutations located in
N-terminal domain of S1 region (T95I, D138H, *D142G, Y145H and A222V) and S2 region
(T719I and *N950D). The most predominant cluster was T95I mutation, with the highest
frequencies (71.1% - 83.9%) in Wales, England and Scotland, and the least frequencies (8.9% -
12.1%) in Germany. Two mutations, *D142G and *N950D here described as *reverse mutations
and T719I mutation, were largely unique to Germany. In a month, frequencies of D142G had
increased from 55.6% to 67.8 % in Germany. Additionally, a cluster of D142G+T719I/T mutation
went up from 27.7% to 34.1%, while a T95I+ D142G+N950D/N cluster rose from 19.2% to
26.2%. Although, two distinct clusters of T95I+D138H (2.6% - 3.8%) and T95I+Y145H+A222V
(2.5% - 8.5%) mutations were present in all the countries, they were most predominant in Wales
and Scotland respectively. Results suggest divergent evolutionary trajectories between the clusters
of T95I mutation and those of D142G mutation. These findings provide insights into underlying
dynamics of evolution of the delta variant. Future studies may evaluate the epidemiological and
biological implications of these sublineages.

Introduction

SARS-CoV-2 (Severe acute respiratory syndrome coronavirus type 2) is a coronavirus that caused the Covid-19 disease outbreak in late 2019 in Wuhan China[1–3]. By early 2020, the disease had rapidly spread across the world and was declared a global pandemic[4]. Concurrently, the first Covid-19 genome from Wuhan, which became the official reference genome was published[3]. The genome consists of around 30000 letters of single stranded positive sense RNA molecule[1,5]. The genome codes for four structural proteins: S - spike; E - envelop; M - membrane and N - nucleoprotein, and eight non - structural proteins for RNA replication: Open reading frame (orf)1a, orf1ab; orf3a; orf6; orf7a; orf7b; orf8 and orf10[1].

The global spread of Covid-19 was compounded by emergence of polymorphisms in the coding sequences across its genome, which resulted in new variants of concern (VOC)[6–8]. Delta (B.1.617.2) variant[9] was first reported in Indian in late 2020. It spread globally and effectively outcompeted the alpha, B.1.1.7 variant[8,10–12]. Consequently, the delta variant became the most transmissible and virulent of all the variants that have emerged to date[13,14]. Key amino acid mutations that define the delta variant relative to the Wuhan reference genome include:- **orf1ab:** P4715L, P5401L and G5063S; **S:** T19R, G142D, E156-, F157-, R158G, L452R, T478K, D614G, P681R and D950N; **orf3a:** S26L; **M:** I82T; **orf7a:** V82A and T120I; **orf8:** D119- and F120-; and **N:** R203M and D377Y[15].

Genomic surveillance and open sharing of genomic data[16] has guided the global scientific community to monitor, detect and characterize new variants[6,7,9,17], develop vaccine[5,18], develop and continually review performance of diagnostic tools[19,20] and research on biological implications of the emerging mutations[6,16]. In addition, early detection of emerging Covid-19 mutations is important for monitoring their prevalence and spread for prompt deployment of control measures, as well as designing experiments for assessment of efficacy of vaccines and addressing epidemiological concerns of the emerging variants[12,17].

Here, complete genome sequences for Covid-19 delta variant originating from Germany and United Kingdom (England, Scotland, Northern Ireland and Wales) were characterized for genetic diversity. First, two easier methods for retrieving coding gene sequences and variant calling directly from large datasets of unaligned SARS-CoV-2 complete genome sequences were streamlined for use instead of doing the computationally intensive multiple sequence alignments. These methods were validated using SARS-CoV-2 genome sequences, which were downloaded

from the NCBI GenBank[21] and the GISAID platform[22]. To this end, positions of the mutations in

each variant were renamed with respect to positions of unaligned self (individual) variant, and not

relative to the reference genome (Table 1). The validated methods were applied to analyze a total

of 169315 SARS-CoV-2 complete genome sequences (GISAID submissions from 2021.07.23 to

2021.08.30), and effectively simplifying genomic information of large SARS-CoV-2 genomic

dataset for faster visualization and interpretation.

Results

Spike gene sequences were retrieved from the unaligned genome sequences (N = 169315) and

processed to 95684 high quality sequences by cleaning to remove ambiguous base calls. By

exploiting the spike marker mutations that define each variant, whose positions were renamed in

Table 1, variant calling of 13 different variants was executed. The frequency of the previously

predominant B.1.1.7 alpha variant had significantly reduced to 1%, 991 sequences. The delta

variant, which was called using all the delta markers in Table 1, was the most dominant variant

with 92.4%, 88418 sequences (Fig.1a). However, the total number of all the delta lineages

accounted for 98.2% since 5547 out of 6193 sequences that were categorized as 'other' were of

the delta lineage but without the G142D and/or D950N markers, and therefore had not been called

to any of the 13 variants. This group of sequences had key positions with new mutations at 95,

142, 222, 719 and 950 (Fig. S1a), while the delta variant with 88418 sequences, had mutations at

positions 95,138, 145 and 222 (Fig. S1b). Together, specific positions of new main mutations in

the delta variant were 95, 138, 142, 145 and 222 (Fig.1b). Amino acid spike substitutions at all

these positions were T95I, D138H, D142G, Y145H, A222V, T719I and N950D (Table S1). Of

these, *D142G and/or *N950D are suspected to be *reverse mutation changes from G142D and/or

D950N in the parental delta variant back to the wild type amino acids, which are present in the

Wuhan reference genome[3].

To further interrogate these amino acid substitutions, the genome sequences of the delta variant

were clustered into 6 main spike mutation subgroups (Table S1). For ease of description in this

study, these subgroups were designated as follows; delta, delta2, delta3, delta4, delta5 and delta6

respectively (Table S1). The delta variant (Fig. 1b) was split into five subgroups: Delta

representing the parental delta without T95I, Y145H and A222V mutations (n = 15324, 16.5%);

delta2 with T95I mutations (n = 75307, 81.2%); delta4 with A222V mutations (n = 3749, 4%);

delta5 with Y145H mutations (n = 1664, 1.8%); and delta6 with D138H mutations (n = 1314,

1.4%). The 'other' group (Fig.S1a) was left as delta3 subgroup with D142G reverse mutation (n =

5508, 6%).   A combination of T95I and A222V spike substitutions were detected in delta2, delta4
and delta5. Delta3 with D142G reverse mutations segregated further into two main subgroups
with T95I and A222V mutations. Notably, T719I new mutation was present in delta3. Delta6 had
T95I mutation in which Y145H and A222V sites were conserved.

To reveal the extend of mutation changes in the rest of the SARS-CoV-2 genes, similar analyses
were extended to all the gene coding sequences in each of the six subgroups (Table S1). All the
key mutations that define the parental delta variant in orf1ab, spike, orf3a, M, orf7a, and N genes
were present in all the delta subgroups. The orf6 protein in all the delta subgroups were the most
conserved followed by the E protein. Orf1ab, orf7b, N and orf10 genes showed signatures of new
mutations. Although orf10 protein was the third most conserved gene, it showed emerging
mutation sites at positions L16P in delta2 and T38I in delta3 and delta5. Orf1ab had fixed
substitutions at positions A1306S, P2046L, P2287S, A2529V, V2930L, T3255I, T3646A and
A6319V. New fixed substitutions in orf7b and N genes at positions T40I and G215C respectively
were observed.

Both orf8 and orf7a protein sequences in the reference genome, are 121 amino acid long[3].
However, orf8 and orf7a gene sequences in these delta sublineages were characterized by complex
polymorphisms that included substitutions, deletions (-) and stop codons (!). Some of the orf8
sequences had deletions at positions G66-, S67-, F120- and I121-, and stop codons in many
positions such as Q18!, E19!, and E106!. Majority of orf8 sequences, had mutations at positions
D119I, F120! and I121T at the levels of fixation in the gene, effectively almost replacing deletions
D119- and F120-, which were characteristic of the parental delta variant. In addition, orf8 had the
lowest sequencing coverage of its genome, which forced many sequences (n >896) to be discarded
from the analysis, suggesting that increasing polymorphism in this gene may be responsible for
the low sequencing coverage. In orf7a protein sequences, there were deletions at positions F63-
and V104-, and stop codons in many positions including G38!, Q62!, Q90!, E91!, E92!, Q94! and
E95!.

To check the extend of geographical spread of the individual spike mutations, T95I, D138H,
D142G, Y145H, A222V, T719I and N950D mutations were mapped to their respective countries
(Fig. 2a). As sequencing may not be random and/or standardized across nations, different nations
may have under- and/or over representation of genome sequences. To correct for over- and/or
under representation of genome sequences, frequencies of the mutations were calculated relative
to the total numbers of all the sequences coming from the respective countries.

Delta_ D142G and delta_ N950D mutations were absent in Scotland. Delta_ N950D mutation was

also not detected in Northern Ireland. Delta_T95I mutations were the most prevalent mutation

with highest frequencies being observed in Wales (83.7%), followed by England (81%) and

Scotland (76.9%), while the lowest frequencies (9.1%) were observed in Germany. Interestingly,

the highest frequencies of the delta_ D142G (55.6%) and delta_ N950D (4.3%) 'reverse'

mutations as well as T719I (4.6%) were most prevalent in Germany, suggesting that these

mutations may be driven by selective pressures different from those of T95I mutations in England,

Scotland and Wales.

To understand genetic diversity among these seven delta sublineages, phylogeny was inferred

(Fig. 2b). First, representative sequences for phylogenetic analysis were selected. To do this, all

the genome sequences for each of the seven groups were processed and resolved to haplotype

level (Table S2). From each group, the first ten sequences representing ten of the most abundant

haplotypes (with exception of N950D with only 4 representatives) in each group were selected

(Table S2). Results of maximum likelihood phylogenetic analysis showed seven distinct clusters

of mutations (Fig. 2b). Of these, clusters of delta, delta+T95I, delta+T95I+D138H,

delta+T95I+Y145H+A222V mutations were detected in all the five countries. Signatures of

Delta+D142G+A222V+N950D/N mutations were present in England, Germany and Northern

Ireland. Clusters of Delta+D142G+T719I/T and delta+T95I+D142G+N950D/N mutations were

present in England, Germany, Northern Ireland and Wales. Since the start of the pandemic, the

SARS-CoV-2 has been evolving differently in various jurisdictions worldwide[10].

To track how frequencies of these mutations may have tilted over the succeeding one-month,

similar analyses was done on a new dataset consisting of 214766 complete genome sequences

submitted to the GISAID platform from 2021.08.31 to 2021.09.30. Frequencies of mutations were

compared between the first submission (2021.07.23 to 2021.08.30) and the second submission

(31.08.2021 to 2021.09.30) data sets (Fig. 3a and Fig. 3b). Synonymous mutations at positions

163A, 410I, 856N, 1122V, 1147S and 1264V were observed (Fig. S2a). The same positions of

non-synonymous mutations at positions T95I, D138H, D142G, Y145H and A222V, which were

revealed in the first dataset, were still present in the second dataset (Fig. S2a, Fig. S2b).

Consistently, the T95I cluster of mutations in Wales, England, and Scotland maintained the

highest frequencies in the ranges between 71.2% and 80.8%. Delta+T95I+D142G+N950D/N

mutations in Germany had increased from 19.2% to 26.2%. In addition, delta+D142G+T719I/T in

Germany had also increased from 27.7% and 34.1%. Single delta_D142G mutation, in overall,

increased in frequency from 55.7% (Fig. 2a) to 67.8% (Fig. S3a). In both submissions, England

had the highest number of sequences, while sample size from Northern Ireland in the second

submission suffered significantly from the lowest (N = 76) representation of sequences (Fig. S3b),

which was a drastic drop from 1085 sequences in the first submission (Fig. 2a).

Discussion

Considering good sample sizes of genome sequences analyzed in this study, and the observed

wide spread of these mutations, may suggest that natural selection and not chance events drives

the emergence of these mutations[23]. Mutations T95I, D138H, D142G, Y145H and A222V are

clustered in the N-terminal domain (NTD) in S1 region (Fig. S3b). The T719I position is located

in the S2 region just before the fusion peptide, while N950D is located in the central helix in the

S2 domain[24] (Fig. S2b). Human neutralizing antibody recognizes an epitope of the NTD

suggesting that it has some immunogenic properties[25,26]. Some studies reviewed in[27], have

correlated deletions at positions 69 -70 and other mutations (occurring at positions 141-146 and

242 -244) with prolonged SARS-CoV-2 infections and interference with the neutralizing

antibodies respectively. Mutations at spike involving T95I, was reported in Mu - B.1.621 variant

in Colombia, alongside other mutations located in the NTD (ins146N, Y144T and Y145S), and

receptor binding domain (RBD) (R346K, E484K and N501Y) and S1/S2 cleavage region (P681H)

mutations[28]. Many other VOC variants of interest (VOI) such as Eta - B.1.525, Iota - B.1.526 and

some Kappa – B.1.617.1 also share T95I mutations[15,27]. P681H substitution was also present in

B.I.1.7 alpha variant in which position 145 was deleted[10], suggesting that positions 95 and 145

across the variants, may be under high selective pressures. Worldwide, there has been a

significant reduction in frequencies of many of VOC, including the B.1.1.7 alpha variant. Due to

this, USA recently de-escalated their classification and definition from VOC or VOI to variants

being monitored (VBM)[29].

Indeed, spike protein has been used in vaccine development[5] because it induces neutralizing

antibodies[30]. Delta variant, however, has ability to evade the neutralizing antibodies[31]. Although,

vaccinated people may still get infected with the delta variant, vaccination prevents severe illness

and critical hospitalization[13,32]. Mutations may increase or reduce fitness and adaptiveness[33] by

influencing its transmissibility and virulence[23,34]. Delta variant mutations at T478K and L452R

located in RBD and P681R (Fig. S2b), enhance virus transmissibility[35]. Specifically, P681R

mutation enhances cleavage of the protein at the S1/S2 site[36], while L452R/T478K alter

conformation of the RBD[31] and enhance affinity to bind to mink angiotensin-converting enzyme

2 (ACE2) receptor[31,37], reviewed in[27]. Spike D614G mutation emerged during the early periods of

the pandemic, and was rapidly fixed[23,27]. The D614G enhances the activity of proteases at the

S1/S2 cleavage site[38], suggesting that it works in synergy with P681R mutation to promote higher

rates of virulence[39,40] and efficiency in transmission[41].

In this context, the T95I mutation being the most predominant mutation with a wide geographical

spread, may confer more transmissive ability or fitness and/or adaptiveness to the virus[25].

Evidently, the reducing frequencies of the parental delta variant observed in this survey, may be a

pointer that the parental delta may soon be phased out by its emerging descendants, especially by

T95I and/or *D142G mutations. These T95I and D142G mutations appear to evolve independently

as seen by their clustering with D138H, Y145H+A222V, D142G+N950D/N and D142G+T719I/T.

The unique splitting of delta variant into distinct clusters of emerging delta sublineages may be

hypothesized that the parental delta variant is evolving into new genetic variants. A speculation,

that needs to be tested on the basis of their phenotype differences in transmissibility and/or

epidemiology under real SARS-CoV-2 public health infections. Notably, mutations in N gene and

non- structural genes; orf1ab, orf3a, orf7a and orf8 genes (Table S1), revealed evident signatures

of polymorphic differences, which may have some consequences in viral packaging and

replication. Whether these substitutions are associated with roles of L452R/T478K and/or

D614G/P681R mutations remains unknown. In addition, outstanding questions on adaptive

benefits of the new mutations, and the implications they have on transmissibility, antigenicity, or

virulence of the virus remain to be understood.

In summary, this study has simplified SARS-CoV-2 genomic information for efficient

interpretation, an aspect that may be handy in easing the complexity of visualizing and

interpretating large genomic dataset in routine genomic surveillance and variant diagnosis. The

findings provide insights into the current, and possible future dynamics of evolution of the delta

variant in the face of emerging sublineages under different selective pressures, including those

driven by immune responses in the vaccinated populace. This study was limited to assessing

emergence and characterization of sublineages of the delta variant in a limited geographical region

and time points. Future research may highlight epidemiological and functional impacts of these

clusters of mutations, especially the single mutations that are widespread and are increasing in

frequencies and/or are persisting in the circulation.

Methods

1. Sample size and origin of SARS-CoV-2 genome sequences

A total of 169315 complete SARS-CoV-2 genome sequences from Germany and United Kingdom that were submitted to the GISAID platform from 2021.07.23 to 2021.08.30, and a subsequent, 214766 latest genome sequences, which were submitted from 2021.08.31 to 2021.09.30 were downloaded for analysis in this study from the GISAID platform[22] on 2021.08.30 and 2021.09.30 respectively. These nations and their respective research communities are among the geographical regions, which have invested in genomic surveillance as part of their routine monitoring of the SARS-CoV-2, hence their genomic data are reliable in estimating the actual Covid-19 circulation in their respective territories[45,46]. The downloaded sequences were read to v4.1.1 R software[47] using the readDNAStringSet function of v2.60.2 Biostrings R package[48]. DECIPHER R package v2.20.0[49] was used to Browse and align the sequences.

2. Streamlining the retrieval of SARS-CoV-2 gene sequences

To streamline a faster and easier method to retrieve gene sequences without doing computationally intensive process of SARS-CoV-2 sequence alignments of large sequence datasets with respect to the Wuhan NC_045512 reference genome, patterns of short (between 8 and 40 bases) sequences flanking all the genes, similar length ranges for patterns at the start of the genes, and similar length ranges for patterns at the end of the genes were identified from the Wuhan reference genome[3]. The patterns specific to the flanking regions and those specific to spike gene were used to trim off the spike gene regions out of the nucleotide string sets of the genome sequences using the sub function of R Documentation. For example:

To trim off the flanking region and keep gene sequences but excluding the two patterns, which are not part of the spike gene, a code like this was used:

```
S <- DNAStringSet(sub(".*ACAACTAAACGAACA(.*?)TAAACGAACTTAT.*", "\\1", S))
```

To trim off the flanking regions and keep gene sequences including the two patterns because they are part of the spike gene, a code similar to this was used:

```
S <- DNAStringSet(sub(".*(ATGTTTGTTTTTCTTGT.*?TCAAATTACATTACACA).*", "\\1", S))
```

To trim off the flanking regions and keep gene sequences including the first pattern, which is part of the gene, while at the same time discarding the second pattern, which is not part of the coding gene sequences, a code like this was used:

```
S <- DNAStringSet(sub(".*(ATGTTTGT.*)TAAACGAACT.*","\\1", S))
```

Note that multiple patterns were selected to capture all the sequences, especially gene sequences

with unambiguous nucleotide mutations in regions that match the selected pattern. To clarify this,

a few selected sequences, in which the used patterns did not capture the gene due to mutations,

were filtered and selected for alignment with the complete genome of Wuhan reference sequence.

To inspect and identify additional patterns required to capture all the sequences, the results of

sequence alignment were browsed in the browser.

```
S <- S[width(S) >3819,]
S <- DNAStringSet(c(ref,S))
Salign <- AlignSeqs(S)
BrowseSeq(Salign, highlight = 1)
```

260   The genomic range of the spike gene of the Wuhan reference genome were used to locate and

261   analyze the spike genes from the sequences as follows:

```
refS <- DNAStringSet(substr(ref, start=21563, stop=25384))
Salign2 <- DNAStringSet(substr(Salign, start=21563, stop=25384))
```

Adjustment on this range was made to include immediate flanking regions of the alignment so that

presence of mutations in the flanking regions that render the trimming of the spike gene a failure

was inspected. Using blindly downloaded SARS-CoV-2 complete genome sequences from the

GISAID platform[22] and the NCBI GenBank[21] , all possible patterns were validated for trimming

genome sequences and retrieving the spike gene sequences without having to do multiple

sequence alignments. To this end, the spike gene sequences were successfully retrieved from

thousand complete genomic sequences, while excluding the stop codon at the end of the spike

gene. Widths were used to assess the efficiency of trimming and to check variations in lengths of

the retrieved spike gene sequence delta variant.

```
all_S1 <- all_S[!width(all_S) >3813,]
all_S2 <- all_S[!width(all_S) ==3813,]
all_S3 <- all_S[width(all_S) ==3813,]
all_S4 <- all_S[!width(all_S) <3813,]
```

3. Easing the approach of variant calling

Next, a method for variant calling was also simplified. To do this, the workflow for variant calling

was done by numbering positions of the spike mutation markers that define individual variants

relative to self, instead of the Wuhan reference[3] spike positions (Table 1). The retrieved spike

sequences were processed by cleaning to remove non-nucleotide characters using clean function

of v1.50.0 ShortRead R package[50].

all_S <- clean(all_S)

Cleaned sequences were translated to protein amino acid sequences using translate function of

Biostrings R package[48].

all_Sa <- translate(all_S)

The strings of amino acid sequences were processed for calling the variants in data frame format,

where all the strings of sequences should subsequently be split into individual amino acid letters.

Before converting them to data frame, all the sequences were made to have the same lengths.

Edges from the ends of all the sequences regions were slightly trimmed to retain 1 to 1195,

effectively forcing them to have same lengths. This range was chosen because all the positions for

calling the variants were within this range (Table 1). This was done using substr function of R

Documentation.

Sa <- AAStringSet(substr(Sa, start=1, stop=1195)).

The trimmed sequences were subsequently transformed to data frame dataset.

```
dfSa <- data.frame(width=width(Sa), seq=as.character(Sa), names=names(Sa))
                    #or
dfSa <- data.frame(Sa)
```

294　　To get individual amino acid characters in their respective positions, stringsets of amino acid were

295　　split using stri_extract_all_regex function of v1.7.4 stringi R package[51].

296　　For amino acid, at protein level:

```
dfSa   <- data.frame(stri_extract_all_regex(dfSa$Sa, '.{1,1}'))
```

298　　For codon, at nucleotide level:

```
dfSc   <- data.frame(stri_extract_all_regex(dfSc$Sc, '.{1,3}'))
```

300　　To call the variants by exploiting the positions of the amino acid from the split amino acid

301　　stringsets, key genetic markers specific to each variant (Table 1), were used. For example, below

302　　is the code, which was used to call the parental delta variant:

```
                 dfSa$B.1.617.2_Delta <- ifelse(dfSa$X19=="R" &dfSa$X142=="D" &
                 dfSa$X156=="G&dfSa$X157=="V"&dfSa$X415=="K"&dfSa$X450=="R"&
                 dfSa$X476=="K" & dfSa$X482=="E"& dfSa$X612=="G"& dfSa$X679=="R"&
                 dfSa$X948=="N" ,"yes", "no")
```

This was repeated for each individual variant named in Table 1. Sequences which could not be

classified into any of the thirteen variants were categorized as 'other' for further interrogation.

Data were processed further using v1.4.4 reshape2[52], v1.14.0 data.table[53], v1.3.1 tidyverse[54],

v1.1.3 tidyr[55], v0.6.5 xlsx[56], and v1.4.0 writexl[57] R packages. Plots were visualized using v3.3.5

ggplot2 R package[58] and v0.4.13 circlize R package[59]. Plots were further refined in v1.1.1

Inkscape[60]. Mendeley Deskstop was used as a reference manager for Bibliography[61].

4. *Frequency of codons or amino acids per position*

To summarize frequency matrix of codons or amino acids per position in all the gene sequences,

the excised sequences from the complete genomes were analyzed both at the codon and the amino

acid levels. Delta spike protein sequences have protein lengths of 1271 amino acids, which are

characterized by two key fixed deletion mutations of two codons (amino acids) in the positions

between 156 and 158 (See comment* for explanation below the Table S1). The sequences were

transformed to data frame and split to positions 1 to 1271 of individual amino acids. Note that

these sequences were not aligned relative to the Wuhan reference genome, and as such they were

two-amino acid shorter in lengths than the reference genome. To make them have a full-size of

1273 amino acid long equivalent to the positions in the reference genome, 2 instances of gaps "-"

both at the codon and the amino acid levels were introduced into each sequence at positions 156

and 157 in the data frame of the sequences. This was done as follows:

```
                 Sa$"156" <- "-"
                 Sa$"157" <- "-"
                 Sa<- Sa[, c(1:155,1272,1273,156:1271)]
                 Sa <- data.frame(t(Sa))
                 rownames(Sa) <-1:1273
                 Sa <- data.frame(t(Sa))
```

323 To enable visualization of positions with mutations, the split positions were transformed by

324 transposing the rows (sequences) to columns, which in turn made the columns (amino acid

325 positions) as rows:

```
                 Sa <- data.frame(t(Sa))
```

This offered an opportunity to count in a row wise the codons (nucleotide level) and amino acids

(protein level) competing for each individual position across all the sequences. To do this, unite

function of v1.0.7 dplyr R package[62] was used to unite the rows (codons or amino acids) with

commas as separators.

For instance, consider 180000 as the number of sequences to be analyzed. The unite code would

look like this:

```
Sdelta   <- unite(Sdelta, "seq", c("X1":"X180000"), sep = ",")
```
333

334 To count the frequencies of individual codons or amino acids across all the sequences present per

335 position, a code below was used:

```
Freq.Sdelta   <- cbind(Sdelta, as.data.frame.matrix(
  +   table(
    stack(
      setNames(
        strsplit(as.character(Sdelta$seq), ','), 1:nrow(Sdelta))
      )[2:1])))
Freq.Sdelta$seq = NULL
Freq.Sdelta <- as.data.frame(Freq.Sdelta)
```
336

337 *5. Visualizing positions of mutations*

338 To count the number of codons or amino acids competing for an individual position, a new

339 column was introduced to the above Freq.Sdelta data frame as follows:

```
Freq.Sdelta$Variability   <- rowSums(Freq.Sdelta!=0)
```
340

For a conserved codon or amino acid in a given position, in a total of 180000 sequences, it is

expected that all the 180000 sequences will have same codon or amino acid at this position.

However, if there is a synonymous or non-synonymous mutation at codon level, the maximum

number of sequences, in this case 180000, will be shared by the wild type codon and the new

codon bearing a mutation. At the amino acid level, only non-synonymous mutation will compete

for the position, effectively reducing the total number of 180000 sequences from the wild type

amino acid. As a consequence, no position will have the maximum counts of amino acids to be

180000 sequences in total. To show the most predominant codon and/or amino acid in each

position, which in this case will have highest maximum number of sequences in each position,

new column was introduced by running the following code:

```
Freq.Sdelta$Max_n <- apply(Freq.Sdelta, 1, max)
```
351

To enable visualization in a graph indicating position of the sequence, a new variable 'Position' in

the last column was introduced as:

Freq.Sdelta$Position <- 1:1273

Positions of codons or amino acids, were visualized in a plot showing the number of sequences

'Max_n' against the position 'Position' in the sequence with 'variability' using ggplot2 R

package. To reveal sites that are undergoing mutations, sites with threshold of <99% of the

sequences were labelled. Note that this graph did not discriminate between the fixed mutations and

the conserved sites in the gene. To interrogate amino acid sites that have fixed mutations and/or

are undergoing fixation in the spike sequences (all_Sa) relative to the NC_045512 reference

sequence, the Wuhan wild-type genome was included as the first sequence in the amino acid

sequences dataset. The sequences were browsed using the BrowseSeq function of DECIPHER R

package as mentioned above, while highlighting the reference gene.

BrowseSeq(all_Sa, highlight = 1)

In the final dataset, the reference gene sequences for codons and/or amino acids were included for

comparison with the codons and/or amino acids in the sequences. For codon frequency, codons

used in the sequences were translated to show their respective amino acids, so that synonymous

and non-synonymous codons can easily be identified and compared.

To observe genetic diversity in the remaining genes; Orf1ab, orf3a, E, M, orf6, orf7a, orf7b, orf8,

N and orf10, similar analyses that were done on the spike gene were extended to all the genes.

Note that for the orf1ab gene, the reading was corrected so that part orf1a of the gene joins with

the second part orf1b to give the correct reading frame for the entire orf1ab gene. Therefore,

reading frame of orf1ab gene in all the orf1ab sequences was corrected by running the following

code:

all_orf1ab <- DNAStringSet(str_replace (all_orf1ab, "AAACGGGTT", "AAACCGGGTT"))

6. Grouping of single mutations

Sequences were grouped into groups with different single non-synonymous mutations using the

genetic markers for the delta variant in Table 1 as well as new emerging single amino acid

mutations. By looking at the patterns of these new single mutations at the spike protein sequences,

and to some extend the rest of the genes, further groups consisting of clusters of one or more

combinations of mutations were created, and confirmed by phylogenetic analysis.

7. Phylogenetic analysis

To select representative genomes from 93647 sequences (GISAID submissions from 2021.07.23

to 2021.08.30) for use to infer phylogeny, complete genome sequences were trimmed. The

sequences were trimmed to exclude the sequence portions before the first gene (orf1ab) and the

sequence portions after the last gene (orf10) using the previously described pattern matching

method above. Sequences were cleaned to ensure quality of sequence coverage on the entire

genome. Frequency of identical genomes (haplotypes) were counted from the trimmed sequences,

and those sequences which appeared once were discarded to remain with 27993 genomes.

Frequencies of haplotypes in each mutation category were counted and arranged from the most

dominant to the least dominant haplotypes. From these grouped haplotypes, representative

sequence from each of the first 10 haplotypes per mutation category (except for delta_ N950D,

which had only 4 representatives), were used to infer phylogeny (Table S2).

Phylogeny was inferred using IQTREE maximum likelihood[42] (http://iqtree.cibiv.univie.ac.at/)

applying a GTR+R6 model with 1000 rapid bootstraps. Two similar Wuhan reference genomes (

GenBank ID: MN908947.3 and NC_045512.2)[3] and previously tested delta isolate (GISAID ID:

EPI_ISL_2378732)[43] were included in the analysis to assess the extent of genetic divergence.

Phylogeny annotations, including geographic distributions of clusters of mutations were done

using v3.1.4.991 ggtree R package[63].

8. Clustering, mapping and tracking of mutations

Beyond the haplotypes, clusters of mutations were retraced back to the whole dataset to confirm

the divergence of emerging delta sublineages. These clusters of mutations were mapped to their

respective countries, which the sequences originated from. To detect changes in frequencies of

clusters of mutations that had been identified, the most recent set of genomic data (N = 214766)

submitted to the GISAID platform from the same countries in the previous month from

2021.08.31 to 2021.09.30, were downloaded on 2021.09.30 and analysed in a similar way as

described above. Results were compared with those of genome sequences (N = 169315) from the

submissions of the period from 2021.07.23 to 2021.08.30.

Additional information

Supplementary figures (Fig. S1, S2 and S3) and supplementary tables (Table S1 and S2) are

included at end of the manuscript.

Data availability

Under terms and conditions of use, genome sequences used in this study cannot be circulated here

or elsewhere. Supplementary data files: Data 1 (Fig. 1 and Fig. S1); Data 2 to data 12 (Table S1);

and Data 13 (Table S2, Fig. 2, Fig. 3, Fig. S2 and Fig. S3) are provided. Any other additional data

and methods are available from the author upon request.

Funding

The author did not receive any funding towards this work.

Disclosure statement

The author has no conflict of interest to disclose.

Author contributions

The author conceived the study, designed and validated the methods, downloaded and analyzed

the data, prepared and submitted the manuscript.

Acknowledgements

Many thanks to all the researchers and the five Nations: England, Germany, Northern Ireland,

Scotland and Wales, for investing in SARS-CoV-2 genome sequencing and openly sharing their

genomic data via the GISAID platform. Great appreciation to the NCBI and the GISAID for the

access of the SARS-CoV-2 genomic data.

References

1.   Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N. Engl.*

J. Med. **382**, 727–733 (2020).

2.   Gorbalenya, A. *et al.* The species Severe acute respiratory syndrome-related coronavirus:

classifying 2019-nCoV and naming it SARS-CoV-2. *Nat. Microbiol.* **5**, (2020).

3.   Wu, F. *et al.* A new coronavirus associated with human respiratory disease in China. *Nature*

579, 265–269 (2020).

4.   Cucinotta, D. & Vanelli, M. WHO Declares COVID-19 a Pandemic. *Acta Biomed.* **91**, 157–

160 (2020).

5.   Jamil, S. *et al.* Genetic variants of SARS-CoV-2: What do we know so far? *Am. J. Respir.*

Crit. Care Med. **203**, (2021).

6.   Tegally, H. *et al.* Sixteen novel lineages of SARS-CoV-2 in South Africa. *Nat. Med.* **27**,

440–446 (2021).

7.   Tegally, H. *et al.* Detection of a SARS-CoV-2 variant of concern in South Africa. *Nature*
592, 438–443 (2021).

8.   CDC. Emerging SARS-CoV-2 variants. https://www.cdc.gov/coronavirus/2019-
ncov/science/science-briefs/scientific-brief-emerging-
variants.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F201
9-ncov%2Fmore%2Fscience-and-research%2Fscientific-brief-emerging-variants.html
(2021).

9.   GISAID. Delta variant. https://www.gisaid.org/hcov19-variants/ (2021).

10.   WHO. Tracking SARS-CoV-2 variants. https://www.who.int/en/activities/tracking-SARS-
CoV-2-variants/ (2021).

11.   RKI. Bericht zu Virusvarianten von SARS-CoV-2 in Deutschland, insbesondere zur Variant
of Concern (VOC) B.1.1.7.
rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/DESH/Bericht_VOC_2021-03-
17.pdf?__blob=publicationFile (2021).

12.   Abdool Karim, S. S. & de Oliveira, T. New SARS-CoV-2 Variants — Clinical, Public
Health, and Vaccine Implications. *N. Engl. J. Med.* **384**, 1866–1868 (2021).

13.   Sheikh, A., McMenamin, J., Taylor, B. & Robertson, C. SARS-CoV-2 Delta VOC in
Scotland: demographics, risk of hospital admission, and vaccine effectiveness. *Lancet*
(London, England) vol. 397 2461–2462 (2021).

14.   Fisman, D. N. & Tuite, A. R. Progressive Increase in Virulence of Novel SARS-CoV-2
Variants in Ontario, Canada. *medRxiv* 2021.07.05.21260050 (2021)
doi:10.1101/2021.07.05.21260050.

15.   CoVariants. Variant: 21A (Delta). https://covariants.org/variants/21A.Delta (2021).

16.   Elbe, S. & Buckland-Merrett, G. Data, disease and diplomacy: GISAID's innovative
contribution to global health. *Glob. Challenges* **1**, 33–46 (2017).

17.   ECDC. Sequencing of SARS-CoV-2 - first update.
https://www.ecdc.europa.eu/en/publications-data/sequencing-sars-cov-2 (2021).

18.   WHO. The effects of virus variants on COVID-19 vaccines. https://www.who.int/news-
room/feature-stories/detail/the-effects-of-virus-variants-on-covid-19-
vaccines?gclid=CjwKCAjw-sqKBhBjEiwAVaQ9azeXfBJUkJAMRUAYSG-
Z9mQziqRWzpkDVBD8-wFLoykiLqqng0YBCBoCKN0QAvD_BwE (2021).

19.   Wang, R., Hozumi, Y., Yin, C. & Wei, G. W. Mutations on COVID-19 diagnostic targets.
Genomics **112**, 5204–5213 (2020).

20.   CDC. Why Strain Surveillance is Important for Public Health.

https://www.cdc.gov/coronavirus/2019-ncov/science/science-briefs/scientific-brief-

emerging-

variants.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F201

9-ncov%2Fmore%2Fscience-and-research%2Fscientific-brief-emerging-variants.html

(2021).

21.   NCBI. NCBI. https://www.ncbi.nlm.nih.gov/ (2021).

22.   GISAID. GISAID. https://www.gisaid.org/ (2021).

23.   Lauring, A. & Hodcroft, E. Genetic Variants of SARS-CoV-2—What Do They Mean?

JAMA **325**, (2021).

24.   Lan, J. *et al.* Structure of the SARS-CoV-2 spike receptor-binding domain bound to the

ACE2 receptor. *Nature* **581**, 215–220 (2020).

25.   Liu, L. *et al.* Potent neutralizing antibodies against multiple epitopes on SARS-CoV-2

spike. *Nature* **584**, 450–456 (2020).

26.   Chi, X. *et al.* A neutralizing human antibody binds to the N-terminal domain of the Spike

protein of SARS-CoV-2. *Science* **369**, 650–655 (2020).

27.   Tao, K. *et al.* The biological and clinical significance of emerging SARS-CoV-2 variants.

Nat. Rev. Genet. (2021) doi:10.1038/s41576-021-00408-x.

28.   Laiton-Donato, K. *et al.* Characterization of the emerging B.1.621 variant of interest of

SARS-CoV-2. *Infect. Genet. Evol.* **95**, 105038 (2021).

29.   CDC. SARS-CoV-2 Variant Classifications and Definitions.

https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-

info.html?ACSTrackingID=USCDC_2157-DM66375&ACSTrackingLabel=CDC Updates

SARS-CoV-2 Variant Classifications&deliveryName=USCDC_2157-

DM66375#anchor_1632150752495 (2021).

30.   Wu, K. *et al.* mRNA-1273 vaccine induces neutralizing antibodies against spike mutants

from global SARS-CoV-2 variants. *bioRxiv* 2021.01.25.427948 (2021)

doi:10.1101/2021.01.25.427948.

31.   Baral, P. *et al.* Mutation-induced changes in the receptor-binding interface of the SARS-

CoV-2 Delta variant B.1.617.2 and implications for immune evasion. *Biochem. Biophys.*

Res. Commun. **574**, 14–19 (2021).

32.   Zaveri, L. *et al.* Genomic analysis of SARS-CoV-2 breakthrough infections from Varanasi,

India. *medRxiv* 2021.09.19.21262487 (2021) doi:10.1101/2021.09.19.21262487.

33.   Plante, J. A. *et al.* Spike mutation D614G alters SARS-CoV-2 fitness. *Nature* **592**, 116–121

(2021).

34.   Li, Q. *et al.* The Impact of Mutations in SARS-CoV-2 Spike on Viral Infectivity and
Antigenicity. *Cell* **182**, 1284-1294.e9 (2020).

35.   Starr, T. N., Greaney, A. J., Dingens, A. S. & Bloom, J. D. Complete map of SARS-CoV-2
RBD mutations that escape the monoclonal antibody LY-CoV555 and its cocktail with LY-
CoV016. *Cell Reports Med.* **2**, 100255 (2021).

36.   Peacock, T. P. *et al.* The SARS-CoV-2 variants associated with infections in India, B.1.617,
show enhanced spike cleavage by furin. *bioRxiv* **44**, 2021.05.28.446163 (2021).

37.   Motozono, C. *et al.* SARS-CoV-2 spike L452R variant evades cellular immunity and
increases infectivity. *Cell Host Microbe* **29**, 1124-1136.e11 (2021).

38.   Gobeil, S. M. C. *et al.* D614G Mutation Alters SARS-CoV-2 Spike Conformation and
Enhances Protease Cleavage at the S1/S2 Junction. *Cell Rep.* **34**, (2021).

39.   Becerra-Flores, M. & Cardozo, T. SARS-CoV-2 viral spike G614 mutation exhibits higher
case fatality rate. *Int. J. Clin. Pract.* **74**, (2020).

40.   Korber, B. *et al.* Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases
Infectivity of the COVID-19 Virus. *Cell* **182**, 812-827.e19 (2020).

41.   Hou, Y. J. *et al.* SARS-CoV-2 D614G variant exhibits efficient replication ex vivo and
transmission in vivo. *Science (80-.).* **370**, 1464–1468 (2020).

42.   Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

43.   Saito, A. *et al.* SARS-CoV-2 spike P681R mutation enhances and accelerates viral fusion.
bioRxiv 2021.06.17.448820 (2021) doi:10.1101/2021.06.17.448820.

44.   WHO. WHO announces simple, easy-to-say labels for SARS-CoV-2 Variants of Interest
and Concern. (2021).

45.   GOV.UK. Investigation of SARS-CoV-2 variants of concern: technical briefings.
https://www.gov.uk/government/publications/investigation-of-novel-sars-cov-2-variant-
variant-of-concern-20201201 (2021).

46.   RKI. MF 2: Genome Sequencing and Genomic Epidemiology.
https://www.rki.de/EN/Content/Institute/DepartmentsUnits/MF/MF2/mf2_node.html
(2021).

47.   Team, R. C. R: A language and environment for statistical computing. *R Foundation for*
Statistical Computing, Vienna, Austria (2021).

48.   Pagès H., Aboyoun, P., Gentleman, R. & DebRoy, S. Biostrings: Efficient manipulation of
biological strings. R package version 2.60.2. (2021).

49.   Erik, S. Using DECIPHER v2.0 to Analyze Big Biological Sequence Data in R. *R J.* **8**, 352

(2016).

50.   Morgan, M. *et al.* ShortRead: a bioconductor package for input, quality assessment and
exploration of high-throughput sequence data. *Bioinformatics* **25**, 2607–2608 (2009).

51.   Gagolewski, M. stringi: Fast and portable character string processing in R_. R package
version 1.7.4. (2021).

52.   Wickham, H. Reshaping Data with the reshape Package. *J. Stat. Softw.* **21**, 1–20 (2007).

53.   Srinivasan, M. D. and A. data.table: Extension of \`data.frame\`. R package version 1.14.0.
https://cran.r-project.org/package=data.table%0D (2021).

54.   Wickham, H. *et al.* Welcome to the Tidyverse. *J. Open Source Softw.* **4**, 1686 (2019).

55.   Wickham, H. tidyr: Tidy Messy Data. R package version 1.1.3. (2021).

56.   Arendt, A. D. and C. Read, Write, Format Excel 2007 and Excel 97/2000/XP/2003 Files. R
package version 0.6.5. (2020).

57.   Ooms, J. writexl: Export Data Frames to Excel 'xlsx' Format. R package version 1.4.0.
(2021).

58.   Wickham, H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York,
2016. *Springer-Verlag New York.* (2016).

59.   Gu, Z., Gu, L., Eils, R., Schlesner, M. & Brors, B. circlize implements and enhances
circular visualization in R . *Bioinformatics* **30**, 2811–2812 (2014).

60.   Project, I. Inkscape. https://inkscape.org (2021).

61.   Mendeley. Stay on top of your research. https://www.mendeley.com/reference-
management/mendeley-desktop (2021).

62.   Hadley Wickham, Romain François, L. & Henry, K. M. dplyr: A Grammar of Data
Manipulation. R package version 1.0.7. (2021).

63.   Yu, G. Using ggtree to Visualize Data on Tree-Like Structures. *Curr. Protoc. Bioinforma.*
69, (2020).

Table 1. Spike amino acids and positions relative to individual variant: Genetic markers for variant calling directly from unaligned SARS-CoV-2 complete genome sequences

1	2	3	4	5	6	7	8	9	10	11	12	13
69S	18F	417T	484K	19R	19R	19R	19R	452R	13I	5L	69S	141Y
70G	80A	484K	614G	142D	142D	142D	142D	614G	152C	95I	70G	142H
78D	215G	501Y		156G	156G	156G	156G		452R	253G	95I	143K
80P	242H			157V	157V	157V	157V		614G	253G	253G	481K
145N	243I			450R	415K	415N	450R			477N	477N	498Y
501G	244S			476T	450R	450R	476T			484K	484K	611G
611G	246L			482Q	476K	476K	482Q			614G	614G	678H
570T	414N			612G	482E	482E	612G			701V	701V	1173F
614C	481K			679R	612G	612G	679R				888L	1089K
678H	498Y				679R	679R	948N					1098Y
681A	501G				948N	948N						
713I	611G											
716T	614C											
979A	698V											
982D	701S											
1115H												
1118F												

1 = B.1.1.7 (Alpha); **2** = B.1.351 (Beta); **3** = B.1.1.248 (P1, Gamma); **4** = P.2 (Zeta); **5** = B.1.617.1 (Kappa); **6** = B.1.617.2 (Delta); **7** = B.1.617.2 (Delta Plus); **8** = B.1.617.3; **9** = B.1.427 (Epsilon); **10** = B.1.429 (Epsilon); **11** = B.1.526 (Iota); **12** = B.1.525 (Eta); **13** = P.3 (Theta)[10,44]. Sequences, which could not be called into any of the 13 variants were categorized as 'other' for further interrogation. The initial cases of the variants were first reported in 1 = United Kingdom, 2 = South Africa, 3 = Brazil, 4 = Brazil, 5 = India, 6 = India, 7 = India, 8 = India, 9 = USA, 10 = USA, 11 = USA, 12 = USA/Denmark, and 13 = Philippines[10,44].

575
576

Fig. 1 | Variant calling and revelation of positions of mutations
The total number of sequences were *n* = 93992, sieved from *N* = 169315 by removing non-nucleotide characters from the spike sequences. **a).** Variant-calling using the marker mutations specific to each variant of concern (VOC) in Table 1. Wuhan reference sequence was included as a wild type sequence. The most dominant sequence was the delta variant. By using all the delta markers in Table1, sequences grouped under 'other', did not fall into any of the groups of the variants. **b).** Visualization of amino acid positions of the delta variant from sequences called using the 156- and 157- fixed deletion markers for the delta variant. The variability indicates the number of different amino acid molecules competing for each position. Positions are numbered relative to the Wuhan reference sequence. Each plotted data point represents the total number of sequences sharing the most dominant amino acid in each position. The labeling threshold for position of amino acid was placed at <99% of the total number of sequences. Positions 95, 138, 142, 145, and 222 were revealed to be accumulating mutations.

Fig. 2 | Diversity and clustering of the emerging delta sublineages
a) Mapping geographical distribution of new mutations. Sample sizes for total sequences (N) and delta variant (n) were: England (N = 80443, n = 79674); Germany (N =7128, n = 6235); Northern Ireland (N = 1115, n = 1085); Scotland (N = 5411, n = 5381); Wales (N = 1587, n = 1569) from GISAID submissions from 2021.07.23 to 2021.08.30. The frequencies were determined relative to the total number of sequences from individual country. **b)** Phylogenetic analysis. The tree shows 74 delta variant sequences representing 10 major haplotypes per group in each of the 7 groups with exception of N950D/N, which had 4 representatives. Phylogeny was inferred using IQTREE maximum likelihood using a GTR+R6 model with 1000 rapid bootstraps[42]. Two similar Wuhan reference genomes (GenBank ID: MN908947.3 and NC_045512.2)[3] and one previously tested delta isolate (GISAID ID: EPI_ISL_2378732)[43] were included. Seven wide spread clusters of mutations were evident from the tree.

Fig. 3 | Emergence and spread of delta clusters of mutations
a) Frequencies of cluster of mutations from the sequence batch from 2021.07.23 to 2021.08.30. Sample sizes are as listed in Fig. 2a. **b)** Frequencies of cluster of mutations from the sequence batch from 2021.08.31 to 2021.09.30. Sample sizes for total sequences (N) and delta variant (n) were: England (N = 87668, n = 87195); Germany (N =17847, n = 17596); Northern Ireland (N = 76, n = 75); Scotland (N = 13757, n = 13716); Wales (N = 5339, n = 5303). Frequencies per country were calculated relative to the total (N) number of sequences coming from respective countries.

Additional Information

Supplementary figures and tables

Fig. S1 | Resolving sublineages of the delta variant at positions of mutations
a). Revealing positions of mutations, from the 'other' group in Fig. 1a. The delta variant deletion mutations at positions 156- and 157- were used as conserved markers. Positions 95, 142, 222, 719 and, 950 were revealed as the main hotspot sites for mutations. The total number of sequences were $n = 5547$. Out of these, 5242 and 370 sequences had *reverse mutations at positions *G142D and *N950D respectively. The rest of the mutations in all the sequences within the 'other' delta group were the typical mutations of the delta variant. **b).** Position of mutations in the delta variant sequences called using all the key marker mutations present in the parental delta variant. Positions of mutations 95, 138, 145, and 222 were also observed to be under high selective pressures. The total number of sequences were $n = 88418$. Variability is as was defined in Fig. 1b.

Fig. S2 | Synonymous and non-synonymous mutations in the delta sublineages
a). Revealing positions of mutations at codon level, from the sequence batch submitted from 2021.08.31 to 2021.09.30. Positions 163, 410, 856, 1122 and 1147 are synonymous mutations. The rest of the labelled positions were non-synonymous mutations, which included position 5. **b).** Non-synonymous mutations cluster at the NTD. The total number of the delta sequences were $n = 123867$. The inset on top shows a schematic representation of spike protein showing domains[24] and approximate positions of amino acids: N-terminal domain; NTD, receptor binding domain; RBD, receptor binding motif; RBM, subdomain 1; SD1, subdomain 2; SD2, fusion peptide; FP, heptad repeat 1; HR1, central helix; CH, connector domain; CD, heptad repeat 1; heptad repeat 2; HR2, cytoplasmic tail; CT. Variability is as was defined before.

Fig. S3 | Distribution of single and clusters of mutations
a) Frequencies of single delta mutations from the sequence batch from 2021.08.31 to 2021.09.30. Sample sizes are as listed in Fig. 3b. **b)** Frequencies of cluster of mutations from the sequence submission batch from 2021.08.31 to 2021.09.30. Sample sizes are as listed in Fig. 3b. Frequencies per country were calculated relative to the total (N) number of sequences coming from respective countries.

Table S1. Amino acid mutations in all the coding sequences of 6 subgroups of the delta sublineages

Protein (Length)	Delta key mutations	Delta (16.5%) n = 15324	Delta2 +T95I (81.2%) n = 75307	Delta3 +D142G (6%) n = 5508	Delta4 +A222V (4%) n = 3749	Delta5 +Y145H (1.8%) n = 1664	Delta6 +D138H (1.4%) n = 1314
Spike (1271)	T19R G142D *E156- *F157- *R158G L452R T478K D614G P681R D950N	T19R G142D E156-F157-R158G L452R T478K D614G P681R D950N T22T/I P251P/L D1127D/G A67A/V I1115I/V G1219G/V	T19R G142D E156-F157-R158G L452R T478K D614G P681R D950N T95I D138D/H Y145Y/H A222A/V P1162P/S	T19R D142G E156-F157-R158G L452R T478K D614G P681R D950N/D T29T/A T95T/I A222A/V T250T/I P251P/L T719T/I L5L/F R21R/T H66H/Y G181G/V D253D/A I850I/L D979D/E G1167G/V	T19R G142D E156-F157-R158G L452R T478K D614G P681R D950N A222V V36V/F T95T/I Y145Y/H V1264V/L L5L/F R21R/T D253D/A D979D/E	T19R G142D E156-F157-R158G L452R T478K D614G P681R D950N Y145H T95I A222V V36V/F L5L/F V1264V/L	T19R G142D E156-F157-R158G L452R T478K D614G P681R D950N D138H T95I L5L/F A522A/S T827T/I I1114I/T T29T/A T547T/I T1120T/I G1124G/V
Orf1ab (7096)	P4715L P5401L G5063S	P4715L P5401L G5063S A1306S P2046L P2287S V2930L T3255I T3646A A6319V E87E/D K261K/N D691D/N L5230L/I E5689E/D	P4715L P5401L G5063S A1306S P2046L P2287S V2930L T3255I T3646A A6319V A2529V V665V/I G661G/S T814T/I E1909E/A S2048S/F V2766V/F L3606L/F H5005H/Y D5271D/N/Y	P4715L P5401L G5063S A1306S P2046L P2287S V2930L T3255I T3646A A6319V E87E/D K261K/N P309P/L V665V/I P1640P/L E1724E/D A2529A/V A3209A/V L3606L/F V3718V/A T3750T/I R4589R/Q D5216D/Y L5230L/I T5941T/I K6958K/R	P4715L P5401L G5063S A1306A/S P1640L P2046P/L P2287P/S A2529A/V V2930V/L A3209V V3718A T3750I A6319A T708T/I T2906T/I T3255T/A Y3502Y/C T3646T/A T4161T/I R4589R/Q T5941T/I D6249D/Y R7014R/N	P4715L P5401L G5063S A1306S P2046L P2287S A2529V V2930L T3255I T3646A A6319V M5900M/I R7014R/N	P4715L P5401L G5063S A1306S P2046L P2287S A2529V V2930L T3255I T3646A A6319V I695I/V T1168T/I S1520S/F Y1859Y/H L2329L/F A3392A/V L3606L/F V3690V/L K3832K/T P4624P/S
ORF3a (275)	S26L	S26L A99A/V S180S/F	S26L K235K/T	S26L K16K/T D27D/Y V88V/L K235K/T	S26L A23A/S L83L/F S171S/L	S26L	S26L K16K/N K66K/I

Table S1. Amino acid mutations in all the coding sequences of 6 subgroups of the delta sublineages

Protein (Length)	Delta key mutations	Delta (16.5%) n = 15324	Delta2 +T95I (81.2%) n = 75307	Delta3 +D142G (6%) n = 5508	Delta4 +A222V (4%) n = 3749	Delta5 +Y145H (1.8%) n = 1664	Delta6 +D138H (1.4%) n = 1314
E (75)	-	-	-	V58F	-	-	-
M (222)	I82T	I82T	I82T	I82T	I82T A2A/S	I82T	I82T
Orf6 (61)	-	-	-	-	-	-	-
Orf7a (121)	V82A T120I	V82A T120I V24V/F P45P/L C58C/F A79A/D L116L/F	V82A T120I H73H/Y	V82A T120I P45P/L L116L/F R118R/G	V82A T120I L112L/I	V82A T120I	V82A T120I G38G/E
Orf7b (43)		T40I H42H/S A43A/P	T40I	T40I/T	T40T/I	T40I	T40I
Orf8 (121) (120) (119)	D119- F120-	D119I F120!(stop) I121T V33V/F A65A/S	D119I F120! I121T S67S/F	D119I F120! I121T A65A/S R115R/C	D119I F120! I121T	D119I F120! I121T	D119I F120! I121T
N (419)	D63G R203M D377Y	D63G R203M D377Y G215C Q9Q/L S327S/L G96G/C G238G/C P383P/S D402D/Y	D63G R203M D377Y G215C	D63G R203M D377Y G215C Q9Q/L R209R/I M210M/E S327S/L W330W/L	D63G R203M D377Y G215G/C S202S/S R209R/I M210M/S	D63G R203M D377Y G215C A55A/S S202S/I	D63G R203M D377Y G215C S327S/L
Orf10 (38)	-	-	L16L/P	T38T/I	-	T38T/I	-

Bright green colour highlights fixed mutations. Turquoise colour highlights new mutations undergoing fixation. Grey colour shows emerging reverse-mutations. Yellow colour represents mutations that have significantly increased in frequencies. The rest mutations (not highlighted) are candidates for future genomic surveillance. Cut off for highlighting was placed at >1% prevalence, that is, more than 1 sequence in every 100 sequences sharing same mutation per site. '/' for example, in T22T/I, means that the site has two amino acids, but in this case, there are more 'T's than 'I's. *These three positions can also be captured in these two ways: Either as deletions at F157- and R158-, and substitution at E156G or deletions at E156- and R158-, and substitution at F157G. In all three cases, the final markers that define an unaligned delta variant sequence at these positions are 156G and 157V, and therefore all the three ways have no effect on variant calling.

590   **Table S2.** Resolving the genome sequences into clusters of haplotypes in each delta lineage

Order of haplotype dominance	Frequency of Spike delta haplotypes per mutation							
	Delta	Delta + T95I	Delta + D138H	Delta + D142G	Delta + Y145H	Delta + A222V	Delta + T719I	Delta + N950D
1st	215	401	36	54	188	188	54	2
2nd	123	188	34	37	33	33	16	1
3rd	102	187	30	25	24	28	4	1
4th	96	146	21	21	21	28	3	1
5th	87	131	20	20	21	24	3	NA
6th	67	127	16	18	18	23	3	NA
7th	63	122	13	18	15	21	2	NA
8th	52	103	10	16	14	21	2	NA
9th	47	84	10	15	14	18	2	NA
10th	41	83	10	13	9	16	1	NA