Speech sound categorization: beyond auditory cortices

Basil Preisig* (1,2,3,5), Lars Riecke (4) & Alexis Hervais-Adelman (3,5)

Affiliations

1) Donders Institute for Cognitive Neuroimaging, Radboud University, Nijmegen, the Netherlands

2) Max Planck Institute for Psycholinguistics, Nijmegen, the Netherlands

3) Department of Psychology, Neurolinguistics, University of Zurich, Zurich, Switzerland

4) Department of Cognitive Neuroscience, Faculty of Psychology and Neuroscience, Maastricht University, the Netherlands

5) Neuroscience Center Zurich, University of Zurich and Eidgenössische Technische Hochschule Zurich, 8057 Zurich, Switzerland

*Address for correspondence: Basil C. Preisig, PhD, basilpreisig@gmail.com

Abstract

Which processes in the human brain lead to the categorical perception of speech sounds? Investigation of this question is hampered by the fact that categorical speech perception is normally confounded by acoustic differences in the stimulus. By using ambiguous sounds, however, it is possible to dissociate acoustic from perceptual stimulus representations. Twenty-seven normally hearing individuals took part in an fMRI study in which they were presented with an ambiguous syllable (intermediate between /da/ and /ga/) in one ear and with disambiguating acoustic feature (third formant) in the other ear. Multi-voxel pattern searchlight analysis was used to identify brain areas that consistently differentiated between response patterns associated with different syllable reports. By comparing responses to different stimuli with identical syllable reports and identical stimuli with different syllable reports, we disambiguated whether these regions primarily differentiated the acoustics of the stimuli or the syllable report. We found that BOLD activity patterns in left perisylvian regions (STG, SMG), left inferior frontal regions (vMC, IFG, AI), left supplementary motor cortex (SMA/pre-SMA), and right motor and somatosensory regions (M1/S1) represent listeners' syllable report irrespective of stimulus acoustics. Most of these regions are outside of what is traditionally regarded as auditory or phonological processing areas. Our results indicate that the process of speech sound categorization implicates decision-making mechanisms and auditory-motor transformations.

Keywords: Speech perception; Auditory; MVPA; fMRI; Dichotic listening

1   Introduction

The mapping of sensory information onto common categories is a fundamental feature of human cognition. For example, we can identify a familiar person on different photographs even if taken from different angles. Likewise, we can map speech sounds onto common words even if uttered from different speakers. Categorical perception in speech was described first by Liberman and colleagues (1957) who showed that synthetic syllables along the continuum between prototypes (e.g., /ba/ vs /da/) were perceived categorically despite their incremental acoustic variations. However, it remains unclear how the brain maps the large variety of sensory signals to a limited number of invariant categories.

Speech sound categorization can be conceptualized as the graded and continuous process from sensory perception to action. Consistent with this notion, previous studies have identified multiple auditory and non-auditory brain regions, where speech sound categories can be segregated (Myers et al., 2009). Several studies identified categorical phoneme representations in the auditory association cortex including the superior temporal gyrus (STG) and the superior temporal (STS) using fMRI (Formisano et al., 2008; Kilian-Hütten et al., 2011; Arsenault and Buchsbaum, 2015; Levy and Wilson, 2020) and intracranial recordings (Chang et al., 2010; Mesgarani et al., 2014; Yi et al., 2019). Other studies identified phoneme representations in the motor cortex using fMRI (Lee et al., 2012; Chevillet et al., 2013; Du et al., 2014; Evans and Davis, 2015) and intracranial recordings (Cheung et al., 2016). Evidence that is consistent with the role of motor cortical areas in speech sound categorization is provided by non-invasive brain stimulation studies which perturbed phoneme perception when applied directly over the motor cortex (D'Ausilio et al., 2009; Möttönen and Watkins, 2009; Smalle et al., 2015). Further studies found neural response patterns consistent with categorical encoding in frontal areas like the inferior frontal gyrus possibly involved in executive and decision-making processes (Hasson et al., 2007; Myers et al., 2009; Lee et al., 2012). Lastly, categorical responses to speech sounds have been detected in regions associated with phonological processing such as the left angular gyrus (Blumstein et al., 2005) and the left supramarginal gyrus (Caplan et al., 1995; Zevin and McCandliss, 2005; Raizada and Poldrack, 2007).

A major methodological challenge is to identify brain regions in which neural activity tracks the perceived speech rather than its sensory properties. For instance, if physically different stimuli are used as exemplars of different categories, the perceptual representation of the stimulus is confounded with its physical properties. Under such circumstances, it is thus

difficult to disentangle whether observed neural responses represent the different stimuli, their different percepts, or both. One approach to overcoming this problem is to present two stimuli that are different at the level of sensory encoding, but result in the same speech percept (Formisano et al., 2008; Evans and Davis, 2015). In an fMRI study, Hasson and colleagues (2007) exploited the McGurk effect (Mcgurk & Macdonald, 1976) to present physically different stimuli (audiovisual /ta/ vs. auditory /ka/ + visual /pa/) that elicit the same phonemic percept (/ta/). Using this approach, they identified brain regions that show neural facilitation for stimuli with shared perceptual, but different sensory, encoding. This results were confirmed by Skipper and colleagues (2007) who showed that patterns in these frontal motor areas resulting from the illusory /ta/ percept are more similar to the activity patterns evoked by audiovisual /ta/ than they are to patterns evoked by audiovisual /pa/ or audiovisual /ka/. However, this approach does not easily allow differentiation between areas implicated in audiovisual integration from those involved in categorical speech perception. Another approach is to employ ambiguous stimuli that can elicit different percepts in order to identify brain regions that represent the perceptual report of the participant given the same acoustic stimulus (Kilian-Hütten et al., 2011; Lee et al., 2012). This approach, however, weights perceptual representations resulting from sensory uncertainty more strongly, which may not generalize well to situations with higher sensory evidence.

We presented ambiguous (intermediate between /ga/ and /da/) and unambiguous stimuli (clear /ga/ vs. /da/) together with a disambiguating speech feature (high vs low F3) in a binaural integration paradigm. In the ambiguous condition, the input to each ear was incomplete but complementary so that the perceived syllable identity emerged from the combination of the inputs into a single percept. In the unambiguous condition, the stimulus could be interpreted based on monaural input alone (see Fig. 1).

We used fMRI and searchlight MVPA (Kriegeskorte et al., 2006) to identify brain areas which differentiated between different syllable reports (/ga/ and /da/). In contrast to previous studies (Kilian-Hütten et al., 2011; Lee et al., 2012), we aimed to identify regions that consistently differentiate perceptual reports of both unambiguous and ambiguous stimuli. We tested whether the identified brain regions carry information about the perceived speech (/da/ vs /ga/ response) alone, or its acoustic properties (high vs low F3).

2 Material & Methods

2.1 Participants

Twenty-seven right-handed listeners with no history of hearing impairment (M=21.89 years, SD=3.14, 8 male) took part. The present analysis is based on a dataset collected as part of our previous research on the influence of transcranial brain stimulation on binaural integration (Preisig et al., 2021). All participants had normal or corrected-to-normal visual acuity. The participants reported no history of neurological, psychiatric, nor hearing disorders. All participants had normal hearing (hearing thresholds of less than 25 dB HL at 250, 500, 750, 1000, 1500, 3000, and 4000 Hz, tested on each ear using pure tone audiometry) and no threshold difference between the left ear (LE) and the right ear (RE) larger than 5dB for any of the tested frequencies. All participants gave written informed consent prior to the experiment. This study was approved by the local research ethics committee (CMO region Arnhem-Nijmegen) and was conducted in accordance with the principles of the latest Declaration of Helsinki.

2.2 Auditory Stimuli

The auditory stimuli were based on stimuli reported in previous work (Preisig and Sjerps, 2019; Preisig et al., 2020, 2021). A /da/ to /ga/ stimulus continuum was created in Praat software (Boersma and Weenink, 2019) by shifting the third formant (F3) in 17 equidistant steps from ~2.9 to ~2.5 kHz. The intermediate step at which each participant reported perceiving the stimulus as /da/ or /ga/ in ~50% of the trials was selected as the ambiguous stimuli for RE presentation. To generate the F3 stimulus for LE presentation, the third formant (F3) was extracted from the endpoints of the continuum (from /da/ and /ga/) using a bandpass filter with frequencies between 2100 and 3300 Hz. For a schematic representation of the stimuli see Fig. 1.

2.3 Experimental Design and Task

The dataset reported in this article comprised four task fMRI runs and one fMRI run with passive listening. The data from four additional task runs during which participants underwent non-invasive brain stimulation are reported elsewhere (Preisig et al., 2021).

Each task fMRI run comprised 128 trials, of which 88 trials included the presentation of an auditory stimulus (four trials were discarded because they occurred during a time period where non-invasive brain stimulation was turned on and off to evoke sensations associated with the initial onset and offset of stimulation). Each task fMRI run included 60 binaural integration

trials, for which the F3 frequency of the RE stimulus was set at the individual category boundary, and 24 unambiguous control trials, for which the F3 component of the RE stimulus supported 12 times a clear /da/ and 12 times a clear /ga/ interpretation (see Fig. 1). For binaural integration trials, the LE stimulus was 30 times the high F3 and 30 times the low F3. For control trials, the LE stimulus included a F3 cue with the same F3 frequency as the RE stimulus. Control trials did not require binaural integration for disambiguation because they could be readily identified based on monaural input alone, i.e., the unambiguous /da/ or /ga/ stimulus presented to the RE.

During task fMRI runs, each trial was 3 s long (equal to the repetition time of the fMRI sequence) and started with the acquisition of a single fMRI volume (TA = 930 ms). The auditory stimulus was presented approximately 1750 ms after the trial onset (Preisig et al., 2021). The presentation of the auditory stimulus lasted 250ms. The participant's response window corresponded to the interval from the auditory stimulus onset to 70 ms before the onset of the next trial.

The passive listening fMRI runs consisted of 336 trials, of which 96 trials included auditory stimuli: 48 binaural integration trials and 48 unambiguous trials. Passive listening trials were 2 s long (equal to the repetition time of the fMRI sequence). The auditory stimulus was presented between 1450 and 1550 ms after trial onset. The presentation of the auditory stimulus lasted 250ms.

For additional details on the stimulus presentation see Preisig et al. (2021).

2.4   MRI data acquisition and preprocessing

Anatomical and functional MRI data were acquired with a 3-Tesla Siemens Prisma scanner using a 64-channel head coil. A 3-dimensional high-resolution T1-weighted anatomical volume was acquired using a 3D MPRAGE sequence with the following parameters: repetition time (TR) / inversion time (TI) / echo time (TE) = 2300/1100/3ms, 8° flip angle, FOV 256x216x176 and a 1x1x1 mm isotropic resolution. Parallel imaging (iPAT = GRAPPA) was used to accelerate the acquisition. The acquisition time (TA) of the T1-weighted images was 5 min and 21 sec.

Whole-brain fMRI data was acquired with sparse imaging to minimize the impact of EPI gradient noise during presentation of auditory stimuli (Hall et al., 1999). For this purpose, a delay was introduced in the TR during which the auditory stimuli were presented. This delay was 2070ms during task fMRI and 1070 ms during the passive listening run. For task fMRI

each run included 128 echo planar imaging (EPI) volumes. The passive listening run included 336 EPI volumes. Each scan comprised 66 slices of 2mm thickness which were acquired using a interleaved acquisition sequence with multi-band acceleration (TR$_{task}$: 3000 ms, TR$_{passive listening}$: 2000 ms, TA: 930 ms, TE: 34 ms, flip angle: 90 deg, matrix size: 104x104x66, in-plane resolution: 2x2mm, Multi-band accel. factor: 6x).

fMRI data were pre-processed in SPM12 (http://www.fil.ion.ucl.ac.uk/spm). For multivariate analyses, preprocessing included (I) functional realignment and unwarping and (II) co-registration of the structural image to the mean EPI. Univariate analyses included the additional preprocessing steps (III) normalization of the structural image to a standard template, (IV) application of the normalization parameters to all EPI volumes, and (V) spatial smoothing using a Gaussian kernel with a full-width at half maximum of 8 mm.

2.5 Multi voxel pattern analysis (MVPA)

The MVPA analysis was carried out in subjects' native image space using realigned and unwarpped, but unsmoothed EPI images (Kriegeskorte et al., 2006). A first-level design matrix was specified including one regressor per condition: unambiguous /da/ report, unambiguous /ga/ report, ambiguous /da/ report, ambiguous /ga/ report. Further, the model included a regressor coding the onset of the participants' button presses. As above, six realignment regressors were further included to account for movement-related effects and a constant term per functional imaging run.

We conducted a whole-brain MVPA searchlight analysis (sphere radius 8mm, equivalent to 251 voxels) to identify brain regions in which different syllable reports (/da/ vs /ga/) elicited distinct spatial BOLD response patterns. To evaluate the representational consistency across unambiguous and ambiguous stimuli, an encoding model was specified using the TDT toolbox (Hebart et al., 2015) such that the fitted regression coefficients (beta values) from the unambiguous trials were used for the training dataset and the beta values from ambiguous trials were used for the test dataset.

For statistical inference, we computed the crossnobis distance between the response patterns associated with /da/ and /ga/ syllable reports. The crossnobis distance is the cross-validated version of the Mahalanobis distance (multivariate noise normalized Euclidean distance) (Kriegeskorte et al., 2006; Walther et al., 2016).

The crossnobis distance between /da/ and /ga/ reports was computed as the arithmetic product

of the distances within stimulus class (unambiguous and ambiguous stimulus trials).

d(/da/ report $_{\text{unambiguous}}$, /ga/ report $_{\text{unambiguous}}$) $\times$ d(/da/ report $_{\text{ambiguous}}$, /ga/ report $_{\text{ambiguous}}$)

Like this, BOLD response patterns which consistently differentiated /da/ vs. /ga/ reports in both

unambiguous and ambiguous stimuli yielded greater distances, while inconsistent BOLD

patterns yielded smaller distances.

Crossnobis distances were computed using a leave-one-run-out procedure (Allefeld and

Haynes, 2014). In each of four cross-validation folds, the beta-value maps derived from

unambiguous trials from three fMRI runs were cross-validated against the beta-value maps

derived from ambiguous trials from the left-out run. Cross-validation ensures that the distance

is zero if two voxel patterns are not statistically different from each other, making crossnobis

distance a suitable summary statistic for group-level inference (e.g. with the one-sample t-test).

Note that because of this cross-validation, the crossnobis distance can take negative values if

its true value is close to zero in the presence of noise (Sohoglu et al., 2020).

For group-level inference, individual crossnobis distance maps were normalized and smoothed

(using a Gaussian kernel with a full-width at half maximum of 8 mm) and then entered into a

group level random-effects analysis using permutation-based nonparametric statistics in SNPM

(http://www2.warwick.ac.uk/fac/sci/statistics/staff/academic-research/nichols/software). False

discovery rate (FDR) correction for multiple comparisons across voxels was applied at a

threshold of $p < .05$.

2.6   Univariate analyses

Voxel-wise BOLD activity was modeled by means of a single subject first-level General Linear

Model (GLM) using normalized and spatially smoothed images. The model included one

regressor coding the onsets of the auditory stimuli and one regressor coding the onset of the

participants' button presses during task fMRI. The onsets of the button presses during task fMRI

were modelled to account for the BOLD signal variability resulting from different response

latencies. For each run, six realignment regressors accounting for movement-related effects and

a constant term per functional imaging run were included in the model.

For each participant, T-contrasts (all auditory stimuli > implicit baseline) were computed to

identify brain regions that responded significantly to auditory stimuli during passive listening

and task (Pernet, 2014). Contrast maps from each subject were summarized at the group level
using a one-sample t-test (Fig. 2B).

3 Results

3.1 Binaural integration

Twenty-seven participants listened to ambiguous syllables (intermediate between /ga/ and /da/),
whose perceived identity depends upon binaural integration, and to unambiguous syllables
(clear /ga/ vs. /da/) that could be readily interpreted based on monaural input (for details see
Material & Methods) (Fig. 1). During task fMRI, participants reported on every trial whether
they heard a /da/ or a /ga/ syllable.

Behavioral analyses indicated that participants reliably integrated the speech feature (high vs
low F3) in the binaural integration condition. Participants answered on average with $25.40 \pm$
2.80% (mean $\pm$ SEM) /ga/ responses and $70.60\% \pm 3.00\%$ /da/ responses to ambiguous syllables
combined with the high F3 and $75.00\% \pm 3.5\%$ /ga/ responses and $20.70\% \pm 3.10\%$ /da/
responses for ambiguous syllables combined with the low F3.

The probability of reporting an unambiguous stimulus with low or high F3 respectively as a
/ga/ or /da/ syllable was high: Participants gave on average $84.4\% \pm 2.2\%$ /da/ responses to
unambiguous /da/ stimuli, and $87.4 \pm 1.8\%$ (mean $\pm$ SEM) /ga/ responses were registered for
unambiguous /ga/ stimuli.

247
Fig. 1. Adapted from (Preisig et al., 2021). Schematic illustration of the auditory stimuli
presented during (A) binaural integration and (B) unambiguous control trials. (Left side)
Sound pressure waveform and corresponding sound spectrogram of the non-speech
acoustic feature (F3) presented to the LE in both binaural integration and unambiguous
control trials. (Upper left in A+B) High F3 supporting a /da/ interpretation. (Lower left in
A+B) Low F3 supporting a /ga/ interpretation. (Right) Sound pressure waveform and
corresponding spectrogram of the syllables presented to the right ear. (A: Right side)
Ambiguous syllable intermediate to the syllables /da/ and /ga/ was presented in binaural
integration trials. (B: Right side) Unambiguous /ga/ and /da/ syllables presented to the RE
in control trials. The red lines in the spectrogram highlight the spectral area of stimulus
manipulation.

3.2 BOLD patterns differentiating /da/ vs /ga/ syllable reports

We conducted an MVPA searchlight analysis to identify BOLD response patterns which consistently differentiated /da/ vs. /ga/ reports in unambiguous and ambiguous stimuli. Group-level analysis revealed significant ($p < .05$ FDR-corrected) BOLD activity patterns in auditory and non-auditory brain regions that reliably represented the participants' syllable report. One cluster extended from the anterior insula (AI) over the left inferior frontal gyrus (IFG), into the left ventral motor cortex (vMC). Another cluster was identified in the perisylvian region extending from the left supramarginal gyrus (SMG) into the left superior temporal gyrus (STG). A similar, but smaller cluster was also observed in the right hemisphere. Further, we found a cluster that included the left SMA and pre-SMA and a cluster that included part of the right somatosensory cortex (M1/S1) (see Fig. 2). The analysis also revealed significant clusters in the right anterior cingular cortex (ACC), the left superior frontal gyrus (SFG), the right cerebellum, the right middle frontal gyrus (MFG), and the left corpus callosum (CC) (see Table S1).

Fig. 2. Results of the MVPA searchlight analyses projected onto a canonical MNI single-subject brain. (A) In the highlighted regions, the average crossnobis distance associated with different syllable reports was found to be significantly larger than zero at $p_{FDR} < .05$. (B) The clusters presented in (A) overlaid on regions activated during auditory stimulus presentation and passive listening (light blue color) and regions activated during auditory stimulus presentation and task (dark blue color).

3.3 Acoustic or phonemic representation?

As physical stimulus and its perceptual interpretation were confounded in the MVPA analysis reported in section 3.2, the observed categorical response patterns could be driven by the stimulus acoustics, the syllable percept, or both. Thus, we tested in a follow-up analysis whether categorical patterns derived from the unambiguous stimuli in the localized regions generalize better to the stimulus percept (/da/ vs /ga/) within the same acoustic stimulus, or to the presented acoustic stimulus (high vs low F3) within the same stimulus percept.

For this purpose, we recomputed the crossnobis distance between syllable report-evoked patterns within each stimulus class (high F3; low F3). The converse was also calculated – the distance between acoustic stimulus-evoked patterns, within each syllable report (/da/; /ga/). In both cases the distances were cross-validated as previously, against the syllable reports in unambiguous stimuli.

A group-level analysis (random-effects analysis using permutation-based nonparametric statistics) of normalized and smoothed distance maps revealed significant BOLD activity patterns ($p < .05$ *FDR*-corrected) that differentiated between /da/ vs /ga/ percepts in the aforementioned areas (see Fig. 3A). In contrast, we found no significant clusters for the crossnobis distance maps between high vs low F3 acoustic stimulus at the same statistical threshold.

In order to test for significant differences between syllable report-evoked patterns and acoustic stimulus-evoked patterns, we contrasted perceptual and acoustic maps at the group level (paired t-test using permutation-based nonparametric statistics). The analysis revealed that the crossnobis distance was significantly larger between different syllable percepts than between different stimulus acoustics in the left vMC, the left IFG (part triangularis), in the left Te 2.2, and the left SMA (see Fig 3B).

Fig. 3. (A) Follow-up MVPA analysis constrained to the regions presented in Fig. 2. In green, crossnobis distance between /da/ vs /ga/ percept within the same acoustic stimulus (*p* < .05 *FDR*-corrected) is shown in green, overlaid on the clusters that discriminate different syllable reports across all trials. At the same threshold, we found no significant clusters for the crossnobis distance maps between high vs low F3 acoustic stimulus. (B) The contrast between perceptual and acoustic maps (paired t-test, *p* < .05 *FDR*-corrected) is shown in blue, overlaid on the clusters presented in Fig. 2A.

3.4 Auditory activation during passive listening and task

In order to see whether the identified syllable report-related regions co-localize with areas being activated by the syllable stimuli in the absence of any task and motor response, we mapped auditory-evoked activity during passive listening. Further, we also mapped the areas that were additionally activated by the task (see Fig. 2B). For this purpose, we examined BOLD responses on all trials, contrasted with the baseline (see Methods). During passive listening and task, whole-brain analysis revealed a bilateral brain network extending from the bilateral supratemporal plane into the IFG and the ventral AI. Further, we see activation in the right inferior parietal lobe extending into the right SMG and a cluster in-between the middle cingulum extending into the ventral pre-SMA (see Fig. 2B). In the task blocks, we observed more extensive activation in these cortical areas and additional somatosensory, occipital as well as subcortical activation. As illustrated in Fig. 2B, regions which discriminate different syllable reports co-localize with areas activated during passive listening in the left and right STG, in the left vMC, in the left inferior frontal operculum and the left AI. The other BOLD activity patterns including those in the SMA and right M1/S1 co-localize with regions that were activated during the task.

4 Discussion

In the present study, we aimed to identify the level of neural representation at which speech sounds are coded as abstract categorical perceptual units that are invariant to the sensory signals from which they are derived. For this purpose, we first localized brain regions which consistently discriminated the syllable reports evoked by different unambiguous and ambiguous stimuli. We then assessed whether these regions discriminated primarily the acoustics of the stimuli or the syllable report itself. Our results show that the syllable report was reflected in a set of auditory and non-auditory cortical regions which include regions in the bilateral perisylvian region (STG, SMG), left inferior frontal regions (vMC, IFG, AI), SMA/pre-SMA, and the right M1/S1. Most of these regions are outside of those traditionally regarded as auditory or phonological processing areas, such as the auditory cortex and superior temporal cortex. When interpreting these findings, it is important to keep in mind that the task, as is typical of perceptual experiments, embeds perception in a context of decision- and response-making. We have divided the discussion of our results in four subsections below. Subsections 1-3 each represent a different hierarchy level: (1) auditory perception, (2) categorical perception of syllables, (3) motor response planning and execution. In subsection (4) we argue that there

346 must be a level at which perception and motor planning must integrate to allow performing the
347 task and that this integration involves also higher order cognitive processes such as decision
348 making.

4.1 Auditory perception

One possibility is that the identified neural representation reflects auditory perception. Auditory
speech perception is typically seen as a hierarchical process where simple acoustic features are
extracted in early auditory areas (Schönwiesner and Zatorre, 2009; Leaver and Rauschecker,
2010; Norman-Haignere et al., 2015), which are later transformed into more complex and
speech-specific representations in nonprimary cortex in the STG (Rauschecker and Scott, 2009;
Brodbeck et al., 2018), possibly via parallel streams of auditory processing (Hackett et al., 2001;
Jasmin et al., 2019; Hamilton et al., 2021). Indeed, the relatively subtle acoustic difference
between our stimuli may suggest early auditory processing. However, we found no area where
the crossnobis distance between different acoustic stimuli (high vs low F3) was larger than zero.
It is possible that the acoustic difference was too subtle to be resolved from the fMRI data. Our
analyses show that the identified BOLD patterns rather reflect the syllable percept than the
acoustic difference between the stimuli. Therefore, we think it is likely that the identified BOLD
patterns reflect processes beyond auditory stimulus perception.

4.2 Categorical perception of syllables

Another interpretation is that the identified patterns reflect the categorical perception of
syllables. In line with existing data, we found that the bilateral pSTG/STS (Formisano et al.,
2008; Chang et al., 2010; Kilian-Hütten et al., 2011; Mesgarani et al., 2014; Arsenault and
Buchsbaum, 2015; Yi et al., 2019; Levy and Wilson, 2020), left SMG (Caplan et al., 1995;
Dehaene-Lambertz et al., 2005; Zevin and McCandliss, 2005; Raizada and Poldrack, 2007) for
a meta-analysis see (Turkeltaub and Branch Coslett, 2010), left IFG (Hasson et al., 2007; Myers
et al., 2009; Lee et al., 2012; Chevillet et al., 2013; Du et al., 2014), the left vMC (Du et al.,
2014; Evans and Davis, 2015; Cheung et al., 2016) and premotor cortex (Chevillet et al., 2013)
discriminate between syllable reports in ambiguous and unambiguous stimuli. We found that
all these areas also discriminate between different syllable reports when stimulus acoustics are
kept constant. This effect was most visible in the left vMC, the left IFG (*pars triangularis*), and
the left superior temporal lobe (Te 2.2), in which the crossnobis distances between syllable
percepts were significantly greater than the distances between different stimulus acoustics (see
Fig. 3B).

Further, the analysis revealed areas discriminating syllable reports that have rarely been associated with categorical speech perception: the left SMA and the left AI. In the midline motor areas, the strongest responses to auditory stimuli have been reported in pre-SMA and at the boundary area between pre-SMA and SMA (Lima et al., 2016). According to studies, which localized pre-SMA and SMA with fMRI (Mayka et al., 2006; Kim et al., 2010), the area identified in our study overlaps with the boundary area between pre-SMA and SMA. The SMA receives direct projections from the basal ganglia (Lehéricy et al., 2004; Akkal et al., 2007), the STG/STS (Luppino et al., 1993, 2001; Reznik et al., 2015), and inferior-parietal (Luppino et al., 1993) and inferior-frontal cortices (Catani et al., 2012; Vergani et al., 2014). The SMA is not typically considered as a part of the speech processing network in the brain (Hickok and Poeppel, 2007; Friederici, 2011; Hagoort, 2014), despite its connections with this network. Although SMA activity in response to speech and non-speech sounds has been reported in several studies (for reviews see Hertrich et al., 2016; Lima et al., 2016), the functional role of the SMA in auditory speech processing has remained elusive, possibly because SMA and pre-SMA are traditionally conceptualized as being linked to action-related processes, like speech motor control (Tourville and Guenther, 2011), rather than auditory processes (Lima et al., 2016).

The left AI cluster that we identified includes a portion of the dorsal anterior insula. Insula activity during speech perception has been reported in several studies of sublexical processing of speech (Benson et al., 2001; Golestani and Zatorre, 2004; Falkenberg et al., 2011) Particularly, the dorsal AI seems to be involved in speech perception, whereas other parts of the insula seem to contribute more to speech production (for a meta-analysis see Oh et al., 2014). Similar to the SMA, the insula has extensive connections with the auditory cortex, temporal pole, and superior temporal sulcus (Augustine, 1996; Oh et al., 2014), and is not typically considered in speech perception models (Hickok and Poeppel, 2007; Friederici, 2011; Hagoort, 2014).

Despite their contribution to auditory speech perception the SMA and the AI have rarely been associated with categorical speech perception. Possibly because previous studies either applied passive listening without task (Formisano et al., 2008; Lee et al., 2012), active listening with catch-up trials (Myers et al., 2009) or did not analyze brain signals from the SMA and the AI (Chang et al., 2010; Kilian-Hütten et al., 2011; Mesgarani et al., 2014). Moreover, the contribution of motor action-related regions like the SMA may indicate that additional

410   processes may contribute to speech sound categorization that go beyond categorical encoding

411   as such.

4.3   Motor response planning and execution

Another explanation is that the identified response patterns reflect task-related motor or

domain-general cognitive processes, such as button presses or response selection (Zatorre et al.,

1992; Kawashima et al., 1996). This view is partially supported by our observation that the

right M1/S1 differentiated between different syllable percepts, which we presume to be the

result of the different motor patterns and tactile feedback for each response alternative.

Nonetheless, it is likely that additional processes, other than responding, are responsible for the

effect in these and other regions outside primary sensorimotor areas. Indeed, other regions in

the identified network like the SMA, the left IFG or the left AI have been associated with motor

planning of volitional movements (Tanji and Shima, 1994; Cunnington et al., 2005) and motor

aspects of speech production (Ackermann and Riecker, 2004; Indefrey and Levelt, 2004; Oh et

al., 2014). However, SMA (Benson et al., 2001; Scott et al., 2004; Warren et al., 2006; Jardri

et al., 2007) and insula (Ackermann et al., 2001; Benson et al., 2001; Steinbrink et al., 2009;

Hervais-Adelman et al., 2012) activation has also been reported during passive listening under

various conditions. Further, there are studies showing involvement of SMA and insula during

auditory tasks, which goes beyond merely motor-execution related activity, suggesting a role

of these regions in accent imitation (Adank et al., 2013), vocal affect differentiation

(Bestelmeyer et al., 2014), and categorization of different prosodic contours (statement vs

question) (Sammler et al., 2015). Further, our results show that the regions which discriminate

different syllable reports co-localize with areas activated during passive listening (in the left

and right STG, in the left vMC, in the left inferior frontal operculum and the left AI). This

suggests that other processes unrelated to any task or motor response contribute to the neural

representation that we identified.

4.4   Speech sound categorization: A combination of different processes including higher order cognitive processes

The categorization of auditory information into syllables may involve various processes

including auditory processing, categorical encoding and higher order cognitive processes such

as decision making. In a perceptual task, decision making relies on sensory evidence (Gold and

Shadlen, 2007). This decision process is not trivial because the sensory evidence provided by

neural activation patterns is a mixture of signal and noise. A recent study found that SMA plays

an important role for the transformation of auditory information into categorical responses

during auditory decisions. Morán and colleagues (2021) recorded extracellular activity directly from SMA neurons in rhesus monkeys that were trained to categorize numerous complex sounds, including words. When the authors presented sounds along the continuum between two categories they observed that the neural response during auditory stimulus presentation varied gradually as a function of the changing sensory stimulus, while the neural responses during the button responses showed a categorical pattern that was consistent with the behavioral choices of the animal. This suggests that the SMA integrates acoustic information in order to form categorical signals that drive the behavioral response. Interestingly, the population activity in error trials reflected the behavioral decision, rather than the presented physical stimulus. In line with this findings, our follow-up analysis reported in section 3.3 revealed that the categorical patterns derived from unambiguous stimuli generalize better to the reported syllable (/da/ vs /ga/) than to the presented acoustic stimulus (high vs low F3). This indicates that higher order top-down influences may determine the activation patterns in auditory and somatosensory cortices.

It is possible that the left vMC has a similar function in speech sound categorization as the SMA. Skipper and colleagues (2007) investigated the role of the motor regions that are responsible for speech production during audiovisual speech perception. For their experiment, they used incongruent audiovisual stimuli (auditory /pa/ + visual /ka/) that elicit the phonemic percept /ta/ (Mcgurk & Macdonald, 1976). They found that "illusory" /ta/ percepts evoked activation patterns in the left vMC that were more similar to the activation pattern evoked by the audiovisual /ta/ stimulus than the activation pattern evoked by the audiovisual /pa/ or the audiovisual /ka/ stimulus. This suggests that left vMC activation represented the perceived stimulus rather than auditory or visual stimulus features. Most interestingly, the authors did an additional analysis where they compared the similarity of activations patterns over the entire time course of the hemodynamic response. For the left vMC, their analysis revealed that the activity patterns evoked by the "illusory" /ta/ percept showed the highest similarity with the activity evoked by the audiovisual /ta/ over the entire time course in the left vMC. However, the activation patterns in auditory and somatosensory cortices showed the highest similarity with audiovisual /pa/ for the first 1.5s of the hemodynamic response and were only afterwards more similar to the "illusory" /ta/ percept. This suggests that higher order top-down influences may determine the activation patterns in auditory and somatosensory cortices and the phoneme percept as well as the behavioral response.

The left AI has been associated with decision making in a number of studies (for a review see Droutman et al., 2015). Particularly, the dorsal AI seems to be functionally implicated in cognitive control processes associated with decision-making, including attention re-focusing, evaluation, action, and outcome processing (Droutman et al., 2015). Further, dorsal AI activation has been related to decision ambiguity (Huettel et al., 2006) and error awareness (Ullsperger et al., 2010). Further recent evidence indicates that the anterior insula could serve as a gate for conscious access to sensory information (Huang et al., 2021).

We found two interesting distinctions between the BOLD patterns in the left SMA, left vMC, and the left AI. First, only the cluster in the left AI and left vMC overlapped with areas which were activated during passive listening to the same stimuli (Fig. 2B). Second, only in the left SMA and the left vMC, but not in the left AI, the crossnobis distances between syllable percepts were larger than distances between different stimulus acoustics. This indicates that the responses in the left SMA and left vMC, but not left AI, were invariant to the physical stimulus properties of the stimuli. Taken together, one may speculate about the processing hierarchy and the functional differences between the processing in the AI, the vMC and the SMA. The AI may be more relevant for the integration of auditory evidence and the preparation of a perceptual representation, which is then strengthened to become a stimulus-syllable representation in the vMC and SMA.

4.5   Conclusion

Our finding that BOLD patterns discriminating different syllable reports occur in brain regions contributing to auditory processing, categorical encoding and higher order cognitive processes suggests that the identified representations reflect a combination of various processes along this hierarchy.

Our finding that areas whose responses to speech stimuli discriminate between phonemic categories exist outside auditory cortical areas is consistent with the possibility that higher-order areas are instrumental in determining the syllables we hear, and that these regions feed abstract categorical representations back into the auditory association cortex (Formisano et al., 2008; Chang et al., 2010; Kilian-Hütten et al., 2011) and even to earlier auditory areas (Levy and Wilson, 2020).

5   Acknowledgement

This work was supported by the Swiss National Science Foundation [P2BEP3_168728 / PP00P1_163726] and the Janggen-Pöhn Stiftung. The authors would like to thank Benjamin Kop, Brigit Knudsen, Iris Schmits, Uriel Plones, and Paul Gaalman for their assistance as well as Martin Hebart for the methodological advice with regard to the MVPA analysis.

6   References

Ackermann H, Riecker A (2004) The contribution of the insula to motor aspects of speech production: A review and a hypothesis. Brain Lang 89:320–328.

Ackermann H, Riecker A, Mathiak K, Erb M, Grodd W, Wildgruber D (2001) Rate-dependent activation of a prefrontal-insular-cerebellar network during passive listening to trains of click stimuli: An fMRI study. NeuroReport 12:4087–4092.

Adank P, Rueschemeyer S-A, Bekkering H (2013) The role of accent imitation in sensorimotor integration during processing of intelligible speech. Front Hum Neurosci 7 Available at: https://www.frontiersin.org/articles/10.3389/fnhum.2013.00634/full [Accessed July 8, 2021].

Akkal D, Dum RP, Strick PL (2007) Supplementary Motor Area and Presupplementary Motor Area: Targets of Basal Ganglia and Cerebellar Output. J Neurosci 27:10659–10673.

Allefeld C, Haynes J-D (2014) Searchlight-based multi-voxel pattern analysis of fMRI by cross-validated MANOVA. NeuroImage 89:345–357.

Arsenault JS, Buchsbaum BR (2015) Distributed Neural Representations of Phonological Features during Speech Perception. J Neurosci 35:634–642.

Augustine JR (1996) Circuitry and functional aspects of the insular lobe in primates including humans. Brain Res Rev 22:229–244.

Benson RR, Whalen DH, Richardson M, Swainson B, Clark VP, Lai S, Liberman AM (2001) Parametrically Dissociating Speech and Nonspeech Perception in the Brain Using fMRI. Brain Lang 78:364–396.

Bestelmeyer PEG, Maurage P, Rouger J, Latinus M, Belin P (2014) Adaptation to Vocal Expressions Reveals Multistep Perception of Auditory Emotion. J Neurosci 34:8098–8105.

Blumstein SE, Myers EB, Rissman J (2005) The Perception of Voice Onset Time: An fMRI Investigation of Phonetic Category Structure. J Cogn Neurosci 17:1353–1366.

Boersma P, Weenink D (2019) Praat: doing phonetics by computer [Computer program]. Available at: http://www.praat.org/ [Accessed February 17, 2019].

Brodbeck C, Hong LE, Simon JZ (2018) Rapid Transformation from Auditory to Linguistic Representations of Continuous Speech. Curr Biol 28:3976-3983.e5.

Caplan D, Gow D, Makris N (1995) Analysis of lesions by MRI in stroke patients with acoustic-phonetic processing deficits. Neurology 45:293–298.

Catani M, Dell'Acqua F, Vergani F, Malik F, Hodge H, Roy P, Valabregue R, Thiebaut de Schotten M (2012) Short frontal lobe connections of the human brain. Cortex 48:273–291.

Chang EF, Rieger JW, Johnson K, Berger MS, Barbaro NM, Knight RT (2010) Categorical
speech representation in human superior temporal gyrus. Nat Neurosci 13:1428–1432.

Cheung C, Hamilton LS, Johnson K, Chang EF (2016) The auditory representation of speech
sounds in human motor cortex Shinn-Cunningham BG, ed. eLife 5:e12577.

Chevillet MA, Jiang X, Rauschecker JP, Riesenhuber M (2013) Automatic Phoneme Category
Selectivity in the Dorsal Auditory Stream. J Neurosci 33:5208–5215.

Cunnington R, Windischberger C, Moser E (2005) Premovement activity of the pre-
supplementary motor area and the readiness for action: Studies of time-resolved event-
related functional MRI. Hum Mov Sci 24:644–656.

D'Ausilio A, Pulvermüller F, Salmas P, Bufalari I, Begliomini C, Fadiga L (2009) The Motor
Somatotopy of Speech Perception. Curr Biol 19:381–385.

Dehaene-Lambertz G, Pallier C, Serniclaes W, Sprenger-Charolles L, Jobert A, Dehaene S
(2005) Neural correlates of switching from auditory to speech perception. NeuroImage
24:21–33.

Droutman V, Bechara A, Read SJ (2015) Roles of the Different Sub-Regions of the Insular
Cortex in Various Phases of the Decision-Making Process. Front Behav Neurosci 9
Available at: https://www.frontiersin.org/articles/10.3389/fnbeh.2015.00309/full
[Accessed May 21, 2019].

Du Y, Buchsbaum BR, Grady CL, Alain C (2014) Noise differentially impacts phoneme
representations in the auditory and speech motor systems. Proc Natl Acad Sci
111:7126–7131.

Evans S, Davis MH (2015) Hierarchical Organization of Auditory and Motor Representations
in Speech Perception: Evidence from Searchlight Similarity Analysis. Cereb Cortex
25:4772–4788.

Falkenberg LE, Specht K, Westerhausen R (2011) Attention and cognitive control networks
assessed in a dichotic listening fMRI study. Brain Cogn 76:276–285.

Formisano E, Martino FD, Bonte M, Goebel R (2008) "Who" Is Saying "What"? Brain-Based
Decoding of Human Voice and Speech. Science 322:970–973.

Friederici AD (2011) The brain basis of language processing: From structure to function.
Physiol Rev 91:1357–1392.

Gold JI, Shadlen MN (2007) The neural basis of decision making. Annu Rev Neurosci 30:535–
574.

Golestani N, Zatorre RJ (2004) Learning new sounds of speech: reallocation of neural
substrates. NeuroImage 21:494–506.

Hackett TA, Preuss TM, Kaas JH (2001) Architectonic identification of the core region in
auditory cortex of macaques, chimpanzees, and humans. J Comp Neurol 441:197–222.

Hagoort P (2014) Nodes and networks in the neural architecture for language: Broca's region
and beyond. Curr Opin Neurobiol 28:136–141.

Hall DA, Haggard MP, Akeroyd MA, Palmer AR, Summerfield AQ, Elliott MR, Gurney EM,
Bowtell RW (1999) "sparse" temporal sampling in auditory fMRI. Hum Brain Mapp
7:213–223.

Hamilton LS, Oganian Y, Hall J, Chang EF (2021) Parallel and distributed encoding of speech
across   human   auditory   cortex.   Cell   Available   at:
https://www.sciencedirect.com/science/article/pii/S0092867421008783   [Accessed
August 19, 2021].

Hasson U, Skipper JI, Nusbaum HC, Small SL (2007) Abstract Coding of Audiovisual Speech:
Beyond Sensory Representation. Neuron 56:1116–1126.

Hebart MN, Görgen K, Haynes J-D (2015) The Decoding Toolbox (TDT): a versatile software
package for multivariate analyses of functional imaging data. Front Neuroinformatics 8
Available   at: https://www.frontiersin.org/articles/10.3389/fninf.2014.00088/full
[Accessed February 25, 2019].

Hertrich I, Dietrich S, Ackermann H (2016) The role of the supplementary motor area for
speech and language processing. Neurosci Biobehav Rev 68:602–610.

Hervais-Adelman AG, Carlyon RP, Johnsrude IS, Davis MH (2012) Brain regions recruited for
the effortful comprehension of noise-vocoded words. Lang Cogn Process 27:1145–
1166.

Hickok G, Poeppel D (2007) The cortical organization of speech processing. Nat Rev Neurosci
8:393–402.

Huang Z, Tarnal V, Vlisides PE, Janke EL, McKinney AM, Picton P, Mashour GA, Hudetz AG
(2021) Anterior insula regulates brain network transitions that gate conscious access.
Cell Rep 35:109081.

Huettel SA, Stowe CJ, Gordon EM, Warner BT, Platt ML (2006) Neural Signatures of
Economic Preferences for Risk and Ambiguity. Neuron 49:765–775.

Indefrey P, Levelt WJM (2004) The spatial and temporal signatures of word production
components. Cognition 92:101–144.

Jardri R, Pins D, Bubrovszky M, Despretz P, Pruvo J-P, Steinling M, Thomas P (2007) Self
awareness and speech processing: An fMRI study. NeuroImage 35:1645–1653.

Jasmin K, Lima CF, Scott SK (2019) Understanding rostral–caudal auditory cortex
contributions to auditory perception. Nat Rev Neurosci 20:425.

Kawashima R, Satoh K, Itoh H, Ono S, Furumoto S, Gotoh R, Koyama M, Yoshioka S,
Takahashi T, Takahashi K, Yanagisawa T, Fukuda H (1996) Functional anatomy of
GO/NO-GO discrimination and response selection — a PET study in man. Brain Res
728:79–89.

619 Kilian-Hütten N, Valente G, Vroomen J, Formisano E (2011) Auditory Cortex Encodes the
620   Perceptual Interpretation of Ambiguous Sound. J Neurosci 31:1715–1720.

621 Kim J-H, Lee J-M, Jo HJ, Kim SH, Lee JH, Kim ST, Seo SW, Cox RW, Na DL, Kim SI, Saad
622   ZS (2010) Defining functional SMA and pre-SMA subregions in human MFC using
623   resting state fMRI: Functional connectivity-based parcellation method. NeuroImage
624   49:2375–2386.

625 Kriegeskorte N, Goebel R, Bandettini P (2006) Information-based functional brain mapping.
626   Proc Natl Acad Sci 103:3863–3868.

627 Leaver AM, Rauschecker JP (2010) Cortical Representation of Natural Complex Sounds:
628   Effects of Acoustic Features and Auditory Object Category. J Neurosci 30:7604–7612.

629 Lee Y-S, Turkeltaub P, Granger R, Raizada RDS (2012) Categorical Speech Processing in
630   Broca's Area: An fMRI Study Using Multivariate Pattern-Based Analysis. J Neurosci
631   32:3942–3948.

632 Lehéricy S, Ducros M, Krainik A, Francois C, Van de Moortele P-F, Ugurbil K, Kim D-S
633   (2004) 3-D Diffusion Tensor Axonal Tracking shows Distinct SMA and Pre-SMA
634   Projections to the Human Striatum. Cereb Cortex 14:1302–1309.

635 Levy DF, Wilson SM (2020) Categorical Encoding of Vowels in Primary Auditory Cortex.
636   Cereb Cortex 30:618–627.

637 Liberman AM, Harris KS, Hoffman HS, Griffith BC (1957) The discrimination of speech
638   sounds within and across phoneme boundaries. J Exp Psychol 54:358–368.

639 Lima CF, Krishnan S, Scott SK (2016) Roles of Supplementary Motor Areas in Auditory
640   Processing and Auditory Imagery. Trends Neurosci 39:527–542.

641 Luppino G, Calzavara R, Rozzi S, Matelli M (2001) Projections from the superior temporal
642   sulcus to the agranular frontal cortex in the macaque. Eur J Neurosci 14:1035–1040.

643 Luppino G, Matelli M, Camarda R, Rizzolatti G (1993) Corticocortical connections of area F3
644   (SMA-proper) and area F6 (pre-SMA) in the macaque monkey. J Comp Neurol
645   338:114–140.

646 Mayka MA, Corcos DM, Leurgans SE, Vaillancourt DE (2006) Three-dimensional locations
647   and boundaries of motor and premotor cortices as defined by functional brain imaging:
648   A meta-analysis. NeuroImage 31:1453–1474.

649 Mcgurk H, Macdonald J (1976) Hearing lips and seeing voices. Nature 264:746–748.

650 Mesgarani N, Cheung C, Johnson K, Chang EF (2014) Phonetic feature encoding in human
651   superior temporal gyrus. Science 343:1006–1010.

652 Morán I, Perez-Orive J, Melchor J, Figueroa T, Lemus L (2021) Auditory decisions in the
653   supplementary motor area. Prog Neurobiol 202:102053.

654 Möttönen R, Watkins KE (2009) Motor representations of articulators contribute to categorical
655   perception of speech sounds. J Neurosci Off J Soc Neurosci 29:9819–9825.

Myers EB, Blumstein SE, Walsh E, Eliassen J (2009) Inferior Frontal Regions Underlie the Perception of Phonetic Category Invariance. Psychol Sci 20:895–903.

Norman-Haignere S, Kanwisher NG, McDermott JH (2015) Distinct Cortical Pathways for Music and Speech Revealed by Hypothesis-Free Voxel Decomposition. Neuron 88:1281–1296.

Oh A, Duerden EG, Pang EW (2014) The role of the insula in speech and language processing. Brain Lang 135:96–103.

Pernet CR (2014) Misconceptions in the use of the General Linear Model applied to functional MRI: a tutorial for junior neuro-imagers. Front Neurosci 8 Available at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3896880/ [Accessed May 1, 2020].

Preisig BC, Riecke L, Sjerps MJ, Kösem A, Kop BR, Bramson B, Hagoort P, Hervais-Adelman A (2021) Selective modulation of interhemispheric connectivity by transcranial alternating current stimulation influences binaural integration. Proc Natl Acad Sci 118 Available at: https://www.pnas.org/content/118/7/e2015488118 [Accessed February 11, 2021].

Preisig BC, Sjerps MJ (2019) Hemispheric specializations affect interhemispheric speech sound integration during duplex perception. J Acoust Soc Am 145:EL190–EL196.

Preisig BC, Sjerps MJ, Hervais-Adelman A, Kösem A, Hagoort P, Riecke L (2020) Bilateral Gamma/Delta Transcranial Alternating Current Stimulation Affects Interhemispheric Speech Sound Integration. J Cogn Neurosci 32:1242–1250.

Raizada RDS, Poldrack RA (2007) Selective Amplification of Stimulus Differences during Categorical Processing of Speech. Neuron 56:726–740.

Rauschecker JP, Scott SK (2009) Maps and streams in the auditory cortex: nonhuman primates illuminate human speech processing. Nat Neurosci 12:718–724.

Reznik D, Ossmy O, Mukamel R (2015) Enhanced Auditory Evoked Activity to Self-Generated Sounds Is Mediated by Primary and Supplementary Motor Cortices. J Neurosci 35:2173–2180.

Sammler D, Grosbras M-H, Anwander A, Bestelmeyer PEG, Belin P (2015) Dorsal and Ventral Pathways for Prosody. Curr Biol 25:3079–3085.

Schönwiesner M, Zatorre RJ (2009) Spectro-temporal modulation transfer function of single voxels in the human auditory cortex measured with high-resolution fMRI. Proc Natl Acad Sci 106:14611–14616.

Scott SK, Rosen S, Wickham L, Wise RJS (2004) A positron emission tomography study of the neural basis of informational and energetic masking effects in speech perception. J Acoust Soc Am 115:813–821.

Skipper JI, van Wassenhove V, Nusbaum HC, Small SL (2007) Hearing lips and seeing voices: how cortical areas supporting speech production mediate audiovisual speech perception. Cereb Cortex 17:2387–2399.

Smalle EHM, Rogers J, Möttönen R (2015) Dissociating Contributions of the Motor Cortex to
Speech Perception and Response Bias by Using Transcranial Magnetic Stimulation.
Cereb Cortex 25:3690–3698.

Sohoglu E, Kumar S, Chait M, Griffiths TD (2020) Multivoxel codes for representing and
integrating acoustic features in human cortex. NeuroImage 217:116661.

Steinbrink C, Ackermann H, Lachmann T, Riecker A (2009) Contribution of the anterior insula
to temporal auditory processing deficits in developmental dyslexia. Hum Brain Mapp
30:2401–2411.

Tanji J, Shima K (1994) Role for supplementary motor area cells in planning several
movements ahead. Nature 371:413–416.

Tourville JA, Guenther FH (2011) The DIVA model: A neural theory of speech acquisition and
production. Lang Cogn Process 26:952–981.

Turkeltaub PE, Branch Coslett H (2010) Localization of sublexical speech perception
components. Brain Lang 114:1–15.

Ullsperger M, Harsay HA, Wessel JR, Ridderinkhof KR (2010) Conscious perception of errors
and its relation to the anterior insula. Brain Struct Funct 214:629–643.

Vergani F, Lacerda L, Martino J, Attems J, Morris C, Mitchell P, Schotten MT de, Dell'Acqua
F (2014) White matter connections of the supplementary motor area in humans. J Neurol
Neurosurg Psychiatry 85:1377–1385.

Walther A, Nili H, Ejaz N, Alink A, Kriegeskorte N, Diedrichsen J (2016) Reliability of
dissimilarity measures for multi-voxel pattern analysis. NeuroImage 137:188–200.

Warren JE, Sauter DA, Eisner F, Wiland J, Dresner MA, Wise RJS, Rosen S, Scott SK (2006)
Positive emotions preferentially engage an auditory-motor "mirror" system. J Neurosci
26:13067–13075.

Yi HG, Leonard MK, Chang EF (2019) The Encoding of Speech Sounds in the Superior
Temporal Gyrus. Neuron 102:1096–1110.

Zatorre RJ, Evans AC, Meyer E, Gjedde A (1992) Lateralization of phonetic and pitch
discrimination in speech processing. Science 256:846–849.

Zevin JD, McCandliss BD (2005) Dishabituation of the BOLD response to speech sounds.
Behav Brain Funct 1:4.

7 Supplementary Table

Table S1: The results of the group-level analysis (random-effects analysis using permutation-based nonparametric statistics) of normalized and smoothed individual distance maps (p< .05 FDR-corrected)

	Coordinates			T score	k
	x	y	z		
right postcentral gyrus (PoCG)	42	-26	44	6.47	2552
left anterior insula (AI)	-30	20	12	6.25	2368
left ventral motor cortex (vMC)	-38	2	42	5.51	
left precentral gyrus (PCG)	-40	10	42	5.34	
right anterior cingular cortex (ACC)	4	34	-4	5.49	141
left superior frontal gyrus (SFG)	-22	-2	66	5.48	75
right cerebellum	16	-68	-44	5.31	118
left supplementarymotor cortex (SMA)	-6	-8	54	5.05	817
left pre-SMA	-6	12	62	3.26	
left supramarginal gyrus / left superior temporal gyrus (STG)	-52	-26	22	4.87	1588
left postcentral gyrus (PoG)	-48	-32	52	4.44	
left angular gyrus	-40	-60	42	3.72	
superior medial frontal gyurs (at the border to the pre-SMA)	0	28	50	4.68	244
right superior medial frontal gyurs	4	48	44	3.90	
right middle frontal gyrus (MFG)	22	44	32	4.62	198
left corpus callosum (CC)	-10	14	24	4.16	107

Coordinates are in MNI space.