Age-related change in task-evoked amygdala-prefrontal circuitry: a multiverse approach with an accelerated longitudinal cohort aged 4-22 years

Paul Alexander Bloom[1*], Michelle VanTieghem[2], Laurel Gabard-Durnam[3], Dylan G. Gee[4], Jessica Flannery[5], Christina Caldera[6], Bonnie Goff[6], Eva H. Telzer[7], Kathryn L. Humphreys[8], Dominic S. Fareri[9], Mor Shapiro[10], Sameah Algharazi[11], Niall Bolger[1], Mariam Aly[1], & Nim Tottenham[1]

Affiliations:
[1]Columbia University
[2]Virta Health
[3]Northeastern University
[4]Yale University
[5]Limbix Health
[6]University of California Los Angeles
[7]University of North Carolina at Chapel Hill
[8]Vanderbilt University
[9]Adelphi University
[10]Kaiser Permanente
[11]City College of New York

Please address correspondence to:
Paul Alexander Bloom
Columbia University Psychology, New York, NY
Email: paul.bloom@columbia.edu

Abstract
There has been considerable interest in the development of the amygdala and its connections with medial prefrontal cortex (mPFC) given the central role of these brain regions in emotional processes. While several studies have suggested that this circuitry exhibits functional changes across the first two decades of life, they have typically employed cross-sectional designs, and findings have been mixed. Additionally, analytic choices may contribute to discrepancies across studies. Here we used an accelerated longitudinal design to examine task-evoked changes in amygdala–mPFC circuitry from 4-22 years of age (N=98; 183 total scans; 1-3 scans per participant). Participants were recruited from the greater Los Angeles area, and completed an event-related emotional face (fear, neutral) task designed to be appropriate for the wide age range. 'Multiverse' analyses examined the robustness of our findings to fMRI analysis choices. 2808 parallel analyses varying in preprocessing and modeling choices found evidence for average age-related decreases in amygdala reactivity to faces. Greater amygdala reactivity at younger ages was attributable to elevated responses during the first few trials relative to later trials. Within-participant changes in amygdala reactivity with age could not be differentiated from between-participant differences, however. Across analysis decision points, we did not find consistent evidence of age-related change in amygdala–mPFC connectivity through generalized psychophysiological interaction (gPPI) or beta-series correlation (BSC) methods. We also did not find evidence for associations between separation anxiety behaviors and amygdala reactivity or amygdala-mPFC connectivity. Within the context of this faces task and age range, age-related changes in amygdala reactivity were more robust to processing pipeline than were task-evoked functional connectivity measures, particularly those using gPPI. These findings highlight both the challenges in estimating developmental change in longitudinal cohorts and the value of multiverse approaches in developmental neuroimaging for assessing robustness of results.

Rodent models and human neuroimaging have provided converging evidence for the

importance of the amygdala and medial prefrontal cortex (mPFC) in the development of threat

processing[1,2], emotion regulation[3–5], and affective learning[6,7]. Characterizing growth

trajectories of these regions may provide insight into neural constructions underlying

emotional development[8]. To probe amygdala–mPFC circuitry across development, face

stimuli are frequently employed because they effectively engage this circuitry while being

child-appropriate[9]. Though a number of studies have examined age-related changes from

childhood to young adulthood in amygdala responses and amygdala–mPFC functional

connectivity (FC) associated with emotional face stimuli, findings have varied widely. Several

studies have found age-related change in amygdala reactivity, including decreases as a

function of age in response to emotional faces[10–15] (as well as other images)[16–18], increases in

amygdala reactivity with age[19,20], or peaks during adolescence[21,22]. Others have observed no

age-related changes[23–28]. With task-evoked amygdala–mPFC FC, several studies have found

age-related decreases from childhood to young adulthood[10,23,26,29], while others have found

increases[16,18,30], or little age-related change[28].

While the small sample sizes examined in many studies on amygdala–mPFC

development likely contribute to differences in findings[31], important methodological differences

also exist across studies. Differences in age range or sample demographics, stimuli, task

(e.g. passive viewing vs. emotion labeling or matching[32]), task design (block vs. event-

related[33]), or contrast used (faces > shapes vs. faces > baseline) may also contribute to

discrepancies. The brain regions under investigation also differ across studies; for example,

prefrontal clusters with which amygdala connectivity was assessed. Interpreting discrepancies

across studies without appreciation for these methodological differences would be

inappropriate, and in fact, incorrect. Yet, such differences do not account for all discrepancies

in findings across studies. Variation in processing pipelines is another source of differences

across studies, as varying analytic decisions can produce qualitatively different findings, even

between putatively identical analyses of the same dataset[34]. Choices including software

package[35], spatial smoothing[36], treatment of head motion[37], parcellation[38], and functional

connectivity approach[39] can also impact results and qualitatively change findings[40].

Additionally, the majority of developmental investigations of amygdala–mPFC function have

studied cross-sectional samples. Because cross-sectional studies are susceptible to cohort

effects and cannot measure within-participant change, longitudinal work has been

recommended for better charting of developmental trajectories[41,42].

Here, we examined task-related amygdala–mPFC functional development in an

accelerated longitudinal sample ranging from ages 4-22 years. We selected a task that was

designed to be appropriate for young ages to characterize developmental change in

amygdala–mPFC responses to fear and neutral faces across childhood and adolescence, and

we asked whether findings were robust to analytical choices. This accelerated longitudinal

design is an extension of the sample reported on by Gee et al. (2013). We preregistered two

hypotheses (https://osf.io/8nyj7/) predicting that both amygdala reactivity (1) and amygdala–

mPFC connectivity (2) as measured with generalized psychophysiological interaction models

(gPPI), would decrease as a function of age during viewing of fearful faces (relative to

baseline), and that gPPI would show a shift from positive to negative valence to fear faces

during early adolescence (see Table 1 Aims 1a & 2a). Although we did not preregister further

hypotheses, we also investigated age-related changes in within-scan differences in amygdala

responses across trials and FC using beta series correlations. As previous work identified

associations between amygdala–mPFC FC and separation anxiety[10,43], we asked whether any

amygdala–mPFC measures were associated longitudinally with separation anxiety behaviors

(see Table 1 Aim 3). We used 'multiverse' analyses and specification curves to examine

whether results were robust to many analytical decisions.

51
52 **Figure 1. A.** Schematic showing study inclusion criteria. **B.** Included scans at each study wave,
53 with each dot representing one scan, and horizontal lines connecting participants across study
54 waves.
55

Results

Age-related change in amygdala reactivity

We used multilevel regression models and specification curve analyses to examine age-

related changes in amygdala reactivity to faces in an accelerated longitudinal sample ranging

from ages 4-22 years (Figure 2). Across specifications, we found consistent evidence for

negative age-related change in anatomically-defined (Harvard-Oxford atlas and Freesurfer-

defined) amygdala reactivity to fear faces > baseline, such that the vast majority of analysis

specifications (99.6%) estimated linear slopes at the group level that were negative in sign, and

the majority (60.0%) of 95% posterior intervals about these slopes excluded 0 (Figure 2A;

interactive version at https://pbloom.shinyapps.io/amygdala_mpfc_multiverse). Thus, over half

of models indicated that on average, increases in age were associated with decreases in

amygdala reactivity to fear faces > baseline. Because the timepoint 1 data in the current study

included the 42 scans used by Gee et al. (2013) to age-related changes in amygdala—mPFC

circuitry for the fear > baseline contrast, results including these scans may have been more

likely to find similar change (particularly for fear > baseline, see sFigures 11-13, & 25).

Estimated age-related change was on average weaker, though still largely negative (98.1%

negative, 25.3% of posterior intervals excluding 0) when 42 previously analyzed scans were

excluded to provide stricter independence from previously analyzed data (see sFigure 11, Gee

et al., 2013).

Parallel multiverse analyses found consistent age-related decreases in neutral faces >

baseline amygdala reactivity (see Figure 2C for an example pipeline & sFigure 9 for

specification curve), but no consistent evidence for age-related change for the fear > neutral

contrast (see sFigure 10). However, there was consistent evidence for higher reactivity for fear

faces > neutral on average as well as each emotion compared to baseline (sFigures 6-8),

indicating that while the amygdala responses were robust and generally stronger for fear faces

compared to neutral, such fear > neutral differences did not change with age. Across contrasts,

inclusion of block order or scanner covariates, inclusion of random intercepts, and use of robust

regression models had little impact on age-related change estimates (see Figure 2B sFigures 6-

8).

While group-level estimates of average age-related change were relatively consistent

across specifications, the estimated age terms in these models could be influenced by both

within-participant change and between-participant differences[44,45]. A smaller separate

specification curve indicated that when models were parametrized to differentiate within-

participant change and between-participant differences, average within-participant change was

not consistent across specifications and could not be estimated with precision (Figure 2D). In

contrast, estimates of between-participant differences largely indicated negative age-related

change in concurrence with our initial model parametrization. At the same time, within-

participant versus between-participant terms were not reliably different from one another,

indicating that models could not distinguish them despite higher precision for estimating

between-participant differences (see sFigure 18). We did not find consistent evidence for

quadratic age-related changes in amygdala reactivity (see sFigures 14-17).

100
Figure 2. Multiverse analyses of age-related change in amygdala reactivity. **A.** Specification
curve of age-related change in fear > baseline amygdala reactivity. Points represent estimated
linear age-related change and lines are corresponding 95% posterior intervals (PIs). Models are
ordered by age-related change estimates, with the dotted line representing the median estimate
across all specifications. Color indicates sign of beta estimates and whether respective posterior
intervals include 0 (red = negative excluding 0; blue = negative including 0, green = positive
including 0, black = median across all specifications). **B.** Model specification information
corresponding to each model in A. Variables on the y-axis represent analysis choices,
corresponding color-coded marks indicate that a choice was made, and blank space indicates
that the choice was not made in a given analysis. Within each category panel (amygdala ROI,
Group-Level Model, and Participant-Level Model), decision points are ordered from top to
bottom by the median model rank when the corresponding choice is made (i.e. choices at the
top of each panel tend to have more negative age-related change estimates). Black points with
error bars represent the median and IQR ranks of specifications making the choice indicated on
the corresponding line. **C.** Example participant-level data and model predictions for age-related
related change in amygdala reactivity for both the fear > baseline (green) and neutral-baseline
(orange) contrasts. Data are shown for a preregistered pipeline using a native space bilateral
amygdala mask, 24 motion regressors, t-statistics, high-pass filtering, and participant-level

GLMs in FSL. Points represent participant-level estimates, light lines connect estimates from
participants with multiple study visits, and dark lines with shaded area represent model
predictions and 95% posterior intervals. **D.** Specification curves for a subset of models
separately parametrizing within-participant (right) vs. between-participant (left) age-related
change for both the fear > baseline (green) and neutral > baseline (orange) contrasts, as well as
the median across specifications (black). See
https://pbloom.shinyapps.io/amygdala_mpfc_multiverse/ for interactive visualizations.
126

Age-related differences in within-scan amygdala reactivity change

To ask whether age-related changes in amygdala reactivity could be due to

developmental changes in patterns of amygdala reactivity across face trials (within a run), we

examined whether within-scan change in amygdala reactivity varied with age (see Table 1 Aim

1b). Analyses included 42 specifications (3 amygdala regions of interest [ROIs] x 2 global signal

correction options x 7 group-level models). Across both fear and neutral trials, linear slopes of

amygdala reactivity were negative on average, indicating higher amygdala reactivity at the

beginning of the run (Figure 3A, sFigure 30). Across specifications, for both fear (100% of

estimates had the same sign, 95.2% of posterior intervals excluding 0 in the same direction) and

neutral trials (100% of estimates in the same direction, 38.1% of posterior intervals excluding 0),

there was evidence that these within-scan slopes were steeper (i.e., more negative) at younger

ages, though evidence did not meet our rule-of-thumb criteria for 'robustness' (>=50% of 95%

posterior intervals excluding 0 in the same direction) for neutral trials (Figure 3D-E).

Specifications with a global signal subtraction step also tended to find stronger age-related

change.

Similarly, when splitting trials into the first half (trials 1-12) versus second half (trials 13-

24), there was consistent evidence (100% of estimates had the same sign, 69.2% with posterior

interval excluding 0) for an interaction between age and trial half, such that average reactivity to

fear faces > baseline in the first half of trials decreased as a function of age more so than did

average reactivity during the second half of trials (see Figure 3B, sFigure 32). This interaction

was in the same direction for neutral trials across most specifications (88.5% of estimates), but

was typically not as strong (3.8% of posterior intervals excluding 0). Single-trial models

indicated similar age-related change in within-scan amygdala dynamics (see Figure 3C,

sFigures 33-34). Mean group-level amygdala reactivity was higher for the first half of trials for

fear faces > baseline across several specifications, though there were not consistent differences

between trial halves for mean amygdala reactivity to neutral faces (sFigure 31).

Figure 3. Age-related change in amygdala reactivity across trials. **A.** An example model of
estimated age-related change in slopes of beta estimates across both fear (green) and neutral
(orange) trials. Negative slopes represent higher amygdala activity in earlier trials relative to
later trials. **B.** Example models of estimated age-related change in amygdala reactivity for the
fear > baseline (left) and neutral > baseline (right) contrasts for both the first (red) and second
(blue) halves of trials. In both A and B, points represent participant-level estimates, light lines
connect estimates from participants with multiple study visits, and dark lines with shaded area
represent model predictions and 95% posterior intervals. **C.** Example single-trial model
predictions of estimated amygdala reactivity for fear (left) and neutral (right) faces as a function

of age and trial number. Age was modeled as a continuous variable, and average predictions for
participants of age 6 (red), 12 (green) and 18 (blue) years are shown for visualization purposes.
All estimates in A-C shown are from an example analysis pipeline using bilateral amygdala
estimates and without global signal correction. **D.** Specification curve for age-related change in
slopes across fear trials (i.e., many parallel analyses for the fear trials in subplot B). **E.**
Specification curve for age-related change in slopes across neutral trials (i.e., neutral trials in
plot B). For both D and E, color indicates sign of beta estimates and whether respective
posterior intervals include 0 (green = positive including 0, purple = positive excluding 0, black =
median across all specifications), and horizontal dotted lines represent median estimates across
all analysis decisions. Variables on the y-axis represent analysis choices, corresponding color-
coded marks indicate that a choice was made, and blank space indicates that the choice was
not made in a given analysis.
176

Age-related change in task-evoked amygdala–mPFC functional connectivity

We used multilevel regression modelling and specification curve analyses to examine

age-related change in task-evoked amygdala–mPFC functional connectivity within the

accelerated longitudinal cohort (see Table 1 Aims 2a-b). For the fear > baseline contrast, a

specification curve with 288 total specifications (4 definitions of participant-level gPPI estimates

x 4 mPFC ROIs x 18 group-level models) of amygdala–mPFC gPPI did not find consistent

evidence of age-related change, as 59.0% of models found point estimates in the positive

direction, and 23% of posterior intervals excluded 0 (Figure 4C-D, interactive version at

https://pbloom.shinyapps.io/amygdala_mpfc_multiverse/). Through specification curve analyses,

we found that the sign of the estimated age-related change depended almost entirely on

deconvolution, such that most specifications including a deconvolution step resulted in negative

age-related change estimates never distinguishable from 0 (78.5% of point estimates negative,

0% of posterior intervals excluding 0), and most specifications not including a deconvolution

step resulted in positive age-related change estimates (96.5% of point estimates positive, 47.9%

of posterior intervals excluding 0). A visualization of the effects of the deconvolution step on

amygdala FC with each of four mPFC ROIs is presented in Figure 4B. While mPFC ROI

definition and other analysis decision points also influenced estimates of age-related change in

gPPI (Figure 4D), follow-up regression models indicated that the effect of including the

deconvolution step was several times larger for the fear > baseline contrast (see sFigures 41-

43).

Through equivalent multiverse analyses we also found no evidence of consistent linear

age-related change in amygdala–mPFC gPPI for the neutral > baseline and fear > neutral

contrasts (see sFigures 39-40), or quadratic change for any contrast (see sFigures 44-46). In

addition, we did not see consistent evidence for group average amygdala–mPFC gPPI for any

contrast, though such results often differed as a function of whether a deconvolution step was

included (see sFigure 38). Though we included gPPI analysis specifications excluding the 42

scans at timepoint 1 studied by Gee et al.[10], exclusion of these scans had little impact on age-

related change results (see Figure 4D).

Figure 4. Multiverse analyses of age-related change in amygdala–mPFC connectivity using
gPPI methods. **A.** MNI space mPFC ROIs used in connectivity analyses. **B.** Example
participant-level data and model predictions for age-related related change in amygdala–mPFC
gPPI for analysis pipelines with a deconvolution step (red), or without (blue) for each of the four
regions shown in A. **C.** Specification curve of age-related change in fear > baseline amygdala–
mPFC gPPI. Points represent estimated linear age-related change and lines are corresponding
95% posterior intervals. Models are ordered by age-related change estimates, and the dotted

line represents the median estimate across all specifications. Color indicates sign of beta
estimates and whether respective posterior intervals include 0 (blue = negative including 0,
green = positive including 0, purple = positive excluding 0, black = median across all
specifications). Black points with error bars represent the median and IQR ranks of
specifications making the choice indicated on the corresponding line. **D.** Model specification
information corresponding to each model in C. Variables on the y-axis represent analysis
choices, corresponding color-coded marks indicate that a choice was made, and blank space
indicates that the choice was not made in a given analysis. Within each category (Group-Level
Model, mPFC ROI, and Participant-Level Model) respectively, decision points are ordered from
top to bottom by the median model rank when the corresponding choice is made (i.e., choices at
the top of each panel tend to have more negative age-related change estimates). See
https://pbloom.shinyapps.io/amygdala_mpfc_multiverse/ for interactive visualizations.
228

In addition to gPPI analyses, we used beta series correlation (BSC) analyses to examine

age-related changes in task-evoked amygdala–mPFC connectivity (see Table 1 Aim 2b). As

with gPPI, multiverse analyses of amygdala–mPFC BSC (168 total specifications; 3 amygdala

ROI definitions x 4 mPFC ROI definitions x 2 global signal options x 7 group-level models) for

fear trials (vs baseline) did not yield consistent evidence of age-related change across pipelines

(84.5% of point estimates in the same direction as the median estimate, 24.4% of posterior

interval excluding 0; Figure 5A, interactive version at

https://pbloom.shinyapps.io/amygdala_mpfc_multiverse). Unlike gPPI analyses, however,

choice of mPFC ROI (as well as amygdala ROI, though this was not examined for gPPI) most

impacted age-related change in BSC estimates, rather than preprocessing or modeling

analytical choices (Figure 5B, sFigures 53-55). Accordingly, while global signal subtraction

resulted in weaker amygdala–mPFC BSC on average, inclusion of this step did not consistently

affect age-related change estimates (Figure 4C). We did not find consistent evidence for age-

related change in amygdala–mPFC BSC for neutral trials (vs baseline), or for fear > neutral

trials (sFigures 51-52). We did not find consistent evidence for quadratic age-related change for

any contrast (sFigures 56-58).

Additionally, we constructed a correlation matrix using rank-order correlations of scan-

level BSC and gPPI estimates for the fear (vs baseline) condition. Across scans, there was little

evidence of correspondence between BSC and gPPI metrics for amygdala–mPFC connectivity

(Figure 5D, sFigures 60-63). Further, FC estimates tended to be positively correlated within a

method type (BSC, gPPI) across mPFC ROIs, though less strongly for gPPI estimates with

versus without a deconvolution step.

In addition to gPPI and BSC methods for functional connectivity, we also explored

between-scan associations between amygdala reactivity and mPFC reactivity (sFigures 28-29).

Multilevel models indicated that amygdala reactivity for fear faces > baseline was positively

associated with mPFC reactivity for fear faces > baseline for all mPFC ROIs, though we did not

find consistent evidence for age-related changes in associations between amygdala and mPFC

reactivity to fear faces > baseline (see sFigure 29).

258
Figure 5. Multiverse analyses of age-related change in amygdala–mPFC connectivity using
beta-series correlation (BSC) methods. **A.** Specification curve of age-related change in
amygdala–mPFC BSC for fear trials. Points represent estimated linear age-related change and
lines are corresponding 95% posterior intervals. Models are ordered by age-related change
estimates, and the dotted line represents the median estimate across all specifications. Color
indicates sign of beta estimates and whether respective posterior intervals include 0 (blue =
negative including 0, green = positive including 0, purple = positive excluding 0, black = median

across all specifications). **B.** Model specification information corresponding to each model in A.
Variables on the y-axis represent analysis choices, corresponding color-coded marks indicate
that a choice was made, and blank space indicates that the choice was not made in a given
analysis. Within each category (amygdala ROI, group-level model, global signal subtraction, and
mPFC ROI) respectively, decision points are ordered from top to bottom by the median model
rank when the corresponding choice is made (i.e., choices at the top of each panel tend to have
more negative age-related change estimates). Black points with error bars represent the median
and IQR ranks of specifications making the choice indicated on the corresponding line. **C.**
Example model predictions for age-related change in amygdala–mPFC BSC for fear trials for
analysis pipelines with a global signal subtraction (GSS, post-hoc mean centering) step (red), or
without (blue) for each of the four mPFC regions (see Figure 4A) with the left and right
amygdala. Pipelines shown have random slopes, no covariates for task block or scanner, and
no quadratic age term. **D.** Between-scan rank-order correlations between amygdala–mPFC
connectivity measures. All gPPI measures are for the fear > baseline contrast, and BSC
measures are for fear trials. See https://pbloom.shinyapps.io/amygdala_mpfc_multiverse/ for
interactive visualizations.
282
283

Amygdala–mPFC Measures & Separation Anxiety

We conducted multiverse analyses of associations between several amygdala–mPFC

measures (amygdala reactivity, amygdala–mPFC FC, within-scan changes in amygdala

reactivity) and separation anxiety behaviors (see Table 1 Aim 3). Separation anxiety behaviors

on average decreased with age, as indicated by the RCADS-P and SCARED-P raw scores

(Figure 6A-C). Neither specification curves for amygdala reactivity (18 total specifications, 56%

of point estimates in the same direction as median estimate, 0% of posterior intervals excluding

0), amygdala–mPFC FC (90 total specifications, 51% of point estimates in the same direction as

median estimate, 1% of posterior intervals excluding 0), nor slope of amygdala responses

across trials (12 total specifications, 75% of point estimates in the same direction as median

estimate, 17% of posterior intervals excluding 0), found consistent evidence for associations

between brain measures and separation anxiety. All specifications controlled for age (see

supplemental methods p.26).

To more specifically follow up on previous work reporting associations between

separation anxiety behaviors and amygdala–mPFC gPPI for fear > baseline, we plotted model

299   predictions for such models from the above multiverse analysis for each of the four mPFC ROIs,

300   across all three separation anxiety outcome measures, and both with and without a

301   deconvolution step (Figure 6E). We did not find consistent evidence for associations with

302   separation anxiety, and results showed high sensitivity to the deconvolution step, mPFC ROI,

303   and outcome measure used.

304
Figure 6. Multiverse analyses of associations between amygdala–mPFC circuitry and
separation anxiety. **A-C.** Age-related change in SCARED and RCADS raw and t-scores for
parent-reported separation anxiety subscales. The red dotted line in B represents the clinical
threshold. **D.** Separate specification curves for associations of amygdala reactivity (left),
amygdala–mPFC connectivity (both gPPI and BSC; center), and amygdala reactivity slopes
across trials (right) with the three separation anxiety outcomes shown in A-C. Points represent
estimated associations between brain measures and separation anxiety (controlling for mean
FD and age) and lines are corresponding 95% posterior intervals. Models are ordered by beta
estimates, and the dotted line represents the median estimate across all specifications. Color
indicates sign of beta estimates and whether respective posterior intervals include 0 (red =
negative excluding 0, blue = negative including 0, green = positive including 0). Scores on each
separation anxiety outcome were z-scored for comparison. **E.** Example model predictions for
associations between fear > baseline amygdala–mPFC gPPI and each separation anxiety
measure. Predictions and 95% posterior intervals are plotted for each separation anxiety
measure separately for each mPFC region, and for gPPI pipelines with and without a
deconvolution step. Pipelines shown use robust regression, have random slopes, no covariates
for task block or scanner, and no quadratic age term.
322

Discussion
Our accelerated longitudinal design and multiverse analysis approach afforded unique

opportunities to examine age-related changes in task-evoked amygdala-prefrontal activity

within a cohort recruited from the Los Angeles area. While there were differences across

model specifications, the majority of pipelines supported our hypothesis that amygdala

reactivity to fearful faces decreases with age from early childhood through early adulthood

(see Table 1 Aim 1a). We did not find evidence for our second hypothesis, as neither gPPI nor

BSC analyses indicated consistent evidence of age-related change in amygdala–mPFC

functional connectivity (see Table 1 Aims 2a-2b). Specification curves also revealed that gPPI

analyses were highly sensitive to analysis methods (deconvolution). We did not find

associations between any task-related amygdala–mPFC measures (reactivity or functional

connectivity) and separation anxiety behaviors (see Table 1 Aim 3).

Amygdala reactivity
Across specifications, we found evidence for age-related decreases in amygdala

reactivity to both fearful and neutral faces (Figure 2A). Yet, findings also varied considerably

across specifications. For example, only 60% of pipelines produced results that would be

individually labeled as 'significant' (under α = .05), indicating that multiple investigations of this

dataset could likely lead to qualitatively different conclusions. While over half of analyses

found evidence consistent with studies indicating greater amygdala reactivity to fear faces >

baseline in younger children[2,10,11,14], the other 40% of specifications would have been

consistent with investigations that found little age-related change[23,26,28].

Models also found evidence for between-participant differences, but could neither

identify within-participant change (Figure 2D) nor differentiate between-participant from within-

participant estimates. As such, interpretation of the age-related change reported here is

subject to many of the same limitations that apply to cross-sectional designs[46], where age-

related changes may not necessarily indicate 'true' developmental growth. High uncertainty in

estimating average within-participant change could be driven by several factors, including true

heterogeneity in individual trajectories, low measurement reliability[47–49], scanner differences

across longitudinal timepoints, or unmodeled variables impacting amygdala reactivity.

Additionally, the within-participants terms represent a smaller age range (a maximum of 4

years for any given participant), relative to the broader age range assessed by the between-

participants terms (18 years), which may have placed additional limits on identifying reliable

within-participant change.

Age-related change in amygdala responses to fear faces over baseline seemed largely

the result of earlier trials in the task (see sFigures 32-34). While differences in task design and

contrast across studies have been highlighted as potential sources of discrepant findings on the

development of amygdala function[14,32,50], this result indicates that attention to trial structure and

task duration may also be necessary in comparing studies. Because the paradigm used in the

current study involved a task requiring participants to press for one face ('neutral') and not press

for 'fear' faces, findings specific to fear faces over baseline under the current paradigm may also

be driven by behavioral task demands.

Amygdala–mPFC Functional Connectivity

Multiverse analyses did not find consistent evidence for age-related change in

amygdala–mPFC connectivity using gPPI or BSC (Figures 4-5). Crucially, however,

specification curves did not find strong evidence against such age-related change, as we did

not observe precise and consistent 'null' estimates across specifications. This absence of

evidence for age-related changes in task-evoked amygdala–mPFC connectivity contrasts prior

work suggesting a developmental shift in such connectivity[10,23,26], but is consistent with recent

work[28].

gPPI results were sensitive to whether a deconvolution step had been included in the

preprocessing pipeline, such that we mostly found age-related decreases in amygdala–mPFC

connectivity with a deconvolution step included, and age-related increases without it (see

Figure 4B). While deconvolution has been argued to be a necessary step for event-related

PPI analyses[51], recent work has shifted guidelines on its use, and it may not be recommended

for block designs [39,52]. Because the true 'neuronal' signal underlying the BOLD timeseries

within a given ROI cannot be directly measured, deconvolution algorithms are difficult to

validate. Further, deconvolution may cause PPI results to be driven by baseline connectivity

depending on the centering of the task regressors[53]. Within the current study, small tweaks to

AFNI's 3dTfitter algorithm for deconvolution resulted in vastly different regressors (see

sFigure 36), suggesting the potential for high analytic variability even between gPPI analyses

ostensibly using deconvolution. While the present study does not provide evidence that can

inform whether or not deconvolution is recommended, further work is needed to optimize and

validate applications of gPPI methods and selection of appropriate task designs. gPPI may be

better equipped for block-designs and particularly ill-posed for rapid event-related tasks due to

both difficulties in resolving which times within the BOLD timeseries reflect functional

connectivity evoked by rapid (350ms) events and low statistical power in estimating such task-

evoked connectivity[54] (see supplemental results p.62-66, sFigures 35-37). Concurrent with

previous work, beta series correlation analyses may have higher statistical power for

identifying task-related connectivity signal than gPPI within event-related designs more

generally[40].

Amygdala–mPFC circuitry and separation anxiety

Specification curve analyses did not yield consistent evidence of associations between

any measures of amygdala–mPFC function (amygdala reactivity & change in amygdala

reactivity across trials, in addition to FC) and separation anxiety behaviors. This finding stands

in contrast to associations between amygdala–mPFC connectivity and anxiety identified in

previous developmental work[10,23,55,56]. However, given that analyses of brain-behavior

associations may require imaging cohorts much larger than the current sample [31,57], the

absence of relationships here may not be strong evidence against the existence of potential

associations between amygdala–mPFC circuitry and developing anxiety-related behaviors.

Advantages and pitfalls of the multiverse approach

407   Our findings contribute to a body of work demonstrating that preprocessing and

modeling choices can meaningfully influence results[34]. Indeed, most studies involving many

analytical decision points could benefit from multiverse analyses. Such specification curves

can help to examine the stability of findings in both exploratory and confirmatory research[58].

Particularly when methodological 'gold standards' have not been determined, specification

curves may be informative for examining the impacts of potential analysis decisions[59]. Further,

wider use of specification curves might help to resolve discrepancies between study findings

stemming from different analysis pipelines.

415   While specification curve analyses may benefit much future research, we also note

that multiverses are only as comprehensive as the included specifications[60], and such

analyses alone do not solve problems related to unmodeled confounds, design flaws,

inadequate statistical power, circular analyses, or non-representative sampling. Further,

unless all specifications are decided *a priori*, analyses are vulnerable to problems of analytic

flexibility[61], and inclusion of less justified specifications can bias results[62]. Because

specification curves can include hundreds or thousands of individual analyses, rigorous

evaluation of individual models can be difficult. To this end, we created interactive

visualizations for visual exploration of individual analysis specifications.

Limitations

The present study is subject to several limitations that may be addressed in future

investigations. Perhaps most crucially, our conclusions (along with those of many

developmental fMRI studies) are limited by the number of participants studied[31,63], the number

of longitudinal study sessions per participant[44], and the duration of the task[64]. In particular, the

low statistical power of our rapid event-related task design may be a major contributor to the

variance in outcomes across analysis specifications, as low-powered studies can yield

increased rates of both false positive and false negative results (as well as errors of the sign

and magnitude of estimates[65,66]). Work with larger sample sizes, more study sessions per

participant, and more task data collected per session will be necessary for charting functional

amygdala–mPFC development and examining heterogeneity across individuals (although

collecting task-based fMRI will continue to be challenging for studies including younger

children). The generalizability of the current findings may also be limited by the fact that this

cohort was skewed towards high incomes and not racially or ethnically representative of the

Los Angeles or United States population.

Findings are also somewhat limited by the fact that the present study is not wholly

confirmatory, despite preregistration. Because our multiverse analysis approaches expanded

significantly beyond the methods we preregistered, most of the present analyses, while

hypothesis-driven, must be considered exploratory[58]. The fact that some specifications used

data included in previous similar analyses of the same cohort[10] also limits the confirmatory

power of the present study[67]. This may be especially true because longitudinal models could

not identify within-person change as distinct from between-participant differences (see Figure

2D), indicating that our age-related change estimates may be influenced by cross-sectional

information similar to that investigated by Gee et al.[10].

Though the current study aimed to estimate longitudinal age-related changes in

amygdala–mPFC functional circuitry evoked by fear and neutral faces, the current findings

may not be specific to these stimuli[9]. Because our task did not include non-face foils or probe

specific emotion-related processes, results may be driven by attention, learning, or visual

processing, rather than affective or face processing. Findings for the fear faces > baseline and

neutral > baseline contrasts also may not be valence-specific in the absence of a different

emotional face as part of the contrast. Further, because all faces were adult White women, the

current results may not generalize to faces more broadly[68,69].

Finally, our findings on age-related change in amygdala and mPFC function may be

biased or confounded by age-related differences in head motion[70], anatomical image quality

and alignment[71,72], signal dropout, and physiological artifacts[73–75]. While our multiverse

analyses included preprocessing and group-level modeling specifications designed to

minimize some of such potential issues, future work is still needed to optimize discrimination

of developmental changes of interest from such potential confounds.

Despite these limitations, the present study concurs with prior investigations in finding

evidence for age-related decreases in amygdala reactivity to fear faces over baseline. Though

our work did not find evidence for previously reported developmental 'shifts' in positive to

negative task-related amygdala–mPFC functional connectivity, our results highlight that both

longitudinal study designs and specification curve analyses can be valuable tools for

examining the robustness of such analyses, as well as for studies of human brain

development more broadly.

Table 1: Summary of main aims, hypotheses, methods, and findings

Aim	Preregistered Hypothesis	Analysis Methodology	Key Findings
1a. Age-related change in amygdala reactivity to fear faces	Amygdala reactivity to fearful faces will decrease with age, such that younger children will have more positive amygdala reactivity (higher BOLD response to fear faces relative to implicit baseline) than older youth.	Multiverse amygdala ROI (anatomically-defined) analysis using multilevel linear regression at the group level. *Multiverse decision points:* Preprocessing software, GLM software, GLM nuisance regressors, amygdala ROI definition, contrast estimate type (t-stat vs. beta estimate), HRF shape, group-level model covariates, exclusion of previously analyzed scans	• Across decision points, weak but consistent negative age-related change in amygdala reactivity to fear > baseline and neutral > baseline contrasts • No consistent evidence for age-related change in fear > neutral contrast • Longitudinal models could identify consistent between-participant differences but not within-participant age-related change
1b. Age-related change in patterns of amygdala responses across task trials	None	Multiverse analysis of slopes of amygdala reactivity across trials, and amygdala reactivity in each half of trials using multilevel linear regression at the group level, single trial models *Multiverse decision points:* Global signal subtraction, amygdala ROI definition, group-level model covariates	• On average, amygdala reactivity decreased across trials (for both fear and neutral faces) • Amygdala reactivity for earlier trials was higher at younger ages • Age-related change in amygdala reactivity to fear faces in the first half of trials, but not the second half • Similar, but somewhat weaker age-related change for neutral faces
2a. Age-related change in amygdala–mPFC functional connectivity (FC) to fear faces, as measured by generalized psychophysiological interaction (gPPI)	Amygdala–mPFC FC will decrease as a function of age such that as age increases, the valence of FC will shift from positive to negative.	Multiverse gPPI analysis with anatomically defined bilateral amygdala seed and mPFC target ROIs using multilevel linear regression at the group level. *Multiverse decision points:* Deconvolution step, mPFC ROI definition, contrast estimate type (t-stat vs. beta estimate), group-level model covariates	• No consistent evidence for age-related change in gPPI for any contrast • gPPI estimates extremely sensitive to deconvolution step in creation of regressors
2b. Age-related change in amygdala–mPFC functional connectivity to fear faces, as measured by beta-series correlation (BSC)	None for BSC specifically	Multiverse BSC analysis between amygdala and mPFC using multilevel linear regression at the group level. *Multiverse decision points:* Global signal subtraction, amygdala ROI definition, mPFC ROI definition, group-level model covariates	• No consistent evidence for age-related change in BSC for any condition • Amygdala–mPFC BSC was most sensitive to selection of mPFC ROI • Global signal subtraction reduced average amygdala–mPFC BSC, but impacts on age-related changes were small • BSC estimates were not strongly associated with gPPI estimates
3. Associations of amygdala reactivity, change in amygdala reactivity across trials, or amygdala–mPFC FC with separation anxiety	None	Multiverse multilevel linear regressions with brain measures as predictors for separation anxiety behaviors, controlling for age *Multiverse decision points:* Separation anxiety measure, FC measure, mPFC ROI (FC only), amygdala ROI, contrast, deconvolution step (gPPI only)	• No evidence that amygdala reactivity, amygdala–mPFC connectivity, or change in amygdala reactivity across trials were associated with separation anxiety behaviors

Methods

Before completing analyses, we preregistered methods for the current study through the

Open Science Framework at https://osf.io/8nyj7/. Only analyses for age-related changes in

amygdala reactivity and amygdala–mPFC gPPI were preregistered in detail (see Table 1 Aims

1a & 2a), and we did not preregister multiverse analyses. Methods detailed below include both

information included in the preregistration and additional information and analyses not

preregistered. Analysis code & documentation can be found at

https://github.com/pab2163/amygdala_mpfc_multiverse.

Participants

Participants were recruited as part of a larger study examining brain development as a

function of early life caregiving experiences. The current sample (N=98; 55 female, 43 male)

included typically developing children, adolescents, and young adults covering the ages 4-22

years-old (M = 11.9 years old) who enrolled to participate in a study on emotional development.

All participants were reported to be physically and psychiatrically healthy (no medical or

psychiatric disorders), as indicated by a telephone screening before participation, and free of

MRI contraindications. All except 4 participants fell below clinical cutoffs (see sFigure 2) on the

Child Behavior Checklist (CBCL) Total Problems, Internalizing Problems, and Externalizing

Problems scales[76]. The larger study also included youths with a history of institutional and/or

foster care outside of the United States, who are not included here. Participants from the greater

Los Angeles area were recruited through flyers, state birth records, community events, online

advertising, lab website and newsletters, psychologists' offices, psychology courses at a local

university (participants ages 18-22 years old only), and word-of-mouth. Each participant

completed up to 3 MRI scans spaced at an average interval of 18 months between visits.

Parents provided written consent, children 7+ years old gave written assent, and children under

7 years old gave verbal assent. Study protocols were approved by the local university

institutional review board. These data were collected between 2009 and 2015.

An accelerated longitudinal design was used such that participants' starting ages at scan

1 comprised the entire range of sample ages (4-22 years old), and coverage was approximately

balanced across the entire age range (see Figure 1B). The design was structured into 3 study

'waves' corresponding with recruitment efforts and visit protocols. Participants were

overenrolled at wave 1 to account for anticipated attrition (e.g., braces, relocation, etc) to

achieve the desired sample size across the three waves. While most participants were recruited

such that their first scan occurred at wave 1, a smaller set of participants were recruited at wave

2, such that some participants completed their first scan at wave 2 (see Figure 1). For these

participants, only 2 scans were planned.

Of the 191 participants participating in the longitudinal study, 140 completed at least one

MRI scan. After exclusions for incomplete task runs (including falling asleep), computer errors

resulting in missing stimulus timing files, high head motion, and failed visual QA

(scanner/motion artifacts), a final sample of 98 participants (N = 183 total scans) was included

for analysis (see Figure 1). Exclusion criteria were preregistered after conducting preliminary

preprocessing, but before construction of group-level models and multiverse analysis plans.

This sample included 40 participants with 1 scan, 31 with 2 scans, and 27 with 3 scans (one

more participant than preregistered due to an initial coding error). Wave 1 data from forty-two of

these participants were reported on by Gee et al.[10].

The median annual household income for participating families was $85,001-$100,000

(for reference, median annual household income in Los Angeles County from 2015-2019 was

$68,044; US Census Bureau, 2021). Epidemiological methods were not used to recruit a

sample representative of the Los Angeles or United States populations[78], and Hispanic or Latinx

participants were particularly underrepresented. Further sample demographics can be found in

the supplementary materials (see sTables 1-2, sFigures 1-2).

Separation Anxiety

For each participant (except for 10 adults 18-22 years), a parent completed both the

Revised Children's Anxiety and Depression Scale (RCADS-P) and the Screen for Child Anxiety

Related Emotional Disorders (SCARED-P) to assess the frequency of symptoms of anxiety and

low mood[79,80]. Following prior work suggesting associations between task-evoked amygdala–

mPFC functional connectivity and separation anxiety[10,43], we used the separation anxiety

subscales from both the SCARED-P and RCADS-P as measures of anxiety-related behaviors in

asking whether such functional connectivity may be linked to anxiety levels during childhood

and adolescence. For 11 participants who had missing items on the SCARED-P, indicating

parents had skipped or forgotten to answer a question, we imputed responses using 5-Nearest

Neighbor imputation using only the other items included in the SCARED-P separation anxiety

subscale[81]. As expected, raw separation anxiety scores on both measures decreased as a

function of age, while standardized scores were consistent across development with few

children at or near clinical threshold (see Figure 6, and supplemental results p.31 for intraclass

correlation coefficients).

Emotion Discrimination Task

Participants completed either two (at wave 3) or three (at waves 1 and 2) runs of a

modified 'go/no-go' task with emotional faces during fMRI scanning. Runs varied by emotional

expression (fear, happy, sad), and within each run participants viewed emotional faces

interspersed with neutral faces. To ensure that participants were paying attention, they were

asked to press a button whenever they saw a neutral face (no response was required for any

other face expression). The order of the runs was counterbalanced across participants; the

stimuli within each run were pseudorandomized[82] to allow for event-related estimates of the

hemodynamic response, and fixed across participants. For the present analysis, only the fear

run of the task was used. The other two runs, which used happy and sad faces in place of fear,

are not included in the present analysis as these conditions were not present at all waves of

data collection. As 50% of trials were 'go' trials under this paradigm, we refer to the task as an

emotion discrimination task, rather than a true 'go/no-go' paradigm since there was no strong

prepotent motor response. Stimuli within each run were presented with a jittered ITI (3-10s,

Median = 4.93s]) according to a genetic algorithm with a fixation cross on the screen[82]. Face

images were adult White female faces from the Karolinska Directed Emotional Faces database

[83]. Each run (130 TRs, duration of 4:20) consisted of 48 trials (24 neutral faces, 24 fearful

faces), each presented for 350ms. All fMRI analyses of this task used event-related designs.

MRI Acquisition

Participants under 18-years-old completed a mock scanning session before the MRI

scan to acclimate to the scanner environment and practice lying still for data collection. Waves 1

and 2 were collected on a Siemens 3T TIM Trio MRI scanner using a standard radiofrequency

head coil. A 2D spin echo image (TR, 4000 ms; TE, 40 ms; matrix size, 256 x 256; 4 mm thick;

0mm gap) was acquired in the oblique plane to guide slice configuration in both structural and

functional scans. A whole-brain high-resolution T1-weighted anatomical scan (MPRAGE; 256 x

256 in-plane resolution; 256mm FOV; 192 x 1 mm sagittal slices) was acquired for each

participant for registration of functional data. The task was presented through MR-compatible

goggles during scanning. T2*-weighted echoplanar images (interleaved slice acquisition) were

collected at an oblique angle of ~30 degrees (130 volumes/run; TR=2000ms; TE=30 ms; flip

angle=90°; matrix size=64 x 64; FOV=192 mm; 34 slices; 4 mm slice thickness; skip=0 mm).

Wave 3 was collected on a Siemens 3T TIM Trio MRI scanner at a different location using

identical acquisition parameters.

Behavioral Analyses

We used multilevel logistic regression models to estimate age-related changes in

several task performance metrics. We fit separate models for the d' performance metric, overall

accuracy (probability of a correct response on any trial), hit rate (on neutral face trials), and false

alarm rate (on fear face trials) as the respective outcomes, and included nested random effects

for task sessions within participants (models were not nested for d' as this analysis used only 1

metric per session rather than trial-wise outcomes, but still included random effects for

participants). Additionally, to model age-related change in reaction times during correct hit trials,

we fit linear, quadratic, and cubic regressions with identical random effects structures. Model

equations and results for all behavioral analyses can be found in the supplement (see

supplemental methods p.8-9, sFigures 3-4).

Preregistered fMRI Pipeline

MPRAGE scans were skull-stripped using 3dSkullStrip from the Analysis of Functional

NeuroImages (AFNI) software package[84] or FSL's brain extraction tool (BET)[85], depending on

visual inspection of image quality. For each scan, an experimenter visually compared skull-

stripped images using both software packages and chose the one judged to be more accurate.

Slice-time correction was not used. Timeseries of the 6 motion parameters were calculated and

subsequent spatial realignment of BOLD volumes was completed using MCFLIRT in FSL[86].

Scans over a threshold of >40 volumes with > .9mm framewise displacement were excluded

from analysis (12 out of an initial 195, or 6.2%). The mean age of participants with excluded

scans was 7.16 years and 8/12 were male. Registration matrices were calculated for registration

of functional images to high-resolution structural T1 images using FSL's FLIRT with boundary-

based registration. Registration matrices for standard MNI space were also calculated using

both FLIRT (linear registration) and FNIRT (nonlinear registration) with 12 DOF and a warp

resolution of 10mm. Data were high-pass filtered at .01Hz and smoothed with an isotropic

Gaussian kernel with FWHM of 6mm before running general linear models (GLMs), and 4d

volumes were grand mean scaled such that the average intensity value was 10000.

Following preprocessing, we ran scan-level GLMs using FSL's FEAT. We included

event-related regressors for fear and neutral faces (convolved with a double-gamma HRF), their

temporal derivatives[87], and 24 head motion nuisance regressors (the 6 head realignment

parameters, their temporal derivatives, and their squares[88]. Volumes with FD > .9mm were

downweighed to 0 in the GLM. Pre-whitening was used to estimate and remove temporal

autocorrelation[89]. For each scan, we calculated fear > baseline, neutral > baseline, and fear >

neutral contrasts. We used native-space bilateral amygdala masks generated using Freesurfer

(v6.0[90]) by VanTieghem et al.[91].

Multiverse Analyses and Specification Curves

In addition to the preregistered pipelines, we conducted multiverse analyses to address

all aims in Table 1 and constructed sets of separate specification curves for each aim (see

Table 2). In general, multiverse analyses aim to probe the consistency of results across all

'reasonable' possible combinations of analysis decisions (i.e. simultaneously taking all possible

'forking paths')[60]. Because analyzing fMRI data using all reasonable specifications was

infeasible (i.e., possibilities are virtually infinite), we took the approach of 'sampling' from the

many reasonable or commonly-used analysis choices for each multiverse. Despite not being

completely comprehensive, this approach still allowed for thorough investigation into the

robustness of results. For all multiverse analyses, we constructed specification curves by

ranking models by their beta estimates (ascending) for parameters of interest for interpretation

and visualization[92–96]. Because specification choices were not preregistered, we did not conduct

formal null hypothesis testing of specification curves. Instead, we used a rough rule-of-thumb for

whether a finding was 'robust', or consistent across specification choices, by asking whether

more than 50% of the specifications resulted in a 95% posterior interval excluding 0 in the same

direction. In addition, to analyze in more detail the impact of specific choices, we submitted point

estimates for parameters of interest across all specifications to multiple regression models.

From these models, we examined the conditional effects of each analysis decision point on the

parameter of interest (see supplemental methods p.26, sFigures 11-13, 41-43, & 53-55).

Multiverse amygdala reactivity analyses

For amygdala reactivity analyses, we examined the robustness of age-related change

estimates to a variety of analytical decisions. In addition to the preregistered FSL-based

pipeline, we preprocessed data using C-PAC software (v1.4.1[97]). We used C-PAC to take

advantage of features supporting running multiple pipeline 'forks' in parallel (for example

multiple nuisance regression forks using the same registration). No spatial smoothing was used

in C-PAC pipelines (see supplemental methods p.10). Following C-PAC and FSL

preprocessing, we examined the impact of different sets of commonly-used analysis methods

on age-related change in amygdala reactivity. We varied analysis choices of GLM software,

hemodynamic response function, nuisance regressors, first-level GLM estimates, amygdala

ROI, exclusion criteria (exclude vs. include scans analyzed by Gee et al.[10]), group-level model

outlier treatment, and group-level model covariates (see Table 2 & supplemental methods p.10-

13). Multiverse analyses of amygdala reactivity included a total of 2808 model specifications

(156 ways of defining participant-level amygdala reactivity x 18 group-level model

specifications) for each contrast. We analyzed all specifications in parallel. In addition, we

examined quadratic age-related changes (see sFigures 14-17) and ran a smaller set of

analyses (24) to ask whether we could differentiate within-participant change over time from

between-participant differences through alternative model parametrization (see supplemental

methods p.14).

For all specifications, individual-level amygdala reactivity estimates were submitted to a

group-level multilevel regression model for estimation of age-related changes. All models

allowed intercepts to vary by participant, and some specifications also allowed for varying

slopes (see supplemental methods p.15 for model syntax). All models also included a scan-level

covariate for head motion (mean FD[88,98,99]). Consistent with prior work, head motion was higher

on average in younger children, and decreased with age (see sFigure 5), though head motion

was not associated with amygdala reactivity estimates for most specifications (see sFigure 26).

Age-related change findings examined for the preregistered pipeline also remained consistent

under more stringent exclusion thresholds based on mean FD (see sFigure 27).

Across preprocessing specifications, we also examined within-scan similarity of

amygdala and whole-brain voxelwise reactivity patterns (see sFigures 19-20) and between-scan

correlations of average amygdala reactivity estimates (sFigures 21-23). To understand the

proportion of variance in amygdala reactivity explained by repeated measurements of the same

participants across study timepoints, we estimated intraclass correlation coefficients (ICC) for

amygdala reactivity for each preprocessing specification (see sFigure 23).

672 Table 2: Summary of forking pipelines used in analyses for each aim[1]

Aim/Analysis	Decision Point	Choices
1a. Age-related change in amygdala reactivity to fear faces > baseline	Preprocessing Software	**FSL FEAT**, C-PAC
	GLM Software	**FSL FEAT**, AFNI 3dDeconvolve
	Hemodynamic Response Function	**Double Gamma**, Single Gamma
	Nuisance Regressors	**24 motion regressors**, 6 motion regressors, 18 motion regressors + WM + CSF
	Low-frequency artifact removal	**High-pass filter (.01Hz)**, Quadratic drift regressor
	First-level GLM Estimates	**Beta Estimates**, T-statistics
	Native vs. Standard MNI Space	**Native Space (Freesurfer)**, Harvard-Oxford Atlas in MNI
	Amygdala ROI	**Bilateral**, Left, Right, High Signal, Low Signal
	Inclusion of 45 previously analyzed scans	**Include**, Exclude
	Outlier treatment	**Exclude +/-3SD from mean**, Exclude +/-3SD from mean + robust regression
	Group-level model covariates	**Mean FD**, Mean FD + run, Mean FD + scanner, Mean FD + run + scanner
	Group-level model quadratic term	Yes, **No**
	Group-level model random slopes	**Yes**, No
1b. Age-related change in patterns of amygdala responses across task trials *FSL preproc & GLM, high-pass filter, 24 motion regressors, 2G HRF, beta estimates, included previously analyzed scans, and robust group-level regression*	Method of quantifying within-scan change	Slopes across trials, trials split into halves, single-trial models
	Global Signal Subtraction	Yes, No
	Amygdala ROI (all MNI space)	Bilateral, Left, Right
	Group-level model covariates	Mean FD, Mean FD + run, Mean FD + scanner, Mean FD + run + scanner
	Group-level model quadratic term	Yes, No
	Group-level model random slopes	Yes, No
2a Age-related change in amygdala–mPFC functional connectivity (FC) to fear faces > baseline, as measured by (gPPI) *FSL preproc & GLM, high-pass filter, 24 motion regressors, 2G HRF, bilateral amygdala ROI in MNI space*	Deconvolution step	Yes, **No**
	mPFC ROI (all MNI space)	**3 different 5mm spheres**, large vmPFC mask
	Outlier treatment	**Exclude +/-3SD from mean**, Exclude +/-3SD from mean + robust regression
	Inclusion of 45 previously analyzed scans	**Include**, Exclude
	Group-level model covariates	**Mean FD**, Mean FD + run, Mean FD + scanner, Mean FD + run + scanner
	Group-level model quadratic term	Yes, **No**
	Group-level model random slopes	**Yes,** No
2b. Age-related change in amygdala–mPFC functional connectivity to fear faces > baseline, as measured by (BSC) *FSL preproc & GLM, high-pass filter, 24 motion regressors, 2G HRF, beta estimates, robust group-level regression, included previously analyzed scans*	Amygdala ROI (all MNI space)	Bilateral, Left, Right
	mPFC ROI (all MNI space)	3 different 5mm spheres, large vmPFC mask
	Global Signal Subtraction	Yes, No
	Group-level model covariates	Mean FD, Mean FD + run, Mean FD + scanner, Mean FD + run + scanner
	Group-level model quadratic term	Yes, No
	Group-level model random slopes	Yes, No
3. Associations of amygdala reactivity, change in amygdala reactivity across trials, or amygdala–mPFC FC with separation anxiety *See supplemental methods p. 25 for details on included pipelines.*	Brain measure	Amygdala reactivity, amygdala reactivity slopes, amygdala–mPFC gPPI, amygdala–mPFC BSC
	Global Signal Subtraction (amygdala reactivity slopes & BSC only)	Yes, No
	Deconvolution step (gPPI only)	Yes, No
	mPFC ROI (all MNI space, gPPI & BSC only)	3 different 5mm spheres, large vmPFC mask
	Separation anxiety outcome variable	RCADS, SCARED raw scores, SCARED t-scores

673 [1]**Bolded** choices indicate those most closely matching preregistered pipelines.

Change in amygdala reactivity across trials

676　To probe whether amygdala reactivity exhibited within-scan change in an age-dependent

manner, we modeled reactivity to each face trial using a Least Squares Separate method

(LSS[100]). After preprocessing, we used FEAT to fit 48 separate GLMs corresponding to each

trial in each scan. A given trial was modeled with its own regressor and the remaining 47 trials

were modeled with a single regressor. Each GLM also included 24 head motion nuisance

regressors and had TRs with framewise displacement > .9mm downweighted to 0. BOLD data

were high-pass filtered at .01Hz before the GLM. From each of the 48 GLMs, we extracted the

mean amygdala beta estimates corresponding to a contrast for each single trial > baseline.

684　We constructed separate multiverse analyses using three different methods for

measuring change in amygdala reactivity across trials. For method 1 (*slopes*), we measured

rank-order correlations between trial number and trial-wise amygdala betas. For method 2 (*trial

halves*), we split trials into the first half (trials 1-12) and second half (trials 13-24), and modeled

age-related change in each half. For method 3 (*single-trial models*), we constructed larger

multilevel models with individual trials as the unit of observation. We conducted several analysis

specifications for each method (see Table 2 & supplemental methods p.17-19), and generated

corresponding specification curves.

Multiverse amygdala–mPFC functional connectivity (FC) analyses

694　We applied multiverse analysis techniques towards examining age-related changes in

amygdala–mPFC FC using gPPI and beta-series correlation (BSC) methods. Briefly, gPPI

estimates functional connectivity by constructing an interaction term between the timecourse in

a seed region of interest and a stimulus (task) regressor. Voxels whose activity is well fit by this

interaction term (a psychological-physiological interaction, or PPI) are assumed to be

"functionally coupled" with the seed region in a way that depends on the behavioral task [54,101].

BSC offers a different way of estimating functioning connectivity, by constructing 'timeseries' of

beta values (i.e., a beta series) in a condition of interest for two regions of interest, and

calculating the product-moment correlation between those beta series.

We constructed separate specification curves for age-related change in gPPI and BSC

for each contrast. Across gPPI specifications, we varied whether to use a deconvolution step in

creating interaction regressors[51,52], as well as several other analysis decision points (see Table

2 & supplemental methods p.19). The deconvolution step applies to the preprocessed BOLD

data from the seed timecourse: these data are first deconvolved to estimate the "underlying

neural activity" that produced the BOLD signal[51], then these deconvolved signals are multiplied

with the task regressor (e.g., for fear faces). Finally, this new interaction term is convolved with

a hemodynamic response to produce the BOLD functional connectivity regressor of interest. We

preregistered constructing an mPFC ROI containing 120 voxels centered at the peak

coordinates reported by Gee at al.[10] for age-related change in fear > baseline gPPI (Talairach

2,32,8; or MNI 3,35,8). However, after preregistration we discovered that these peak

coordinates were quite close to the corpus collosum, and this ROI would have contained a high

proportion of white matter voxels relative to cortical voxels (though this was not true for the

mPFC ROI identified by Gee et al.[10]). To address this issue, we instead constructed three

spherical ROIs with 5mm radii; the first centered at the above peak coordinates, the second

shifted slightly anterior, and the third shifted slightly ventral relative to the second (see Figure 4).

Lastly, to examine amygdala functional connectivity with a more broadly-defined mPFC, we also

used a 'large vmPFC' mask encompassing many of the areas within the ventromedial prefrontal

cortex derived from Mackey & Petrides[102].

For BSC analyses, we used beta estimates from the LSS GLMs described above for

analyses of within-scan change in amygdala reactivity. Across BSC specifications we varied

analyses across several decision points (see Table 2 & supplemental methods p.21), including

whether to include a correction for global signal (post-hoc distribution centering[103]). We

extracted mean beta estimates for amygdala and mPFC ROIs for each trial, then calculated

product-moment correlations between the timeseries of betas across trials (neutral and fear

separately) for both regions[39]. These correlation coefficients were transformed to z-scores, then

submitted to group-level models.

Age-related changes in gPPI and BSC were estimated using multilevel regression

models as described for the amygdala reactivity analyses (see supplemental methods p.22). We

focused primarily on linear age-related change, but also examined quadratic associations (see

sFigures 44-46 & 56-58). We separately examined group mean gPPI and BSC for each contrast

(see sFigures 38 & 50), as well as associations between mean FD and both FC measures

across specifications (see sFigures 47 & 59). Additionally, we examined mean estimates and

age-related change in 'task-independent' FC as measured by beta weight of the 'physio' term

from the seed amygdala timeseries within the gPPI model (representing baseline amygdala–

mPFC functional connectivity controlling for task-induced variance; sFigures 48-49).

Multiverse analyses of associations between amygdala–mPFC circuitry and separation anxiety

behaviors.

We used further multiverse analyses to ask whether amygdala reactivity, change in

amygdala reactivity over the course of the task, or amygdala–mPFC FC were associated with

separation anxiety behaviors. Separate specification curves were created for each brain

measure type (amygdala reactivity, amygdala reactivity change across trials, amygdala–mPFC

FC). All analyses used multilevel regression models with covariates for age, and specification

curves included both RCADS and SCARED separation anxiety subscales as outcomes (see

Table 2 & supplemental methods p.25).

Model-fitting

All statistical models fit at the group level were run in the R (v 3.6.1) computing

environment [104]. In order to most accurately model age-related changes in each of our

measures, we attempted to take into account both between-participants information and

repeated measurements within participants over time. Unless otherwise indicated, models were

estimated using Hamiltonian Monte Carlo sampling as implemented in the Stan programming

language through the brms package in R[105,106]. Unless otherwise indicated, all models used

package default weakly-informative priors and were run with 4 chains of 2000 sampling

iterations (1000 warmup) each (see supplemental methods p.14 and p.26 for syntax).

Interactive visualizations

Because static plots visualizing the model predictions for all models in each multiverse

would require far more page space than available, we created web-based interactive

visualization tools for exploring different model specifications and viewing the corresponding raw

(participant-level) data and fitted model predictions using R and Shiny[107]. These visualizations

can be found at https://pbloom.shinyapps.io/amygdala_mpfc_multiverse/

Deviations from preregistration

Although we largely completed the preregistered analyses, the current study includes

many analyses beyond those proposed in the initial preregistration. Because the additional

analyses (i.e., all multiverses) conducted here give us substantial analytical flexibility over that

initially indicated by preregistration, we consider all results here to be at least in part exploratory

(rather than completely confirmatory), despite the preregistered hypotheses. Additionally, we

note that BSC analyses, analyses of change in amygdala reactivity across trials, and analyses

of associations between all brain measures and separation anxiety were exploratory, and

conducted after we had seen the results of the preregistered reactivity and gPPI analyses. In

addition, to avoid possible selection bias introduced by the analytical flexibility inherent in

running many parallel analyses, we consider all analysis specifications simultaneously,

emphasizing that without further methodological work, we consider all such choices in tandem

as providing equal evidential value.

CRediT Author Statement:

783
Paul Alexander Bloom: Conceptualization, Methodology, Formal Analysis, Writing - Original
Draft, Visualization **Michelle VanTieghem:** Methodology, Writing – Review & Editing **Laurel
Gabard-Durnam:** Investigation, Methodology, Writing – Review & Editing, **Dylan Gee:**
Investigation, Methodology, Writing – Review & Editing **Jessica Flannery:** Investigation**,** Writing
– Review & Editing **Christina Caldera:** Investigation**,** Writing – Review & Editing **Bonnie Goff:**
Investigation**,** Writing – Review & Editing **Eva Telzer:** Investigation**,** Writing – Review & Editing
Kathryn L. Humphreys: Investigation**,** Writing – Review & Editing **Dominic Fareri:**
Investigation**,** Writing – Review & Editing **Mor Shapiro:** Investigation**,** Writing – Review &
Editing **Sameah Algharazi:** Validation**,** Writing – Review & Editing **Niall Bolger:** Methodology,
Formal Analysis, Writing – Review & Editing **Mariam Aly:** Methodology, Formal Analysis,
Supervision, Writing – Review & Editing **Nim Tottenham:** Conceptualization, Methodology,
Investigation, Resources, Data Curation, Writing - Original Draft, Supervision, Funding
Acquisition

797
798   **Data Availability**
799   Because participants and their parents did not consent to data sharing at the time of
800   participation, we cannot make data from the current study publicly available.

801
802   **Code Sharing & Materials Availability**

803
Code for preprocessing, analysis, and data visualizations for this manuscript is
available at https://github.com/pab2163/amygdala_mpfc_multiverse. While unfortunately this
code cannot be run as written without data, we have attempted to document analysis steps
clearly. In addition, we provide publicly available simulated data structured similarly to the
study data on amygdala reactivity, such that interested readers can view multiverse analysis
walkthroughs (https://pab2163.github.io/amygdala_mpfc_multiverse) and experiment with
analysis code. Additional materials, including MNI space masks and preregistration
documentation, are available at https://osf.io/hvdmx/

812
813

References

1.   Adolphs R. Fear, Faces, and the Human Amygdala. *Curr Opin Neurobiol*. 2008;18(2):166-
172. doi:10.1016/j.conb.2008.06.006

2.   Forbes EE, Phillips ML, Silk JS, Ryan ND, Dahl RE. Neural Systems of Threat Processing in
Adolescents: Role of Pubertal Maturation and Relation to Measures of Negative Affect.
Developmental Neuropsychology. 2011;36(4):429-452.
doi:10.1080/87565641.2010.550178

3.   Pozzi E, Vijayakumar N, Rakesh D, Whittle S. Neural correlates of emotion regulation in
adolescents and emerging adults: A meta-analytic study. *Biological Psychiatry*. 2020;0(0).
doi:10.1016/j.biopsych.2020.08.006

4.   Silvers JA, Shu J, Hubbard AD, Weber J, Ochsner KN. Concurrent and Lasting Effects of
Emotion Regulation on Amygdala Response in Adolescence and Young Adulthood.
Developmental Science. 2015;18(5):771-784. doi:10.1111/desc.12260

5.   Sullivan RM, Perry RE. Mechanisms and Functional Implications of Social Buffering in
Infants: Lessons from Animal Models. *Social Neuroscience*. 2015;10(5):500-511.
doi:10.1080/17470919.2015.1087425

6.   Pattwell SS, Liston C, Jing D, et al. Dynamic Changes in Neural Circuitry during Adolescence
Are Associated with Persistent Attenuation of Fear Memories. *Nature Communications*.
2016;7:11475. doi:10.1038/ncomms11475

7.   Moriceau S, Sullivan RM. Maternal presence serves as a switch between learning fear and
attraction in infancy. *Nature Neuroscience; New York*. 2006;9(8):1004-1006.
doi:http://dx.doi.org.ezproxy.cul.columbia.edu/10.1038/nn1733

8.   Meyer HC, Lee FS. Translating Developmental Neuroscience to Understand Risk for
Psychiatric Disorders. *AJP*. 2019;176(3):179-185. doi:10.1176/appi.ajp.2019.19010091

9.   Hariri AR, Tessitore A, Mattay VS, Fera F, Weinberger DR. The Amygdala Response to
Emotional Stimuli: A Comparison of Faces and Scenes. *NeuroImage*. 2002;17(1):317-323.
doi:10.1006/nimg.2002.1179

10.   Gee DG, Humphreys KL, Flannery J, et al. A Developmental Shift from Positive to Negative
Connectivity in Human Amygdala–Prefrontal Circuitry. *J Neurosci*. 2013;33(10):4584-4593.
doi:10.1523/JNEUROSCI.3446-12.2013

11.   Guyer AE, Monk CS, McClure-Tone EB, et al. A Developmental Examination of Amygdala
Response to Facial Expressions. *J Cogn Neurosci*. 2008;20(9):1565-1582.
doi:10.1162/jocn.2008.20114

12.   Killgore WDS, Oki M, Yurgelun-Todd DA. Sex-specific developmental changes in amygdala
849   responses to affective faces. *NeuroReport*. 2001;12(2):427.

850   13.   Passarotti AM, Sweeney JA, Pavuluri MN. Neural Correlates of Incidental and Directed
851   Facial Emotion Processing in Adolescents and Adults. *Soc Cogn Affect Neurosci*.
852   2009;4(4):387-398. doi:10.1093/scan/nsp029

853   14.   Swartz JR, Carrasco M, Wiggins JL, Thomason ME, Monk CS. Age-Related Changes in the
854   Structure and Function of Prefrontal Cortex–Amygdala Circuitry in Children and
855   Adolescents: A Multi-Modal Imaging Approach. *NeuroImage*. 2014;86:212-220.
856   doi:10.1016/j.neuroimage.2013.08.018

857   15.   Telzer EH, Flannery J, Humphreys KL, et al. "The Cooties Effect": Amygdala Reactivity to
858   Opposite-versus Same-Sex Faces Declines from Childhood to Adolescence. *J Cogn
859   Neurosci*. 2015;27(9):1685-1696. doi:10.1162/jocn_a_00813

860   16.   Decety J, Michalska KJ, Kinzler KD. The Contribution of Emotion and Cognition to Moral
861   Sensitivity: A Neurodevelopmental Study. *Cereb Cortex*. 2012;22(1):209-220.
862   doi:10.1093/cercor/bhr111

863   17.   Silvers JA, Insel C, Powers A, et al. The Transition from Childhood to Adolescence Is
864   Marked by a General Decrease in Amygdala Reactivity and an Affect-Specific Ventral-to-
865   Dorsal Shift in Medial Prefrontal Recruitment. *Developmental Cognitive Neuroscience*.
866   2017;25:128-137. doi:10.1016/j.dcn.2016.06.005

867   18.   Vink M, Derks JM, Hoogendam JM, Hillegers M, Kahn RS. Functional Differences in
868   Emotion Processing during Adolescence and Early Adulthood. *NeuroImage*. 2014;91:70-76.
869   doi:10.1016/j.neuroimage.2014.01.035

870   19.   Joseph JE, Zhu X, Gundran A, et al. Typical and Atypical Neurodevelopment for Face
871   Specialization: An fMRI Study. *J Autism Dev Disord*. 2015;45(6):1725-1741.
872   doi:10.1007/s10803-014-2330-4

873   20.   Todd RM, Evans JW, Morris D, Lewis MD, Taylor MJ. The Changing Face of Emotion: Age-
874   Related Patterns of Amygdala Activation to Salient Faces. *Soc Cogn Affect Neurosci*.
875   2011;6(1):12-23. doi:10.1093/scan/nsq007

876   21.   Hare TA, Tottenham N, Galvan A, Voss HU, Glover GH, Casey BJ. Biological Substrates of
877   Emotional Reactivity and Regulation in Adolescence During an Emotional Go-Nogo Task.
878   *Biological Psychiatry*. 2008;63(10):927-934. doi:10.1016/j.biopsych.2008.03.015

879   22.   Vijayakumar N, Pfeifer JH, Flournoy JC, Hernandez LM, Dapretto M. Affective reactivity
880   during adolescence: Associations with age, puberty and testosterone. *Cortex*.
881   2019;117:336-350. doi:10.1016/j.cortex.2019.04.024

23.   Kujawa A, Wu M, Klumpp H, et al. Altered Development of Amygdala-Anterior Cingulate
Cortex Connectivity in Anxious Youth and Young Adults. *Biological Psychiatry: Cognitive*
Neuroscience and Neuroimaging. 2016;1(4):345-352. doi:10.1016/j.bpsc.2016.01.006

24.   Pfeifer JH, Masten CL, Moore WE, et al. Entering Adolescence: Resistance to Peer
Influence, Risky Behavior, and Neural Changes in Emotion Reactivity. *Neuron*. 2011;69(5).
doi:10.1016/j.neuron.2011.02.019

25.   Pine DS, Grun J, Zarahn E, et al. Cortical Brain Regions Engaged by Masked Emotional Faces
in Adolescents and Adults: An fMRI Study. *Emotion*. 2001;1(2):137-147. doi:10.1037/1528-
3542.1.2.137

26.   Wu M, Kujawa A, Lu LH, et al. Age-related changes in amygdala–frontal connectivity during
emotional face processing from childhood into young adulthood. *Human Brain Mapping*.
2016;37(5):1684-1695. doi:10.1002/hbm.23129

27.   Yurgelun-Todd DA, Killgore WDS. Fear-related activity in the prefrontal cortex increases
with age during adolescence: A preliminary fMRI study. *Neuroscience Letters*.
2006;406(3):194-199. doi:10.1016/j.neulet.2006.07.046

28.   Zhang Y, Padmanabhan A, Gross JJ, Menon V. Development of human emotion circuits
investigated using a Big-Data analytic approach: Stability, reliability, and robustness. *J*
Neurosci. Published online July 22, 2019:0220-19. doi:10.1523/JNEUROSCI.0220-19.2019

29.   Silvers JA, Insel C, Powers A, et al. vlPFC–vmPFC–Amygdala Interactions Underlie Age-
Related Differences in Cognitive Regulation of Emotion. *Cereb Cortex*. 2017;27(7):3502-
3514. doi:10.1093/cercor/bhw073

30.   Perlman SB, Pelphrey KA. Developing Connections for Affective Regulation: Age-Related
Changes in Emotional Brain Connectivity. *Journal of Experimental Child Psychology*.
2011;108(3):607-620. doi:10.1016/j.jecp.2010.08.006

31.   Marek S, Tervo-Clemmens B, Calabro FJ, et al. Towards Reproducible Brain-Wide
Association Studies. *bioRxiv*. Published online August 22, 2020:2020.08.21.257758.
doi:10.1101/2020.08.21.257758

32.   Lieberman MD, Eisenberger NI, Crockett MJ, Tom SM, Pfeifer JH, Way BM. Putting feelings
into words: Affect labeling disrupts amygdala activity in response to affective stimuli.
Psychological Science. 2007;18(5):421-428. doi:10.1111/j.1467-9280.2007.01916.x

33.   Sergerie K, Chochol C, Armony JL. The role of the amygdala in emotional processing: A
quantitative meta-analysis of functional neuroimaging studies. *Neuroscience &*
Biobehavioral Reviews. 2008;32(4):811-830. doi:10.1016/j.neubiorev.2007.12.002

34.   Botvinik-Nezer R, Holzmeister F, Camerer CF, et al. Variability in the analysis of a single
916   neuroimaging dataset by many teams. *Nature*. 2020;582(7810):84-88.
917   doi:10.1038/s41586-020-2314-9

918   35.   Bowring A, Maumet C, Nichols TE. Exploring the impact of analysis software on task fMRI
919   results. *Human Brain Mapping*. 2019;40(11):3362-3384. doi:10.1002/hbm.24603

920   36.   Jo HJ, Lee J-M, Kim J-H, et al. Spatial accuracy of fMRI activation influenced by volume- and
921   surface-based spatial smoothing techniques. *NeuroImage*. 2007;34(2):550-564.
922   doi:10.1016/j.neuroimage.2006.09.047

923   37.   Achterberg M, van der Meulen M. Genetic and environmental influences on MRI scan
924   quantity and quality. *Developmental Cognitive Neuroscience*. 2019;38:100667.
925   doi:10.1016/j.dcn.2019.100667

926   38.   Bryce NV, Flournoy JC, Guassi Moreira JF, et al. Brain parcellation selection: An overlooked
927   decision point with meaningful effects on individual differences in resting-state functional
928   connectivity. *NeuroImage*. 2021;243:118487. doi:10.1016/j.neuroimage.2021.118487

929   39.   Di X, Zhang Z, Biswal BB. Understanding psychophysiological interaction and its relations to
930   beta series correlation. *bioRxiv*. Published online January 31, 2020:322073.
931   doi:10.1101/322073

932   40.   Cisler JM, Bush K, Steele JS. A Comparison of Statistical Methods for Detecting Context-
933   Modulated Functional Connectivity in fMRI. *NeuroImage*. 2014;84:1042-1052.
934   doi:10.1016/j.neuroimage.2013.09.018

935   41.   Crone EA, Elzinga BM. Changing brains: how longitudinal functional magnetic resonance
936   imaging studies can inform us about cognitive and social-affective growth trajectories.
937   *WIREs Cognitive Science*. 2015;6(1):53-63. doi:10.1002/wcs.1327

938   42.   Madhyastha T, Peverill M, Koh N, et al. Current Methods and Limitations for Longitudinal
939   fMRI Analysis across Development. *Developmental Cognitive Neuroscience*. Published
940   online November 22, 2017. doi:10.1016/j.dcn.2017.11.006

941   43.   Carpenter KLH, Angold A, Chen N-K, et al. Preschool Anxiety Disorders Predict Different
942   Patterns of Amygdala-Prefrontal Connectivity at School-Age. *PLOS ONE*.
943   2015;10(1):e0116854. doi:10.1371/journal.pone.0116854

944   44.   King KM, Littlefield AK, McCabe CJ, Mills KL, Flournoy J, Chassin L. Longitudinal modeling in
945   developmental neuroimaging research: Common challenges, and solutions from
946   developmental psychology. *Developmental Cognitive Neuroscience*. 2018;33:54-72.
947   doi:10.1016/j.dcn.2017.11.009

45. Madhyastha T, Peverill M, Koh N, et al. Current methods and limitations for longitudinal
fMRI analysis across development. *Developmental Cognitive Neuroscience*. 2018;33:118-
128. doi:10.1016/j.dcn.2017.11.006

46. Glenn ND. Distinguishing Age, Period, and Cohort Effects. In: Mortimer JT, Shanahan MJ,
eds. *Handbook of the Life Course*. Handbooks of Sociology and Social Research. Springer
US; 2003:465-476. doi:10.1007/978-0-306-48247-2_21

47. Elliott ML, Knodt AR, Ireland D, et al. What Is the Test-Retest Reliability of Common Task-
Functional MRI Measures? New Empirical Evidence and a Meta-Analysis: *Psychological*
Science. Published online June 3, 2020. doi:10.1177/0956797620916786

48. Sauder CL, Hajcak G, Angstadt M, Phan KL. Test-retest reliability of amygdala response to
emotional faces. *Psychophysiology*. 2013;50(11):1147-1156. doi:10.1111/psyp.12129

49. Zuo X-N, Xu T, Milham MP. Harnessing reliability for neuroscience research. *Nat Hum*
Behav. 2019;3(8):768-771. doi:10.1038/s41562-019-0655-x

50. Killgore WDS, Yurgelun-Todd DA. Unconscious Processing of Facial Affect in Children and
Adolescents. *Social Neuroscience*. 2007;2(1):28-47. doi:10.1080/17470910701214186

51. Gitelman DR, Penny WD, Ashburner J, Friston KJ. Modeling regional and psychophysiologic
interactions in fMRI: the importance of hemodynamic deconvolution. *NeuroImage*.
2003;19(1):200-207. doi:10.1016/S1053-8119(03)00058-2

52. Di X, Biswal BB. Psychophysiological Interactions in a Visual Checkerboard Task:
Reproducibility, Reliability, and the Effects of Deconvolution. *Front Neurosci*. 2017;11.
doi:10.3389/fnins.2017.00573

53. Di X, Reynolds RC, Biswal BB. Imperfect (de)Convolution May Introduce Spurious
Psychophysiological Interactions and How to Avoid It. *Human Brain Mapping*.
2017;38(4):1723-1740. doi:10.1002/hbm.23413

54. O'Reilly JX, Woolrich MW, Behrens TEJ, Smith SM, Johansen-Berg H. Tools of the Trade:
Psychophysiological Interactions and Functional Connectivity. *Soc Cogn Affect Neurosci*.
2012;7(5):604-609. doi:10.1093/scan/nss055

55. Jalbrzikowski M, Larsen B, Hallquist MN, Foran W, Calabro F, Luna B. Development of
White Matter Microstructure and Intrinsic Functional Connectivity Between the Amygdala
and Ventromedial Prefrontal Cortex: Associations With Anxiety and Depression. *Biological*
Psychiatry. 2017;82(7):511-521. doi:10.1016/j.biopsych.2017.01.008

56. Qin S, Young CB, Duan X, Chen T, Supekar K, Menon V. Amygdala Subregional Structure
and Intrinsic Functional Connectivity Predicts Individual Differences in Anxiety During Early
Childhood. *Biological Psychiatry*. 2014;75(11):892-900.
doi:10.1016/j.biopsych.2013.10.006

57. Grady CL, Rieck JR, Nichol D, Rodrigue KM, Kennedy KM. Influence of sample size and
analytic approach on stability and interpretation of brain-behavior correlations in task-
related fMRI data. *Human Brain Mapping*. Published online 2020.
doi:https://doi.org/10.1002/hbm.25217

58. Flournoy JC, Vijayakumar N, Cheng TW, Cosme D, Flannery JE, Pfeifer JH. Improving
practices and inferences in developmental cognitive neuroscience. *Developmental*
Cognitive Neuroscience. 2020;45:100807. doi:10.1016/j.dcn.2020.100807

59. Bridgeford EW, Wang S, Yang Z, et al. Big Data Reproducibility: Applications in Brain
Imaging. *bioRxiv*. Published online January 18, 2020:802629. doi:10.1101/802629

60. Steegen S, Tuerlinckx F, Gelman A, Vanpaemel W. Increasing Transparency Through a
Multiverse Analysis. *Perspect Psychol Sci*. 2016;11(5):702-712.
doi:10.1177/1745691616658637

61. Gelman A, Loken E. The statistical crisis in science: data-dependent analysis--a 'garden of
forking paths'--explains why many statistically significant comparisons don't hold up.
American Scientist. 2014;102(6):460-466.

62. Del Giudice M, Gangestad SW. A Traveler's Guide to the Multiverse: Promises, Pitfalls, and
a Framework for the Evaluation of Analytic Decisions. *Advances in Methods and Practices*
in Psychological Science. 2021;4(1):2515245920954925. doi:10.1177/2515245920954925

63. Bossier H, Roels SP, Seurinck R, et al. The empirical replicability of task-based fMRI as a
function of sample size. *NeuroImage*. 2020;212:116601.
doi:10.1016/j.neuroimage.2020.116601

64. Nee DE. fMRI replicability depends upon sufficient individual-level data. *Communications*
Biology. 2019;2(1):1-4. doi:10.1038/s42003-019-0378-6

65. Button KS, Ioannidis JPA, Mokrysz C, et al. Power failure: why small sample size
undermines the reliability of neuroscience. *Nature Reviews Neuroscience*. 2013;14(5):365-
376. doi:10.1038/nrn3475

66. Gelman A, Carlin J. Beyond Power Calculations: Assessing Type S (Sign) and Type M
(Magnitude) Errors. *Perspect Psychol Sci*. 2014;9(6):641-651.
doi:10.1177/1745691614551642

67. Kriegeskorte N, Simmons WK, Bellgowan PSF, Baker CI. Circular analysis in systems
neuroscience: the dangers of double dipping. *Nat Neurosci*. 2009;12(5):535-540.
doi:10.1038/nn.2303

68. Richeson JA, Todd AR, Trawalter S, Baird AA. Eye-Gaze Direction Modulates Race-Related
Amygdala Activity. *Group Processes & Intergroup Relations*. 2008;11(2):233-246.
doi:10.1177/1368430207088040

69.   Telzer EH, Humphreys KL, Shapiro M, Tottenham N. Amygdala Sensitivity to Race Is Not
Present in Childhood but Emerges over Adolescence. *Journal of Cognitive Neuroscience*.
2012;25(2):234-244. doi:10.1162/jocn_a_00311

70.   Ciric R, Wolf DH, Power JD, et al. Benchmarking of participant-level confound regression
strategies for the control of motion artifact in studies of functional connectivity.
Neuroimage. 2017;154:174-187. doi:10.1016/j.neuroimage.2017.03.020

71.   Gilmore A, Buser N, Hanson JL. Variations in Structural MRI Quality Significantly Impact
Commonly-Used Measures of Brain Anatomy. *bioRxiv*. Published online May 7,
2020:581876. doi:10.1101/581876

72.   Rorden C, Bonilha L, Fridriksson J, Bender B, Karnath H-O. Age-specific CT and MRI
templates for spatial normalization. *NeuroImage*. 2012;61(4):957-965.
doi:10.1016/j.neuroimage.2012.03.020

73.   Boubela RN, Kalcher K, Huf W, et al. fMRI measurements of amygdala activation are
confounded by stimulus correlated signal fluctuation in nearby veins draining distant brain
regions. *Scientific Reports*. 2015;5(1):10499. doi:10.1038/srep10499

74.   Fair DA, Miranda-Dominguez O, Snyder AZ, et al. Correction of respiratory artifacts in MRI
head motion estimates. *NeuroImage*. 2020;208:116400.
doi:10.1016/j.neuroimage.2019.116400

75.   Gratton C, Dworetsky A, Coalson RS, et al. Removal of high frequency contamination from
motion estimates in single-band fMRI saves data without biasing functional connectivity.
bioRxiv. Published online April 21, 2020:837161. doi:10.1101/837161

76.   Achenbach TM. *Integrative Guide for the 1991 CBCL/4-18, Ysr, and Trf Profiles*. 1st US-1st
Printing edition. Univ Vermont/Dept Psychiatry; 1991.

77.   US Census Bureau. U.S. Census Bureau QuickFacts: Los Angeles County, California.
Published 2021. Accessed March 24, 2021.
https://www.census.gov/quickfacts/losangelescountycalifornia

78.   Heeringa SG, Wagner J, Torres M, Duan N, Adams T, Berglund P. Sample designs and
sampling methods for the Collaborative Psychiatric Epidemiology Studies (CPES).
International Journal of Methods in Psychiatric Research. 2004;13(4):221-240.
doi:https://doi.org/10.1002/mpr.179

79.   Birmaher B, Brent DA, Chiappetta L, Bridge J, Monga S, Baugher M. Psychometric
Properties of the Screen for Child Anxiety Related Emotional Disorders (SCARED): A
Replication Study. *Journal of the American Academy of Child & Adolescent Psychiatry*.
1999;38(10):1230-1236. doi:10.1097/00004583-199910000-00011

80.   Chorpita BF, Yim L, Moffitt C, Umemoto LA, Francis SE. Assessment of Symptoms of DSM-
1053   IV Anxiety and Depression in Children: A Revised Child Anxiety and Depression Scale.
1054   *Behaviour Research and Therapy*. 2000;38(8):835-855. doi:10.1016/S0005-7967(99)00130-
1055   8

1056   81.   Beretta L, Santaniello A. Nearest Neighbor Imputation Algorithms: A Critical Evaluation.
1057   *BMC Med Inform Decis Mak*. 2016;16(Suppl 3). doi:10.1186/s12911-016-0318-z

1058   82.   Wager TD, Nichols TE. Optimization of experimental design in fMRI: a general framework
1059   using a genetic algorithm. *Neuroimage*. 2003;18(2):293-309.

1060   83.   Calvo MG, Lundqvist D. Facial expressions of emotion (KDEF): Identification under
1061   different display-duration conditions. *Behav Res*. 2008;40(1):109-115.
1062   doi:10.3758/BRM.40.1.109

1063   84.   Cox RW. AFNI: Software for Analysis and Visualization of Functional Magnetic Resonance
1064   Neuroimages. *Comput Biomed Res*. 1996;29(3):162-173.

1065   85.   Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM. FSL. *NeuroImage*.
1066   2012;62(2):782-790. doi:10.1016/j.neuroimage.2011.09.015

1067   86.   Jenkinson M, Bannister P, Brady M, Smith S. Improved Optimization for the Robust and
1068   Accurate Linear Registration and Motion Correction of Brain Images. *NeuroImage*.
1069   2002;17(2):825-841. doi:10.1006/nimg.2002.1132

1070   87.   Pernet CR. Misconceptions in the use of the General Linear Model applied to functional
1071   MRI: a tutorial for junior neuro-imagers. *Front Neurosci*. 2014;8:1.
1072   doi:10.3389/fnins.2014.00001

1073   88.   Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE. Spurious but systematic
1074   correlations in functional connectivity MRI networks arise from subject motion.
1075   *NeuroImage*. 2012;59(3):2142-2154. doi:10.1016/j.neuroimage.2011.10.018

1076   89.   Woolrich MW, Ripley BD, Brady M, Smith SM. Temporal autocorrelation in univariate
1077   linear modeling of FMRI data. *Neuroimage*. 2001;14(6):1370-1386.
1078   doi:10.1006/nimg.2001.0931

1079   90.   Fischl B. FreeSurfer. *NeuroImage*. 2012;62(2):774-781.
1080   doi:10.1016/j.neuroimage.2012.01.021

1081   91.   VanTieghem M, Korom M, Flannery J, et al. Longitudinal changes in amygdala,
1082   hippocampus and cortisol development following early caregiving adversity.
1083   *Developmental Cognitive Neuroscience*. 2021;48:100916. doi:10.1016/j.dcn.2021.100916

1084   92.   Cosme D, Lopez R. Neural indicators of food cue reactivity, regulation, and valuation and
1085   their associations with body composition and daily eating behavior. Published online
1086   March 7, 2020. doi:10.31234/osf.io/23mu5

1087   93.   Klapwijk E, van den Bos W, Tamnes CK, Mills KL, Raschle N. *Opportunities for Increased*
1088   *Reproducibility and Replicability of Developmental Cognitive Neuroscience*. PsyArXiv; 2019.
1089   doi:10.31234/osf.io/fxjzt

1090   94.   Orben A, Przybylski AK. The association between adolescent well-being and digital
1091   technology use. *Nat Hum Behav*. 2019;3(2):173-182. doi:10.1038/s41562-018-0506-1

1092   95.   Simonsohn U, Simmons J, Nelson L. Specification Curve: Descriptive and Inferential
1093   Statistics on All Reasonable Specifications. *Marketing Papers*. Published online January 1,
1094   2015. doi:10.2139/ssrn.2694998

1095   96.   Simonsohn U, Simmons JP, Nelson LD. Specification curve analysis. *Nat Hum Behav*.
1096   2020;4(11):1208-1214. doi:10.1038/s41562-020-0912-z

1097   97.   Craddock C, Sikka S, Cheung B, et al. Towards Automated Analysis of Connectomes: The
1098   Configurable Pipeline for the Analysis of Connectomes (C-PAC). In: ; 2013.
1099   doi:10.3389/conf.fninf.2013.09.00042

1100   98.   Satterthwaite TD, Wolf DH, Loughead J, et al. Impact of in-scanner head motion on
1101   multiple measures of functional connectivity: Relevance for studies of neurodevelopment
1102   in youth. *NeuroImage*. 2012;60(1):623-632. doi:10.1016/j.neuroimage.2011.12.063

1103   99.   Satterthwaite TD, Wolf DH, Ruparel K, et al. Heterogeneous Impact of Motion on
1104   Fundamental Patterns of Developmental Changes in Functional Connectivity during Youth.
1105   *Neuroimage*. 2013;83:45-57. doi:10.1016/j.neuroimage.2013.06.045

1106   100. Abdulrahman H, Henson RN. Effect of trial-to-trial variability on optimal event-related
1107   fMRI design: Implications for Beta-series correlation and multi-voxel pattern analysis.
1108   *NeuroImage*. 2016;125:756-766. doi:10.1016/j.neuroimage.2015.11.009

1109   101. McLaren DG, Ries ML, Xu G, Johnson SC. A generalized form of context-dependent
1110   psychophysiological interactions (gPPI): A comparison to standard approaches.
1111   *NeuroImage*. 2012;61(4):1277-1286. doi:10.1016/j.neuroimage.2012.03.068

1112   102. Mackey S, Petrides M. Architecture and Morphology of the Human Ventromedial
1113   Prefrontal Cortex. *Eur J Neurosci*. 2014;40(5):2777-2796. doi:10.1111/ejn.12654

1114   103. Fox MD, Zhang D, Snyder AZ, Raichle ME. The Global Signal and Observed Anticorrelated
1115   Resting State Brain Networks. *Journal of Neurophysiology*. 2009;101(6):3270-3283.
1116   doi:10.1152/jn.90777.2008

104. R Core Team. R: The R Project for Statistical Computing. Published 2017. Accessed October
30, 2019. https://www.r-project.org/

105. Gelman A, Lee D, Guo J. Stan: A Probabilistic Programming Language for Bayesian
Inference and Optimization. *Journal of Educational and Behavioral Statistics*.
2015;40(5):530-543. doi:10.3102/1076998615606113

106. Bürkner P-C. *Brms: Bayesian Regression Models Using "Stan."*; 2019. Accessed December
6, 2019. https://CRAN.R-project.org/package=brms

107. Beeley C. *Web Application Development with R Using Shiny*. Packt Publishing Ltd; 2013.

1125