1

3 **Time-resolved, integrated analysis of clonally evolving genomes**

6 Carine Legrand[1,2,*], Ranja Andriantsoa[1], Peter Lichter[3,4], Frank Lyko[1]

1 Division of Epigenetics, DKFZ-ZMBH Alliance, German Cancer Research Center, 69120 Heidelberg, Germany

2 Université Paris Cité, Génomes, biologie cellulaire et thérapeutique U944, INSERM, CNRS, F-75010 Paris,

France

3 Division of Molecular Genetics, German Cancer Research Consortium (DKTK), German Cancer Research

Center (DKFZ), Im Neuenheimer Feld 280, 69120 Heidelberg, Germany

4 Molecular Precision Oncology, National Center for Tumor Diseases, Im Neuenheimer Feld 460, 69120

Heidelberg, Germany

* Corresponding author

E-mail: carine.legrand@inserm.fr

Abstract

Clonal genome evolution is a key aspect for parthenogenetic species and cancer. While many studies describe precise landscapes of clonal evolution in cancer, few determine the underlying evolutionary parameters from molecular data, and even fewer integrate theory with data. We derived theoretical results linking mutation rate, time, expansion dynamics, and clinical parameters. We then inferred time-resolved estimates of evolutionary parameters from mutation accumulation, mutational signatures and selection. Using this framework, we traced the speciation of the clonally evolving marbled crayfish population to a time window between 1986 and 1990, which is consistent with biological records. We also used our framework to analyze a published dataset of glioblastoma samples, which identified tumor expansion patterns, cell survival at resection, and selective forces as important factors for tumor development. In conclusion, our framework allowed a time-resolved, integrated analysis of key parameters in clonally evolving genomes, and provided novel insights into the evolutionary age of marbled crayfish and the progression of glioblastoma.

Introduction

The evolution of genomes is shaped by many factors, among which the random accumulation of mutations over time plays a fundamental role [1,2]. Far from being homogeneous, the probability of a mutation depends on many factors such as the genomic location [3], mutator alleles, local nucleotide context or mutagenic exposures [4]. Other genomic modifications include recombination in sexual reproduction, copy number variants and genomic rearrangements, gene transfers and hybridization. The capacity of any genomic modification to be inherited is partly stochastic, for instance through genetic drift [5], but can be favored or disfavored by positive or negative selection. Genome evolution was historically observed through the analysis of phenotypes [6], and can now be determined more precisely using high-throughput sequencing in parallel with experimental or cohort settings, such as mutation accumulation experiments, or the analysis of genetic trios [7,8].

Clonal genome evolution is shaped by a more limited set of mechanisms. Mutation rate, growth and variant frequencies are key parameters, which determine the speed of evolution, and which function under the influence of selection pressure. Due to its particular mode of asexual reproduction, the marbled crayfish (*Procambarus virginalis*) represents an ideal animal model to study clonal genome evolution [9]. The animals are currently colonizing diverse habitats in a process that is associated with emerging genetic differentiation [10]. Interestingly, marbled crayfish appears to be an evolutionary young species, as their first emergence can be traced back to a specific event in 1995 [11]. If confirmed, this exceptionally young evolutionary age would represent a highly distinctive feature of the model system.

Clonal genome evolution also plays an important role in cancer formation. Cancer is a

disease of clonal evolution, characterized by the accumulation of somatic mutations

into a pathogenic tumoral genome. Several authors have described the critical role of

mutational patterns and selection in cancer [1,3,12], while neutral evolution is still

debated [13,14]. In glioblastoma, the analysis of tumor trajectories revealed a tumor

initiation years before diagnosis [15]. Consequently, it would be of great interest to

infer evolutionary parameters over the course of tumor progression.

In this study, we aimed to develop an integrated analysis of clonal genome evolution.

To this end, we reformulated the dependence of mutation accumulation on variant

allele frequency, and used this formulation to determine the links between the

mutation rate, growth and survival rates. We further integrated these parameters with

selection estimates, obtained from the non-synonymous to synonymous ratio. Finally,

we integrated time estimates in our model, based on clock-like mutational signatures.

We applied our approach to the clonally evolving marbled crayfish. We provided a

detailed view of mutation accumulation, selection, and time, and estimated the time of

speciation. We further applied our framework to clonal genome evolution in cancer,

using recently published samples of primary and recurrent glioblastoma [15].

Results

The genetic near-monoclonality of the marbled crayfish population [9,10] establishes

this species as an excellent model system for studying clonal genome evolution. In

order to assess the mutation rate of the *P. virginalis* genome, we used whole-genome

sequencing of a line of direct descendants from our laboratory colony of *P. virginalis*,

that were sampled over a period of seven years (Fig 1A). The mutation rate was

calculated as the average number of de novo mutations in animals 34 and 35 as

compared to animal 1, per nucleotide and per year. From these samples, we obtained a

mutation rate equal to $\mu=3.51 \cdot 10^{-8}/nt/y$ (95% confidence interval (CI):

$\left[1.67 \cdot 10^{-8}; 5.35 \cdot 10^{-8}\right]/nt/y$). This mutation rate of *P. virginalis* is comparable to

known mutation rates from other arthropods (Fig 1B).

Fig 1. Mutation rate of *P. virginalis* and coalescent. (A) Genealogy of laboratory

animals, with sequenced animals marked in grey. (B) Mutation rate in *P. virginalis*,

in other arthropods and in *Homo sapiens* [16–27]. HSC: Hematopoietic Stem Cells,

CCRCC: Clear Cell Renal Cell Carcinoma, NSCLC: Non-Small Cell Lung

Carcinoma. Error bars correspond to reported 95% confidence intervals. (C)

Coalescent tree based on a constant mutation rate and sequences of sampled animals.

The posterior probability of each branch is indicated in red.

Based on this mutation rate, we made an evaluation of the evolutionary age of *P.*

virginalis, using a Markov Chain Monte Carlo with Bayesian evolutionary analysis

[28] on whole-genome sequencing datasets from 13 animals (Fig 1C). We generated

10 million states, which allowed convergence of the sampled states, and led to

adequately large effective sample sizes (see Methods for details). The resulting

coalescent tree showed that animals 1, 34 and 35 correctly clustered together, as well

as animals from German wild populations (Hannover, Reilingen, Moosweiher) and

from the likely foundational laboratory lineage of the German wild populations

(Heidelberg). Furthermore, samples from Madagascar formed a separate branch.

Interestingly, Petshop 2 [9] was nested in the branch of animals from Madagascar.

This is consistent with the notion that the Malagasy population was founded by an

animal that was originally obtained from a German pet shop. Posterior probabilities

(Fig 1C, red annotations) indicate highly probable branching for all but the top

coalescent event, which has 0.5206 probability. From this tree, the most recent

common ancestor of the 13 animals occured in 1988.0 (95% CI: [1986.1 ; 1989.8]).

This is anterior, and therefore broadly consistent, with the first documented

appearance of *P. virginalis* in 1995 [11].

We next modeled mutation accumulation under a fast growth scenario. This included

the patterns of mutation accumulation as a function of time, the dependence of allele

frequency on time, and the number of mutations as a function of allelic frequency. We

found that mutation accumulation gave information on the mutation rate, but not on

selection (S1 File, Eq. 12). This provided the rationale for examining the dynamics of

the mutation rate in *P. virginalis* using the *M(1/f)* curve, where *M* is the number of

mutations and *f* is the allelic frequency (Fig 2A). The resulting curve suggested that

the mutation rate changed over time, with 4 phases delineated by a segmented

regression (Fig 2A; *p*=0.06). The mutation rate was reduced in phase 3, as compared

to phases 1 and 2, and increased in phase 4 (Fig 2A). Under our model, selection *s* is

not observable using *M(1/f)* (S1 File, Eq. 12). We therefore used the ratio of non-

synonymous to synonymous mutations to estimate *s* (Fig 2B). The resulting values

were close to unity, for phases 1 and 2, suggesting the absence of selection. During

phases 3 and 4, we detected *s* values >1, and <1, suggesting phases of positive and

negative selection respectively, in the more recent past of *P. virginalis*.

Fig 2. Mutation accumulation, selection and time course of *P. virginalis* genome evolution. (A) Mutation accumulation as a function of the inverse allele frequency *1/f* (black) and phases from automated segmentation (breakpoints in grey, segments in blue). (B) Non-synonymous to synonymous ratio (*dNdS*). The smoothed ratio is shown in red. (C) Comparison of clock-like and non-clock-like mutational signatures. (D) Mutation accumulation as a function of time. Smoothened mutation accumulation is shown in red.

In order to obtain time-resolved estimates, we then used previously established clock-like mutational single-base signatures (SBS1 and SBS5) [29–31] as a proxy for the time course of mutation accumulation (Fig 2C). We further assumed that the arrow of time from past to present corresponds to the arrow of increasing *1/f*. To obtain a time course in arbitrary units, we calculated the integral of the clock-like components of mutation accumulation (Fig 2D, Methods). According to the mathematical model, the

slope of this curve is proportional to the mutation rate as a function of time (S1 File,

Eq. 12). The results (Fig 2D) showed that this mutation rate exhibited less variation

than the mutation rate per division (Fig 2A). Because the temporal and per-division

mutation rates differ essentially by the growth rate, this might indicate fluctuations in

the growth rate of the marbled crayfish population. As a whole, our analyses

suggested distinct phases, detected significant variations of evolutionary parameters in

the recent past of *P. virginalis*, and allowed to trace its speciation to a time point that

is consistent with biological records.

In *P. virginalis*, we developed a framework to analyse the evolution of a clonal

genome, which is driven by germline mutations. This framework can in principle also

be applied to analyse the clonal evolution of a tumor genome, which is driven by

somatic mutations. Since glioblastoma is a high grade tumor with systematic

recurrence and poor patient survival, a better understanding of evolutionary

parameters is important. We therefore applied our framework to a published set of

whole-genome sequencing data of primary and recurrent glioblastoma tumors [15].

This study also estimated the age of primary tumors, allowing further data integration.

Based on the curve *M(1/f)*, we generated mutation rate profiles (Fig. 4A, see S1 Fig A

for individual samples), which we further segmented into phases (Fig. 4A, $p < 2.2 \times 10^{-16}$

). The results indicated distinct variations in the mutation rate in primary and

recurrent samples (Fig 3; S1 Table). In the exemplary sample 1 in Fig 3A, the

segmentation separates 5 phases significantly. After we excluded the outermost

phases 1 and 5, where changes in mutation frequencies are more likely to be

influenced by artifacts, the mutation rate per division decreased steadily in phases 2-4.

Fig 3. Mutation accumulation, selection and time dynamics of a representative

glioblastoma tumor (patient 1, primary tumor), expansion patterns and survival

ratio. (A) Mutation accumulation as a function of the inverse allele frequency *1/f*

(black) and phases from automated segmentation (breakpoints are indicated as dashed

vertical lines, segments are indicated in blue). The confidence band at 95% level is

shown in grey. (B) Non-synonymous to synonymous ratio *dNdS*. In the lower inest,

purple and blue bars show non-synonymous and synonymous mutations, respectively.

The smoothened ratio is shown in red. (C) Clock-like and non-clock-like mutational

signatures. (D) Mutation accumulation as a function of time. (E-H) Stratification of

expansion curves $\omega\gamma N$ into 4 subtypes: 1: convex, 2: horizontal then convex, 3:

complex, 4: increase (showing exemplary primary tumor samples GBM30

(GlioBlastoma Multiforme, sample number 30), GBM26, GBM17, GBM14,

respectively). (I) Dependence of time to recurrence on the γ_R/γ_P ratio. Fit1 corresponds

to a linear regression of time versus $\log_{10}(\gamma_R/\gamma_P)$, with intercept=19.511 (standard error

SE=2.544) and slope=-5.819 (SE=1.455), fit2 corresponds to a linear regression of

time versus $\log_{10}(\log_{10}(\gamma_R/\gamma_P))$, with intercept=18.922 (SE=1.806) and slope=-29.321

(SE=5.285).

We next looked at selection using the *dN/dS* ratio. Taking confidence bounds into

account, the results were compatible with neutral selection for most tumors (Fig. 4B,

S1 Fig B per sample). However, 11 primary tumor samples showed evidence of

negative selection during intervals, for instance sample 35 (S1 Fig B for sample 35).

We also observed evidence for positive selection in two primary tumor samples

(Samples 2 and 7, S1 Fig B). Interestingly, 7 out of 9 recurrent tumor samples

underwent prolonged phases of negative selection (for example, sample 4, 1/f in

[1/0.5; 1/0.1], S1 Fig B), while 2 samples exhibited short phases of negative selection.

No recurrent tumor sample showed any significant phase of positive selection.

We next determined the timeline of tumor evolution, by comparing the prevalence of

the stable, clock-like SBS1 signature to the other, non-clock like SBS signatures (Fig

3C, see S1 Fig for individual samples). This includes SBS5, which is non clock-like

in glioblastoma according to [29]. This was confirmed by low correlation coefficients

between SBS1 and SBS5 in the samples under study (ρ_P=-0.060 +/- 0.143 in primary

tumors; ρ_P=0.067 +/- 0.136 in recurrences). Using the information on the clock-like

signature SBS1, and using equation (9) (Methods), we reconstructed *M* as a function

of time (Fig 3D, S1 Fig D per sample), in arbitrary units. The slope of this curve is

proportional to the mutation rate per time unit. Similar to the mutation rate per

division in Fig 3A, the mutation rate per time unit decreased from phase 2 to phase 4,

though less markedly. Furthermore, similar to the situation for *P. virginalis*, the

difference observed in sample 1 might indicate fluctuating growth during phases 2-4.

More specifically, differences between per-division and temporal mutation rates

corresponded in our model to the growth rate, the survival rate, and the number of

cells (S1 File, Eq. 8).

We next asked if these terms could be used to characterize the set of 42 primary and

recurrent tumor pairs. Using temporal and per division mutation rates, we could

reconstruct these terms, which correspond to the product $\omega\gamma N$ (growth rate ω, tumor

cell survival rate γ and number of cells N), and which we termed expansion

parameters. We examined the corresponding curves and found that they allowed us to

classify the tumors into four subtypes: (1) convex, (2) stable then convex, (3) double

convexity and (4) increase (Fig 3E).We then looked at a possible association between

the patterns of the $\omega\gamma N$ curve in the primary tumors, and the time to the recurrence,

but the results were inconclusive (p=0.0367, p_{adj}=0.0734, n=20). However, the time

difference between the resection of the primary tumor and the resection of the

recurrence is known for a subset of samples, and the age of tumors was estimated

previously [15]. This allowed us to transform the time course from arbitrary units into

real units (Methods, Eq. 2, S2 Fig). Furthermore, we extended our modelling to be

able to express the transition from the primary to the recurrent tumor (Methods, Eq.

4). With this, we could determine the tumor survival ratio from time estimates. Using

the previously established values of 2 years and 7 years [15] as the lowest and highest

limits for the time course of the primary tumor, we could determine a range for the

value of the tumor survival ratio γ_R/γ_P for each individual sample (Methods, Eq. 8, S2

Table). As a result, the lowest value of the ratio γ_R/γ_P, corresponding to a tumor

emergence about 2 years before diagnosis, was always higher than 1 (median =27.8

[17.4; 54.0] for the lower bound, median=97,5 [60.9; 189.0] for the upper bound,

n=20 samples). These results indicated that tumor cell survival was higher at the start

of the recurrence than at the end of the primary tumor growth. Not surprisingly, γ_R/γ_P

ratios were associated with the time to recurrence (Fig 3F, $p_{adj}=1.258\times10^{-3}$ and

$p_{adj}=8.649\times10^{-4}$), with higher γ_R/γ_P ratios corresponding to shorter time to recurrence.

Collectively, these results uncover substantial variations of evolutionary parameters

among glioblastoma samples, and provide an improved understanding of growth and

survival in tumor subtypes.

Discussion

In this study, we presented an integrated framework to analyse the evolution of clonally evolving genomes. We first determined the mutation rate of *P. virginalis* to be 3.51×10^{-8}/nt/y, which is comparable to the mutation rates of other arthropods. We detected separated evolutionary phases, during which the mutation rate varied significantly, and which suggested selection events in the recent past of the species. The variations of the mutation rate, observed as well in the tumor samples, support the argument that the mutation rate should not be considered constant [1,32,33]. Known mechanisms can explain these variations, including a temporarily more pronounced effect of error-prone mechanisms, hypoxia-induced mutagenesis, or transcription-associated mutagenesis [1,34]. We further noted that the temporal mutation rate seemed more stable than the per-division mutation rate, and that the difference, according to our modelling approach, could be explained by variations of growth. Finally, we traced the speciation of *P. virginalis* to 1988 (95% confidence limits: [1986 ; 1990]), in agreement with first reports of this animal in 1995 [11], and consistent with a very young evolutionary age of the species.

In tumor samples, our approach allowed a single patient-level analysis of evolutionary parameters, and similarly revealed the presence of different phases, variations of the mutation rate, and a few significant events of selection. Utilizing the difference between temporal and per-division mutation rate, we could stratify the samples into 4 subtypes. Building on previously estimated tumor age, we could also derive the survival ratio for tumor cells in the recurrence, relatively to the primary tumor. We

found that tumor cells survive better at the start of the recurrence, though with

important variations. This supports the notion that tumor regrowth can be more

aggressive after surgical resection of primary glioblastoma tumors [35,36], possibly

because resection-induced astrocyte injury can support faster growth [36], or because

the tumor microenvironment can promote tumor regrowth [37–39]. Conversely, a

stronger immune response might prevent tumor regrowth.

In conclusion, this integrated analysis of mutation accumulation, dNdS ratio and

mutational signatures provided a detailed landscape of evolutionary parameters, and

delivered important insights for two paradigms of clonal genome evolution,

demonstrating its potential for further applications.

Materials and Methods

***Procambarus virginalis* samples.** Freshwater crayfish samples from [9] were used. Additionally, samples from Madagascar 1 sample and Moosweiher sample were resequenced (S3 Table). Animal 1 corresponds to the lab strain, acquired from a pet shop. New genomic DNA samples were taken from animal 34 and animal 35, which, as animal 1, also correspond to lab strains animals, and which are direct offsprings of animal 1. These new samples were prepared and submitted for whole genome sequencing following the protocol already described. The genealogy and birth date of animals were retrieved from laboratory records and field records (S3 Table). Sequence data was trimmed using Trimmomatic v0.32 (settings: LEADING:3 TRAILING:3 SLIDINGWINDOW:4:20 MINLEN:40, adapter sequence: TruSeq3-PE). Next, trimmed data was mapped to Pvir genome assembly v04 (https://www.ncbi.nlm.nih.gov, Bioproject Accession: PRJNA356499), using Bowtie2 (v2.2.6, setting: --sensitive). Aligned reads were sorted, cleared from duplicates, sorted and indexed using samtools. Subsequently, variant calling was performed using Freebayes v0.9.21-g7dd41db (parameters: --report-all-haplotype-alleles -P 0.7 -p 3 --min-mapping-quality 30 --min-base-quality 20 --min-coverage 6 --report-genotype-likelihood-max).

Glioblastoma Multiforme samples. The glioblastoma primary and recurrent tumor samples correspond to the WGS cohort already described in [15]. In particular, summary information can be found in supplementary table 1 of [15]. After approval of the research project, access to the SNP data of primary and recurrent tumor samples, as well as time to recurrence when available, was granted.

Mutation rate of *P. virginalis*. Mutation accumulation between animals 1 and

descendant animals 34 and 35 was used to calculate the mutation rate. SNP variants

were examined in terms of quality and coverage. Variants with quality≥35,

coverage≥50 and ≤200 were retained for the main estimate of the mutation rate.

Coverages 200 and higher exhibited altered SNP distribution and were thus excluded

because possibly corresponding to a distinct part of *P. virginalis* genome (possibly highly

repetitive and variable domains). Subsequently, the mutation rate per nucleotide per year

was calculated as the count of biallelic mutated nucleotides in animal 34

(respectively, animal 35) as compared to animal 1, divided by the count of nucleotides

in the triploid genome of *P. virginalis* ($N=10.5 \times 10^9$), and divided by the time (5.75

and 6.08 years), between the birth dates of animal 1 and 34 (respectively, birth date of

animal 35). We assumed that counts of new mutations follow a Poisson distribution,

and with this we determined the standard deviation on the count of mutations

observed (genotyping uncertainty). Second, we assumed that the standard deviation

for the date of animal birth equated to a third of the total uncertainty on time of birth.

Third, we calculated the standard deviation between mutation rates for animal 34 and

animal 35 (biological variability). Finally, we took the total standard deviation as the

quadratic sum of these three components (assuming that the different sources of

variability follow a normal distribution)

Coalescent time. Time to most recent common ancestor for *P. virginalis* samples was

determined using Bayesian evolutionary analysis by sampling trees (BEAST v1.10.4

[28]). Mutation data with quality >35 and coverage depth >15 was used in this

analysis (a coverage cutoff of 25 was not justified here because samples other than

animals 1, 34 and 35 possessed a notably lower average sequencing depth). Samples

birth dates were used as tip dates. Further BEAST parameters used were: simple

substitution model with estimated base frequencies, strict clock, skyride coalescent

prior. The length of chain for the Markov chain Monte Carlo was 10M. These

parameters were built into the BEAST input file, using the utility BEAUTi. The

outputs were analyzed using the utility TRACER. In particular, convergence was read

from the sampled states curves of the different parameters, and effective sample sizes

were adequately >100 (2984 or more), indicating sufficiently decorrelated sampled

states.

Study of mutation accumulation. An infinitesimal increment of mutations was

defined from the mutation rate, ploidy, cell survival, growth and number of cells (see

S1 File for a detailed description). We stratified this expression for each subclone. We

have then determined the relationship between the observable allele frequency of a

mutation, which is the one obtained after sequencing and SNP calling, and the

features of the subclone where this mutation appeared. Next, we have determined a

formula for the increment of the number of cells dN and for the increment of inverse

allele frequency $d(1/f)$. For this latter increment, we have made the assumption that

ploidy was constant. We could then deduce the mutation accumulation dM as a

function of inverse allele frequency $1/f$, in each subclone i. Finally, the equation for

mutation accumulation over all subclones dM was obtained by summing the

individual contributions dM of each subclone. Mutation accumulation $M(1/f)$, which

slope corresponds to $dM/d(1/f)$, was plotted from SNP data (filtered by quality phred

score QUAL ≥ 30, depth ≥ 10, depth of alternate allele ≥ 3), and from the corresponding

allele frequencies. For uncertainties, we used bins of +/-0.25 over 1/f. In each bin, the

mutation count was subjected to bootstrap resampling, which yielded a bootstrap

distribution. The confidence interval at 95% was taken as the interval bounded by the

2.5% and 97.5% quantiles of the bootstrap distribution.

Mutation annotation and dNdS ratio. Mutations were annotated as synonymous or

non-synonymous (including splice or stopgain mutations) using SNPdat v1.0.5. The

dNdS ratio was calculated as the quotient of non-synonymous mutations by

synonymous mutations in a sample, divided by the average quotient in the full

genome. The average quotient of non-synonymous to synonymous in humans was

taken equal to 3.34951759 (hg19). Uncertainties were determined by bootstrap

resampling of mutations, and calculations of the dNdS ratio for each bootstrap

sample.

Mutational signatures. Mutational signatures for human subjects were downloaded

from the COSMIC database (https://cancer.sanger.ac.uk/signatures/ ; version 3.1 as of

11.08.2020). Mutation data was binned using a bin half-width of 0.5 on the inverse

allele frequency. Exposure of binned data was determined using R 3.5.2 with package

YAPSA (version 1.8.0). Uncertainty on mutational signatures was determined by

bootstrap resampling of mutations and generation of the binned data and YAPSA

exposures on the resampled data. We have used 1000 bootstrap replicates as a

compromise between an ideally larger (1M) number of replicates, and reasonable

computing time. Large mutation sets (>100,000 mutations) were subsampled to

50,000-60,000 mutations for the bootstrap analysis. Mean, median, percentiles and

95% confidence bounds were determined using the resulting bootstrap distribution.

Time course. We made the assumption that clock-like mutational signatures SBS1

and SBS5 were a surrogate indicator for time (SBS1 only for glioblastoma, in

agreement with Alexandrov et al. 2015). We further assumed that the arrow of time,

could be identified with the arrow of inverse allele frequency $1/f$. Under these

assumptions, an increment of time can be deteremined in arbitrary units, by

integrating θ, which denotes the exposure to clock-like mutations SBS1 or SBS5, over an

increment of inverse allele frequency $1/f$. Since the exposure θ is also proportional to

the number of cells in the tumor, it is necessary to normalize θ, by dividing its value

by the number of cells N. Since N is proportional to $1/f$ under some assumptions (S1

File), N can be replaced, up to an unknown constant, by $1/f$. This yielded the formula

for determining time t in arbitrary units, over an interval of time which is unknown,

but identifiable with an interval over inverse allele frequency $1/f$:

378
$$t_{a.u.} = \int_{(1/f)\,min}^{(1/f)\,max} \theta \cdot f \cdot d(1/f), \qquad (1)$$

where $t_{a.u.}$ is the time in arbitrary units (a.u.), and θ corresponds to the exposure to clock-like

mutational signatures. By computing $t_{a.u.}$ at all values of $1/f$, we obtained a vector of values

$T_{a.u.}$ for the time in arbitrary units, from its minimum value, $\min(T_{a.u.})$ (0 by definition), to its

maximum value, $\max(T_{a.u.})$ (which corresponds to integration from $(1/f)_{min}$ to $(1/f)_{max}$.

Confidence limits at 95% for time were calculated using the confidence bounds for mutational

signature SBS1, taken as exposure θ.

Time calibration in recurrent glioblastomas. In a subset of glioblastoma samples, we

additionally know the time-to-relapse, denoted T, in months. We thereby identify the time

course of the relapse [0 ; T] in months, with the time course in arbitrary units [0 ; max($T_{a.u.}$)]

(see the preceding paragraph "Time course" for the calculation of $T_{a.u.}$). To obtain t, the time

in real units at any instant t in the interval [0 ; T], $t_{a.u.}$ is multiplied by the scaling factor

T/max($T_{a.u.}$):

$$t = t_{a.u.} \cdot T/max\left(T_{a.u.}\right). \tag{2}$$

Time propagation from primary to recurrent glioblastomas. To obtain a link between the

time course in the primary tumor, and the time course in the recurrent tumor, we studied the

ratio of mutation accumulation between the end of primary tumor (subscripted 'P', taken as

the last 5% time points) and start of recurrence (subscript 'R', first 5% time points), over the

entire tumor. This is justified by the observation that the passage from the primary tumor to

the recurrence corresponds to the instant of primary tumor resection. Using Eq. 1 (S1 File),

this ratio could be written as follows:

$$\frac{\left(dM/dt\right)_P}{\left(dM/dt\right)_R} = \frac{\left(\mu(t) \cdot \pi \cdot \omega(t) \cdot \gamma(t) \cdot N(t)\right)_P}{\left(\mu(t) \cdot \pi \cdot \omega(t) \cdot \gamma(t) \cdot N(t)\right)_R}, \tag{3}$$

where subscript i per subclone is not used, because we considered here that evolution

parameters are taken over the whole tumor. The constant term π can be normalized out of this

ratio. Further, we have assumed that the mutation rate μ and division rate ω stay constant over

this short period, because they are intrinsic features of the tumor cells. However, the count of

tumor cells $N(t)$, and the tumor cell survival rate $\gamma(t)$ could not be considered constant.

Regarding $N(t)$, we expressed it as the ratio of inverse allele frequency, since it is proportional

to N [13]:

407
$$\frac{N(t)_P}{N(t)_R} = \frac{(1/f)_P}{(1/f)_R}. \tag{4}$$

Of note, expression (11) is biased in practice by mutations which are not de novo in the

recurrence, but inherited from the primary tumor. Ideally, only de novo mutations should be

included to perform this calculation. Finally, the survival rate of tumor cells, γ, also could not

be considered constant, and we had no indicator or surrogate for this value. For this reason,

we have set an arbitrary value for the survival at end of primary tumor, relatively to the start

of recurrence, $\gamma_P/\gamma_R = 1/300$.

Using the above, we could obtain expression (5) for dM/dt at end of primary tumor:

415
$$\left(\frac{dM}{dt}\right)_P = \frac{(1/f)_P}{(1/f)_R} \cdot \frac{\gamma_P}{\gamma_R} \cdot \left(\frac{dM}{dt}\right)_R. \tag{5}$$

416 From this, and since the number of mutations at the end of the primary tumor, as well as the

417 rest of parameters, was known, the time in real units at the end of primary tumor could be

418 determined as follows:

$$419 \qquad dt_P = \frac{(dM)_P}{\dfrac{(1/f)_P}{(1/f)_R} \cdot \dfrac{\gamma_P}{\gamma_R} \cdot \left(\dfrac{dM}{dt}\right)_R} . \qquad (6)$$

420 Similarly to the time course in the recurrence, the fact that the last instant of the recurrence,

421 dt_P, was known, allowed to calibrate the time course t in the primary at each instant, by

422 multiplying by the scaling factor $dt_P / dt_{a.u.,P}$:

$$423 \qquad dt = dt_{a.u.} \cdot \frac{dt_P}{dt_{a.u.,P}} . \qquad (7)$$

Tumor cell survival ratio. For time calibration to real units, we have made an

assumption on tumor cell survival ratio γ_R/γ_P to determine real time in the primary

tumor. To quantify the survival ratio, we have proceeded the other way around, using

an assumption on the time course in the primary tumor, in order to determine the ratio

γ_R/γ_P. We have taken the assumption that the time between the most recent common

ancestor (TMRCA) lies either 2 years or 7 years before primary tumor resection.

These durations correspond to the shortest and longest time spans from [15]. The

calculation of the tumor cell survival ratio was done by reformulating expression (6)

into the following expression:

$$433 \qquad \frac{\gamma_R}{\gamma_P} = \frac{\left(\dfrac{dM}{dt}\right)_R}{\dfrac{(1/f)_R}{(1/f)_P} \cdot \left(\dfrac{dM}{dt}\right)_P} , \qquad (8)$$

434 where all terms on the right hand side are known, either from the data (dM, f, dt_R) or

435 from assumptions (dt_P).

Tumor expansion profile. From equation (8) in S1 File, time and *1/f* are

proportional, with, as modulators, the growth rate ω, the tumor cell survival rate γ,

and the number of tumor cells *N*:

$$d\left(1/f\right) \propto \omega \cdot \gamma \cdot N \cdot dt, \qquad (9)$$

where the increment *d(1/f)* is known from the data, and increment *dt* could be

determined above, in section "Time course". As a consequence, one obtains the

product $\omega\gamma N$, by dividing the increment *d(1/f)* by the increment *dt*. The expansion

parameters $\omega\gamma N$ are known within the constants of expression (8) from S1 File, which

are $\pi/K_{i,r}$. The curves giving $\omega\gamma N$ as a function of time are denoted expansion curves

or expansion profiles.

Curve segmentation. Segments for curves M(1/f) were determined using R package

segmented (v1.1-0), using an objective adjusted R^2 set to 0.995 (P. virginalis) and

0.9995 (GBM), and using the lowest number of segments which attained this

objective, limited to a maximum of 20 breakpoints. P-values of significant changes

between segments were evaluated using Davies' test (implementation from R package

segmented). Because the regions before the first breakpoint, and after the last

breakpoint, display marked and consistent differences with the general profile of the

curve, we hypothesized that the automated segmentation revealed clonal mutations, or

mutations which could originate from contamination by normal tissue (mutations with

allele frequency lower than the first breakpoint), as well as mutations likely affected

by the limit of detection of SNVs on sequencing data (mutations with allele frequency

higher than the last breakpoint), We exclude those mutations, and restrict the accepted

range to the interval between the first and last breakpoints.

Classification of expansion profiles. Expansion profiles $\omega\gamma N(t)$ were visually

inspected. This led to the identification of four curve categories (Fig 3). The curves

were then manually curated to one of these four categories.

Statistical analyses. R [40] and Python [41] were used for all statistical analyses.

Confidence intervals at 95% probability for the tree root in *P. virginalis* are taken as

the 95% highest posterior density (HPD) interval. All statistical tests were unpaired

and two-sided, with a level of significance set at 5%. Segmentation p-values

correspond to the test of significant difference between segments (Davies' test, R

package segmented). Correlation coefficients between SBS1 and SBS5 were

determined using Pearson method, and summarized by their median and IQR over

GBM samples. A comparison between groups was made using an unpaired Wilcoxon

rank-sum test. Differential time to recurrence between subgroups in the manually

sorted $\omega\gamma N$ curves was assessed using a wilcoxon rank-sum test against curve type 3.

Differences on the time to recurrence in the systematic analysis was explored by

analysis of variance against the number of maxima in the primary tumors and in the

recurrence, with Bonferroni adjustment of *p*-values. The difference of time to

recurrence between subgroups based on the number of maxima in the primary tumors

was further explored using a non-parametric wilcoxon test and using a F-test of

comparison of variances. The association of the γ_R/γ_P ratio (n=20) with the pattern of

$\omega\gamma N$ curves was investigated using a F-test of comparison of variances. The

association of the γ_R/γ_P ratio with the time to recurrence was assessed with a linear

regression, using a simple or double $\log_{10}$ scale on the γ_R/γ_P ratio, with Bonferroni

adjustment.

Acknowledgements

We would like to thank Verena Körber and Thomas Höfer for helpful discussions and for providing data, and Julian Gutekunst for discussions about the methods. We would also like to thank Katharina Hanna for data and for crayfish culture, and Sina Tönges for sample processing. We further acknowledge the German Cancer Research Center Genomics and Proteomics Core Facility for whole-genome sequencing.

493 **References**

1.   Lynch M, Ackerman MS, Gout JF, Long H, Sung W, Thomas WK, et al. Genetic drift, selection and the evolution of the mutation rate. Nat Rev Genet. 2016 Oct 14;17(11):704–14.

2.   Kent DG, Green AR. Order Matters: The Order of Somatic Mutations Influences Cancer Evolution. Cold Spring Harb Perspect Med. 2017 Apr 3;7(4):a027060.

3.   Martincorena I, Raine KM, Gerstung M, Dawson KJ, Haase K, Van Loo P, et al. Universal Patterns of Selection in Cancer and Somatic Tissues. Cell. 2017 Nov 16;171(5):1029-1041.e21.

4.   Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SAJR, Behjati S, Biankin AV, et al. Signatures of mutational processes in human cancer. Nature. 2013 Aug 22;500(7463):415–21.

5.   Tataru P, Simonsen M, Bataillon T, Hobolth A. Statistical Inference in the Wright-Fisher Model Using Allele Frequency Data. Syst Biol. 2017 Jan 1;66(1):e30–46.

6.   Benton ML, Abraham A, LaBella AL, Abbot P, Rokas A, Capra JA. The influence of evolutionary history on human health and disease. Nat Rev Genet. 2021 May;22(5):269–83.

7.   Martincorena I, Campbell PJ. Somatic mutation in cancer and normal cells. Science. 2015 Sep 25;349(6255):1483–9.

8.   Katju V, Bergthorsson U. Old Trade, New Tricks: Insights into the Spontaneous Mutation Process from the Partnering of Classical Mutation Accumulation Experiments with High-Throughput Genomic Approaches. Genome Biol Evol. 2018 Nov 26;11(1):136–65.

9. Gutekunst J, Andriantsoa R, Falckenhayn C, Hanna K, Stein W, Rasamy J, et al. Clonal genome evolution and rapid invasive spread of the marbled crayfish. Nat Ecol Evol. 2018 Mar;2(3):567–73.

10. Maiakovska O, Andriantsoa R, Tönges S, Legrand C, Gutekunst J, Hanna K, et al. Genome analysis of the monoclonal marbled crayfish reveals genetic separation over a short evolutionary timescale. Commun Biol. 2021 Jan 18;4(1):1–7.

11. Lyko F. The marbled crayfish (Decapoda: Cambaridae) represents an independent new species. Zootaxa. 2017 Dec 12;4363(4):544–52.

12. Alexandrov LB, Kim J, Haradhvala NJ, Huang MN, Tian Ng AW, Wu Y, et al. The repertoire of mutational signatures in human cancer. Nature. 2020 Feb;578(7793):94–101.

13. Williams MJ, Werner B, Barnes CP, Graham TA, Sottoriva A. Identification of neutral tumor evolution across cancer types. Nat Genet. 2016 Mar;48(3):238–44.

14. Balaparya A, De S. Revisiting signatures of neutral tumor evolution in the light of complexity of cancer genomic data. Nat Genet. 2018 Dec;50(12):1626–8.

15. Körber V, Yang J, Barah P, Wu Y, Stichel D, Gu Z, et al. Evolutionary Trajectories of IDHWT Glioblastomas Reveal a Common Path of Early Tumorigenesis Instigated Years ahead of Initial Diagnosis. Cancer Cell. 2019 Apr 15;35(4):692-704.e12.

16. Yang S, Wang L, Huang J, Zhang X, Yuan Y, Chen JQ, et al. Parent-progeny sequencing indicates higher mutation rates in heterozygotes. Nature. 2015 Jul 23;523(7561):463–7.

17. Liu H, Jia Y, Sun X, Tian D, Hurst LD, Yang S. Direct Determination of the Mutation Rate in the Bumblebee Reveals Evidence for Weak Recombination-Associated Mutation and an Approximate Rate Constancy in Insects. Mol Biol Evol. 2017 Jan;34(1):119–30.

18. Oppold AM, Pedrosa JAM, Bálint M, Diogo JB, Ilkova J, Pestana JLT, et al. Support for the evolutionary speed hypothesis from intraspecific population genetic data in the non-biting midge Chironomus riparius. Proc Biol Sci. 2016 Feb 24;283(1825):20152413.

19. Flynn JM, Chain FJJ, Schoen DJ, Cristescu ME. Spontaneous Mutation Accumulation in Daphnia pulex in Selection-Free vs. Competitive Environments. Mol Biol Evol. 2017 Jan;34(1):160–73.

20. Ho EKH, Macrae F, Latta LC, McIlroy P, Ebert D, Fields PD, et al. High and Highly Variable Spontaneous Mutation Rates in Daphnia. Mol Biol Evol. 2020 Jun 10;37(11):3258–66.

21. Keightley PD, Ness RW, Halligan DL, Haddrill PR. Estimation of the spontaneous mutation rate per nucleotide site in a Drosophila melanogaster full-sib family. Genetics. 2014 Jan;196(1):313–20.

22. Keightley PD, Pinharanda A, Ness RW, Simpson F, Dasmahapatra KK, Mallet J, et al. Estimation of the spontaneous mutation rate in Heliconius melpomene. Mol Biol Evol. 2015 Jan;32(1):239–43.

23. Ohno M. Spontaneous de novo germline mutations in humans and mice: rates, spectra, causes and consequences. Genes Genet Syst. 2019 Apr 9;94(1):13–22.

24. Lee-Six H, Øbro NF, Shepherd MS, Grossmann S, Dawson K, Belmonte M, et al. Population dynamics of normal human blood inferred from somatic mutations. Nature. 2018 Sep;561(7724):473–8.

25. Bae T, Tomasini L, Mariani J, Zhou B, Roychowdhury T, Franjic D, et al. Different mutational rates and mechanisms in human cells at pregastrulation and neurogenesis. Science. 2018 Feb 2;359(6375):550–5.

26. Gerlinger M, Horswell S, Larkin J, Rowan AJ, Salm MP, Varela I, et al. Genomic architecture and evolution of clear cell renal cell carcinomas defined by multiregion sequencing. Nat Genet. 2014 Mar;46(3):225–33.

27. Jamal-Hanjani M, Wilson GA, McGranahan N, Birkbak NJ, Watkins TBK, Veeriah S, et al. Tracking the Evolution of Non–Small-Cell Lung Cancer. N Engl J Med. 2017 Jun;376(22):2109–21.

28. Drummond AJ, Rambaut A. BEAST: Bayesian evolutionary analysis by sampling trees. BMC Evol Biol. 2007 Nov 8;7:214.

29. Alexandrov LB, Jones PH, Wedge DC, Sale JE, Campbell PJ, Nik-Zainal S, et al. Clock-like mutational processes in human somatic cells. Nat Genet. 2015 Dec;47(12):1402–7.

30. Petljak M, Alexandrov LB, Brammeld JS, Price S, Wedge DC, Grossmann S, et al. Characterizing Mutational Signatures in Human Cancer Cell Lines Reveals Episodic APOBEC Mutagenesis. Cell. 2019 Mar 7;176(6):1282-1294.e20.

31. Baez-Ortega A, Gori K, Strakova A, Allen JL, Allum KM, Bansse-Issa L, et al. Somatic evolution and global expansion of an ancient transmissible cancer lineage. Science. 2019 Aug 2;365(6452):eaau9923.

32. Rubanova Y, Shi R, Harrigan CF, Li R, Wintersinger J, Sahin N, et al. Reconstructing evolutionary trajectories of mutation signature activities in cancer using TrackSig. Nat Commun. 2020 Feb 5;11(1):731.

33. DeWitt WS, Harris KD, Ragsdale AP, Harris K. Nonparametric coalescent inference of mutation spectrum history and demography. Proc Natl Acad Sci U S A. 2021 May 25;118(21):e2013798118.

34. Kondo A, Safaei R, Mishima M, Niedner H, Lin X, Howell SB. Hypoxia-induced enrichment and mutagenesis of cells that have lost DNA mismatch repair. Cancer Res. 2001 Oct 15;61(20):7603–7.

35. Pirzkall A, McGue C, Saraswathy S, Cha S, Liu R, Vandenberg S, et al. Tumor regrowth between surgery and initiation of adjuvant therapy in patients with newly diagnosed glioblastoma. Neuro-Oncol. 2009 Dec;11(6):842–52.

36. Okolie O, Bago JR, Schmid RS, Irvin DM, Bash RE, Miller CR, et al. Reactive astrocytes potentiate tumor aggressiveness in a murine glioma resection and recurrence model. Neuro-Oncol. 2016 Dec;18(12):1622–33.

37. Scott JN, Rewcastle NB, Brasher PM, Fulton D, MacKinnon JA, Hamilton M, et al. Which glioblastoma multiforme patient will become a long-term survivor? A population-based study. Ann Neurol. 1999 Aug;46(2):183–8.

38. Monteiro AR, Hill R, Pilkington GJ, Madureira PA. The Role of Hypoxia in Glioblastoma Invasion. Cells. 2017 Nov 22;6(4):45.

39. Sabelström H, Quigley DA, Fenster T, Foster DJ, Fuchshuber CAM, Saxena S, et al. High density is a property of slow-cycling and treatment-resistant human glioblastoma cells. Exp Cell Res. 2019 May 1;378(1):76–86.

40. R Core Team. R: A Language and Environment for Statistical Computing. Vienna: R Foundation for Statistical Computing; 2018.

41. Van Rossum G, Drake Jr FL. Python reference manual. Amsterdam: Centrum voor Wiskunde en Informatica; 1995.

Data availability

Sequence data for marbled crayfish data have been deposited as a National Center for

Biotechnology Information BioProject (accession number: PRJNA356499).

Glioblastoma data were accessed from the European Genome-phenome Archive

(EGA) database, with accession number: EGAS00001003184 (glioblastoma).

Supporting Information

S1 Fig. Mutation accumulation, selection and time dynamics of GBM tumors. Panels A1-D1 show the primary tumor T1, panels A2-D2 show the recurrent tumor T2. (A) Mutation accumulation as a function of the inverse of allele frequency $1/f$ (black) and phases from automated segmentation (breakpoints (grey) and segments (blue)). The confidence band at 95% level is indicated in grey. (B) Nonsynonymous to synonymous ratio. Purple and blue stars show nonsynonymous and synonymous mutations, respectively. The smoothened ratio is shown in red. (C) Clock-like and non-clock-like mutational signatures. (D) Mutation accumulation as a function of time.

S2 Fig. Transition of primary tumor to recurrent tumor. (A) Dynamics of growth rate ω times tumor cell survival rate γ times number of cells N, for (P) the primary tumor and (R) the recurrence. (B) Time-resolved mutation accumulation for primary tumor and recurrence.

S1 File. Supplementary Methods.

S2 File. Supplementary Tables.