Complete pan-plastome sequences enable high resolution phylogenetic classification of sugar beet and closely related crop wild relatives

Katharina Sielemann[1,2], Boas Pucker[1,3,4], Nicola Schmidt[5], Prisca Viehöver[1], Bernd Weisshaar[1], Tony Heitkam[5*], Daniela Holtgräwe[1*]

[1]Genetics and Genomics of Plants, Center for Biotechnology (CeBiTec) & Faculty of Biology, Bielefeld University, 33615 Bielefeld, Germany

[2]Graduate School DILS, Bielefeld Institute for Bioinformatics Infrastructure (BIBI), Bielefeld University, 33615 Bielefeld, Germany

[3]Evolution and Diversity, Department of Plant Sciences, University of Cambridge, Cambridge CB2 3EA, United Kingdom

[4]Institute of Plant Biology, TU Braunschweig, Braunschweig, Germany

[5]Faculty of Biology, Institute of Botany, Technische Universität Dresden, 01069 Dresden, Germany

* Correspondence (These authors share senior authorship.):
Tony Heitkam, Daniela Holtgräwe
tony.heitkam@tu-dresden.de, dholtgra@cebitec.uni-bielefeld.de

Keywords: sugar beet, *Beta vulgaris*, *Beta*, *Corollinae*, *Patellifolia*, chloroplast, plastome assembly, phylogeny, phylogenomics

Short title: Pan-plastome of cultivated and wild beets

Abstract

Background

As the major source of sugar in moderate climates, sugar-producing beets (*Beta vulgaris* subsp. *vulgaris*) have a high economic value. However, the low genetic diversity within cultivated beets requires introduction of new traits, for example to increase their tolerance and resistance attributes – traits that often reside in the crop wild relatives. For this, genetic information of wild beet relatives and their phylogenetic placements to each other are crucial. To answer this need, we sequenced and assembled the complete plastome sequences from a broad species spectrum across the beet genera *Beta* and *Patellifolia*, both embedded in the Betoideae (order Caryophyllales). This pan-plastome dataset was then used to determine the wild beet phylogeny in high-resolution.

Results

We sequenced the plastomes of 18 closely related accessions representing 11 species of the Betoideae subfamily and provided high-quality plastome assemblies which represent an important resource for further studies of beet wild relatives and the diverse plant order Caryophyllales. Their assembly sizes range from 149,723 bp (*Beta vulgaris* subsp. *vulgaris*) to 152,816 bp (*Beta nana*), with most variability in the intergenic sequences. Combining plastome-derived phylogenies with read-based treatments based on mitochondrial information, we were able to suggest a unified and highly confident phylogenetic placement of the investigated Betoideae species.

Our results show that the genus *Beta* can be divided into the two clearly separated sections *Beta* and *Corollinae*. Our analysis confirms the affiliation of *B. nana* with the other *Corollinae* species, and we argue against a separate placement in the *Nanae* section. Within the *Patellifolia* genus, the two diploid species *Patellifolia procumbens* and *Patellifolia webbiana* are, regarding the plastome sequences, genetically more similar to each other than to the tetraploid *Patellifolia patellaris*. Nevertheless, all three *Patellifolia* species are clearly separated.

Conclusion

In conclusion, our wild beet plastome assemblies represent a new resource to understand the molecular base of the beet germplasm. Despite large differences on the phenotypic

56 level, our pan-plastome dataset is highly conserved. For the first time in beets, our whole
57 plastome sequences overcome the low sequence variation in individual genes and provide
58 the molecular backbone for highly resolved beet phylogenomics. Hence, our plastome
59 sequencing strategy can also guide genomic approaches to unravel other closely related
60 taxa.

Background

As the crop plant *Beta vulgaris* (sugar beet) has a high economic value (1), continuous crop
development is essential to enhance stress tolerances and resistances against pathogens.
The White Silesian Beet provided the germplasm for sugar beet (2) and is a derivative of
wild sea beet (*B. vulgaris subsp. maritima*). This leaves, similar to the situation in many
domesticated crops, only a narrow genetic base for sugar beet breeding (3). Additionally,
early sugar beet breeding has focused mainly on increasing yield. This caused strong
domestication bottle necks and removed many useful traits that may benefit plant fitness
(3,4). The higher genetic variation in crop wild relatives of sugar beet offers potential that
might be harnessed to introduce desired traits. Thus, giving insight into the genomic basis
of wild beets is progressively moving into the focus of beet breeding research (5,6).

Phylogenetically, wild beets belong to the Betoideae (order Caryophyllales) and are
separated in the genera *Beta* and *Patellifolia* (7,8), with all cultivated beets belonging to the
genus *Beta* (9). The genus *Beta* is then further subdivided into at least two sections, *Beta*
and *Corollinae*. In general, the section *Beta* is widespread across Western Europe, whereas
Corollinae species are generally distributed across the eastern Mediterranean area and
South-West Asia (1,8) (Additional file S1A). Despite the long history of different systematic
treatments (1,7–12), the phylogenetic relationships of *Beta* and *Patellifolia* species are still a
matter of ongoing debate (as reviewed in (7)). Especially the subdivision of the genus *Beta*
into three sections (*Beta*, *Corollinae*, and *Nanae*) is discussed, with the pending suggestion
to integrate *B. nana* into the section *Corollinae*, hence disbanding the section *Nanae*
(7,8,13). Similarly, as the *Beta* section *Corollinae* harbors a highly variable polyploid/hybrid
complex, including di-, tri-, tetra-, penta-, and hexaploid forms, the species boundaries are
far from resolved. Regarding the sister genus, there is still an ongoing discussion on
whether the morphologically variable *Patellifolia* comprise three distinct species (*P.*

patellaris, *P. procumbens*, and *P. webbiana*) or only two or even one (8,10,12). Resolving
the unclear wild beet relationships may inform beet improvement programs and contribute
to the development of new, better equipped beets.

The plastome is well-suited for the reconstruction of phylogenies due to high structural
conservation, a conserved evolutionary rate, uniparental inheritance, and high abundance
of DNA across all species (14,15). Historically, systematic information was obtained from
plastome sequence restriction site variants, inversions, single nucleotide variants (SNVs), or
spacers in single genes. Although this has led to a range of wild beet phylogenies resolving
relationships on the level of genera and sections, these are often based on only a few
species and contain collapsed branches due to low genetic variation. In contrast, the
investigation of whole plastome sequences may enhance the resolution of phylogenetic
relationships (14,16–18). As gene sequences and intergenic regions can be included and
combined, whole plastome sequence analyses enable the detection of well-supported
phylogenetic relations on the species- and even on the accession level. Thus, plastid
genomics may offer a route to clarify many of the pending questions regarding the wild beet
phylogeny.

The plastome sequence of most angiosperms comprises a total of 79 protein-coding genes,
4 rRNA genes, and 30 tRNA genes (19). The quadripartite structure is characteristic for
plastome sequences comprising a large (LSC) and a small single-copy region (SSC) as well
as two inverted repeats (IRs) (20,21), all contributing to a total length of 120 kb to 210 kb
(20). This difference in size can be mainly attributed to the IRs that range from 6 kb to 76 kb
in length (21–23). The relative orientation of the SSC between the IRs differentiates two
structural variants which occur simultaneously in a single cell and might have been
previously mistakenly annotated as differences between species (24).

The Caryophyllales, including *B. vulgaris*, contain canonical plastomes, harboring all
hallmarks typical for angiosperm plastome sequences as described above (25,26). For the
wild beet species, until now, no plastome assembly is available, and of our investigated
species, only the plastome sequence of *B. vulgaris* subsp. *vulgaris* was published
previously (27). A detailed, plastome- and mitogenome-based evolutionary positioning of
species outside of the section *Beta* is still missing but needed to answer some of the
unresolved issues in beet systematics.

Here, we resolve the phylogenetic relationships within the Betoideae at high resolution
through genome-wide comparison based on complete plastome assemblies and reads from
both, the plastome and the mitogenome. Eleven different *Beta* and *Patellifolia* members,
spanning the previously neglected plastome sequences of the *Corollinae* section and the
Patellifolia genus, are included in our analyses. For this, whole plastome sequences of up to
two accessions per species are sequenced, assembled, and compared. This novel
contribution to the Betoideae pan-plastome intends to clarify the phylogenetic relationships
of wild beets on a species-level and provides an important resource for further studies of
beet wild relatives.

Results

Our pan-plastome dataset comprises 18 different accessions, including a biological
replicate of *B. corolliflora*, which leads to 19 plastome assemblies in total (see Methods,
Table 2). To provide a basis for comparative plastome analysis, all plastome sequences
were fully assembled. Out of those, 17 were split into three scaffolds (LSC, SSC, and IR),
apart from Bmar1 (four scaffolds) and Bnan2 (six scaffolds). Collapsed IR regions were
confidently identified in all plastome assemblies based on a doubled average read coverage
in comparison to the single copy regions as well as a gene content that is characteristic and
expected for the IR region. Average read coverage and assembly length are shown in Table
1. The distribution of these values is shown in Additional file S1B. Circular and linear plots
of a representative selection of plastome assembly sequences are provided in Additional file
S1C.

Table 1: Plastome assembly statistics. Average read coverage values and assembly
lengths (in bp) for each region and the complete assemblies are shown. Abbreviations: LSC
= Long single copy region; SSC = Short single copy region; IR = Inverted repeats.

Total coverage	LSC coverage	SSC coverage	IR coverage	Total length	LSC length	SSC length	IR length

396 ± 73	261 ± 81	265 ± 98	662 ± 100	150,519 ± 892	83,496 ± 621	17,845 ± 257	24,588 ± 307

Comparing the plastome assemblies of all *Beta* vs. *Patellifolia* species, the average length
of the four *Patellifolia* plastome sequences (avg. 151,621 bp) is higher than for the 15 *Beta*
plastome sequences (avg. 150,225 bp). This length difference can be mainly assigned to
the LSC (avg. *Beta* 83,401 bp; avg. *Patellifolia* 83,853 bp) and to the IRs (avg. *Beta* 24,435
bp; avg. *Patellifolia* 25,166 bp). However, the SSC is longer in *Beta* plastome sequences
(avg. 17,954 bp) when compared to the plastome sequences of all *Patellifolia* accessions
(avg. 17,437 bp).

Interestingly, plastome assemblies of *B.* section *Corollinae* (avg. 36.67 %) show a higher
GC content when compared to *B.* section *Beta* plastome sequences (avg. 35.81 %). The
total length of *B.* section *Corollinae* plastome sequences (avg. 150,504 bp from nine
species) is higher in comparison to *B.* section *Beta* plastome assemblies (avg. 149,808 bp
from six species). This length difference is visible for all regions of the plastome sequence
(LSC, SSC and IRs).

The final plastome assemblies were subsequently annotated and the alignment identity of
all regions included in the phylogenetic analysis was assessed for gene regions and
intergenic regions, respectively (Figure 1). The alignment identity is significantly higher for
gene sequences when compared to intergenic regions. This significant difference was
obtained when amaranth, quinoa, and spinach were included as outgroups (Additional file
S2A) (avg. gene/intergenic regions 90.73/83.55 %; Mann-Whitney-U test; p 2e-10) as well
as without outgroup reference sequences (Additional file S2B) (avg. gene/intergenic regions
97.26/94.93 %; Mann-Whitney-U test; p ≈ 1e-09). *Rrn* genes show high similarity among all
plastome genes, whereas *ycf1* and *rpl22* show the greatest variance between all
investigated accessions. The intergenic region between the genes *ycf4* and *cema*
contributes most to the differences in the alignment.

Figure 1: Alignment identities of gene sequences (A) and intergenic regions (B). The sequences (x-axis) are ordered based on the alignment identity represented by the y-axis. The alignment identity including amaranth, quinoa, and spinach as outgroup (light blue) as well as the alignment identity without the reference plastome sequences (dark blue) are shown.

The distributions of SNVs (Figure 2A) and InDels (Figure 2B) throughout the plastome sequences were further investigated. InDels are mostly absent from gene regions and SNVs are in general more frequent than InDels. Further, some clear hotspots of SNVs and

InDels can be detected in the intergenic regions *psbK-psbI*, *ycf4-cema*, *rpl33-rps18*, *psaj-*
rpl33 and *rpl32-ccsa*.

Figure 2: Number of SNVs and InDels in the assembled plastome sequences. Number
of SNVs (A) and number of InDels (B) in the sequence alignments normalised by the length
of the respective gene sequence/intergenic region and by the number of
species/accessions. The gene names and intergenic regions on the x-axis are ordered
based on the arrangement in the plastome assembly. Amaranth, quinoa, and spinach were
included as outgroup.

As InDels with a length, which is a multiple of three, do not influence the reading frame (28),
we expected that the proportion of these InDels (which are multiples of three) is higher in
gene regions compared to the proportion in intergenic regions. Indeed, we observed that
43.4 % of the InDels in gene regions were a multiple of three, whereas this applies to only
29.6 % of the InDels in intergenic regions (Fisher's exact test; p ≈ 3e-8; Figure 3 [arrows]).

Figure 3: Number of InDels based on size in gene sequences (A) and intergenic
regions (B). The arrows represent InDels with a size of a multiple of three. Please mind the
variable y-axis.

Phylogenetic relations of the Betoideae subfamily were inferred from colored de Bruijn

graph-based splits (Figure 4) as well as by an alignment-based maximum likelihood (ML)

analysis (Figure 5). Mitochondrial and plastid read derived kmers were used to calculate

phylogenetic splits, and annotated gene sequences as well as intergenic regions derived

from the plastome assemblies were used for the ML analysis. A clear separation of

Patellifolia, *B.* section *Beta* and *B.* section *Corollinae* samples is visible in all four

phylogenetic trees. In comparison to the ML tree based on fully assembled plastome

sequences (Figure 4C, black), the tree based on splits derived from the same dataset (fully

assembled plastome sequences) (Figure 4C, dark green) shows only one difference among

the *B. intermedia* and *B. corolliflora* accessions. The phylogenetic relationships derived from

kmers show a few additional differences (Figure 4C, light green and pink). These

differences comprise for example the assignment of the four *Patellifolia* accessions/species

to a clade consisting of both *P. patellaris* accessions and a separate clade formed by *P.*

procumbens and *P. webbiana* for the assembly-based phylogenies (Figure 4C, black and

dark green), whereas the kmer based phylogenies show a separate clade for *P. webbiana*

and a second clade comprising the other three *Patellifolia* accessions/species (Figure 4C,

light green and pink). The calculation of the weighted F1 score, the weighted symmetric set

distance and the Robinsons-Foulds distance shows that there is a high identity between all

splitstree results (based on cp_reads, mt_reads and cp_assemblies) (Additional file S1D).

The usage of different input data formats (reads vs. assemblies) has a larger impact than

the usage of different datasets (chloroplasts vs. mitochondria) meaning that these splitstree

results show a higher divergence.

Figure 4: Phylogenetic relationships of 18 different *Beta*/*Patellifolia* accessions
derived from different strategies and datasets. Kmer-based trees were constructed
using raw sequencing reads (A: mitochondrial reads, B: chloroplastic reads) as well as the
final chloroplast assemblies as input (not shown as only used for comparison). The
splitstree results (green and pink) were then compared to the ML analysis (black) (C).
Discordance between the phylogenetic trees is shown in the respective color.
(Abbreviations: cp=chloroplast; mt=mitochondria).

For the ML-based tree (Alignment sites / patterns: 216442 / 2441; Gaps: 0.44 %; Invariant
sites: 86.73 %), the phylogenetic relationships on the species level are highly supported
(high bootstrap values) (Figure 5). A phylogenetic tree based on the diagnostic set of 53
gene sequences and intergenic regions matches this phylogeny (Additional file S1E).

Figure 5: Plastome phylogeny for Betoideae. The tree can be divided into three groups: *B.* section *Corollinae* (8 accessions), *B.* section *Beta* (6 accessions), and *Patellifolia* (3 accessions). The plastome sequences of three Caryophyllales species (amaranth, quinoa, and spinach) were used as outgroup. Bootstrap support values are shown above each branch. The resulting phylogeny is based on the variation in 83 genes and 76 intergenic regions from the plastid genomes of 18 accessions and species (plus outgroup). Different background colors represent the sampling location and the origin of the breeding line (Bvul), respectively. Inset: Actual branch lengths based on the ML analysis.

To investigate the contribution of different regions to the phylogeny, in addition to the whole plastome sequences (genic and intergenic regions combined), sequence matrices for (I) all gene regions, (II) all intergenic regions, (III) whole coding sequences, (IV) first and second codon position and (V) third codon position only were constructed and used for the inference of phylogenetic relationships. Even though the topology of the phylogeny derived from whole coding sequences is highly similar (Additional file S1F), especially for the codon position-based matrices, alternative branches can be observed. However, substantially more nodes are only poorly supported with bootstrap values up to below 20.

The alignment of the 18s rRNA gene sequence for *B. corolliflora* (Bcor1) as representative for the *Corollinae* and *B. vulgaris subs. vulgaris* as member of the section *Beta* revealed not a single genomic difference and therefore no signal which could be used to resolve their phylogenetic relations.

The results presented here show that especially for closely related species of the same subfamily, a higher number of gene sequences and more variable intergenic regions provides greater phylogenetic resolution.

Discussion

Our proposed beet whole-plastome phylogeny is superior to single gene phylogenies

Here, we present the plastome sequence assemblies of 18 accessions covering most of the species' diversity within the beet genera *Beta* and *Patellifolia* and representing an important resource for future studies. All newly generated plastome sequences are highly similar including 79 protein coding genes and four rRNA genes distributed across a mean length of

150,519 bp (± 892bp). A previously published *B. vulgaris* subsp. *vulgaris* (KR230391)

plastome sequence comprises a length of 149,722 bp (27). This is almost identical to our

Bvul assembly which differs by only 1 bp in length (149,723 bp). This difference occurs in a

stretch of five or six guanines in the IR region between the genes *rrn23* and *rrn16* – either a

sequencing error or a biological difference. As expected, all Betoideae assemblies show

high similarity as all angiosperm plastomes are highly conserved and the species

investigated here are closely related, while most differences are located in intergenic

regions.

Between the cultivated beet and wild beet accessions, most chloroplast genes are highly

conserved, one example being *rpl2* with only a low number of polymorphisms (Figure 1)

(22). The intergenic regions are significantly less conserved containing more InDels and are

therefore more suitable for phylogenies on a lower taxonomic level (22). Among the beet

plastomes, *ycf1* is the most informative genic region (Figure 1A), which was also detected in

other plant groups, such as the Tropaeolaceae, the orchids, and the Malvales (29–31).

Additionally, the *rpl22*, *matk*, *clpP1*, *ycf2*, *psaC* and *ndhF* genes were reported to be highly

divergent (29,31,32), which is mainly consistent with our findings. Here, the rrn-genes, *ndhB*

(already classified as gene with low divergence (29,31)), *rps12* and *rps7* can be found

among the gene loci with lower variance among the investigated species.

Plastomes in general show very similar sequences with most differences occurring in non-

coding regions (33). For beet and wild beets, the most informative intergenic regions are

ycf4-cema, *accd-psai* and *rpl32-ccsa* (Figure 1B). However, for *ycf4-cema* the high

difference between 'reference' and 'without reference plastome-' alignment identity should

be noticed. For the Malvales, the regions *psaB-psaA*, *psbF-psbE*, *rpl2-rpl23* and *ndhH-ndhA*

were identified as lowly divergent regions (31). We confirmed these for beets, except for the

latter (*ndhH-ndhA*) that showed higher divergence among the wild beet plastomes. In

contrast, the intergenic regions with the highest divergence in the Malvales (*ndhD-ccsA* and

rps19-rpl2) (31) accounted for less differences among the wild beets. Nevertheless, the

percent identity values among all intergenic regions are relatively similar in beets, especially

when excluding the polymorphisms in the outgroup reference genomes.

As high variability among the investigated sequences is required to resolve phylogenetic

relations of closely related species (34), the retrieved plastome sequences from beets and

wild beets provide an excellent resource to approach systematic treatment of the Betoideae.
To further increase sequence variability, we also integrated the more diverged intergenic
regions into the analysis.

In addition to the plastome-inferred ML-based phylogeny, a mitogenome- and plastome-
inferred read-based phylogenetic tree was constructed based on phylogenetic splits. The
resulting trees can be considered reliable due to three distinct, robust properties: (I) all splits
networks show a tree-like appearance (Figure 4A, 4B), (II) the usage of different kmers
leads to the same tree and (III) the usage of the geometric mean leads to the removal of
samples in case no kmer occurs in the sample and both, geom and geom2 lead to the same
results (with the exception of k=11, which results in a cloud-shaped network).

Differences between the phylogenetic trees derived from different strategies and datasets
(Figure 4C) may be explained by chance and/or by the use of the method (all splitstree
results show high identities as indicated by all comparison metrics (Additional file S1D)
while differences are larger when using different strategies [reads vs assemblies] in
comparison to different datasets [chloroplasts vs. mitochondria]) or by real differences in the
biological nature of mitochondria and chloroplasts. Mitochondrial DNA shows a low
nucleotide substitution rate when compared to chloroplast DNA (25,35). Reasons for this
may be recent species hybridization or incomplete lineage sorting (36,37). Therefore, the
mitogenome seems to be mostly useful at higher taxonomic levels (25) and might not be the
most suitable system for the beet and wild beet accessions investigated in this study. In
summary, the trees based on plastome assemblies (ML and splits) are likely the most
reliable as the same phylogenetic relations are the outcome of different established
strategies including the widely used ML method.

Compared to our pan-plastome assembly, the information derived from individual genic and
intergenic regions are insufficient to fully resolve the beet phylogeny, highlighting the power
of our plastome approach:

I) Investigation of specific regions of the plastome sequence (genic and intergenic regions,
coding sequences, codon positions) revealed a few alternative branches for the codon
position-based sequence matrix (Additional file S1F). These are marked by short internal
branch lengths due to the close relationships of the species within this subfamily.
Nevertheless, these conflicting relationships are only weakly supported. This is explained by

the lower genetic diversity and therefore insufficient phylogenetic signal when using a

smaller amount of sequences and total sequence length.

II) An approach based on high-quality single-copy nuclear genes would require a minimum

coverage of about 10x (25). Moreover, nuclear genes are often part of gene families and

influenced by whole genome duplication events (14). Using our available data, nrDNA

sequences were selected for the phylogenetic reconstruction. Especially the ITS and ETS

regions were previously used for the investigation of phylogenetic relationships, but entire

nrDNA repeats (18S-ITS1-5.8S-ITS2-26/28S) were also already assembled for multiple

phylogenetic studies (15,38). Unfortunately, the assembled nrDNA sequences constructed

here are not useful to infer confident phylogenetic relationships as the coverage is very low

(1.9x - 8.5x) and intragenomic polymorphisms of different nrDNA repeat sequences might

limit the reliability of the phylogeny (15). The low bootstrap values and the low coverage

make this phylogenetic tree unreliable. Therefore, we do not show these results here.

Another possible explanation, apart from the low coverages, is that the biparental nature of

the nuclear genome may be problematic for the inference of phylogenetic relationships (14).

Previous studies already suggest that phylogenies based on nrDNA and few selected

plastid sequences only weakly support relationships (30). The 18s rRNA gene sequences of

representatives of the sections *Corollinae* and *Beta* are completely identical containing no

phylogenetic signal to separate them.

Summarizing, our plastome-derived phylogeny benefits from the incorporation of genic and

intergenic regions as well as the 'nature' of the plastome itself (as described in the

Background section). Despite the low available read coverage and the low genetic diversity

within our beet dataset, this leads to a highly confident phylogenetic tree. Further, the use of

higher alignment lengths and the use of nucleotides instead of amino acids are favored to

construct well supported phylogenies (39). Therefore, we conclude that for resolving the

relationships of cultivated and wild beets, our whole-plastome-based approach is the most

reliable.

Implications for the systematic placements within the Betoideae subfamily

With efforts tracing back half a century, resolving the phylogeny of the subfamily Betoideae

has been already a major undertaking (1,8–11,40). However, in most cases only few

selected sequence regions were targeted, leading to unresolved relations at shallower
taxonomic levels or with focus on specific species or sections, i.e.:

I) In a study by Hohmann et al. (2006), the *Beta* species *B. vulgaris*, *B. corolliflora*, *B. nana*
and *B. trigyna* were investigated using ITS, *trnL-trnF* spacer and *ndhF* sequences (10).
Kadereit et al. (2006) provide a comprehensive analysis of a high number of different
Betoideae species, finding that, *B.* section *Beta* was clearly separated from *B.* section
Corollinae, which contained *B. nana*, *B. trigyna*, *B. macrorhiza*, *B. corolliflora* and *B.*
lomatogona. As the analysis was based on ITS sequences comprising only 251 characters
of which 147 were invariable, relations between species in both sections could not be
resolved (8). Our study confirms the deep separation of the sections *Beta* and *Corollinae*
and refines the resolution on the species level.

II) A recent comprehensive study by Romeiras et al. (2016) of phylogenetic relationships in
the Betoideae is based on ITS and *matK*, *trnH-psbA*, *trnL* intron and *rbcL* sequences and
investigates a high number of different species leading to the following main result: *Beta*
and *Patellifolia* species are two clearly separated monophyletic groups (1). In total, three
monophyletic lineages were identified: *B.* section *Beta* (*B. vulgaris* subsp. *vulgaris*, *B.*
vulgaris subsp. *maritima*, *B. macrocarpa*, *B. patula*), *B.* section *Corollinae* (*B. nana*, *B.*
corolliflora, *B. trigyna*) and *Patellifolia* (*P. patellifolia*, *P. procumbens* and *P. webbiana*).
Exact relations within the Betoideae on a lower taxonomic level remain unclear as the
branches are not well supported and collapsed. We also identify the three monophyletic
groups as proposed and manage to resolve many of the previously collapsed branches.

III) Recently, Touzet et al. (2018) investigated the relationship of a wide range of *B. vulgaris*
subsp. *maritima*, *B. macrocarpa* and *B. vulgaris* subsp. *adanensis* accessions based on a
3,742 bp alignment of plastome sequences and a 1,715 bp alignment of selected nuclear
sequences (11). They find, based on a representative geographical sampling, that *B.*
macrocarpa is a distinct lineage from the two investigated *B. vulgaris* subspecies. Despite
this interesting finding, the suggested phylogeny did not focus on other important species
and accessions of the Betoideae subfamily and might be further improved by the analysis of
sequences with higher diversity to reach higher bootstrap values, which we achieved using
intergenic and genic regions of the whole plastome sequences.

With our pan-plastome-informed datasets, we have been able to confirm many of the
observations before and added an unprecedented resolution at the species-level. More in
detail, we conclude that:

I) Among the section *Beta*, the plastome sequences of *B. patula*, *B. vulgaris* subsp. *vulgaris*
and *B. vulgaris* subsp. *maritima* are highly similar as indicated by the slightly lower
bootstrap values (96/100) for these three beets. As this section harbors wild beets in
relatively close geographical proximity across the coastal Mediterranean area, the detected
similarity can be explained by (natural) crossing and gene flow due to close geographical
proximity or accidental cross-pollination during cultivation as wild beet and cultivated beet
groups are easily cross-compatible (3). Further, *B. vulgaris* subsp. *vulgaris* and some *B.*
vulgaris subsp. *maritima* accessions are even phenotypically highly similar (41). The
phylogenetic relationships among species can also be influenced by the geographical
distribution, mating systems and polyploidization (11). Allogamy and self-incompatibility are
characteristics of *B. vulgaris* subsp. *maritima*, whereas *B. macrocarpa* and *B. vulgaris*
subsp. *adanensis* are self-compatible leading to lower divergence and higher homozygosity.
Cross-compatibility can lead to hybridization by facilitating gene flow between individual
species (40), especially between *B. patula* and *B. vulgaris* subsp. *maritima*, which may
explain the lower bootstrap support and the more unclear relations in the phylogenetic tree
presented here. Although previous studies found low divergence between *B. vulgaris*
subspecies (11), *B. vulgaris* subsp. *adanensis* seems to be clearly separated from the other
subspecies in our analysis, possibly explained by the geographical distance to the other
investigated samples.

II) We confidently assigned specific species, including *B. nana*, to the section *Corollinae*: *B.*
corolliflora, B. intermedia, B. macrorhiza, B. lomatogona and *B. nana* cluster together and
form this *s*ection (also suggested by (8)). Here, we particularly focused on the *B. s*ection
Corollinae by analysing the plastome sequences of eight accessions from five different
species plus a biological replicate of the eponymous species *B. corolliflora. B. nana*, which
is endemic to Greece (1,8,42), was previously considered a separate *B.* section *Nanae*. Our
results, however, combined with multiple other studies, clearly show that *B. nana* falls within
the *B.* section *Corollinae*, which is distinct from *B.* section *Beta* (8,10,13). In addition to our
plastome-based phylogenetic analysis, further genomic evidence points to high genomic
similarities between *B. nana* and other *Corollinae***:** For example, these species are marked

by similar repeat accumulation profiles as shown for many individual transposable element
types (43–46). Regarding plant characteristics, frost tolerance and seed hardiness are
useful traits in the section *Corollinae*, including *B. nana*, but do not occur in any species of
the section *Beta* (47). Thus, frost tolerance is specific to the *Corollinae* when compared to
the *Beta* and *Patellifolia* species. These points lead to the classification of *B. nana* as a
member of the *Corollinae*. Considering the highly variable polyploid/hybrid status complexes
within the *Corollinae*, our plant set encompassed three diploids (*B. macrorhiza*, *B.*
lomatogona, and *B. nana*), a tetraploid (*B. corolliflora*), as well as a pentaploid (*B.*
intermedia). Although the hybrid status and parental contributions of the polyploids remain
unresolved (1,40), we present convincing evidence that *B. intermedia* and *B. corolliflora* are
closely related. Thus, our plastome sequence analysis brings new evidence supporting the
hypothesis that *B. corolliflora* and *B. intermedia* belong to a highly variable polyploid hybrid
complex (summarised by (7); Figure 5). The investigation of the whole genome sequences
of these polyploid species may help to resolve these parental contributions.

III) Among the *Patellifolia* members, *P. procumbens* and *P. webbiana* can be
phylogenetically distinguished: *Patellifolia* was previously classified as *B.* section
Procumbentes and there is still an ongoing taxonomic debate whether *P. patellaris*, *P.*
procumbens and *P. webbiana* can be considered as separate species. However, due to
molecular and morphological traits, *Patellifolia* are now mostly considered a separate genus
which is divided into three distinct species (8,10,12). The relationships among the
Patellifolia species could not be resolved in previous studies (1). In the phylogenetic tree
presented here, *P. procumbens* and *P. webbiana* seem to be closely related (however still
distinguished with high support) and clearly separated from the two *P. patellaris* accessions.
The branch lengths distinguishing *B. patula* and the *B. vulgaris* subsp. *vulgaris*/*maritima*,
which are both considered separate species, are highly similar (0.0001). The same branch
length separates *P. procumbens* and *P. webbiana*. Therefore, our phylogenetic analysis
indicates that the three *Patellifolia* species are distinguishable on a molecular level.

Comparing the results presented here with earlier studies, the previous investigation of
Betoideae was substantially extended and refined. The phylogenetic relationships were
resolved in more detail and not only based on the monophyletic groups. This is especially
important for the species of the *B.* section *Corollinae* which were investigated in depth.
Using the whole plastome sequences, including intergenic regions, it was possible to further

resolve the phylogenetic relationships with higher bootstrap support due to the extraction of

higher sequence variance and phylogenetic signal within the subfamily.

Conclusions

We provide 19 plastome assemblies for 18 different beet and wild beet accessions, which

can also be re-used for future investigations of beets and other Caryophyllales species, and

harnessed these to revisit systematic issues within the genera *Beta* and *Patellifolia*. This

analysis advanced our understanding of the phylogenetic relationships of the subfamily

Betoideae in four ways: I) Analysing sequences of intergenic regions of the whole plastome

assemblies made it possible to reveal the phylogeny of closely related species with high

reliability. Our phylogenetic tree shows a clear separation of the wild beet genera *Beta* and

Patellifolia, as well as of the two sections *Beta* and *Corollinae*. II) *B. vulgaris* subsp.

adanensis and *B. macrocarpa* can be clearly distinguished from *B. vulgaris* subsp. *vulgaris*,

B. vulgaris subsp. *maritima* and *B. patula*. A clear split of *B. patula* from the two *B. vulgaris*

subsp. (*B. vulgaris* subsp. *vulgaris* and *B. vulgaris* subsp. *maritima*) was not observed, likely

due to the high sequence identity possibly explained by the close geographical proximity

and the fact that these species are easily cross-compatible. III) All three *Patellifolia* species

are clearly separated in our phylogenetic analysis, while *P. procumbens* and *P. webbiana*

are more closely related to each other than to *P. patellaris*. These results, including the

investigation of the branch lengths, point to a molecular separation within the *Patellifolia*

species. IV) Finally, the taxonomic classification of *B. nana* as a member of the *Corollinae*

was further supported.

Methods

Plant material, genomic DNA extraction, and DNA sequencing

Seeds of Betoideae species were obtained from KWS Saat SE, Einbeck, Germany (*B.*

vulgaris subsp. *vulgaris* genotype KWS 2320) and from the Leibniz Institute of Plant

Genetics and Crop Plant Research Gatersleben (IPK), Germany (all other accessions with

accession numbers listed in Table 2 and Additional file 2). The material of the KWS Saat

SE, Einbeck and IPK Gatersleben was transferred under the regulations of the standard

material transfer agreement (SMTA) of the International Treaty.

Apart from *B. vulgaris* subsp. *vulgaris*, 17 other *Beta* and *Patellifolia* accessions shown in

Figure 6 were analysed. The exact sampling location of the investigated accessions was

extracted from the GBIS/I (Genebank information system; IPK) (48) (Figure 6).

Table 2: Abbreviation, species name, accession number, genus, and section of the

investigated accessions of the Betoideae subfamily.

Abbreviation	Species name	Accession number	Genus	Section
Bmar1	*Beta vulgaris* subsp. *maritima*	BETA 1101	*Beta*	*Beta*
Bada	*Beta vulgaris* subsp. *adanensis*	BETA 1233	*Beta*	*Beta*
Bmar2	*Beta vulgaris* subsp. *maritima*	BETA 2322	*Beta*	*Beta*
Bvul	*Beta vulgaris* subsp. *vulgaris*	KWS 2320	*Beta*	*Beta*
Bptu	*Beta patula*	BETA 548	*Beta*	*Beta*
Bmca	*Beta macrocarpa*	BETA 881	*Beta*	*Beta*
Bnan1	*Beta nana*	BETA 546	*Beta*	*Corollinae*, formerly *Nanae*
Bnan2	*Beta nana*	BETA 570	*Beta*	*Corollinae*, formerly *Nanae*
Bint1	*Beta intermedia*	BETA 431	*Beta*	*Corollinae*
Bint2	*Beta intermedia*	BETA 923	*Beta*	*Corollinae*
Bcor1	*Beta corolliflora*	BETA 408	*Beta*	*Corollinae*
Bcor2	*Beta corolliflora*	BETA 408	*Beta*	*Corollinae*
Bmrh1	*Beta macrorhiza*	BETA 830	*Beta*	*Corollinae*
Bmrh2	*Beta macrorhiza*	BETA 576	*Beta*	*Corollinae*
Blom	*Beta lomatogona*	BETA 674	*Beta*	*Corollinae*
Pweb	*Patellifolia webbiana*	BETA 526	*Patellifolia*	-
Ppro	*Patellifolia procumbens*	BETA 951	*Patellifolia*	-
Ppat1	*Patellifolia patellaris*	BETA 534	*Patellifolia*	-
Ppat2	*Patellifolia patellaris*	BETA 892	*Patellifolia*	-

Figure 6: Geographic distribution of the *Beta* and *Patellifolia* species. The exact sampling locations of the investigated species are shown. The black lines represent the distribution area of the respective species and sections (see legend). The distribution areas of *B. vulgaris* subsp. *maritima*, *B. vulgaris* subsp. *adanensis*, and *B. macrocarpa* are not shown as these species occur along the whole coastline of Western Eurasia (1,10,11,49).

The plants were grown under long day conditions in a greenhouse and were obtained and grown in accordance with German legislation. Genomic DNA was isolated from young leaves using the NucleoSpin ® Plant II protocols from Macherey & Nagel. Each high-quality gDNA (200 ng) was fragmented by sonication using a Bioruptor (Fa. Diagenode) and subsequently used for library preparation with the TruSeq Nano DNA library preparation kit (Fa. Illumina). End repaired fragments were size selected by AmpureXp Beads (Fa. Beckmann-Coulther) to an average size of around 700 bp. After end repair, A-tailing and ligation of barcoded adapters, fragments were enriched by eight cycles of PCR. The final libraries were quantified using PicoGreen (Fa. Quant-iT) on a FLUOstar plate reader (Fa. BMG labtech) and quality checked by HS-Chip on a 2100 Bioanalyzer (Fa. Agilent Technologies). Before sequencing all libraries were pooled depending on the genome size and ploidy of each accession and sequenced 2 x 250 nt on a HiSeq1500 in rapid mode over two lanes using onboard cluster generation. Processing and demultiplexing of raw data was performed by bcl2fastq-v2.19.1 to generate FASTQ files for each accession.

Plastome assemblies and annotation

Trimmomatic (v0.39) (50) was applied to remove adapter sequences
(ILLUMINACLIP:adapters.fa:2:30:10:2:keepBothReads) and to ensure high quality of the
reads (SLIDINGWINDOW:4:15 MINLEN:50 TOPHRED33). FastQC (v0.11.9) (51) was used
for quality checks. The trimmed reads were subjected to GetOrganelle (v1.7.0) (52) to
generate plastome assemblies as suggested for Embryophyta plant plastome sequences (-
R 15; -F embplant_pt). The SPAdes (53) kmer settings were set to -k 21, 45, 65, 85, or 105.
The contig coverage information and other graph characteristics are used by GetOrganelle
to construct the final assembly graphs, which were plotted and visually assessed using
Bandage (v0.8.1) (54). The assemblies suggested a circular sequence, however, circular
plastome molecules might only comprise a small proportion of all molecules in the cell,
whereas other plastome molecules may occur in branched or linear configurations (55–57).
The assemblies were submitted in the FASTA format, retaining the possibility to reuse the
submitted assemblies as circular or linear sequences. The complex assembly graph of
Bmar1 was not automatically resolved. Therefore, single contigs of Bmar1 were sorted
manually based on the structure of the other assemblies to enable comparative analyses as
described in the following section.

Structural annotation of all plastome assemblies was performed with GeSeq (v2.01) (58).
The BLAT (59) search parameters 'Annotate plastid trans-spliced rps12' and 'Ignore genes
annotated as locus tags' were used together with a 'Protein search identity' of 25 and a
'rRNA, tRNA, DNA search identity' of 85. For HMMER profile search 'Embryophyta
chloroplast (CDS+rRNA)' was selected and 'MPI-MP chloroplast references (Embryophyta
CDS + rRNA)' was chosen as reference. The resulting annotation files in the gff format were
directly used for further analyses. To avoid confusion, we want to make aware of the fact
that *psbN* and *pbf1* are two different names for the same gene.

Construction of phylogenetic trees

The workflow for the alignment-based phylogenetic analysis is available in Additional file
S1G. The position of each gene was extracted from the GFF files obtained through GeSeq.
Next, adjacent genes with conserved microsynteny across all investigated samples
(including amaranth, quinoa, and spinach as outgroup) were identified and the interleaved

intergenic regions of these neighbouring genes were extracted. For overlapping genes, the
extraction of an intergenic region was not possible.

Using the gene sequences and intergenic regions of all samples, gene/region specific
alignments were performed using MAFFT (v7.299b) (60). High accuracy was ensured using
the L-INS-I method. To align sequences with different orientations, the parameter '--
adjustdirection' was used. The alignments were trimmed using trimAl (v1.4.rev22) (61). The
gap threshold was set to '-gt 0.8', whereas the threshold for the minimum average similarity
was set to '-st 0.001'. Then, the single alignments were concatenated and the resulting
alignment matrix was inspected using SeaView (62). Manual adjustment was not
necessary.

RAxML-NG (v1.0.0) (63) was used for ML analysis together with bootstrapping (Model:
GTR+FC+G8m). The substitution matrix GTR (for DNA) was applied together with the
model parameter 'G8' and 'F'. The parsimony-based randomised stepwise addition
algorithm was selected for the starting tree (--tree pars{10}). The number of replicate trees
for bootstrapping was set to 200. The resulting tree was visualised using FigTree (v1.4.4)
(64).

Location-based clustering of the clades in the tree was performed manually based on the
sampling locations (Additional file S2C).

To identify a reduced set of gene sequences and intergenic regions for the construction of a
phylogenetic tree distinguishing all accessions, the sequences were iteratively added with
increasing alignment identities until all species were separated by informative positions.

To investigate the region dependent phylogeny different additional data matrices were
constructed for (I) all gene regions, (II) all intergenic regions, (III) complete coding
sequences, (IV) first and second codon position and (V) third codon position only.
Therefore, coding sequences for all 79 protein coding genes were extracted from the
Genbank annotation files of our plastome assemblies. Start and stop codons were removed
and extracted sequences were processed as described above for ML analysis.

To extract the 18s rRNA gene sequence from *B. corolliflora* (Bcor1) as representative of the
Corollinae, SOAPdenovo2 assemblies were generated using the trimmed reads as input.
SOAPdenovo2 (v2.04) (65) was tested with different kmer sizes ranging from 67-127 in

steps of 10. The resulting assembly with the highest N50 length was used for further
investigations. The reference 18s rRNA gene sequence for *B. vulgaris* subsp. *vulgaris* was
retrieved from the NCBI (GeneID=809573). The 18s rRNA gene sequence for Bcor1 was
identified via BLAST and then extracted from the SOAPdenovo2 assembly, consecutively
adding the following overlapping BLAST hit with the smallest e-value. Next, these extracted
sequences were combined for a 18s gene sequence reconstruction. The assembled 18s
rRNA gene sequence and the corresponding reference gene sequence were aligned via
MAFFT and inspected using SeaView.

Mitogenome and plastome phylogeny based on kmer-derived phylogenetic splits

SANS serif (v2.1_04B) (66,67), a method based on colored de Bruijn graphs, was selected
for the reconstruction of additional phylogenies using variable input data (mitochondrial
reads, chloroplastic reads, and full plastome assemblies). This method does not require
prior assembly of the reads and is therefore especially suitable for the mitochondrial
sequences which could not be fully assembled using GetOrganelle due to the relatively low
available sequencing depth and also higher complexity of the mitogenome in comparison to
the plastome.

Reads were assigned to the plastome or the mitogenome, respectively, after mapping with
BWA-MEM (v0.7.13) (68) against the sugar beet reference genome sequence, including the
respective sugar beet chloroplast (KR230391.1) and sugar beet mitochondrial sequence
(BA000009.3), which were published independently from this study. This enabled the
extraction of reads mapping with higher confidence to the chloroplast/mitochondrial
sequence in contrast to mapping to the nucleome (with e.g. a few mismatches) and *vice*
versa. Therefore, 'samtools view', with the -b and -h options, was used after indexing the
BAM file. The resulting BAM file was then further processed using 'samtools collate' and
converted to the FASTQ format to extract the corresponding reads (chloroplast or
mitochondria, respectively) using 'samtools fastq'. After that, for the read-derived
phylogenetic analyses, a colored de Bruijn graph was constructed using Bifrost (v1.0.5) (69)
to filter kmers which only occur once in the dataset. This graph was then used as input for
SANS serif. In addition, a phylogeny was reconstructed using the newly constructed
plastome assemblies as direct input for SANS serif.

Different parameters were applied to test the robustness of the results. These arguments
include different mean weight functions (-m; geom vs. geom2), the number of splits in the
output list (-t; all vs. 10n) as well as various kmer sizes (11, 21, 31 and 61). The SANS serif
output file was then converted to the nexus format (sans2nexus.py) and subsequently
visualised using Splitstree5 (70). To analyse the discrepancies between trees derived from
different methods, SANS serif was used with the option 'strict' to generate an output file in
the newick format which was then visualised using FigTree (64). Further, the SANS serif
script 'comp.py' was used to calculate weighted (length of the edges/size of the splits is
taken into account) precision and recall (combined in F1 scores) while using each tree as
reference/ground truth in an 'all vs. all' comparison. In this use case, precision means 'the
total weight of all correctly predicted splits divided by the total weight of all predicted splits'.
Further, weighted symmetric set distances and Robinsons-Foulds distances were
calculated for each comparison. Details can be found in Additional file S1D. For this
analysis, the trees constructed with different input data (cp_reads, mt_reads, and
cp_assemblies) and the fixed parameters '-m geom2, -t 10n, -k 31' were compared.

Investigation of alignment identities

The alignment identities for each plastome gene sequence or intergenic region were
calculated to infer the phylogenetic information of all sequences. The events (SNV or InDel)
were detected by iteration over each position in the sequence. The identity score (percent
identity) was calculated by division of conserved positions (number of residues [position in
alignment] with the same nucleotide in all accessions) by the number of residues in the
alignment ('sequence length'). Alignment identities were calculated (i) for all accessions and
(ii) for all accessions including outgroup reference plastome sequences (amaranth, quinoa,
and spinach). Visualisation of the results was performed using matplotlib (71). Next,
potential hotspots for SNVs and InDels in the plastome sequences were investigated.

Declarations

Ethics approval and consent to participate

The material of the KWS Saat SE, Einbeck, and IPK Gatersleben was transferred under the
regulations of the standard material transfer agreement (SMTA) of the International Treaty.
Plants were grown in accordance with German legislation

Consent for publication

631 Not applicable.

Availability of data and materials

634 Sequence reads have been submitted to the European Nucleotide Archive (ENA; Additional
635 file S2D). The plastome assemblies and the corresponding annotations are available at
636 ENA/GenBank (PRJEB45680). Additional information used in phylogenetic analyses are
637 included in the supplementary files.

Competing interests

639 The authors declare that the research was conducted in the absence of any commercial or
640 financial relationships that could be construed as a potential conflict of interest.

Funding

642 Open Access funding enabled and organized by Projekt DEAL. KS is funded by Bielefeld
643 University through the Graduate School DILS (Digital Infrastructure for the Life Sciences).

Authors' contributions

645 BP, BW, TH and DH designed the study. NS selected and cultivated the plants and
646 performed DNA extraction. PV designed the layout for sequencing. BP and KS developed
647 and implemented the bioinformatic methodology. KS analysed the data and prepared the
648 figures and tables. KS, BP, NS and TH wrote the manuscript. All authors read and approved
649 the final manuscript.

Acknowledgements

651 We acknowledge support for the Article Processing Charge by the Deutsche
652 Forschungsgemeinschaft (German Research Foundation) and the Open Access Publication

653 Fund of Bielefeld University. We thank the CeBiTec Bioinformatic Resource Facility team for
654 great technical support and Dr. Roland Wittler for great support with the SANS serif
655 software.

References

1. Romeiras MM, Vieira A, Silva DN, Moura M, Santos-Guerra A, Batista D, et al. Evolutionary
and Biogeographic Insights on the Macaronesian Beta-Patellifolia Species (Amaranthaceae)
from a Time-Scaled Molecular Phylogeny. Robillard T, editor. PLoS ONE. 2016 Mar
31;11(3):e0152456.

2. Fischer HE. Origin of the 'Weisse Schlesische Rübe' (white Silesian beet) and resynthesis of
sugar beet. Euphytica. 1989 Apr;41(1–2):75–80.

3. Panella L, Lewellen RT. Broadening the genetic base of sugar beet: introgression from wild
relatives. Euphytica. 2007 Mar 7;154(3):383–400.

4. Biancardi E, Lewellen RT. History and Current Importance. In: Biancardi E, Panella LW,
McGrath JM, editors. Beta maritima [Internet]. Cham: Springer International Publishing; 2020
[cited 2021 Jul 28]. p. 1–48. Available from: http://link.springer.com/10.1007/978-3-030-
28748-1_1

5. Capistrano-Gossmann GG, Ries D, Holtgräwe D, Minoche A, Kraft T, Frerichmann SLM, et al.
Crop wild relative populations of Beta vulgaris allow direct mapping of agronomically
important genes. Nat Commun. 2017 Aug;8(1):15708.

6. Rodríguez del Río Á, Minoche AE, Zwickl NF, Friedrich A, Liedtke S, Schmidt T, et al.
Genomes of the wild beets *Beta patula* and *Beta vulgaris* ssp. *maritima*. Plant J. 2019
Sep;99(6):1242–53.

7. Frese L, Ford-Lloyd B. Taxonomy, Phylogeny, and the Genepool. In: Biancardi E, Panella LW,
McGrath JM, editors. Beta maritima [Internet]. Cham: Springer International Publishing; 2020
[cited 2021 Jul 28]. p. 121–51. Available from: http://link.springer.com/10.1007/978-3-030-
28748-1_6

8. Kadereit G, Hohmann S, Kadereit JW. A Synopsis of Chenopodiaceae Subfam. Betoideae and
Notes on the Taxonomy of Beta. Willldenowia. 2006 Apr 20;Bd. 36, H. 1(Special Issue:
Festschrift Werner Greuter):9–19.

9. Ford-Lloyd BV, Williams JT. A revision of Beta section Vulgares (Chenopodiaceae), with new
light on the origin of cultivated beets. Botanical Journal of the Linnean Society. 1975
Sep;71(2):89–102.

10. Hohmann S, Kadereit JW, Kadereit G. Understanding Mediterranean-Californian disjunctions:
molecular evidence from Chenopodiaceae-Betoideae. Taxon. 2006 Feb;55(1):67–78.

11. Touzet P, Villain S, Buret L, Martin H, Holl A-C, Poux C, et al. Chloroplastic and nuclear
diversity of wild beets at a large geographical scale: Insights into the evolutionary history of the
Beta section. Ecol Evol. 2018 Mar;8(5):2890–900.

12. Frese L, Nachtigall M, Iriondo JM, Rubio Teso ML, Duarte MC, Pinheiro de Carvalho MÂA.
Genetic diversity and differentiation in Patellifolia (Amaranthaceae) in the Macaronesian
archipelagos and the Iberian Peninsula and implications for genetic conservation programmes.
Genet Resour Crop Evol. 2019 Jan;66(1):225–41.

13. Shen Y, Ford-lloyd BV, Newbury HJ. Genetic relationships within the genus Beta determined
using both PCR-based marker and DNA sequencing techniques. Heredity. 1998
May;80(5):624–32.

14. Gitzendanner MA, Soltis PS, Yi T-S, Li D-Z, Soltis DE. Plastome Phylogenetics: 30 Years of
Inferences Into Plant Evolution. In: Advances in Botanical Research [Internet]. Elsevier; 2018
[cited 2021 Jul 28]. p. 293–313. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0065229617300885

15. Liu B-B, Ma Z-Y, Ren C, Hodel RGJ, Sun M, Liu X-Q, et al. Capturing single-copy nuclear
genes, organellar genomes, and nuclear ribosomal DNA from deep genome skimming data for
plant phylogenetics: A case study in Vitaceae [Internet]. Evolutionary Biology; 2021 Feb [cited
2021 Jul 28]. Available from: http://biorxiv.org/lookup/doi/10.1101/2021.02.25.432805

16. Palmer JD, Zamir D. Chloroplast DNA evolution and phylogenetic relationships in
Lycopersicon. Proceedings of the National Academy of Sciences. 1982 Aug 1;79(16):5006–10.

17. Givnish TJ, Spalink D, Ames M, Lyon SP, Hunter SJ, Zuluaga A, et al. Orchid phylogenomics
and multiple drivers of their extraordinary diversification. Proc R Soc B. 2015 Sep
7;282(1814):20151553.

18. Orton LM, Burke SV, Duvall MR. Plastome phylogenomics and characterization of rare
genomic changes as taxonomic markers in plastome groups 1 and 2 Poeae (Pooideae; Poaceae).
PeerJ. 2019 Jun 3;7:e6959.

19. Guo X, Liu J, Hao G, Zhang L, Mao K, Wang X, et al. Plastome phylogeny and early
diversification of Brassicaceae. BMC Genomics. 2017 Dec;18(1):176.

20. Singh BP, Kumar A, Kaur H, Singh H, Nagpal AK. CpGDB : A Comprehensive Database of
Chloroplast Genomes. Bioinformation. 2020 Feb 29;16(2):171–5.

21. Wang M, Wang X, Sun J, Wang Y, Ge Y, Dong W, et al. Phylogenomic and evolutionary
dynamics of inverted repeats across Angelica plastomes. BMC Plant Biol. 2021 Dec;21(1):26.

22. Zurawski G, Clegg M. Evolution of higher-plant chloroplast DNA-encoded genes: implications
for structure-function and phylogenetic studies. Annual review of plant physiology.
1987;38:391–418.

23. Sugiura M. The chloroplast genome. Plant Molecular Biology (Netherlands). 1992;

24. Wang W, Lanfear R. Long-reads reveal that the chloroplast genome exists in two distinct
versions in most plants. Gaut B, editor. Genome Biology and Evolution. 2019 Nov 21;evz256.

25. Chen Y, Yang Z. Characterization of the complete plastome of *Dysphania botrys* , a candidate
plant for cancer treatment. Mitochondrial DNA Part B. 2018 Jul 3;3(2):1214–5.

26. Yao G, Jin J-J, Li H-T, Yang J-B, Mandala VS, Croley M, et al. Plastid phylogenomic insights
into the evolution of Caryophyllales. Molecular Phylogenetics and Evolution. 2019
May;134:74–86.

27. Stadermann KB, Weisshaar B, Holtgräwe D. SMRT sequencing only de novo assembly of the
sugar beet (Beta vulgaris) chloroplast genome. BMC Bioinformatics. 2015 Dec;16(1):295.

28. Williams LE, Wernegreen JJ. Sequence Context of Indel Mutations and Their Effect on Protein
Evolution in a Bacterial Endosymbiont. Genome Biology and Evolution. 2013 Mar;5(3):599–
605.

29. Gomes Pacheco T, Morais da Silva G, de Santana Lopes A, de Oliveira JD, Rogalski JM,
Balsanelli E, et al. Phylogenetic and evolutionary features of the plastome of Tropaeolum
pentaphyllum Lam. (Tropaeolaceae). Planta. 2020 Aug;252(2):17.

30. Serna-Sánchez MA, Pérez-Escobar OA, Bogarín D, Torres-Jimenez MF, Alvarez-Yela AC,
Arcila-Galvis JE, et al. Plastid phylogenomics resolves ambiguous relationships within the
orchid family and provides a solid timeframe for biogeography and macroevolution. Sci Rep.
2021 Dec;11(1):6858.

31. Wang J-H, Moore MJ, Wang H, Zhu Z-X, Wang H-F. Plastome evolution and phylogenetic
relationships among Malvaceae subfamilies. Gene. 2021 Jan;765:145103.

32. de Santana Lopes A, Pacheco TG, Santos KG dos, Vieira L do N, Guerra MP, Nodari RO, et al.
The Linum usitatissimum L. plastome reveals atypical structural evolution, new editing sites,
and the phylogenetic position of Linaceae within Malpighiales. Plant Cell Rep. 2018
Feb;37(2):307–28.

33. Qiu T, Cui S. Evolutionary analysis for Phragmites ecotypes based on full-length plastomes.
Aquatic Botany. 2021 Mar;170:103349.

34. Igea J, Juste J, Castresana J. Novel intron markers to study the phylogeny of closely related
mammalian species. BMC Evol Biol. 2010;10(1):369.

35. Palmer JD, Herbon LA. Plant mitochondrial DNA evolved rapidly in structure, but slowly in
sequence. Journal of Molecular evolution. 1988;28(1):87–97.

36. Heckenhauer J, Paun O, Chase MW, Ashton PS, Kamariah AS, Samuel R. Molecular
phylogenomics of the tribe Shoreeae (Dipterocarpaceae) using whole plastid genomes. Annals
of Botany. 2019 May 20;123(5):857–65.

37. Olmstead RG, Bedoya AM. Whole genomes: the holy grail. A commentary on: 'Molecular
phylogenomics of the tribe Shoreeae (Dipterocarpaceae) using whole plastidgenomes'. Annals
of Botany. 2019 May 20;123(5):iv–v.

38. Kim Y-K, Jo S, Cheon S-H, Joo M-J, Hong J-R, Kwak M, et al. Plastome Evolution and
Phylogeny of Orchidaceae, With 24 New Sequences. Front Plant Sci. 2020 Feb 21;11:22.

39. Walker JF, Walker-Hale N, Vargas OM, Larson DA, Stull GW. Characterizing gene tree
conflict in plastome-inferred phylogenies. PeerJ. 2019 Sep 24;7:e7747.

40. Coons GH. The wild species of Beta. Proc Am Soc Sugar Beet Technol. 1954;8(2).

41. Biancardi E, de Biaggi M. Morphology. In: Biancardi E, Panella LW, McGrath JM, editors.
Beta maritima [Internet]. Cham: Springer International Publishing; 2020 [cited 2021 Jul 29]. p.
61–86. Available from: http://link.springer.com/10.1007/978-3-030-28748-1_3

42. Frese L, de Carvalho MAP, Duarte C. Crop case study Beta L.(including Patellifolia AJ Scott et
al.). AEGRO project. Julius Kühn-Institut, Bundesforschungsinstitut für Kulturpflanzen, Institut
für Züchtungsforschung an landwirtschaftlichen Kulturen; 2011.

43. Gao D, Schmidt T, Jung C. Molecular characterization and chromosomal distribution of
species-specific repetitive DNA sequences from *Beta corolliflora* , a wild relative of sugar beet.
Genome. 2000 Dec 1;43(6):1073–80.

44. Heitkam T, Holtgräwe D, Dohm JC, Minoche AE, Himmelbauer H, Weisshaar B, et al.
Profiling of extensively diversified plant LINEs reveals distinct plant-specific subclades. Plant
J. 2014 Aug;79(3):385–97.

45. Maiwald S, Weber B, Seibt KM, Schmidt T, Heitkam T. The Cassandra retrotransposon
landscape in sugar beet (*Beta vulgaris*) and related Amaranthaceae: recombination and re-
shuffling lead to a high structural variability. Annals of Botany. 2021 Jan 1;127(1):91–109.

46. Weber B, Wenke T, Frömmel U, Schmidt T, Heitkam T. The Ty1-copia families SALIRE and
Cotzilla populating the Beta vulgaris genome show remarkable differences in abundance,
chromosomal distribution, and age. Chromosome Res. 2010 Feb;18(2):247–63.

47. Panella LW, Stevanato P, Pavli O, Skaracis G. Source of Useful Traits. In: Biancardi E, Panella
LW, McGrath JM, editors. Beta maritima [Internet]. Cham: Springer International Publishing;
2020 [cited 2021 Jul 29]. p. 167–218. Available from: http://link.springer.com/10.1007/978-3-
030-28748-1_8

48. Oppermann M, Weise S, Dittmann C, Knüpffer H. GBIS: the information system of the German
Genebank. Database [Internet]. 2015 Jan 1 [cited 2021 Jul 28];2015. Available from:
https://academic.oup.com/database/article/doi/10.1093/database/bav021/2433153

49. Castro S, Romeiras MM, Castro M, Duarte MC, Loureiro J. Hidden diversity in wild Beta taxa
from Portugal: Insights from genome size and ploidy level estimations using flow cytometry.
Plant Science. 2013 Jun;207:72–8.

50. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014;30(15):2114–20.

51. Andrews S. FastQC: a quality control tool for high throughput sequence data. [Internet]. 2020.
Available from: http://www.bioinformatics.babraham.ac.uk/projects/fastqc

52.   Jin J-J, Yu W-B, Yang J-B, Song Y, dePamphilis CW, Yi T-S, et al. GetOrganelle: a fast and
799   versatile toolkit for accurate de novo assembly of organelle genomes. Genome Biol. 2020
800   Dec;21(1):241.

801   53.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: A
802   New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. Journal of
803   Computational Biology. 2012 May;19(5):455–77.

804   54.   Wick RR, Schultz MB, Zobel J, Holt KE. Bandage: interactive visualization of *de novo* genome
805   assemblies: Fig. 1. Bioinformatics. 2015 Oct 15;31(20):3350–2.

806   55.   Oldenburg DJ, Bendich AJ. Most Chloroplast DNA of Maize Seedlings in Linear Molecules
807   with Defined Ends and Branched Forms. Journal of Molecular Biology. 2004 Jan;335(4):953–
808   70.

809   56.   Oldenburg DJ, Bendich AJ. DNA maintenance in plastids and mitochondria of plants. Front
810   Plant Sci [Internet]. 2015 Oct 29 [cited 2021 Jul 28];6. Available from:
811   http://journal.frontiersin.org/Article/10.3389/fpls.2015.00883/abstract

812   57.   Shaver JM, Oldenburg DJ, Bendich AJ. The Structure of Chloroplast DNA Molecules and the
813   Effects of Light on the Amount of Chloroplast DNA during Development in *Medicago
814   truncatula*. Plant Physiology. 2008 Mar 3;146(3):1064–74.

815   58.   Tillich M, Lehwark P, Pellizzer T, Ulbricht-Jones ES, Fischer A, Bock R, et al. GeSeq –
816   versatile and accurate annotation of organelle genomes. Nucleic Acids Research. 2017 Jul
817   3;45(W1):W6–11.

818   59.   Kent WJ. BLAT---The BLAST-Like Alignment Tool. Genome Research. 2002 Mar
819   20;12(4):656–64.

820   60.   Katoh K. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier
821   transform. Nucleic Acids Research. 2002 Jul 15;30(14):3059–66.

822   61.   Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. trimAl: a tool for automated alignment
823   trimming in large-scale phylogenetic analyses. Bioinformatics. 2009 Aug 1;25(15):1972–3.

824   62.   Gouy M, Guindon S, Gascuel O. SeaView Version 4: A Multiplatform Graphical User Interface
825   for Sequence Alignment and Phylogenetic Tree Building. Molecular Biology and Evolution.
826   2010 Feb 1;27(2):221–4.

827   63.   Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. RAxML-NG: a fast, scalable and
828   user-friendly tool for maximum likelihood phylogenetic inference. Wren J, editor.
829   Bioinformatics. 2019 Nov 1;35(21):4453–5.

830   64.   Rambaut A. FigTree [Internet]. 2009. Available from:
831   http://evomics.org/resources/software/molecular-evolution-software/figtree/

832   65.   Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, et al. Erratum: SOAPdenovo2: an empirically
833   improved memory-efficient short-read de novo assembler. GigaSci. 2015 Dec;4(1):30.

834   66.   Wittler R. Alignment- and reference-free phylogenomics with colored de Bruijn graphs.
835   Algorithms Mol Biol. 2020 Dec;15(1):4.

836   67.   Rempel A, Wittler R. SANS serif: alignment-free, whole-genome-based phylogenetic
837   reconstruction. Schwartz R, editor. Bioinformatics. 2021 Jun 16;btab444.

838   68.   Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
839   arXiv:13033997 [q-bio] [Internet]. 2013 May 26 [cited 2021 Jul 28]; Available from:
840   http://arxiv.org/abs/1303.3997

841   69.   Holley G, Melsted P. Bifrost: highly parallel construction and indexing of colored and
842   compacted de Bruijn graphs. Genome Biol. 2020 Dec;21(1):249.

843   70.   Huson DH, Bryant D. Application of Phylogenetic Networks in Evolutionary Studies.
844   Molecular Biology and Evolution. 2006 Feb 1;23(2):254–67.

71.   Hunter JD. Matplotlib: A 2D Graphics Environment. Comput Sci Eng. 2007;9(3):90–5.

Supplementary Material

Additional file S1A: Geographic distribution of the Betoideae species as described in the
literature.

Additional file S1B: Distribution of coverage values and assembly length (in bp) for each
region and the total assemblies.

Additional file S1C: Circular and linear plots of selected plastome assembly sequences.

Additional file S1D: Distance metrics for the comparison of splitstree results.

Additional file S1E: Reduced phylogenetic tree based on 53 gene and intergenic regions.

Additional file S1F: Phylogenetic trees based on different sequence matrices.

Additional file S1G: Workflow for the construction of phylogenetic trees.

Additional file S2A: Sequence identities [%] of all investigated plastome gene sequences
and intergenic regions including outgroup reference sequences.

Additional file S2B: Sequence identities [%] of all investigated plastome gene sequences
and intergenic regions excluding outgroup reference sequences.

Additional file S2C: Accession IDs, taxonomy, read and assembly statistics and geographic
location of the investigated accessions of the Betoideae subfamily.

Additional file S2D: SRA-IDs of the processed read datasets.