1   **Low heritability and high phenotypic plasticity of salivary cortisol in response to**

2   **environmental heterogeneity in a wild pinniped**

4   Rebecca Nagel[1], Sylvia Kaiser[2], Claire Stainfield[3], Camille Toscani[3], Cameron Fox-
5   Clarke[3], Anneke J. Paijmans[1], Camila Costa Castro[1], David L. J. Vendrami[1], Jaume
6   Forcada[3,*], Joseph I. Hoffman [1,3,*]
         1. Department of Animal Behaviour, Bielefeld University, 33501 Bielefeld,
7   Germany
8
         2. Department of Behavioural Biology, University of Münster, 48149 Münster,
9   Germany
10
         3. British Antarctic Survey, Cambridge CB3 OET, UK
11   * Joint senior authors
12

13   Corresponding author: Rebecca Nagel, Department of Animal Behaviour, Bielefeld
14   University, 33501 Bielefeld, Germany, rebecca.nagel@uni-bielefeld.de
15
16
17

Abstract

Individuals are unique in how they interact with and respond to their

environment. Correspondingly, unpredictable challenges or environmental stressors

often produce an individualized response of the hypothalamic-pituitary-adrenal axis

and its downstream effector cortisol. We used a fully crossed, repeated measures

design to investigate the factors shaping individual variation in baseline cortisol in

Antarctic fur seal pups and their mothers. Saliva samples were collected from focal

individuals at two breeding colonies, one with low and the other with high density,

during two consecutive years of contrasting food availability. Mothers and pups were

sampled concurrently at birth and shortly before weaning, while pups were

additionally sampled every 20 days. We found that heritability was low for baseline

cortisol, while within-individual repeatability and among-individual variability were

high. A substantial proportion of the variation in baseline cortisol could be explained

in pups and mothers by a combination of intrinsic and extrinsic factors including sex,

weight, day, season, and colony of birth. Our findings provide detailed insights into

the individualization of endocrine phenotypes and their genetic and environmental

drivers in a wild pinniped. Furthermore, the strong associations between cortisol and

life history traits that we report in fur seals could have important implications for

understanding the population dynamics of species impacted by environmental change.

Keywords

Introduction

Despite similarities in age, sex, or social status, individuals often differ in how they interact with their environment (Réale and Dingemanse 2010; Dall et al. 2012). Once perceived as a statistical nuisance, an ever growing body of evidence now suggests that these individual differences are often consistent and stable across time and contexts, with profound implications for understanding phenotypic variation, niche specialization, and animal personality (Sih et al. 2012; Wolf and Weissing 2012). The past decade has thus witnessed an increased awareness of individualization as a fundamentally important and compelling aspect of evolutionary biology, ecology, and animal behavior (Bolnick et al. 2003; Trillmich et al. 2018; Krüger et al. 2021).

Consistent differences among individuals are likely to be mediated by a combination of intrinsic and extrinsic factors and can be understood as the interaction between an individual's phenotype, genotype, and its ecological context such that its fitness is maximized (Dingemanse and Réale 2005; Vessey and Drickamer 2010; Lihoreau et al. 2021). At the proximate level, individualized phenotypic adjustments to environmental factors may be governed by, among other things, variation in the concentrations of circulating hormones (Müller et al. 2020). In particular, individual variation in cortisol, an important hormone in the physiological stress response, appears to play a major role in shaping individual responses to environmental conditions (Wingfield and Romero 2011).

Cortisol is a steroid hormone that belongs to the class of glucocorticoids. Its release is regulated by the hypothalamic-pituitary-adrenal (HPA) axis. Glucocorticoids play an essential role in maintaining metabolic and homeostatic functions (Kuo et al. 2015). Under predictable conditions, cortisol is released continuously at baseline levels that vary naturally throughout the day and over an individual's lifetime (Lightman and Conway-Campbell 2010). In the face of unpredictable challenges, however, activation of the HPA axis results in increased levels of secreted cortisol for the duration of the stressor, before levels return to baseline (Bellavance and Rivest 2014).

While the physiology of the HPA axis is largely conserved across mammals (Romero and Butler 2007; Wingfield and Romero 2011), empirical studies have

documented significant differences in cortisol concentrations among species and
individuals (Schoenemann and Bonier 2018; Taff et al. 2018), as well as notable
variation in the HPA axis response across time and space (Romero and Gormally
2019). Such phenotypic variation in response to intrinsic and environmental
differences could potentially facilitate adaptation to different habitats and conditions
(Sih et al. 2012). For example, cortisol levels have been shown to vary significantly
among individuals experiencing different densities (Meise et al. 2016) and levels of
nutritional stress (Kitaysky et al. 2007). Variation in cortisol among individuals has
also been linked to intrinsic factors such as age (Pavitt et al. 2015), sex (Azevedo et
al. 2019), and weight (Jeanniard du Dot et al. 2009). Nonetheless, when such factors
are accounted for, empirical studies have shown that cortisol levels are often highly
repeatable within individuals (Schoenemann and Bonier 2018; Taff et al. 2018). In
guinea pigs, for example, cortisol responsiveness and to a lesser extent baseline
cortisol levels are highly repeatable from late adolescence to adulthood (Mutwill et al.
2021).

Given that cortisol levels often exhibit high among-individual variability and
within-individual repeatability, it has been argued that genetic rather than
environmental factors could explain much of the observed phenotypic variance, which
would imply a high evolvability of this endocrine trait (Boake 1989; Jenkins et al.
2014). Correspondingly, several empirical studies have reported moderate to high
levels of cortisol heritability in free-living vertebrate populations (Jenkins et al. 2014;
Stedman et al. 2017; Bairos-Novak et al. 2018). However, persistent intrinsic and / or
extrinsic factors might also produce repeatable phenotypes regardless of the
underlying genotype (Taff et al. 2018), a possibility that has often been overlooked in
the literature (Bonier and Martin 2016). This is particularly true when genetically
similar individuals experience a similar temporal and spatial environment leading to,
for example, phenotypic similarity in the endocrine phenotype in response to current
internal and environmental stimuli (contextual plasticity) or stimuli encountered in the
past (developmental plasticity) (Stamps and Biro 2016). If unaccounted for, such
phenotype-environment correlations may upwardly bias the estimated additive genetic
variance for phenotypic traits (Kruuk et al. 2003).

Pinnipeds, and otariids in particular, are ideally suited to investigate the effects

of internal and environmental factors on cortisol levels. First, otariids are colonially

breeding, with males competing to establish and maintain harems on densely packed

breeding beaches (Forcada and Staniland 2018). Cortisol may play an important role

in how individuals adapt to this dynamic environment by restoring homeostasis after

unpredictable challenges such as territorial bouts or unwanted mating attempts.

Second, while many breeding beaches do not differ appreciably in qualities such as

substrate type or topology, the density of individuals often varies from one place to

another, setting up a spatial dynamic that tends to remain stable over time (Cassini

1999). Consequently, as pups are born on land and remain ashore throughout much of

their early ontogeny (Payne 1979; McCafferty et al. 1998), cortisol might play an

important role in mediating individual responses to variation in density. Finally,

cortisol levels have been investigated in several pinniped species in relation to

ontogeny (Ortiz et al. 2003; Atkinson et al. 2011), environmental conditions

(DeRango et al. 2019), and handling regimes (Engelhard et al. 2002; Harcourt et al.

2010; Bennett et al. 2012; Champagne et al. 2012). Methodologies for collecting and

assessing cortisol in pinnipeds are therefore well established in the literature.

Our model otariid species, the Antarctic fur seal (*Arctocephalus gazella*), has

been extensively studied by the British Antarctic Survey (BAS) on Bird Island, South

Georgia since the 1980s. Two breeding colonies on the island provide a unique

"natural experiment" for investigating individual responses to population density.

Freshwater Beach (FWB) and Special Study Beach (SSB) are situated less than 200

meters apart (Figure 1a), meaning they are exposed to comparable climatic

conditions. Breeding females from both locations also likely forage in the same areas

(Hunt et al. 1992) and do not differ significantly in quality traits such as body size and

condition (Nagel et al. 2021). Despite these similarities, the two colonies differ in the

density of conspecifics. Direct counts of individuals ashore suggest that the density of

breeding females is almost four times higher at SSB than FWB (Meise et al. 2016)

and the modal local density of focal pups across the entire breeding season is also

higher for pups born at SSB (Nagel et al. 2021).

We took advantage of this unique natural setup to investigate the intrinsic and

extrinsic factors shaping individual variation in baseline cortisol in Antarctic fur seal

pups and their mothers. We used a fully crossed, repeated measures design (Figure 1b,
c) comprising longitudinal data from mother-offspring pairs from the two colonies
over two consecutive breeding seasons, the first of which was coincidentally one of
the worst years on record with respect to food availability (Nagel et al. 2021).
Specifically, we collected saliva samples and accompanying biometric data from 25
randomly selected focal pairs from each colony in both seasons. To quantify baseline
cortisol levels, saliva was collected immediately, within three minutes of capture.
Mothers were sampled twice during the breeding season, while pups were sampled
every 20 days from birth until just before molting at around 60 days of age.

We used animal models to obtain heritability estimates for baseline cortisol
using both a simple pedigree and a genomic relatedness matrix obtained from a high
density single nucleotide polymorphism (SNP) array (Humble et al. 2020). We then
used linear mixed models to evaluate the within- and among-individual variability of
cortisol levels in pups and their mothers. Included in each model as explanatory
variables were multiple intrinsic (sex, weight, body condition, and days after initial
sampling) and extrinsic (density and year) variables. In line with previous studies of
wild vertebrate populations, we hypothesized that cortisol levels would be heritable
and repeatable within individuals. We further hypothesized that baseline cortisol
would be higher in pups and mothers from the high-density colony and in the season
of low food availability.

Materials and Methods

Field study

This study was conducted during the Antarctic fur seal breeding seasons (December to March) of 2018–19 (hereafter 2019) and 2019–20 (hereafter 2020) at Bird Island, South Georgia (54°00'24.8□ S, 38°03'04.1□ W). Each season, we sampled 25 unique mother-pup pairs from two neighboring breeding colonies, one of low (FWB) and the other of high (SSB) density (Figure 1a). Sampling at both locations was randomized with respect to pup sex, resulting in a final sample size of 51 male and 49 female pups. Pup mortality was higher at FWB than SSB (32% *versus* 12%, respectively), and averaged 25.6% over the two colonies and seasons.

Each mother and her pup were captured concurrently on two separate occasions: 2–3 days postpartum (December) and again as the pups began to moult shortly before weaning (March). Pups were additionally recaptured every 20 days. For the capture, restraint, and sampling of individuals, we employed protocols that have been established and refined over 30 consecutive years of the BAS long-term monitoring and survey program. Briefly, adult females were captured with a noosing pole and held on a restraint board during processing. Pups were captured with a slip noose or by hand and were restrained by hand. After sampling, individuals were released as close to their capture site as possible and, when present, pups were reunited with their mothers.

At first sampling, focal individuals were fitted with VHF transmitters to the dorsal side of the neck between the shoulder blades with epoxy glue (pups: Sirtrack core marine glue-on V2G 152A; mothers: Sirtrack core marine glue-on V2G 154C). Transmitter signals were monitored throughout the season using a hand-held VHF receiver (AOR LTD., AR8200). Focal individuals were also given cattle ear tags (Dalton Supplies, Henley on Thames, UK) in the trailing edge of each foreflipper (Gentry and Holt 1982) for identification. Tissue plugs were collected and stored in 20% dimethyl sulfoxide (DMSO) saturated with salt at -20°C for subsequent genetic analysis.

At every capture, weight and length measurements were taken from which a scaled mass index was calculated according to (Peig and Green 2009). This condition metric serves as a reliable indicator of overall fitness as it has been correlated with,

among other things, offspring survival (Milenkaya et al. 2015; Gélin et al. 2016) and

mating success (Gastón and Vaira 2020). At every capture, the saliva sample was

taken within three minutes of capture to provide data on baseline cortisol levels

(Bozovic et al. 2013). Saliva was collected by rotating sterile cotton tip applicators

fitted in polypropylene tubes (ROTH, Art. No. XC10.2) in the cheek pouch and under

the tongue. Samples were centrifuged and stored at -20°C for subsequent cortisol

analysis.

Hormone quantification

Saliva samples were thawed and centrifuged for ten minutes to separate the

mucins. The clear supernatant was then used for the determination of cortisol

concentrations. Samples contaminated with blood (reddish supernatant) were

discarded ($n = 30$ in 2019 and $n = 31$ in 2020), as cortisol values are often falsely

elevated in such samples. Cortisol concentrations were determined in duplicate using

enzyme-linked immunosorbent assays (cortisol free in saliva DES6611, Demeditec

Diagnostics GmbH, Kiel, Germany). We calculated the average coefficient of

variation (CV) resulting from individual CVs for all duplicates in an assay. Mean

intra-assay CV for a total of 12 assays was 3.89%. All samples were determined in

duplicate and if CV was larger than 10%, determination of the sample was repeated.

Furthermore, two samples of different concentrations were run in duplicate on a total

of 12 plates to assess inter-assay variation, which was on average 4.36%. The

antibody showed the following cross-reactivities: cortisol 100%, 11-desoxycortisol

50%, corticosterone 6.2%, 11-desoxycorticosterone 2.6%, 17α-oh-progesterone 1.3%,

cortisone and prednisone < 1%, testosterone, estradiol, and androstendione < 0.1%.

For additional information on kit validation as per the linearity and recovery rate, see

the respective Supplementary Tables S1 and S2.

SNP genotyping and genomic relatedness matrix construction

For the 95 focal individuals sampled in 2019, we extracted total genomic

DNA from tissue samples using a standard chloroform-isoamylalcohol protocol (for a

description of the full protocol, see the Supplementary Materials). SNP genotyping

was performed on these samples using a custom Affymetrix SNP array as described

by (Humble et al. 2020). Quality control of the raw output data and genotyping were

implemented using the Axiom Analysis Suite (5.0.1.38, Affymetrix) based on

parameter thresholds set to their default values for diploid organisms. SNPs initially

classified as "off target variants" (OTV) were recovered using the "Run OTV caller"

function. Of the 85,359 SNPs tiled on the array, 77,873 were retained for further

analysis representing SNPs classified as "PolyHighResolution" (SNPs that passed all

of the Axiom Analysis Suite quality controls) and "NoMinorHomozygote" (SNPs that

passed all quality controls but no homozygote genotypes for the minor allele were

found). An additional 3,423 SNPs with minor allele frequencies below 0.01 and 2,096

SNPs that departed significantly from Hardy Weinberg equilibrium (HWE) were

removed using PLINK version 1.9 (Purcell et al. 2007). Departures from HWE were

identified based on an alpha level of 0.01 after implementing mid-p adjustment

(Graffelman and Moreno 2013). After filtering, a total of 72,354 SNPs were retained

and used to produce a genomic relatedness matrix using the --make-grm option in

GCTA version 1.93.1 (Yang et al. 2011).

Heritability of cortisol levels

To quantify the proportion of the total variance in baseline cortisol attributable

to genetic differences among individuals, we fitted two multivariate generalized linear

mixed models (GLMMs) in *MCMCglmm* (Hadfield 2010) with baseline cortisol as the

dependent variable and individual ID and relatedness as random effects. For the first

model, a simple pedigree (comprising mother-offspring pairs) was built for the full

dataset providing an estimate of maternal genetic effects on cortisol concentrations.

The second model incorporated the SNP relatedness matrix from individuals sampled

only in the first season (2019) and, although smaller in sample size, provides a less

biased estimate of genetic relatedness by including not only mother-offspring pairs

but also detecting, for example, half-siblings. We used weak but informative priors

(0.05 of the observed phenotypic variance) in both models. Markov chains were run

for 9,000,000 iterations and we retained every 8,500[th] value after removing 150,000

iterations of burn-in to generate posterior distributions of the random parameters. The

posterior distribution of the model intercept and autocorrelation were checked to

assess model fit. We obtained estimates of baseline cortisol heritability by dividing

the additive genetic variance by the total phenotypic variance ($h^2 = V_A/V_P$) for each

sample of the posterior distribution.

Estimating intrinsic and extrinsic factors influencing baseline cortisol

257   To determine whether an individual's weight, body condition, age, sex,

season, colony, and an interaction between season and colony explained a significant

proportion of the variation in baseline cortisol among pups, we fitted a GLMM with a

log-link gamma error distribution in *lme4* (Bates et al. 2015). A second GLMM with

maternal cortisol as the dependent variable was used to determine the proportion of

variation explained by an individual's weight, body condition, the number of days

postpartum, season, colony, and an interaction between season and colony. To

account for both structural and data multicollinearity among weight, condition, and

age (pups) / days postpartum (mothers), these variables were rescaled and centered by

subtracting the mean from all observed values. In preliminary analyses, we tested for

the presence of heterogeneous variance by allowing individual slopes to vary by age

(pups) or days postpartum (mothers). For both models, random intercepts were

included for each individual to account for repeated measures. The residuals of the

full models were inspected for linearity and equality of error variances (using plots of

residuals versus fits), normality (using Q – Q plots), and homogeneity of variance

(using Levene's test). A backward elimination based on the chi-squared statistic was

implemented to simplify the full models such that, in each iteration, the fixed effect

with the lowest chi-squared value was removed from the model until we only tested

the null model. The best fitting models were then taken to be those with the lowest

AIC values. We present only the best model from each analysis in the Results, but

model reduction and AIC scores for all models are available in the Supporting R

Markdown file. The statistical significance of fixed predictors was assessed using

Wald tests. We determined the marginal R^2 (variance explained by fixed effects) and

conditional R^2 (variance explained by fixed and random effects) according to

(Nakagawa and Schielzeth 2013).

Animal handling, ethics and permits

Sampling was carried out by the BAS under permits from the Government of

South Georgia and the South Sandwich Islands (Wildlife and Protected Areas

Ordinance (2011), RAP permit numbers 2018/024 and 2019/032). The samples were

imported into the UK under permits from the Department for Environment, Food, and

Rural Affairs (Animal Health Act, import license number ITIMP18.1397) and from

the Convention on International Trade in Endangered Species of Wild Fauna and

Flora (import numbers 578938/01-15 and 590196/01-18). All procedures used were

approved by the BAS Animal Welfare and Ethics Review Body (AWERB

applications 2018/1050 and 2019/1058).

Results

We used a fully crossed, repeated measures design incorporating saliva

samples from 96 unique pups and 93 unique mothers from two colonies of contrasting

density across two consecutive years of contrasting food availability (Figure 1).

Sample sizes were balanced between the colonies ($n = 93$ from FWB and $n = 96$ from

SSB) and seasons ($n = 95$ from 2019 and $n = 94$ from 2020). Each season, pups were

sampled every 20 days from birth until weaning, amounting to a total of 290 analyzed

saliva samples. Mothers were sampled twice each season, once shortly after birth and

again shortly before molting, which amounted to a total of 145 analyzed saliva

samples .

Cortisol heritability estimates

We estimated heritability of baseline cortisol using two animal models, the

first incorporating known pedigree relationships (i.e. mother-offspring pairs from both

years) and the second incorporating a SNP relatedness matrix, which was only

available for the first year of the study. Narrow-sense heritability (h^2) estimates from

both models were low with overlapping 95% credible intervals (pedigree model: $h^2 =$

0.013, 95% highest posterior density 0.004 – 0.045; SNP relatedness model: $h^2 =$

0.018, 95% highest posterior density 0.004 – 0.062) (Figure 2). The additive genetic

(V_A) and residual (V_R) variance estimates of the two models were also comparable

(pedigree model: 95% highest posterior density of $V_A = 0.1 – 1.6$ and $V_R = 30.6 –$

39.9; SNP relatedness model: 95% highest posterior density of $V_A = 0.1 – 2.3$ and V_R

$= 28.5 – 41.7$).

Intrinsic and extrinsic factors influencing baseline cortisol

The best supported model of pup baseline cortisol contained individual age (p

< 0.001), weight ($p < 0.001$), sex ($p = 0.004$), season ($p = 0.003$), body condition ($p =$

0.093), and colony ($p = 0.107$) as fixed effects (Table 1a, Figure 3a). The total amount

of variance explained by this model was high (conditional $R^2 = 0.657$), as was the

repeatability of baseline cortisol values across individuals (ICC = 0.39). Including ID

as a random effect significantly improved the fit of the model, indicating appreciable

among-individual variability in baseline cortisol ($p < 0.001$; Supplementary Table

S1a). Allowing individual slopes to vary between age groups also significantly
improved model fit ($p < 0.001$; Supplementary Table S3) suggesting that individuals
responded to the covariates differently depending on their age. Overall, baseline
cortisol decreased with increasing pup age (Figure 3b) and was higher among pups
born in 2020, the year of higher food availability (Figure 3c). Baseline cortisol
decreased significantly as pup weight increased (Figure 3d), although the slope of the
regression between cortisol and weight approached zero as pups approached their
moult. Finally, baseline cortisol tended to be higher in males than females (Figure 3e).
The best-supported model of maternal baseline cortisol contained days
postpartum ($p < 0.001$), season ($p = 0.004$), and colony ($p = 0.068$) as fixed effects
(Table 1b, Figure 3f). Neither individual weight nor body condition were retained in
the model. The total variance explained by the model was again high (conditional R^2
$= 0.774$), as was repeatability of baseline cortisol within individuals (ICC $= 0.43$).
Including ID as a random effect significantly improved model fit, indicating
appreciable among-individual variability in baseline cortisol ($p < 0.001$;
Supplementary Table S3). The concentration of cortisol in maternal saliva decreased
as the season progressed (Figure 3g) and tended to be higher in 2020, the year of
higher food availability (Figure 3h).
344
345

Discussion

We used a fully crossed, repeated measures design to characterize individual

variation in baseline cortisol levels in a wild population of Antarctic fur seals. We

found that baseline cortisol was only marginally explained by genetic factors, while

high within-individual repeatability and among-individual variability in both pups and

mothers could be largely explained by a combination of intrinsic and extrinsic factors.

Our results provide detailed insights into the individualization of an endocrine

phenotype in a wild pinniped population.

Heritability estimates

We quantified the narrow sense heritability of baseline cortisol using animal

models based on a simple pedigree and a SNP array. The former approach assumes

that individuals of unknown parentage are unrelated to all other individuals in the

population, which can lead to heritability being underestimated (Kruuk 2004). By

contrast, genomic approaches are capable of quantifying unbiased relatedness for all

sampled individuals, but can be time-consuming and costly to produce (Frentiu et al.

2008). Despite these differences, we found that both approaches produced

consistently low heritability estimates for baseline cortisol in Antarctic fur seals.

Heritability can be low because of low additive genetic variation, high environmental

variance, or cross-environment genetic correlations (i.e. the genetic basis of the trait

varies between different environments) (Charmantier and Garant 2005). Our low

estimates, which stand in contrast to our original expectations and previously

published empirical estimates from other species (Jenkins et al. 2014; Stedman et al.

2017; Bairos-Novak et al. 2018), might therefore be reflective of the extreme

heterogeneity of the environmental conditions encountered by the study population at

Bird Island. This explanation would be in line with other empirical studies of wild

vertebrate populations showing a decrease in heritability under poor environmental

conditions (e.g. (Wilson et al. 2006)). Alternatively, low heritability may reflect the

different life stages at which cortisol concentrations were measured (i.e. mothers

verses pups).

Intrinsic and extrinsic factors affecting baseline cortisol

For both pups and mothers, we were able to explain a substantial proportion of
the total phenotypic variance in baseline cortisol by including individual-based and
environmental variables in our models (conditional $R^2 = 0.66$ for pups and $R^2 = 0.77$
for mothers). Similar results have been obtained for a variety of species (e.g. (Joly and
Cameron 2018; Uchida et al. 2021), but see (Azevedo et al. 2019)), suggesting that
contextual and developmental phenotypic plasticity in response to environmental
heterogeneity may be a widespread phenomenon. In addition, we found that baseline
cortisol levels were consistent within individuals, suggesting that individualized
endocrine phenotypes become established during early ontogeny and persist at least
until nutritional independence. Baseline cortisol might therefore represent a stable
attribute by which fur seals adapt to spatial or temporal heterogeneity in their
environment (Réale and Dingemanse 2010).

Our models uncovered a strong influence of age and days postpartum on
baseline cortisol levels in pups and mothers, respectively, with salivary cortisol
decreasing over time. One explanation for this may be the shifting environmental
conditions that individuals experience as the season progresses. Pregnant females
arrive ashore in December and give birth on crowded breeding beaches. Mothers
continue to suckle their pups on the beach until about 30 days postpartum, when most
females transition into the more sheltered and less crowded tussock grass that covers
most of the island's interior (Doidge et al. 1984). Also during this time, adult males
begin to abandon their territories and migrate to higher latitudes around the Antarctic
ice shelf (Forcada and Staniland 2018). Consequently, the frequency of unpredictable
challenges for both pups and mothers likely declines as the season progresses. A
corresponding reduction in baseline cortisol is therefore in line with previous research
suggesting that cortisol levels are lower under stable environmental conditions
compared with environments associated with frequent disturbances (Fairbanks et al.
2011).

We detected significantly higher cortisol concentrations among pups and
mothers sampled in 2020, the year of higher food availability. This was surprising
given the many empirical studies that have linked elevated cortisol concentrations to
food shortages and periods of nutritional stress (e.g. (Kitaysky et al. 2007; Behie et al.
2010; Bryan et al. 2013; Garber et al. 2020)). We can think of two possible

explanations for this pattern. On the one hand, our results could be explained by
higher population densities in the second year of our study, as significantly more
females bred in 2020 compared to 2019 (Nagel et al. 2021). This would be in line
with the small, albeit non-significant, effect of colony on baseline cortisol, with
hormone concentrations being marginally higher in both pups and mothers at SSB
compared to FWB. On the other hand, circulating levels of cortisol are essential for
the maintenance of metabolic functions (Kuo et al. 2015), and food-induced cortisol
secretions have been documented in the literature (Quigley and Yen 1979; Gibson et
al. 1999; Stimson et al. 2014). Shorter foraging trip durations (Nagel et al. 2021) and
consequently more frequent meals for pups and mothers in 2020 may have resulted in
higher average baseline cortisol concentrations, which facilitate protein and
carbohydrate metabolism.

More indicative of the hypothesized correlation between cortisol and
nutritional stress, we found a significant negative relationship between baseline
cortisol levels and weight in pups. Elevated cortisol levels provide individuals with a
source of energy by stimulating gluconeogenesis, which increases the delivery of
glucose into the bloodstream (Wingfield and Romero 2011). Furthermore, cortisol can
enhance fat oxidation by other mechanisms, like promoting production of hormone
sensitive lipase (Samra et al. 1996). Given that pups must tolerate bouts of fasting
lasting up to 11 days while their mothers forage at sea (Forcada and Staniland 2018),
our results may reflect a physiological response to prolonged periods of natural food
limitation (Ortiz et al. 2001; Jeanniard du Dot et al. 2009). In other words, fasting
pups may increase baseline cortisol to release energy, resulting in a reduction of
absolute body fat and overall weight. This would also explain why we do not see a
similar relationship in mothers, who remain ashore between foraging trips for as little
as 24 hours and on average only two days (Boyd et al. 1991). Alternatively, pups may
be more susceptible to environmental stressors than adult females, with lighter pups
requiring more energy to maintain homeostasis under, for example, unfavorable
climatic conditions.

In pups, we also uncovered a significant association between baseline cortisol
and sex, with hormone concentrations being moderately higher in males than females.
Previous studies of the sex-specific secretion of cortisol have produced diverse

results, including conflicting evidence from within a single species (Steller sea lion

pups: (Keogh et al. 2010; Myers et al. 2010). These contrasting findings highlight the

complexity of interactions between cortisol and the sex hormones, which can vary

with the reproductive system, phase, and cycle (Levine 2002). In addition, the social

and environmental stressors associated with growth and reproduction in each sex are

likely to vary among species. For example, male Antarctic fur seal pups engage more

often in social interactions and risk prone behaviors (Jones et al. 2020) such that the

number of "stressful" events encountered may be higher in males than females.

Contrary to our initial expectations, colony only had a marginal, non-

significant effect on baseline salivary cortisol in pups and their mothers. This is in

contrast to a previous study where cortisol concentrations, measured from hair, were

higher in mothers (but not offspring) from the high density colony (Meise et al. 2016).

We can think of two non-mutually exclusive explanations for this discrepancy. First,

hair cortisol concentrations are thought to reflect events in the recent past (hours or

days) (Kalliokoski et al. 2019), whereas salivary cortisol captures circulating hormone

levels at the immediate time of sampling (Lewis 2006). Consequently, hair

concentrations of baseline cortisol might integrate a larger number of stressful events,

allowing density-dependent differences to more readily accumulate. Second, the first

study was conducted in 2011 when population densities were much higher at SSB

(approximately 568 breeding females were observed at SSB in 2011 compared with

282 in 2019 and 409 in 2020, a reduction of 50% and 28%, respectively (Forcada and

Hoffman 2014; Nagel et al. 2021), which may have accentuated differences between

the two colonies.

Strengths, limitations, and future directions

The past decade has witnessed a shift in our understanding of individual

differences from a perceived statistical nuisance to a fundamental and compelling

aspect of behavioral ecology and evolutionary biology (Bolnick et al. 2003; Trillmich

et al. 2018; Krüger et al. 2021). Our study contributes towards this narrative by

decomposing individual variation in endocrine phenotypes across different life history

stages and environments in Antarctic fur seal pups and their mothers. Our results are

indicative of substantial developmental plasticity in the HPA axis of pups, with

individualized endocrine phenotypes becoming established during early ontogeny and

persisting at least until nutritional independence. Baseline cortisol may therefore help

to facilitate the match between an individual's phenotype and the environment. That

we find a higher relative contribution of the environment compared to genetic factors

to phenotypic variation further suggests that the endocrine phenotype may be highly

adaptable to unpredictable environmental conditions. This has implications for our

understanding of population dynamics, both in the declining Antarctic fur seal

population (Forcada and Hoffman 2014) and other in systems impacted by climate

change.

However, further research is needed to elucidate both the fitness consequences

of endocrine variability and how this may respond to environmental heterogeneity

over longer timescales. To meet the strictest definition of phenotypic plasticity,

individual variation in the focal trait must be demonstrated in different environments,

ideally across multiple years (Boutin and Lane 2014). For developmental plasticity,

an individual's phenotype must persist into later life stages, while changing reaction

norms over an individual's lifetime would suggest contextual plasticity (Nettle and

Bateson 2015; Trillmich et al. 2015). Our conclusions are also limited by the

comparison of only two colonies across two seasons. While including additional

colonies in the study would be logistically prohibitive given the inaccessibility of

most breeding beaches, a continuation of this study across additional breeding seasons

would certainly be feasible. Increasing the duration of this study would be particularly

relevant given the ever increasing occurrence and intensity of severe weather events

in the sub-Antarctic region (Turner et al. 2005).

Overall, our work builds upon the existing literature on the heritability and

phenotypic plasticity of baseline cortisol in response to environmental heterogeneity.

Although our results are framed in the context of the Antarctic fur seal, our study has

implications for understanding the importance of endocrine mechanisms in other

populations. We can contribute towards a growing body of evidence showing that

endocrine phenotypes are ecologically important traits that can potentially affect

population dynamics through their influence on life history traits. Expanding our

understanding of those extrinsic and intrinsic factors that influence baseline cortisol

therefore gives insight into the factors that potentially limit or improve population

persistence in a changing environment.

References

Atkinson S, Arnould JPY, Mashburn KL. 2011. Plasma cortisol and thyroid hormone

concentrations in pre-weaning Australian fur seal pups. Gen Comp Endocrinol.

172:277–281. doi:10.1016/j.ygcen.2011.03.014.

Azevedo A, Bailey L, Bandeira V, Dehnhard M, Fonseca C, Sousa L De, Jewgenow

K. 2019. Age, sex and storage time influence hair cortisol levels in a wild mammal

population. PLoS One. 14(8):e0221124. doi:10.1371/journal. pone.0221124.

Bairos-Novak KR, Ryan CP, Freeman AR, Gary Anderson W, Hare JF. 2018. Like

mother, like daughter: Heritability of female Richardson's ground squirrel

Urocitellus richardsonii cortisol stress responses. Curr Zool. 64(2):153–163.

doi:10.1093/cz/zox014.

Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models

using lme4. J Stat Softw. 67(1):1–48. doi:10.18637/jss.v067.i01.

Behie AM, Pavelka MSM, Chapman CA. 2010. Sources of variation in fecal cortisol

levels in howler monkeys in Belize. Am J Primatol. 72:600–606.

doi:10.1002/ajp.20813.

Bellavance MA, Rivest S. 2014. The HPA-immune axis and the immunomodulatory

actions of glucocorticoids in the brain. Front Immunol. 5:1–13.

doi:10.3389/fimmu.2014.00136.

Bennett KA, Moss SEW, Pomeroy P, Speakman JR, Fedak MA. 2012. Effects of

handling regime and sex on changes in cortisol, thyroid hormones and body mass in

fasting grey seal pups. Comp Biochem Physiol Part A. 161:69–76.

doi:10.1016/j.cbpa.2011.09.003. 10.1016/j.cbpa.2011.09.003.

Boake CRB. 1989. Repeatability: Its role in evolutionary studies of mating behavior.

Evol Ecol. 3(2):173–182. doi:10.1007/BF02270919.

Bolnick DI, Svanbäck R, Fordyce JA, Yang LH, Davis JM, Hulsey CD, Forister ML.

2003. The ecology of individuals: Incidence and implications of individual

specialization. Am Nat. 161(1):1–28. doi:10.1086/343878.

Bonier F, Martin PR. 2016. How can we estimate natural selection on endocrine

traits? Lessons from evolutionary biology. Proc R Soc B. 283:20161887.

doi:10.1098/rspb.2016.1887.

Boutin S, Lane JE. 2014. Climate change and mammals: Evolutionary versus plastic
responses. Evol Appl. 7(1):29–41. doi:10.1111/eva.12121.

Boyd IL, Lunn NJ, Barton T. 1991. Time budgets and foraging characteristics of
lactating Antarctic fur seals. J Anim Ecol. 60(2):577–592. doi:10.2307/5299.

Bozovic D, Racic M, Ivkovic N. 2013. Salivary cortisol levels as a biological marker
of stress reaction. Med Arch. 67(5):371–374. doi:10.5455/medarh.2013.67.374-377.

Bryan HM, Darimont CT, Paquet PC, Wynne-Edwards KE, Smits JEG. 2013. Stress
and reproductive hormones in grizzly bears reflect nutritional benefits and social
consequences of a salmon foraging niche. PLoS One. 8(11):e80537.
doi:10.1371/journal.pone.0080537.

Cassini MH. 1999. The evolution of reproductive systems in pinnipeds. Behav Ecol.
10(5):612–616. doi:10.1093/beheco/10.5.612.

Champagne CD, Houser DS, Costa DP, Crocker DE. 2012. The effects of handling
and anesthetic agents on the stress response and carbohydrate metabolism in
Northern elephant seals. PLoS One. 7(5):e38442.
doi:10.1371/journal.pone.0038442.

Charmantier A, Garant D. 2005. Environmental quality and evolutionary potential:
Lessons from wild populations. Proc R Soc B. 272(1571):1415–1425.
doi:10.1098/rspb.2005.3117.

Dall SRX, Bell AM, Bolnick DI, Ratnieks FLW. 2012. An evolutionary ecology of
individual differences. Ecol Lett. 15(10):1189–1198. doi:10.1111/j.1461-
0248.2012.01846.x.

DeRango EJ, Prager KC, Greig DJ, Hooper AW, Crocker DE. 2019. Climate
variability and life history impact stress, thyroid, and immune markers in California
sea lions (*Zalophus californianus*) during El Niño conditions. Conserv Physiol. 7:1–
15. doi:10.1093/conphys/coz010.

Dingemanse NJ, Réale D. 2005. Natural selection and animal personality. Behaviour.
142(9):1159–1184. doi:10.1163/156853905774539445.

Doidge DW, Croxall JP, Baker JR. 1984. Density-dependent pup mortality in the

Antarctic fur seal *Arctocephalus gazella* at South Georgia. J Zool London. 202:449–

460. doi:10.1111/j.1469-7998.1984.tb05095.x.

Engelhard GH, Brasseur SMJM, Hall AJ, Burton HR, Reijnders PJH. 2002.

Adrenocortical responsiveness in southern elephant seal mothers and pups during

lactation and the effect of scientific handling. J Comp Physiol B. 172:315–328.

doi:10.1007/s00360-002-0257-0.

Fairbanks LA, Jorgensen MJ, Bailey JN, Breidenthal SE, Grzywa R, Laudenslager

ML. 2011. Heritability and genetic correlation of hair cortisol in vervet monkeys in

low and higher stress environments. Psychoneuroendocrinology. 36:1201–1208.

doi:10.1016/j.psyneuen.2011.02.013.

http://dx.doi.org/10.1016/j.psyneuen.2011.02.013.

Forcada J, Hoffman JI. 2014. Climate change selects for heterozygosity in a declining

fur seal population. Nature. 511(7510):462–465. doi:10.1038/nature13542.

Forcada J, Staniland IJ. 2018. Antarctic Fur Seal. In: Würsig B, Thewissen JGM,

Kovacs KM, editors. Encyclopedia of Marine Mammals. 3rd ed. Academic Press. p.

25–27.

Frentiu FD, Clegg SM, Chittock J, Burke T, Blows MW, Owens IPF. 2008. Pedigree-

free animal models: The relatedness matrix reloaded. Proc R Soc B. 275(1635):639–

647. doi:10.1098/rspb.2007.1032.

Garber PA, McKenney A, Bartling-John E, Bicca-Marques JC, De la Fuente MF,

Abreu F, Schiel N, Souto A, Phillips KA. 2020. Life in a harsh environment: The

effects of age, sex, reproductive condition, and season on hair cortisol concentration

in a wild non-human primate. PeerJ. 8:e9365. doi:10.7717/peerj.9365.

Gastón MS, Vaira M. 2020. Male mating success is related to body condition and

stress-induced leukocyte response in an anuran with scramble competition. Can J

Zool. 98(6):391–398. doi:10.1139/cjz-2019-0193.

Gélin U, Coulson G, Festa-Bianchet M. 2016. Heterogeneity in reproductive success

explained by individual differences in bite rate and mass change. Behav Ecol.

27(3):777–783. doi:10.1093/beheco/arv209.

Gentry RL, Holt JR. 1982. Equipment and techniques for handling northern fur seals.
NOAA Tech Rep NMFS SSRF-758.:1–17.

Gibson EL, Checkley S, Papadopoulos A, Poon L, Daley S, Wardle J. 1999. Increased
salivary cortisol reliably induced by a protein-rich midday meal. Psychosom Med.
61(2):214–224. doi:10.1097/00006842-199903000-00014.

Graffelman J, Moreno V. 2013. The mid p-value in exact tests for Hardy-Weinberg
equilibrium. Stat Appl Genet Mol Biol. 12(4):433–448. doi:10.1515/sagmb-2012-
0039.

Hadfield JD. 2010. MCMC methods for multi-response generalized linear mixed
models: The MCMCglmm R package. J Stat Softw. 33(2):1–22.
doi:10.1002/ana.22635.

Harcourt RG, Turner E, Hall A, Waas JR, Hindell M. 2010. Effects of capture stress
on free-ranging, reproductively active male Weddell seals. J Comp Physiol A.
196:147–154. doi:10.1007/s00359-009-0501-0.

Humble E, Paijmans AJ, Forcada J, Hoffman JI. 2020. A 85K SNP array uncovers
inbreeding and cryptic relatedness in an Antarctic fur seal breeding colony. G3.
10:2787–2799. doi:10.1101/2020.04.01.020123.

Hunt GL, Heinemann D, Everson I. 1992. Distributions and predator-prey interactions
of macaroni penguins, Antarctic fur seals, and Antarctic krill near Bird Island, South
Georgia. Mar Ecol Prog Ser. 86(1):15–30. doi:10.3354/meps086015.

Jeanniard du Dot T, Rosen DAS, Richmond JP, Kitaysky AS, Zinn SA, Trites AW.
2009. Changes in glucocorticoids, IGF-I and thyroid hormones as indicators of
nutritional stress and subsequent refeeding in Steller sea lions (*Eumetopias jubatus*).
Comp Biochem Physiol Part A. 152:524–534. doi:10.1016/j.cbpa.2008.12.010.

Jenkins BR, Vitousek MN, Hubbard JK, Safran RJ. 2014. An experimental analysis of
the heritability of variation in glucocorticoid concentrations in a wild avian
population. Proc R Soc B. 281:20141302. doi:10.1098/rspb.2014.1302.

Joly K, Cameron MD. 2018. Early fall and late winter diets of migratory caribou in
northwest Alaska. Rangifer. 38(1):27–38. doi:10.7557/2.38.1.4107.

Jones KA, Wood H, Jonathan P, Ratcliffe N, Votier SC, Staniland IJ. 2020. Risk

exposure trade-offs in the ontogeny of sexual segregation in Antarctic fur seal pups.

Behav Ecol.:1–12. doi:10.1093/beheco/araa018.

Kalliokoski O, Jellestad FK, Murison R. 2019. A systematic review of studies

utilizing hair glucocorticoids as a measure of stress suggests the marker is more

appropriate for quantifying short-term stressors. Sci Rep. 9:11997.

doi:10.1038/s41598-019-48517-2.

Keogh MJ, Maniscalco JM, Atkinson S. 2010. Steller sea lion (*Eumetopias jubatus*)

pups undergo a decrease in circulating white blood cells and the ability of T cells to

proliferate during early postnatal development. Vet Immunol Immunopathol.

137(3–4):298–304. doi:10.1016/j.vetimm.2010.05.008.

http://dx.doi.org/10.1016/j.vetimm.2010.05.008.

Kitaysky AS, Piatt JF, Wingfield JC. 2007. Stress hormones link food availability and

population processes in seabirds. Mar Ecol Prog Ser. 352:245–258.

doi:10.3354/meps07074.

Krüger O, Anaya-Rojas J, Caspers B, Chakarov N, Elliot-Graves A, Fricke C, Gadau

J, Gossmann T, Hoffman J, Kaiser MI, et al. 2021. Individualised niches: An

integrative conceptual framework across behaviour, ecology, and evolution.

EcoEvoRxiv. preprint. doi:10.32942/osf.io/7h5xq.

Kruuk LEB. 2004. Estimating genetic parameters in natural populations using the

"animal model." Philos Trans R Soc B. 359(1446):873–890.

doi:10.1098/rstb.2003.1437.

Kruuk LEB, Merilä J, Sheldon BC. 2003. When environmental variation short-circuits

natural selection. Trends Ecol Evol. 18(5):207–209. doi:10.1016/S0169-

5347(03)00073-9.

Kuo T, McQueen A, Chen T-C, Wang Jen-Chywan. 2015. Regulation of glucose

homeostasis by glucocorticoids. In: Wang J.-C., Harris C, editors. Glucocorticoid

Signaling. Vol. 872. New York: Springer. p. 99–126.

http://www.ncbi.nlm.nih.gov/pubmed/26215998.

Levine JE. 2002. Editorial: Stressing the importance of sex. Endocrinology.

143(12):4502–4504. doi:10.1210/en.2002-221041.

Lewis JG. 2006. Steroid analysis in saliva: An overview. Clin Biochem Rev. 27:139–
146.

Lightman SL, Conway-Campbell BL. 2010. The crucial role of pulsatile activity of
the HPA axis for continuous dynamic equilibration. Nat Rev Neurosci. 11(10):710–
718. doi:10.1038/nrn2914.

Lihoreau M, Kaiser S, Resende B, Rödel HG, Châline N. 2021. Editorial: Context-
dependent plasticity in social species: Feedback loops between individual and social
environment. Front Psychol. 12:645191. doi:10.3389/fpsyg.2021.645191.

McCafferty DJ, Boyd IL, Taylor RI. 1998. Diving behaviour of Antarctic fur seal
(*Arctocephalus gazella*) pups. Can J Zool. 76(3):513–520. doi:10.1139/z97-219.

Meise K, Von Engelhardt N, Forcada J, Hoffman JI. 2016. Offspring hormones reflect
the maternal prenatal social environment: Potential for foetal programming? PLoS
One. 11(1):e0145352. doi:10.1371/journal.pone.0145352.

Milenkaya O, Catlin DH, Legge S, Walters JR. 2015. Body condition indices predict
reproductive success but not survival in a sedentary, tropical bird. PLoS One.
10(8):e0136582. doi:10.1371/journal.pone.0136582.

Müller C, Caspers BA, Gadau J, Kaiser S. 2020. The power of infochemicals in
mediating individualized niches. Trends Ecol Evol. 35(11):981–989.
doi:10.1016/j.tree.2020.07.001.

Mutwill AM, Schielzeth H, Zimmermann TD, Richter SH, Kaiser S, Sachser N. 2021.
Individuality meets plasticity: Endocrine phenotypes across male dominance rank
acquisition in guinea pigs living in a complex social environment. Horm Behav.
131:104967. doi:10.1016/j.yhbeh.2021.104967.

Myers MJ, Litz B, Atkinson S. 2010. The effects of age, sex, season and geographic
region on circulating serum cortisol concentrations in threatened and endangered
Steller sea lions (*Eumetopias jubatus*). Gen Comp Endocrinol. 165(1):72–77.
doi:10.1016/j.ygcen.2009.06.006. http://dx.doi.org/10.1016/j.ygcen.2009.06.006.

Nagel R, Stainfield C, Fox-Clarke C, Toscani C, Forcada J, Hoffman J. 2021.

Evidence for an Allee effect in a declining fur seal population. Proc R Soc B.

288:20202882. doi:https://doi.org/10.1098/rspb.2020.2882.

Nakagawa S, Schielzeth H. 2013. A general and simple method for obtaining R2 from

generalized linear mixed-effects models. Methods Ecol Evol. 4(2):133–142.

doi:10.1111/j.2041-210x.2012.00261.x.

Nettle D, Bateson M. 2015. Adaptive developmental plasticity: what is it, how can we

recognize it and when can it evolve? Proc R Soc B. 282:20151005.

doi:10.1098/rspb.2015.1005.

Ortiz RM, Houser DS, Wade CE, Ortiz CL. 2003. Hormonal changes associated with

the transition between nursing and natural fasting in northern elephant seals

(*Mirounga angustirostris*). Gen Comp Endocrinol. 130:78–83. doi:10.1016/S0016-

6480(02)00572-5.

Ortiz RM, Wade CE, Ortiz CL. 2001. Effects of prolonged fasting on plasma cortisol

and TH in postweaned northern elephant seal pups. Am J Physiol - Regul Integr

Comp Physiol. 280:R790–R795. doi:10.1006/gcen.2000.7514.

Pavitt AT, Walling CA, Möstl E, Pemberton JM, Kruuk LEB. 2015. Cortisol but not

testosterone is repeatable and varies with reproductive effort in wild red deer stags.

Gen Comp Endocrinol. 222:62–68. doi:10.1016/j.ygcen.2015.07.009.

Payne MR. 1979. Growth in the Antarctic fur seal *Arctocephalus gazella*. J Zool

London. 187:1–20.

Peig J, Green AJ. 2009. New perspectives for estimating body condition from

mass/length data: The scaled mass index as an alternative method. Oikos.

118:1883–1891. doi:10.1111/j.1600-0706.2009.17643.x.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J,

Sklar P, de Bakker PIW, Daly MJ, et al. 2007. PLINK: A tool set for whole-genome

association and population-based linkage analyses. Am J Hum Genet. 81(3):559–

575. doi:10.1086/519795.

Quigley ME, Yen SSC. 1979. A mid-day surge in cortisol levels. J Clin Endocrinol

Metab. 49(6):945–947. doi:10.1210/jcem-49-6-945.

Réale D, Dingemanse NJ. 2010. Personality and individual social specialization. In:

Székely T, Moore AJ, Komdeur J, editors. Social Behaviour: Genes, Ecology and

Evolution. Cambridge University Press. p. 417–441.

Romero LM, Gormally BMG. 2019. How truly conserved is the "well-conserved"

vertebrate stress response? Integr Comp Biol. 59(2):273–281.

doi:10.1093/icb/icz011.

Romero ML, Butler LK. 2007. Endocrinology of stress. Int J Comp Psychol. 20:89–

95. https://escholarship.org/uc/item/87d2k2xz.

Samra JS, Clark ML, Humphreys SM, Macdonald IA, Matthews DR, Frayn KN.

1996. Effects of morning rise in cortisol concentration on regulation of lipolysis in

subcutaneous adipose tissue. Am J Physiol. 271:E996-1002.

doi:10.1152/ajpendo.1996.271.6.E996.

Schoenemann KL, Bonier F. 2018. Repeatability of glucocorticoid hormones in

vertebrates: A meta-analysis. PeerJ. 6:e4398. doi:10.7717/peerj.4398.

Sih A, Cote J, Evans M, Fogarty S, Pruitt J. 2012. Ecological implications of

behavioural syndromes. Ecol Lett. 15(3):278–289. doi:10.1111/j.1461-

0248.2011.01731.x.

Stamps JA, Biro PA. 2016. Personality and individual differences in plasticity. Curr

Opin Behav Sci. 12:18–23. doi:10.1016/j.cobeha.2016.08.008.

http://dx.doi.org/10.1016/j.cobeha.2016.08.008.

Stedman JM, Hallinger KK, Winkler DW, Vitousek MN. 2017. Heritable variation in

circulating glucocorticoids and endocrine flexibility in a free-living songbird. J Evol

Biol. 30:1724–1735. doi:10.1111/jeb.13135.

Stimson RH, Mohd-Shukri NA, Bolton JL, Andrew R, Reynolds RM, Walker BR.

2014. The postprandial rise in plasma cortisol in men is mediated by macronutrient-

specific stimulation of adrenal and extra-adrenal cortisol production. J Clin

Endocrinol Metab. 99(1):160–168. doi:10.1210/jc.2013-2307.

Taff CC, Schoenle LA, Vitousek MN. 2018. The repeatability of glucocorticoids: A

review and meta-analysis. Gen Comp Endocrinol. 260:136–145.

doi:10.1016/j.ygcen.2018.01.011.

Trillmich F, Günther A, Müller C, Reinhold K, Sachser N. 2015. New perspectives in
behavioural development: Adaptive shaping of behaviour over a lifetime? Front
Zool. 12:S1. doi:10.1186/1742-9994-12-S1-S1.

Trillmich F, Müller T, Müller C. 2018. Understanding the evolution of personality
requires the study of mechanisms behind the development and life history of
personality traits. Biol Lett. 14(2):20170740. doi:10.1098/rsbl.2017.0740.

Turner J, Colwell SR, Marshall GJ, Lachlan-Cope TA, Carleton AM, Jones PD,
Lagun V, Reid PA, Iagovkina S. 2005. Antarctic climate change during the last 50
years. Int J Climatol. 25(3):279–294. doi:10.1002/joc.1130.

Uchida K, Ng R, Vydro SA, Smith JE, Blumstein DT. 2021. The benefits of being
dominant: Health correlates of male social rank and age in a marmot. Curr Zool.:1–
8. doi:10.1093/cz/zoab034.

Vessey SH, Drickamer LC. 2010. Integration of proximate and ultimate causes. In:
Breed M, Moore J, editors. Encyclopedia of Animal Behavior. 1st ed. Academic
Press. p. 180–185.

Wilson AJ, Pemberton JM, Pilkington JG, Coltman DW, Mifsud D V., Clutton-Brock
TH, Kruuk LEB. 2006. Environmental coupling of selection and heritability limits
evolution. PLoS Biol. 4(7):1270–1275. doi:10.1371/journal.pbio.0040216.

Wingfield JC, Romero LM. 2011. Adrenocortical responses to stress and their
modulation in free-living vertebrates. In: Comprehensive Physiology. Hoboken, NJ,
USA: John Wiley & Sons, Inc. p. 211–234.

Wolf M, Weissing FJ. 2012. Animal personalities: Consequences for ecology and
evolution. Trends Ecol Evol. 27(8):452–461. doi:10.1016/j.tree.2012.05.001.

Yang J, Lee SH, Goddard ME, Visscher PM. 2011. GCTA: A tool for genome-wide
complex trait analysis. Am J Hum Genet. 88:76–82.
doi:10.1016/j.ajhg.2010.11.011.

Acknowledgements

The authors would like to thank Ana Bertoldi Carneiro, Freya Blockley, Jamie Coleman, Alexandra Dodds, Vicki Foster, Derren Fox, Iain Angus Gordon, Pauline Goulet, Rosie Hall, Cary Jackson, Adam Lowndes, Elizabeth Morgan, Rachael Orben, Jessica Ann Philips, David Reid, and Mark Whiffin for additional help in the field. We kindly thank Sabine Kruse for conducting the endocrine analyses. We are also grateful to Jonas Schwartz for tips on how to collect saliva samples from young pups in the field. A special thanks to Océane Salles for producing the fur seal cartoon in Figure 1c.

Conflict of Interest

The authors declare no conflict of interest.

Author contributions

RN, CS, CT, and CF-C collected the data. SK advised on field protocols and coordinated cortisol quantification. RN analyzed the data. AJP generated the SNP genotype dataset and DLJV built the genomic relatedness matrix. RN and CC performed the molecular laboratory work. JIH, SK, and JF conceived and developed the project. RN drafted the manuscript. All of the authors commented on and approved the final manuscript.

Data Availability Statement

Our code and accompanying documentation are provided in the form of an R Markdown file (Supplementary Information). The data used for this study are available via Zenodo, https://doi.org/10.5281/zenodo.6336716.

Funding

This work was supported by the German Research Foundation (DFG) as part of the SFB TRR 212 (NC³) (project numbers 316099922, 396774617). It was also supported

801 by the priority program "Antarctic Research with Comparative Investigations in
802 Arctic Ice Areas" SPP 1158 (project number 424119118) and core funding from the
803 Natural Environment Research Council to the British Antarctic Survey's Ecosystems
804 Program.

Figures

806
Figure 1: Location and study design. (a) Map of Bird Island, South Georgia, a sub-Antarctic island in the southern Atlantic Ocean. The

inset shows an enlarged view of the two study colonies from which mother-pup pairs were sampled. Freshwater Beach (FWB, shown in

blue) and Special Study Beach (SSB, shown in red) are separated by approximately 200 meters. (b) We employed a fully crossed

sampling scheme involving the collection of saliva samples from a total of 100 pairs from the two colonies in two successive breeding

seasons, the first of which was coincidentally a year of particularly low food availability. (c) Each focal mother was sampled twice in a

season while pups were sampled every 20 days from birth until weaning.

814
815 Figure 2: Posterior distributions of heritability (h^2) estimates for baseline cortisol. A

816 simple pedigree (mother-offspring pairs) for the entire dataset and a custom 85K SNP

817 array for the 95 individuals sampled in the 2019 season were used to calculate the

818 relatedness matrix. The modes and highest posterior density intervals of the posterior

819 distributions are shown as points and bars, respectively.

821
Figure 3: Generalized linear mixed models for pup (a – e) and mother (f – h) baseline

cortisol values. Estimates ± 95% confidence intervals for all fixed effects included in

the best fit models for pups and mothers are shown in panels a and f, respectively.

Significant main effects for both models are shown in panels b – e and panels g – h,

respectively. Boxes show median values ± 75% percentiles with the vertical lines

indicating 95% confidence intervals. Further details of the model output can be found

in Table 1.

 Tables

Table 1: Parameter estimates from the best fit generalized linear mixed models of (a) pup and

(b) maternal baseline cortisol. Random intercepts were included for each individual to account

for repeated measures. Estimates together with their 95% confidence intervals (CI) as well as

Wald *t*-values and chi-squared values are presented. Significant *p*-values are in bold. The

mean squared error (σ^2), between group variance ($\tau00$), Intraclass Correlation Coefficient

(ICC; the consistency within an individual across multiple measurements), the sample size (*n*)

and total number of observations, as well as the variance explained by the fixed effects

(marginal R^2) and variance explained by both fixed and random effects (conditional R^2) are

given.

(a) Pup baseline cortisol					
Fixed effects	Estimates	CI	*t*	*chi-squared*	*p*
(Intercept)	1.58	1.40 – 1.75	17.75		**<0.001**
Age	-0.29	-0.44 – -0.14	-3.70	13.68	**<0.001**
Season [2020]	0.25	0.09 – 0.41	3.00	8.99	**0.003**
Weight	-0.36	-0.52 – -0.20	-4.44	19.71	**<0.001**
Sex [male]	0.24	0.08 – 0.41	0.09	8.11	**0.004**
Body condition	0.14	-0.02 – 0.31	1.68	2.83	0.093
Colony [SSB]	0.13	-0.03 – 0.29	0.08	2.60	0.107
Random effect					
σ^2	0.18				
$\tau_{00\ ID}$	0.05				
ICC	0.39				
n $_{ID}$	96				
Observations	290				
Marginal R^2 / conditional R^2	0.440 / 0.657				
(b) Maternal baseline cortisol					
Fixed effects	Estimates	CI	*t*	*chi-squared*	*p*
(Intercept)	1.39	1.18 – 1.60	12.93		**<0.001**
Days postpartum	-0.66	-0.73 – -0.59	-18.62	347.01	**<0.001**
Season [2020]	0.35	0.11 – 0.59	2.87	8.25	**0.004**
Colony [SSB]	0.22	-0.02 – 0.47	1.82	3.32	0.068

Random effect	
σ^2	0.16
$\tau_{00\ \text{ID}}$	0.12
ICC	0.43
n_{ID}	92
Observations	145
Marginal R^2 / conditional R^2	0.603 / 0.774