1   Low heritability and high phenotypic plasticity of cortisol in response to environmental

2   heterogeneity in fur seals

Running header: Response of the cortisol phenotype to environmental heterogeneity

Rebecca Nagel1,†, Sylvia Kaiser2, Claire Stainfield3, Camille Toscani3, Cameron Fox-

Clarke3, Anneke J. Paijmans1, Camila Costa Castro1, David L. J. Vendrami1, Jaume

Forcada3,*, Joseph I. Hoffman 1,3,*

1. Department of Animal Behaviour, Bielefeld University, 33501 Bielefeld, Germany

2. Department of Behavioural Biology, University of Münster, 48149 Münster, Germany

3. British Antarctic Survey, Cambridge CB3 OET, UK

* Joint senior authors

† Corresponding author: rebecca.nagel@uni-bielefeld.de

Lay summary

Stress is ubiquitous, but individuals differ. In Antarctic fur seals, individual stress levels

can be explained by sex, weight, day, density and food availability. Stress can influence

how an individual behaves and interacts with its environment, so gaining a better

understanding of those factors that explain variation in the stress response is important

for population monitoring and conservation.

Title

Low heritability and high phenotypic plasticity of salivary cortisol in response to

environmental heterogeneity in fur seals

Abstract

Individuals are unique in how they interact with and respond to their environment.

Correspondingly, unpredictable challenges or stressors often produce an individualized

response of the hypothalamic-pituitary-adrenal axis and its downstream effector cortisol.

We used a fully crossed, repeated measures design to investigate the factors shaping

individual variation in baseline cortisol and cortisol responsiveness in Antarctic fur seal

pups and their mothers. Saliva samples were collected from focal individuals at two

breeding colonies, one with low and the other with high population density, during two

consecutive years of contrasting food availability. Mothers and pups were sampled

concurrently at birth and shortly before weaning, while pups were additionally sampled

every 20 days. We found that heritability was low for both baseline cortisol and cortisol

responsiveness, while within-individual repeatability and among-individual variability

were high. A substantial proportion of the variation in baseline cortisol could be

explained in pups and mothers by a combination of intrinsic and extrinsic factors

including sex, weight, day, season, and colony of birth. However, the same variables

explained little of the variation in cortisol responsiveness. Our findings provide detailed

insights into the individualization of endocrine phenotypes in a wild pinniped.

Keywords

Antarctic fur seal, baseline cortisol, cortisol responsiveness, individualization, phenotypic

plasticity, pinniped

Introduction

Despite similarities in age, sex, or social status, individuals often differ in how they interact with their environment (Réale and Dingemanse 2010; Dall et al. 2012). Once perceived as a statistical nuisance, an ever growing body of evidence now suggests that these individual differences are often consistent and stable across time and contexts, with profound implications for understanding phenotypic variation, niche specialization, and animal personality (Sih et al. 2012; Wolf and Weissing 2012). The past decade has thus witnessed an increased awareness of individualization as a fundamentally important and compelling aspect of evolutionary biology, ecology, and animal behavior (Bolnick et al. 2003; Trillmich et al. 2018; Krüger et al. 2021).

Consistent differences among individuals are likely to be mediated by a combination of intrinsic and extrinsic factors and can be understood as the interaction between an individual's phenotype, genotype, and its ecological context such that its fitness is maximized (Dingemanse and Réale 2005; Vessey and Drickamer 2010; Lihoreau et al. 2021). At the proximate level, individualized phenotypic adjustments to environmental factors may be governed by, among other things, variation in the concentrations of circulating hormones (Müller et al. 2020). In particular, individual variation in cortisol, commonly known as one of the most important stress hormones, appears to play a major role in shaping individual responses to environmental conditions (Wingfield and Romero 2011).

Cortisol is a steroid hormone that belongs to the class of glucocorticoids. Its release is regulated by the hypothalamic-pituitary-adrenal (HPA) axis. Glucocorticoids

play an essential role in maintaining metabolic and homeostatic functions (Kuo et al.

2015). Under predictable conditions, cortisol is released continuously at baseline levels

that vary naturally throughout the day and over an individual's lifetime (Lightman and

Conway-Campbell 2010). In the face of unpredictable challenges, however, activation of

the HPA axis results in increased levels of secreted cortisol for the duration of the

stressor, before levels return to baseline (Bellavance and Rivest 2014).

While the physiological effects of circulating cortisol are conserved across

individuals and species (Wingfield and Romero 2011), empirical studies have

documented significant differences in baseline cortisol and cortisol responsiveness

among individuals (Schoenemann and Bonier 2018; Taff et al. 2018). These differences

could potentially facilitate adaptation to different habitats and environmental conditions

(Sih et al. 2012). For example, cortisol levels have been shown to vary significantly

among individuals experiencing different population densities (Meise et al. 2016) and

levels of nutritional stress (Kitaysky et al. 2007). Variation in cortisol among individuals

has also been linked to intrinsic factors such as age (Pavitt et al. 2015), sex (Azevedo et

al. 2019), and weight (Jeanniard du Dot et al. 2009). Furthermore, when such factors are

accounted for, empirical studies have shown that cortisol levels are often highly

repeatable within individuals (Schoenemann and Bonier 2018; Taff et al. 2018). In guinea

pigs, for example, cortisol responsiveness and to a lesser extent baseline cortisol levels

are highly repeatable from late adolescence to adulthood (Mutwill et al. 2021).

Given that cortisol levels often exhibit high among-individual variability and

within-individual repeatability, it has been argued that genetic rather than environmental

factors could explain much of the observed phenotypic variance, which would imply a

high evolvability of this endocrine trait (Boake 1989; Jenkins et al. 2014).

Correspondingly, several empirical studies have reported moderate to high levels of

cortisol heritability in free-living vertebrate populations (Jenkins et al. 2014; Stedman et

al. 2017; Bairos-Novak et al. 2018). However, persistent intrinsic and / or extrinsic

factors might also produce repeatable phenotypes regardless of the underlying genotype

(Taff et al. 2018), a possibility that has often been overlooked in the literature (Bonier

and Martin 2016). Such examples may include variation in the endocrine phenotype in

response to current internal and environmental stimuli (contextual plasticity) or stimuli

from the past (developmental plasticity) (Stamps and Biro 2016).

Pinnipeds, and otariids in particular, are ideally suited to investigate the effects of

internal and environmental factors on cortisol levels. First, otariids are colonially

breeding, with males competing to establish and maintain harems on densely packed

breeding beaches (Forcada and Staniland 2018). Cortisol may play an important role in

how individuals adapt to this dynamic environment by restoring homeostasis after

unpredictable challenges such as territorial bouts or unwanted mating attempts. Second,

while many breeding beaches do not differ appreciably in qualities such as substrate type

or topology, the density of individuals often varies from one place to another, setting up a

spatial dynamic that tends to remain stable over time (Cassini 1999). Consequently, as

pups are born on land and remain ashore throughout much of their early ontogeny (Payne

1979; McCafferty et al. 1998), cortisol might play an important role in mediating

individual responses to variation in density. Finally, baseline cortisol levels and cortisol

responsiveness have been investigated in several pinniped species in relation to ontogeny

(Ortiz et al. 2003; Atkinson et al. 2011), environmental conditions (DeRango et al. 2019),

and handling regimes (Engelhard et al. 2002; Harcourt et al. 2010; Bennett et al. 2012;

Champagne et al. 2012). Methodologies for collecting and assessing baseline cortisol and

cortisol responsiveness in pinnipeds are therefore well established in the literature.

Our model otariid species, the Antarctic fur seal (*Arctocephalus gazella*), has

been extensively studied by the British Antarctic Survey (BAS) on Bird Island, South

Georgia since the 1980s. Two breeding colonies on the island provide a unique "natural

experiment" for investigating individual responses to population density. Freshwater

Beach (FWB) and Special Study Beach (SSB) are situated less than 200 meters apart

(Figure 1a), meaning they are exposed to comparable climatic conditions. Breeding

females from both locations also likely forage in the same areas (Hunt et al. 1992) and do

not differ significantly in quality traits such as body size and condition (Nagel et al.

2021). Despite these similarities, the two colonies differ in the density of conspecifics.

Direct counts of individuals ashore suggest that the density of breeding females is almost

four times higher at SSB than FWB (Meise et al. 2016) and the modal local density of

focal pups across the entire breeding season is also higher for pups born at SSB (Nagel et

al. 2021).

We took advantage of this unique natural setup to investigate the intrinsic and

extrinsic factors shaping individual variation in baseline cortisol and cortisol

responsiveness in Antarctic fur seal pups and their mothers. We used a fully crossed,

repeated measures design (Figure 1b, c) comprising longitudinal data from mother-

offspring pairs from the two colonies over two consecutive breeding seasons, the first of

which was coincidentally one of the worst years on record with respect to food

availability (Forcada and Hoffman 2014; Nagel et al. 2021). Specifically, we collected

saliva samples and accompanying biometric data from 25 randomly selected focal pairs

from each colony in both seasons. To quantify baseline and cortisol responsiveness

levels, saliva was collected once immediately after capture and again after approximately

ten minutes of "capture stress", a procedure known to elicit a response of the HPA axis

(reviewed in (Wingfield and Romero 2011)). Mothers were sampled twice during the

breeding season, while pups were sampled every 20 days from birth until just before

molting at around 60 days of age.

We used animal models to obtain heritability estimates for baseline cortisol and

cortisol responsiveness using both a simple pedigree and a genomic relatedness matrix

obtained from a high density single nucleotide polymorphism (SNP) array (Humble et al.

2020). We then used linear mixed models to evaluate the within- and among-individual

variability of cortisol levels in pups and their mothers. Included in each model as

explanatory variables were multiple intrinsic (sex, weight, body condition, and days after

initial sampling) and extrinsic (density and year) variables. In line with previous studies

of wild vertebrate populations, we hypothesized that cortisol levels would be heritable

and repeatable within individuals. We further hypothesized that baseline cortisol and

cortisol responsiveness would be higher in mothers and pups from the high-density

colony and in the season of low food availability.

Methods

Field study

This study was conducted during the Antarctic fur seal breeding seasons
(December to March) of 2018–19 (hereafter 2019) and 2019–20 (hereafter 2020) at Bird
Island, South Georgia (54°00'24.8□ S, 38°03'04.1□ W). Each season, we sampled 25
unique mother-pup pairs from two neighboring breeding colonies, one of low (FWB) and
the other of high density (SSB) (Figure 1a). Sampling at both locations was randomized
with respect to pup sex, resulting in a final sample size of 51 male and 49 female pups.
Pup mortality was 25.6% averaged over the two colonies and seasons.

Each mother and her pup were captured concurrently on two separate occasions:
2–3 days postpartum (December) and again as the pups began to moult shortly before
weaning (March). Pups were additionally recaptured every 20 days. For the capture,
restraint, and sampling of individuals, we employed protocols that have been established
and refined over 30 consecutive years of the BAS long-term monitoring and survey
program. Briefly, adult females were captured with a noosing pole and held on a restraint
board during processing. Pups were captured with a slip noose or by hand and were
restrained by hand. After sampling, individuals were released as close to their capture site
as possible and, when present, pups were reunited with their mothers.

At the first sampling, focal individuals were fitted with VHF transmitters to the
dorsal side of the neck between the shoulder blades with epoxy glue (mothers: Sirtrack
core marine glue-on V2G 154C; pups: Sirtrack core marine glue-on V2G 152A).
Transmitter signals were monitored throughout the season using a hand-held VHF

receiver (AOR LTD., AR8200). Focal individuals were also given cattle ear tags (Dalton

Supplies, Henley on Thames, UK) in the trailing edge of each foreflipper (Gentry and

Holt 1982) for identification. Tissue plugs were collected and stored in 20% dimethyl

sulfoxide (DMSO) saturated with salt at -20°C for subsequent genetic analysis.

At every capture, weight and length measurements were taken from which a

scaled mass index was calculated according to (Peig and Green 2009). This condition

metric serves as a reliable indicator of overall fitness as it has been correlated with,

among other things, offspring survival (Milenkaya et al. 2015; Gélin et al. 2016) and

mating success (Gastón and Vaira 2020). Two saliva samples were collected at every

capture, the first of which was taken within three minutes of capture to provide data on

baseline cortisol levels (Bozovic et al. 2013). The second saliva sample was taken 10 ± 2

minutes thereafter to provide data on cortisol responsiveness. Saliva was collected by

rotating sterile cotton tip applicators fitted in polypropylene tubes (ROTH, Art. No.

XC10.2) in the cheek pouch and under the tongue. Samples were centrifuged and stored

at -20°C for subsequent cortisol analysis.

Hormone quantification

Saliva samples were thawed and centrifuged for ten minutes to separate the

mucins. The clear supernatant was then used for the determination of cortisol

concentrations. Samples contaminated with blood (reddish supernatant) were discarded (*n*

= 30 in 2019 and *n* = 31 in 2020), as cortisol values are often falsely elevated in such

samples. Cortisol concentrations were determined in duplicate using an enzyme-linked

immunosorbent assays (cortisol free in saliva DES6611, Demeditec Diagnostics GmbH,

Kiel, Germany). The intra-assay variances were on average CV = 5.2% and the inter-

assay variances were on average CV = 6.9%. The antibody showed the following ross-

reactivities: cortisol 100%, 11-desoxycortisol 50%, corticosterone 6.2%, 11-

Deoxycorticosterone 2.6%, 17α-oh-progesterone 1.3%, cortisone and prednisone < 1%,

testosterone, estradiol, and androstendione < 0.1%.

SNP genotyping and genomic relatedness matrix construction

For the 96 focal individuals sampled in 2019, we extracted total genomic DNA

from tissue samples using a standard chloroform-isoamylalcohol protocol (for a

description of the full protocol, see the Supplementary methods). SNP genotyping was

performed on these samples from 2019 using a custom Affymetrix SNP array as

described by (Humble et al. 2020). Quality control of the raw output data and genotyping

were implemented using the Axiom Analysis Suite (5.0.1.38, Affymetrix) based on

parameter thresholds set to their default values for diploid organisms. SNPs that were

initially classified as "off target variants" (OTV) were recovered by using the "Run OTV

caller" function. Of the 85,359 SNPs tiled on the array, 77,873 were retained for further

analysis representing SNPs classified as "PolyHighResolution" (SNPs that passed all of

the Axiom Analysis Suite quality controls) and "NoMinorHomozygote" (SNPs that

passed all quality controls but no homozygote genotypes for the minor allele were found).

An additional 3,423 SNPs with minor allele frequencies below 0.01 and 2,096 SNPs that

departed significantly from Hardy Weinberg equilibrium (HWE) were removed using

PLINK version 1.9 (Purcell et al. 2007). Departures from HWE were identified based on

an alpha level of 0.01 after implementing mid-*p* adjustment (Graffelman and Moreno

2013). After filtering, a total of 72,354 SNPs were retained and used to produce a

genomic relatedness matrix using the --make-grm option in GCTA version 1.93.1 (Yang

et al. 2011).

Heritability of cortisol levels

To quantify the proportion of the total variance in baseline cortisol attributable to

genetic differences among individuals, we fitted two multivariate generalized linear

mixed models (GLMMs) in *MCMCglmm* (Hadfield 2010) with baseline cortisol as the

dependent variable and individual ID and relatedness as random effects. For the first

model, a simple pedigree (comprising mother-offspring pairs) was built for the full

dataset, while the second model incorporated the SNP relatedness matrix from

individuals sampled only in the first season (2019). We used weak but informative priors

(0.05 of the observed phenotypic variance) in both models. Markov chains were run for

9,000,000 iterations and we retained every 8,500[th] value after removing 150,000

iterations of burn-in to generate posterior distributions of the random parameters. The

posterior distribution of the model intercept and autocorrelation were checked to assess

model fit. We obtained estimates of baseline cortisol heritability by dividing the additive

genetic variance by the total phenotypic variance ($h^2 = V_A/V_P$) for each sample of the

posterior distribution. We used the same approach to estimate the proportion of variance

in cortisol responsiveness, calculated as the difference between baseline and response

cortisol concentrations.

Estimating intrinsic and extrinsic factors influencing baseline cortisol

To determine whether weight, body condition, age, sex, season, colony, and an

interaction between season and colony explained a significant proportion of the variation

in baseline cortisol among pups, we fitted a GLMM with a log-link gamma error

distribution in *lme4* (Bates et al. 2015). A second GLMM with maternal cortisol as the

dependent variable was used to determine the proportion of variation explained by

weight, body condition, the number of days postpartum, season, colony, and an

interaction between season and colony. To account for both structural and data

multicollinearity among weight, condition, and age (pups) / days postpartum (mothers),

these variables were rescaled and centered by subtracting the mean from all observed

values. In preliminary analyses, we tested for the presence of heterogeneous variance by

allowing individual slopes to vary by age (pups) or days postpartum (mothers). For both

models, random intercepts were included for each individual to account for repeated

measures. The residuals of the full models were inspected for linearity and equality of

error variances (using plots of residuals versus fits), normality (using Q – Q plots), and

homogeneity of variance (using Levene's test). A backward elimination based on the chi-

squared statistic was implemented to simplify the full models. The best fitting models

were selected based on AIC values. The statistical significance of fixed predictors was

assessed using Wald tests. We determined the marginal R^2 (variance explained by fixed

effects) and conditional R^2 (variance explained by fixed and random effects) according to

(Nakagawa and Schielzeth 2013).

Estimating intrinsic and extrinsic factors influencing cortisol responsiveness

To determine whether weight, body condition, age, sex, season, colony, and an

interaction between season and colony explained a significant proportion of the variation

in cortisol responsiveness among pups, we fitted a generalized additive model (GAM) in

mgcv (Wood et al. 2016). A second GAM with maternal cortisol responsiveness as the

dependent variable was fitted to determine the proportion of variation explained by

weight, body condition, days postpartum, season, colony, and an interaction between

season and colony. In our dataset, cortisol responsiveness was "heavy-tailed" compared

to a Normal distribution so we used a scaled *t*-distribution. To account for both structural

and data multicollinearity, weight, body condition, and age (pups) / days postpartum

(mothers) were rescaled and centered. Since cortisol response samples were collected

between 8 and 12 minutes after the baseline, in preliminary analyses we tested if

accounting for variation in the amount of time between sampling significantly improved

model fit. In both models, random intercepts were included for each individual to account

for repeated measures. The residuals of the full models were inspected for linearity and

equality of error variances (using plots of residuals versus fits), normality (using Q – Q

plots), and homogeneity of variance (using Levene's test). A backward elimination based

on the *z*-values of explanatory variables was implemented to simplify the models. The

best fitting model for pups and mothers was determined based on AIC values. The

statistical significance of fixed predictors was assessed using Wald tests. Conditional R^2

was calculated according to (Nakagawa and Schielzeth 2013). All analyses and

visualizations were implemented in R version 4.0.2 (R Core Team 2020) using the

integrated development environment RStudio version 1.3.1093 (RStudio Team 2020).

Animal handling, ethics and permits

Sampling was carried out by the BAS under permits from the Government of

South Georgia and the South Sandwich Islands (Wildlife and Protected Areas Ordinance

(2011), RAP permit numbers 2018/024 and 2019/032). The samples were imported into

the UK under permits from the Department for Environment, Food, and Rural Affairs

(Animal Health Act, import license number ITIMP18.1397) and from the Convention on

International Trade in Endangered Species of Wild Fauna and Flora (import numbers

578938/01-15 and 590196/01-18). All procedures used were approved by the BAS

Animal Welfare and Ethics Review Body (AWERB applications 2018/1050 and

2019/1058).

Results

We used a fully crossed, repeated measures design incorporating saliva samples

from 98 unique pups and 93 unique mothers from two colonies of contrasting density

across two consecutive years of contrasting food availability (Figure 1). Sample sizes

were balanced between the colonies (n = 94 from FWB and n = 97 from SSB) and

seasons (n = 96 from 2019 and n = 95 from 2020). Each season, pups were sampled every

20 days from birth until weaning, amounting to a total of 551 analyzed saliva samples

(290 baseline and 221 response samples) or an average of six samples per individual.

Mothers were sampled twice each season, once shortly after birth and again shortly

before molting, which amounted to a total of 240 analyzed saliva samples (145 baseline

and 95 response samples) or an average of three samples per individual.

Cortisol heritability estimates

We estimated heritability using two animal models, the first incorporating known

pedigree relationships (i.e. mother-offspring pairs from both years) and the second

incorporating a SNP relatedness matrix, which was only available for the first year of the

study. For baseline cortisol, narrow-sense heritability (h^2) estimates from both models

were low with overlapping 95% credible intervals (pedigree model: h^2 = 0.013, 95%

highest posterior density 0.004 – 0.045; SNP relatedness model: h^2 = 0.018, 95% highest

posterior density 0.004 – 0.062) (Figure 2). The additive genetic (V_A) and residual (V_R)

variance estimates of the two models were also comparable (pedigree model: 95%

highest posterior density of V_A = 0.1 – 1.6 and V_R = 30.6 – 39.9; SNP relatedness model:

95% highest posterior density of V_A = 0.1 – 2.3 and V_R = 28.5 – 41.7). We obtained

similarly low h^2 estimates for cortisol responsiveness, but the 95% credible intervals were

somewhat broader (pedigree model: h^2 = 0.015, 95% highest posterior density 0.004 –

0.101; SNP relatedness model: h^2 = 0.035, 95% highest posterior density 0.010 – 0.225)

(Figure 2). The V_A and V_R variance estimates of the two models were also comparable

(pedigree model: 95% highest posterior density of V_A = 0.1 – 1.9 and V_R = 16.0 – 22.3;

SNP relatedness model: 95% highest posterior density of V_A = 0.1 – 3.5 and V_R = 9.7 –

17.8).

Intrinsic and extrinsic factors influencing baseline cortisol

The best supported model of pup baseline cortisol contained age ($p < 0.001$),

weight ($p < 0.001$), sex ($p = 0.004$), season ($p = 0.003$), body condition ($p = 0.093$), and

colony ($p = 0.107$) as fixed effects (Table 1a, Figure 3a). The total amount of variance

explained by this model was high (conditional R^2 = 0.657), as was the repeatability of

baseline cortisol values across individuals (ICC = 0.39). Including ID as a random effect

significantly improved the fit of the model, indicating appreciable among-individual

variability in baseline cortisol ($p < 0.001$; Supplementary Table S1a). Allowing

individual slopes to vary between age groups also significantly improved model fit ($p <$

0.001; Supplementary Table S1a) suggesting that individuals responded to the covariates

differently depending on their age. Overall, baseline cortisol decreased with increasing

pup age (Figure 3b) and was lower among pups born in 2020, the year of higher food

availability (Figure 3c). Baseline cortisol decreased significantly as pup weight increased

(Figure 3d), although the slope of the regression between cortisol and weight approached

zero as pups approached their moult. Finally, baseline cortisol tended to be higher in

males than females (Figure 3e).

The best-supported model of maternal baseline cortisol contained days

postpartum ($p < 0.001$), season ($p = 0.004$), and colony ($p = 0.068$) as fixed effects (Table

1b, Figure 3f). Neither weight nor body condition were retained in the model. The total

variance explained by the model was again high (conditional $R^2 = 0.774$), as was

repeatability of baseline cortisol within individuals (ICC = 0.43). Including ID as a

random effect significantly improved model fit, indicating appreciable among-individual

variability in baseline cortisol ($p < 0.001$; Supplementary Table S1b). The concentration

of cortisol in maternal saliva decreased as the season progressed (Figure 3g) and tended

to be higher in 2020, the year of higher food availability (Figure 3h).

Intrinsic and extrinsic factors influencing cortisol responsiveness

For the majority of observations (80.0% in pups and 89.5% in mothers),

individual cortisol levels increased from the first to the second sampling time point,

suggesting that the capture and handling procedure induced a measurable stress response

(Figure 4a, e). Including the amount of time between samples as a random intercept did

not significantly improve the fit of the model (Supplementary Table S2). The best-

supported model of pup cortisol responsiveness contained age ($p < 0.001$) and sex ($p =$

0.010) as fixed effects (Table 2a, Figure 4b). Compared to the baseline cortisol model,

the total amount of explained variance was small (conditional $R^2 = 0.080$). The fixed

effects of the cortisol responsiveness model were also in the opposite direction to the

equivalent fixed effects in the baseline cortisol model, with cortisol responsiveness

increasing with pup age and being higher in female pups (Figure 4c, d). For maternal

cortisol responsiveness, the best-supported model contained only days postpartum ($p =$

0.001) as a fixed effect (Table 2b, Figure 4f). The total amount of explained variance was

again small (conditional $R^2 = 0.056$) and maternal cortisol responsiveness decreased as

the season progressed (Figure 4g).

Discussion

We used a fully crossed, repeated measures design to characterize individual

variation in baseline cortisol levels and their response to capture stress in Antarctic fur

seals. We found that baseline cortisol was only marginally explained by genetic factors,

while high within-individual repeatability and among-individual variability in both pups

and mothers could be largely explained by a combination of intrinsic and extrinsic

factors. For cortisol responsiveness, we found that neither genetic nor environmental

factors explained a substantial proportion of the phenotypic variation. Our results provide

detailed insights into the individualization of an endocrine phenotype in a wild pinniped

population.

Baseline cortisol

We quantified the narrow sense heritability of baseline cortisol using animal

models based on a simple pedigree and a SNP array. The former approach assumes that

individuals of unknown parentage are unrelated to all other individuals in the population,

which can lead to heritability being underestimated (Kruuk 2004). By contrast, genomic

approaches are capable of quantifying unbiased relatedness for all sampled individuals,

but can be time-consuming and costly to produce (Frentiu et al. 2008). Despite these

differences, we found that both approaches produced consistently low heritability

estimates for baseline cortisol in Antarctic fur seals. Heritability can be low because of

low additive genetic variation, high environmental variance, or cross-environment

genetic correlations (i.e. the genetic basis of the trait varies between different

environments) (Charmantier and Garant 2005). Our low estimates, which stand in

contrast to our expectations and previously published empirical results (Jenkins et al.

2014; Stedman et al. 2017; Bairos-Novak et al. 2018), might therefore be reflective of the

extreme heterogeneity of the environmental conditions encountered by the study

population at Bird Island.

For both pups and mothers, we were able to explain a substantial proportion of the

total phenotypic variance in baseline cortisol by including individual-based and

environmental variables in our models (conditional $R^2 = 0.66$ for pups and $R^2 = 0.77$ for

mothers). Similar results have been obtained for a variety of species (e.g. (Joly and

Cameron 2018; Uchida et al. 2021), but see (Azevedo et al. 2019)), suggesting that

contextual and developmental phenotypic plasticity in response to environmental

heterogeneity may be a widespread phenomenon. In addition, we found that baseline

cortisol levels were highly consistent within individuals, suggesting that individualized

endocrine phenotypes become established during early ontogeny and persist at least until

nutritional independence. Baseline cortisol might therefore represent a stable attribute by

which fur seals adapt to spatial or temporal heterogeneity in their environment (Réale and

Dingemanse 2010).

Our models uncovered a strong influence of age and days postpartum on baseline

cortisol levels in pups and mothers, respectively, with salivary cortisol decreasing from

birth to moult. One explanation for this may be the shifting environmental conditions that

individuals experience as the season progresses. Pregnant females arrive ashore in

December and give birth on crowded breeding beaches. Mothers continue to suckle their

pups on the beach until about 30 days postpartum, when most females transition into the

tussock grass that covers most of the island's interior (Doidge et al. 1984). Also during

this time, adult males begin to abandon their territories and migrate to higher latitudes

around the Antarctic ice shelf (Forcada and Staniland 2018). Consequently, the frequency

of unpredictable challenges for both pups and mothers likely declines as the season

progresses.

We detected significantly higher cortisol concentrations among pups and mothers

sampled in 2020, the year of higher food availability. This was surprising given the many

empirical studies that have linked elevated cortisol concentrations to food shortages and

periods of nutritional stress (e.g. (Kitaysky et al. 2007; Behie et al. 2010; Bryan et al.

2013; Garber et al. 2020)). We can think of two possible explanations for this pattern. On

the one hand, our results could be explained by higher population densities in the second

year of our study, as significantly more females bred in 2020 compared to 2019 (Nagel et

al. 2021). This would be in line with the small, albeit non-significant, effect of colony on

baseline cortisol, with hormone concentrations being marginally higher in both pups and

mothers at SSB compared to FWB. On the other hand, circulating levels of cortisol are

essential for the maintenance of metabolic functions (Kuo et al. 2015), and food-induced

cortisol secretions have been documented in the literature (Quigley and Yen 1979;

Gibson et al. 1999; Stimson et al. 2014). Shorter foraging trip durations (Nagel et al.

2021) and consequently more frequent meals for pups and mothers in 2020 may have

resulted in higher average baseline cortisol concentrations, which facilitate protein and

carbohydrate metabolism.

More indicative of the hypothesized correlation between cortisol and nutritional

stress, we found a significant negative relationship between baseline cortisol levels and

weight in pups. Elevated cortisol levels stimulate gluconeogenesis and thereby fat

oxidation, which provides individuals with a source of energy (Wingfield and Romero

2011). Given that pups must tolerate bouts of starvation lasting up to 11 days while their

mothers forage at sea (Forcada and Staniland 2018), our results may reflect a

physiological response to prolonged periods of natural fasting (Ortiz et al. 2001;

Jeanniard du Dot et al. 2009). In other words, fasting pups may increase baseline cortisol

to release energy, resulting in a reduction of absolute body fat and overall weight. This

would also explain why we do not see a similar relationship in mothers, who remain

ashore between foraging trips for as little as 24 hours and on average only two days

(Boyd et al. 1991). Alternatively, pups may be more susceptible to environmental

stressors than adult females, with lighter pups requiring more energy to maintain

homeostasis under, for example, extreme climatic conditions.

In pups, we also uncovered a significant association between baseline cortisol and

sex, with hormone concentrations being moderately higher in males than females.

Previous studies of the sex-specific secretion of cortisol have produced diverse results,

reporting higher concentrations in males (Egyptian mongoose: Azevedo et al. 2019),

females (caribou: Joly and Cameron 2018; marmoset: Garber et al. 2020), or comparable

concentrations in both sexes (grizzly bear: Macbeth et al. 2010). These contrasting

findings highlight the complexity of interactions between cortisol and the sex hormones,

which can vary with the reproductive system, phase, and cycle (Levine 2002). In

addition, the social and environmental stressors associated with growth and reproduction

in each sex are likely to vary among species. For example, male Antarctic fur seal pups

engage more often in social interactions and risk prone behaviors (Jones et al. 2020) such

that the number of "stressful" events encountered may be higher in males than females.

Contrary to our initial expectations, colony only had a marginal, non-significant

effect on baseline salivary cortisol in pups and their mothers. This is in contrast to a

previous study where cortisol concentrations, measured from hair, were higher in mothers

(but not offspring) from the high density colony (Meise et al. 2016). We can think of two

non-mutually exclusive explanations for this discrepancy. First, hair cortisol

concentrations are thought to reflect events in the recent past (hours or days) (Kalliokoski

et al. 2019), whereas salivary cortisol captures circulating hormone levels at the time of

sampling (Lewis 2006). Consequently, hair concentrations of baseline cortisol might

integrate a larger number of stressful events, allowing density-dependent differences to

more readily accumulate. Second, the first study was conducted in 2011 when population

densities were much higher at SSB (Forcada and Hoffman 2014), which may have

accentuated differences between the two colonies.

Cortisol responsiveness

When measuring baseline cortisol in the wild, there is typically little control over

the stimuli that focal individuals experience prior to sampling. By contrast, "capture

stress" protocols elicit a repeatable and uniform acute stress response, providing a

measure of the activity of the HPA axis that can be compared across individuals,

populations, and time points (Wingfield and Romero 2011). Cortisol responsiveness,

measured here as the difference between cortisol concentrations recorded at baseline and

after ten minutes of induced handling stress, thus represents an individual's secretory

capacity and potential to respond to an immediate stressor.

Given its important role in initiating appropriate physiological and behavioral

responses to unpredictable challenges, it has been hypothesized that cortisol

responsiveness may be under stronger selection than baseline cortisol (Jenkins et al.

2014; Bairos-Novak et al. 2018). However, our heritability estimates for cortisol

responsiveness were of a similar magnitude to those obtained for baseline cortisol,

although the associated credible intervals were broader. One factor that may contribute

towards this finding is our experimental design. While our ten-minute capture stress

protocol did elicit a measurable HPA response, peak cortisol concentrations in pinnipeds

may occur anywhere from 30 to 60 minutes after the cessation of stress (Engelhard et al.

2002; Champagne et al. 2012). Consequently, if we could have applied a longer period of

capture stress, we might have obtained more accurate estimates of the HPA response.

This was not possible in the current study due to both logistic constraints and ethical

concerns (Richter and Hintze 2019), but could potentially be investigated in the future.

Although our models did not explain a large proportion of the variation in cortisol

responsiveness, they did reveal significant effects of age and days postpartum, which

were in the opposite direction for pups and mothers. While responsiveness increased

from birth to moult in pups, it decreased over the same period in mothers. For pups, an

increase in cortisol responsiveness over time is likely related to the corresponding

decrease in baseline concentrations. Additionally, pups may be more sensitive to HPA

axis activation or mount a stronger HPA response as the adrenocortical system develops

with age (Walker and Scribner 1991; Engelhard et al. 2002). For mothers, a decrease in

cortisol responsiveness over time might be a reflection of acclimation to capture stress.

However, mothers were captured only twice during the season, whereas pups were

captured on average five times, so we would expect to observe greater habituation in

pups than mothers. Alternatively, exposure to chronic stress has been shown to

downregulate cortisol responses to an acute stressor (Rich and Romero 2005; DeRango et

al. 2019). Maternal provisioning is costly, with offspring survival and growth often being

traded off against parental condition (Smiseth et al. 2013). Prolonged energetic

investment in offspring care might therefore have depressed the ability of mothers to

mount a strong cortisol response to capture and handling stress.

Finally, we also uncovered a significant association between cortisol

responsiveness and pup sex, with the magnitude of the stress response being moderately

higher in females than males. This may be linked to elevated levels of testosterone in

males, which have been shown in other mammals to reduce cortisol responsiveness (e.g.

(Lürzel et al. 2011)). For example, artificially increased testosterone levels in female rats

resulted in decreased activity of the HPA axis in response to stress (Seale et al. 2005).

Consequently, although males have higher baseline levels, cortisol secretion in response

to a stressor may be depressed due to the inhibiting effect of testosterone on the HPA

axis.

Conclusions

The past decade has witnessed a shift in our understanding of individual

differences from a perceived statistical nuisance to a fundamental and compelling aspect

of behavioral ecology and evolutionary biology (Bolnick et al. 2003; Trillmich et al.

2018; Krüger et al. 2021). Our study contributes towards this narrative by decomposing

individual variation in endocrine phenotypes across different life-history stages and

environments in Antarctic fur seal pups and their mothers. Our results are indicative of

substantial contextual and developmental plasticity in the HPA axis, which may help to

facilitate the match between an individual's phenotype and the environment. However,

further research is needed to elucidate both the fitness consequences of endocrine

variability and how this may respond to environmental heterogeneity over longer

timescales.

References

Atkinson S, Arnould JPY, Mashburn KL. 2011. Plasma cortisol and thyroid hormone

concentrations in pre-weaning Australian fur seal pups. Gen Comp Endocrinol.

172:277–281. doi:10.1016/j.ygcen.2011.03.014.

Azevedo A, Bailey L, Bandeira V, Dehnhard M, Fonseca C, Sousa L De, Jewgenow K.

2019. Age, sex and storage time influence hair cortisol levels in a wild mammal

population. PLoS One. 14(8):e0221124. doi:10.1371/journal. pone.0221124.

Bairos-Novak KR, Ryan CP, Freeman AR, Gary Anderson W, Hare JF. 2018. Like

mother, like daughter: Heritability of female Richardson's ground squirrel *Urocitellus*

richardsonii cortisol stress responses. Curr Zool. 64(2):153–163.

doi:10.1093/cz/zox014.

Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models

using lme4. J Stat Softw. 67(1):1–48. doi:10.18637/jss.v067.i01.

Behie AM, Pavelka MSM, Chapman CA. 2010. Sources of variation in fecal cortisol

levels in howler monkeys in Belize. Am J Primatol. 72:600–606.

doi:10.1002/ajp.20813.

Bellavance MA, Rivest S. 2014. The HPA-immune axis and the immunomodulatory

actions of glucocorticoids in the brain. Front Immunol. 5:1–13.

doi:10.3389/fimmu.2014.00136.

Bennett KA, Moss SEW, Pomeroy P, Speakman JR, Fedak MA. 2012. Effects of

handling regime and sex on changes in cortisol, thyroid hormones and body mass in

fasting grey seal pups. Comp Biochem Physiol Part A. 161:69–76.

doi:10.1016/j.cbpa.2011.09.003. 10.1016/j.cbpa.2011.09.003.

Boake CRB. 1989. Repeatability: Its role in evolutionary studies of mating behavior.

Evol Ecol. 3(2):173–182. doi:10.1007/BF02270919.

Bolnick DI, Svanbäck R, Fordyce JA, Yang LH, Davis JM, Hulsey CD, Forister ML.

2003. The ecology of individuals: Incidence and implications of individual

specialization. Am Nat. 161(1):1–28. doi:10.1086/343878.

Bonier F, Martin PR. 2016. How can we estimate natural selection on endocrine traits?

Lessons from evolutionary biology. Proc R Soc B. 283:20161887.

doi:10.1098/rspb.2016.1887.

Boyd IL, Lunn NJ, Barton T. 1991. Time budgets and foraging characteristics of lactating

Antarctic fur seals. J Anim Ecol. 60(2):577–592. doi:10.2307/5299.

Bozovic D, Racic M, Ivkovic N. 2013. Salivary cortisol levels as a biological marker of

stress reaction. Med Arch. 67(5):371–374. doi:10.5455/medarh.2013.67.374-377.

Bryan HM, Darimont CT, Paquet PC, Wynne-Edwards KE, Smits JEG. 2013. Stress and

reproductive hormones in grizzly bears reflect nutritional benefits and social

consequences of a salmon foraging niche. PLoS One. 8(11):e80537.

doi:10.1371/journal.pone.0080537.

Cassini MH. 1999. The evolution of reproductive systems in pinnipeds. Behav Ecol.

10(5):612–616. doi:10.1093/beheco/10.5.612.

Champagne CD, Houser DS, Costa DP, Crocker DE. 2012. The effects of handling and

anesthetic agents on the stress response and carbohydrate metabolism in Northern

elephant seals. PLoS One. 7(5):e38442. doi:10.1371/journal.pone.0038442.

Charmantier A, Garant D. 2005. Environmental quality and evolutionary potential:

Lessons from wild populations. Proc R Soc B. 272(1571):1415–1425.

doi:10.1098/rspb.2005.3117.

Dall SRX, Bell AM, Bolnick DI, Ratnieks FLW. 2012. An evolutionary ecology of

individual differences. Ecol Lett. 15(10):1189–1198. doi:10.1111/j.1461-

0248.2012.01846.x.

DeRango EJ, Prager KC, Greig DJ, Hooper AW, Crocker DE. 2019. Climate variability

and life history impact stress, thyroid, and immune markers in California sea lions

(*Zalophus californianus*) during El Niño conditions. Conserv Physiol. 7:1–15.

doi:10.1093/conphys/coz010.

Dingemanse NJ, Réale D. 2005. Natural selection and animal personality. Behaviour.

142(9):1159–1184. doi:10.1163/156853905774539445.

Doidge DW, Croxall JP, Baker JR. 1984. Density-dependent pup mortality in the

Antarctic fur seal *Arctocephalus gazella* at South Georgia. J Zool London. 202:449–

460. doi:10.1111/j.1469-7998.1984.tb05095.x.

Engelhard GH, Brasseur SMJM, Hall AJ, Burton HR, Reijnders PJH. 2002.

Adrenocortical responsiveness in southern elephant seal mothers and pups during

lactation and the effect of scientific handling. J Comp Physiol B. 172:315–328.

doi:10.1007/s00360-002-0257-0.

Forcada J, Hoffman JI. 2014. Climate change selects for heterozygosity in a declining fur

seal population. Nature. 511(7510):462–465. doi:10.1038/nature13542.

Forcada J, Staniland IJ. 2018. Antarctic Fur Seal. In: Würsig B, Thewissen JGM, Kovacs

KM, editors. Encyclopedia of Marine Mammals. 3rd ed. Academic Press. p. 25–27.

Frentiu FD, Clegg SM, Chittock J, Burke T, Blows MW, Owens IPF. 2008. Pedigree-free

animal models: The relatedness matrix reloaded. Proc R Soc B. 275(1635):639–647.

doi:10.1098/rspb.2007.1032.

Garber PA, McKenney A, Bartling-John E, Bicca-Marques JC, De la Fuente MF, Abreu

F, Schiel N, Souto A, Phillips KA. 2020. Life in a harsh environment: The effects of

age, sex, reproductive condition, and season on hair cortisol concentration in a wild

non-human primate. PeerJ. 8:e9365. doi:10.7717/peerj.9365.

Gastón MS, Vaira M. 2020. Male mating success is related to body condition and stress-

induced leukocyte response in an anuran with scramble competition. Can J Zool.

98(6):391–398. doi:10.1139/cjz-2019-0193.

Gélin U, Coulson G, Festa-Bianchet M. 2016. Heterogeneity in reproductive success

explained by individual differences in bite rate and mass change. Behav Ecol.

27(3):777–783. doi:10.1093/beheco/arv209.

Gentry RL, Holt JR. 1982. Equipment and techniques for handling northern fur seals.

NOAA Tech Rep NMFS SSRF-758.:1–17.

Gibson EL, Checkley S, Papadopoulos A, Poon L, Daley S, Wardle J. 1999. Increased

salivary cortisol reliably induced by a protein-rich midday meal. Psychosom Med.

61(2):214–224. doi:10.1097/00006842-199903000-00014.

Graffelman J, Moreno V. 2013. The mid *p*-value in exact tests for Hardy-Weinberg

equilibrium. Stat Appl Genet Mol Biol. 12(4):433–448. doi:10.1515/sagmb-2012-0039.

Hadfield JD. 2010. MCMC methods for multi-response generalized linear mixed models:

The MCMCglmm R package. J Stat Softw. 33(2):1–22. doi:10.1002/ana.22635.

Harcourt RG, Turner E, Hall A, Waas JR, Hindell M. 2010. Effects of capture stress on

free-ranging, reproductively active male Weddell seals. J Comp Physiol A. 196:147–

154. doi:10.1007/s00359-009-0501-0.

Humble E, Paijmans AJ, Forcada J, Hoffman JI. 2020. A 85K SNP array uncovers

inbreeding and cryptic relatedness in an Antarctic fur seal breeding colony. G3.

10:2787–2799. doi:10.1101/2020.04.01.020123.

Hunt GL, Heinemann D, Everson I. 1992. Distributions and predator-prey interactions of

macaroni penguins, Antarctic fur seals, and Antarctic krill near Bird Island, South

Georgia. Mar Ecol Prog Ser. 86(1):15–30. doi:10.3354/meps086015.

Jeanniard du Dot T, Rosen DAS, Richmond JP, Kitaysky AS, Zinn SA, Trites AW. 2009.

Changes in glucocorticoids, IGF-I and thyroid hormones as indicators of nutritional

stress and subsequent refeeding in Steller sea lions (*Eumetopias jubatus*). Comp

Biochem Physiol Part A. 152:524–534. doi:10.1016/j.cbpa.2008.12.010.

Jenkins BR, Vitousek MN, Hubbard JK, Safran RJ. 2014. An experimental analysis of

the heritability of variation in glucocorticoid concentrations in a wild avian population.

Proc R Soc B. 281:20141302. doi:10.1098/rspb.2014.1302.

Joly K, Cameron MD. 2018. Early fall and late winter diets of migratory caribou in

northwest Alaska. Rangifer. 38(1):27–38. doi:10.7557/2.38.1.4107.

Jones KA, Wood H, Jonathan P, Ratcliffe N, Votier SC, Staniland IJ. 2020. Risk

exposure trade-offs in the ontogeny of sexual segregation in Antarctic fur seal pups.

Behav Ecol.:1–12. doi:10.1093/beheco/araa018.

Kalliokoski O, Jellestad FK, Murison R. 2019. A systematic review of studies utilizing

hair glucocorticoids as a measure of stress suggests the marker is more appropriate for

quantifying short-term stressors. Sci Rep. 9:11997. doi:10.1038/s41598-019-48517-2.

Kitaysky AS, Piatt JF, Wingfield JC. 2007. Stress hormones link food availability and

population processes in seabirds. Mar Ecol Prog Ser. 352:245–258.

doi:10.3354/meps07074.

Krüger O, Anaya-Rojas J, Caspers B, Chakarov N, Elliot-Graves A, Fricke C, Gadau J,

Gossmann T, Hoffman J, Kaiser MI, et al. 2021. Individualised niches: An integrative

conceptual framework across behaviour, ecology, and evolution. EcoEvoRxiv. preprint.

doi:10.32942/osf.io/7h5xq.

Kruuk LEB. 2004. Estimating genetic parameters in natural populations using the

"animal model." Philos Trans R Soc B. 359(1446):873–890.

doi:10.1098/rstb.2003.1437.

Kuo T, McQueen A, Chen T-C, Wang Jen-Chywan. 2015. Regulation of glucose

homeostasis by glucocorticoids. In: Wang J.-C., Harris C, editors. Glucocorticoid

Signaling. Vol. 872. New York: Springer. p. 99–126.

http://www.ncbi.nlm.nih.gov/pubmed/26215998.

Levine JE. 2002. Editorial: Stressing the importance of sex. Endocrinology.

143(12):4502–4504. doi:10.1210/en.2002-221041.

Lewis JG. 2006. Steroid analysis in saliva: An overview. Clin Biochem Rev. 27:139–146.

Lightman SL, Conway-Campbell BL. 2010. The crucial role of pulsatile activity of the

HPA axis for continuous dynamic equilibration. Nat Rev Neurosci. 11(10):710–718.

doi:10.1038/nrn2914.

Lihoreau M, Kaiser S, Resende B, Rödel HG, Châline N. 2021. Editorial: Context-

dependent plasticity in social species: Feedback loops between individual and social

environment. Front Psychol. 12:645191. doi:10.3389/fpsyg.2021.645191.

Lürzel S, Kaiser S, Krüger C, Sachser N. 2011. Inhibiting influence of testosterone on

stress responsiveness during adolescence. Horm Behav. 60:691–698.

doi:10.1016/j.yhbeh.2011.09.007.

Macbeth BJ, Cattet MRL, Stenhouse GB, Gibeau ML, Janz DM. 2010. Hair cortisol

concentration as a noninvasive measure of long-term stress in free-ranging grizzly bears

(*Ursus arctos*): Considerations with implications for other wildlife. Can J Zool.

88:935–949. doi:10.1139/Z10-057.

McCafferty DJ, Boyd IL, Taylor RI. 1998. Diving behaviour of Antarctic fur seal

(*Arctocephalus gazella*) pups. Can J Zool. 76(3):513–520. doi:10.1139/z97-219.

Meise K, Von Engelhardt N, Forcada J, Hoffman JI. 2016. Offspring hormones reflect

the maternal prenatal social environment: Potential for foetal programming? PLoS One.

11(1):e0145352. doi:10.1371/journal.pone.0145352.

Milenkaya O, Catlin DH, Legge S, Walters JR. 2015. Body condition indices predict

reproductive success but not survival in a sedentary, tropical bird. PLoS One.

10(8):e0136582. doi:10.1371/journal.pone.0136582.

Müller C, Caspers BA, Gadau J, Kaiser S. 2020. The power of infochemicals in

mediating individualized niches. Trends Ecol Evol. 35(11):981–989.

doi:10.1016/j.tree.2020.07.001.

Mutwill AM, Schielzeth H, Zimmermann TD, Richter SH, Kaiser S, Sachser N. 2021.

Individuality meets plasticity: Endocrine phenotypes across male dominance rank

acquisition in guinea pigs living in a complex social environment. Horm Behav.

131:104967. doi:10.1016/j.yhbeh.2021.104967.

Nagel R, Stainfield C, Fox-Clarke C, Toscani C, Forcada J, Hoffman J. 2021. Evidence

for an Allee effect in a declining fur seal population. Proc R Soc B. 288:20202882.

doi:https://doi.org/10.1098/rspb.2020.2882.

Nakagawa S, Schielzeth H. 2013. A general and simple method for obtaining R2 from

generalized linear mixed-effects models. Methods Ecol Evol. 4(2):133–142.

doi:10.1111/j.2041-210x.2012.00261.x.

Ortiz RM, Houser DS, Wade CE, Ortiz CL. 2003. Hormonal changes associated with the

transition between nursing and natural fasting in northern elephant seals (*Mirounga*

angustirostris). Gen Comp Endocrinol. 130:78–83. doi:10.1016/S0016-6480(02)00572-

5.

Ortiz RM, Wade CE, Ortiz CL. 2001. Effects of prolonged fasting on plasma cortisol and

TH in postweaned northern elephant seal pups. Am J Physiol - Regul Integr Comp

Physiol. 280:R790–R795. doi:10.1006/gcen.2000.7514.

Pavitt AT, Walling CA, Möstl E, Pemberton JM, Kruuk LEB. 2015. Cortisol but not

testosterone is repeatable and varies with reproductive effort in wild red deer stags. Gen

Comp Endocrinol. 222:62–68. doi:10.1016/j.ygcen.2015.07.009.

Payne MR. 1979. Growth in the Antarctic fur seal *Arctocephalus gazella*. J Zool London.

187:1–20.

Peig J, Green AJ. 2009. New perspectives for estimating body condition from

mass/length data: The scaled mass index as an alternative method. Oikos. 118:1883–

1891. doi:10.1111/j.1600-0706.2009.17643.x.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar

P, de Bakker PIW, Daly MJ, et al. 2007. PLINK: A tool set for whole-genome

association and population-based linkage analyses. Am J Hum Genet. 81(3):559–575.

doi:10.1086/519795.

Quigley ME, Yen SSC. 1979. A mid-day surge in cortisol levels. J Clin Endocrinol

Metab. 49(6):945–947. doi:10.1210/jcem-49-6-945.

R Core Team. 2020. R: A language and environment for statistical computing.

http://www.r-project.org/.

Réale D, Dingemanse NJ. 2010. Personality and individual social specialization. In:

Székely T, Moore AJ, Komdeur J, editors. Social Behaviour: Genes, Ecology and

Evolution. Cambridge University Press. p. 417–441.

Rich EL, Romero LM. 2005. Exposure to chronic stress downregulates corticosterone

responses to acute stressors. Am J Physiol - Regul Integr Comp Physiol. 288:R1628–

R1636. doi:10.1152/ajpregu.00484.2004.

Richter SH, Hintze S. 2019. From the individual to the population – and back again?

Emphasising the role of the individual in animal welfare science. Appl Anim Behav

Sci. 212:1–8. doi:10.1016/j.applanim.2018.12.012.

RStudio Team. 2020. RStudio: Integrated development environment for R.

http://www.rstudio.com/.

Schoenemann KL, Bonier F. 2018. Repeatability of glucocorticoid hormones in

vertebrates: A meta-analysis. PeerJ. 6:e4398. doi:10.7717/peerj.4398.

Seale J V., Wood SA, Atkinson HC, Harbuz MS, Lightman SL. 2005. Postnatal

masculinization alters the HPA axis phenotype in the adult female rat. J Physiol.

563(1):265–274. doi:10.1113/jphysiol.2004.078212.

Sih A, Cote J, Evans M, Fogarty S, Pruitt J. 2012. Ecological implications of behavioural

syndromes. Ecol Lett. 15(3):278–289. doi:10.1111/j.1461-0248.2011.01731.x.

Smiseth PT, Kölliker M, Royle NJ. 2013. What is parental care? In: Royle NJ, Smiseth

PT, Kölliker M, editors. The Evolution of Parental Care. 1st ed. Oxford: Oxford

University Press. p. 1–17.

Stamps JA, Biro PA. 2016. Personality and individual differences in plasticity. Curr Opin

Behav Sci. 12:18–23. doi:10.1016/j.cobeha.2016.08.008.

http://dx.doi.org/10.1016/j.cobeha.2016.08.008.

Stedman JM, Hallinger KK, Winkler DW, Vitousek MN. 2017. Heritable variation in

circulating glucocorticoids and endocrine flexibility in a free-living songbird. J Evol

Biol. 30:1724–1735. doi:10.1111/jeb.13135.

Stimson RH, Mohd-Shukri NA, Bolton JL, Andrew R, Reynolds RM, Walker BR. 2014.

The postprandial rise in plasma cortisol in men is mediated by macronutrient-specific

stimulation of adrenal and extra-adrenal cortisol production. J Clin Endocrinol Metab.

99(1):160–168. doi:10.1210/jc.2013-2307.

Taff CC, Schoenle LA, Vitousek MN. 2018. The repeatability of glucocorticoids: A

review and meta-analysis. Gen Comp Endocrinol. 260:136–145.

doi:10.1016/j.ygcen.2018.01.011.

Trillmich F, Müller T, Müller C. 2018. Understanding the evolution of personality

requires the study of mechanisms behind the development and life history of

personality traits. Biol Lett. 14(2):20170740. doi:10.1098/rsbl.2017.0740.

Uchida K, Ng R, Vydro SA, Smith JE, Blumstein DT. 2021. The benefits of being

dominant: Health correlates of male social rank and age in a marmot. Curr Zool.:1–8.

doi:10.1093/cz/zoab034.

Vessey SH, Drickamer LC. 2010. Integration of proximate and ultimate causes. In: Breed

M, Moore J, editors. Encyclopedia of Animal Behavior. 1st ed. Academic Press. p.

180–185.

Walker CD, Scribner KA. 1991. The pituitary-adrenocortical system of neonatal rats is

responsive to stress throughout development in a time-dependent and stressor-specific

fashion. Endocrinology. 128(3):1385–1395. doi:10.1210/endo-128-3-1385.

Wingfield JC, Romero LM. 2011. Adrenocortical responses to stress and their

modulation in free-living vertebrates. In: Comprehensive Physiology. Hoboken, NJ,

USA: John Wiley & Sons, Inc. p. 211–234.

Wolf M, Weissing FJ. 2012. Animal personalities: Consequences for ecology and

evolution. Trends Ecol Evol. 27(8):452–461. doi:10.1016/j.tree.2012.05.001.

Wood SN, Pya N, Säfken B. 2016. Smoothing parameter and model selection for general

smooth models. J Am Stat Assoc. 111(516):1548–1563.

doi:10.1080/01621459.2016.1180986.

Yang J, Lee SH, Goddard ME, Visscher PM. 2011. GCTA: A tool for genome-wide

complex trait analysis. Am J Hum Genet. 88:76–82. doi:10.1016/j.ajhg.2010.11.011.

 Figure Legends

Figure 1: Location and sampling protocol for the study. (a) Map of Bird Island, South

Georgia, a sub-Antarctic island in the southern Atlantic Ocean. The inset shows an

enlarged view of the two study colonies from which mother-pup pairs were sampled.

Freshwater Beach (FWB, shown in blue) and Special Study Beach (SSB, shown in red)

are separated by approximately 200 meters. (b) We employed a fully crossed sampling

scheme involving the collection of saliva samples from a total of 100 pairs from the two

colonies in two successive breeding seasons, the first of which was coincidentally a year

of particularly low food availability. (c) Each focal mother was sampled twice in a season

while pups were sampled every 20 days from birth until weaning.

Figure 2: Posterior distributions of heritability (h^2) estimates for baseline cortisol and

cortisol responsiveness. A simple pedigree (mother-offspring pairs) for the entire dataset

and a custom 85K SNP array for the 96 individuals sampled in the 2019 season were used

to calculate the relatedness matrix. The modes and highest posterior density intervals of

the posterior distributions are shown in yellow (baseline) and green (responsiveness).

Figure 3: Generalized linear mixed models for pup (a – e) and mother (f – h) baseline

cortisol values. Estimates ± 95% confidence intervals for all fixed effects included in the

best fit models for pups and mothers are shown in panels a and f, respectively. Significant

main effects for both models are shown in panels b – e and panels g – h, respectively.

Boxes show median values ± 75% percentiles with the vertical lines indicating 95%

confidence intervals. Further details of the model output can be found in Table 1.

Figure 4: Generalized linear mixed models for pup (a – d) and mother (e – g) cortisol

responsiveness. The change in an individuals' cortisol level from baseline (t = 0 minutes)

to response (t = 10 ± 2 minutes) sampling is shown for pups and mothers in panels a and

e, respectively. Estimates ± 95% confidence intervals for all fixed effects included in the

best fit models are respectively shown in panels b and f. Significant main effects are also

shown for pups (panels c, d) and mothers (panel g). Boxes show median values ± 75%

percentiles, with the vertical lines indicating 95% confidence intervals. Further details of

the model output can be found in Table 2.

 Tables

Table 1: Parameter estimates from the best fit generalized linear mixed models of (a) pup

and (b) maternal baseline cortisol. Random intercepts were included for each individual

to account for repeated measures. Estimates together with their 95% confidence intervals

(CI) are presented. Significant p-values are in bold. The mean squared error (σ^2), between

group variance ($\tau 00$), Intraclass Correlation Coefficient (ICC; the consistency within an

individual across multiple measurements), the sample size (n) and total number of

observations, as well as the variance explained by the fixed effects (marginal R^2) and

variance explained by both fixed and random effects (conditional R^2) are given.

(a)	Pup baseline cortisol		
Fixed effects	Estimates	CI	p
(Intercept)	1.58	1.40 – 1.75	**<0.001**
Age	-0.29	-0.44 – -0.14	**<0.001**
Season [2020]	0.25	0.09 – 0.41	**0.003**
Weight	-0.36	-0.52 – -0.20	**<0.001**
Sex [male]	0.24	0.08 – 0.41	**0.004**
Body condition	0.14	-0.02 – 0.31	0.093
Colony [SSB]	0.13	-0.03 – 0.29	0.107
Random effect			
σ^2	0.18		
$\tau_{00\ ID}$	0.05		

ICC	0.39
n_{ID}	96
Observations	290
Marginal R^2 / conditional R^2	0.440 / 0.657

(b)	Maternal baseline cortisol		
Fixed effects	Estimates	CI	*p*
(Intercept)	1.39	1.18 – 1.60	**<0.001**
Days postpartum	-0.66	-0.73 – -0.59	**<0.001**
Season [2020]	0.35	0.11 – 0.59	**0.004**
Colony [SSB]	0.22	-0.02 – 0.47	0.068
Random effect			
σ^2	0.16		
$\tau_{00\ ID}$	0.12		
ICC	0.43		
n_{ID}	92		
Observations	145		
Marginal R^2 / conditional R^2	0.603 / 0.774		

Table 2: Parameter estimates from the best fit generalized additive models of (a) pup and
(b) maternal cortisol responsiveness. Random intercepts were included for each
individual to account for repeated measures. Estimates together with their 95%
confidence intervals (CI) are presented. Significant p-values are in bold. The sample size
(n) and total number of observations, as well as the variance explained by the fixed and
random effects in the model (conditional R^2) are given.

(a)	Pup cortisol responsiveness		
Fixed effects	Estimates	CI	p
(Intercept)	2.91	2.26 – 3.55	**<0.001**
Age	0.88	0.41 – 1.35	**<0.001**
Sex [male]	-1.18	-2.07 – -0.28	**0.010**
Random effect	13.42		**0.032**
n $_{ID}$	91		
Observations	221		
Conditional R^2	0.080		
(b)	**Maternal cortisol responsiveness**		
Fixed effects	Estimates	CI	p
(Intercept)	3.33	2.62 – 4.04	**<0.001**
Days postpartum	-0.96	-1.54 – -0.38	**0.001**
Random effect	0.00		0.660
n $_{ID}$	71		
Observations	95		

Conditional R^2 0.056

841 **Figures**

842 Figure 1

(a) Map of Bird Island

(b) Fully crossed design

n = 25 mother-pup pairs per colony per season
n = 100 mother-pup pairs total in the dataset

(c) Repeated measures

843
844

Figure 2

Figure 3

Figure 4

Data Availability Statement

Raw data will be uploaded to an open-access data repository upon acceptance of this

manuscript. R code is available as a Markdown file and included in the Supplementary

Material.

Conflict of Interest

The authors declare that they have no conflict of interest.

Funding

This work was supported by the German Research Foundation (DFG) as part of the SFB TRR

212 (NC³) (project numbers 316099922, 396774617). It was also supported by the priority

program "Antarctic Research with Comparative Investigations in Arctic Ice Areas" SPP 1158

(project number 424119118) and core funding from the Natural Environment Research

Council to the British Antarctic Survey's Ecosystems Program.

Acknowledgements

The authors would like to thank Ana Bertoldi Carneiro, Freya Blockley, Jamie Coleman,

Alexandra Dodds, Vicki Foster, Derren Fox, Iain Angus Gordon, Pauline Goulet, Rosie Hall,

Cary Jackson, Adam Lowndes, Elizabeth Morgan, Rachael Orben, Jessica Ann Philips, David

Reid, and Mark Whiffin for additional help in the field. We kindly thank Sabine Kruse for

conducting the endocrine analyses. We are also grateful to Jonas Schwartz for tips on how to

collect saliva samples from young pups in the field. A special thanks to Océane Salles for

producing the fur seal cartoon in Figure 1c.

Author contributions

RN, CS, CT, and CF-C collected the data. SK advised on field protocols and coordinated

cortisol quantification. RN analyzed the data. AJP generated the SNP genotype dataset and

DLJV built the genomic relatedness matrix. RN and CC performed the molecular laboratory

work. JIH, SK, and JF conceived and developed the project. RN drafted the manuscript. All of

the authors commented on and approved the final manuscript.

50