Artificial reefs geographical location matters more than its age and depth for sessile invertebrate

colonization in the Gulf of Lion (NorthWestern Mediterranean Sea)

Sylvain Blouet[1,2], Lorenzo Bramanti[1], Katell Guizien[1]

[1] CNRS-Sorbonne Université, Laboratoire d'Ecogéochimie des Environnements Benthiques, LECOB,

Observatoire Océanologique de Banyuls Sur Mer, 66650 Banyuls sur Mer, France

[1 2] Ville d'Agde, Aire marine protégée de la côte agathoise, 34300 Agde, France.

ABSTRACT

Artificial reefs (ARs) have been used to support fishing activities. Sessile invertebrates are essential

components of trophic networks within ARs, supporting fish productivity. However, colonization by

sessile invertebrates is possible only after effective larval dispersal from source populations, usually in

natural habitat. We tested the relevance of geographic location, duration of immersion and depth on

ARs colonization processes in the Gulf of Lion. Five species sessile invertebrates species, with

contrasting life history traits and regional distribution in the natural rocky habitat, were inventoried

on ARs deployed during two immersion periods (1985 and 2000-2009) and at different depths. At the

local level, neither depth nor immersion duration differentiated ARs assemblages. At the regional

scale, colonization patterns differed between species, resulting in diverse assemblages.

This study highlights the primacy of geographical positioning over immersion duration and depth in

ARs colonization, suggesting it should be accounted for in maritime spatial planning.

KEY WORDS: ARTIFICIAL REEF, BENTHIC INVERTEBRATES, DEPTH, IMMERSION DURATION,

GEOGRAPHICAL LOCATION, LIFE TRAITS

INTRODUCTION

The decline of fish stocks and natural marine habitat degradation resulting from human exploitation have been documented worldwide for decades (Claudet and Fraschetti, 2010; Jackson, 2001; Pauly et al., 2002).

Artificial reefs (ARs) have been primarily implemented to reduce the pressure of fisheries in coastal areas, complementing other management tools such as marine protected areas or regulatory measures such as fishing licenses (Claudet and Pelletier, 2004; Seaman, 2007; Wilson, 2002). Moreover, ARs could provide economic benefits linked to recreational activities such as recreational fishing and scuba diving (Chen et al., 2013). Beneficial effects such as fish biomass and capture efficiency increase near ARs have been reported (reviewed by Bohnsack and Sutherland, 1985; Tessier et al., 2014). However, the quick colonization by fish and mobile invertebrates (Powers et al., 2003; Relini, 2002; Santos and Monteiro, 2007) led to a debate opposing attraction vs production regarding the effects of ARs on fishery (Grossman et al., 1997). The fish production argument is based on the hypotheses of a better protection against predators and an increase in available substrate area for larval establishment thanks to habitat complexification and an increase of available trophic resource (Bohnsack, 1989). In natural rocky habitats, benthic invertebrates play an essential role in fish trophic networks (Ardizzone et al., 1996; Martens et al., 2006), and ARs trophic network showed similarity with natural rocky habitat one, with dominance of filter-feeders using phytoplanktonic primary production and fish predation on crustacean colonizing the ARs (Cresson, 2013). Moreover, AR deployed in sandy areas are expected to enhance fish productivity given that epifauna secondary production per ARs unit surface has been estimated to be 30 times greater than that of natural sandy infauna (Steimle, 2002). However, those estimates were made shortly after immersion and do not prove the long-term fish production in ARs and supporting this argument would require extending, in the long-term, data on colonization by benthic invertebrates (Svane and Petersen, 2001). Indeed, the age since deployment has been described as a key factor to explain ARs coverage by benthic invertebrates (Svane and Petersen, 2001). The assemblages of benthic communities are expected to

change over time in a succession between pioneer and specialist species (Connell and Slatyer, 1977).

In contrast with pioneer species, specialist ones have slower colonization dynamics, because of their

lower fecundity (Fava et al., 2016). However, after colonization, specialists are expected to

outcompete pioneer species due to their more efficient use of environmental resources (Connell and

Slatyer, 1977). Among these, light availability is an essential factor shaping marine benthic

communities across the water depth gradient (Odum, 1971). Several studies have shown a decrease

in the density of benthic invertebrates with depth on ARs (Lewbel et al., 1986.; Moura et al., 2007;

Shinn and Wicklund, 1989; van der Stap et al., 2016) explained by the decrease in light intensity (Relini

et al., 1994). The structural complexity has also been put forward as important characteristics linked

to ARs efficiency in ecological restoration (Strain et al., 2018). Structural complexity increases available

surface for colonization and niches diversity with various shelter and light exposure conditions, the

latter being related to different benthic assemblage compositions (Glasby, 2000; T. M. Glasby, 1999)

and higher productivity (Vivier et al., 2021). The recent 3D printing techniques using concrete, allow

the design of ARs mimicking natural habitats (Ly et al., 2021). However, those studies concerned short-

term colonization (<3.5 years) (Wendt et al., 1989) hence based on pioneer species with high dispersal

capacities which colonization is likely mainly regulated by post-settlement processes such as

competition, predation and physical disturbance (Todd, 1998). In contrast to mobile species,

sessile benthic invertebrates can only colonize reefs after larval dispersal which is limited by

reproduction frequency (Thorson, 1950). Colonization implies thus an effective dispersal between

natural areas and ARs, which depends on source population spatial distribution, species fecundity,

dispersive larval traits and ocean circulation. Nevertheless, until now dispersal drivers have been

disregarded while colonization disparities among ARs may result from differences in both larval

connectivity (which in turn depends on fecundity, dispersal capacities and adult distribution in the

natural habitat) and post-recruitment processes.

The objective of the present study was to test the hypothesis that the geographical location of ARs

deployment with respect to the natural habitat can condition ARs colonization in the long term. To this

aim we investigated the effects of local (depth and immersion duration) and regional (geographic area)

factors on the presence and abundance of five species of sessile invertebrates with different life history

traits, endemic to the Gulf of Lion (GDL) (Northwestern Mediterranean Sea) and frequently found on

natural hard substrates. Data were collected on ARs immersed for more than 10 years in the GDL and

assemblage composition analysis at three nested spatial scales (~ 1kms; ~5kms; >30 kms) were

combined.

MATERIALS AND METHODS

<u>**Study area and spatially stratified sampling design**</u>

The study area extended along 160 km of the GDL coastline (Figure 1). The GDL is a wide micro-tidal

continental shelf dominated by soft-bottom habitat with few small rocky habitat patches of less than

20 km^2. The GDL is a homogeneous and isolated hydrodynamic unit (Rossi et al., 2014), delimited by

the northern current (Millot, 1990).

Between 1985 and 2009, 763 ARs with different shapes or material and a total volume of 37 575 m^3

(Tessier et al., 2015) have been deployed along the GDL coastline over 66 km^2 of state concessions

(Cepralmar, 2015) between 10 and 35 m depth (Figure 1, Blouet et al., 2021). The ARs deployed in GDL

represent 40% of the total AR volume in France (Tessier et al., 2015).

Deployment was performed during three major time periods: 1985, 1992-1999 and 2004-2009, but for

the present study we examined only the oldest (1985) and the youngest (after 2000) ones. For the

analysis of data, we followed a stratified sampling design. To this aim, the GDL coastline was regularly

divided into 6 geographical sectors separated by a distance ranging from 12 to 117 km, with a median

value of 49 km (AGM, AGD, VLR, GRU, LEU, CST, Figure 1). Each sector included ARs deployed during

either the first (1985, CST geographical sector), the last (2004-2009, AGM, VLR, GRU, LEU geographical

sectors) or both periods of immersion (AGD). In each sector, two sites separated by a distance ranging

from 2.1 to 11.7 km (median value of 7.5 km) were defined, except in AGD where 5 sites were defined.

In each site, three sampling units separated by a distance between 4m and 3.6 Km (median distance

of 251 m) were set out by pooling neighboring ARs reaching a minimum surface of 89 m^2 per sampling

unit and totalling a minimum developed surface of 306 m^2 per site. Such large continuous sampling

units in each site aimed at limiting the effect of recruitment spatial variability over distances from 100s

meters to kilometers arising from the non-uniformity of the flow of larvae (Daigle et al., 2014; Glasby,

2000; Simpson et al., 2017; Smale, 2012). Such a spatial scale is consistent with the spatial scale of flow

homogeneity obtained in simulations over GDL soft-bottom habitat (Briton et al., 2018). This inventory

methodology enabled us to test for the existence of structuring factors at the local and regional scale.

In the geographical sector GRU, in one of the two sites, only two sampling units could be defined given

the ARs shapes, yielding a total surface of 600 m^2 (Supplementary Material Table 1). Developed reef

surface was calculated on the basis of technical specifications data present in the state concession

documents taking into account only the colonizable surface (surfaces in contact with the sediment

were excluded).

Colonization assessment

Assessment of ARs colonization was carried out by autonomous scuba-diving in 2020 by counting the

number of individuals of the five target species in the 44 sampling units (Supplementary Material

Table1, totalling 80 ARs fully inventoried). Among the species listed in previous ARs inventories in the

GDL, we selected five species that were present in most inventories, easy to identify by scuba diving

and spanning different phyla with contrasting life-history traits (Créocean, 2003 & 2004; Table 1). We

selected two gorgonians *Eunicella singularis* (Esper, 1791) and *Leptogorgia sarmentosa* (Esper, 1789),

one bryozoan *Pentapora fascialis* (Pallas, 1766), one annelida *Sabella spallanzanii* (Gmelin, 1791) and

one ascidian *Halocynthia papillosa* (Linnaeus, 1797) (Figure 2). The five species have a similar wide

natural repartition area along European coasts ranging from 1m to 250m depth (Giangrande et al.,

2005; Gori et al., 2011; Ponti et al., 2019; Turon, 1990; Weinberg and Weinberg, 1979). In addition, *S.*

spallanzanii has been recorded along the coasts of Brazil, Australia and New Zealand where it is

classified as an invasive non-indigenous species (Currie et al., 2000).

The five species are present in the rocky habitat of the NW Mediterranean Sea (Laubier, 1966; True,

1970; Hong, 1980). However, in the GDL, where natural rocky habitat covers uneven surfaces within

the 6 geographical sectors (from 3,123 10^7 m^2 for the AGM sector to 5 10^5 m^2 for the LEU sector), the

five species display different spatial distributions (Dutrieux et al., 2005; Dalias et al., 2011; Padron et

al., 2018; S. Blouet personal observation). *E. singularis* is frequently observed and abundant

throughout the GDL (from the AGM sector to the CST sector), while *L. sarmentosa,* less abundant, is

present mainly in the center of the GDL (AGD, VLR, LEU, CST). *P. fascialis* is abundant in the west of the

GDL (AGD, LEU, CST, and south of CST). The distribution of *H. papillosa* is not well known, however the

species has been observed in all the rocky areas of the GDL. *S. spallanzanii* is present but rare in natural

rocky habitat. Nevertheless, *S. spallanzanii* is very abundant in lagoons, ports and marinas of the GDL

(S. Blouet personal observation) which have been indicated as preferred habitat of the species (Currie

et al., 2000).

The five species display different life-history traits. All five species reproduce once a year in different

seasons and with different strategies (Table 1).

S. spallanzani reproduces in January-February, when water temperature is the coldest. The species

displays multiple reproductive strategies: internal fertilization, with larvae brooded either inside or

outside the mineral tube secreted around the body, and external fertilization broadcast spawning

(Giangrande et al., 2000). In addition, asexual reproduction by fission has been reported (Read et al.,

2011). *S. spallanzani* releases lecithotrophic larvae with a planktonic larval duration (PLD) of about 4

weeks (Giangrande et al., 2000). It's life span can exceed 5 years, with sexual maturity after one year

(Giangrande and Petraroli 1994; Giangrande et al., 2000). Like most gorgonians, *E. singularis* releases

lecithotrophic larvae in early summer (June to August). Even though larval competency period can

reach up to 2 months (Guizien et al., 2020; Zelli et al., 2020), PLDs ranging from 7 to 14 days best

explain gene flow among *E. singularis* natural populations dwelling in the fragmented rocky habitat of

the GDL (Padron et al. 2018). *E. singularis* life span can reach 25-30 years with sexual maturity before

6 years (Gori et al., 2007; Weinberg and Weinberg, 1979) (Weinberg and Weinberg 1979, Gori et al.

2007).

The other gorgonian, *L. sarmentosa* also releases lecithotrophic larvae but in the late summer

(September to October) and the PLD is unknown (Rossi and Gili 2009). *L. sarmentosa* life span can

reach 20 years with female sexual maturity within 2-3 years after settlement (Rossi and Gili 2009).

H. papillosa is a simultaneous hermaphrodite which releases larvae in late summer (September-

October; Becerro and Turon, 1992), presumably lecithotrophic. The PLD of *H. papillosa* larvae is

unknown but PLD shorter than 12 hours has been consistently reported for other solitary ascidian

species (Ayre et al., 1997). We did not find any data about the age at sexual maturity and the life span

of *H. papillosa*. However, the ascidians are considered as highly invasive, particularly because of their

rapid growth and early sexual maturity (Zhan et al., 2015), with some species such as C*iona*

intestionalis complex and *Ciona savigniy,* reaching sexual maturity at the age of 2 months (Zhan et al.,

2015) and continuous spawning (Carver et al. 2003).

P. fascialis displays both sexual and asexual reproduction. During sexual reproduction, most bryozoans

release lecithotrophic larvae which settle after a few minutes or a few hours, rarely beyond several

days (Keough, 1983). *P. fascialis* larval release has been inferred to happen in June based on

recruitment observations (Cocito et al., 1998a). Asexual reproduction happens by colony

fragmentation or budding extension (Cocito et al., 1998b). Individual life span is estimated to be about

10 years with early sexual maturity after 2 years (Cocito et al., 1998b).

Statistical   analysis

We examined to which extent ARs colonization is affected by the location and timing of deployment.

To do so, we tested the effect of 4 factors (geographical sector being the regional factor, site being the

local factor, depth and age) on the presence/absence of the five species and on the dissimilarity

between their co-occurrence assemblages. In all analyses, site was a random factor, nested either in

year, age or the geographical sector factor. Due to AR deployment set up, the effect of factors age (2

levels, 1985, 2002-2009) and depth (2 levels, >20 m, <20 m depth) was tested within the AGD

geographical sector only. Factor age was tested in 3 sites at <20m and factor depth was tested in 4

sites deployed in 2009. To avoid any confounding effect due to age or depth, the regional factor vs

local factor were tested on the 5 geographical sectors where AR were deployed during the 2002-2009

immersion phase and at <20 m depth only (AGM, AGD, VLR, GRU, LEU: 5x2 levels). A Jaccard similarity

matrix was built on presence/absence data across all pairwise sampling units. Two multivariate

analyses were performed. Non-parametric multivariate analysis of variance with permutation was

applied to test for the effects of age, depth and geographical sector on species assemblages (NP-

manova: Anderson, 2001; Zar, 1999). Another multivariate analysis was performed to cluster most

similar species assemblages in the sector of AGD (SIMPROF : Clarke et al., 2008).

When significant differences between the five species co-occurrence assemblages were detected for

a factor, a non-parametric univariate analysis (ANOVA Kruskall-Wallis) was performed for each species

independently to detect the species driving the difference. Fisher post-hoc test was used to identify

the site where the difference arose. A same p-value of 0.05 was taken for detecting significant

differences. Analyses were performed with Matlab software using the Fathom package for multivariate

analyses (Jones, 2014) and the Matlab statistics toolbox for univariate analyses.

RESULTS

Artificial reefs colonization by the five target species at **regional scale**

Among the five target species, *S. spallanzanii* was the only one whose presence was recorded in all the

sampling units and geographical sectors (Figure 4). In only one out of 15 sites, it was the only species

detected. *H. papillosa* was detected in all geographic sectors but not in all the sampling units. *L.

sarmentosa* was detected in five of the six geographical sectors (not present in AGM) and *P. fascialis*

was detected in 3 of the 6 geographical sectors (CST, LEU and AGD). Finally, *E.singularis* was the least

frequently observed species, being detected in only three geographical sectors (AGM, AGD and LEU).

In all geographical sectors, at least three of the five target species were detected. Assemblages of two

species were found in only one site out of 15 (sector AGD), assemblages of three species were found

in 8 sites, assemblages of four species were found in 2 sites and assemblages of five species were found

in 3 sites.

Age and depth effects on five target species at local scale

Despite all five target species being detected on ARs in the AGD sector, assemblage composition

among sites differed (Figure 3). Clustering of sampling units within the 5 sites (A, B, C, D, E) in AGD

identified 2 clusters (SIMPROF: P <0.05; Figure 4). The two sites (B and C) with same age (2009), depth

range (less than 20 m) and reef shape (pipe) were attributed to different clusters.

In fact, one cluster grouped ARs of different age at a same depth (1985 in site A and 2009 in site B)

while the other cluster grouped ARs of the same age but at different depths (less than 20 m in site C

and more than 20 m in sites D and E). In both clusters, different AR shapes were found (steel cage and

pipes in one cluster, pipes, Comin and Bonna in the other cluster, Figure 4). The geographic distance

between the two clusters (A-B) and (C-D-E) was 7.5 km and the median value of the intra-cluster

geographic distance was 3 km.

Multivariate analysis of variance confirmed that neither age (NP-MANOVA, F=1.43 P<0.05 ; Table 2)

nor depth (NP-MANOVA, F=1.37 P<0.05 ; Table 3) explained site differences in the five species

assemblages found on ARs in AGD (P<0.05 ; Tables 2 and 3). Differences among the five sites in AGD

were due to different ARs colonization by three species (Kruskall-wallis: *E. singularis*, *P. fascialis*, *H.*

papillosa: all P<0.05; Table 4). Site C differed from other sites by the presence of *E. singularis*, and sites

D and E differed from sites A and B due to the presence of *P. fascialis* and *H. papillosa* (Post-hoc tests).

Geographical effect on five target species at local and regional scales

The five species co-occurrence assemblages on ARs deployed in the same period and at same depth

were significantly different at both regional and local scales (NP-MANOVA : geographical sector F=3.19

P<0.05; site (geographical sector) F=3.64 P<0.05; Table 5). These differences were due to different

colonization of ARs by three of the five species, *E. singularis*, *L. sarmentosa* and *P. fascialis*. For the

latter two species, regional differences (Kruskall-wallis: geographical sector P=0.0002) were more

significant than local differences (Kruskall-wallis: site P=0.001 for *L. sarmentosa* and P=0.003 for *P.*

fascialis; Table 6). Both species were not detected in the north of the GDL (AGM). In contrast, for *E.*

singularis, local differences (Kruskall-wallis: site P=0.01) were more significant than regional ones

(geographical sector P=0.036; Table 6), the species being detected in geographical sectors in the north,

center and south of GDL.

DISCUSSION

The study of effective integration of ARs into the rocky habitat network through their colonization by

species building up their natural biodiversity is essential to assess their ecosystem functions, prior and

after deployment. In the present study, we showed that ARs spread along the GDL coastline had been

colonized by five species found in the natural rocky habitat of the region, ten years after their

deployment. However, spatial colonization patterns differed among species, resulting in diverse

assemblages in different geographical sectors of the GDL. Locally, neither age, immersion depth or reef

shape significantly affected colonization patterns.

Colonization of ARs are expected to evolve toward a stable state comparable to that of the natural

environment, through the succession of opportunistic species (wide dispersal, high fertility, low

tolerance of reduced resource levels, short life-spans, minimal dietary specialisation) followed by

specialized species (limited dispersal, slow growth to a large size at maturity, delayed and limited

reproduction, optimization to reduced resources and long life-spans; (Platt and Connell, 2003; Faurie

et al., 2003). Monitoring of ARs short-term colonization (<3 years) have indeed shown a dominance of

pioneer species (hydroids, serpulids, barnacles and bivalves), most of them having life history traits

typical of opportunistic species (Fariñas-Franco et al., 2013; T. Glasby, 1999; Moura et al., 2007;

Pamintuan and Ali, 1994; Ponti, 2015; Relini et al., 1994; Spagnolo et al., 2014; Toledo et al., 2020).

Long-term studies confirmed successions in ARs colonization (Burt et al., 2011; Nicoletti et al., 2007;

Perkol-Finkel and Benayahu, 2005; Whomersley and Picken, 2003), but none have described saturation

(Svane and Petersen, 2001). In the Tyrrhenian Sea, Nicolleti et al., (2007) described colonization in 5

distinct phases: (i) A first recruitment by pioneer species (hydroids, serpulids, barnacles and bivalves)

during the first months after immersion, followed by phases of (ii) cover dominance, (iii) regression

and (iv) absence of *Mytilus galloprovincialis* for more than 10 years.The installation of diverse bio-

builders bryozoans characteristic of the natural environment was recorded after 20 years only (v). Our

study shows that bio-builder engineering species such as bryozoans (*P. fascialis*) and gorgonians (*E.*

singularis, L. sarmentosa) colonized ARs as early as 10 years after their deployment, without significant

difference between 10 years and 35 years old ARs. However, the presence of *S. spallanzanii* described

as an opportunistic species (sexual precocity, various reproduction modes, rapid growth, short lived;

Giangrande et al., 2000) on all ARs independently of their age of deployment suggests that ARs did not

yet reach a stable state comparable to the natural environment. Thus, the presence of bio-builders is

not a sufficient indicator of the ARs naturalization to the local biodiversity.

The GDL's ARs being located in the sandy coastal zone are likely regularly disturbed by sediment

deposits due to river delivery or/and their resuspension by either trawling activities or the mechanical

action of the swell (Dufois et al., 2014; Durrieu de Madron et al., 2005; Ulses et al., 2008). Testing the

impact of swell and sediment deposit on ARs requires exploring the colonization of ARs along a

gradient of depth and distance from the coast (van der Stap et al., 2016). However, current ARs

deployment in the GDL ranged from 15 to 30 m depth and within 3 miles from the coast and did not

allow testing for differential effect of sediment disturbances as swell impact occurs every year in this

area (Guizien, 2009). Testing the impact of sediment disturbance on ARs colonization would require

exploring reefs deployed deeper than 50 m, such as the anchorages of the floating wind farm that will

be placed in the GDL in the next future (https://info-efgl.fr/le-projet/le-parc/#). Light is also expected

to be an important factor structuring benthic assemblages, along a depth gradient in natural and

artificial environments (T. Glasby, 1999a; T. M. Glasby, 1999b; Svane and Petersen, 2001) . Absence

of depth effect in the present study, although in the GDL light intensity strongly attenuates within the

upper 30 m of the water column (Durrieu de Madron et al., 2011) was potentially a bias due to the five

species selected in the present study whose distributions are not strongly structured by light intensity.

Another factor which has been shown to drive the intensity of ARs colonization is structural complexity

(see Bohnsack and Sutherland, 1985 for a review). Nevertheless, there is no consensus about the

relationship between complexity and subtidal benthic invertebrates abundance, due to potential bias

in controlling the surface and scale in ARs of different complexity (Rouse et al., 2019; Strain et al.,

2018). The similarity in the 5 species co-occurrence between different reef shapes at the same depth

found in the present study suggests structural complexity is less important than the geographical

location in AR colonization by benthic invertebrates.

Benthic invertebrate assemblages result from complex processes that operate at multiple spatial and

temporal scales (Smale, 2012). At the regional scale, larval availability can become a major factor

explaining colonization success. Change in the composition of assemblages during the early

colonization of artificial substrates by benthic invertebrates has been attributed to the availability and

abundance of larvae during the seasons rather than a sequence of distinct succession (Bramanti et al.,

2003; Turner and Todd, 1993). The larval behaviour (buoyancy and motility) and the characteristics of

the biological cycle of the species (spawning timing and PLD) can play a key role in determining the

dispersal distance (Todd, 1998), and consequently the possibility to reach habitat suitable for

settlement. Dispersion distance is generally correlated with PLD, thus, a species with a long PLD is

supposed to colonize habitats further away than species with a shorter PLD (Shanks, 2009). In this

study, the five species were chosen among different phyla known for their contrasting planktonic

durations, swimming abilities and larval dispersal periods, although these larval traits are only known

accurately for *E. singularis* (Guizien et al., 2020; Zelli et al., 2020). *P. fascialis* and *H. papillosa*, the two

species with short PLD (<24h and <48, respectively) colonized ARs located close to the natural habitat

where they are present (< 4.8 km and <10 km, respectively). The coastal circulation of the GDL allows

such dispersal distance over periods of a few days (Guizien et al., 2012). *S. spallanzanii*, which has a

PLD of 3 weeks, colonized all the inventoried ARs, in line with a dispersal distance of 40 km after 3

weeks (Guizien et al., 2012). In contrast, *E. singularis* did not colonize all ARs within geographical

sectors of 30 km width where the species is present in the natural habitat, although a 2-week PLD was

expected to enable such dispersal (Padron et al., 2018). The other gorgonian species, *L. sarmentosa*

colonized more ARs located within distances of less than 30 km from its natural habitat than *E.*

singularis while the PLD of the two species are presumably the same. This suggests that other factors

influence the colonization failure of *E. singularis*.

Reproductive traits are another key to the success in colonising new settings (Stearns, 2000). In this

regard, *E. singularis* colonization potential could be limited by its low fecundity (~25-40 larvae.cm^{-1} of

colony branch, Ribes et al., 2007; Theodor, 1967) compared to that of *L. sarmentosa* (~75 larvae.cm^{-1}

of colony branch, (Rossi et al., 2011; Rossi and Gili, 2009). The wide colonization of ARs by *S.*

spallanzanii is in line with its reproductive traits typical of opportunistic species (early sexual maturity,

high fecundity with more than 50 000 eggs per female, Currie et al., 2000, a fertilization close to 100%,

Giangrande et al., 2000). Since arriving in the Pacific Ocean, *S. spallanzanii* has been declared one of

the ten priority pest species in the marine environment by the Australian authorities and classified as

an invasive species (Hayes et al., 2005). Similarly to *S. spallanzanii*, *H. papillosa* colonized nearly all ARs

located within its 10-km dispersal distance from the natural habitat. Within the ascidian class, a wide

disparity in species fecundity has been reported (Pandian, 2018). This suggests *H. papillosa*

reproductive traits would be close to those of invasive ascidians (Zhan et al., 2015).

Ultimately, understanding ARs colonization requires a precise mapping of source populations in the

natural environment. To this respect, the abundance of *S. spallanzan*i on ARs is surprising, as the

species is not abundant in the natural rocky habitat of the GDL. For this species, other sources of larval

supply than natural settings should be considered, such as the numerous ports and marinas along the

coast of the GDL, as *S. spallanzanii* is very tolerant to environmental conditions (Currie et al., 2000). In

this case of intense colonization by an endemic benthic invertebrate species, ARs apparently extended

its metapopulation, acting as stepping stones for further larval dispersal beyond its natural current

colonization limits (Bishop et al., 2017; Wang et al., 2020). In the GDL, the 14 500 m^3 of ARs deployed

30 years ago are now decommissioned and the relevance of their removal is currently debated.

Connectivity between natural populations has been shown to support species resilience after

disturbances in fragmented habitat, and could be extended to ARs (Fahrig, 2003). However, ARs may

also facilitate the spread of non-indigenous species introduced with maritime traffic in ports (Glasby

et al., 2007). The present study advocates accounting for the geographical arrangement in planning

ARs deployment to enhance fish productivity while avoiding the spread of invasive species.

*Acknowledgement*s. This work was funded by the Agence de l'Eau Rhône-Méditerranée-Corse under

project ICONE - Impacts actuels et potentiels de la CONnectivité Ecologique ajoutée par les récifs

artificiels sur la biodiversité fixée des substrats durs du Golfe du Lion (PI, K. Guizien, AAP 2016). The

authors gratefully acknowledge the helpful assistance during sampling of the staff of the Aire Marine

Protégée Agathoise.

Authors contributions

SB and KG conceived the study, SB carried out sampling and statistical analysis. All contributed to

manuscript writing.

LITERATURE CITED

Anderson, M., 2001. A new method for non-parametric multivariate analysis of variance. Austral
Ecology 26, 32–46. https://doi.org/10.1111/j.1442-9993.2001.01070.pp.x
Ardizzone, G.D., Belluscio, A., Gravina, M.F., Somaschini, A., 1996. Colonization and Disappearance of
Mytilus galloprovincialis Lam. on an Artificial Habitat in the Mediterranean Sea. Estuarine,
Coastal and Shelf Science 43, 665–676. https://doi.org/10.1006/ecss.1996.0095
Ayre, D.J., Davis, A.R., Billingham, M., Llorens, T., Styan, C., 1997. Genetic evidence for contrasting
patterns of dispersal in solitary and colonial ascidians. Marine Biology 130, 51–61.
https://doi.org/10.1007/s002270050224

Becerro, M.A., Turon, X., 1992. Reproductive Cycles of the Ascidians Microcosmus sabatieri and
Halocynthia papillosa in the Northwestern Mediterranean. Marine Ecology 13, 363–373.
https://doi.org/10.1111/j.1439-0485.1992.tb00360.x
Bishop, M.J., Mayer-Pinto, M., Airoldi, L., Firth, L.B., Morris, R.L., Loke, L.H.L., Hawkins, S.J., Naylor,
L.A., Coleman, R.A., Chee, S.Y., Dafforn, K.A., 2017. Effects of ocean sprawl on ecological
connectivity: impacts and solutions. Journal of Experimental Marine Biology and Ecology 492,
7–30. https://doi.org/10.1016/j.jembe.2017.01.021
Blouet, S., Quittet, L., Agin, G., Thorin, S., Dalias, N., Marobin, D., Lenfant, P., Guizien, K., 2021.
Database of the location and typology of artificial reefs in the Gulf of Lion (NW
Mediterranean Sea). https://doi.org/10.17882/81407
Bohnsack, J.A., 1989. Are High Densities of Fishes at Artificial Reefs the Result of Habitat Limitation or
Behavioral Preference? Bulletin of marine science 44, 15.
Bohnsack, J.A., Sutherland, D.L., 1985. Artificial reef research. A review with recommendations for
future priorities. Bulletin of marine science 37, 30.
Bramanti, L., Magagnini, G., Santangelo, G., 2003. Settlement and recruitment: The first stages in the
life cycle of two epibenthic suspension feeders (*Corallium rubrum* and *Anomia ephippium)*.
Italian Journal of Zoology 70, 175–178. https://doi.org/10.1080/11250000309356512
Briton, F., Cortese, D., Duhaut, T., Guizien, K., 2018. High-resolution modelling of ocean circulation
can reveal retention spots important for biodiversity conservation. Aquatic Conserv: Mar
Freshw Ecosyst 28, 882–893. https://doi.org/10.1002/aqc.2901
Burt, J., Bartholomew, A., Sale, P.F., 2011. Benthic development on large-scale engineered reefs: A
comparison of communities among breakwaters of different age and natural reefs. Ecological
Engineering 37, 191–198. https://doi.org/10.1016/j.ecoleng.2010.09.004
Carver, CE., Chisholm, A., Mallet, AL., 2003. Strategies to mitigate the impact of Ciona intestinalis (L.)
biofouling on shellfish production. J Shellfish Res 22:621–631
Cépralmar, Région Languedoc-Roussillon., 2015. Guide pratique d'aide à l'élaboration, l'exploitation
et la gestion des récifs artificiels en Languedoc-Roussillon : 236 pages.
Chen, J.-L., Chuang, C.-T., Jan, R.-Q., Liu, L.-C., Jan, M.-S., 2013. Recreational Benefits of Ecosystem
Services on and around Artificial Reefs: A Case Study in Penghu, Taiwan. Ocean & Coastal
Management 85, 58–64. https://doi.org/10.1016/j.ocecoaman.2013.09.005
Clarke, K.R., Somerfield, P.J., Gorley, R.N., 2008. Testing of null hypotheses in exploratory community
analyses: similarity profiles and biota-environment linkage. Journal of Experimental Marine
Biology and Ecology 366, 56–69. https://doi.org/10.1016/j.jembe.2008.07.009
Claudet, J., Fraschetti, S., 2010. Human-driven impacts on marine habitats: A regional meta-analysis
in the Mediterranean Sea. Biological Conservation 143, 2195–2206.
https://doi.org/10.1016/j.biocon.2010.06.004
Claudet, J., Pelletier, D., 2004. Marine protected areas and artificial reefs: A review of the
interactions between management and scientific studies. Aquat. Living Resour. 17, 129–138.
https://doi.org/10.1051/alr:2004017
Cocito, Silvia, Ferdeghini, F., Sgorbini, S., 1998a. Pentapora fascialis (Pallas) [Cheilostomata:
Ascophora] colonization of one sublittoral rocky site after sea-storm in the northwestern
Mediterranean, in: Baden, S., Phil, L., Rosenberg, R., Strömberg, J.-O., Svane, I., Tiselius, P.
(Eds.), Recruitment, Colonization and Physical-Chemical Forcing in Marine Biological Systems.
Springer Netherlands, Dordrecht, pp. 59–66. https://doi.org/10.1007/978-94-017-2864-5_5
Cocito, S., Sgorbini, S., Bianchi, C.N., 1998b. Aspects of the biology of the bryozoan Pentapora
fascialis in the northwestern Mediterranean. Marine Biology 131, 73–82.
https://doi.org/10.1007/s002270050298
Connell, J.H., Slatyer, R.O., 1977. Mechanisms of Succession in Natural Communities and Their Role in
Community Stability and Organization. The American Naturalist 111, 1119–1144.
https://doi.org/10.1086/283241
CREOCEAN., 2003. Impact des récifs artificiels sur le milieu marin et la pêche professionnelle dans le
golfe d'Aigues- Mortes, CREOCEAN, France, 53 pp.

CREOCEAN., 2004. Suivi scientifique des récifs artificiels immergés au large de Gruissan—Rapport Annuel 2003— Peuplements benthiques, CREOCEAN, France, 32 pp.

Currie, D.R., McArthur, M.A., Cohen, B.F., 2000. Reproduction and distribution of the invasive European fanworm Sabella spallanzanii (Polychaeta: Sabellidae) in Port Phillip Bay, Victoria, Australia. Marine Biology 136, 645–656. https://doi.org/10.1007/s002270050724

Daigle, R.M., Metaxas, A., deYoung, B., 2014. Fine-Scale Distribution and Spatial Variability of Benthic Invertebrate Larvae in an Open Coastal Embayment in Nova Scotia, Canada. PLOS ONE 9, e106178. https://doi.org/10.1371/journal.pone.0106178

Dalias, N., Astruch, P., Foulquié, M., Rouanet, E., 2012. Suivi à long terme de l'impact potentiel de la plongée sous-marine sur les peuplements d'invertébrés benthiques au sein de la Réserve Naturelle Marine de Cerbère-Banyuls (2007 - 2011) – Rapport final. Contrat Conseil Général des Pyrénées Orientales & SEANEO - GIS Posidonie. SEANEO - GIS Posidonie publ. Fr. : 136 P

Dufois, F., Verney, R., Le Hir, P., Dumas, F., Charmasson, S., 2014. Impact of winter storms on sediment erosion in the Rhone River prodelta and fate of sediment in the Gulf of Lions (North Western Mediterranean Sea). Continental Shelf Research 72, 57–72. https://doi.org/10.1016/j.csr.2013.11.004

Durrieu de Madron, X., Ferré, B., Le Corre, G., Grenz, C., Conan, P., Pujo-Pay, M., Buscail, R., Bodiot, O., 2005. Trawling-induced resuspension and dispersal of muddy sediments and dissolved elements in the Gulf of Lion (NW Mediterranean). Continental Shelf Research 25, 2387–2409. https://doi.org/10.1016/j.csr.2005.08.002

Durrieu de Madron, X., Guieu, C., Sempéré, R., Conan, P., Cossa, D., D'Ortenzio, F., Estournel, C., Gazeau, F., Rabouille, C., Stemmann, L., Bonnet, S., Diaz, F., Koubbi, P., Radakovitch, O., Babin, M., Baklouti, M., Bancon-Montigny, C., Belviso, S., Bensoussan, N., Bonsang, B., Bouloubassi, I., Brunet, C., Cadiou, J.-F., Carlotti, F., Chami, M., Charmasson, S., Charrière, B., Dachs, J., Doxaran, D., Dutay, J.-C., Elbaz-Poulichet, F., Eléaume, M., Eyrolles, F., Fernandez, C., Fowler, S., Francour, P., Gaertner, J.C., Galzin, R., Gasparini, S., Ghiglione, J.-F., Gonzalez, J.-L., Goyet, C., Guidi, L., Guizien, K., Heimbürger, L.-E., Jacquet, S.H.M., Jeffrey, W.H., Joux, F., Le Hir, P., Leblanc, K., Lefèvre, D., Lejeusne, C., Lemé, R., Loÿe-Pilot, M.-D., Mallet, M., Méjanelle, L., Mélin, F., Mellon, C., Mérigot, B., Merle, P.-L., Migon, C., Miller, W.L., Mortier, L., Mostajir, B., Mousseau, L., Moutin, T., Para, J., Pérez, T., Petrenko, A., Poggiale, J.-C., Prieur, L., Pujo-Pay, M., Pulido-Villena, Raimbault, P., Rees, A.P., Ridame, C., Rontani, J.-F., Ruiz Pino, D., Sicre, M.A., Taillandier, V., Tamburini, C., Tanaka, T., Taupier-Letage, I., Tedetti, M., Testor, P., Thébault, H., Thouvenin, B., Touratier, F., Tronczynski, J., Ulses, C., Van Wambeke, F., Vantrepotte, V., Vaz, S., Verney, R., 2011. Marine ecosystems' responses to climatic and anthropogenic forcings in the Mediterranean. Progress in Oceanography 91, 97–166. https://doi.org/10.1016/j.pocean.2011.02.003

Dutrieux, E., Thorin, S., Jouvenel, J.-Y., 2005. 100 belles plongées en Languedoc-Roussillon, Edition GAP, 336 p.

Fahrig, L., 2003. Effects of Habitat Fragmentation on Biodiversity. Annu. Rev. Ecol. Evol. Syst. 34, 487–515. https://doi.org/10.1146/annurev.ecolsys.34.011802.132419

Fariñas-Franco, J.M., Allcock, L., Smyth, D., Roberts, D., 2013. Community convergence and recruitment of keystone species as performance indicators of artificial reefs. Journal of Sea Research 78, 59–74. https://doi.org/10.1016/j.seares.2012.10.008

Fava, F., Ponti, M., Abbiati, M., 2016. Role of Recruitment Processes in Structuring Coralligenous Benthic Assemblages in the Northern Adriatic Continental Shelf. PLOS ONE 11, e0163494. https://doi.org/10.1371/journal.pone.0163494

Giangrande, A., Petraroli, A., 1994. Observations on reproduction and growth of Sabella spallanzanii (Polychaeta, Sabellidae) in the Mediterranean Sea. In: Dauvin JC, Laubier L, Reish DJ (eds) Actes de la 4éme Conférence Internationale des Polychetes. Mus Hist nat Paris 162: 51±56

Giangrande, A., Cavallo, A., Licciano, M., Mola, E., Pierri, C., Trianni, L., 2005. Utilization of the filter
feeder polychaete Sabella. Aquacult Int 13, 129–136. https://doi.org/10.1007/s10499-004-
9025-3
Giangrande, A., Licciano, M., Pagliara, P., Gambi, M.C., 2000. Gametogenesis and larval development
in Sabella spallanzanii (Polychaeta: Sabellidae) from the Mediterranean Sea. Marine Biology
136, 847–861. https://doi.org/10.1007/s002279900251
Glasby, T.M., 1999a. Interactive effects of shading and proximity to the seafloor on the development
of subtidal epibiotic assemblages. Mar. Ecol. Prog. Ser. 190, 113–124.
https://doi.org/10.3354/meps190113
Glasby, T.M., 1999b. Effects of shading on subtidal epibiotic assemblages. Journal of Experimental
Marine Biology and Ecology 234, 275–290. https://doi.org/10.1016/S0022-0981(98)00156-7
Glasby, T.M., Connell, S., Holloway, M., Hewitt, C., 2007. Nonindigenous biota on artificial structures:
Could habitat creation facilitate biological invasions? Marine Biology 151, 887–895.
https://doi.org/10.1007/s00227-006-0552-5
Glasby, T.M., 2000. Surface composition and orientation interact to affect subtidal epibiota. Journal
of Experimental Marine Biology and Ecology 248, 177–190. https://doi.org/10.1016/S0022-
0981(00)00169-6
Gori, A., Linares, C., Rossi, S., Coma, R., Gili, J.-M., 2007. Spatial variability in reproductive cycle of the
gorgonians Paramuricea clavata and Eunicella singularis (Anthozoa, Octocorallia) in the
Western Mediterranean Sea. Mar Biol 151, 1571–1584. https://doi.org/10.1007/s00227-006-
0595-7
Gori, A., Rossi, S., Berganzo, E., Pretus, J.L., Dale, M.R.T., Gili, J.-M., 2011. Spatial distribution patterns
of the gorgonians Eunicella singularis, Paramuricea clavata, and Leptogorgia sarmentosa
(Cape of Creus, Northwestern Mediterranean Sea). Mar Biol 158, 143–158.
https://doi.org/10.1007/s00227-010-1548-8
Grossman, G.D., Jones, G.P., Seaman, W.J., 1997. Do Artificial Reefs Increase Regional Fish
Production? A Review of Existing Data 22, 7.
Guizien, K., 2009. Spatial variability of wave conditions in the Gulf of Lions (NW Mediterranean Sea).
Vie Milieu 11.
Guizien, K., Belharet, M., Marsaleix, P., Guarini, J.M., 2012. Using larval dispersal simulations for
marine protected area design: Application to the Gulf of Lions (northwest Mediterranean).
Limnology and Oceanography 57, 1099–1112. https://doi.org/10.4319/lo.2012.57.4.1099
Guizien, K., Viladrich, N., Martínez-Quintana, Á., Bramanti, L., 2020. Survive or swim: different
relationships between migration potential and larval size in three sympatric Mediterranean
octocorals. Scientific Reports 10, 18096. https://doi.org/10.1038/s41598-020-75099-1
Hayes, K.R., Australia, Department of the Environment and Heritage, CSIRO (Australia), Division of
Marine Research, 2005. National priority pests. Part II, Part II,. Australian Government Dept.
of the Environment and Heritage, Parkes A.C.T.
Hong, J.S., 1980. Étude faunistique d'un fond de concrétionnement de type coralligène soumis à un
gradient de pollution en Méditerranée nord-occidentale (Golfe de Fos). Thèse de Doctorat.
Université d'Aix -Marseille II. 134 pp.

Jackson, J.B.C., 2001. Historical Overfishing and the Recent Collapse of Coastal Ecosystems. Science
293, 629–637. https://doi.org/10.1126/science.1059199
Jones, D. L., 2014. The Fathom Toolbox for Matlab: multivariate ecological and oceanographic data
analysis. College of Marine Science, University of South Florida, St. Petersburg, Florida, USA.
Available from: http://www.marine.usf.edu/user/djones/

Keough, M.J., 1983. Patterns of recruitment of sessile invertebrates in two subtidal habitats. Journal
of Experimental Marine Biology and Ecology 66, 213–245. https://doi.org/10.1016/0022-
0981(83)90162-4
Laubier, L., 1966. Le coralligène des Albères. Monographie biocénotique. Ann. Inst. Océanogr. : 137-
316.

Lewbel, G.S., Howard, R.L., Gallaway, B.J., n.d. Zonation of Dominant Fouling Organisms on Northern
Gulf of Mexico Petroleum Platforms 26.
Ly, O., Yoris-Nobile, A.I., Sebaibi, N., Blanco-Fernandez, E., Boutouil, M., Castro-Fresno, D., Hall, A.E.,
Herbert, R.J.H., Deboucha, W., Reis, B., Franco, J.N., Teresa Borges, M., Sousa-Pinto, I., van
der Linden, P., Stafford, R., 2021. Optimisation of 3D printed concrete for artificial reefs:
Biofouling and mechanical analysis. Construction and Building Materials 272, 121649.
https://doi.org/10.1016/j.conbuildmat.2020.121649
Martens, K., Queiroga, H., Cunha, M.R., Cunha, A., Moreira, M.H., Quintino, V., Rodrigues, A.M.,
Seroôdio, J., Warwick, R.M. (Eds.), 2006. Marine Biodiversity: Patterns and Processes,
Assessment, Threats, Management and Conservation. Springer Netherlands, Dordrecht.
https://doi.org/10.1007/1-4020-4697-9
Millot, C., 1990. The Gulf of Lions' hydrodynamics. Continental Shelf Research 10, 885–894.
https://doi.org/10.1016/0278-4343(90)90065-T
Moura, A., Boaventura, D., Cúrdia, J., Carvalho, S., da Fonseca, L.C., Leitão, F.M., Santos, M.N.,
Monteiro, C.C., 2007. Effect of depth and reef structure on early macrobenthic communities
of the Algarve artificial reefs (southern Portugal). Hydrobiologia 580, 173–180.
https://doi.org/10.1007/s10750-006-0455-y
Nicoletti, L., Marzialetti, S., Paganelli, D., Ardizzone, G.D., 2007. Long-term changes in a benthic
assemblage associated with artificial reefs, in: Relini, G., Ryland, J. (Eds.), Biodiversity in
Enclosed Seas and Artificial Marine Habitats, Developments in Hydrobiology. Springer
Netherlands, Dordrecht, pp. 233–240. https://doi.org/10.1007/978-1-4020-6156-1_20
Odum, E.P., n.d. An understanding of ecological succession provides a basis for resolving man's
conflict with nature. 164, 9.
Padron, M., Costantini, F., Baksay, S., Bramanti, L., Guizien, K., 2018. Passive larval transport explains
recent gene flow in a Mediterranean gorgonian. Coral Reefs 37.
https://doi.org/10.1007/s00338-018-1674-1
Pamintuan, I.S., Ali, P.M., n.d. Early Successional Patterns of Invertebrates in Artificial Reefs
Established at Clear and Silty Areas in Bolinao, Pangasinan, Northern Philippines 11.
Pandian, T.J., 2018. Reproduction and Development in Echinodermata and Prochordata. CRC Press,
Boca Raton. https://doi.org/10.1201/9780815364733
Pauly, D., Christensen, V., Guénette, S., Pitcher, T.J., Sumaila, U.R., Walters, C.J., Watson, R., Zeller,
D., 2002. Towards sustainability in world fisheries. Nature 418, 689–695.
https://doi.org/10.1038/nature01017
Perkol-Finkel, S., Benayahu, Y., 2005. Recruitment of benthic organisms onto a planned artificial reef:
shifts in community structure one decade post-deployment. Marine Environmental Research
59, 79–99. https://doi.org/10.1016/j.marenvres.2004.03.122
Platt, W.J., Connell, J.H., 2003. Natural Disturbances and Directional Replacement of Species.
Ecological Monographs 73, 507–522. https://doi.org/10.1890/01-0552
Ponti, M., 2015. Benthic assemblages on artificial reefs in the northwestern Adriatic Sea: Does
structure type and age matter? Marine Environmental Research 10.
Ponti, M., Turicchia, E., Costantini, F., Gori, A., Bramanti, L., Di Camillo, C., Linares, C., Rossi, S.,
Abbiati, M., Garrabou, J., Cerrano, C., 2019. Mediterranean gorgonian forests: Distribution
patterns and ecological roles.
Powers, S., Grabowski, J., Peterson, C., Lindberg, W., 2003. Estimating enhancement of fish
production by offshore artificial reefs: uncertainty exhibited by divergent scenarios. Mar.
Ecol. Prog. Ser. 264, 265–277. https://doi.org/10.3354/meps264265

Read, G., Inglis, G., Stratford, P., Ahyong, S., 2011. Arrival of the alien fanworm Sabella spallanzanii
(Gmelin, 1791) (Polychaeta: Sabellidae) in two New Zealand harbours. AI 6, 273–279.
https://doi.org/10.3391/ai.2011.6.3.04
Relini, G., 2002. Trophic relationships between fishes and an artificial reef. ICES Journal of Marine
Science 59, S36–S42. https://doi.org/10.1006/jmsc.2002.1212
Relini, G., Zamboni, N., Tixi, F., Torchia, G., n.d. Patterns of Sessile Macrobenthos Community
Development on an Artificial Reef in the Gulf of Genoa (Northwestern Mediterranean) 27.
Ribes, M., Coma, R., Rossi, S., Micheli, M., 2007. Cycle of gonadal development in Eunicella singularis
(Cnidaria: Octocorallia): trends in sexual reproduction in gorgonians: Gametogenic cycle in a
temperate gorgonian. Invertebrate Biology 126, 307–317. https://doi.org/10.1111/j.1744-
7410.2007.00101.x
Rossi, S., Gili, J.-M., 2009. Reproductive features and gonad development cycle of the soft bottom-
gravel gorgonian *Leptogorgia sarmentosa* (Esper, 1791) in the NW Mediterranean Sea.
Invertebrate Reproduction & Development 53, 175–190.
https://doi.org/10.1080/07924259.2009.9652304
Rossi, S., Gili, J.-M., Garrofé, X., 2011. Net negative growth detected in a population of Leptogorgia
sarmentosa: quantifying the biomass loss in a benthic soft bottom-gravel gorgonian. Mar Biol
158, 1631–1643. https://doi.org/10.1007/s00227-011-1675-x
Rossi, V., Ser-Giacomi, E., López, C., Hernández-García, E., 2014. Hydrodynamic provinces and oceanic
connectivity from a transport network help designing marine reserves. arXiv:1407.6920
[physics]. https://doi.org/10.1002/2014GL059540
Rouse, S., Porter, J.S., Wilding, T.A., n.d. Artificial reef design affects benthic secondary productivity
and provision of functional habitat 9.
Santos, M.N., Monteiro, C.C., 2007. A fourteen-year overview of the fish assemblages and yield of the
two oldest Algarve artificial reefs (southern Portugal) 7.
Seaman, W., 2007. Artificial habitats and the restoration of degraded marine ecosystems and
fisheries, in: Relini, G., Ryland, J. (Eds.), Biodiversity in Enclosed Seas and Artificial Marine
Habitats, Developments in Hydrobiology. Springer Netherlands, Dordrecht, pp. 143–155.
https://doi.org/10.1007/978-1-4020-6156-1_13
Shanks, A., 2009. Barnacle settlement versus recruitment as indicators of larval delivery. II. Time-
series analysis and hypothesized delivery mechanisms. Mar. Ecol. Prog. Ser. 385, 217–226.
https://doi.org/10.3354/meps08002
Shinn, E.A., Wicklund, R.I., n.d. Artificial Reef Observations from a Manned Submersible off Southeast
Florida 10.
Simpson, T.J.S., Smale, D.A., McDonald, J.I., Wernberg, T., 2017. Large scale variability in the
structure of sessile invertebrate assemblages in artificial habitats reveals the importance of
local-scale processes. Journal of Experimental Marine Biology and Ecology 494, 10–19.
https://doi.org/10.1016/j.jembe.2017.05.003
Smale, D.A., 2012. Extreme spatial variability in sessile assemblage development in subtidal habitats
off southwest Australia (southeast Indian Ocean). Journal of Experimental Marine Biology
and Ecology 438, 76–83. https://doi.org/10.1016/j.jembe.2012.10.002
Spagnolo, A., Cuicchi, C., Punzo, E., Santelli, A., Scarcella, G., Fabi, G., 2014. Patterns of colonization
and succession of benthic assemblages in two artificial substrates. Journal of Sea Research
88, 78–86. https://doi.org/10.1016/j.seares.2014.01.007
Stearns, S.C., n.d. Life history evolution: successes, limitations, and prospects 11.
Steimle, F., 2002. Benthic macrofauna productivity enhancement by an artificial reef in Delaware
Bay, USA. ICES Journal of Marine Science 59, S100–S105.
https://doi.org/10.1006/jmsc.2002.1268
Strain, E.M.A., Olabarria, C., Mayer-Pinto, M., Cumbo, V., Morris, R.L., Bugnot, A.B., Dafforn, K.A.,
Heery, E., Firth, L.B., Brooks, P.R., Bishop, M.J., 2018. Eco-engineering urban infrastructure
for marine and coastal biodiversity: Which interventions have the greatest ecological
benefit? J Appl Ecol 55, 426–441. https://doi.org/10.1111/1365-2664.12961

Svane, I., Petersen, J.K., 2001. On the Problems of Epibioses, Fouling and Artificial Reefs, a Review.
Marine Ecology 22, 169–188. https://doi.org/10.1046/j.1439-0485.2001.01729.x
Tessier, A., Francour, P., Charbonnel, E., Dalias, N., Bodilis, P., Seaman, W., Lenfant, P., 2015.
Assessment of French artificial reefs: due to limitations of research, trends may be
misleading. Hydrobiologia 753, 1–29. https://doi.org/10.1007/s10750-015-2213-5
Tessier, A., Verdoit-Jarraya, M., Blouet, S., Dalias, N., Lenfant, P., 2014. A case study of artificial reefs
as a potential tool for maintaining artisanal fisheries in the French Mediterranean Sea. Aquat.
Biol. 20, 255–272. https://doi.org/10.3354/ab00563
Theodor, J., 1967. Contribution à l'études des gorgones (VII): écologie et comportement de la
planula. Vie Milieu, 18 (2A): 291-301.
Thorson, G., 1950. Reproductive and larval ecology of marine bottom invertebrates. Biological
Reviews 25, 1–45. https://doi.org/10.1111/j.1469-185X.1950.tb00585.x
Todd, C.D., 1998. Larval supply and recruitment of benthic invertebrates: do larvae always disperse
as much as we believe? Hydrobiologia 375, 1–21. https://doi.org/10.1023/A:1017007527490
Toledo, M.-I., Torres, P., Díaz, C., Zamora, V., López, J., Olivares, G., 2020. Ecological succession of
benthic organisms on niche-type artificial reefs. Ecol Process 9, 38.
https://doi.org/10.1186/s13717-020-00242-9
True, M.A., 1970. Étude quantitative de quatre peuplements sciaphiles sur substrat rocheux dans la
région marseillaise. Bulletin Institut Océanographique Monaco, 69(1401): 1- 48.

Turner, S.J., Todd, C.D., 1993. The early development of epifaunal assemblages on artificial substrata
at two intertidal sites on an exposed rocky shore in St. Andrews Bay, N.E. Scotland. Journal of
Experimental Marine Biology and Ecology 166, 251–272. https://doi.org/10.1016/0022-
0981(93)90223-B
Turon, X., 1990. Distribution and Abundance of Ascidians from a Locality on the Northeast Coast of
Spain. Marine Ecology 11, 291–308. https://doi.org/10.1111/j.1439-0485.1990.tb00385.x
Ulses, C., Estournel, C., Durrieu de Madron, X., Palanques, A., 2008. Suspended sediment transport in
the Gulf of Lions (NW Mediterranean): Impact of extreme storms and floods. Continental
Shelf Research 28, 2048–2070. https://doi.org/10.1016/j.csr.2008.01.015
van der Stap, T., Coolen, J.W.P., Lindeboom, H.J., 2016. Marine Fouling Assemblages on Offshore Gas
Platforms in the Southern North Sea: Effects of Depth and Distance from Shore on
Biodiversity. PLoS ONE 11, e0146324. https://doi.org/10.1371/journal.pone.0146324
Vivier, B., Dauvin, J.-C., Navon, M., Rusig, A.-M., Mussio, I., Orvain, F., Boutouil, M., Claquin, P., 2021.
Marine artificial reefs, a meta-analysis of their design, objectives and effectiveness. Global
Ecology and Conservation e01538. https://doi.org/10.1016/j.gecco.2021.e01538
Wang, W., Wang, J., Choi, F.M.P., Ding, P., Li, X., Han, G., Ding, M., Guo, M., Huang, X., Duan, W.,
Cheng, Z., Chen, Z., Hawkins, S.J., Jiang, Y., Helmuth, B., Dong, Y., 2020. Global warming and
artificial shorelines reshape seashore biogeography. Global Ecol Biogeogr 29, 220–231.
https://doi.org/10.1111/geb.13019
Weinberg, S., Weinberg, F., 1979. The Life Cycle of a Gorgonian: Eunicella Singularis (Esper, 1794).
Contributions to zoology Bijdragen tot de dierkunde 48, 127–140.
https://doi.org/10.1163/26660644-04802003
Wendt, P.H., Knott, D.M., Dolah, R.F.V., n.d. COMMUNITY STRUCTURE OF THE SESSILE BIOTA ON FIVE
ARTIFICIAL REEFS OF DIFFERENT AGES 18.
Whomersley, P., Picken, G.B., 2003. Long-term dynamics of fouling communities found on offshore
installations in the North Sea. J. Mar. Biol. Ass. 83, 897–901.
https://doi.org/10.1017/S0025315403008014h
Wilson, K., 2002. Restoration of Hong Kong fisheries through deployment of artificial reefs in marine
protected areas. ICES Journal of Marine Science 59, S157–S163.
https://doi.org/10.1006/jmsc.2002.1186
Zar, J., 1999. Biostatistical analysis.

Zelli, E., Quéré, G., Lago, N., Di Franco, G., Costantini, F., Rossi, S., Bramanti, L., 2020. Settlement
dynamics and recruitment responses of Mediterranean gorgonians larvae to different
crustose coralline algae species. Journal of Experimental Marine Biology and Ecology 530–
531, 151427. https://doi.org/10.1016/j.jembe.2020.151427
Zhan, A., Briski, E., Bock, D.G., Ghabooli, S., MacIsaac, H.J., 2015. Ascidians as models for studying
invasion success. Mar Biol 162, 2449–2470. https://doi.org/10.1007/s00227-015-2734-5

Figure captions:

Figure 1: Map showing the layout of the 6 geographical sectors and 15 sites where artificial reef
sampling was carried out.

Figure 2: Photographs of the five species inventoried on AR *a) Sabella spallanzanii b) Eunicella*
singularis c) Leptogorgia sarmentosa d) Halocynthia papillosa e) *Pentapora fascialis; all © Blouet*
sylvain

Figure 3: Map showing the five species co-occurence assemblages inventoried on ARs in the 15 sites
in the Gulf of Lion.

Figure 4: Composite showing in the upper part the diversity of species assemblage in the sampling
units of the five sites (A, B, C, D, E) in AGD sector together with the type of AR, depth and years of
deployment and in the lower part, the dendrogram obtained by group average clustering based on
the Jaccard dissimilarity index using the presence /absence of species (P=0.04 at 62% of dissimilarity).
The red dotted line delineates the two clusters identified by the analysis.

Figures :

Figure 1 :

Figure 2 :

A)   **B)**   **C)**

D)   **E)**

752   **Figure 3 :**

760 **Figure 4:**

Table captions

Table 1: Larval traits (Planktonic larval duration (PLD), reproduction period, and larval trophic regime)

for the five species inventoried on ARs.

Table 2: Results of multivariate NP-Manova testing the interactive effects of the year (of deployment

and site (nested in year) on the presence/absence assemblage. Significant ($P<0.05$) values in bold.

Table 3: Results of multivariate NP-Manova testing the interactive effects of the depth (of

deployment) and site (nested in depth) on the presence/absence assemblage. Significant ($P<0.05$)

values in bold.

Table 4: Results of univariate Kruskall_wallis testing the effects of year and depth of deployment on

the presence/absence by species. Significant ($P<0.05$) values in bold.

Table 5: Results of multivariate NP-Manova testing the interactive effects of geographical sector and

site (nested in geographical sector) on the presence/absence assemblage. Significant ($P<0.05$) values

in bold.

Table 6: Results of univariate Kruskall_wallis testing the effects of geographical sector and site on the

presence/absence by species. Significant ($P<0.05$) values in bold.

Tables:

Table 1:

Species	Spawning period	Larval type	Larval duration	Ref
Leptogorgia sarmentosa	June - August	Lecithotrophic	? (supposed 1-4 weeks)	(Rossi and Gili 2009)
Eunicella singularis	July - August	Lecithotrophic	1-4 weeks (in aquarium)	(Guizien et al. 2020)
Pentapora fascialis	June	Lecithotrophic	? (supposed <1 days)	(Cocito et al. 1998)
Sabella spallanzani	January - February	Lecithotrophic	21 days (in aquarium)	(Giangrande et al. 2000)
Halocynthia papillosa	September - October	? lecithotrophic	? (supposed <2 days)	(Becerro and Turon 1992)

789   **Table 2:**

Multivariate measures					
Source	Df	SS	Ms	F	Pvalue
Assemblage (Presence/Absence)					
Year	1	0.4825	0.4825	1.4330	0.3330
Site(Year)	1	0.3367	0.3367	7.5551	**0.0130**
Residual	6	0.2674	0.0446		
Total	8	1.0866			

791   **Table 3:**

Multivariate measures					
Source	Df	SS	Ms	F	Pvalue
Assemblage (Presence/Absence)					
Depth	1	0.2491	0.2491	1.3715	0.3280
Site(Depth)	2	0.3633	0.1816	8.5969	**0.001**
Residual	8	0.1690	0.0211		
Total	11	0.7816			

797 **Table 4:**

Univariate measures					
Presence/absence					
	Df	SS	Ms	Chi_sq	Pvalue
Eunicella singularis					
	4	127.5	31.875	10.82	**0.0287**
Leptogorgia sarmentosa					
	4	60	15	6.22	0.1832
Pentapora fascialis					
	4	172.5	43.125	11.5	**0.0215**
Sabella spallazanii					
	4	0	0	nan	nan
Halocynthia papillosa					
	4	150	37.5	11.2	**0.0244**

800 **Table 5:**

Multivariate measures					
Source	**Df**	**SS**	**Ms**	**F**	**Pvalue**
Assemblage (Presence/Absence)					
Geographical sector	4	1.9921	0.4980	3.1961	0.0380
Site(Geographical sector)	5	0.7791	0.1558	3.6434	0.0010
Residual	19	0.8125	0.0427		
Total	28	3.5838			

Table 6:

Presence/absence					
Source	Df	SS	Ms	Chi_sq	Pvalue
Eunicella singularis					
Geographical sector	4	447.08	111.77	10.28	**0.036**
Site	9	937.66	104.18	21.56	**0.0104**
Leptogorgia sarmentosa					
Geographical sector	4	965.7	241.42	22.2	**0.0002**
Site	9	1218	135.33	28	**0.001**
Pentapora fascialis					
Geographical sector	4	937.66	234.41	21.56	**0.0002**
Site	9	1077.83	119.75	24.78	**0.0032**
Sabella spallazanii					
Geographical sector	4	0	0	nan	nan
Site	9	0	0	nan	nan
Halocynthia papillosa					
Geographical sector	4	323.59	80.89	6.01	0.1985
Site	9	561.87	62.43	10.43	0.3166