Artificial reefs geographical location matters more than shape, age and depth for sessile invertebrate

colonization in the Gulf of Lion (NorthWestern Mediterranean Sea)

Sylvain Blouet[1,2], Lorenzo Bramanti[1], Katell Guizien[1]

[1] CNRS-Sorbonne Université, Laboratoire d'Ecogéochimie des Environnements Benthiques, LECOB,

Observatoire Océanologique de Banyuls Sur Mer, 66650 Banyuls sur Mer, France

[2] Ville d'Agde, Aire marine protégée de la côte agathoise, 34300 Agde, France.

ABSTRACT

Artificial reefs (ARs) have been used to support fishing activities. Sessile invertebrates are essential

components of trophic networks within ARs, supporting fish productivity. However, colonization by

sessile invertebrates is possible only after effective larval dispersal from source populations, usually in

natural habitat. While most studies focused on short term colonization by pioneer species, we propose

to test the relevance of geographic location, shape, duration of immersion and depth on the ARs long

term colonization by species found in natural stable communities in the Gulf of Lion. We recorded the

presence of five sessile invertebrates species, with contrasting life history traits and regional

distribution in the natural rocky habitat, on ARs with different shapes deployed during two immersion

time periods (1985 and the 2000s) and in two depth ranges (<20m and >20m). At the local level

(~5kms), neither shape, depth nor immersion duration differentiated ARs assemblages. At the regional

scale (>30kms), colonization patterns differed between species, resulting in diverse assemblages. This

study highlights the primacy of geographical positioning over shape, immersion duration and depth in

ARs colonization, suggesting it should be accounted for in maritime spatial planning.

KEY WORDS: ARTIFICIAL REEF, BENTHIC INVERTEBRATES, SHAPE, DEPTH, IMMERSION DURATION,

GEOGRAPHICAL LOCATION, LIFE TRAITS

INTRODUCTION

The decline of fish stocks and natural marine habitat degradation resulting from human exploitation have been documented worldwide for decades (Claudet and Fraschetti, 2010; Jackson, 2001; Pauly et al., 2002).

Artificial reefs (ARs) have been primarily implemented to reduce the pressure of fisheries in coastal areas, complementing other management tools such as marine protected areas or regulatory measures such as fishing licenses (Claudet and Pelletier, 2004; Seaman, 2007; Wilson, 2002). Moreover, ARs could provide economic benefits such as recreational and traditional fishing and scuba diving (Chen et al., 2013). Beneficial effects such as increase in fish biomass and capture efficiency near ARs have been reported (reviewed by Bohnsack and Sutherland, 1985; Tessier et al., 2014) but led to a debate on the effects of ARs fishery, opposing attraction vs production (Grossman et al., 1997). The fish attraction argument is based on the quick colonization by fish and mobile invertebrates (Powers et al., 2003; Relini, 2002; Santos and Monteiro, 2007). The fish production argument is based on the hypotheses of a better protection against predators and an increase in available substrate area for larval establishment thanks to habitat complexification and an increase of available trophic resource (Bohnsack, 1989). In natural rocky habitats, benthic invertebrates play an essential role in fish trophic networks (Ardizzone et al., 1996; Martens et al., 2006), and ARs trophic network showed similarity with natural rocky habitat, with dominance of filter-feeders using phytoplanktonic primary production and fish predation on crustacean colonizing the ARs (Cresson, 2013). Moreover, AR deployed in sandy areas are expected to enhance fish productivity given that epifauna secondary production per ARs unit surface has been estimated to be 30 times greater than that of natural sandy infauna (Steimle, 2002). However, those estimates were made shortly after immersion and do not prove the long-term fish production in ARs and supporting this argument would require extending, data on colonization by benthic invertebrates in the long-term (Svane and Petersen, 2001). Indeed, age since deployment has been described as a key factor to explain ARs coverage by benthic invertebrates (Svane and Petersen, 2001). The assemblages of benthic communities are expected to change over time in a succession

between pioneer and specialist species (Connell and Slatyer, 1977). In contrast with pioneer species,

specialist ones have slower colonization dynamics, because of their lower fecundity (Fava et al., 2016).

However, after colonization, specialists are expected to outcompete pioneer species due to their more

efficient use of environmental resources (Connell and Slatyer, 1977). Among these, light availability is

an essential factor shaping marine benthic communities across the water depth gradient (Odum,

1971). Several studies have shown a decrease in the density of benthic invertebrates with depth on

ARs (Lewbel et al., 1986.; Moura et al., 2007; Shinn and Wicklund, 1989; van der Stap et al., 2016)

explained by the decrease in light intensity (Relini et al., 1994). The structural complexity has also

been put forward as important characteristics linked to ARs efficiency in ecological restoration (Strain

et al., 2018). Structural complexity increases available surface for colonization and niches diversity with

various shelter and light exposure conditions, the latter being related to different benthic assemblage

compositions (Glasby, 2000; Glasby, 1999) and higher productivity (Vivier et al., 2021). The recent 3D

printing techniques using concrete, allow the design of ARs mimicking natural habitats (Ly et al., 2021).

However, those studies concerned short-term colonization (<3.5 years) (Wendt et al., 1989) hence

based on pioneer species with high dispersal capacities which colonization is likely mainly regulated

by post-settlement processes such as competition, predation and physical disturbance (Todd, 1998).

In contrast to mobile species, sessile benthic invertebrates can only colonize reefs after larval dispersal

which is limited by reproduction frequency (Thorson, 1950). Colonization implies thus an effective

dispersal between natural areas and ARs, which depends on source population spatial distribution,

species fecundity, dispersive larval traits and ocean circulation. Nevertheless, until now dispersal

drivers have been disregarded while colonization disparities among ARs may result from differences in

both larval connectivity (which in turn depends on fecundity, dispersal capacities and adult distribution

in the natural habitat) and post-recruitment processes.

The objective of the present study was to test the hypothesis that the geographical location of ARs

deployment with respect to the natural habitat can condition ARs colonization after more than 10

years. To this aim we investigated the effects of local (shape, depth and immersion duration) and

regional (geographic area) factors on the presence of five species of sessile invertebrates with different

life history traits, endemic to the Gulf of Lion (GoL) (Northwestern Mediterranean Sea) and frequently

found on natural hard substrates.

MATERIALS AND METHODS

<u>**Study area and spatially stratified sampling design**</u>

The study area extended along 160 km of the GoL coastline (Figure 1). The GoL is a wide micro-tidal

continental shelf dominated by soft-bottom habitat with few small rocky habitat patches of less than

20 km^2. The GoL is a homogeneous and isolated hydrodynamic unit (Rossi et al., 2014), delimited by

the northern current (Millot, 1990).

Between 1985 and 2009, 763 ARs with different shapes or material and a total volume of 37 575 m^3

(Tessier et al., 2015) have been deployed along the GoL coastline over 66 km² of state concessions

(Cepralmar, 2015) between 10 and 35 m depth (Figure 1, Blouet et al., 2021). The ARs deployed in GoL

represent 40% of the total AR volume in France (Tessier et al., 2015).

In the present study we examined the oldest (in the 80s) and the youngest (in the 2000s) ARs with the

most common shapes (namely pipe, steel cage, Bonna, Comin, pyramid, heap of telegraph poles, and

concrete box, Figure 1).   For the analysis of data, we followed a stratified sampling design. To this aim,

the GoL coastline was regularly divided into 6 geographical sectors separated by a distance ranging

from 12 to 117 km, with a median value of 49 km (AGM referring to Aigues-Mortes, AGD to Agde, VLR

to Valras, GRU to Gruissan, LEU to Leucate, CST to Canet/Saint-Cyprien sectors, Figure 1). Each sector

included ARs deployed during either the first (1985, CST geographical sector), the last (2002-2009,

AGM, VLR, GRU, LEU geographical sectors) or both periods of immersion (AGD), and of different

shapes. In each geographical sector, two sites were defined except in AGD (5 sites, AGD1, AGD2, AGD3,

AGD4, AGD5) and GRU (3 sites, GRU1, GRU2, GRU3). To test for location effect, sites were separated

by a distance ranging from 2.1 to 11.7 km (median value of 7.5 km) except in GRU, where two of the

three sites (GRU1 and GRU2) were defined in the same location to account for different shapes. Hence,

depending on the geographical sector, sites may differ either by AR shape, immersion depth or timing

of deployment (Figure 1). Due to this set up, the effect of these three factors was only tested locally

(see Statistical analysis). In each site (except in the geographical sector GRU), three sampling units

separated by a distance between 4m and 3.6 Km (median distance of 251 m) were set out by pooling

neighboring ARs reaching a minimum surface of 89 m^2 per sampling unit and totalling a minimum

developed surface of 306 m^2 per site (Table 1: Supplementary Material). In the geographical sector

GRU, GRU2 and GRU3 included only one sampling unit because the surface of a single AR in these sites

already yielded 459 m2 (heap of telegraph poles; Table 1: Supplementary Material). Such large

continuous sampling units in each site aimed at limiting the effect of recruitment spatial variability

over distances from 100s meters to kilometers arising from the non-uniformity of the flow of larvae

(Daigle et al., 2014; Glasby, 2000; Simpson et al., 2017; Smale, 2012). Such a spatial scale is consistent

with the spatial scale of flow homogeneity obtained in simulations over GoL soft-bottom habitat

(Briton et al., 2018). This inventory methodology enabled us to test for the existence of structuring

factors at the local and regional scale. Developed reef surface was calculated on the basis of technical

specifications data present in the state concession documents taking into account only the colonizable

surface (surfaces in contact with the sediment were excluded).

Colonization assessment and species selection

Assessment of ARs colonization was carried out by autonomous scuba-diving in 2020 by direct visual

census counting the number of individuals of the five target species on the entire surface of ARs (7295

m²; Table1: Supplementary Material). On the ARs, all individuals older than one year (size > 2 cm) have

been recorded. Among the species listed in previous ARs inventories in the GoL, we selected five

species that were present in most inventories, easy to identify by direct visual census and spanning

different phyla with contrasting life-history traits (Créocean, 2003 & 2004; Table 1). We selected two

gorgonians *Eunicella singularis* (Esper, 1791) and *Leptogorgia sarmentosa* (Esper, 1789), one bryozoan

Pentapora fascialis (Pallas, 1766), one annelida *Sabella spallanzanii* (Gmelin, 1791) and one ascidian

Halocynthia papillosa (Linnaeus, 1797) (Figure 2). The five species have a similar wide natural

repartition area along European coasts ranging from 1m to 250m depth (Giangrande et al., 2005; Gori

et al., 2011; Ponti et al., 2019; Turon, 1990; Weinberg and Weinberg, 1979). In addition, *S. spallanzanii*

has been recorded along the coasts of Brazil, Australia and New Zealand where it is classified as an

invasive non-indigenous species (Currie et al., 2000).

The five species are present in the rocky habitat of the NW Mediterranean Sea (Laubier, 1966; True,

1970; Hong, 1980). However, in the GoL, where natural rocky habitat covers uneven surfaces within

the 6 geographical sectors (from 3,123 10^7 m^2 for the AGM sector to 5 10^5 m^2 for the LEU sector), the

five species display different spatial distributions (Dutrieux et al., 2005; Dalias et al., 2011; Guizien et

al., 2022; S. Blouet personal observation). *E. singularis* is frequently observed and abundant

throughout the GoL (from the AGM sector to the CST sector), while *L. sarmentosa,* less abundant, is

present mainly in the center of the GoL (AGD, VLR, LEU, CST). *P. fascialis* is abundant in the west of the

GoL (AGD, LEU, CST, and south of CST). The distribution of *H. papillosa* is not well known, however the

species has been observed in all the rocky areas of the GoL.

S. spallanzanii is present but rare in natural rocky habitat. Nevertheless, *S. spallanzanii* is very abundant

in lagoons, ports and marinas of the GoL (S. Blouet personal observation) which have been indicated

as preferred habitat of the species (Currie et al., 2000).

The five species display different life-history traits. All five species reproduce once a year in different

seasons and with different strategies (Table 1).

S. spallanzani reproduces in January-February, when water temperature is the coldest. The species

displays multiple reproductive strategies: internal fertilization, with larvae brooded either inside or

outside the mineral tube secreted around the body, and external fertilization broadcast spawning

(Giangrande et al., 2000). In addition, asexual reproduction by fission has been reported (Read et al.,

2011). *S. spallanzani* releases lecithotrophic larvae with a planktonic larval duration (PLD) of about 4

weeks (Giangrande et al., 2000). It's life span can exceed 5 years, with sexual maturity after one year

(Giangrande and Petraroli 1994; Giangrande et al., 2000). Like most gorgonians, *E. singularis* releases

lecithotrophic larvae in early summer (June to August). Even though larval competency period can

reach up to 2 months (Guizien et al., 2020; Zelli et al., 2020), PLDs ranging from 7 to 14 days best

explain gene flow among *E. singularis* natural populations dwelling in the fragmented rocky habitat of

the GoL (Padron et al. 2018). *E. singularis* life span can reach 25-30 years with sexual maturity before

6 years (Gori et al., 2007; Weinberg and Weinberg, 1979) (Weinberg and Weinberg 1979, Gori et al.

2007).

The other gorgonian, *L. sarmentosa* also releases lecithotrophic larvae but in the late summer

(September to October) and the PLD is unknown (Rossi and Gili 2009). *L. sarmentosa* life span can

reach 20 years with female sexual maturity within 2-3 years after settlement (Rossi and Gili 2009).

H. papillosa is a simultaneous hermaphrodite which releases larvae in late summer (September-

October; Becerro and Turon, 1992), presumably lecithotrophic. The PLD of *H. papillosa* larvae is

unknown but PLD shorter than 12 hours has been consistently reported for other solitary ascidian

species (Ayre et al., 1997). We did not find any data about the age at sexual maturity and the life span

of *H. papillosa*. However, the ascidians are considered as highly invasive, particularly because of their

rapid growth and early sexual maturity (Zhan et al., 2015), with some species such as C*iona*

intestionalis complex and *Ciona savigniy,* reaching sexual maturity at the age of 2 months (Zhan et al.,

2015) and continuous spawning (Carver et al. 2003).

P. fascialis displays both sexual and asexual reproduction. During sexual reproduction, most bryozoans

release lecithotrophic larvae which settle after a few minutes or a few hours, rarely beyond several

days (Keough, 1983). *P. fascialis* larval release has been inferred to happen in June based on

recruitment observations (Cocito et al., 1998a). Asexual reproduction happens by colony

fragmentation or budding extension (Cocito et al., 1998b). Individual life span is estimated to be about

10 years with early sexual maturity after 2 years (Cocito et al., 1998b).

Statistical   analysis

We examined to which extent ARs colonization measured by the presence/absence of the five species

and on the dissimilarity between their co-occurrence assemblages is affected by the AR shape, location

and timing of deployment (age). Due to the GoL AR deployment set up, the effect of some factors was

tested only locally. We first verified that shape was not affecting colonization, by testing the effect of

shape in one geographical sector where different shapes were deployed at same depth and time

(VLR1/VLR2, Table 1: Supplementary Material). Second, the effect of factors age (2 levels: 1985 and

the 2000s) and depth (2 levels, >20 m, <20 m depth) was tested within the AGD geographical sector

only. Factor age was tested in 3 sites at <20m (AGD1/ AGD2 & AGD3) and factor depth was tested in 4

sites deployed in 2009 (AGD2&AGD3 / AGD4&AGD5).   Third, factor location was tested both at

regional and local level (geographical sector being the regional factor, site being the local factor). To

avoid any confounding effect due to age or depth, the regional factor vs local factor were tested on

the 5 geographical sectors where AR were deployed during the 2000s immersion phase and at <20 m

depth only (AGM, AGD, VLR, GRU, LEU: 5x2 levels).   A Jaccard similarity matrix was built on

presence/absence data across all pairwise sampling units used in each test (Table 2: supplementary

material). Two multivariate analyses were performed. Non-parametric multivariate analysis of

variance with permutation was applied to test for the effects of shape, age, depth and geographical

sector on species assemblages (NP-manova: Anderson, 2001; Zar, 1999). Site was considered as a

random factor, nested either in depth, age or the geographical sector. Another multivariate analysis

was performed to cluster most similar species assemblages in the sector of AGD (SIMPROF : Clarke et

al., 2008) (Table 2 : supplementary material).

When significant differences between the five species co-occurrence assemblages were detected for

a factor, a non-parametric univariate analysis (Kruskall-Wallis) was performed for each species

independently to detect the species driving the difference (Table 2 : supplementary material). Fisher

post-hoc test was used to identify the site where the difference arose. A same p-value of 0.05 was

taken for detecting significant differences. Analyses were performed with Matlab software using the

Fathom package for multivariate analyses (Jones, 2014) and the Matlab statistics toolbox for univariate

analyses.

RESULTS

Artificial reefs colonization by the five target species at **regional scale**

Among the five target species, *S. spallanzanii* was the only one whose presence was recorded in all the

sampling units and geographical sectors (Figure 3). In only one out of 16 sites, it was the only species

detected. *H. papillosa* and *L. sarmentosa* were detected in five of the six geographical sectors (not

present in GRU and AGM, respectively). *P. fascialis* and *E. singularis* were detected in 3 of the 6

geographical sectors (CST, LEU, and AGD for *P. fascialis* and AGM, AGD and LEU, for *E. singularis*).

Finally, *E. singularis* was the least frequently observed species, being detected in only 5 out of the 16

sites. In all geographical sectors, at least three of the five target species were detected, except GRU

where only two of the five species were detected. Assemblages of two species were found in two sites

out of 16 (sector AGD and GRU), assemblages of three species were found in 7 sites, assemblages of

four species were found in 2 sites and assemblages of five species were found in 3 sites.

Shape effect on five target species at local scale

No differences in species assemblages were found in VLR between pipe and steel cage (NP-MANOVA,

F=9.08 e-19, P>0.05; Table 2).

Similarly, in the geographical sector GRU, the same assemblages were detected on sites differing by

ARs shape (pipes and a heap of telegraphical poles) in the same location (GRU1 and GRU2). Conversely,

different assemblages were detected between two sites (GRU2 and GRU3) separated by 9 km although

ARs shape was the same (heaps of telegraphical poles). Due to the absence of replication in unit

samplings, it was not possible to perform a statistical test on the effect of shape in this geographical

sector.

Age and depth effects on five target species at local scale

Despite all five target species being detected on ARs in the AGD sector, assemblage composition among sites differed (Figure 3). Clustering of sampling units within the 5 sites (AGD1, AGD2, AGD3, AGD4, AGD5) in AGD identified 2 clusters (SIMPROF: P <0.05; Figure 4). The two sites (AGD2 and AGD3) with same age (2009), depth range (less than 20 m) and reef shape (pipe) were attributed to different clusters.

In fact, one cluster grouped ARs of different age at a same depth (1985 in site AGD1 and 2009 in site AGD2) while the other cluster grouped ARs of the same age but at different depths (less than 20 m in site AGD3 and more than 20 m in sites AGD4 and AGD5). In both clusters, different AR shapes were found (steel cage and pipes in one cluster, pipes, Comin and Bonna in the other cluster) (Figure 4). The geographic distance between the two clusters (AGD4-AGD5) and (AGD3-AGD4-AGD5) was 7.5 km and the median value of the intra-cluster geographic distance was 3 km.

Multivariate analysis of variance confirmed that neither age (NP-MANOVA, F=1.43 P<0.05; Table 3) nor depth (NP-MANOVA, F=1.37 P<0.05; Table 4) explained site differences in the five species assemblages found on ARs in AGD (P<0.05; Tables 3 and 4). Differences among the five sites in AGD were due to different ARs colonization by three species (Kruskall-wallis: *E. singularis*, *P. fascialis*, *H. papillosa*: all P<0.05; Table 5). Site AGD3 differed from sites AGD5, AGD1 and AGD2 by the presence of E. singularis, the sites AGD4 and AGD5 differed from sites AGD1 and AGD2 due to the presence of P. fascialis and the site AGD1 differed from sites AGD5, AGD3 and AGD4 to the presence of H. papillosa (Post-hoc tests, Table 3: supplementary material).

Geographical effect on five target species at local and regional scales

The five species co-occurrence assemblages on ARs deployed in the same period and at same depth were significantly different at both regional and local scales (NP-MANOVA : geographical sector F=5.09 P<0.05; site (geographical sector) F=2.78 P<0.05; Table 6). These differences were due to different colonization of ARs by three of the five species, *E. singularis*, *L. sarmentosa* and *P. fascialis*. For the

latter two species, regional differences (Kruskall-wallis: geographical sector P=0.0002) were more

significant than local differences (Kruskall-wallis: site P=0.001 for *L. sarmentosa* and P=0.005 for *P.*

fascialis; Table 7). Both species were not detected in the north of the GoL (AGM). In contrast, for *E.*

singularis, local differences (Kruskall-wallis: site P=0.017) were more significant than regional ones

(geographical sector P=0.036; Table 7), the species being detected in geographical sectors in the north,

center and south of GoL.

DISCUSSION

The resilience of marine natural biodiversity, including benthic invertebrates, is essential to ensure the

sustainability of ecosystem functions. Increasing habitat is key to support the natural biodiversity

resilience, which in the case of rocky habitats can be achieved by effectively integrated ARs. The

effective integration of ARs into the rocky habitat network starts with their colonization by species

from the natural habitat. In the present study the five species selected from the natural rocky habitat

of the GoL colonized differently the ARs spread along the GoL coastline ten years after their

deployment. Locally, neither age, immersion depth or shape of the ARs significantly affected

colonization patterns.

Colonization of ARs are expected to evolve toward a stable state comparable to that of the natural

environment, through the succession of opportunistic species (wide dispersal, high fertility, low

tolerance of reduced resource levels, short life-spans, minimal dietary specialisation) followed by

specialized species (limited dispersal, slow growth to a large size at maturity, delayed and limited

reproduction, optimization to reduced resources and long life-spans; (Platt and Connell, 2003; Faurie

et al., 2003). Monitoring of ARs short-term colonization (<3 years) have indeed shown a dominance of

pioneer species (hydroids, serpulids, barnacles and bivalves), most of them having life history traits

typical of opportunistic species (Fariñas-Franco et al., 2013; T. Glasby, 1999; Moura et al., 2007;

Pamintuan and Ali, 1994; Ponti, 2015; Relini et al., 1994; Spagnolo et al., 2014; Toledo et al., 2020).

Long-term studies confirmed successions in ARs colonization (Burt et al., 2011; Nicoletti et al., 2007;

Perkol-Finkel and Benayahu, 2005; Whomersley and Picken, 2003), but none have described saturation

(Svane and Petersen, 2001). In the Tyrrhenian Sea, Nicolleti et al., (2007) described colonization in 5

distinct phases: (i) A first recruitment by pioneer species (hydroids, serpulids, barnacles and bivalves)

during the first months after immersion, followed by phases of (ii) cover dominance, (iii) regression

and (iv) absence of *Mytilus galloprovincialis* for more than 10 years. The installation of diverse bio-

builders bryozoans characteristic of the natural environment was recorded after 20 years only (v). Our

study shows that bio-builder engineering species such as bryozoans (*P. fascialis*) and gorgonians (*E.*

singularis, L. sarmentosa) colonized ARs as early as 10 years after their deployment, without significant

difference between 10 years and 35 years old ARs. However, the presence of *S. spallanzanii* described

as an opportunistic species (sexual precocity, various reproduction modes, rapid growth, short lived;

Giangrande et al., 2000) on all ARs independently of their age of deployment suggests that ARs did not

yet reach a stable state comparable to the natural environment. Thus, the presence of bio-builders is

not a sufficient indicator of the ARs naturalization to the local biodiversity.

The GoL's ARs being located in the sandy coastal zone are likely regularly disturbed by sediment

deposits due to river delivery or/and their resuspension by either trawling activities or the mechanical

action of the swell (Dufois et al., 2014; Durrieu de Madron et al., 2005; Ulses et al., 2008). Testing the

impact of swell and sediment deposit on ARs requires exploring the colonization of ARs along a

gradient of depth and distance from the coast (van der Stap et al., 2016). However, current ARs

deployment in the GoL ranged from 15 to 30 m depth and within 3 miles from the coast and did not

allow testing for differential effect of sediment disturbances as swell impact occurs every year in this

area (Guizien, 2009). Testing the impact of sediment disturbance on ARs colonization would require

exploring reefs deployed deeper than 50 m, such as the anchorages of the floating wind farm that will

be placed in the GoL in the next future (https://info-efgl.fr/le-projet/le-parc/#). Light is also expected

to be an important factor structuring benthic assemblages, along a depth gradient in natural and

artificial environments (T. Glasby, 1999a; T. M. Glasby, 1999b; Svane and Petersen, 2001) . Absence

of depth effect in the present study, although in the GoL light intensity strongly attenuates within the

upper 30 m of the water column (Durrieu de Madron et al., 2011) was potentially a bias due to the five

species selected in the present study whose distributions are not strongly structured by light intensity.

Another factor which has been shown to drive the intensity of ARs colonization is structural complexity

(see Bohnsack and Sutherland, 1985 for a review). Nevertheless, there is no consensus about the

relationship between complexity and subtidal benthic invertebrates abundance, due to potential bias

in controlling the surface and scale in ARs of different complexity (Rouse et al., 2019; Strain et al.,

2018). The similarity in the 5 species co-occurrence between different reef shapes at the same depth

and of the same age found in the present study suggests that shape is less important than the

geographical location in AR colonization by benthic invertebrates when controlling the colonized

surface. However, shape is an imprecise measure of structural complexity. The latter is rarely assessed

quantitatively and can be described by different metrics which may be similar for apparently different

shapes (such as steel cages and pipes, (Riera, 2020).

Benthic invertebrate assemblages result from complex processes that operate at multiple spatial and

temporal scales (Smale, 2012). At the regional scale, larval availability can become a major factor

explaining colonization success (Padron et al., 2018). Change in the composition of assemblages during

the early colonization of artificial substrates by benthic invertebrates has been attributed to the

availability and abundance of larvae during the seasons rather than a sequence of distinct succession

(Bramanti et al., 2003; Turner and Todd, 1993). The larval behaviour (buoyancy and motility) and the

characteristics of the biological cycle of the species (spawning timing and PLD) can play a key role in

determining the dispersal distance (Todd, 1998), and consequently the possibility to reach habitat

suitable for settlement. Dispersion distance is generally correlated with PLD, thus, a species with a long

PLD is supposed to colonize habitats further away than species with a shorter PLD (Shanks, 2009). In

this study, the five species were chosen among different phyla known for their contrasting planktonic

durations, swimming abilities and larval dispersal periods, although these larval traits are only known

accurately for *E. singularis* (Guizien et al., 2020; Zelli et al., 2020). *P. fascialis* and *H. papillosa*, the two

species with short PLD (<24h and <12h, respectively) colonized ARs located close to the natural habitat

where they are present (< 4.8 km and <10 km, respectively). The coastal circulation of the GoL allows

such dispersal distance over periods of a few days (Guizien et al., 2012). *S. spallanzanii*, which has a

PLD of 3 weeks, colonized all the inventoried ARs, in line with a dispersal distance of 40 km after 3

weeks (Guizien et al., 2012). In contrast, *E. singularis* did not colonize all ARs within geographical

sectors of 30 km width where the species is present in the natural habitat, although a 2-week PLD was

expected to enable such dispersal (Padron et al., 2018). The other gorgonian species, *L. sarmentosa*

colonized more ARs located within distances of less than 30 km from its natural habitat than *E.*

singularis while the PLD of the two species are presumably the same. This suggests that other factors

influence the colonization failure of *E. singularis*.

Reproductive traits are another key to the success in colonising new settings (Stearns, 2000). In this

regard, *E. singularis* colonization potential could be limited by its low fecundity (~25-40 larvae.cm^{-1} of

colony branch, Ribes et al., 2007; Theodor, 1967) compared to that of *L. sarmentosa* (~75 larvae.cm^{-1}

of colony branch, (Rossi et al., 2011; Rossi and Gili, 2009). The wide colonization of ARs by *S.*

spallanzanii is in line with its reproductive traits typical of opportunistic species (early sexual maturity,

high fecundity with more than 50 000 eggs per female, Currie et al., 2000, a fertilization close to 100%,

Giangrande et al., 2000). Since arriving in the Pacific Ocean, *S. spallanzanii* has been declared one of

the ten priority pest species in the marine environment by the Australian authorities and classified as

an invasive species (Hayes et al., 2005). Similarly to *S. spallanzanii*, *H. papillosa* colonized nearly all ARs

located within its 10-km dispersal distance from the natural habitat. Within the ascidian class, a wide

disparity in species fecundity has been reported (Pandian, 2018). This suggests *H. papillosa*

reproductive traits would be close to those of invasive ascidians (Zhan et al., 2015).

Ultimately, understanding ARs colonization requires a precise mapping of source populations in the

natural environment. To this respect, the abundance of *S. spallanzani*i on ARs is surprising, as the

species is not abundant in the natural rocky habitat of the GoL. For this species, other sources of larval

supply than natural settings should be considered, such as the numerous ports and marinas along the

coast of the GoL, as *S. spallanzanii* is very tolerant to environmental conditions (Currie et al., 2000). In

this case of intense colonization by an endemic benthic invertebrate species, ARs apparently extended

its metapopulation, acting as stepping stones for further larval dispersal beyond its natural current

colonization limits (Bishop et al., 2017; Wang et al., 2020). In the GoL, the 14 500 m^3 of ARs deployed

30 years ago are now decommissioned and the relevance of their removal is currently debated.

Connectivity between natural populations has been shown to support species resilience after

disturbances in fragmented habitat, and could be extended to ARs (Fahrig, 2003). However, ARs may

also facilitate the spread of non-indigenous species introduced with maritime traffic in ports (Glasby

et al., 2007).

The present study advocates accounting for the geographical arrangement in planning ARs deployment

to extend the habitat of hard bottom benthic invertebrate natural populations.

*Acknowledgement*s. This work was funded by the Agence de l'Eau Rhône-Méditerranée-Corse under

project ICONE - Impacts actuels et potentiels de la CONnectivité Ecologique ajoutée par les récifs

artificiels sur la biodiversité fixée des substrats durs du Golfe du Lion (PI, K. Guizien, AAP 2016). The

authors gratefully acknowledge the helpful assistance during sampling of the staff of the Aire Marine

Protégée Agathoise.

Authors contributions

383 SB and KG conceived the study, SB carried out sampling and statistical analysis. All contributed to

384 manuscript writing.

Conflict of interest disclosure

The authors of this manuscript declare that they have no financial conflict of interest with the content

of this article.

LITERATURE CITED

Anderson, M., 2001. A new method for non-parametric multivariate analysis of variance. Austral
Ecology 26, 32–46. https://doi.org/10.1111/j.1442-9993.2001.01070.pp.x
Ardizzone, G.D., Belluscio, A., Gravina, M.F., Somaschini, A., 1996. Colonization and Disappearance of
Mytilus galloprovincialis Lam. on an Artificial Habitat in the Mediterranean Sea. Estuarine,
Coastal and Shelf Science 43, 665–676. https://doi.org/10.1006/ecss.1996.0095
Ayre, D.J., Davis, A.R., Billingham, M., Llorens, T., Styan, C., 1997. Genetic evidence for contrasting
patterns of dispersal in solitary and colonial ascidians. Marine Biology 130, 51–61.
https://doi.org/10.1007/s002270050224
Becerro, M.A., Turon, X., 1992. Reproductive Cycles of the Ascidians *Microcosmus sabatieri* and
Halocynthia papillosa in the Northwestern Mediterranean. Marine Ecology 13, 363–373.
https://doi.org/10.1111/j.1439-0485.1992.tb00360.x
Bishop, M.J., Mayer-Pinto, M., Airoldi, L., Firth, L.B., Morris, R.L., Loke, L.H.L., Hawkins, S.J., Naylor,
L.A., Coleman, R.A., Chee, S.Y., Dafforn, K.A., 2017. Effects of ocean sprawl on ecological
connectivity: impacts and solutions. Journal of Experimental Marine Biology and Ecology 492,
7–30. https://doi.org/10.1016/j.jembe.2017.01.021
Blouet, S., Quittet, L., Agin, G., Thorin, S., Dalias, N., Marobin, D., Lenfant, P., Guizien, K., 2021.
Database of the location and typology of artificial reefs in the Gulf of Lion (NW
Mediterranean Sea). https://doi.org/10.17882/81407
Bohnsack, J.A., 1989. Are High Densities of Fishes at Artificial Reefs the Result of Habitat Limitation or
Behavioral Preference? BULLETIN OF MARINE SCIENCE 44, 15.
Bohnsack, J.A., Sutherland, D.L., 1985. ARTIFICIAL REEF RESEARCH: A REVIEW WITH
RECOMMENDATIONS FOR FUTURE PRIORITIES. BULLETIN OF MARINE SCIENCE 37, 30.
Bramanti, L., Magagnini, G., Santangelo, G., 2003. Settlement and recruitment: The first stages in the
life cycle of two epibenthic suspension feeders (Corallium *rubrum* and *Anomia ephippium)*.
Italian Journal of Zoology 70, 175–178. https://doi.org/10.1080/11250000309356512
Briton, F., Cortese, D., Duhaut, T., Guizien, K., 2018. High-resolution modelling of ocean circulation
can reveal retention spots important for biodiversity conservation. Aquatic Conserv: Mar
Freshw Ecosyst 28, 882–893. https://doi.org/10.1002/aqc.2901
Burt, J., Bartholomew, A., Sale, P.F., 2011. Benthic development on large-scale engineered reefs: A
comparison of communities among breakwaters of different age and natural reefs. Ecological
Engineering 37, 191–198. https://doi.org/10.1016/j.ecoleng.2010.09.004
Chen, J.-L., Chuang, C.-T., Jan, R.-Q., Liu, L.-C., Jan, M.-S., 2013. Recreational Benefits of Ecosystem
Services on and around Artificial Reefs: A Case Study in Penghu, Taiwan. Ocean & Coastal
Management 85, 58–64. https://doi.org/10.1016/j.ocecoaman.2013.09.005
Clarke, K.R., Somerfield, P.J., Gorley, R.N., 2008. Testing of null hypotheses in exploratory community
analyses: similarity profiles and biota-environment linkage. Journal of Experimental Marine
Biology and Ecology 366, 56–69. https://doi.org/10.1016/j.jembe.2008.07.009
Claudet, J., Fraschetti, S., 2010. Human-driven impacts on marine habitats: A regional meta-analysis
in the Mediterranean Sea. Biological Conservation 143, 2195–2206.
https://doi.org/10.1016/j.biocon.2010.06.004
Claudet, J., Pelletier, D., 2004. Marine protected areas and artificial reefs: A review of the
interactions between management and scientific studies. Aquat. Living Resour. 17, 129–138.
https://doi.org/10.1051/alr:2004017
Cocito, Silvia, Ferdeghini, F., Sgorbini, S., 1998. *Pentapora fascialis* (Pallas) [Cheilostomata:
Ascophora] colonization of one sublittoral rocky site after sea-storm in the northwestern
Mediterranean, in: Baden, S., Phil, L., Rosenberg, R., Strömberg, J.-O., Svane, I., Tiselius, P.
(Eds.), Recruitment, Colonization and Physical-Chemical Forcing in Marine Biological Systems.
Springer Netherlands, Dordrecht, pp. 59–66. https://doi.org/10.1007/978-94-017-2864-5_5

Cocito, S., Sgorbini, S., Bianchi, C.N., 1998. Aspects of the biology of the bryozoan *Pentapora fascialis*
in the northwestern Mediterranean. Marine Biology 131, 73–82.
https://doi.org/10.1007/s002270050298
Connell, J.H., Slatyer, R.O., 1977. Mechanisms of Succession in Natural Communities and Their Role in
Community Stability and Organization. The American Naturalist 111, 1119–1144.
https://doi.org/10.1086/283241
Currie, D.R., McArthur, M.A., Cohen, B.F., 2000. Reproduction and distribution of the invasive
European fanworm *Sabella spallanzanii* (Polychaeta: Sabellidae) in Port Phillip Bay, Victoria,
Australia. Marine Biology 136, 645–656. https://doi.org/10.1007/s002270050724
Daigle, R.M., Metaxas, A., deYoung, B., 2014. Fine-Scale Distribution and Spatial Variability of Benthic
Invertebrate Larvae in an Open Coastal Embayment in Nova Scotia, Canada. PLOS ONE 9,
e106178. https://doi.org/10.1371/journal.pone.0106178
Dufois, F., Verney, R., Le Hir, P., Dumas, F., Charmasson, S., 2014. Impact of winter storms on
sediment erosion in the Rhone River prodelta and fate of sediment in the Gulf of Lions (North
Western Mediterranean Sea). Continental Shelf Research 72, 57–72.
https://doi.org/10.1016/j.csr.2013.11.004
Durrieu de Madron, X., Ferré, B., Le Corre, G., Grenz, C., Conan, P., Pujo-Pay, M., Buscail, R., Bodiot,
O., 2005. Trawling-induced resuspension and dispersal of muddy sediments and dissolved
elements in the Gulf of Lion (NW Mediterranean). Continental Shelf Research 25, 2387–2409.
https://doi.org/10.1016/j.csr.2005.08.002
Durrieu de Madron, X., Guieu, C., Sempéré, R., Conan, P., Cossa, D., D'Ortenzio, F., Estournel, C.,
Gazeau, F., Rabouille, C., Stemmann, L., Bonnet, S., Diaz, F., Koubbi, P., Radakovitch, O.,
Babin, M., Baklouti, M., Bancon-Montigny, C., Belviso, S., Bensoussan, N., Bonsang, B.,
Bouloubassi, I., Brunet, C., Cadiou, J.-F., Carlotti, F., Chami, M., Charmasson, S., Charrière, B.,
Dachs, J., Doxaran, D., Dutay, J.-C., Elbaz-Poulichet, F., Eléaume, M., Eyrolles, F., Fernandez,
C., Fowler, S., Francour, P., Gaertner, J.C., Galzin, R., Gasparini, S., Ghiglione, J.-F., Gonzalez,
J.-L., Goyet, C., Guidi, L., Guizien, K., Heimbürger, L.-E., Jacquet, S.H.M., Jeffrey, W.H., Joux, F.,
Le Hir, P., Leblanc, K., Lefèvre, D., Lejeusne, C., Lemé, R., Loÿe-Pilot, M.-D., Mallet, M.,
Méjanelle, L., Mélin, F., Mellon, C., Mérigot, B., Merle, P.-L., Migon, C., Miller, W.L., Mortier,
L., Mostajir, B., Mousseau, L., Moutin, T., Para, J., Pérez, T., Petrenko, A., Poggiale, J.-C.,
Prieur, L., Pujo-Pay, M., Pulido-Villena, Raimbault, P., Rees, A.P., Ridame, C., Rontani, J.-F.,
Ruiz Pino, D., Sicre, M.A., Taillandier, V., Tamburini, C., Tanaka, T., Taupier-Letage, I., Tedetti,
M., Testor, P., Thébault, H., Thouvenin, B., Touratier, F., Tronczynski, J., Ulses, C., Van
Wambeke, F., Vantrepotte, V., Vaz, S., Verney, R., 2011. Marine ecosystems' responses to
climatic and anthropogenic forcings in the Mediterranean. Progress in Oceanography 91, 97–
166. https://doi.org/10.1016/j.pocean.2011.02.003
Fahrig, L., 2003. Effects of Habitat Fragmentation on Biodiversity. Annu. Rev. Ecol. Evol. Syst. 34, 487–
515. https://doi.org/10.1146/annurev.ecolsys.34.011802.132419
Fariñas-Franco, J.M., Allcock, L., Smyth, D., Roberts, D., 2013. Community convergence and
recruitment of keystone species as performance indicators of artificial reefs. Journal of Sea
Research 78, 59–74. https://doi.org/10.1016/j.seares.2012.10.008
Fava, F., Ponti, M., Abbiati, M., 2016. Role of Recruitment Processes in Structuring Coralligenous
Benthic Assemblages in the Northern Adriatic Continental Shelf. PLOS ONE 11, e0163494.
https://doi.org/10.1371/journal.pone.0163494
Giangrande, A., Cavallo, A., Licciano, M., Mola, E., Pierri, C., Trianni, L., 2005. Utilization of the filter
feeder polychaete Sabella. Aquacult Int 13, 129–136. https://doi.org/10.1007/s10499-004-
9025-3
Giangrande, A., Licciano, M., Pagliara, P., Gambi, M.C., 2000. Gametogenesis and larval development
in *Sabella spallanzanii* (Polychaeta: Sabellidae) from the Mediterranean Sea. Marine Biology
136, 847–861. https://doi.org/10.1007/s002279900251

Glasby, T., 1999. Interactive effects of shading and proximity to the seafloor on the development of
subtidal epibiotic assemblages. Mar. Ecol. Prog. Ser. 190, 113–124.
https://doi.org/10.3354/meps190113
Glasby, T., Connell, S., Holloway, M., Hewitt, C., 2007. Nonindigenous biota on artificial structures:
Could habitat creation facilitate biological invasions? Marine Biology 151, 887–895.
https://doi.org/10.1007/s00227-006-0552-5
Glasby, T.M., 2000. Surface composition and orientation interact to affect subtidal epibiota. Journal
of Experimental Marine Biology and Ecology 248, 177–190. https://doi.org/10.1016/S0022-
0981(00)00169-6
Glasby, T.M., 1999. Effects of shading on subtidal epibiotic assemblages. Journal of Experimental
Marine Biology and Ecology 234, 275–290. https://doi.org/10.1016/S0022-0981(98)00156-7
Gori, A., Linares, C., Rossi, S., Coma, R., Gili, J.-M., 2007. Spatial variability in reproductive cycle of the
gorgonians *Paramuricea clavata* and *Eunicella singularis* (Anthozoa, Octocorallia) in the
Western Mediterranean Sea. Mar Biol 151, 1571–1584. https://doi.org/10.1007/s00227-006-
0595-7
Gori, A., Rossi, S., Berganzo, E., Pretus, J.L., Dale, M.R.T., Gili, J.-M., 2011. Spatial distribution patterns
of the gorgonians *Eunicella singularis*, *Paramuricea clavata*, and *Leptogorgia sarmentosa*
(Cape of Creus, Northwestern Mediterranean Sea). Mar Biol 158, 143–158.
https://doi.org/10.1007/s00227-010-1548-8
Grossman, G.D., Jones, G.P., Seaman, W.J., 1997. Do Artificial Reefs Increase Regional Fish
Production? A Review of Existing Data 22, 7.
Guizien, K., 2009. Spatial variability of wave conditions in the Gulf of Lions (NW Mediterranean Sea).
Vie Milieu 11.
Guizien, K., Belharet, M., Marsaleix, P., Guarini, J.M., 2012. Using larval dispersal simulations for
marine protected area design: Application to the Gulf of Lions (northwest Mediterranean).
Limnology and Oceanography 57, 1099–1112. https://doi.org/10.4319/lo.2012.57.4.1099
Guizien, K., Viladrich, N., Martínez-Quintana, Á., Bramanti, L., 2020. Survive or swim: different
relationships between migration potential and larval size in three sympatric Mediterranean
octocorals. Scientific Reports 10, 18096. https://doi.org/10.1038/s41598-020-75099-1
Guizien, K., Bramanti, L., Roca, J-C., Hesse, B., Blouet, S., Charbonnel, E., Payrot, J., Ferrari, B., Bricout,
R., Lescure, L., Chere, E., Hartmann, V., Rivoal, R., Planque, J-F., Cadene, F., Laffon, J-F.,
Monin, M., Bretton, O., Dumontier, M., Labrune, C., Fanfard, S., Cornette, F., 2022. Database
of sea fans spatial distribution in the northwestern Mediterranean sea.
https://doi.org/10.17882/86176
Hayes, K.R., Australia, Department of the Environment and Heritage, CSIRO (Australia), Division of
Marine Research, 2005. National priority pests. Part II, Part II,. Australian Government Dept.
of the Environment and Heritage, Parkes A.C.T.
Jackson, J.B.C., 2001. Historical Overfishing and the Recent Collapse of Coastal Ecosystems. Science
293, 629–637. https://doi.org/10.1126/science.1059199
Keough, M.J., 1983. Patterns of recruitment of sessile invertebrates in two subtidal habitats. Journal
of Experimental Marine Biology and Ecology 66, 213–245. https://doi.org/10.1016/0022-
0981(83)90162-4
Lewbel, G.S., Howard, R.L., Gallaway, B.J., n.d. Zonation of Dominant Fouling Organisms on Northern
Gulf of Mexico Petroleum Platforms 26.
Ly, O., Yoris-Nobile, A.I., Sebaibi, N., Blanco-Fernandez, E., Boutouil, M., Castro-Fresno, D., Hall, A.E.,
Herbert, R.J.H., Deboucha, W., Reis, B., Franco, J.N., Teresa Borges, M., Sousa-Pinto, I., van
der Linden, P., Stafford, R., 2021. Optimisation of 3D printed concrete for artificial reefs:
Biofouling and mechanical analysis. Construction and Building Materials 272, 121649.
https://doi.org/10.1016/j.conbuildmat.2020.121649
Martens, K., Queiroga, H., Cunha, M.R., Cunha, A., Moreira, M.H., Quintino, V., Rodrigues, A.M.,
Seroôdio, J., Warwick, R.M. (Eds.), 2006. Marine Biodiversity: Patterns and Processes,

Assessment, Threats, Management and Conservation. Springer Netherlands, Dordrecht.
https://doi.org/10.1007/1-4020-4697-9

Millot, C., 1990. The Gulf of Lions' hydrodynamics. Continental Shelf Research 10, 885–894.
https://doi.org/10.1016/0278-4343(90)90065-T

Moura, A., Boaventura, D., Cúrdia, J., Carvalho, S., da Fonseca, L.C., Leitão, F.M., Santos, M.N.,
Monteiro, C.C., 2007. Effect of depth and reef structure on early macrobenthic communities
of the Algarve artificial reefs (southern Portugal). Hydrobiologia 580, 173–180.
https://doi.org/10.1007/s10750-006-0455-y

Nicoletti, L., Marzialetti, S., Paganelli, D., Ardizzone, G.D., 2007. Long-term changes in a benthic
assemblage associated with artificial reefs, in: Relini, G., Ryland, J. (Eds.), Biodiversity in
Enclosed Seas and Artificial Marine Habitats, Developments in Hydrobiology. Springer
Netherlands, Dordrecht, pp. 233–240. https://doi.org/10.1007/978-1-4020-6156-1_20

Odum, E.P., n.d. An understanding of ecological succession provides a basis for resolving man's
conflict with nature. 164, 9.

Padron, M., Costantini, F., Baksay, S., Bramanti, L., Guizien, K., 2018. Passive larval transport explains
recent gene flow in a Mediterranean gorgonian. Coral Reefs 37.
https://doi.org/10.1007/s00338-018-1674-1

Pamintuan, I.S., Ali, P.M., n.d. Early Successional Patterns of Invertebrates in Artificial Reefs
Established at Clear and Silty Areas in Bolinao, Pangasinan, Northern Philippines 11.

Pandian, T.J., 2018. Reproduction and Development in Echinodermata and Prochordata. CRC Press,
Boca Raton. https://doi.org/10.1201/9780815364733

Pauly, D., Christensen, V., Guénette, S., Pitcher, T.J., Sumaila, U.R., Walters, C.J., Watson, R., Zeller,
D., 2002. Towards sustainability in world fisheries. Nature 418, 689–695.
https://doi.org/10.1038/nature01017

Perkol-Finkel, S., Benayahu, Y., 2005. Recruitment of benthic organisms onto a planned artificial reef:
shifts in community structure one decade post-deployment. Marine Environmental Research
59, 79–99. https://doi.org/10.1016/j.marenvres.2004.03.122

Platt, W.J., Connell, J.H., 2003. Natural Disturbances and Directional Replacement of Species.
Ecological Monographs 73, 507–522. https://doi.org/10.1890/01-0552

Ponti, M., 2015. Benthic assemblages on artificial reefs in the northwestern Adriatic Sea: Does
structure type and age matter? Marine Environmental Research 10.

Ponti, M., Turicchia, E., Costantini, F., Gori, A., Bramanti, L., Di Camillo, C., Linares, C., Rossi, S.,
Abbiati, M., Garrabou, J., Cerrano, C., 2019. Mediterranean gorgonian forests: Distribution
patterns and ecological roles.

Powers, S., Grabowski, J., Peterson, C., Lindberg, W., 2003. Estimating enhancement of fish
production by offshore artificial reefs: uncertainty exhibited by divergent scenarios. Mar.
Ecol. Prog. Ser. 264, 265–277. https://doi.org/10.3354/meps264265

Read, G., Inglis, G., Stratford, P., Ahyong, S., 2011. Arrival of the alien fanworm Sabella spallanzanii
(Gmelin, 1791) (Polychaeta: Sabellidae) in two New Zealand harbours. AI 6, 273–279.
https://doi.org/10.3391/ai.2011.6.3.04

Relini, G., 2002. Trophic relationships between fishes and an artificial reef. ICES Journal of Marine
Science 59, S36–S42. https://doi.org/10.1006/jmsc.2002.1212

Relini, G., Zamboni, N., Tixi, F., Torchia, G., n.d. Patterns of Sessile Macrobenthos Community
Development on an Artificial Reef in the Gulf of Genoa (Northwestern Mediterranean) 27.

Ribes, M., Coma, R., Rossi, S., Micheli, M., 2007. Cycle of gonadal development in *Eunicella singularis*
(Cnidaria: Octocorallia): trends in sexual reproduction in gorgonians: Gametogenic cycle in a
temperate gorgonian. Invertebrate Biology 126, 307–317. https://doi.org/10.1111/j.1744-
7410.2007.00101.x

Riera, E., 2020. Vers une construction raisonnée d'une nouvelle génération de récifs artificiels :
Analyses comparatives des facteurs intrinsèques favorisant leur colonisation de la micro à la
macro-échelle 192. Thése.

Rossi, S., Gili, J.-M., 2009. Reproductive features and gonad development cycle of the soft bottom-
gravel gorgonian *Leptogorgia sarmentosa* (Esper, 1791) in the NW Mediterranean Sea.
Invertebrate Reproduction & Development 53, 175–190.
https://doi.org/10.1080/07924259.2009.9652304
Rossi, S., Gili, J.-M., Garrofé, X., 2011. Net negative growth detected in a population of *Leptogorgia*
sarmentosa: quantifying the biomass loss in a benthic soft bottom-gravel gorgonian. Mar Biol
158, 1631–1643. https://doi.org/10.1007/s00227-011-1675-x
Rossi, V., Ser-Giacomi, E., López, C., Hernández-García, E., 2014. Hydrodynamic provinces and oceanic
connectivity from a transport network help designing marine reserves. arXiv:1407.6920
[physics]. https://doi.org/10.1002/2014GL059540
Rouse, S., Porter, J.S., Wilding, T.A., n.d. Artificial reef design affects benthic secondary productivity
and provision of functional habitat 9.
Santos, M.N., Monteiro, C.C., 2007. A fourteen-year overview of the fish assemblages and yield of the
two oldest Algarve artificial reefs (southern Portugal) 7.
Seaman, W., 2007. Artificial habitats and the restoration of degraded marine ecosystems and
fisheries, in: Relini, G., Ryland, J. (Eds.), Biodiversity in Enclosed Seas and Artificial Marine
Habitats, Developments in Hydrobiology. Springer Netherlands, Dordrecht, pp. 143–155.
https://doi.org/10.1007/978-1-4020-6156-1_13
Shanks, A., 2009. Barnacle settlement versus recruitment as indicators of larval delivery. II. Time-
series analysis and hypothesized delivery mechanisms. Mar. Ecol. Prog. Ser. 385, 217–226.
https://doi.org/10.3354/meps08002
Shinn, E.A., Wicklund, R.I., n.d. Artificial Reef Observations from a Manned Submersible off Southeast
Florida 10.
Simpson, T.J.S., Smale, D.A., McDonald, J.I., Wernberg, T., 2017. Large scale variability in the
structure of sessile invertebrate assemblages in artificial habitats reveals the importance of
local-scale processes. Journal of Experimental Marine Biology and Ecology 494, 10–19.
https://doi.org/10.1016/j.jembe.2017.05.003
Smale, D.A., 2012. Extreme spatial variability in sessile assemblage development in subtidal habitats
off southwest Australia (southeast Indian Ocean). Journal of Experimental Marine Biology
and Ecology 438, 76–83. https://doi.org/10.1016/j.jembe.2012.10.002
Spagnolo, A., Cuicchi, C., Punzo, E., Santelli, A., Scarcella, G., Fabi, G., 2014. Patterns of colonization
and succession of benthic assemblages in two artificial substrates. Journal of Sea Research
88, 78–86. https://doi.org/10.1016/j.seares.2014.01.007
Stearns, S.C., n.d. Life history evolution: successes, limitations, and prospects 11.
Steimle, F., 2002. Benthic macrofauna productivity enhancement by an artificial reef in Delaware
Bay, USA. ICES Journal of Marine Science 59, S100–S105.
https://doi.org/10.1006/jmsc.2002.1268
Strain, E.M.A., Olabarria, C., Mayer-Pinto, M., Cumbo, V., Morris, R.L., Bugnot, A.B., Dafforn, K.A.,
Heery, E., Firth, L.B., Brooks, P.R., Bishop, M.J., 2018. Eco-engineering urban infrastructure
for marine and coastal biodiversity: Which interventions have the greatest ecological
benefit? J Appl Ecol 55, 426–441. https://doi.org/10.1111/1365-2664.12961
Svane, I., Petersen, J.K., 2001. On the Problems of Epibioses, Fouling and Artificial Reefs, a Review.
Marine Ecology 22, 169–188. https://doi.org/10.1046/j.1439-0485.2001.01729.x
Tessier, A., Francour, P., Charbonnel, E., Dalias, N., Bodilis, P., Seaman, W., Lenfant, P., 2015.
Assessment of French artificial reefs: due to limitations of research, trends may be
misleading. Hydrobiologia 753, 1–29. https://doi.org/10.1007/s10750-015-2213-5
Tessier, A., Verdoit-Jarraya, M., Blouet, S., Dalias, N., Lenfant, P., 2014. A case study of artificial reefs
as a potential tool for maintaining artisanal fisheries in the French Mediterranean Sea. Aquat.
Biol. 20, 255–272. https://doi.org/10.3354/ab00563
Thorson, G., 1950. Reproductive and larval ecology of marine bottom invertebrates. Biological
Reviews 25, 1–45. https://doi.org/10.1111/j.1469-185X.1950.tb00585.x

642 Todd, C.D., 1998. Larval supply and recruitment of benthic invertebrates: do larvae always disperse
643   as much as we believe? Hydrobiologia 375, 1–21. https://doi.org/10.1023/A:1017007527490
644 Toledo, M.-I., Torres, P., Díaz, C., Zamora, V., López, J., Olivares, G., 2020. Ecological succession of
645   benthic organisms on niche-type artificial reefs. Ecol Process 9, 38.
646   https://doi.org/10.1186/s13717-020-00242-9
647 Turner, S.J., Todd, C.D., 1993. The early development of epifaunal assemblages on artificial substrata
648   at two intertidal sites on an exposed rocky shore in St. Andrews Bay, N.E. Scotland. Journal of
649   Experimental Marine Biology and Ecology 166, 251–272. https://doi.org/10.1016/0022-
650   0981(93)90223-B
651 Turon, X., 1990. Distribution and Abundance of Ascidians from a Locality on the Northeast Coast of
652   Spain. Marine Ecology 11, 291–308. https://doi.org/10.1111/j.1439-0485.1990.tb00385.x
653 Ulses, C., Estournel, C., Durrieu de Madron, X., Palanques, A., 2008. Suspended sediment transport in
654   the Gulf of Lions (NW Mediterranean): Impact of extreme storms and floods. Continental
655   Shelf Research 28, 2048–2070. https://doi.org/10.1016/j.csr.2008.01.015
656 van der Stap, T., Coolen, J.W.P., Lindeboom, H.J., 2016. Marine Fouling Assemblages on Offshore Gas
657   Platforms in the Southern North Sea: Effects of Depth and Distance from Shore on
658   Biodiversity. PLoS ONE 11, e0146324. https://doi.org/10.1371/journal.pone.0146324
659 Vivier, B., Dauvin, J.-C., Navon, M., Rusig, A.-M., Mussio, I., Orvain, F., Boutouil, M., Claquin, P., 2021.
660   Marine artificial reefs, a meta-analysis of their design, objectives and effectiveness. Global
661   Ecology and Conservation e01538. https://doi.org/10.1016/j.gecco.2021.e01538
662 Wang, W., Wang, J., Choi, F.M.P., Ding, P., Li, X., Han, G., Ding, M., Guo, M., Huang, X., Duan, W.,
663   Cheng, Z., Chen, Z., Hawkins, S.J., Jiang, Y., Helmuth, B., Dong, Y., 2020. Global warming and
664   artificial shorelines reshape seashore biogeography. Global Ecol Biogeogr 29, 220–231.
665   https://doi.org/10.1111/geb.13019
666 Weinberg, S., Weinberg, F., 1979. The Life Cycle of a Gorgonian: *Eunicella Singularis* (Esper, 1794).
667   Contributions to zoology Bijdragen tot de dierkunde 48, 127–140.
668   https://doi.org/10.1163/26660644-04802003
669 Wendt, P.H., Knott, D.M., Dolah, R.F.V., n.d. Community structure of the sessile biota on five artificial
670   reefs of different ages 18.
671 Whomersley, P., Picken, G.B., 2003. Long-term dynamics of fouling communities found on offshore
672   installations in the North Sea. J. Mar. Biol. Ass. 83, 897–901.
673   https://doi.org/10.1017/S0025315403008014h
674 Wilson, K., 2002. Restoration of Hong Kong fisheries through deployment of artificial reefs in marine
675   protected areas. ICES Journal of Marine Science 59, S157–S163.
676   https://doi.org/10.1006/jmsc.2002.1186
677 Zar, J., 1999. Biostatistical analysis.
678 Zelli, E., Quéré, G., Lago, N., Di Franco, G., Costantini, F., Rossi, S., Bramanti, L., 2020. Settlement
679   dynamics and recruitment responses of Mediterranean gorgonians larvae to different
680   crustose coralline algae species. Journal of Experimental Marine Biology and Ecology 530–
681   531, 151427. https://doi.org/10.1016/j.jembe.2020.151427
682 Zhan, A., Briski, E., Bock, D.G., Ghabooli, S., MacIsaac, H.J., 2015. Ascidians as models for studying
683   invasion success. Mar Biol 162, 2449–2470. https://doi.org/10.1007/s00227-015-2734-5

Figure captions:

Figure 1: Map showing the layout of the 6 geographical sectors (AGM, AGD, VLR, GRU, LEU, CST) and

the 16 sites prospected (indicated in dark blue) for artificial reef sampling. Areas indicated in light

blue corresponded to artificial reef deployment areas. Areas indicated in black corresponded to

natural hard-substrate areas. Some of the AR shape illustrations were taken from Tessier et al., 2014.

Figure 2: Photographs of the five species inventoried on AR *a) Sabella spallanzanii b) Eunicella*

singularis c) Leptogorgia sarmentosa d) Halocynthia papillosa e) *Pentapora fascialis; all © Blouet*

sylvain

Figure 3: Map showing the five species co-occurence assemblages inventoried on ARs in the 16 sites

(AGM1, AGM2, AGD1, AGD2, AGD3, AGD4, AGD5, VLR1, VLR2, GRU1, GRU2, GRU3, LEU1, LEU2, CST1,

CST2) in the Gulf of Lion.

Figure 4: Diversity and dissimilarity. Upper part: diversity of species assemblages in the sampling

units of the five sites (AGD1, AGD2, AGD3, AGD4, AGD5) in AGD sector together with the type of AR,

depth and years of deployment. Lower part: dendrogram obtained by group average clustering

based on the Jaccard dissimilarity index using the presence /absence of species (P=0.04 at 62% of

dissimilarity). The red dotted line separates the two clusters identified by the analysis.

713   **Figures :**

714   **Figure 1 :**

715

716

717

718

719

720

721

722

723

724

725

726

727

728   **Figure 2 :**

A)   B)   C)

D)   E)

742   **Figure 3 :**

Figure 4:

Table captions

Table 1: Life_history traits (spawning period, relative fecundity, life expectancy, age at sexual maturity, larval type, planktonic larval duration (PLD), relative abundance in the GoL) for the five species inventoried on ARs.

Table 2: Results of multivariate NP-Manova testing the effect of the shape between steel cage and pipe on the presence/absence assemblage. Sites tested: VLR1 vs VLR2. Significant (P<0.05) values in bold.

Table 3: Results of multivariate NP-Manova testing the interactive effects of the Age (of deployment) and site (nested in Age) on the presence/absence assemblage. Sites tested: AGD1 vs (AGD2 vs AGD3). Significant (P<0.05) values in bold.

Table 4: Results of multivariate NP-Manova testing the interactive effects of the depth (of deployment) and site (nested in depth) on the presence/absence assemblage. Sites tested: (AGD2 vs AGD3) vs (AGD4 vsAGD5). Significant (P<0.05) values in bold.

Table 5: Results of univariate Kruskall_wallis testing the effect of location of deployment on the presence/absence by species. Sites tested: all sites in AGD. Significant (P<0.05) values in bold.

Table 6: Results of multivariate NP-Manova testing the interactive effects of geographical sector and site (nested in geographical sector) on the presence/absence assemblage. Sites tested: (AGM1 vs AGM2) vs (AGD2 vs AGD3) vs (VLR1 vs VLR2) vs (GRU1 vs GRU2 vs GRU3) vs (LEU1 vs LEU2). Significant (P<0.05) values in bold.

780   Table 7: Results of univariate Kruskall_wallis testing the effects of geographical sector and site on the

781   presence/absence by species. Sites tested: AGM1 vs AGM2 vs AGD2 vs AGD3 vs VLR1 vs VLR2 vs

782   GRU1 vs GRU2 vs GRU3 vs LEU1 vs LEU2. Significant (P<0.05) values in bold.

784   **Tables:**

785   **Table 1:**

Species	Spawning period	Fecundity	Life expectancy	Age at sexual maturity	Larval type	Planktonic larval duration (PLD)	Abundance in the GoL
Leptogorgia sarmentosa	June - August	**	20 years	2-3 years	Lecithotrophic	1-4 weeks (estimated from other gorgonians)	**
Eunicella singularis	July - August	*	25-30 years	6 years	Lecithotrophic	1-4 weeks (in aquarium)	***
Pentapora fascialis	June	**	10 years	2 years	Lecithotrophic	<1 days	**
Sabella spallanzani	January - February	***	5 years	1 years	Lecithotrophic	21 days (in aquarium)	*
Halocynthia papillosa	September - October	*** (estimated from other ascidians)	unknow	2 months (estimated from other ascidians)	lecithotrophic (estimated from other ascidians)	<12 hours (estimated from other ascidians)	**

786   Symbol legend: * = low, **=medium ***=strong

788   **Table 2:**

Multivariate measures					
Source	Df	SS	Ms	F	Pvalue
Assemblage (Presence/Absence)					
Shape	1	3.6408e-19	3.6408e-19	9.8301e-18	0.6090
Residual	4	0.1481	0.0370		
Total	5	0.1481			

792 **Table 3:**

Multivariate measures					
Source	**Df**	**SS**	**Ms**	**F**	**Pvalue**
Assemblage (Presence/Absence)					
Age	1	0.4825	0.4825	1.4330	0.3330
Site(Age)	1	0.3367	0.3367	7.5551	**0.0130**
Residual	6	0.2674	0.0446		
Total	8	1.0866			

795 **Table 4:**

Multivariate measures					
Source	**Df**	**SS**	**Ms**	**F**	**Pvalue**
Assemblage (Presence/Absence)					
Depth	1	0.2491	0.2491	1.3715	0.3280
Site(Depth)	2	0.3633	0.1816	8.5969	**0.001**
Residual	8	0.1690	0.0211		
Total	11	0.7816			

804 **Table 5:**

805

Univariate measures					
Presence/absence					
	Df	SS	Ms	Chi_sq	Pvalue
Eunicella singularis					
	4	127.5	31.875	10.82	**0.0287**
Leptogorgia sarmentosa					
	4	60	15	6.22	0.1832
Pentapora fascialis					
	4	172.5	43.125	11.5	**0.0215**
Sabella spallazanii					
	4	0	0	nan	nan
Halocynthia papillosa					
	4	150	37.5	11.2	**0.0244**

806

807 **Table 6:**

Multivariate measures					
Source	Df	SS	Ms	F	Pvalue
Assemblage (Presence/Absence)					
Geographical sector	4	2.1671	0.5418	5.0994	**0.005**
Site(Geographical sector)	6	0.6375	0.1062	2.7812	**0.007**
Residual	18	0.6876	0.0382		
Total	28	3.4921			

808

809

810

811

812

813

814

815

816 **Table 7:**

	Presence/absence				
Source	Df	SS	Ms	Chi_sq	Pvalue
Eunicella singularis					
Geographical sector	4	447.08	111.77	10.28	**0.036**
Site	9	937.66	93.76	21.56	**0.0175**
Leptogorgia sarmentosa					
Geographical sector	4	965.7	241.42	22.2	**0.0002**
Site	9	1116.5	11.65	28	**0.0018**
Pentapora fascialis					
Geographical sector	4	937.66	234.41	21.56	**0.0002**
Site	9	1077.83	107.78	24.78	**0.0058**
Sabella spallazanii					
Geographical sector	4	0	0	nan	nan
Site	9	0	0	nan	nan
Halocynthia papillosa					
Geographical sector	4	323.59	80.89	6.01	0.1985
Site	9	681.5	68.15	12.53	0.2509

AGM

AGD

VLR

GRU

LEU

CST

Aigues_mortes

Agde

Valras

Gruissan

Leucate

Saint-cyprien

AGM1

AGM2

AGD5

AGD2

AGD3

AGD4

VLR2

VLR1

GRU1

AGD1

GRU3

GRU2

LEU1

LEU2

CST1

CST2

Pentapora fascialis

Eunicella Singularis

Halocynthia papillosa

Leptogorgia sarmentosa

Sabella spallenzanii

GULF OF LION

Artificial reef area prospected

Artificial reef area

Rocky habitat

0 10 20 km

N