Membrane phosphoinositides stabilize GPCR-arrestin complexes and provide temporal control of complex assembly and dynamics

John Janetzko[1], Ryoji Kise[2], Benjamin Barsi-Rhyne[3,4], Dirk H. Siepe[1,5,6], Franziska M. Heydenreich[1], Matthieu Masureel[1,7], Kouki Kawakami[2], K. Christopher Garcia[1,5,6], Mark von Zastrow[3,4], Asuka Inoue[2*], Brian K. Kobilka[1,8*]

Affiliations:

[1]Department of Molecular and Cellular Physiology, Stanford University School of Medicine, Stanford, CA, USA.
[2]Graduate School of Pharmaceutical Sciences, Tohoku University, 6-3, Aoba, Aramaki, Aoba-ku, Sendai, Miyagi, 980-8578 Japan.
[3]Department of Cellular and Molecular Pharmacology, University of California, San Francisco, School of Medicine, San Francisco, CA, USA.
[4]Department of Psychiatry, University of California, San Francisco, School of Medicine, San Francisco, CA, USA.
[5]Department of Structural Biology, Stanford University School of Medicine, Stanford, CA, USA
[6]Howard Hughes Medical Institute, Stanford University School of Medicine, Stanford, CA, USA.
[7]Present address: Department of Structural Biology, Genentech Inc., South San Francisco, CA 94080, USA
[8]Lead contact
*Correspondence: iaska@tohoku.ac.jp (A.I.), kobilka@stanford.edu (B.K.K.)

Keywords: GPCR; arrestin; phosphoinositides; conformational dynamics; endocytosis; fluorescence

Summary:

Binding of arrestin to phosphorylated G protein-coupled receptors (GPCRs) is crucial for modulating signaling. Once internalized some GPCRs may complex with arrestin, while others interact transiently; this difference affects receptor signaling and recycling. Cell-based and in vitro biophysical assays reveal the role of membrane phosphoinositides (PIPs) in arrestin recruitment and GPCR-arrestin complex dynamics. We find that GPCRs broadly stratify into two groups, one requiring PIP-binding for arrestin recruitment and one that does not. Plasma membrane PIPs potentiate an active conformation of arrestin and stabilize GPCR-arrestin complexes by promoting a receptor core-engaged state of the complex. As allosteric modulators of GPCR-arrestin complex dynamics, membrane PIPs allow for additional conformational diversity beyond that imposed by GPCR phosphorylation alone. The dependance on membrane PIPs provides a mechanism for arrestin release from transiently associated GPCRs, allowing their rapid recycling, while explaining how stably associated GPCRs are able to engage G proteins at endosomes.

Introduction:

G protein-coupled receptor (GPCR) activation and deactivation are tightly regulated, allowing them to achieve robust signaling. GPCR deactivation is a complex multi-step process often divided into an acute and a prolonged phase (Rajagopal and Shenoy, 2018). In addition to promoting G protein engagement, agonist stimulation leads to the recruitment of GPCR kinases (GRKs), which phosphorylate the receptor and trigger recruitment of arrestins (Komolov and Benovic, 2018). Arrestin first blocks further G protein engagement, resulting in an acute phase of desensitization, but also mediates the trafficking of activated receptors to clathrin-coated structures (CCSs) and their internalization. Once internalized, receptors can experience markedly

different fates, with some being rapidly recycled to the plasma membrane, while others are retained in intracellular compartments, or directed to lysosomes and degraded (Hanyaloglu and von Zastrow, 2008). In recent years, the discovery that GPCRs can signal from intracellular compartments has led to a re-framing of GPCR signaling to include not only temporal regulation, but also differences that result from spatially distinct receptor populations (Irannejad et al., 2013; Irannejad et al., 2015; Lobingier and von Zastrow, 2019).

There are four human arrestins; arrestins 1 and 4 are dedicated to the visual system, and arrestins 2 and 3, also known as β-arrestin 1 (βarr1) and β-arrestin 2 (βarr2), respectively, are ubiquitously expressed throughout the other tissues. Remarkably, these two β-arrestins are responsible for recognition and desensitization of hundreds of GPCRs. Though most GPCRs recruit arrestin, the nature and duration of this interaction can differ between receptors. Historically GPCRs have been classified as either a "class A" receptor, which interacts transiently with arrestin, or "class B" receptors, which interact more stably with arrestin and co-localize with arrestin in endosomes (Oakley et al., 2001; Oakley et al., 2000; Zhang et al., 1999). This distinction is importantly different from that of family A (rhodopsin-like) and family B (secretin-like) GPCRs. Whether a GPCR interacted transiently or stably with arrestin appears to correlate with rates of re-sensitization, with class A receptors re-sensitizing more rapidly than class B receptors (Oakley et al., 1999). Stable association of arrestin to "class B" GPCRs correlates with the presence of particular phosphorylation site clusters (Oakley et al., 2001); however, it has remained unknown what event precipitates the dissociation of β-arrestins from "class A" receptors to allow their dephosphorylation and recycling.

Early structural studies into GPCR-arrestin complexes suggested that arrestin binds to a GPCR either through only the phosphorylated C-terminus (called tail-engaged), or through both the phosphorylated C-terminus and the transmembrane core of the GPCR (called core-engaged) (Latorraca et al., 2018; Shukla et al., 2014; Staus et al., 2018). However, it remains unclear what determines the equilibrium between these states. While core-engagement is necessary for receptor desensitization (Kumari et al., 2017), it is not required for internalization (Cahill et al., 2017). If, however, complexes can shift from core-engaged to tail-engaged in endosomes it would allow for G proteins to access the receptor core while remaining bound to arrestin. These so-called "megaplex" assemblies (Nguyen et al., 2019; Thomsen et al., 2016) have been implicated in the sustained cAMP signaling produced by endosomal populations of V2R and PTH1R (Feinstein et al., 2013; Ferrandon et al., 2009), both of which stably associate with β-arrestins.

At a molecular level, the prevailing model for arrestin activation, and thus recruitment to an active and phosphorylated GPCR, involves displacement of the auto-inhibitory C-terminus of arrestin by the GPCR phosphorylated C-terminus (or in some cases an intracellular loop). Once the arrestin C-terminus is displaced, additional structural rearrangements occur that allow for arrestin to engage a GPCR (Sente et al., 2018), including insertion of the arrestin finger loop into a cavity formed by the cytoplasmic ends of transmembrane segments. However, arrestin activation functions for more than just GPCR engagement. In its active form, arrestin is able to engage multiple signaling proteins, including JNK3, ERK1/2, p38 (Song et al., 2009) and Src (Pakharukova et al., 2020). It has been suggested that distinct arrestin conformations, which can arise from different inputs (i.e., receptor phosphorylation pattern) may favor interaction with a subset of these signaling partners and affect signaling outcomes downstream of arrestin (Chen et al., 2018; Latorraca et al., 2020). Recently the model for arrestin activation, which suggests a 1:1 interaction, has been challenged by the finding that some "class A" receptors cause the accumulation of super-stoichiometric quantities of arrestin in clathrin-coated structures (CCSs), suggesting an ability to persist at the membrane without an associated GPCR (Eichel et al., 2018;

Nuber et al., 2016). It was speculated that an association with PIP_2 was responsible for retaining
β-arrestins in clathrin coated structures (Eichel et al., 2018); however, based on the established
mechanism for arrestin activation it is unclear how this would be possible, or how arrestin absent
an associated GPCR promoted MAPK signaling from CCSs (Eichel et al., 2016).

Components of the endocytic machinery such as AP2 (Kadlecova et al., 2017), and β-arrestins
(Gaidarov et al., 1999) have been shown to bind to PIPs. These signaling lipids serve critical
functions defining the identity of lipid compartments and acting as coincidence markers for
protein-protein recognition and trafficking to occur only in the appropriate subcellular context (De
Matteis and Godi, 2004; Di Paolo and De Camilli, 2006). While several studies have investigated
the interactions of soluble inositol phosphates with both visual and non-visual arrestins (Chen et
al., 2017; Chen et al., 2021; Milano et al., 2006; Zhuang et al., 2010),there has been only one
where the role of membrane PIPs was explored (Gaidarov et al., 1999). Importantly, this work
suggested that plasma membrane PIPs, such as PI(4,5)P2 and PI(3,4,5)P3, hereafter PIP_2 and
PIP_3, respectively, may function to stabilize GPCR-β-arrestin complexes as they traffic to CCSs.

Recent structural studies showing PIP_2 bound at the interface between the neurotensin type I
receptor (NTSR1) and βarr1 (Huang et al., 2020) prompted us to ask the question: what role do
PIPs serve in mediating GPCR-β-arrestin complex assembly? Here we show that some GPCRs
require PIP binding for β-arrestin recruitment, provided they engage β-arrestins transiently.
Furthermore, we show that the requirement of PIPs depends on specific receptor phosphorylation
sites. Using *in vitro* biochemical and biophysical assays, we demonstrate that phosphoinositide
binding contributes to the stability of the GPCR-β-arrestin complex, where it promotes the core-
engaged state. We also find that PIPs alone promote a partially activated state of arrestin,
providing an explanation for how arrestin can persist in CCSs once dissociated from a GPCR.
Together, these results explain a) how receptors that transiently associate with β-arrestin recruit
and dissociate β-arrestin in a spatiotemporally resolved manner, and b) how strongly coupled
receptors maintain a stable association with arrestin in subcellular compartments yet allow for
further G protein engagement from subcellular structures.

Results and Discussion:

Arrestin PIP-binding is important for desensitization of endogenous β2AR

The PIP-binding-deficient mutant of βarr2 (K233Q/R237Q/K251Q, henceforth 3Q, also used to
denote mutation of the homologous residues in K232Q/R236Q/K250Q in βarr1) was previously
found to be impaired for internalization of β2AR (Gaidarov et al., 1999), with βarr2 (3Q) failing to
traffic to CCSs, though still being recruited from the cytoplasm to the plasma membrane, albeit to
a lesser extent than wild-type (WT) (Eichel et al., 2018). As such, we wondered how this behavior
affects β2AR signaling and specifically whether βarr2 (3Q) is capable of desensitizing β2AR at
the plasma membrane. Using a FRET-based cAMP sensor (Tewson et al., 2016), we monitored
cAMP production in real-time in HEK293 cells lacking both β-arrestins, and endogenously
expressing the β2AR (O'Hayre et al., 2017). In the absence of exogenously expressed βarr2,
isoproterenol (iso) stimulation, via endogenous β2AR, led to a sustained cAMP response, while
expression of βarr2 led to desensitization. However, expression of the 3Q βarr2 mutant resulted
in significantly less desensitization over 30 minutes (Figure S1A); furthermore, this difference was
observed in two independent β-arrestin-deficient cell lines (Luttrell et al., 2018; O'Hayre et al.,
2017). This suggests that the PIP-binding function of β-arrestins plays an important functional

role, in not only internalization (Gaidarov et al., 1999), but also receptor desensitization, and does
so under conditions of endogenous GPCR expression.
152
153   *GPCRs stratify into two groups in their dependence on PIP-binding for arrestin recruitment*
154
That the βarr2 3Q mutant is impaired for recruitment to β2AR, but seemingly not for the chimeric
receptor β2AR-V2C, which bears the C-terminus of the vasopressin V2 receptor (Eichel et al.,
2018), suggested that GPCRs may differ in their dependence on β-arrestin PIP-binding for
recruitment. To investigate a wide range of GPCRs we used a cell-based NanoBiT assay (Dixon
et al., 2016), wherein a plasma membrane localization sequence (CAAX) is fused to the large
subunit of a modified NanoLuc luciferase (LgBiT). Recruitment of either βarr1 or βarr2, which bear
an N-terminal complementary small subunit of NanoLuc (SmBiT), can be monitored by
luminescence changes (Figure 1A). We selected a set of 22 representative GPCRs
(Supplementary Data Table 1), co-expressed the sensors with each receptor of interest in
HEK293 cells, and compared the recruitment of WT β-arrestin to that of the corresponding 3Q β-
arrestin mutant upon agonist stimulation (Figure 1B, top, Supplementary Data Table 1). We used
luminescence fold-change measured over the range of 10-15 minutes post-agonist stimulation for
our end-point values and fit the resulting data to generate concentration response curves and
extract a recruitment amplitude for each receptor-arrestin pair (see methods) (Figure 1B, bottom,
Supplementary Data Figure 1A-B). We then compared the recruitment of WT and 3Q arrestin
using a metric that represented the relative sensitivity of the receptor to loss of arrestin-PIP
binding capacity, we termed the loss of function (LOF) index (see methods). Receptors with a low
LOF value recruit WT and 3Q β-arrestins to the plasma membrane similarly, and are deemed
PIP-independent, while receptors with a high LOF value show greatly diminished recruitment of
3Q β-arrestin and are deemed PIP-dependent (Figure 1C). Both WT and 3Q forms of βarr1 and
βarr2 express similarly (Figure S1B, Supplementary Data Figure 2).
176
Though receptors spanned a continuum of LOF values, they seemed to cluster into two groups
near the ends of the scale. To examine this, we performed k means clustering of plasma
membrane recruitment data for all GPCR-β-arrestin pairs (Supplementary Data Figures 1 and 3,
n = 55 pairs), which suggested that the data is best divided into two clusters (see methods,
clusters marked by dotted ellipses in Figure 1C). We found only a weak inverse correlation
between the amplitude of WT arrestin recruitment and the degree of LOF observed (Pearson
correlation = -0.51; -0.4 when TACR1 and B2R are excluded) (Figure S1C), suggesting that
differences in LOF were not due to lower levels of WT recruitment. Cluster 1 was defined by
receptors that exhibited a high degree of LOF (center LOF = 0.73) and included GPCRs previously
classified as "class A" (Oakley et al., 2000): β2AR, μOR, ETAR, D1R, α1BR. Cluster 2, defined
by receptors with a low degree of LOF (center LOF = 0.06), included GPCRs classified as "class
B" (Oakley et al., 2000): AT1R, NTSR1, V2R, TRHR, and TACR1. We also tested two chimeric
receptors, β2AR-V2C and μOR-V2C (Eichel et al., 2018), both of which showed reduced reliance
on β-arrestin PIP-binding capability for plasma membrane recruitment compared to the respective
parent receptor. The V1AR, which was previously shown to undergo labile phosphorylation and
rapid recycling (Innamorati et al., 1998a; Innamorati et al., 1998b) clusters with the class A
receptors in cluster 1, while the V1BR, bearing a closer similarity in its proximal C-terminus to
V2R clusters with class B receptors in cluster 2, even though it has been found to only associate
transiently with arrestin (Perkovska et al., 2018). In addition, β1AR, S1PR1, and δOR, all three of
which have been shown to either recycle rapidly or interact transiently with arrestin (Martinez-
Morales et al., 2018; Nakagawa and Asahi, 2013; Trapaidze et al., 2000), were assigned to cluster
1. Other receptors known to co-localize with arrestin at endosomes, including PAR2 (DeFea et
al., 2000; Dery et al., 1999; Oakley et al., 2001), B2R (Khoury et al., 2014), and PTH1R (Feinstein

et al., 2011) were also classified into cluster 2. Two receptors, OXTR and HTR2C displayed
unexpected behavior where βarr1 recruitment was dramatically more sensitive to loss of PIP-
binding than βarr2, resulting in these GPCR-β-arrestin pairs being divided between the two
clusters. OXTR was previously classified as a "class B" receptor (Oakley et al., 2001); however,
these studies only examined βarr2 recruitment. For the serotonin 2C receptor (HTR2C), previous
studies showed PIP$_2$-depletion did not block recruitment of βarr2 (Toth et al., 2012), and we
observed an intermediate LOF value for βarr2 plasma membrane recruitment with HTR2C.
Together, these data show that recruitment of β-arrestins is dependent on the PIP-binding
capacity of arrestin for some GPCRs, but not others, and that this distinction is consistent with
the previous class A/B categorization based on microscopy co-localization studies.
210
While our use of a plasma membrane localized LgBiT avoids modifying the receptor of interest,
we wanted to confirm that plasma membrane recruitment is indeed a reliable proxy for arrestin
recruitment to a GPCR of interest. For this, we used a direct NanoBiT assay in which the SmBiT
component is fused to the C-terminus of each GPCR of interest, and the N-terminus of arrestin is
modified with the LgBiT fragment (Figure S1D, left). We found that recruitment measured by direct
complementation largely paralleled recruitment measured using the plasma membrane
bystander, with minor exceptions (Supplementary Data Figure 4-5). Further, directly comparing
LOF as measured by the plasma membrane bystander to that of the direct complementation
showed a strong positive correlation (Pearson correlation = 0.88), suggesting that β-arrestin
recruitment measured through the plasma membrane bystander was indeed a faithful metric
(Figure S1D, right). The most extreme outlier, HTR2C, showed βarr2 recruitment is PIP-binding
independent as measured by the direct recruitment assay, as compared to being partially PIP-
binding-dependent when measured using the plasma membrane bystander. As the direct
recruitment assay seemed to better match prior findings for the HTR2C (Toth et al., 2012), we
wondered why this might be. In addition to HTR2C, α1BR and β1AR also exhibit reduced PIP-
binding sensitivity in the direct recruitment assay for βarr2. Curiously, all three of these receptors
exhibit some level of Gq coupling (Inoue et al., 2019). We speculate that for cluster 1 receptors,
such as HTR2C, which are Gq-coupled, that their dependence on PIP-binding for arrestin
recruitment to the plasma membrane may be amplified by local PIP$_2$-depletion via phospholipase
C upon stimulation.
231
Given that "class B" receptors co-localize with β-arrestins in endosomes, we wondered whether
PIP binding affected this process. We used the FYVE domain of endofin as an endosome
bystander (Endo) (Namkung et al., 2016), which we fused to LgBiT (termed endo-Lg) to monitor
recruitment of arrestin bearing an N-terminal SmBiT (Figure S1E), as was done for plasma
membrane recruitment. Since both βarr1 and βarr2 displayed largely similar behavior in our
plasma membrane recruitment assay, we focused on βarr1 for these experiments (Supplementary
Data Figure 6A); however, βarr2 recruitment was also examined for a subset of receptors
(Supplementary Data Figure 6B). This assay robustly detected endosomal translocation as all
receptors known to co-localize with β-arrestin in endosomes did so (Figure S1E). Consistent with
the stark difference between βarr1 and βarr2 observed for OXTR (Figure 1C), we observed
measurable endosomal association of βarr2, but weak and barely measurable βarr1 endosome
recruitment (Supplementary Data Figure 6C). Our results were consistent with prior microscopy-
based approaches (Oakley et al., 2000), thereby validating our NanoBiT assay. In contrast,
HTR2C showed more robust recruitment of βarr1 than βarr2 (Supplementary Data Figure 6D). As
expected, while cluster 1 receptors whose ability to co-localize with β-arrestins at endosomes had
not yet been described displayed little signal for endosomal translocation for WT and 3Q β-

arrestins, the cluster 2 receptors showed robust signal for recruitment of both WT and 3Q β-
arrestins.
250
Though end-point recruitment of βarr1 to NTSR1 and other cluster 2 GPCRs was largely
unaffected by loss of the PIP-binding site, prior NTSR1 experiments had found that loss of PIP
binding slowed the kinetics of β-arrestin recruitment (Huang et al., 2020), suggesting PIP_2 may
play a role in the complexes formed with cluster 2 receptors, even when end-point recruitment is
unchanged. We fit the rate of β-arrestin translocation to the plasma membrane in response to
stimulation for all GPCRs in cluster 2 using our CAAX bystander NanoBiT assay (Figure 1B, top).
As was seen for NTSR1, other cluster 2 GPCRs showed a slower association for 3Q than WT
(Figure S2A). Though the magnitude of the effect varied across receptors (Figure S2B), these
results clearly show that even recruitment to cluster 2 GPCRs is impacted by loss of PIP-binding
in β-arrestins.
261
Together, these results provide several major findings. First, though the tested GPCRs can be
divided into two groups, the reliance on PIP-binding for arrestin recruitment is very much a
continuum. Generally, GPCRs that co-localize with β-arrestins at endosomes do not require the
PIP-binding capacity of β-arrestins for plasma membrane recruitment and are henceforth referred
to as PIP-independent GPCRs. Secondly, though PIP-independent GPCRs retained the ability to
recruit β-arrestins, the kinetics of recruitment is impaired by loss of PIP binding, suggesting that
PIP-mediated interactions likely function to stabilize GPCR-arrestin complexes across all
receptors. Finally, while βarr1 and βarr2 behave similarly for most GPCRs, there were exceptions;
in much the same way that a continuum of LOF values was observed, this suggests that GPCR-
arrestin complexes are incredibly diverse both in their sensitivity to allosteric inputs and possibly
their conformational landscape.
273
274 *Receptor phosphorylation patterns determine the dependence on PIP-binding by arrestin*
275
The distinction between class A and class B receptors was previously attributed to the presence
of suitably positioned clusters of phosphosites in the receptor C-terminus (Oakley et al., 2001).
We reasoned that there must be a degree of phosphorylation required to overcome the
dependence on arrestin-PIP binding for recruitment to class A receptors. We chose the NTSR1
as a model receptor since WT NTSR1 stably associated with arrestins and the major
phosphorylation cluster responsible for this phenomenon was previously established for the rat
ortholog (Oakley et al., 2001). Using human NTSR1, we designed a set of phosphorylation-
deficient mutants, including both the C-terminus and the third intracellular loop (ICL3) (Figure 2A).
In the recent NTSR1-βarr1 structure (Huang et al., 2020), ICL3 was found to be phosphorylated
and appeared to make contacts to arrestin, though the role of ICL3 phosphorylation in arrestin
recruitment had not been explored. NTSR1 contains four S/T residues in ICL3, three of which are
clustered, and 9 S/T residues in its C-terminus, 6 of which are divided into two clusters. To
compare the PIP-dependence of NTSR1 phosphorylation mutants (Figure 2A) for their ability to
recruit βarr1 to the plasma membrane, we used the CAAX bystander NanoBiT assay (Figure 1A).
We first measured cell-surface expression of the NTSR1 constructs and found similar levels
(Figure S3A), except for ICL3-4A, which showed slightly reduced expression. Regardless, in this
range of receptor expression recruitment signal is saturated with respect to NTSR1 and these
minor differences in expression are unlikely to affect the assay response (Figure S3B-C). Though
WT NTSR1 is classified as a PIP-independent receptor, NTSR1 phosphorylation mutants could
either be classified into cluster 1 or cluster 2 (Figure 2A, Supplementary Data Figure 3),
suggesting that particular phosphorylation mutants rendered arrestin recruitment to NTSR1 PIP-
dependent. Removal of the two C-terminal phosphorylation site clusters (NTSR1-6A, NTSR1-

10A) resulted in a dramatic reduction in arrestin recruitment (Supplementary Data Figure 3), with
remaining arrestin recruitment being largely PIP-dependent (Figure 2A). Removal of the ICL3
phosphorylation sites did not affect PIP-dependence (NTSR1-ICL3-4A); neither did removal of
the proximal phosphorylation cluster (NTSR1-A^{401}VAA), nor removal of any one residue in the
distal cluster (NTSR1-TLSA, NTSR1-ALSS, NTSR1-TLAS). However, removal of the distal
phosphorylation cluster (NTSR1-A^{407}LAA) led to a dramatic reduction in recruitment, and an
increase in PIP-dependence, consistent with findings that the distal cluster in the rat ortholog is
necessary for stable arrestin association (Oakley et al., 2001). NTSR1-5A, bearing a single C-
terminal phosphorylation site in the distal cluster, showed PIP sensitivity comparable to NTSR-
ALAA, while NTSR1-4A with two distal cluster phosphorylation sites showing much less PIP-
dependence, suggesting that two phosphorylation sites are sufficient to overcome the need for
PIP binding. Similarly, NTSR1-TLAA, which differs from NTSR1-5A only in the addition of the
proximal cluster of phosphosites exhibits sensitivity between the NTSR1-5A and NTSR1-4A
constructs, suggesting that a phosphorylation site from the proximal cluster may offer a partial
rescue for the absence of one in the distal cluster.

As the plasma membrane bystander recruitment assay suggested that two phosphorylation sites
were necessary to overcome the PIP-dependence on arrestin recruitment, we wondered whether
this behavior coincided with the ability of arrestin to be recruited to endosomes. We monitored
translocation of arrestin to endosomes using the endosome bystander NanoBiT assay (Figure
S1E). As expected, NTSR1-ALAA (Oakley et al., 2001) as well as NTSR1-6A and NTSR1-10A
failed to recruit arrestin to endosomes (Figure 2B, Supplementary Data Figure 7). A single C-
terminal phosphorylation site (NTSR1-5A) was insufficient to promote arrestin traffic to
endosomes; however, two phosphorylation sites in the distal cluster (NTSR1-4A) were sufficient
to promote endosomal translocation. There was a further increase in recruitment when the
proximal sites were returned (NTSR1-TLSA), suggesting an additional contribution from this
region strengthens the interaction between NTSR1 and arrestin. Further support for a contribution
from the proximal cluster stems from the difference between NTSR1-5A and NTSR1-TLAA, which
differ in the presence of the proximal phosphorylation cluster and show a marked difference in
both targeting of arrestin to endosomes, as well as PIP-dependence (Figure 2B). Within the distal
cluster, any two phosphorylation sites were sufficient, and having the third present appeared to
offer no additional benefit (NTSR1-AVAA compared to NTSR1-ALSS, NTSR1-TLAS and NTSR1-
TLSA) (Figure 2B).

Given that two phosphorylation sites in the distal cluster were sufficient for both PIP-insensitivity
for plasma membrane recruitment, and recruitment of arrestin to endosomes, we asked whether
two phosphorylation sites were also sufficient for receptor internalization. We measured
internalization of the NTSR1 constructs in β-arrestin-deficient HEK293 cells where either WT or
3Q βarr1 was reintroduced. WT NTSR1 was robustly internalized by both WT and 3Q βarr1. In
contrast, NTSR1-5A showed a significant difference in internalization between WT and 3Q βarr1,
while NTSR1-4A showed no difference in internalization between WT and 3Q βarr1. The trend
between NTSR1-5A and NTSR1-4A parallels that seen for β2AR and β2AR-V2C (Figure 2C),
supporting our finding that two phosphorylation sites are sufficient for robust internalization that
is PIP-independent. In addition, the internalization observed for NTSR1-5A by WT βarr1 suggests
that the lack of endosome recruitment observed for this construct (Figure 2B) is due to weakened
GPCR-βarr interaction and not simply a lack of internalization for this receptor (Figure 2C).

Together, these data show that two suitably positioned phosphorylation sites are sufficient to
render β-arrestin recruitment PIP-independent and allow for robust arrestin-dependent
internalization as well as support arrestin translocation to endosomes. Furthermore, they show

that NTSR1, a receptor that recruits β-arrestin in a PIP-independent manner, can become PIP-
dependent by changes in receptor phosphorylation. Given that GPCRs, such as the μOR, have
different phosphorylation patterns depending on the stimulating agonist (Just et al., 2013), we
speculate that the resulting β-arrestin complexes may have drastically different behavior in cells.
352
353   *PIP$_2$ binding affects complex stability and tail-core equilibrium in vitro*
354
As PIP-binding was previously suggested to stabilize the interaction between a GPCR and
arrestin (Gaidarov et al., 1999), based on experiments in cells, we wanted to explicitly test this *in*
vitro. Using NTSR1 as our model receptor, where PIP-binding was not strictly necessary for
recruitment in cells, we compared the ability of GRK5 phosphorylated NTSR1 to form a complex
with βarr1 (WT or 3Q mutant) in the presence of a soluble PIP$_2$ derivative, diC8-PI(4,5)P2
(henceforth PIP$_2$), by size-exclusion chromatography (Figure 3A-B) (Huang et al., 2020). While
complexing with full-length WT βarr1 led to about 25% complex formation, use of 3Q βarr1
resulted in <5% complex formation (Figure 3C). Use of a C-terminally truncated βarr1 (1-382) led
to a more than 2-fold enhancement in complex formation, which was only slightly reduced with
the corresponding 3Q arrestin. Using the LOF metric developed to evaluate the impact of PIP-
binding on arrestin recruitment in cells, we found that full-length arrestin showed a greater degree
of LOF than C-terminally truncated arrestin, suggesting that removal of the arrestin C-terminus is
largely able to overcome the impairment in complexing that results from the 3Q mutation (Figure
S4A). PIP$_2$ affinity *in vitro* was reduced 20x for 3Q βarr1 compared to WT (Figure S4B). Since
arrestin activation is understood to proceed via release of its auto-inhibitory C-terminus (Sente et
al., 2018; Shukla et al., 2013), we wanted to rule-out the possibility that 3Q βarr1 complexing
efficiency is simply reduced due to a lack of arrestin C-terminus release. We designed a Förster
Resonance Energy Transfer (FRET) sensor to report on arrestin C-terminus release (Figure S4C):
using a cysteine-free βarr1 construct, we introduced two new cysteine residues at positions 12
and 387 – βarr1 (12-387) – to allow for selective labeling of these positions with a suitable dye
pair. Given that the expected change in distance between the bound and unbound C-terminus
was ~40 Å (Chen et al., 2017; Kim et al., 2012; Zhuo et al., 2014), we used an AlexaFluor 488/Atto
647N FRET pair, which offers a relatively short Förster radius (R$_0$ ~50 Å). GRK5-phosphorylated
NTSR1 robustly displaced the C-terminus of both WT and 3Q βarr1 (12-387) (Figure S4D).
Displacement was comparable to that seen for a saturating concentration of a peptide
corresponding to the phosphorylated C-terminus of the vasopressin 2 receptor (henceforth
V2Rpp) known to completely displace the arrestin C-terminus (Shukla et al., 2013). GRK5-
phosphorylated NTSR1 fully displaced the βarr1 C-terminus with 10x greater potency than
V2Rpp, suggesting an enhanced affinity for an intact receptor compared to a phosphopeptide
alone. These data show that not only does *in vitro* phosphorylated NTSR1 fully displace the
arrestin C-terminus, but with higher efficacy than an equimolar concentration of phosphopeptide
(even in the presence of unphosphorylated NTSR1), and this is independent of the PIP-binding
ability of arrestin.
388
We reasoned that the reduced complexing efficiency of 3Q βarr1 may be due to differences in the
proportion of core-engaged complex being formed (Figure 3D). To test this hypothesis, we used
an environmentally sensitive bimane fluorophore (bim) site-specifically installed at L68 (L68bim)
on the arrestin finger loop, a region that upon formation of a core-engaged complex with an active
GPCR becomes buried within the receptor TM core. Such a sensor had previously been used to
report on core-engagement for rhodopsin/arrestin-1 (Sommer et al., 2005, 2006), whereupon
receptor core-engagement a blue-shift and an increase in fluorescence emission occurs, owing
to the bimane probe moving into a lower polarity environment within the receptor TM core.
397

While addition of V2Rpp to βarr1 L68bim leads to C-terminus release and a ~50% increase in
bimane fluorescence as seen previously (Latorraca et al., 2020), we speculated that the addition
of receptor may further increase this signal (Sommer et al., 2006). We compared the fluorescence
changes of βarr1 L68bim (WT or 3Q) upon addition of NTSR1 that was either dephosphorylated
or phosphorylated by GRK5 (Figure 3E). In the absence of NTSR1 phosphorylation, there was no
increase in fluorescence; however, phosphorylated NTSR1 led to a ~2-fold enhancement in
fluorescence intensity for WT, but a smaller 1.5-fold enhancement for 3Q. The addition of V2Rpp
at a saturating concentration to the unphosphorylated NTSR1 did not result in a significant
increase over phosphopeptide alone, consistent with the behavior observed for C-terminus
release (Figure S4D). Importantly, for the 3Q βarr1, GRK5 phosphorylated NTSR1 did not elicit a
response different from V2Rpp alone.

Given that the complex exists as a dynamic equilibrium between three states (Figure 3D):
dissociated, tail-bound and core-engaged. We reason that if PIP-binding serves to stabilize the
core-engaged state then loss of PIP binding would bias the equilibrium towards a tail-engaged
state (Figure 3F), which should have a similar spectroscopic signature to V2Rpp alone. Taken
together, these data suggest a model of complex assembly where release of the arrestin C-
terminus by the phosphorylated GPCR C-terminus is rapid, and reversible. The resulting tail-
bound state is in equilibrium with a core-engaged state, where arrestin-PIP binding stabilized this
state and thereby slows dissociation. In the context of full-length arrestin, destabilization of core-
engaged state in the 3Q mutant leads to a reduction in complex stability, presumably due to
arrestin C-tail-mediated dissociation from the tail-bound state. Consistent with these findings
when the arrestin C-terminus is removed the reduced core-engagement of the 3Q mutant does
not impact complexing efficiency due to an increased stability of the tail-bound state (as seen in
Figure 3C, S4A).

PIP$_2$, in the absence of a GPCR, triggers conformational changes in arrestin

The finding that β-arrestins in CCSs, even in the absence of an associated GPCR, signal through
MAPK (Eichel et al., 2016) suggested that arrestin can adopt an active-like conformation without
a GPCR C-terminus to displace its own C-terminus. While PIP$_2$ was proposed to maintain the
membrane association of β-arrestins (Eichel et al., 2018), the impact of this association on the
conformational landscape of β-arrestins, and thus their ability to engage downstream signaling
partners was unknown (Ranjan et al., 2017). Having shown that PIP-binding affects the dynamics
of NTSR1-βarr1 complexes *in vitro*, we wondered whether PIPs in the absence of an associated
GPCR could also affect the conformation of βarr1. We compared the effect of PIP$_2$ to the V2Rpp
for promoting conformational changes in arrestin using FRET and fluorescence reporters on the
finger loop, gate loop, and C-terminus (Figure 4A).

Both the finger loop (Figure 4B, Figure S5A-B) and the gate loop (Figure 4C, Figure S5C-D)
showed saturable conformational changes upon addition of PIP$_2$ which were smaller than those
seen for V2Rpp. Further, the corresponding 3Q mutants did not show PIP$_2$-induced
conformational changes, though they responded to V2Rpp similarly to WT protein. These data
suggest that binding of PIP$_2$ to the arrestin C-lobe allosterically promotes conformational changes
in key arrestin regions involved in GPCR recognition and activation. As the accepted mechanism
for arrestin activation begins with release of its autoinhibitory C-terminus (Sente et al., 2018), we
wondered whether these conformational changes were the result of allosterically promoted C-
terminus release. Using our βarr1 C-terminus FRET sensor (Figure S4C) we found that PIP$_2$
promoted a small movement of the arrestin C-terminus (Figure 4D), but only at concentrations
higher than those needed to saturate the responses seen for either the finger or gate loop sensors

(Figure 4B-C). As was the case for the other sensors, this FRET change in response to PIP$_2$ is
absent in the corresponding 3Q mutant (Figure S5E-F). This finding is consistent with recent
DEER experiments that found little or no C-terminal displacement for βarr1 with IP6 (Chen et al.,
2021). We reason that the conformational changes in the finger and gate loops observed together
with the small FRET change in response to PIP$_2$ could either be due to a change in the equilibrium
of active-inactive βarr1, or a population of an intermediate state of arrestin bearing a change in
position or orientation of the arrestin C-terminus within the arrestin N-lobe. As different membrane
PIPs serve as markers for different subcellular locations, we measured the ability of other PIPs
(Figure S6A) to promote conformational changes in the βarr1 finger loop. Like PI(4,5)P2,
PI(3,4)P2, PI(3,5)P2 and PI(3,4,5)P3 all elicited an increase in bimane fluorescence (Figure S6B-
F). In contrast, PI(4)P showed a weaker response and PI(3)P and PG did not increase
fluorescence of the bimane reporter (Figure S6G-I). Interestingly, these results showed that
plasma membrane resident PIPs (Di Paolo and De Camilli, 2006), PI(4,5)P2, PI(3,4)P2,
PI(3,4,5)P3 and to a lesser extent PI(4)P were able to promote this conformational change in
βarr1, but the early endosomal marker PI(3)P was unable to do so (Figure S6J). PI(3,5)P2 showed
a similar effect to other PIP$_2$s, but is understood to be rare within cells (Hasegawa et al., 2017).
Based on contacts observed in the NTSR1-βarr1 structure (Huang et al., 2020), we speculate that
PIPs bearing adjacent phosphates on the inositol ring may be necessary for chelation of K232
and R236/K250 (Figure S6K); however, a phosphate at the 4-position is sufficient to coordinate
K250 and R236, explaining the small effect seen for PI(4)P. Together, these data show that
different PIP$_2$ derivatives are capable of promoting conformation changes in βarr1, while PIPs
bearing a single phosphate do not, raising the possibility of compartment-specific differences in
the behavior of GPCR-arrestin complexes.
471
472 *PIP$_2$ increases the population of active arrestin*
473
While our fluorescence experiments support PIP$_2$-promoted conformational changes consistent
with arrestin activation, the lack of C-terminus release raised questions of whether these
conformational changes truly reflected an increase in the population of active arrestin, as would
be detected by arrestin binding partners. While the active form of arrestin is understood to mediate
signaling via interactions with a number of protein partners, including MAPK, ERK, SRC (Ranjan
et al., 2017; Reiter et al., 2012), there has been speculation that the binding of a particular partner
might be mediated by a distinct arrestin conformation. We reasoned that the global activation
state of arrestin could be probed using an engineered Fab (Fab30), which has a high-affinity for
the active (V2Rpp-bound) state of βarr1 (Shukla et al., 2013). Fab30 has found utility in a number
of structural studies (Lee et al., 2020; Nguyen et al., 2019; Shukla et al., 2013; Shukla et al., 2014;
Staus et al., 2020), functional studies (Cahill et al., 2017; Ghosh et al., 2019; Kumari et al., 2016;
Latorraca et al., 2020; Thomsen et al., 2016) and more recently it has been adapted as a single-
chain intrabody (IB30) for the detection of active βarr1 in cells (Baidya et al., 2020a; Baidya et al.,
2020b).
488
We used Surface Plasmon Resonance (SPR) to measure binding of Fab30 to immobilized βarr1
(Figure 5A). To confirm the immobilized arrestins behave as expected, we tested binding of
V2Rpp and Fab30+V2Rpp (Figure 5B, Figure S8, Supplementary Data Tables 2-3). Though
selected for binding to the V2Rpp-bound state of βarr1, Fab30 bound to βarr1 weakly in the
absence of V2Rpp, presumably due to a small equilibrium population of active-like arrestin
(Latorraca et al., 2018). Interestingly, binding was enhanced when Fab30 was co-injected with
PIP$_2$ (Figure 5B). This suggested that PIP$_2$ increased the proportion of arrestin in an active-like
state which can be recognized by Fab30, consistent with our fluorescence experiments that
support PIP$_2$ playing a role in arrestin activation. We compared the effect of different additives on

Fab30 binding to WT βarr1, but also a βarr1 3Q mutant, and the pre-activated C-terminally
truncated βarr1(1-382) (Kim et al., 2013). At 1 μM, Fab30 showed 10.2 ± 0.9% (of maximal)
binding to WT βarr1, compared to 56.8 ± 2.0% binding for βarr1 (1-382) (Figure 5C). This suggests
that Fab30 binding is favored by a conformation accessible to WT βarr1, but greatly enhanced by
removal of the arrestin C-terminus. When Fab30 is co-injected with a saturating concentration of
PIP_2 (40 μM), binding to WT βarr1 increased more than 3-fold, to 33.9 ± 1.8%, compared to Fab30
alone. PIP_2 had a smaller effect on the pre-activated βarr1(1-382), but still increased binding from
56.8% to 65.9 ± 0.8%. Titration experiments showed that specific enhancement of Fab30 binding
in the presence of PIP_2 was most pronounced for WT βarr1 (Figure S7G-L). While all three arrestin
constructs showed an increase in Fab30 binding in the presence of PIP_2, the degree of binding
enhancement drastically shifted for WT βarr1 above the K_d for Fab30, but not for either 3Q or (1-
382) βarr1 (Supplementary Data Figure 8). This suggests that while PIP_2 enhanced the population
of active-like βarr1, Fab30 binding remains rate-limiting. Since removal of the βarr1 C-terminus
abrogates the PIP_2-enhancement of Fab30 binding, we reason that PIP_2 acts in *cis* with C-terminal
displacement, consistent with our FRET experiments that showed a PIP_2-induced movement of
the βarr1 C-terminus. To determine whether this effect was specific for PIP_2, we compared the
ability of PG and PI(3)P to enhance binding of Fab30 . Both showed a small enhancement in
Fab30 binding, but significantly less than that seen with PIP_2 (Figure 5C). Further, βarr1 3Q
showed no difference between PG, PI(3)P and PIP_2, suggesting that while anionic lipids weakly
increase Fab30 binding to βarr1, PIP_2 was unique in affecting a specific increase in Fab30 binding.
Both PG and PI(3)P did not enhance Fab30 binding to βarr1 (1-382).
519
Based on these data we propose that spontaneous activation of arrestin to an active-like state
capable of binding Fab30 is possible but rare in the absence of arrestin inputs (Figure 5D). V2Rpp
dramatically shifts the equilibrium towards the active-state by displacement of the arrestin C-
terminus, and removal of the arrestin C-terminus alone is sufficient to greatly enhance the active-
population, even in the absence of V2Rpp or PIP_2. Unlike V2Rpp, which displaces the arrestin C-
terminus, PIP_2 is unable to displace the arrestin C-terminus directly, but able to allosterically move
it. While PIP_2 may stabilize the same active state of arrestin achieved with V2Rpp, albeit to a
lesser extent, it may also act to stabilize an active-like state of arrestin that is on-pathway towards
activation and capable of binding Fab30, though to a lesser extent than V2Rpp-bound βarr1.
Further studies will be necessary to distinguish these possibilities.
530
531   **Conclusions:**
532
533   Our results reveal new molecular details underpinning the regulation of arrestin recruitment to
534   GPCRs, and how spatial and temporal control of GPCR-β-arrestin complexes may occur within a
535   cell.
536
Our findings offer a molecular basis for understanding the phenotypic classification of GPCRs into
"class A" or "class B" for arrestin recruitment. In our model (Figure 6), we refer to "class A" and
"class B" GPCRs as "PIP-dependent" and "PIP-independent", respectively. "PIP-dependent"
GPCRs (Figure 6, left) require the coincident detection of membrane PIPs for recruitment to an
activated and phosphorylated GPCR. We speculate that this is due to an insufficiency in
phosphorylation of these receptors, requiring either an allosteric priming of C-terminus release by
plasma membrane PIPs, or the simultaneous action of both phosphate-mediated contacts and
PIP-mediated contacts to form a sufficiently long-lived complex for effective receptor
desensitization, sequestration and internalization. As some PIP-dependent GPCRs can recruit
arrestin in a C-terminus-independent manner, we consider that release of the arrestin C-terminus
may not be necessary for arrestin function in the context of these receptors. A further trait of these

PIP-dependent GPCRs is that they exhibit, to a varying degree, the "catalytic activation"
phenotype (Eichel et al., 2018) wherein arrestin, after recruitment to an active GPCR, loses
association with the GPCR but remains at the plasma membrane and concentrates at CCSs. This
can be explained by the increasing concentration gradient of PIP_2 leading into the CCS (Sun et
al., 2007) along with our biophysical evidence that PIP_2 promotes conformational transitions
associated with activation. Once a GPCR cargo has been translocated into a CCS, clathrin-
mediated endocytosis (CME) proceeds and PIP_2 levels drop. We suggest that this may serve as
the timing component for arrestin dissociation from these PIP-dependent GPCRs (Zhang et al.,
1999). Presumably, once arrestin has dissociated, the receptor is susceptible to
dephosphorylation, and upon arrival at early endosomes is able to be sorted for rapid recycling
(Krueger et al., 1997). In contrast, "PIP-independent receptors" (Figure 6, right panel) possess
phosphorylation sites which alone can promote a stable association with arrestin, without the need
for membrane PIPs. Since PIP-binding is not necessary to maintain the GPCR-arrestin
association, arrestin co-localizes with these receptors at endosomes. Whether this co-localization
is the product of PIP-independent GPCRs being able to recruit β-arrestins when at endosomes
or by forming a sufficiently stable complex to allow for co-trafficking from the plasma membrane
without exchange remains to be shown.

One question this model raises is: if PIP_2 promotes partial activation of β-arrestins, why is it that
β-arrestins are not basally associated with the plasma membrane? Consistent with our SPR
experiments, and the finding that GPCR C-terminal phosphorylation is required for arrestin
accumulation at CCSs (Eichel et al., 2018), we speculate that PIP_2 binding may occur when
arrestin transitions to an active-like conformation and stabilize this state. We speculate that in
cells, engagement with a GPCR may be necessary to facilitate PIP_2 binding. GPCRs have been
shown to associate with PIP_2 in the local membrane environment (Song et al., 2019; Yen et al.,
2018), and in doing so may act to "load" PIP_2 onto the arrestin, which either can remain associated
with the GPCR or diffuse along the membrane.

These data suggest that while PIP-mediated contacts are not necessary to maintain association,
they likely affect the equilibrium of core vs. tail-engaged states of the complex. Tail-engagement
has been shown to be sufficient for MAPK signaling downstream of β-arrestin (Kumari et al.,
2017). We speculate that this shift in equilibrium, particularly in the context of endosomes defined
by PI(3)P, may explain how PIP-independent receptors, such as V2R and PTH1R are able to
engage and signal through both β-arrestin and G proteins simultaneously in a so-called
"megaplex" assembly (Nguyen et al., 2019; Thomsen et al., 2016).

To-date four structures of GPCR-βarr1 complexes have been described, all of which show arrestin
in a core-engaged state (Huang et al., 2020; Lee et al., 2020; Staus et al., 2020; Yin et al., 2019),
but only one had PIP_2 bound at the interface (Huang et al., 2020). Interestingly, this NTSR1-βarr1
complex with PIP_2 bound used the native NTSR1 C-terminus and did not use Fab30 to stabilize
the complex. We speculate that Fab30 plays a particularly important role in stabilizing the receptor
core-engaged complex (Shukla et al., 2014).

Overall, our data offer a parsimonious explanation for several phenotypic behaviors observed for
GPCR-β-arrestin complexes and provide a biophysical framework for understanding the interplay
between phosphorylation-mediated and PIP-mediated contacts in complex assembly. A reliance
on PIPs for arrestin recruitment offers a robust solution for recruitment of arrestin to receptors
with spatial control, and temporal precision. Given the interplay between PIP-dependent
recruitment and phosphorylation, we believe that distinct signaling outcomes may not only be due
to differences in phosphorylation alone (Latorraca et al., 2020), but rather that these differences

may be further fine-tuned by membrane PIPs that are present in distinct subcellular locations,
adding yet another layer of complexity to our understanding of GPCR signaling.
600

Acknowledgements:

We thank Betsy White for technical laboratory assistance; Shoji Maeda for providing purified
Fab30 protein; Weijiao Huang for assistance with NTSR1 expression and purification; Daniel
Hilger for purified BirA enzyme; Yoon Seok Kim and Eamon Byrne for assistance with FSEC
experiments, Kayo Sato, Yuko Sugamura, Shigeko Nakano and Ayumi Inoue for plasmid
construction and cell based GPCR assays. We thank R. Scott Prosser for additional comments
on the manuscript. This work was supported in part by National Institutes of Health grants
R01NS028471 (B.K.K.), R01 AI125320 (K.C.G), R01DA010711 and R01DA012864 (M.vZ.).
Additional support to both K.C.G and B.K.K. is provided by the Mathers Foundation. B.K.K. is a
Chan-Zuckerberg Biohub Investigator. J.J. is a Damon Runyon Fellow supported by the Damon
Runyon Cancer Research Foundation (DRG-2318-18). B.B.R. is a recipient of an American Heart
Association Predoctoral Fellowship (19PRE34380570). F.M. H. is a recipient of a Marie
Skłodowska-Curie Individual Fellowship from the European Union's Horizon 2020 research and
innovation programme (grant agreement No. 844622) and an American Heart Association
postdoctoral fellowship (19POST34380839). M.M. was supported by an American Heart
Association postdoctoral fellowship (17POST33410958). A.I. was funded by the PRIME
19gm5910013, the LEAP 20gm0010004 and the BINDS JP20am0101095 from the Japan Agency
for Medical Research and Development (AMED); KAKENHI 17K08264, 21H04791 and
21H051130 from by the Japan Society for the Promotion of Science (JSPS); JST Moonshot
Research and Development Program JPMJMS2023 from Japan Science and Technology Agency
(JST); Daiichi Sankyo Foundation of Life Science; Takeda Science Foundation; Ono Medical
Research Foundation; The Uehara Memorial Foundation.

Author Contributions:

627 Conceptualization, J.J. and A.I.; Methodology, J.J., B.B-R., A.I.; Software, J.J., F.M.H.; Formal
628 Analysis, J.J., F.M.H.; Investigation, J.J., R.K., B.B-R., D.H.S., M.M., A.I.; Resources, M.M., K.K.,
629 Data Curation, J.J., F.M.H., A.I.; Writing – Original Draft, J.J.; Writing – Review & Editing, J.J.,
630 R.K., B.B-R., D.H.S., F.M.H., K.K., M.vZ., K.C.G., A.I., B.K.K.; Visualization, J.J., F.M.;
631 Supervision, K.C.G., M.vZ., A.I., B.K.K.; Funding Acquisition, K.C.G., M.vZ., A.I., B.K.K.

Declaration of Interests:

634
635 B.K.K is a cofounder and consultant for ConformetRx, Inc.
636

Figure titles and legends, tables with titles and legends:

638
Figure 1. Arrestin phosphoinositide binding is required for recruitment to some GPCRs A) cartoon
depicting NanoBiT assay for measuring arrestin plasma membrane recruitment upon agonist
stimulation. Upon complementation SmBiT and LgBiT form a functional NanoLuc luciferase. In
key, "Phosphate" denotes phosphorylated Ser/Thr residues and "X" denotes KRK to 3Q mutant
of β-arrestin. B) Two representative GPCRs, β1AR and NTSR1 illustrate data obtained for β-
arrestin recruitment by NanoBiT assay shown in panel A. Data were collected over time after
agonist addition (t=0 min), and values are shown as luminescence fold-change (over vehicle
treatment) ± standard deviation (measured as 2 technical replicates for each of *n*=3 independent
experiments). Colors denote concentrations of agonist used for stimulation. Agonists used were
isoproterenol for β1AR and neurotensin for NTSR1. Grey boxes mark the time region (10-15

minutes post agonist addition) over which luminescence is integrated, for each concentration of
agonist, to produce concentration response curves (bottom). WT and 3Q amplitudes were
determined as the difference of fitted pre- and post-transition plateaus. C) Plot of LOF values for
panel of tested GPCRs. Points represent LOF value obtained as ratio of WT and 3Q recruitment,
and error bars reflect error in LOF derived from standard errors of fits (see methods). Dashed
ellipses denote clusters obtained from k means clustering of data (see methods); AT1R is in
cluster 2 for both βarr1/2, while β2AR-V2C is split with βarr1 (cluster 2) and βarr2 (cluster 1).
Vertical grey lines denote LOF = 0 and LOF =1; vertical purple and orange lines reflect the centers
of the respective clusters from k means and correspond to LOF = 0.06 and LOF = 0.73,
respectively.
659
Figure 2. Receptor phosphorylation patterns govern PIP-dependence for arrestin recruitment. A)
Left, schematic of human NTSR1 showing motifs in receptor ICL3 and C-terminus that are subject
to phosphorylation. Phosphorylation sites examined in this study are shown in red and numbered
1-10 (above). Residue numbers corresponding to the region of human NTSR1 are listed at the
start and end of the shown sequences. Construct key shows possible phosphosites as empty
boxes, which when mutated to alanine are filled with an "X". Plasma membrane recruitment of
arrestin upon stimulation of cells expressing different NTSR1 constructs, measured using the
NanoBiT assay described in Figure 1. Right, points represent LOF value obtained as ratio of WT
and 3Q recruitment, and error bars represent standard error of fits (see methods). Points are
colored based on cluster designation obtained from k means clustering of all receptor-arrestin
recruitment data. B) Translocation of βarr1 to endosomes upon stimulation of cells expressing
different NTSR1 constructs, measured using an endosome bystander NanoBiT assay, as
described in Figure S1. Points represent recruitment (fold chance over basal upon stimulation)
for WT and 3Q recruitment, denoted by circles and triangles, respectively. Points are based on
data from n=3 biological experiments. Error bars represent standard error of fit used to determine
recruitment. Points are colored based on the cluster assignment of that mutant. C) Internalization,
measured by loss of cell-surface receptors upon agonist stimulation, for Δβarr1/2 cells expressing
NTSR1 or β2AR constructs and transfected with arrestin constructs indicated. Values represent
independent experiments (n = 5-10). Internalization by 3Q βarr1 and mock were compared to WT
using a two-tailed paired t-test. ns: $p > 0.05$; *: $p \leq 0.05$; **: $p \leq 0.01$; ***: $p \leq 0.001$; ****: $p \leq$
0.0001.
681
Figure 3. Lipid binding stabilizes core-engaged arrestin complexes. A) cartoon of complexing
efficiency assay. Size-exclusion chromatography (SEC) resolves complex from components. B)
Representative experiment showing SEC chromatograms with vertical dashed lines indicating
free NTSR1, complex, and free arrestins. C) Complexing efficiency, for NTSR1 with indicated
arrestins. Boxplots: center line, median; box range, 25–75th percentiles; whiskers denote
minimum–maximum values. Individual points are shown (n=6 independent experiments). Two-
tailed unpaired t-test used to compare conditions. ns: $p > 0.05$; ****: $p \leq 0.0001$. D) Cartoon
showing equilibrium of NTSR1-arrestin complex. Pink star denotes L68bim probe used for
experiment shown in panel E. E) Bimane spectra for L68bim labeled βarr1 in complex with
NTSR1. All NTSR1 samples contained diC8-PI(4,5)P2 (4.1 μM) Boxplots: center line, median;
box range, 25–75th percentiles; whiskers denote minimum–maximum values. Individual points
are shown (n=3 independent experiments). V2Rpp-NTSR1 (GRK5p) and V2Rpp-NTSR1
(unphos) + V2Rpp were compared by two-tailed unpaired t-test. ns: $p > 0.05$; *: $p \leq 0.05$. Apo
indicates free arrestin; unphos indicates unphosphorylated receptor and GRK5p indicates GRK5-
in vitro phosphorylated receptor. Spectra are normalized to apo (100%) within each experiment
and the fluorescence intensity at lambda max was used as the value. F) Free energy diagram
illustrating how PIP-binding, by stabilizing the core-engaged state of the NTSR1-arrestin complex

slows arrestin dissociation. Loss of the PIP-binding element of arrestin destabilizes the core-
engaged state, shifting equilibrium towards the tail-engaged state leading to a higher degree of
complex disassembly. Removal of the arrestin C-terminus stabilizes the complex in the tail-
engaged state and reduces disassembly even when core-engaged complex is destabilized by
lack of PIP-binding.
704
Figure 4. PIP_2 alone promotes conformational changes in arrestin, including C-terminus
movement, but not release. A) overlay of inactive (PDB: 1G4M) [grey] and active (PDB: 4JQI)
[black] βarr1. The N and C lobes of βarr1 are indicated. Activation leads to reorganization of
several loops, and the gate loop and finger loop are highlighted. Re-orientation of these loops
from inactive (yellow) to active (green) can be monitored by site-specific fluorescence
spectroscopy. In finger loop inset the sphere denotes $C\alpha$ L68C which is labeled with bim. In gate
loop inset, the sphere denotes Ca L293C which is labeled with NBD. An installed W residue
replacing L167 dynamically quenches 293NBD. B) Spectra of bimane labeled (L68C) βarr1 in
response to V2Rpp and PIP_2. Arrow indicates direction of spectral shift with increasing
concentration. Values are mean ± SD (n=3 independent experiments). Spectra were normalized
to the apo condition within a given experiment. C) Spectra of NBD labeled (L167W-L293C) βarr1
in response to V2Rpp and PIP_2. Arrow indicates direction of spectral shift with increasing
concentration. Values are mean ± SD (n=3 independent experiments). Spectra were normalized
to the apo condition within a given experiment. D) Left, cartoon showing how FRET change is
linked to C-terminus release. Right, spectra of AF488/AT647N labeled (A12C-V387C) βarr1 in
response to V2Rpp and PIP_2. Arrow indicates direction of spectral shift with increasing
concentration. Spectra were normalized via donor intensity within a given experiment. Data shown
are for a representative experiment (n=3 independent experiments).
723
Figure 5. PIP_2 enhances Fab30 binding to βarr1. A) Cartoon of surface plasmon resonance (SPR)
experiments, where βarr1 is immobilized via N-terminal biotinylation and a Fab30 binder is
injected in the presence of absence of PIP_2 or V2Rpp. B) Representative sensogram for SPR
binding experiment. With WT βarr1 immobilized, Fab30 (1 μM) was injected alone or together
with V2Rpp (40 μM) or diC8-PI(4,5)P2 (40 μM). The shown sensogram is representative of the
outcome seen for independent experiments (n=3). Dissociation/regeneration phase not shown.
C) Binding of Fab30 to immobilized arrestin constructs in the presence of different additives.
Maximum binding is defined based on normalization of the observed response to the amount of
arrestin immobilized for each construct. Additives: diC8-PG (40 μM), diC8-PI(3)P (40 μM), diC8-
PI(4,5)P2 (40 μM) and V2Rpp (40 μM) were mixed with Fab30 (1 μM) and injected together.
Points reflect independent measurements; open points represent the binding observed for the
additive in the absence of Fab30. Fab30 binding was compared using a two-tailed unpaired t-test.
ns: $p > 0.05$; *: $p \leq 0.05$; **: $p \leq 0.01$; ***: $p \leq 0.001$. D) The proportion of active-like βarr1 increases
in the presence of PIP_2.
738
Figure 6. Model for phosphoinositide regulation of GPCR-β-arrestin complex assembly and
disassembly. GPCRs stratify into two groups with respect to the strength of their interaction with
β-arrestins: one group requires an interaction between β-arrestin and PIP_2 at the plasma
membrane for recruitment (PIP-dependent), while the other does not (PIP-independent). In the
case of PIP-dependent GPCRs, arrestin engagement is unstable and can result in dissociation of
arrestin from the receptor, while maintaining an association with the plasma membrane (left
panel). PIP_2 is enriched at CCSs and in both cases complex assembly can occur. During
endocytosis, PIP_2 is depleted and for PIP-dependent GPCRs, the loss of this PIP_2 contact may
facilitate dissociation of arrestin thereby allowing for receptor recycling. In contrast, a PIP-
independent GPCR will retain the interaction with arrestin even once PIP_2 is depleted owing to

the strong phosphorylation-dependent interactions; however, the full-engaged state of the
complex is less stable in endosomes than at the plasma membrane, thereby allowing further G
protein engagement to occur.
752
753 **Methods**
754
755 **Plasmids**
756
For cell-based assays, we used human, full-length GPCR plasmids cloned into the pCAGGS
vector or the pcDNA3.1 vector derived from a previous study (Inoue et al., 2019). GPCR
constructs were N-terminally FLAG epitope-tagged when they were intended to compare with cell
surface expression levels. Specifically, NTSR1 was fused to the N-terminal FLAG epitope tag with
a linker (M<u>DYKDDDDK</u>GTELGS; the FLAG epitope tag is underlined) and inserted into the
pcDNA3.1 vector. β2AR and μOR were fused to the N-terminal FLAG epitope tag with a preceding
HA-derived   signal   sequence   and   a   flexible   linker
(MKTIIALSYIFCLVFA<u>DYKDDDDK</u>GGSGGGGGSGGSSSGGG) and inserted into the pCAGGS
vector. Unless otherwise noted, other GPCR constructs were untagged. For the bystander
NanoBiT-based β-arrestin assays, human full-length β-arrestin (β-arrestin1 or 2; WT or 3Q) was
N-terminally SmBiT-fused with the flexible linker (M<u>VTGYRLFEEIL</u>GGSGGGGGSGGSSSGG; the
SmBiT is underlined) and inserted into the pCAGGS vector (SmBiT-β-arrestin) (Baidya et al.,
2020a). For the plasma membrane-localizing tag, LgBiT was C-terminally fused to the CAAX motif
derived from human KRAS (SSSGGGKKKKKKSKTKCVIM) through the same flexible linker
(LgBiT-CAAX). For the endosome-localizing tag, LgBiT was N-terminally fused with the human
Endofin FYVE domain (amino-acid regions Gln739-Lys806) again through the same flexible linker
(Endo-LgBiT). For the direct NanoBiT-based β-arrestin assay, human full-length β-arrestin was
N-terminally LgBiT-fused with the same flexible linker and inserted into the pCAGGS vector
(LgBiT-β-arrestin). GPCRs   were   C-terminally   SmBiT-fused   with   the   flexible   linker
(GGSGGGGSGGSSSGG<u>VTGYRLFEEIL</u>; the SmBiT is underlined) and inserted into the
pCAGGS vector (GPCR-SmBiT).
778
779 **Peptides**
780
781 The V2Rpp peptide (ARGRpTPPpSLGPQDEpSCpTpTApSpSpSLAKDTSS) was obtained by
782 custom peptide synthesis (Tufts University Core Facility). Fab30 was expressed and purified as
783 previously described (Shukla et al., 2013). The concentration of V2Rpp stocks were determined
784 by reaction with Ellman's reagent as previously described (Latorraca et al., 2020).
785
786 **NanoBiT-β-arrestin recruitment assays**
787
β-arrestin recruitment to the plasma membrane was measured by the bystander NanoBiT-β-
arrestin assays using the SmBiT-β-arrestin and the LgBiT-CAAX constructs. HEK293A cells
(Thermo Fisher Scientific) were seeded in a 6-cm culture dish (Greiner Bio-One) at a
concentration of 2×10^5 cells per ml (4 ml per dish hereafter) in DMEM (Nissui Pharmaceutical)
supplemented with 10% FBS (Gibco), glutamine, penicillin, and streptomycin, one day before
transfection. The transfection solution was prepared by combining 5 μl of polyethylenimine
solution (1 mg/ml) and a plasmid mixture consisting of 100 ng SmBiT-β-arrestin, 500 ng LgBiT-
CAAX and 200 ng of a test GPCR construct in 200 μl of Opti-MEM (Thermo Fisher Scientific). For
the NTSR1 titration experiment, diluted volume of the FLAG-NTSR1 plasmid (13 ng to 200 ng)
was transfected with 100 ng SmBiT-β-arrestin and 500 ng LgBiT-CAAX with a balanced volume
of the pcDNA3.1 vector (total plasmid volume of 800 ng). After an incubation for one day, the

transfected cells were harvested with 0.5 mM EDTA-containing Dulbecco's PBS, centrifuged, and
suspended in 2 ml of Hank's balanced saline solution (HBSS) containing 0.01% bovine serum
albumin (BSA fatty acid–free grade, SERVA) and 5 mM HEPES (pH 7.4) (assay buffer). The cell
suspension was dispensed in a white 96-well plate (Greiner Bio-One) at a volume of 80 µl per
well and loaded with 20 µl of 50 µM coelenterazine (Carbosynth), diluted in the assay buffer. After
2 h incubation at room temperature, the plate was measured for its baseline luminescence
(SpectraMax L, 2PMT model, Molecular Devices). Thereafter, 20 µl of 6x ligand serially diluted in
the assay buffer were manually added. The ligand used was dependent on the GPCR expressed,
as described in Supplementary Data Table 1. The plate was immediately read for the second
measurement as a kinetics mode and luminescence counts recorded for 15 min with an
accumulation time of 0.18 sec per read and an interval of 20 sec per round. β-arrestin endosomal
translocation was measured by following the same procedure as described above but using the
SmBiT-β-arrestin and the Endo-LgBiT constructs. Similarly, direct recruitment was measured by
the same protocol as described above but using LgBiT-β-arrestin (500 ng) and C-terminally fused-
SmBiT GPCR (500 ng) constructs. For every well, the recorded kinetics data were first normalized
to the baseline luminescence counts.

Analysis of cell-based recruitment data

NanoBiT data were analyzed by converting kinetic data into concentration-response data by
determining an average fold-change (relative to signal pre-stimulation) from 10-15 minutes post-
agonist addition. At least three independent experiments were performed for each receptor-
sensor combination. Concentration-dependent data from two technical replicates for each
independent experiment were collectively fit to a four-parameter log logistic function (LL2.4)
provided in the drc package (v 3.0-1) of the statistical environment R. This equation, of the form:
$f(x) = c + \frac{d-c}{1+e^{(b(\log(x)-\log(e)))}}$ provides pre- and post-transition values, c and d, respectively, that
define the amplitude response for that assay. Cutoffs for bystander NanoBiT experiments were
determined as based on a limit of detection of 3s over the response of mock-transfected cells.
Amplitude values were defined as amplitude = top – bottom of fit, and amplitude error was
calculated as $\delta(amplitude) = \sqrt{(\delta top)^2 + (\delta bottom)^2}$. Converting amplitude to LOF for each
assay was based on the formula: $1 - amplitude(3Q)/amplitude(WT)$. Errors for LOF were
calculated as: $\delta(LOF) = LOF\sqrt{\left(\frac{\delta amplitude(3Q)}{amplitude(3Q)}\right)^2 + \left(\frac{\delta amplitude(WT)}{amplitude(WT)}\right)^2}$. In cases where a fit failed to
converge due to weak recruitment, these amplitudes and errors were set to zero. Recruitment of
βarr1 (3Q) to D1R in both plasma membrane bystander (CAAX) and direct recruitment which was
set to zero. The error amplitude for βarr1 (3Q) endosome translocation assay with D1R was also
set to zero. The error amplitude for βarr1 (3Q) endosome translocation assay with S1PR1 was
set to zero, and the "top" value of the fit was set to 1.2 based on manual inspection. K-means
clustering was performed using pre-built functions in the tidyverse package (v 1.3.1) of R. The
number of clusters was varied from 1 to 10 and an elbow plot of within cluster sum of squares vs
k suggested 2 clusters fit the data well.

For recruitment kinetics, luminescence fold-change was plotted against time, and the values from
zero to five minutes (initial rate) were fit to a logistic function of the form: $f(x) = \frac{L}{1+e^{-k(x-x_0)}}$, where
L is the curve's maximum value, x_0 is the value of the sigmoid midpoint and k is the logistic growth
rate. Fitting was done using the self-starting SSlogis four parameter nls function in the tidyverse
package (v 1.3.1) of R.

GPCR internalization assay

847
GPCR internalization assays was performed as described previously with minor modifications
(Grundmann et al., 2018). Δβarr1/2 double knockout (DKO) cells, previously described (O'Hayre
et al., 2017), were seeded in 6-cm dishes at concentration of 2 x 10^5 cells/ml (4 mL per dish) and
cultured for 1 day before transfection. The cells were transfected with 1 μg of the N-terminally
FLAG-tagged NTSR1 or the β2AR construct, along with 200 ng of the WT or 3Q βarr1 or empty
plasmid, using PEI transfection reagent as described above. After 1-day culture, the transfected
cells were harvested by EDTA-PBS and HEPES-HBSS and, following centrifugation, the cells
were suspended in 500 μL of 0.01% BSA-containing HEPES-HBSS. The cell suspension was
dispensed in a 96-well V-bottom plate (100 μL per well) and mixed with 100 μL of 2× GPCR
solution ligand (2 μM neurotensin for FLAG-NTSR1 or 20 μM Isoproterenol (Sigma-Aldrich) for
FLAG-β2AR). After 30-min incubation in a CO_2 incubator, the plate was centrifuged at 1,500 g for
5 min and the cells were washed twice with D-PBS. The cell pellets were suspended in 2% goat
serum- and 2 mM EDTA-containing D-PBS (blocking buffer; 100 μL per well) and incubated for
30 min on ice. After centrifugation, the cells were stained with anti-FLAG-epitope tag monoclonal
antibody (Clone 1E6, FujiFilm Wako Pure Chemicals; 10 μg mL-1 in the blocking buffer; 25 μL
per well) for 30 min on ice. After washing with D-PBS, the cells were labeled with a goat anti-
mouse IgG secondary antibody conjugated with Alexa Fluor 647 (Thermo Fisher Scientific; 10 μg
mL^{-1} dilution in the blocking buffer; 25 μL per well) for 15 min on ice. The cells were washed once
with D-PBS, resuspended in 100 μL of 2 mM EDTA-containing-D-PBS and filtered through a 40
μm filter. The fluorescently labeled cells (approximately 20,000 cells per sample) were analyzed
by the EC800 flow cytometer (Sony). Fluorescent signal derived from Alexa Fluor 647 was
recorded in the FL3 channel. Mean fluorescence intensity (MFI) from all of the recorded events
was analyzed by a FlowJo software (FlowJo) and used for statistical analysis.
871
872 **Cell-surface expression analysis by flow cytometry**
873
HEK293A cells were seeded in a 6-well culture plate at concentration of 2 x 10^5 cells/ml (2 mL
per dish) and cultured for 1 day before transfection. The cells were transfected with 1 μg of N-
terminally FLAG-tagged GPCR construct using PEI transfection reagent as described above and
cultured for 1 day. The cells were collected by adding 200 μl of 0.53 mM EDTA-containing
Dulbecco's PBS (D-PBS), followed by 200 μl of 5 mM HEPES (pH 7.4)-containing Hank's
Balanced Salt Solution (HBSS). The cell suspension was transferred to a 96-well V-bottom plate
in duplicate and fluorescently labeled with the anti-FLAG epitope tag antibody and a goat anti-
mouse IgG secondary antibody conjugated with Alexa Fluor 488 (Thermo Fisher Scientific, 10 μg
per ml diluted in the blocking buffer) as described above. Live cells were gated with a forward
scatter (FS-Peak-Lin) cutoff at the 390 setting, with a gain value of 1.7 and fluorescent signal
derived from Alexa Fluor 488 was recorded in the FL1 channel. For each experiment, the MFI
value of mutants was normalized to that of WT performed in parallel.
886
887 **cAMP desensitization**
888
HEK293 Δβarr1/2 (DKO) cells that endogenously express β2AR were seeded into 6-well plates and
transiently transfected after 24 hours with mApple, βarr2-mApple, or βarr2(3Q)-mApple. Twenty-four
hours after transfection, cells were transduced with CMV cADDis Green Upward cAMP sensor
according to manufacturer instructions without addition of sodium butyrate (Montana Molecular
#U0200G) and seeded in triplicate in a black clear-bottom 96-well plate (Corning cat# 3340). Twenty-
four hours after transduction, the cells were washed once with 37°C assay buffer [135 mM NaCl, 5
mM KCl, 0.4 mM Mg2Cl, 1.8 mM CaCl2, 5 mM glucose, 20 mM HEPES pH 7.4], loaded into the pre-
warmed 37°C plate reader (Biotek Synergy H4), and equilibrated for five minutes. Prior to beginning

the kinetic assay, mApple was read using monochromoters set to Ex:568/9.0 and Em:592/13.5. Then
cADDis was read using monochromoters set to Ex:500/9.0 and Em:530/20.0. Three cADDis
timepoints were collected to establish baseline, the plate was ejected, isoproterenol in 37°C assay
buffer was added to a final concentration of 100 nM, and the plate was returned to continue collection.
Thirty minutes after isoproterenol addition, 3-isobutyl-1-methylxanthine (IBMX) and forskolin (Fsk) in
37°C assay buffer were added to a final concentrations of 300 µM and 10 µM respectively. Responses
were averaged across technical replicates, normalized to the maximum Fsk/IBMX response, and then
averaged across independent experiments. Expression levels for cADDis and βarr2 were
normalized based on fluorescence.

Western blotting

HEK293A cells were transfected with the SmBiT-β-arrestin and the LgBiT-CAAX constructs by
following the procedure described in the NanoBiT-based β-arrestin assay. After 1-day culture, the
transfected cells were lysed by SDS-PAGE sample buffer (62.5 mM Tris-HCl (pH 6.8), 50 mM
dithiothreitol, 2% SDS, 10% glycerol and 4 M urea) containing 1 mM EDTA and 1 mM
phenylmethylsulfonyl fluoride. Lysates derived from an equal number of cells were separated by
8% SDS-polyacrylamide gel electrophoresis. Subsequently, the proteins were transferred to
PVDF membrane. The blotted membrane was blocked with 5% skim milk-containing blotting
buffer (10 mM Tris-HCl (pH 7.4), 190 mM NaCl and 0.05% Tween 20), immunoblot with primary
(1 µg per mL, unless otherwise indicated) and secondary antibodies conjugated with horseradish
peroxidase (1:2000 dilution). Primary antibodies used in this study were: anti-β-arrestin1 (rabbit
monoclonal; CST, #12697, D8O3J), anti-β-arrestin2 antibody (rabbit monoclonal; CST, #3857,
C16D9) and anti-α-tubulin antibody (mouse monoclonal, clone DM1A; Santa Cruz
Biotechnologies, sc-32293; 1:2000 dilution). Secondary antibodies were anti-rabbit IgG (GE
Healthcare, NA9340) and anti-mouse IgG (GE Healthcare, NA9310). Membrane was soaked with
an ImmunoStar Zeta reagent (FujiFilm Wako Pure Chemical). Chemiluminescence image of the
membrane was acquired, and band intensity was analyzed with Amersham Imager 680 (Cytiva).

NTSR1 expression and purification

Full length human NTSR1 was modified with an N-terminal Flag tag followed by an octa-histidine
tag and cloned into pFastBac1 vector. NTSR1 was expressed in Sf9 insect cells (Expression
Systems) using a FastBac-derived baculovirus. Cells were infected at a density of $4x10^6$ cells/mL
and harvested 60 hrs post infection. Cells were lysed in hypotonic buffer (10 mM HEPES, pH 7.4,
and protease inhibitors) and solubilized at 4 °C for 2 hours in a buffer containing 1% lauryl maltose
neopentyl glycol (LMNG, Anatrace), 0.1% cholesteryl hemisuccinate tris salt (CHS, Steraloids),
0.3% sodium cholate (Sigma), 20 mM HEPES 7.4, 500 mM NaCl, 25% glycerol, iodoacetamide
(to cap cysteine residues) and protease inhibitors. Insoluble debris was removed by centrifugation
and the supernatant was incubated with Ni-NTA (Qiagen) resin for 1 hour at 4 °C. The resin was
washed in batch with buffer containing 0.01% LMNG, 0.001% CHS, 0.003% sodium cholate, 20
mM HEPES pH 7.4, 500 mM NaCl, 10 mM imidazole and eluted with the same buffer
supplemented with 200 mM imidazole, 2 mM $CaCl_2$ and 10 µM NTS_{8-13} (Acetate salt, Sigma). The
eluate was loaded onto M1 FLAG immunoaffinity resin and washed with buffer containing 0.01%
LMNG, 0.001% CHS, 0.003% sodium cholate, 20 mM HEPES pH 7.4, 500 mM NaCl, 10 mM
imidazole, 0.1 µM NTS_{8-13} and 2 mM $CaCl_2$. The receptor was eluted with buffer containing 100
mM NaCl, 20 mM HEPES pH 7.4, 0.005% LMNG, 0.005% CHS, 1 µM NTS_{8-13}, 0.2 mg/mL flag
peptide (DYKDDDDK) and 5 mM EDTA. Elution fractions containing receptor were pooled and
subjected to polishing by SEC on a Superdex 200 Increase 10/300 GL column (GE Healthcare)
in 20 mM HEPES, pH 7.4, 100 mM NaCl, 0.0025% LMNG, 0.00025% CHS, and 0.1 µM NTS_{8-13}.

Peak fractions were pooled and concentrated to 200 μM and aliquots were flash-frozen and stored
at -80 °C until use.
949
GRK5 expression and purification
951
Full length human GRK5 was modified with a C-terminal hexa-histidine tag and cloned into
pVL1392 vector for baculovirus production. GRK5 was expressed and purified as previously
published (Beyett et al., 2019). Briefly, Sf9 insect cells (Expression Systems) were infected with
a BestBac-derived baculovirus at a density of 3.5 x 10^6 cells/mL and harvested 48 hours post
infection. Cells were resuspended, lysed by sonication and the supernatant was applied to Ni-
NTA resin. The resin was washed with lysis buffer and GRK5 eluted with lysis buffer
supplemented with 200 mM imidazole. The combined eluate was then subjected to cation-
exchange chromatography using a MonoS 10/100 column (GE healthcare) and eluted with a
linear gradient of NaCl. Fractions containing GRK5 were combined and run on a Superdex 200
10/300 GL column (GE healthcare). GRK5 was aliquoted, flash frozen, and stored at -80 ˚C until
use.
963
Arrestin expression and purification
965
The parent construct for β-arrestin 1 (βarr1) is the long splice variant of human, cysteine-free
(C59V, C125S, C140L, C150V, C242V, C251V, C269S) β-arrestin 1. This construct is modified
with an N-terminal 6x Histidine tag, followed by a 3C protease site, a GG linker, AviTag and
GGSGGS linker. The sequence was codon-optimized for expression in *E. coli* and cloned into a
pET-15b vector. Point mutations were prepared using site-directed mutagenesis. β-arrestin 1 (1-
382) was prepared by truncating β-arrestin 1 at residue 382. All arrestin constructs used were
prepared as follows: NiCo21(DE3) competent *E. coli* (NEB) were transformed, and large-scale
cultures were grown in TB + ampicillin at 37°C until an OD_{600} of 1.0. Cells were then transferred
to room temperature and induced with 25 μM IPTG when the OD_{600} reached 2.0. Cells were
harvested 20 h post induction and resuspended in lysis buffer [50 mM Hepes pH 7.4, 500 mM
NaCl, 15% glycerol, 7.13 mM β-mercaptoethanol (BME)] to a final volume of 40 mL/L of cells.
Cells were lysed by sonication and the clarified lysate applied to nickel sepharose and batch
incubated for 1.5h at 4 °C. The resin was washed with 10 column volumes of wash buffer (20 mM
HEPES pH 7.4, 500 mM NaCl, 10% glycerol, 7.13 mM BME) + 20 mM imidazole, followed by 10
column volumes of wash buffer + 40 mM imidazole. The protein was then eluted with 5 column
volumes of wash buffer + 200mM imidazole and dialyzed overnight in 100x volume of dialysis
buffer (20 mM Hepes 7.4, 200 mM NaCl, 2 mM BME, 10% glycerol) in the presence of 1:10 (w:w)
of 3C protease. The digested protein was then subjected to reverse-Nickel purification and diluted
with dialysis buffer containing no NaCl to bring the NaCl concentration to 75mM. The protein was
then purified by ion exchange chromatography (mono Q 10/100 GL, GE Healthcare), followed by
SEC using a Superdex 200 increase 10/300 GL column (GE Healthcare) with SEC buffer (20 mM
HEPES pH 7.4, 300 mM NaCl, 10% glycerol). Purified protein was concentrated to between 100-
300 μM using a 30 kDa spin concentrator and aliquots were flash-frozen in liquid nitrogen and
stored at -80 °C until use.
990
Arrestin labeling and biotinylation
992
Following SEC, elution peak fractions were pooled to a concentration of 10-20 μM and labeled
with fluorophore(s): monobromobimane (mBBr), Thermo Fisher Scientific M1378; N,N'-Dimethyl-
N-(Iodoacetyl)-N'-(7-Nitrobenz-2-Oxa-1,3-Diazol-4-yl)Ethylenediamine (IANBD amide), Thermo
Fisher Scientific D2004; or a 1:3 mixture of Alexa Fluor 488 C5 Maleimide, Thermo Fisher

Scientific A10254, and Atto647N Maleimide, ATTO TEC AD647N-41, respectively. Fluorophores
were dissolved to in DMSO and added at 10x molar excess over protein, then allowed to react for
1 h at room temperature prior to quenching with cysteine (10x molar excess over fluorophore).
The labeling reaction was further incubated for 10 minutes after cysteine addition, after which
samples were spin filtered and subjected to a second round of size-exclusion chromatography,
as detailed above, to remove free dye. The purified, was concentrated to between 100-300 μM
using a 30 kDa spin concentrator and aliquots were flash-frozen in liquid nitrogen and stored at -
80 °C until use.
Arrestins (SEC-pure) were biotinylated using recombinant BirA enzyme, according to commercial
protocols (Avidity), with exception that biotinylation was carried out for 12 h at 4 °C, rather than
30 °C. After biotinylation was complete, the reaction was flowed over 100 μL (packed) of nickel
Sepharose, equilibrated in arrestin SEC buffer and supplemented with 10 mM imidazole, then
washed with 200 μL of the equilibration buffer. The combined flow-through and wash fractions
were then purified by size-exclusion as described above.

NTSR1 phosphorylation

NTSR1 (2.5 μM) was equilibrated in phosphorylation buffer (20 mM bis-tris propane (BTP) pH
7.5, 35 mM NaCl, 5 mM MgCl2, 20 μM NTS$_{8-13}$, 20 μM C8-PI(4,5)P2, 0.05 mM TCEP, 0.002%
MNG, 0.0002% CHS) at 25 °C with gentle mixing for 1 h. GRK5 was added to the reaction to a
final concentration of 200 nM, and briefly incubated while the reaction was warmed from 25 °C to
30 °C. ATP was added to a final concentration of 1 mM. Upon completion, the reaction was
supplemented with CaCl$_2$ to a final concentration of 2 mM and applied to an equilibrated M1 FLAG
immunoaffinity resin and washed with buffer containing 0.004% LMNG, 0.004% CHS, 20 mM
HEPES pH 7.4, 100 mM NaCl, 0.2 μM NTS$_{8-13}$, 2 mM CaCl$_2$. The receptor was eluted with buffer
containing 100 mM NaCl, 20 mM HEPES pH 7.4, 0.004% LMNG, 0.004% CHS, 0.2 μM NTS$_{8-13}$,
0.2 mg/mL 1x flag peptide (DYKDDDDK), 5 mM EDTA), followed by SEC using a Superdex 200
increase 10/300 GL column (GE Healthcare) with SEC buffer (20 mM HEPES pH 7.4, 100 mM
NaCl, 0.004% LMNG, 0.0004% CHS).

Analytical fluorescence-detection size-exclusion chromatography

In a final volume of 20 μL, NTSR1 (4.5 μM), the respective arrestin construct (9 μM), NTS$_{8-13}$
peptide (50 μM) and diC8-PI(4,5)P2 (5 μM) were incubated in buffer containing 20 mM HEPES
pH 7.4, 100 mM NaCl, 0.004% LMNG, 0.0004% CHS and 0.2 μM NTS$_{8-13}$. Using a Prominence-i
LC autosampler (Shimadzu), 10 μL was injected onto a ENrich size-exclusion chromatography
650 10 × 300 column (Bio-rad) pre-equilibrated in 20 mM HEPES pH 7.4 100 mM NaCl, 0.004 %
LMNG, 0.004% CHS and 0.2 μM NTS$_{8-13}$, and run at a flow rate of 0.8 ml/min. Tryptophan
fluorescence was monitored at λ(EX) of 280 nm and λ(EM) of 340 nm. Peaks in the obtained size-
exclusion chromatograms were modeled as gaussians, deconvolved and quantified (AUC) using
Magic Plot 3 (Magic Plot).

Surface plasmon resonance measurements

SPR experiments were performed using a GE Biacore T100 instrument. Approximately 300-400
resonance units (RU) of FPLC-purified biotinylated arrestin in HBS-P+ Buffer (GE Healthcare)
were captured on an SA-chip (GE Healthcare), including a reference channel for online
background subtraction of bulk solution refractive index and for evaluation of non-specific binding
of analyte to the chip surface (Biacore T100 Control Software; GE Healthcare). All measurements
were performed with 2-fold serial dilutions using 60 s association followed by a dissociation time

of more than 240 s at 25 °C with a flow rate of at 30 µl min^{-1}. Measurement of titrations at
equilibrium were used to determine K_d values using Biacore Analysis Software (v.2.0.4, GE
Healthcare) and fits to a total binding model were performed in GraphPad Prism 9. Regeneration
was performed by 2 injections of 2 M MgCl$_2$ for 10 s at 50 µl min^{-1} flow rate. In all cases
regeneration resulted in a complete return to baseline. Single cycle measurements were
performed as described above. All single cycle measurements were performed as triplicates and
quantifications calculated to the RU_{max} of the individual immobilized ligands (arrestin proteins).
1054
1055 **Fluorescence anisotropy measurements**
1056
BODIPY-TMR phosphatidylinositol 4,5-bisphosphate (Echelon Biosciences) was dissolved to a
stock concentration of 1 mM in 50 mM Hepes pH 7.4 and used at a final concentration of 4 nM in
the assay. For the arrestin measurements, a two-fold dilution series of was made from a stock of
βarr1 (1-382), yielding fourteen samples with final concentrations ranging from 150 µM to 0.02
µM. A control sample containing buffer only was included to measure the free anisotropy of
BODIPY-PIP$_2$. After mixing the BODIPY-PIP$_2$ with arrestin or buffer, samples were incubated for
1h at room temperature prior to measurements. Samples were measured in five 20 µL replicates
in a 384-well plate on a Tecan Infinite M1000 (Tecan Life Sciences), using an excitation
wavelength of 530 nm, an emission wavelength of 573 nm and bandwidths of 5 nm. The obtained
data was fit using to a one-site total binding model $Y = Bottom + (top - bottom)/1 +$
$10^{HS*log(EC50-X)}$ where HS denotes the hill-slope.
1068
1069 **Bulk fluorescence measurements**
1070
Bulk fluorescence measurements were performed on either a Fluorolog instrument (Horiba) using
FluorEssence v3.8 software and operating in photon-counting mode, or a Tecan Infinite M1000
PRO multimodal microplate reader (Tecan). Fluorolog measurements of bimane-labeled βarr1
constructs (NTSR1 experiments) were performed at final concentration of 0.4 µM [arrestin] in
buffer containing 20 mM HEPES pH 7.4, 100 mM NaCl and 0.004% LMNG (w/v)/0.0004% CHS
(w/v) supplemented with 4 µM NTS(8-13). For NTSR1 experiments the following concentrations
were used: 4 µM NTSR1, 4.1 µM diC8-PI(4,5)P2, 50 µM V2Rpp (depending on condition).
Samples were incubated for 1 h in the dark before measurement. Fluorescence data were
collected in a quartz cuvette with 135 mL of sample. Bimane fluorescence was measured by
excitation at 370 nm with excitation and emission bandwidth passes of 3 nm, and emission spectra
were recorded from 400 to 550 nm in 2 nm increments with 0.1 s integration time. Care was taken
to extensively rinse and argon-dry the cuvette between individual measurements. To remove
background fluorescence, buffer spectra were collected using the same settings, and subtracted
from each sample spectrum.
FRET measurements of AF488-AT647N-labeled βarr1 constructs were performed as described
for bimane measurements, with the following differences: samples were excited at 476 nm with 3
nm excitation and 4 nm emission slit widths. Spectra were collected from 485 nm to 750 nm in 1
nm increments with 0.1 s integration time. FRET measurements in the absence of NTSR1 were
performed in buffer containing 20 mM HEPES pH 7.4, 100 mM NaCl and 0.004% LMNG
(w/v)/0.0004% CHS (no NTS). FRET measurements with NTSR1 were done with 0.5 µM NTSR1
and 0.5 µM diC8-PI(4,5)P2.
NBD spectra measured on the Tecan Infinite M1000 PRO were collected using 384- or 96-well
(1/2 area) flat black Greiner plates with 50 or 100 µL of sample, respectively, at a final
concentration of 0.5 µM βarr1 in buffer containing 20 mM HEPES pH 7.4, 100 mM NaCl and
0.004% LMNG (w/v)/0.0004% CHS. For NBD the following instrument settings were used:
excitation: 490 nm, emission 510-580 nm (1 nm steps) with 20 s read time and 400 Hz flash mode.

Gain and z-position were optimized prior to reading. Bimane spectra were collected in white plates
using the following instrument settings: excitation: 370 nm, emission 420-500 nm (1 nm steps)
with 20 s read time and 400 Hz flash mode.
Efret values for FRET experiments were calculated as $Efret = \frac{A}{(D+A)}$ and normalized to donor
intensity within a given experiment. Scaled FRET values (apo = 100, min(FRET) = 0) were fit to
a single exponential decay function $Y = (Y0 - NS) * e^{-K*x} + NS$ using the nls function in R for
EC$_{50}$ values (obtained as t$_{1/2}$ for decay). NS denotes concentration-dependent non-specific signal.
L167W-293NBD was fit using the same function. L68bim data was fit to a total binding model $Y =$
$Bmax * \frac{X}{(Kd+X)} + NS * X + background$, where background is a constant value. Fitting was
independently performed both in R and with GraphPad Prism 9 for corroboration, values reported
are from Prism 9.
1108
1109 **Supplemental Information titles and legends**
1110
Figure S1. Arrestin phosphoinositide binding is required for plasma membrane recruitment to
some GPCRs. A) cAMP response in HEK293 cells devoid of β-arrestins upon stimulation of
endogenous β2AR with 100 nM isoproterenol (iso). Clone 1 (CL1) and Clone 2 (CL2) are
independent βarr1/2 knock-out cell lines (O'Hare et al. 2017). Data are normalized to response
with Forskolin (Fsk)/3-isobutyl-1-methylxanthine (IBMX) and show mean with 95% confidence
intervals (*n*=3 independent experiments). Two-way analysis of variance (ANOVA), Tukey's
multiple comparison test. For CL2 * denotes p < 0.05 for WT vs. mApple over the interval of 17-
32 minutes, while 3Q vs. mApple was not significant. For CL1 * denotes p < 0.05 for WT vs.
mApple over the interval of 19-29 minutes, while 3Q vs. mApple was not significant. B)
Quantification of expression for βarr1 and βarr2 (both WT and 3Q) NanoBiT constructs, as
determined by western blot (Supplementary data figure 2). Mean values of 3-4 independent
experiments were compared by a two-tailed unpaired t-test, where ns denotes p > 0.05, * P ≤
0.05. Boxplots: center line, median; box range, 25–75th percentiles; whiskers denote minimum–
maximum values. Individual points are shown. C) LOF is only weakly correlated with recruitment
of WT β-arrestins. Data are mean LOF and mean WT βarr1/2 recruitment. βarr1 recruitment is
shown as circles and βarr2 recruitment is shown as triangles. Data are colored based on assigned
cluster. Dashed line shows expected linear relationship and R is the Pearson coefficient, with -
0.51 reflecting a weak negative correlation. D) Plot of LOF data for plasma membrane bystander
(CAAX) vs. LOF for direct recruitment. βarr1 recruitment is shown as circles and βarr2 recruitment
is shown as triangles. Data are colored based on assigned cluster. Dashed line shows expected
linear relationship and R is the Pearson coefficient, with 0.88 reflecting a very strong positive
correlation. E) NanoBiT assay for measuring endosome translocation of βarr1. Cartoon of
endosome bystander assay (left). βarr1 endosome recruitment data (right) with dashed ellipses
to indicate clusters based on CAAX data. β-arrestin endosome recruitment determined by span
of luminescence fold change. Data are mean ± SEM (*n*=3 independent experiments). Dashed line
indicates three times the maximum signal measured in mock (receptor) transfected cells.
1137
Figure S2. Loss of PIP binding slows β-arrestin recruitment to cluster 2 GPCRs. A) initial rate (0-
5 minutes post-agonist stimulation) expressed as luminescence fold-change (FC)/min. Data from
n=3 independent experiments fit independently (see methods). Boxplots: center line, median; box
range, 25–75th percentiles; whiskers denote minimum–maximum values. For each receptor, and
for each βarr1 and βarr2 WT and 3Q were compared by a two-tailed unpaired t-test, where ns
denotes p > 0.05, * P ≤ 0.05, ** P ≤ 0.01, **** P ≤ 0.0001. B) Data from A) expressed as a
difference in rate shows that with the exception of βarr1-TACR1 all cluster 2 receptors show faster

recruitment of WT β-arrestin1/2 than corresponding 3Q mutant. Data are mean ± SEM (n=3
independent experiments).

1147

Figure S3. Arrestin recruitment to NTSR1 mutants can be measured by NanoBiT recruitment
assay. A) Expression of NTSR1 constructs in HEK293A cells used for NanoBiT assays. Boxplots:
center line, median; box range, 25–75th percentiles; whiskers denote minimum–maximum values.
Individual points are shown. Values are mean, relative to NTSR1-WT (n=4 independent
experiments). For each construct, a comparison to NTSR1-WT by a two-tailed unpaired Wilcoxon
test was performed, where ns denotes p > 0.05, * P ≤ 0.05. B) Direct complementation NanoBiT
assay Emax for Sm-βarr1 interaction with Lg-CAAX for cells expressing NTSR1-WT as a function
of mean fluorescence intensity (MFI), as determined by cell-surface staining. Amount of NTSR1-
WT DNA transfected is written; blue arrow denotes 200 ng, the amount used in recruitment assays
in Figure 2. C) As B, except the pEC_{50} of recruitment response upon NTS stimulation is plotted
vs. MFI, instead of Emax. In both B and C, points represent mean values and error bars indicate
95% CI (n=3 independent experiments).

1160

Figure S4. PIP binding stabilizes core-engaged arrestin complexes. A) LOF in complexing
efficiency as determined by SEC. LOF = 1 corresponds to complete loss of complex formation,
while LOF = 0 corresponds to no difference in complexing efficiency between WT and 3Q βarr1
(n=5 independent experiments). Boxplots: center line, median; box range, 25–75th percentiles;
whiskers denote minimum–maximum values. Individual points are shown. compared by a two-
tailed unpaired t-test, where **** P ≤ 0.0001. B) Binding of BODIPY-TMR PI(4,5)P2 to βarr1 (1-
382) protein (WT or 3Q). Points are mean and error bars reflect 95% CI (n=5 independent
experiments). Data were fit to a logistical function as described in methods and best fit values for
B_{max} and K_d are provided with 95% CI in parentheses. C) Structure of transition from inactive
(PDB: 1G4M) to active (PDB: 4JQI) βarr1 involves displacement of the βarr1 C-terminus (dark
grey) by V2Rpp (blue). Two cysteine residues were added to a cys-less βarr1 background at
positions A12 and V387 (pink spheres). These positions were labeled with fluorophores that,
through FRET, allow for monitoring the position of the C-terminus. D) When labeled with a FRET
pair, βarr1-12C/387C shows a high-FRET state in the absence of V2Rpp, and a low-FRET state
when the βarr1 C-terminus is displaced by V2Rpp. FRET measured when βarr1 (WT or 3Q)-
12C/387C-AF488-AT647N is bound to V2Rpp (0.5 μM), NTSR1 (GRK5p, 0.5 μM), or NTSR1
(unphosphorylated, 0.5 μM)+V2Rpp (0.5 μM). All samples containing NTSR1 were supplemented
with diC8-PI(4,5)P2 (0.5 μM). Apo βarr1 (WT or 3Q)-12C/387C-AF488-AT647N was normalized
to 1.0 and βarr1 (WT or 3Q)-12C/387C-AF488-AT647N + V2Rpp (10 μM) was normalized 0.0 for
each experiment (n=3 independent measurements) (right). Boxplots: center line, median; box
range, 25–75th percentiles; whiskers denote minimum–maximum values. Individual points are
shown.

1183

Figure S5. PIP_2 allosterically triggers movement of the arrestin C-tail, but not release. A-B) Finger
loop (L68C-bim) responses. %apo is scaled such that the fluorescence intensity (at λmax) for apo
arrestin is 100% and each condition is scaled as a factor of apo. ND denotes not determined
values. Values for B_{max} (max response) and K_d (based on single-site binding fitting) are provided
and ranges in parentheses correspond to 95% CI. Points are mean and error bars reflect 95% CI
(n=3 independent experiments). C-D) Gate loop (L167W-293C-NBD) responses. %apo is scaled
such that the fluorescence intensity (at λmax) for apo arrestin is 100% and each condition is
scaled as a factor of apo. Values for EC_{50} (half maximal response) and k (rate constant based on
single exponential decay) are provided and ranges in parentheses correspond to 95% CI. Points
are mean and error bars reflect 95% CI (n=3 independent experiments). E-F) C-terminus release
(A12C-V387C-AF488-AT647N) responses. %FRET is scaled such that apo arrestin is 100% and

the highest concentration of V2Rpp (100 uM) is 0%. INF denotes infinite upper bound. ND denotes
not determined values. Range of EC_{50} values is indicated in parentheses and represents 95% CI.
Points represent mean and error bars reflect 95% CI (n=3 independent experiments).
1198
Figure S6. Plasma membrane PIPs promote conformational changes in arrestin. A) Structure of
soluble lipid derivatives examined in this work. B-I) Concentration response curves for L68bim
βarr1. Values for B_{max} (max response) and EC_{50} (based on single-site binding fitting) are provided
and ranges in parentheses correspond to 95% CI. Points are mean and error bars reflect 95% CI
(n=3 independent experiments). ND denotes not determined values. J) Summary of effect size
(B_{max}) for different lipids with L68bim βarr1. Values represent B_{max} obtained from fitting
independent experiments. ND is used for PG and PI(3)P as data could not be fit. Bars represent
mean B_{max} and error bars denote standard deviation across the fits. K) A PI(4,5)P2 derivative
bound in the C-terminal domain of βarr1 (PDB: 6UP7). The side chain of K232 was modeled
based on PDB: 4JQI as it was not ordered in PDB: 6UP7.
1209
Figure S7. Titration of interactions with immobilized βarr1 by SPR. Immobilized βarr1 construct
(ligand) is listed as column heading. Rows represent analyte flowed during titration. K_d and B_{max}
values obtained as described in methods are listed with 95% CI range in parentheses. ND denotes
value not determined. PIP_2 binding was not fit (D-F), whereas B_{max} for Fab30+V2Rpp (M-O) failed
to converge. In all cases the darkest blue curve corresponds to the highest concentration and the
darkest red curve corresponds to the lowest concentration. Titrations, as described in methods,
for V2Rpp and PIP_2 (A-F) ranged from 40 μM to 78.1 nM. Titrations of Fab30 ranged from 2000
nM to 3.9 nM with fixed concentration of V2Rpp (40 μM) or PIP_2 (40 μM), as indicated. Vertical
axes are raw RU, and not corrected for channel loading.
1219

References

Baidya, M., Kumari, P., Dwivedi-Agnihotri, H., Pandey, S., Chaturvedi, M., Stepniewski, T.M., Kawakami, K., Cao, Y., Laporte, S.A., Selent, J.*, et al.* (2020a). Key phosphorylation sites in GPCRs orchestrate the contribution of beta-Arrestin 1 in ERK1/2 activation. EMBO Rep *21*, e49886.

Baidya, M., Kumari, P., Dwivedi-Agnihotri, H., Pandey, S., Sokrat, B., Sposini, S., Chaturvedi, M., Srivastava, A., Roy, D., Hanyaloglu, A.C.*, et al.* (2020b). Genetically encoded intrabody sensors report the interaction and trafficking of beta-arrestin 1 upon activation of G-protein-coupled receptors. J Biol Chem *295*, 10153-10167.

Beyett, T.S., Fraley, A.E., Labudde, E., Patra, D., Coleman, R.C., Eguchi, A., Glukhova, A., Chen, Q., Williams, R.M., Koch, W.J.*, et al.* (2019). Perturbation of the interactions of calmodulin with GRK5 using a natural product chemical probe. Proc Natl Acad Sci U S A *116*, 15895-15900.

Cahill, T.J., 3rd, Thomsen, A.R., Tarrasch, J.T., Plouffe, B., Nguyen, A.H., Yang, F., Huang, L.Y., Kahsai, A.W., Bassoni, D.L., Gavino, B.J.*, et al.* (2017). Distinct conformations of GPCR-beta-arrestin complexes mediate desensitization, signaling, and endocytosis. Proc Natl Acad Sci U S A *114*, 2562-2567.

Chen, Q., Iverson, T.M., and Gurevich, V.V. (2018). Structural Basis of Arrestin-Dependent Signal Transduction. Trends Biochem Sci *43*, 412-423.

Chen, Q., Perry, N.A., Vishnivetskiy, S.A., Berndt, S., Gilbert, N.C., Zhuo, Y., Singh, P.K., Tholen, J., Ohi, M.D., Gurevich, E.V.*, et al.* (2017). Structural basis of arrestin-3 activation and signaling. Nat Commun *8*, 1427.

Chen, Q., Zhuo, Y., Sharma, P., Perez, I., Francis, D.J., Chakravarthy, S., Vishnivetskiy, S.A., Berndt, S., Hanson, S.M., Zhan, X.*, et al.* (2021). An Eight Amino Acid Segment Controls Oligomerization and Preferred Conformation of the two Non-visual Arrestins. J Mol Biol *433*, 166790.

De Matteis, M.A., and Godi, A. (2004). PI-loting membrane traffic. Nat Cell Biol *6*, 487-492.

DeFea, K.A., Zalevsky, J., Thoma, M.S., Dery, O., Mullins, R.D., and Bunnett, N.W. (2000). beta-arrestin-dependent endocytosis of proteinase-activated receptor 2 is required for intracellular targeting of activated ERK1/2. J Cell Biol *148*, 1267-1281.

Dery, O., Thoma, M.S., Wong, H., Grady, E.F., and Bunnett, N.W. (1999). Trafficking of proteinase-activated receptor-2 and beta-arrestin-1 tagged with green fluorescent protein. beta-Arrestin-dependent endocytosis of a proteinase receptor. J Biol Chem *274*, 18524-18535.

Di Paolo, G., and De Camilli, P. (2006). Phosphoinositides in cell regulation and membrane dynamics. Nature *443*, 651-657.

Dixon, A.S., Schwinn, M.K., Hall, M.P., Zimmerman, K., Otto, P., Lubben, T.H., Butler, B.L., Binkowski, B.F., Machleidt, T., Kirkland, T.A.*, et al.* (2016). NanoLuc Complementation Reporter Optimized for Accurate Measurement of Protein Interactions in Cells. ACS Chem Biol *11*, 400-408.

Eichel, K., Jullie, D., Barsi-Rhyne, B., Latorraca, N.R., Masureel, M., Sibarita, J.B., Dror, R.O.,
and von Zastrow, M. (2018). Catalytic activation of beta-arrestin by GPCRs. Nature *557*, 381-386.

Eichel, K., Jullie, D., and von Zastrow, M. (2016). beta-Arrestin drives MAP kinase signalling from
clathrin-coated structures after GPCR dissociation. Nat Cell Biol *18*, 303-310.

Feinstein, T.N., Wehbi, V.L., Ardura, J.A., Wheeler, D.S., Ferrandon, S., Gardella, T.J., and
Vilardaga, J.P. (2011). Retromer terminates the generation of cAMP by internalized PTH
receptors. Nat Chem Biol *7*, 278-284.

Feinstein, T.N., Yui, N., Webber, M.J., Wehbi, V.L., Stevenson, H.P., King, J.D., Jr., Hallows,
K.R., Brown, D., Bouley, R., and Vilardaga, J.P. (2013). Noncanonical control of vasopressin
receptor type 2 signaling by retromer and arrestin. J Biol Chem *288*, 27849-27860.

Ferrandon, S., Feinstein, T.N., Castro, M., Wang, B., Bouley, R., Potts, J.T., Gardella, T.J., and
Vilardaga, J.P. (2009). Sustained cyclic AMP production by parathyroid hormone receptor
endocytosis. Nat Chem Biol *5*, 734-742.

Gaidarov, I., Krupnick, J.G., Falck, J.R., Benovic, J.L., and Keen, J.H. (1999). Arrestin function in
G protein-coupled receptor endocytosis requires phosphoinositide binding. EMBO J *18*, 871-881.

Ghosh, E., Dwivedi, H., Baidya, M., Srivastava, A., Kumari, P., Stepniewski, T., Kim, H.R., Lee,
M.H., van Gastel, J., Chaturvedi, M.*, et al.* (2019). Conformational Sensors and Domain Swapping
Reveal Structural and Functional Differences between beta-Arrestin Isoforms. Cell Rep *28*, 3287-
3299 e3286.

Grundmann, M., Merten, N., Malfacini, D., Inoue, A., Preis, P., Simon, K., Ruttiger, N., Ziegler, N.,
Benkel, T., Schmitt, N.K.*, et al.* (2018). Lack of beta-arrestin signaling in the absence of active G
proteins. Nat Commun *9*, 341.

Hanyaloglu, A.C., and von Zastrow, M. (2008). Regulation of GPCRs by endocytic membrane
trafficking and its potential implications. Annu Rev Pharmacol Toxicol *48*, 537-568.

Hasegawa, J., Strunk, B.S., and Weisman, L.S. (2017). PI5P and PI(3,5)P2: Minor, but Essential
Phosphoinositides. Cell Struct Funct *42*, 49-60.

Huang, W., Masureel, M., Qu, Q., Janetzko, J., Inoue, A., Kato, H.E., Robertson, M.J., Nguyen,
K.C., Glenn, J.S., Skiniotis, G.*, et al.* (2020). Structure of the neurotensin receptor 1 in complex
with beta-arrestin 1. Nature *579*, 303-308.

Innamorati, G., Sadeghi, H., and Birnbaumer, M. (1998a). Transient phosphorylation of the V1a
vasopressin receptor. J Biol Chem *273*, 7155-7161.

Innamorati, G., Sadeghi, H.M., Tran, N.T., and Birnbaumer, M. (1998b). A serine cluster prevents
recycling of the V2 vasopressin receptor. Proc Natl Acad Sci U S A *95*, 2222-2226.

Inoue, A., Raimondi, F., Kadji, F.M.N., Singh, G., Kishi, T., Uwamizu, A., Ono, Y., Shinjo, Y.,
Ishida, S., Arang, N.*, et al.* (2019). Illuminating G-Protein-Coupling Selectivity of GPCRs. Cell
177, 1933-1947 e1925.

Irannejad, R., Tomshine, J.C., Tomshine, J.R., Chevalier, M., Mahoney, J.P., Steyaert, J.,
Rasmussen, S.G., Sunahara, R.K., El-Samad, H., Huang, B.*, et al.* (2013). Conformational
biosensors reveal GPCR signalling from endosomes. Nature *495*, 534-538.

Irannejad, R., Tsvetanova, N.G., Lobingier, B.T., and von Zastrow, M. (2015). Effects of
endocytosis on receptor-mediated signaling. Curr Opin Cell Biol *35*, 137-143.

Just, S., Illing, S., Trester-Zedlitz, M., Lau, E.K., Kotowski, S.J., Miess, E., Mann, A., Doll, C.,
Trinidad, J.C., Burlingame, A.L.*, et al.* (2013). Differentiation of opioid drug effects by hierarchical
multi-site phosphorylation. Mol Pharmacol *83*, 633-639.

Kadlecova, Z., Spielman, S.J., Loerke, D., Mohanakrishnan, A., Reed, D.K., and Schmid, S.L.
(2017). Regulation of clathrin-mediated endocytosis by hierarchical allosteric activation of AP2. J
Cell Biol *216*, 167-179.

Khoury, E., Nikolajev, L., Simaan, M., Namkung, Y., and Laporte, S.A. (2014). Differential
regulation of endosomal GPCR/beta-arrestin complexes and trafficking by MAPK. J Biol Chem
289, 23302-23317.

Kim, M., Vishnivetskiy, S.A., Van Eps, N., Alexander, N.S., Cleghorn, W.M., Zhan, X., Hanson,
S.M., Morizumi, T., Ernst, O.P., Meiler, J.*, et al.* (2012). Conformation of receptor-bound visual
arrestin. Proc Natl Acad Sci U S A *109*, 18407-18412.

Kim, Y.J., Hofmann, K.P., Ernst, O.P., Scheerer, P., Choe, H.W., and Sommer, M.E. (2013).
Crystal structure of pre-activated arrestin p44. Nature *497*, 142-146.

Komolov, K.E., and Benovic, J.L. (2018). G protein-coupled receptor kinases: Past, present and
future. Cell Signal *41*, 17-24.

Krueger, K.M., Daaka, Y., Pitcher, J.A., and Lefkowitz, R.J. (1997). The role of sequestration in
G protein-coupled receptor resensitization. Regulation of beta2-adrenergic receptor
dephosphorylation by vesicular acidification. J Biol Chem *272*, 5-8.

Kumari, P., Srivastava, A., Banerjee, R., Ghosh, E., Gupta, P., Ranjan, R., Chen, X., Gupta, B.,
Gupta, C., Jaiman, D.*, et al.* (2016). Functional competence of a partially engaged GPCR-beta-
arrestin complex. Nat Commun *7*, 13416.

Kumari, P., Srivastava, A., Ghosh, E., Ranjan, R., Dogra, S., Yadav, P.N., and Shukla, A.K.
(2017). Core engagement with beta-arrestin is dispensable for agonist-induced vasopressin
receptor endocytosis and ERK activation. Mol Biol Cell *28*, 1003-1010.

Latorraca, N.R., Masureel, M., Hollingsworth, S.A., Heydenreich, F.M., Suomivuori, C.M., Brinton,
C., Townshend, R.J.L., Bouvier, M., Kobilka, B.K., and Dror, R.O. (2020). How GPCR
Phosphorylation Patterns Orchestrate Arrestin-Mediated Signaling. Cell *183*, 1813-1825 e1818.

Latorraca, N.R., Wang, J.K., Bauer, B., Townshend, R.J.L., Hollingsworth, S.A., Olivieri, J.E., Xu,
H.E., Sommer, M.E., and Dror, R.O. (2018). Molecular mechanism of GPCR-mediated arrestin
activation. Nature *557*, 452-456.

Lee, Y., Warne, T., Nehme, R., Pandey, S., Dwivedi-Agnihotri, H., Chaturvedi, M., Edwards, P.C.,
Garcia-Nafria, J., Leslie, A.G.W., Shukla, A.K.*, et al.* (2020). Molecular basis of beta-arrestin
coupling to formoterol-bound beta1-adrenoceptor. Nature *583*, 862-866.

Lobingier, B.T., and von Zastrow, M. (2019). When trafficking and signaling mix: How subcellular
location shapes G protein-coupled receptor activation of heterotrimeric G proteins. Traffic *20*, 130-
136.

Luttrell, L.M., Wang, J., Plouffe, B., Smith, J.S., Yamani, L., Kaur, S., Jean-Charles, P.Y.,
Gauthier, C., Lee, M.H., Pani, B.*, et al.* (2018). Manifold roles of beta-arrestins in GPCR signaling
elucidated with siRNA and CRISPR/Cas9. Sci Signal *11*.

Martinez-Morales, J.C., Romero-Avila, M.T., Reyes-Cruz, G., and Garcia-Sainz, J.A. (2018).
S1P1 receptor phosphorylation, internalization, and interaction with Rab proteins: effects of
sphingosine 1-phosphate, FTY720-P, phorbol esters, and paroxetine. Biosci Rep *38*.

Milano, S.K., Kim, Y.M., Stefano, F.P., Benovic, J.L., and Brenner, C. (2006). Nonvisual arrestin
oligomerization and cellular localization are regulated by inositol hexakisphosphate binding. J Biol
Chem *281*, 9812-9823.

Nakagawa, T., and Asahi, M. (2013). beta1-adrenergic receptor recycles via a membranous
organelle, recycling endosome, by binding with sorting nexin27. J Membr Biol *246*, 571-579.

Namkung, Y., Le Gouill, C., Lukashova, V., Kobayashi, H., Hogue, M., Khoury, E., Song, M.,
Bouvier, M., and Laporte, S.A. (2016). Monitoring G protein-coupled receptor and beta-arrestin
trafficking in live cells using enhanced bystander BRET. Nat Commun *7*, 12178.

Nguyen, A.H., Thomsen, A.R.B., Cahill, T.J., 3rd, Huang, R., Huang, L.Y., Marcink, T., Clarke,
O.B., Heissel, S., Masoudi, A., Ben-Hail, D.*, et al.* (2019). Structure of an endosomal signaling
GPCR-G protein-beta-arrestin megacomplex. Nat Struct Mol Biol *26*, 1123-1131.

Nuber, S., Zabel, U., Lorenz, K., Nuber, A., Milligan, G., Tobin, A.B., Lohse, M.J., and Hoffmann,
C. (2016). beta-Arrestin biosensors reveal a rapid, receptor-dependent activation/deactivation
cycle. Nature *531*, 661-664.

O'Hayre, M., Eichel, K., Avino, S., Zhao, X., Steffen, D.J., Feng, X., Kawakami, K., Aoki, J.,
Messer, K., Sunahara, R.*, et al.* (2017). Genetic evidence that beta-arrestins are dispensable for
the initiation of beta2-adrenergic receptor signaling to ERK. Sci Signal *10*.

Oakley, R.H., Laporte, S.A., Holt, J.A., Barak, L.S., and Caron, M.G. (1999). Association of beta-
arrestin with G protein-coupled receptors during clathrin-mediated endocytosis dictates the profile
of receptor resensitization. J Biol Chem *274*, 32248-32257.

Oakley, R.H., Laporte, S.A., Holt, J.A., Barak, L.S., and Caron, M.G. (2001). Molecular
determinants underlying the formation of stable intracellular G protein-coupled receptor-beta-
arrestin complexes after receptor endocytosis*. J Biol Chem *276*, 19452-19460.

Oakley, R.H., Laporte, S.A., Holt, J.A., Caron, M.G., and Barak, L.S. (2000). Differential affinities
of visual arrestin, beta arrestin1, and beta arrestin2 for G protein-coupled receptors delineate two
major classes of receptors. J Biol Chem *275*, 17201-17210.

Pakharukova, N., Masoudi, A., Pani, B., Staus, D.P., and Lefkowitz, R.J. (2020). Allosteric
activation of proto-oncogene kinase Src by GPCR-beta-arrestin complexes. J Biol Chem *295*,
16773-16784.

Perkovska, S., Mejean, C., Ayoub, M.A., Li, J., Hemery, F., Corbani, M., Laguette, N., Ventura,
M.A., Orcel, H., Durroux, T.*, et al.* (2018). V1b vasopressin receptor trafficking and signaling: Role
of arrestins, G proteins and Src kinase. Traffic *19*, 58-82.

Rajagopal, S., and Shenoy, S.K. (2018). GPCR desensitization: Acute and prolonged phases.
Cell Signal *41*, 9-16.

Ranjan, R., Dwivedi, H., Baidya, M., Kumar, M., and Shukla, A.K. (2017). Novel Structural Insights
into GPCR-beta-Arrestin Interaction and Signaling. Trends Cell Biol *27*, 851-862.

Reiter, E., Ahn, S., Shukla, A.K., and Lefkowitz, R.J. (2012). Molecular mechanism of beta-
arrestin-biased agonism at seven-transmembrane receptors. Annu Rev Pharmacol Toxicol *52*,
179-197.

Sente, A., Peer, R., Srivastava, A., Baidya, M., Lesk, A.M., Balaji, S., Shukla, A.K., Babu, M.M.,
and Flock, T. (2018). Molecular mechanism of modulating arrestin conformation by GPCR
phosphorylation. Nat Struct Mol Biol *25*, 538-545.

Shukla, A.K., Manglik, A., Kruse, A.C., Xiao, K., Reis, R.I., Tseng, W.C., Staus, D.P., Hilger, D.,
Uysal, S., Huang, L.Y.*, et al.* (2013). Structure of active beta-arrestin-1 bound to a G-protein-
coupled receptor phosphopeptide. Nature *497*, 137-141.

Shukla, A.K., Westfield, G.H., Xiao, K., Reis, R.I., Huang, L.Y., Tripathi-Shukla, P., Qian, J., Li,
S., Blanc, A., Oleskie, A.N.*, et al.* (2014). Visualization of arrestin recruitment by a G-protein-
coupled receptor. Nature *512*, 218-222.

Sommer, M.E., Smith, W.C., and Farrens, D.L. (2005). Dynamics of arrestin-rhodopsin
interactions: arrestin and retinal release are directly linked events. J Biol Chem *280*, 6861-6871.

Sommer, M.E., Smith, W.C., and Farrens, D.L. (2006). Dynamics of arrestin-rhodopsin
interactions: acidic phospholipids enable binding of arrestin to purified rhodopsin in detergent. J
Biol Chem *281*, 9407-9417.

Song, W., Yen, H.Y., Robinson, C.V., and Sansom, M.S.P. (2019). State-dependent Lipid
Interactions with the A2a Receptor Revealed by MD Simulations Using In Vivo-Mimetic
Membranes. Structure *27*, 392-403 e393.

Song, X., Coffa, S., Fu, H., and Gurevich, V.V. (2009). How does arrestin assemble MAPKs into
a signaling complex? J Biol Chem *284*, 685-695.

Staus, D.P., Hu, H., Robertson, M.J., Kleinhenz, A.L.W., Wingler, L.M., Capel, W.D., Latorraca,
N.R., Lefkowitz, R.J., and Skiniotis, G. (2020). Structure of the M2 muscarinic receptor-beta-
arrestin complex in a lipid nanodisc. Nature *579*, 297-302.

Staus, D.P., Wingler, L.M., Choi, M., Pani, B., Manglik, A., Kruse, A.C., and Lefkowitz, R.J. (2018).
Sortase ligation enables homogeneous GPCR phosphorylation to reveal diversity in beta-arrestin
coupling. Proc Natl Acad Sci U S A *115*, 3834-3839.

Sun, Y., Carroll, S., Kaksonen, M., Toshima, J.Y., and Drubin, D.G. (2007). PtdIns(4,5)P2
turnover is required for multiple stages during clathrin- and actin-dependent endocytic
internalization. J Cell Biol *177*, 355-367.

Tewson, P.H., Martinka, S., Shaner, N.C., Hughes, T.E., and Quinn, A.M. (2016). New DAG and
cAMP Sensors Optimized for Live-Cell Assays in Automated Laboratories. J Biomol Screen *21*,
298-305.

Thomsen, A.R.B., Plouffe, B., Cahill, T.J., 3rd, Shukla, A.K., Tarrasch, J.T., Dosey, A.M., Kahsai,
A.W., Strachan, R.T., Pani, B., Mahoney, J.P.*, et al.* (2016). GPCR-G Protein-beta-Arrestin
Super-Complex Mediates Sustained G Protein Signaling. Cell *166*, 907-919.

Toth, D.J., Toth, J.T., Gulyas, G., Balla, A., Balla, T., Hunyady, L., and Varnai, P. (2012). Acute
depletion of plasma membrane phosphatidylinositol 4,5-bisphosphate impairs specific steps in
endocytosis of the G-protein-coupled receptor. J Cell Sci *125*, 2185-2197.

Trapaidze, N., Gomes, I., Bansinath, M., and Devi, L.A. (2000). Recycling and resensitization of
delta opioid receptors. DNA Cell Biol *19*, 195-204.

Yen, H.Y., Hoi, K.K., Liko, I., Hedger, G., Horrell, M.R., Song, W., Wu, D., Heine, P., Warne, T.,
Lee, Y.*, et al.* (2018). PtdIns(4,5)P2 stabilizes active states of GPCRs and enhances selectivity
of G-protein coupling. Nature *559*, 423-427.

Yin, W., Li, Z., Jin, M., Yin, Y.L., de Waal, P.W., Pal, K., Yin, Y., Gao, X., He, Y., Gao, J.*, et al.*
(2019). A complex structure of arrestin-2 bound to a G protein-coupled receptor. Cell Res *29*,
971-983.

Zhang, J., Barak, L.S., Anborgh, P.H., Laporte, S.A., Caron, M.G., and Ferguson, S.S. (1999).
Cellular trafficking of G protein-coupled receptor/beta-arrestin endocytic complexes. J Biol Chem
274, 10999-11006.

Zhuang, T., Vishnivetskiy, S.A., Gurevich, V.V., and Sanders, C.R. (2010). Elucidation of inositol
hexaphosphate and heparin interaction sites and conformational changes in arrestin-1 by solution
nuclear magnetic resonance. Biochemistry *49*, 10473-10485.

Zhuo, Y., Vishnivetskiy, S.A., Zhan, X., Gurevich, V.V., and Klug, C.S. (2014). Identification of
receptor binding-induced conformational changes in non-visual arrestins. J Biol Chem *289*,
20991-21002.
1436

Figure 1. Arrestin phosphoinositide binding is required for recruitment to some GPCRs A) cartoon depicting NanoBiT assay for measuring arrestin plasma membrane recruitment upon agonist stimulation. Upon complementation SmBiT and LgBiT form a functional NanoLuc luciferase. In key, "Phosphate" denotes phosphorylated Ser/Thr residues and "X" denotes KRK to 3Q mutant of β-arrestin. B) Two representative GPCRs, β1AR and NTSR1 illustrate data obtained for β-arrestin recruitment by NanoBiT assay shown in panel A. Data were collected over time after agonist addition (t=0 min), and values are shown as luminescence fold-change (over vehicle treatment) ± standard deviation (measured as 2 technical replicates for each of n=3 independent experiments). Colors denote concentrations of agonist used for stimulation. Agonists used were isoproterenol for β1AR and neurotensin for NTSR1. Grey boxes mark the time region (10-15 minutes post agonist addition) over which luminescence is integrated, for each concentration of agonist, to produce concentration response curves (bottom). WT and 3Q amplitudes were determined as the difference of fitted pre- and post-transition plateaus. C) Plot of LOF values for panel of tested GPCRs. Points represent LOF value obtained as ratio of WT and 3Q recruitment, and error bars reflect error in LOF derived from standard errors of fits (see methods). Dashed ellipses denote clusters obtained from k means clustering of data (see methods); AT1R is in cluster 2 for both βarr1/2, while β2AR-V2C is split with βarr1 (cluster 2) and βarr2 (cluster 1). Vertical grey lines denote LOF = 0 and LOF =1; vertical purple and orange lines reflect the centers of the respective clusters from k means and correspond to LOF = 0.06 and LOF = 0.73, respectively.

Figure 2. Receptor phosphorylation patterns govern PIP-dependence for arrestin recruitment. A) Left, schematic of human NTSR1 showing motifs in receptor ICL3 and C-terminus that are subject to phosphorylation. Phosphorylation sites examined in this study are shown in red and numbered 1-10 (above). Residue numbers corresponding to the region of human NTSR1 are listed at the start and end of the shown sequences. Construct key shows possible phosphosites as empty boxes, which when mutated to alanine are filled with an "X". Plasma membrane recruitment of arrestin upon stimulation of cells expressing different NTSR1 constructs, measured using the NanoBiT assay described in Figure 1. Right, points represent LOF value obtained as ratio of WT and 3Q recruitment, and error bars represent standard error of fits (see methods). Points are colored based on cluster designation obtained from k means clustering of all receptor-arrestin recruitment data. B) Translocation of βarr1 to endosomes upon stimulation of cells expressing different NTSR1 constructs, measured using an endosome bystander NanoBiT assay, as described in Figure S1. Points represent recruitment (fold chance over basal upon stimulation) for WT and 3Q recruitment, denoted by circles and triangles, respectively. Points are based on data from n=3 biological experiments. Error bars represent standard error of fit used to determine recruitment. Points are colored based on the cluster assignment of that mutant. C) Internalization, measured by loss of cell-surface receptors upon agonist stimulation, for Δβarr1/2 cells expressing

NTSR1 or β2AR constructs and transfected with arrestin constructs indicated. Values represent independent experiments ($n = 5$-10). Internalization by 3Q βarr1 and mock were compared to WT using a two-tailed paired t-test. ns: $p > 0.05$; *: $p \leq 0.05$; **: $p \leq 0.01$; ***: $p \leq 0.001$; ****: $p \leq 0.0001$.

Figure 3. Lipid binding stabilizes core-engaged arrestin complexes. A) cartoon of complexing efficiency assay. Size-exclusion chromatography (SEC) resolves complex from components. B) Representative experiment showing SEC chromatograms with vertical dashed lines indicating free NTSR1, complex, and free arrestins. C) Complexing efficiency, for NTSR1 with indicated arrestins. Boxplots: center line, median; box range, 25–75th percentiles; whiskers denote minimum–maximum values. Individual points are shown (n=6 independent experiments). Two-tailed unpaired t-test used to compare conditions. ns: $p > 0.05$; ****: $p \leq 0.0001$. D) Cartoon showing equilibrium of NTSR1-arrestin complex. Pink star denotes L68bim probe used for experiment shown in panel E. E) Bimane spectra for L68bim labeled βarr1 in complex with NTSR1. All NTSR1 samples contained diC8-PI(4,5)P2 (4.1 μM) Boxplots: center line, median; box range, 25–75th percentiles; whiskers denote minimum–maximum values. Individual points are shown (n=3 independent experiments). V2Rpp-NTSR1 (GRK5p) and V2Rpp-NTSR1 (unphos) + V2Rpp were compared by two-tailed unpaired t-test. ns: $p > 0.05$; *: $p \leq 0.05$. Apo indicates free arrestin; unphos indicates unphosphorylated receptor and GRK5p indicates GRK5-in vitro phosphorylated receptor. Spectra are normalized to apo (100%) within each experiment and the fluorescence intensity at lambda max was used as the value. F) Free energy diagram illustrating how PIP-binding, by stabilizing the core-engaged state of the NTSR1-arrestin complex slows arrestin dissociation. Loss of the PIP-binding element of arrestin destabilizes the core-engaged state, shifting equilibrium towards the tail-engaged state leading to a higher degree of

complex disassembly. Removal of the arrestin C-terminus stabilizes the complex in the tail-engaged state and reduces disassembly even when core-engaged complex is destabilized by lack of PIP-binding.

Figure 4. PIP$_2$ alone promotes conformational changes in arrestin, including C-terminus movement, but not release. A) overlay of inactive (PDB: 1G4M) [grey] and active (PDB: 4JQI) [black] βarr1. The N and C lobes of βarr1 are indicated. Activation leads to reorganization of several loops, and the gate loop and finger loop are highlighted. Re-orientation of these loops from inactive (yellow) to active (green) can be monitored by site-specific fluorescence spectroscopy. In finger loop inset the sphere denotes Cα L68C which is labeled with bim. In gate loop inset, the sphere denotes Cα L293C which is labeled with NBD. An installed W residue replacing L167 dynamically quenches 293NBD. B) Spectra of bimane labeled (L68C) βarr1 in response to V2Rpp and PIP$_2$. Arrow indicates direction of spectral shift with increasing concentration. Values are mean ± SD (*n*=3 independent experiments). Spectra were normalized to the apo condition within a given experiment. C) Spectra of NBD labeled (L167W-L293C) βarr1 in response to V2Rpp and PIP$_2$. Arrow indicates direction of spectral shift with increasing concentration. Values are mean ± SD (*n*=3 independent experiments). Spectra were normalized to the apo condition within a given experiment. D) Left, cartoon showing how FRET change is linked to C-terminus release. Right, spectra of AF488/AT647N labeled (A12C-V387C) βarr1 in response to V2Rpp and PIP$_2$. Arrow indicates direction of spectral shift with increasing concentration. Spectra were normalized via donor intensity within a given experiment. Data shown are for a representative experiment (*n*=3 independent experiments).

Figure 5. PIP_2 enhances Fab30 binding to βarr1. A) Cartoon of surface plasmon resonance (SPR) experiments, where βarr1 is immobilized via N-terminal biotinylation and a Fab30 binder is injected in the presence of absence of PIP_2 or V2Rpp. B) Representative sensogram for SPR binding experiment. With WT βarr1 immobilized, Fab30 (1 μM) was injected alone or together with V2Rpp (40 μM) or diC8-PI(4,5)P2 (40 μM). The shown sensogram is representative of the outcome seen for independent experiments (*n*=3). Dissociation/regeneration phase not shown. C) Binding of Fab30 to immobilized arrestin constructs in the presence of different additives. Maximum binding is defined based on normalization of the observed response to the amount of arrestin immobilized for each construct. Additives: diC8-PG (40 μM), diC8-PI(3)P (40 μM), diC8-PI(4,5)P2 (40 μM) and V2Rpp (40 μM) were mixed with Fab30 (1 μM) and injected together. Points reflect independent measurements; open points represent the binding observed for the additive in the absence of Fab30. Fab30 binding was compared using a two-tailed unpaired t-test. ns: $p > 0.05$; *: $p \leq 0.05$; **: $p \leq 0.01$; ***: $p \leq 0.001$. D) The proportion of active-like βarr1 increases in the presence of PIP_2.

Figure 6. Model for phosphoinositide regulation of GPCR-β-arrestin complex assembly and disassembly. GPCRs stratify into two groups with respect to the strength of their interaction with β-arrestins: one group requires an interaction between β-arrestin and PIP$_2$ at the plasma membrane for recruitment (PIP-dependent), while the other does not (PIP-independent). In the case of PIP-dependent GPCRs, arrestin engagement is unstable and can result in dissociation of arrestin from the receptor, while maintaining an association with the plasma membrane (left panel). PIP$_2$ is enriched at CCSs and in both cases complex assembly can occur. During endocytosis, PIP$_2$ is depleted and for PIP-dependent GPCRs, the loss of this PIP$_2$ contact may facilitate dissociation of arrestin thereby allowing for receptor recycling. In contrast, a PIP-independent GPCR will retain the interaction with arrestin even once PIP$_2$ is depleted owing to the strong phosphorylation-dependent interactions; however, the full-engaged state of the complex is less stable in endosomes than at the plasma membrane, thereby allowing further G protein engagement to occur.

Figure S1. Arrestin phosphoinositide binding is required for plasma membrane recruitment to some GPCRs. A) cAMP response in HEK293 cells devoid of β-arrestins upon stimulation of endogenous β2AR with 100 nM isoproterenol (iso). Clone 1 (CL1) and Clone 2 (CL2) are

independent βarr1/2 knock-out cell lines (O'Hare et al. 2017). Data are normalized to response with Forskolin (Fsk)/3-isobutyl-1-methylxanthine (IBMX) and show mean with 95% confidence intervals (n=3 independent experiments). Two-way analysis of variance (ANOVA), Tukey's multiple comparison test. For CL2 * denotes $p < 0.05$ for WT vs. mApple over the interval of 17-32 minutes, while 3Q vs. mApple was not significant. For CL1 * denotes $p < 0.05$ for WT vs. mApple over the interval of 19-29 minutes, while 3Q vs. mApple was not significant. B) Quantification of expression for βarr1 and βarr2 (both WT and 3Q) NanoBiT constructs, as determined by western blot (Supplementary data figure 2). Mean values of 3-4 independent experiments were compared by a two-tailed unpaired t-test, where ns denotes $p > 0.05$, * $P ≤ 0.05$. Boxplots: center line, median; box range, 25–75th percentiles; whiskers denote minimum–maximum values. Individual points are shown. C) LOF is only weakly correlated with recruitment of WT β-arrestins. Data are mean LOF and mean WT βarr1/2 recruitment. βarr1 recruitment is shown as circles and βarr2 recruitment is shown as triangles. Data are colored based on assigned cluster. Dashed line shows expected linear relationship and R is the Pearson coefficient, with -0.51 reflecting a weak negative correlation. D) Plot of LOF data for plasma membrane bystander (CAAX) vs. LOF for direct recruitment. βarr1 recruitment is shown as circles and βarr2 recruitment is shown as triangles. Data are colored based on assigned cluster. Dashed line shows expected linear relationship and R is the Pearson coefficient, with 0.88 reflecting a very strong positive correlation. E) NanoBiT assay for measuring endosome translocation of βarr1. Cartoon of endosome bystander assay (left). βarr1 endosome recruitment data (right) with dashed ellipses to indicate clusters based on CAAX data. β-arrestin endosome recruitment determined by span of luminescence fold change. Data are mean ± SEM (n=3 independent experiments). Dashed line indicates three times the maximum signal measured in mock (receptor) transfected cells.

Figure S2. Loss of PIP binding slows β-arrestin recruitment to cluster 2 GPCRs. A) initial rate (0-5 minutes post-agonist stimulation) expressed as luminescence fold-change (FC)/min. Data from *n*=3 independent experiments fit independently (see methods). Boxplots: center line, median; box range, 25–75th percentiles; whiskers denote minimum–maximum values. For each receptor, and for each βarr1 and βarr2 WT and 3Q were compared by a two-tailed unpaired t-test, where ns denotes p > 0.05, * P ≤ 0.05, ** P ≤ 0.01, **** P ≤ 0.0001. B) Data from A) expressed as a difference in rate shows that with the exception of βarr1-TACR1 all cluster 2 receptors show faster recruitment of WT β-arrestin1/2 than corresponding 3Q mutant. Data are mean ± SEM (*n*=3 independent experiments).

Figure S3. Arrestin recruitment to NTSR1 mutants can be measured by NanoBiT recruitment assay. A) Expression of NTSR1 constructs in HEK293A cells used for NanoBiT assays. Boxplots: center line, median; box range, 25–75th percentiles; whiskers denote minimum–maximum values. Individual points are shown. Values are mean, relative to NTSR1-WT (*n*=4 independent experiments). For each construct, a comparison to NTSR1-WT by a two-tailed unpaired Wilcoxon test was performed, where ns denotes p > 0.05, * P ≤ 0.05. B) Direct complementation NanoBiT assay Emax for Sm-βarr1 interaction with Lg-CAAX for cells expressing NTSR1-WT as a function of mean fluorescence intensity (MFI), as determined by cell-surface staining. Amount of NTSR1-WT DNA transfected is written; blue arrow denotes 200 ng, the amount used in recruitment assays in Figure 2. C) As B, except the pEC_{50} of recruitment response upon NTS stimulation is plotted vs. MFI, instead of Emax. In both B and C, points represent mean values and error bars indicate 95% CI (*n*=3 independent experiments).

Figure S4. PIP binding stabilizes core-engaged arrestin complexes. A) LOF in complexing efficiency as determined by SEC. LOF = 1 corresponds to complete loss of complex formation, while LOF = 0 corresponds to no difference in complexing efficiency between WT and 3Q βarr1 (*n*=5 independent experiments). Boxplots: center line, median; box range, 25–75th percentiles; whiskers denote minimum–maximum values. Individual points are shown. compared by a two-tailed unpaired t-test, where **** P ≤ 0.0001. B) Binding of BODIPY-TMR PI(4,5)P2 to βarr1 (1-382) protein (WT or 3Q). Points are mean and error bars reflect 95% CI (*n*=5 independent experiments). Data were fit to a logistical function as described in methods and best fit values for B_{max} and K_d are provided with 95% CI in parentheses. C) Structure of transition from inactive (PDB: 1G4M) to active (PDB: 4JQI) βarr1 involves displacement of the βarr1 C-terminus (dark grey) by V2Rpp (blue). Two cysteine residues were added to a cys-less βarr1 background at positions A12 and V387 (pink spheres). These positions were labeled with fluorophores that, through FRET, allow for monitoring the position of the C-terminus. D) When labeled with a FRET pair, βarr1-12C/387C shows a high-FRET state in the absence of V2Rpp, and a low-FRET state when the βarr1 C-terminus is displaced by V2Rpp. FRET measured when βarr1 (WT or 3Q)-12C/387C-AF488-AT647N is bound to V2Rpp (0.5 μM), NTSR1 (GRK5p, 0.5 μM), or NTSR1 (unphosphorylated, 0.5 μM)+V2Rpp (0.5 μM). All samples containing NTSR1 were supplemented

with diC8-PI(4,5)P2 (0.5 μM). Apo βarr1 (WT or 3Q)-12C/387C-AF488-AT647N was normalized to 1.0 and βarr1 (WT or 3Q)-12C/387C-AF488-AT647N + V2Rpp (10 μM) was normalized 0.0 for each experiment (n=3 independent measurements) (right). Boxplots: center line, median; box range, 25–75th percentiles; whiskers denote minimum–maximum values. Individual points are shown.

Figure S5. PIP$_2$ allosterically triggers movement of the arrestin C-tail, but not release. A-B) Finger loop (L68C-bim) responses. %apo is scaled such that the fluorescence intensity (at λmax) for apo arrestin is 100% and each condition is scaled as a factor of apo. ND denotes not determined

values. Values for B_{max} (max response) and K_d (based on single-site binding fitting) are provided and ranges in parentheses correspond to 95% CI. Points are mean and error bars reflect 95% CI (*n*=3 independent experiments). C-D) Gate loop (L167W-293C-NBD) responses. %apo is scaled such that the fluorescence intensity (at λmax) for apo arrestin is 100% and each condition is scaled as a factor of apo. Values for EC_{50} (half maximal response) and k (rate constant based on single exponential decay) are provided and ranges in parentheses correspond to 95% CI. Points are mean and error bars reflect 95% CI (*n*=3 independent experiments). E-F) C-terminus release (A12C-V387C-AF488-AT647N) responses. %FRET is scaled such that apo arrestin is 100% and the highest concentration of V2Rpp (100 uM) is 0%. INF denotes infinite upper bound. ND denotes not determined values. Range of EC_{50} values is indicated in parentheses and represents 95% CI. Points represent mean and error bars reflect 95% CI (*n*=3 independent experiments).

Figure S6. Plasma membrane PIPs promote conformational changes in arrestin. A) Structure of soluble lipid derivatives examined in this work. B-I) Concentration response curves for L68bim βarr1. Values for B_{max} (max response) and EC_{50} (based on single-site binding fitting) are provided

and ranges in parentheses correspond to 95% CI. Points are mean and error bars reflect 95% CI (n=3 independent experiments). ND denotes not determined values. J) Summary of effect size (B_{max}) for different lipids with L68bim βarr1. Values represent B_{max} obtained from fitting independent experiments. ND is used for PG and PI(3)P as data could not be fit. Bars represent mean B_{max} and error bars denote standard deviation across the fits. K) A PI(4,5)P2 derivative bound in the C-terminal domain of βarr1 (PDB: 6UP7). The side chain of K232 was modeled based on PDB: 4JQI as it was not ordered in PDB: 6UP7.

Figure S7. Titration of interactions with immobilized βarr1 by SPR. Immobilized βarr1 construct (ligand) is listed as column heading. Rows represent analyte flowed during titration. K_d and B_{max} values obtained as described in methods are listed with 95% CI range in parentheses. ND denotes

value not determined. PIP$_2$ binding was not fit (D-F), whereas B$_{max}$ for Fab30+V2Rpp (M-O) failed to converge. In all cases the darkest blue curve corresponds to the highest concentration and the darkest red curve corresponds to the lowest concentration. Titrations, as described in methods, for V2Rpp and PIP$_2$ (A-F) ranged from 40 μM to 78.1 nM. Titrations of Fab30 ranged from 2000 nM to 3.9 nM with fixed concentration of V2Rpp (40 μM) or PIP$_2$ (40 μM), as indicated. Vertical axes are raw RU, and not corrected for channel loading.