Impact of hedonic value of stimuli on sampling dynamics during a preference test

Ben Ballintyn, John Ksander, Donald Katz, and Paul Miller

Brandeis University

Abstract

The decision of whether to continue with a current action or to stop and consider alternatives is ever present in the life of an animal. Such continuous-time decision making lies at the heart of food preference tests whose outcomes are typically quantified by a single variable, the total amount consumed. However, the dynamics that give rise to such a quantity in terms of durations of bouts of sampling at a stimulus before pauses, and the impact of alternative stimuli on those bout durations and subsequent actions following a pause, can contain a richness of behavior that is not captured in a single palatability measure. Here we carry out multiple analyses of these dynamics, with a particular focus on assessing how the hedonic value of one taste stimulus impacts the behavior of a rat sampling a second taste stimulus during a preference test. We find evidence for an explicit competitive interaction between bout durations, such that the more palatable a stimulus the longer the bout durations when the rat samples the stimulus and the shorter the bout durations at the alternative. Such competition is reproduced in a model of a neural circuit that could underlie the continuous decision of when to end a sampling bout. We find that the competitive impact on bout durations is relatively short-lived whereas a competitive impact on the choice of which stimulus to approach following a pause persists. Such a discrepancy in the timescales for the decay of the impact of the alternative stimulus suggests different neural processes are involved in the choice of which stimulus to approach versus the choice of how long to sample from it. Since these two choices together combine to determine net consumption and therefore the inferred palatability or preference of a gustatory stimulus, our results suggest that palatability is not a unitary quantity but the result of at least two distinct, context-dependent neural processes.

Introduction

The ability to efficiently forage for food and other resources is critical to most animals' survival, and evolution has doubtless shaped the neural circuitry responsible for decision-making to optimize this task (Hayden et al., 2011; Pearson et al., 2014). There are many types of questions that an animal must answer during a foraging bout. A first class of questions lies within the domain of perceptual decision-making, where an animal uses its senses to gather information about the state of the environment to answer questions such as "is there a predator present?". A second class of problems constitutes value-based decision-making which uses available information to choose between different action plans such as "Is it worth continuing to gather food here, or should I search for a more abundant area?".

Studies of value-based decision-making have typically fallen into two categories: 1) "self-control" or "delay-discounting" tasks where animals tradeoff waiting times and payoff sizes (Bateson & Kacelnik, 1996; Blanchard et al., 2013; Pearson et al., 2010; Stephens, 2002; Stephens & Anderson, 2001) and 2) "stay-switch" or "patch-leaving" task where animals are presented with a source of reward and must decide when to leave it in search of a better alternative (Barack et al., 2017; Blanchard & Hayden, 2014; Constantino & Daw, 2015; Hayden et al., 2011). Patch-leaving tasks better represent the natural situation where animals have sequential interactions with individual reward sources as opposed to the simultaneous presentation of (cued) alternatives. While these prior patch-leaving studies have helped elucidate the neural circuits responsible for foraging behavior, they ubiquitously utilize a trial-based structure and patch options are presented randomly, as assumed by the Marginal Value

Theorem (Charnov, 1976), as opposed to in the wild, where animals use their experience with

the environment to direct their encounters with patches. We suggest that two- or multi-bottle

preference tests represent a simplified naturalistic foraging scenario where an animal can

rapidly learn about the state of its environment and direct its encounters with reward options.

Preference tests are used widely to measure the relative hedonic values (or palatabilities) of a

set of stimuli, with the degree of preference based on the amount of interaction the subject has

with each stimulus. For example, in taste preference tests, the relative palatabilities of pairs of

substances are measured by the total amounts consumed or the number of licks at each food

source. It is assumed that the decision of the animal to lick (or eat) more of one tastant than

another, given equal opportunity for time with both, reflects an underlying preference. While

these tests have been used to determine the relative preferences of different options, little

attention has been paid to the decision dynamics of animals during such preference tests. The

lack of data quantifying the underlying behaviors that lead to the overall preference precludes

the assessment of models of this type of decision-making. Indeed, one of the goals of our

analyses is to test predictions of our recent modeling paper (Ksander et al., 2021).

Several questions can be asked about the sequences of decisions that animals make during

these preference tests. Perhaps the most obvious and important is: Do animals rapidly settle on

a favorite option or continue to switch between and sample both options? If the animal has a

clear preference between two options (as revealed by the total amount consumed), then an

intuitive and theoretically optimal strategy is to first sample both options to determine a

favorite and then spend all of the remaining time sampling the favored option (or until the

source is exhausted or the animal is sated). In this case, it would be very difficult to quantify the

palatability of a tastant: only a ranking would be possible. If the answer to the above question is

"no" (as we find in our data), then several additional questions can be asked.

The first is, given that the animals switch back and forth between the two options, how do

sampling times at one option depend on that option's palatability—as measured by total

amount consumed in sessions without alternatives—and on the palatability of the alternative?

To answer this question, we analyze durations of bouts of licking, which are comprised of series

of rapid licks without significant pauses, to assess whether and how the behavior at one lick-

spout depends on the contents of the alternative lick-spout.

Separate bouts are demarcated by pauses, following which the animal can either return to a

new bout of licking at the same spout or switch to the alternative spout. Our analyses of bout

durations separately following returns versus switches provides an indication of how the

influence of the alternative sample on current behavior decays over time. Lastly, an analysis of

the choice of which spout to choose after each pause, provides an insight into how the animal

weighs the relative value of the two spouts. Distinct dynamics of that choice probability would

provide evidence that the choice of which sample to taste and the choice of for how long to

taste it are separate and distinct processes.

It is worth noting that competition arises in preference tests without the need for any direct

interaction between the hedonic value of one alternative and the behavior displayed at the

other. The source of implicit competition is the limited time available in most tasks, or even if

time were not limited in a food preference test, the total amount of food desired until satiety

provides a limit. Such limits mean that the more time spent and the greater the consumption at

one sampled stimulus, the less time available at the alternative, even if the behavioral dynamics

were not altered. Indeed, one can hypothesize that an association with a more appetizing

stimulus might boost the perceived hedonic value of a paired neutral stimulus leading to longer

bouts at the neutral alternative, even as total amount consumed at the neutral alternative goes

down due to the fewer visits there. On the contrary, if behavior in preference tests resembles

that during foraging, one would anticipate that the greater the value of the alternatives, the

less time spent at a particular source. A primary goal of this work is to identify the nature of the

across-stimulus interaction.

To summarize, we analyze the behavior of rats engaged in a naturalistic continuous-time taste

preference task. We also compare the behavioral dynamics with the dynamics of a simulated

circuit of model spiking neurons designed to possess two states, one representing the ongoing

choice to sample a stimulus, the other to leave that stimulus. Competition between successive

stimuli can arise in the model from adaptation-like processes, leading to predictions of a

competitive interaction between one bout of sampling a stimulus and the subsequent bout

with the alternative stimulus. We assess our behavioral findings for evidence of such a

competitive interaction.

Results

Measurement of palatability

To study stay-switch decision dynamics, rats were put through two weeks of preference testing

(Figure 1A/B). On each day, they were given one hour to freely sample a random two solutions

drawn (with replacement) from a possible four (0M, 0.01M, 0.1M, or 1M) NaCl solutions,

selected to provide three significantly different palatabilities (Sadacca et al., 2012). Licks at each

solution spout were recorded using a custom circuit and identified using a semi-automated

process (see methods).

As a first look at the rats' preference behavior, we confirmed the rank order of the relative

palatabilities of these solutions (Figure 2A), by measuring the total number of licks to each

solution on days where the solution was paired with itself and dividing by the mean number of

licks to dH2O on dH2O only days. The previously determined palatability ranking (0.1M > 0.01M

> 1M,(Sadacca et al., 2012)) was recapitulated, and no sex-specific differences were found

(0.01M: $z = .17$, $p = .86$; 0.1M: $z = -1.056$ $p = .29$; 1M: $z = .51$, $p = .61$); data from both sexes

were combined for all analyses in which different solutions were pitted against one another

(see Methods and Figure 1).

*Figure 1. Behavioral setup and example behavior. **A)** 1' x 1' custom acrylic chamber had a solution spout available through the*

left and right walls each containing 25mL of 1 of 3 different NaCl solutions (0.01M, 0.1M, 1M) or dH2O. Rats were allowed to

*freely move and sample from either spout over the course of 1 hour. **B)** Preference test timeline. Rats were given 2 habituation*

days with 1 bottle of dH2O on opposite sides across sessions. This was followed by 2 weeks of sessions where each week started

*with a session of dH2O only followed by all 6 combinations of NaCl solutions. **C)** Example licking data. Each rectangular*

deflection is one lick. D) Example sampling data from a session with 0.01M and 0.1M NaCl solutions. Yellow stripes represent

active sampling at the corresponding solution.

Given the observed differences in palatabilities, we expected different distributions of sampling

durations, with more palatable solutions having on average longer durations of lick bouts. This

expectation was borne out: the distributions of bout durations of all solutions were well

approximated by exponential distributions (Figure 2C-E), with decay constants akin to the mean

time of bouts at each solution; these bout duration distributions are mostly commensurate

with the calculated palatability of each solution (dH2O: mean 13.39 ± .79s, .01M: mean 24.33 ±

1.69s, .1M: mean 27.54 ± 1.72s, 1M: mean 4.72 ± .29s).

*Figure 2. Effect of palatabilities on bout duration. **A)** Relative palatabilities of the 3 NaCl solutions relative to water.*

Palatabilities are based on the total number of licks at each solution during sessions where the solution was paired with itself.

*No sex specific differences were found (0.01M: two-tailed: z = .17, p = .86, **.1M:** z = -1.06, p = .29, 1M: z = .51, p = .61). Based on*

*pairwise comparisons (the rank order of palatabilities from highest to lowest is (0.1M, 0.01M, H2O, 1M). **B)** Cumulative*

distribution of lick times. Rats continue to lick throughout the session but less over time. These CDFs were used to distinguish

*early and late bouts for each rat. **C)** Bout duration distributions were fit well by exponential distributions. Frequency of each*

*bout duration for 0.01M NaCl with y-axis on a log-scale. Linear fits to the exponential data are shown in orange. **D)** Same as (C)*

*but for bouts at the 0.1M solution. **E)** Same as (C,D) but for bouts at the 1M solution.*

Impact of relative palatability on bout duration

While a higher palatability of the currently sampled solution translates into longer sampling

bouts, a critical unanswered question is how the palatability of the alternative solution in a

preference test impacts these sampling bout durations. We considered three possibilities: 1) a

high alternative palatability will have an appetitive effect, increasing the perceived palatability

of the current solution and leading to longer sampling bouts; 2) conversely, a higher alternative

palatability could reduce the perceived palatability of the current solution, leading to shorter

sampling bouts; and 3) the palatabilities of alternative choices could have no impact on bouts at

the current solution. Implied in hypotheses 1 or 2 is the maintenance of a memory of the

alternative solution's value (palatability).

To evaluate the above possibilities we performed multilinear regression, predicting bout

duration as a function of the palatability of available alternatives. As suggested by differences in

mean bout durations across solutions, regression coefficients for the current solution's

palatability were significantly positive (z = 4.09, p = 2.15e-5, mean = 15.78 $\pm$ 1.63) – that is, the

more palatable a stimulus the longer the bouts of licking at it (Fig. 3A). Alternative palatability

coefficients were found to be significantly negative (mean -5.69 $\pm$ 1.32), consistent with

possibility 2 above (Fig. 3B)—durations of bouts are shorter when the alternative stimulus is of

higher palatability.

*Figure 3. Effect of current/alternative palatability on duration of lick bouts. **A)** Example results from 1 rat of a multilinear*

regression model for predicting bout duration with the currently sampled solution's and alternative solution's palatabilities as

*factors. **Ai)** Using best fit regression coefficients for 1 rat, bout duration is plotted against the current solution's palatability for*

*all 3 possible alternative solutions. **Aii)** Same as (Ai) but plotting bout duration vs. the alternative solution's palatability for 2*

*levels of the current solution's palatability. **B)** Current and alternative palatability regression coefficients normalized by the*

mean bout duration for each rat. Normalized coefficients for current palatability are significantly positive (right-tailed: z = 4.09,

p = 2.15e-5) and those for alternative palatability are significantly negative (left-tailed: z = -3.57, p = 1.7e-4).

The above results were stable across the course of the session, even though bout durations in

general decreased over time (likely due to satiation). When we split sessions into 'early' and

'late' portions based on a per-animal criterion (we used the 2nd derivatives of each rat's

cumulative distribution of lick times to detect the "kink" in the curve of Fig. 2B, where licking

slowed from a high rate to a lower rate) and performed the same multilinear regression on

early/late bouts separately, we found no significant change in (normalized) regression

coefficients between the early and late portions of the session (Fig. 4A/B).

*Figure 4. Effect of palatability on bout duration is constant across early and late portions of the session. **A)** Normalized*

regression coefficients for current solution palatability for bouts in the early or late portion of the session. Current palatability

coefficients were significantly positive for both the early (right-tailed: z = 4.06, p = 2.47e-5) and late (right-tailed: z = 4.09, p =

2.15e-5) portions of the task. Coefficients were not significantly different across portions of the session (paired: z = -.11, p = .91).

***B)** Same as (A) but for the alternative solution's palatability. Normalized coefficients were significantly negative for both early*

(left-tailed: z = -2.99, p = 1.4e-3) and late portions of the session (left-tailed: z = -2.76, p = 2.9e-3). Coefficients were not

significantly different across portions of the session (paired: z = -.76, p = .45).

We next split bouts into those following stay or switch decisions to ascertain whether the same

decision process was at play for each type of decision. We again repeated the multilinear

regression analysis on these groups individually. We found that regression coefficients for

current palatability are similarly positive following stays (z = 4.09, p = 2.15e-5, mean = 15.46 $\pm$

1.93) and switches (z = 4.06, p = 2.4e-5, mean = 16.04 $\pm$ 2.38), with no significant difference

between the two groups (z = .011, p = .91, Fig. 5A). In contrast, there is a significant difference

in alternative palatability coefficients in the post-stay vs. post-switch bouts: coefficients for the

post-switch bouts were significantly more negative (z = 4.06, p = 2.4e-5) than those following a

stay decision, which were themselves not significantly different from zero (z = -1.7, p = .088,

226   mean = -1.09 ± 1.61, Fig. 5B). This result suggests that information regarding the alternative

227   solution may only factor into decisions about sampling times only following a switch between

228   the two samples.

Figure 5. Difference in impact of current/alternative palatability on bout duration following a stay or switch decision. **A)**

Normalized multilinear regression coefficients for the currently sampled solution's palatability are significantly positive following

both a stay (right-tailed: z = 4.09, p = 2.1e-5) and switch decision (right-tailed: z = 4.06, p = 2.4e-5). Coefficients were not

significantly different across stay/switch conditions (paired: z = .011, p = .91). **B)** Normalized multilinear regression coefficients

for the alternative solution's palatability are not significantly different from zero following a stay decision (two-tailed: z = -1.7, p

= .088) but are significantly negative for bouts following a switch decision (left-tailed: z = -4.09, p = 2.1e-5). Coefficients for bouts

following a switch decision are significantly more negative than those for bouts following a stay decision (paired right-tailed: z =

4.06, p = 2.5e-5).

Lack of dependence of results on bout definition criteria

For the analyses described above, we define a 'licking bout' as sequences of licks which had no

period of >2s of no contact with the lick spout. Here, 'contact' includes brief periods of nose-

poking at the spout. While this definition of a 'licking bout' slightly overestimates the total time

spent licking, brief periods of nose poking in between licks represent active engagement with

the spout rather than a decision to stop sampling or switch to the alternative. An inter-lick-

interval of 2s was used as rats never switched between solutions in <2s.

*Figure 6. Using a 200ms ILI interval criterion to define bouts does not significantly alter the results. **A)** As in Fig 3B, normalized*

multilinear regression coefficients for predicting bout duration using the currently sampled and alternative solutions'

palatabilities as factors are shown. Current palatability coefficients were significantly positive (right-tailed: z = 4.09, p = 2.1e-5)

*and alternative palatability coefficients were significantly negative (left-tailed: z = -3.15, p = 8.2e-4). **B)** As in Fig 5A, normalized*

multilinear regression coefficients for the currently sampled solution's palatability are significantly positive following both a stay

(right-tailed: z = 4.09, p = 2.1e-5) and switch decision (right-tailed: z = 3.8, p = 7.3e-5). Coefficients were not significantly

*different across stay/switch conditions (paired two-tailed: z = -1.7, p = .089). **C)** Multilinear regression coefficients for the*

alternative solution's palatability are shown for models predicting bout durations following a stay or switch decision. Using this

criterion, regression coefficients for bouts following a stay decision are significantly negative (left-tailed: z = -3.08, p = .001) and

following a switch decision (left-tailed: z = -3.25, p = 5.8e-4). Coefficients for bouts following a switch decision are significantly

more negative than those for bouts following a stay decision (paired right-tailed: z = 2.82, p = .0024).

Of course, this is only one dividing line that could be used. Prior studies of licking

microstructure in rats (Davis, 1996; Davis & Smith, 1992) have grouped licks into 'bursts' or

'clusters' based on a <250ms or >500ms inter-lick-interval (ILI) criterion. To test that the results

presented above are not artifacts of our choice of bout definition, we repeated all the above

analyses using a 200ms ILI criterion. In this re-analysis, the magnitudes of the resulting

regression coefficients are much smaller, since bout lengths themselves are much shorter

(Supp. Fig. 1). Nonetheless, all the qualitative results presented above hold (Figs. 6/7):

coefficients for current palatability are significantly positive for early vs. late and stay vs. switch

bouts, and coefficients for alternative palatability in early and late bouts do not differ; while

coefficients for alternative palatability are significantly negative following a stay decision using

this bout criterion ($z = -3.02$, $p = .0012$, mean = $-.61 \pm .19$), they are again significantly more

negative (mean = $-2.6 \pm .68$) following a switch decision ($z = 2.59$, $p = .0047$).

One interesting difference did arise with this more stringent bout length criterion, as revealed

in Figure 6C compared with Figure 5B. The more stringent criterion split many prior single bouts

into multiple bouts of shorter duration. The shorter duration of bouts meant that the time

passed from a sampling of the alternative stimulus would often be less than previously for a

repeated bout of sampling at a stimulus – that is a bout of sampling following a "Stay" decision.

As a result, in Figure 6C we see a small significant impact of the alternative stimulus following a

"Stay" decision that was absent in Figure 5B where bout durations were longer. Such a finding

is consistent with a model in which the impact of the alternative stimulus on a current bout's

duration decays gradually over a period of many seconds after leaving that stimulus.

Indifference of results to change in rank order of palatability

As noted above, we calculated palatabilities using data from days in which identical solutions

were available at both spouts (this was done to separate the data used to compute

palatabilities from those used in the multilinear regressions). Using this method, 0.1M NaCl was

found to be significantly more palatable than 0.01M (Figure 2A). However, on days in which

0.1M NaCl was paired with 0.01M, rats licked more—on average, 1.3x as much—for 0.01M than

0.1M. That is, the 0.01M solution seemed more palatable than the 0.1M solution in direct

comparisons.

To the best of our knowledge, this is the first time a study has compared palatabilities obtained

by comparison to water and through direct comparison. This difference might be explained by

an appetitive effect whereby the palatability of the 0.1M is reduced when paired with the less

palatable 0.01M with the converse being true for the 0.01M (made more palatable by pairing

with 0.1M). Alternatively, these differences could simply represent the inherently context-

dependent nature of palatability.

In either case, we tested whether our prior results were impacted when the palatability of the

0.01M solution was defined as a factor of 1.3 times greater than that of the 0.1M solution. We

find that the results for the alternative palatability regression coefficients do not change. That

is, in aggregate, coefficients for alternative palatability are significantly negative ($z = -3.3$, $p = $

4.5e-4, mean $= -5.69 \pm 1.32$, Fig. 7A), coefficients for bouts following a switch decision were

significantly more negative than for those following a stay decision ($z = 4.09$, $p = 2.15e-5$, Fig.

7C), and coefficients were not significantly different between early and late portions of the

session ($z = -.5$, $p = .61$, Supp Fig. 2B). That is, our qualitative results are robust to whether the

0.01M or 0.1M solution is the more palatable and all conclusions arise from those two solutions

being more palatable than the 1M NaCl solution.

Figure 7. Impact of current/alternative palatability on bout duration when artificially setting Palatability(.01M) = 1.3 x

*Palatability(.1M). **A)** As in Fig 3B, multilinear regression coefficients for predicting bout duration using the currently sampled and*

alternative solutions' palatabilities as factors are shown. Current palatability coefficients were significantly positive (right-tailed:

*z = 4.09, p = 2.1e-5) and alternative palatability coefficients were significantly negative (left-tailed: z = -3.57, p = 1.7e-4). **B)** As in*

Fig 5A, multilinear regression coefficients for the currently sampled solution's palatability are significantly positive following

both a stay (right-tailed: z = 2.17, p = .0148) and switch decision (right-tailed: z = 3.9, p = 4.9e-5). With these artificially altered

palatabilities, the coefficients for bouts following a switch decision were significantly more positive than those for bouts

*following a stay decision (paired left-tailed: z = -2.2, p = .014). **C)** As in Fig 5B, multilinear regression coefficients for the*

alternative solution's palatability are shown for models predicting bout durations following a stay or switch decision. With the

artificially altered palatabilities, regression coefficients for bouts following a stay decision (left-tailed: z = -2.56, p = .0052) and

following a switch decision (left-tailed: z = -4.09, p = 2.15e-5) are significantly negative. Coefficients for bouts following a switch

decision are significantly more negative than those for bouts following a stay decision (paired right-tailed: z = 4.09, p = 2.15e-5).

There are however some minor differences regarding the coefficients for current palatability.

Coefficients for current palatability are significantly more positive following a switch decision (z

= -3.05, p = .0011, Fig. 8B) and normalized regression coefficients in the late portion of the

session were significantly smaller (less positive) than those in the early portion of the session (z

= 3.18, p = 7.3e-4).

Impact of palatability on transition probability

Thus far, our results describe sampling duration as a function of the palatabilities of the two

solutions. To fully understand the impact of palatability on choice dynamics, we also asked

whether the palatability of the current or alternative solution impacted the transition

probabilities between the solutions. A transition could be from a solution back to the same

solution, if following a bout of licking there is a pause then a return to the same solution.

Therefore, we are assessing the degree to which, following a pause in licking, the rat returns to

the same solution or switches to the alternative. As with measurements of bout durations, the

choice to return or switch could depend on both the palatability of the most recently sampled

("current") solution and that of the alternative.

To dissociate the contributing factors, we compared the transition probabilities between pairs

of solutions with either a common source (e.g. 0.01M -> 0.1M and 0.01M -> 1M) or a common

target (e.g. 0.01M -> 0.1M and 1M -> 0.1M). If the palatability of the current solution was to

influence transition probability, this influence would be reflected in a higher probability of

switching to a common target taste from a taste with a low palatability than from a taste with a

high palatability. Similarly, if alternative palatability was to impact transition probability, this

would be reflected in a higher probability of switching from a common source to a solution with

high palatability.

We find evidence that palatability of both the current and the alternative solution impacts the

transition probabilities (Fig. 8A-B). These results are further supported by a logistic regression

model trained to predict switches based on the current and alternative palatability. In the

regression model, both current palatability (p = 4.89e-31, coefficient = -.948, 95% CI = [-1.1 -

.78]) and alternative palatability (p = 3.15e-21, coefficient = .63, 95% CI = [.5 .76]) are found to

be significant factors.

Figure 8. Comparison of transition probabilities for transitions with a common source/target. **A)** Transition probabilities for transitions with a common source (0.01M -> 0.1M and 0.01M -> 1M, 0.1M -> 0.01M and 0.1M -> 1M, 1M -> 0.01M and 1M -> 0.1M) reveal the influence of a memory of the alternative solution's palatability. **Ai)** P(0.01M -> 0.1M) and P(0.01M -> 1M) are not significantly different (paired two-tailed: z = 1.51, p = .13). **Aii)** P(0.1M -> 0.01M) is significantly higher than P(0.1M -> 1M) (paired right-tailed: z = 3.73, p = 9.4e-5). **Aiii)** P(1M -> 0.01M) is significantly lower than P(1M -> 0.1M) (paired left-tailed: z = -2.52, p = 5.8e-3). **B)** Transition probabilities for transitions with a common target (0.1M -> 0.01M and 1M -> 0.01M, 0.01M -> 0.1M and 1M -> 0.1M, 0.01M -> 1M and 0.1M -> 1M) reveal the influence of the last sampled solution's palatability on switch

 *probability. **Bi)** P(0.1M -> 0.01M) is significantly lower than P(1M -> 0.01M) (paired left-tailed: z = -3.63, p = 1.3e-4). **Bii)***

*P(0.01M -> 0.1M) is significantly lower than P(1M -> 0.1M) (paired left-tailed: z = -4.09, p = 2.15e-5). **Biii)** P(0.01M -> 1M) is*

significantly higher than P(0.1M -> 1M) (paired right-tailed: z = 2.43, p = 7.5e-3).

No evidence for memory across days

Lastly, we investigated whether rats held a bias for the first side they visited in a session based

on their experience the prior day. To do this, we counted the number of times rats first visited

the side they preferred (had the most licks at) on the prior day and compared this to the

number expected. Given a null hypothesis of no memory across days, the expected number is

given by the binomial distribution with p = q = 0.5. Our results are consistent with the null

hypothesis that rats did not carry a preference for side across days ($\hat{p} = .485, 95\% \; CI =$

$[.424 \; .547]$, Fig. 9).

*Figure 9. No evidence for preference across days. **A)** Histogram showing the fraction of times rats first sampled from the side*

*they preferred on the prior day. **B)** (blue) Probability density function for the total number of times (across all rats) that rats first*

sampled from the side they preferred on the prior day which is given by the binomial distribution with N = 270, p = q = 0.5. (red

line) Total number of times rats first sampled from the preferred side from the prior day.

Comparison to spiking network models

In our modeling study (Ksander et al., 2021) we found competition in the durations of an

activity state representing bout duration in response to alternating stimuli. The competition

arose from a slow synaptic depression in the model so we hypothesized that the competition

between successive stimuli would diminish over the timescale of recovery from that synaptic

depression. We predicted that the impact of the alternative stimulus on bout duration would,

therefore, be significantly lower, during a second or later successive bout at the same stimulus,

as compared to the first bout at that stimulus following a switch, just as seen in the behavioral

data (Figs. 5-7). Therefore, we adapted the stimulus protocol in our prior study such that

following any state transition indicating the end of a bout of sampling, the subsequent stimulus

presented was chosen randomly, with a 50% probability for each of the two stimuli being

compared in that preference test.

Our results are shown in Figure 10, in which we produced regression coefficients in the

same manner as Figs. 5-7 but based on the state durations arising from three preference tests

(the distinct pairs of three stimuli of different strengths, representing different palatability) for

each of two types of network. Figure 10A-C depicts the results of an "entice-to-stay" network in

which stimuli of greater palatability were modeled by increased excitatory stimulus to neurons

whose activity represented a "stay" state that enhanced bout duration. The results are

qualitatively identical to the behavioral data (Figs. 3 and 5-7) with the alternative stimulus

having a competitive impact on bout duration (a negative regression coefficient, Fig. 10A) but

with the impact diminished following a repeat bout (a "stay" transition, Fig. 10C) at the same

stimulus. On the other hand, the results of a "repel-to-leave" network shown (Fig. 10D-F) did

not match the behavioral data well. The asymmetry between the model networks arises

because only in the "entice-to-stay" network does greater stimulus input correspond to longer

state durations, as needed to maximize the impact of synaptic depression.

*Figure 10. 'Fast' ('entice-to-stay') (A-C) but not 'slow' ('repel-to-leave') (D-F) model networks replicate rat behavior. **A)** As in Fig.*

3B regression coefficients for predicting bout duration as a function of current and alternative palatability are shown for 'fast'

networks. Similar to rats, coefficients for current palatability are significantly positive (right-tailed: z = 4.09, p 2.15e-5) and

*coefficients for alternative palatability are significantly negative (left-tailed: z = -3.83, p = 6.38e-5). **B)** As for rats, palatability*

coefficients for both stay (right-tailed: z = 4.09, p = 2.15e-5) and switch (right-tailed: z = 4.09, p = 2.15e-5) bouts were

*significantly positive and are not significantly different between groups (paired two-tailed: z = -1.8, p = 0.07). **C)** As for rats,*

alternative palatability coefficients for bouts following a stay decision were not significantly different from zero (two-tailed: z = -

1.28, p = 0.19) whereas coefficients for bouts following a switch decision are significantly negative (left-tailed: z = -4.06, p =

*2.47e-5)). **D)** In contrast, 'slow' networks did not replicate the pattern of coefficients of rats. Coefficients for current palatability*

were significantly positive (right-tailed: z = 4.09, p = 2.15e-5) but those for alternative palatability were not significantly

*different from (two-tailed: z = 0.14, p = 0.88). **E)** Palatability coefficients following both stay (right-tailed: z = 4.09, p = 2.15e-5)*

and switch (right-tailed: z = 4.09, p = 2.15e-5) bouts were significantly positive and coefficients for switch bouts were

*significantly more positive than those for stay bouts (paired left-tailed: z = -2.53, p = 0.0057). **F)** Alternative palatability*

Discussion

Palatability is typically measured as the amount of a food or solution consumed. The amount

solution consumed by a rat, for example, is equal to the number of licks taken multiplied by the

mean amount consumed per lick. Given that the mean amount consumed per lick has been

shown to vary little across stimuli or time, we assume that is constant in this study and measure

palatability as the number of licks of a solution. Since rats sample a solution in clearly

demarcated bouts of high frequency (approximately 6 Hz) regular sampling, we can identify two

distinct factors that impact the total number of licks and hence the palatability: the duration of

the bouts and the total number of bouts. In theory, bout durations could be independent of a

stimulus such that its palatability is only evident in the total number of bouts. However, prior

work (Davis, 1996) has shown that the more palatable a stimulus, the longer the bouts, a result

which we recapture here. In a more novel direction, we have assessed how, in a preference

test, the palatability of one stimulus impacts the behavior of an animal at the alternative

stimulus. Our main finding is that the more palatable one stimulus the shorter the bouts of

licking at the alternative as compared to when that alternative is paired with a less palatable

stimulus.

The second factor determining the number of licks at a spout is the total number of times the

spout is visited. Therefore, we also analyzed the transition probability during preference tests,

which indicates that after a bout of sampling from one lick spout how likely is an animal to

return to the same lick spout or to transition to the alternative. We find that the likelihood to

return increases with the palatability of the stimulus just tasted. In of itself such behavior would

produce competition in the total amount consumed, as it would produce more repeated bouts

at sources of high palatability, leading to fewer at an alternative. However, we also found that

following a pause in sampling at one lick spout memory of the alternative also impacted the

likelihood to return to the same spout, such that the more palatable the alternative, the less

likely to return. Thus, the choice of which spout to lick from is impacted by memory, which has

persisted beyond a single bout's duration, of the contents of both lick spouts.

Our findings of the competitive interaction between stimuli on bout durations supports a

recent model (Ksander et al., 2021) in which the duration of a bout is given by the duration of a

particular state of activity in a neural circuit. In the model, noise fluctuations terminate states of

activity, leading to an exponential-like distribution of state durations, just as we find in the

behavioral data. Moreover, the impact of the well-established neural-circuit level process of

synaptic depression in the model leads to a competitive impact between successive stimuli,

such that following a highly palatable stimulus a subsequent bout duration is shorter than

otherwise expected. Since the underlying biological processes have a limited timescale, we

tested and found in the model that such a competitive impact on bout durations diminishes

over time and is much weaker for bouts following a return to a stimulus when the time passed

since the visit to the alternative has increased. Indeed, we find in our behavioral data a similar

fact, with no impact of the alternative stimulus on durations of bouts of sampling that do not

directly follow a switch from that alternative stimulus.

When a food substance or taste stimulus is considered palatable of unpalatable, the implicit

suggestion is that palatability is a property of a substance to be ingested. However, in practice,

palatability is a measure of behavior—typically the total amount of a substance consumed—so

is inherently dependent on the state of an animal and the context in which the animal is

sampling the stimulus. In our study we find that even the rank order of palatability can be

altered depending on context. When a rat has two lick spouts available to it, both of which

contain the same solution, in accordance with prior work (Sadacca et al., 2012) we find rats lick

the spouts more often when the solution is 0.1M NaCl than when the solution is 0.01M NaCl

(Fig 1). Such a finding suggests that 0.1M NaCl is more palatable to rats than is 0.01M NaCl.

However, when one lick spout contains 0.1M NaCl and the other contains 0.01M NaCl, we find

rats lick more often at the spout containing 0.01M NaCl, suggesting a switch in relative

palatability and preference of the two salt solutions in the new context. Such a switch is

intriguing and its cause warrants further investigation. Fortunately, our findings in this paper on

the interactions between stimuli were robust to the switch. That is, whether we assumed 0.1M

NaCl was more or less palatable than 0.01M NaCl did not alter the findings on how the

palatability of the alternative stimulus impacted the behavior at a lick spout.

Our findings of a competitive interaction of palatability of bout durations of alternatives and

our model of the process contribute to the foraging literature, in which behavior is discussed

historically in terms of the Marginal Value Theorem (Charnov, 1976). The theorem prescribes

optimal behavior in an environment with multiple sources, at each of which the rate of reward

diminishes with the time an animal spends at the source. Specifically, an animal should only

stay at a food source until its rate of reward has depleted to the mean rate of reward it would

achieve by moving from alternative source to alternative source and remaining the optimal

time at the alternative sources. Our behavioral findings and model are in qualitative accordance

with such behavior in that the more palatable an alternative (*i.e.,* the greater the mean rate of

reward) the less time spent at a source while the reduction is ameliorated over time (the

greater the time between sources in foraging, the lower the mean rate of reward, so time spent

at a diminishing source increases). However, unlike in foraging studies, in preference tests the

potential rate of reward at a lick spout is constant, so if one spout contains more rewarding

solution than the other, the optimal behavior of an animal would be to stay at the more

rewarding spout as soon as it has sampled both. That the animals do not behave in such a

manner, but continue to sample even aversive stimuli many times, is either an indication of

limited memory duration (*i.e.,* they forget what is in each spout) or a strong drive to explore in

case the environment changes.

Acknowledgements

501 Acknowledgements
502

503   We are grateful for support from NIH (NIDA) NRSA grant DA051155 to BB, NIH (NINDS) grant

504   NS104818 via the Brain Initiative to DBK and PM, and to the Swartz Foundation for support to

505 JK. Computational resources were provided by the Brandeis HPCC which is partially supported

506 by the Brandeis Center for Bioinspired Soft Materials, an NSF MRSEC, DMR-2011846.

Methods

Behavioral apparatus

The preference test was carried out in 1' x 1' x 1' custom acrylic chambers. Each chamber has 3

holes through which rats could lick a stainless steel solution spout. There is one hole on each of

the left, right, and back walls of the chamber. For this study, only the left and right sides ever

had a solution spout. In order to record licks, a custom circuit, based on a published design

(Hayar et al., 2006) was used. A small voltage was applied to the stainless steel floor of the

chamber such that when the rat licked one of the solution spouts, a voltage deflection

(measuring the water-metal junction potential) was recorded. A RaspberryPi was used to both

supply power to the floor and record licks using custom Python software.

Preference test

22 adult Long-Evans rats (14 female, 8 male) from Charles River were water deprived for 22

hours prior to the first habituation session. The preference test timeline consisted of 16 1 hour

sessions of which the first 2 were habituation sessions with only 1 bottle of dH2O available on

one side of the experimental chamber (the side was switched for the second habituation

session). Following each session, rats were given 1 hour of ad lib access to water in their home

cage such that they were deprived of water for 22 hours prior to each session. After the two

habituation days, the first day of the preference test was always a session with 2 bottles of

dH2O. This was followed by 6 consecutive days of pairings of 3 NaCl concentrations (0.01M,

0.1M, 1M) including self pairings. This was then repeated for a second week such that each rat

experienced 2 dH2O only sessions and 2 pairings of each combination of NaCl concentrations.

These concentrations were used because they had been previously measured to have different

palatabilities (Sadacca et al., 2012) covering both palatable and unpalatable (at least relative to

water) tastants.

Lick identification

Licks were identified via a semi-automated process using custom MATLAB software. A simple

threshold could not be used to identify licks because both licks and nose pokes were picked up

as large voltage deflections. Additionally, occasionally a rat would maintain contact with the lick

spout while licking resulting in a sustained voltage deflection on top of which licks could be

seen. As a result, we produced a dataset of hand-identified licks from the data of the first few

rats and used MATLAB's neural network toolbox to train a bidirectional LSTM recurrent neural

network to predict the presence or absence of a lick at any point in time. These automatic

identifiers were then used as a first pass on all future data to capture presumptive licks, which

were then accepted/discarded by eye based on the stereotypical shape and timing of licks.

Lastly, a final pass over the data was made by eye to ensure that no licks were missed by the

neural network.

Lick bout identification

Following identification, licks were grouped together into 'bouts' based on 3 different inter-lick

interval (ILI) criteria. That is, we repeated all analyses described below using bouts defined by 3

different ILI criteria to determine how our results depended on this somewhat subjective

threshold. Based on previous studies of licking dynamics in rats (Davis, 1996) and our own

investigation of ILI distributions, we grouped together licks with 200ms or 500ms ILIs into lick

'bouts' (also referred to as lick clusters). We also used a more nuanced criterion which we

believe better represents active engagement with a lick spout (indicating an ongoing 'stay'

decision). This criterion consisted of grouping together adjacent licks in which there was no

period >2s between them in which there was no activity on the recording channel. This means

that if the rat nose-poked the solution spout in between licks such that the ILI was >2s but there

was intervening activity on the channel such that there was no period of >2s of silence, then

these licks would be grouped together. We included this criterion since we are primarily

concerned in this study with the rats' decisions to leave a solution spout and not on the

microstructure of their licking behavior.

Measurement of palatability

To measure the palatability of each concentration of NaCl, we analyzed data exclusively from

sessions where a solution was paired with itself. The palatabilities were defined relative to

water such that the relative palatability of solution X was:

567
$$Palatability(X) = \frac{N_{licks}(X)}{N_{licks}(dH2O)}$$

568 where $N_{licks}(X)$ is the total number of licks to tastant X across both sessions when X was paired

569 with itself. $N_{licks}(H2O)$ is the same except for H2O only sessions.

Linear and logistic regression models

To assess the impact of the palatability of the currently sampled or alternative solution on the

current bout duration, we performed multilinear regressions using MATLAB's *regress* function

to predict bout duration with both palatabilities as factors. This was done for all bouts together

as well as for subsets of bouts depending on if they were 'early' or 'late' in a session or

following a stay or switch decision.

As one method of measuring the impact of current/alternative palatability on switch

probability, we performed logistic regression using MATLAB's *fitglm* function to predict a switch

(0 or 1) with the last sampled solution's and alternative solution's palatabilities as factors.

Separating early and late bouts

We separated bouts for each session on a per-animal basis into 'early' or 'late' bouts by

analyzing the 2nd derivative of the cumulative distribution of lick times across all sessions. First,

a smoothed probability density function of lick onset times was computed using MATLAB's

ksdensity function with a bandwidth of 200s (controlling the amount of smoothing). The

cumulative density function of this pdf was then computed and its 2nd derivatives

approximated. The time point with the minimum 2nd derivative was then used as the divider

between early and late bouts.

Statistical tests

Unless otherwise stated, all z and p-values reported in this paper are from the Wilcoxon signed

rank test performed using MATLAB's *signrank* function. Tests of whether the median of a

distribution is significantly positive/negative utilized the right/left-tailed test respectively. Tests

of differences between distributions were done using a paired test where data points were

paired by animal or, in the case of the spiking model, points were paired by network.

Simulation protocol

Simulations were carried out using a recently published model (Ksander et al., 2021). In brief,

leaky integrate-and-fire neurons were designated excitatory or inhibitory and assigned either to

a group whose activity promoted a continuous decision of "Stay" at the current stimulus or a

group whose activity promoted a decision to "Leave" the stimulus. In the original paper the

"Leave" group of cells was denoted "Switch" as we had assumed that leaving one stimulus

meant the animal had to switch to the alternative. We produced new simulations for this paper

to investigate consequences of a "Return" to the same stimulus following a "Leave" decision

that ended a bout.

Connections between types of neurons were arranged in a manner of self-excitation and cross-

inhibition such that activity of one type of neurons (*e.g.,* representing "Stay") could maintain

itself while at the same time suppressing activity of the other type of neurons (*e.g.,*

representing "Leave"). Activity of the "Stay" neurons while suppressing the "Leave" neurons

would represent a "Stay" state in which the animal continues to sample a stimulus. Noise

fluctuations would irregularly cause a transition from such a "Stay" state to a "Leave" state. We

would ensure the "Leave" state was transient by reactivating "Stay" neurons to represent the

animal's commencement of the next sampling bout. As in our behavioral data, such noise-

induced transitions to terminate a bout of sampling resulted in an exponential-like distribution

of bout durations.

We assessed two types of model, in one type, the "entice-to-stay" model, the mean bout

durations were determined by stimulus-dependent input to excitatory neurons in the "Stay"

pool, such that the more palatable the represented stimulus, the greater the input. In the other

type, the "repel-to-leave" model, the mean bout durations were determined by stimulus-

dependent input to excitatory neurons in the "Leave" pool, such that the more palatable the

represented stimulus, the lower the input. We also test both types of model in this work.

All synapses in the model include synaptic depression, comprising a fast (300 ms) process

representing docking of new vesicles after vesicle release and a slow (7 sec) process

representing replenishment of a reserve pool of vesicles. Synaptic depression is key in

producing the competitive interaction across time as after a period of strong activity the

connections that sustain activity are weakened, impacting the response of the network to a

subsequent stimulus, until recovery of the supply of vesicles is complete.

Properties of model neurons

Individual neurons were simulated with an exponential leaky integrate-and-fire model

(Fourcaud-Trocmé et al., 2003) following the equation:

$$C_m \frac{dV_m}{dt} = \frac{E_l - V_m + \Delta_{th}\exp\left(\frac{V_m - V_{th}}{\Delta_{th}}\right)}{R_m} + G_{syn}\, S_I\left(E_{rev_I} - V_m\right) + G_{syn}\, S_E\left(E_{rev_E} - V_m\right)$$

$$+ G_{ref}(E_K - V_m) + G_{ext_I}\left(E_{rev_I} - V_m\right) + G_{ext_E}\left(E_{rev_E} - V_m\right)$$

where V_m is the membrane potential, C_m is the total membrane capacitance, E_l is the leak

potential, R_m is the total membrane resistance, Δ_{th} is the spiking range, V_{th} is the spiking

threshold, S is the synaptic input variable, G_{syn} and E_{rev} are the maximal conductance and

reversal potential for synaptic connections, G_{ref} is the dynamic refractory conductance, E_K is

the potassium reversal potential, and G_{ext} is the input conductance. The "E" and "I" subscripts

denote the variables specific to excitatory and inhibitory channels, respectively (e.g. S_E and

E_{rev_E} are the synaptic input and reversal variables for excitatory channels; S_I and E_{rev_I} are the

corresponding inhibitory variables). This equation simulates the neuron's membrane potential

until $V_m > V_{spike}$, at which point the neuron spikes.

When a neuron spikes, V_m is set to the V_{reset} value. Additionally, the neuron's refractory

conductance, synaptic output, s, and synaptic depression (noted as D) are updated according

to the equations:

$$G_{ref} \mapsto G_{ref} + \Delta G_{ref}$$

$$s \mapsto s + p_R D_{fast}(1 - s)$$

$$D_{fast} \mapsto D_{fast}(1 - p_R)$$

where ΔG_{ref} is the increase in refractory conductance, and p_R is the vesicle release probability

following a spike.

In the timestep immediately following a spike, the neuron's membrane potential continues to

follow the exponential leaky integrate-and-fire model equation. In this equation the separate

excitatory ($S_{E,i}$) and inhibitory ($S_{I,i}$) synaptic inputs for cell i are obtained from the sum of all

presynaptic outputs multiplied by the corresponding connection strengths, W_{ij}, from neurons j

(see *Network architecture and connections*):

$$S_i = \sum_j W_{ij} s_j,$$

each of which decay with the appropriate (excitatory or inhibitory) synaptic gating time

constant τ_S according to:

$$\frac{ds_i}{dt} = -\frac{s_i}{\tau_S}.$$

Likewise, refractory conductance decays with the time constant τ_{ref} according to:

$$\frac{dG_{ref}}{dt} = -\frac{G_{ref}}{\tau_{ref}}$$

The G_{ext} input conductance serves as both noisy-background and stimulus inputs in the same

manner. Inputs were modeled as Poisson spike trains with rates r_{noise} and $r_{stimulus}$, which

produce input spikes (from all sources) at timepoints $\{t_{sp}\}$. Please note, the noisy-background

includes both excitatory and inhibitory spiking input (included in G_{ext_I} and G_{ext_E} , respectively);

the r_{noise} parameter specifies the rate for both excitatory and inhibitory background noise. The

input conductance values for a given timepoint, t, are updated as:

$$G_{ext} \mapsto G_{ext} + \Delta G_{ext} \delta(t - t_{sp})$$

where the conductance increases by ΔG_{ext} at the time of each input spike. The input

conductance otherwise decays with the time constant τ_{ext} according to:

677
$$\frac{dG_{ext}}{dt} = -\frac{G_{ext}}{\tau_{ext}}.$$

The cellular parameters with values specific to excitatory neurons (e.g. that differ from

inhibitory values) are: $E_{rev_E} = 0\ mV$, $\tau_s = 50\ ms$, and $\tau_{ext} = 3.5\ ms$. The complementary

values for inhibitory neurons are: $E_{rev_I} = -70\ mV$, $\tau_s = 10\ ms$, and $\tau_{ext} = 2\ ms$. The

remaining parameters applicable to both excitatory and inhibitory neurons are: $G_{syn} = 10\ nS$,

$p_R = .1$, $\tau_{fast} = 300\ ms$, $\tau_{slow} = 7\ s$, $p_{slow} = .5$, $E_l = -70\ mV$, $E_K = -80\ mV$, $V_{reset} =$

$-80\ mV$, $R_m = 100\ M\Omega$, $C_m = 100\ pF$, $V_{spike} = 20\ mV$, $\Delta G_{ext} = 1\ nS$, $V_{th} = -50\ mV$,

$\Delta_{th} = 2\ mV$, $\tau_{ref} = 25\ ms$, and $\Delta G_{ref} = 12.5\ nS$. The Poisson spike-train parameters r_{noise}

and $r_{stimulus}$ are described in the next section. Neurons were simulated with a simulation

timestep $dt = .1\ ms$.

Synaptic depression

We modeled synaptic depression using two separate timescales, noted in the previous spike-

update equations as D_{slow} and D_{fast}. These two variables reflect, respectively, the fraction of

the maximum number of vesicles available in the reserve pool and the release-ready pool.

Following a spike, the variables recover to a value of one with different timescales, because

vesicles regenerate and are replenished slowly in the reserve pool, but may dock and become

release-ready much more quickly once available in the reserve pool

Specifically, D_{slow} represents the ratio of currently available reserve-pool vesicles out of the

maximum possible, that is $D_{slow} = \frac{N_{pool}}{N_{max}}$. These dock quickly at empty docking sites on the

timescale τ_{fast}, but are replaced slowly on the timescale τ_{slow}. D_{fast} represents the ratio of

docked vesicles out of total docking sites, that is $D_{fast} = \frac{N_{docked}}{N_{sites}}$. We also incorporate the

constant parameter, $f_D = 0.05$, which is equal to the ratio of the number of docking sites to the

maximum size of the reserve pool of vesicles, $f_D = \frac{N_{sites}}{N_{max}}$. Only docked vesicles can be released

immediately following a spike, such that upon each spike we update $D_{fast} \mapsto D_{fast}(1 - p_R)$

where p_R is the vesicle release probability.

During sustained spiking, the fast-docking can maintain a firing-rate dependent supply of

docked vesicles until the reserve pool depletes. Vesicles dock at empty sites according to:

708
$$\frac{dD_{fast}}{dt} = \frac{(D_{slow} - D_{fast})}{\tau_{fast}}$$

709   Reserve-pool vesicles fill the empty docking sites on the fast timescale τ_{fast}. On the other hand,

710   the reserve-pool regenerates much more slowly according to:

711
$$\frac{dD_{slow}}{dt} = \frac{(1 - D_{slow})}{\tau_{slow}} - f_D \frac{(D_{slow} - D_{fast})}{\tau_{fast}}$$

The first term represents the reserve-pool vesicle regeneration on timescale τ_{slow}. The second

term $-f_D \frac{(D_{slow} - D_{fast})}{\tau_{fast}}$ accounts for the vesicles lost due to docking.

Our model reflects the empirical evidence showing the effects of synaptic-depression at short

timescales on the order of milliseconds, and longer timescales on the order of seconds (Abbott

et al., 1997; Varela et al., 1997); depression timescales on the order of minutes have even

reported in non-mammalian animals (Tabak et al., 2000). Additional, recent evidence (Kusick et

al., 2020) directly supports our fast-depression mechanism where available vesicles quickly refill

empty docking sites. Our model provides a coherent mechanism for both fast-acting and long-

lasting synaptic depression effects.

Network architecture and connections

Each network consists of 250 individual neurons, split into two populations of 100 excitatory

cells (*i. e.,* "stay" and "switch" populations, E_{stay} and E_{switch}) and two populations of 25 inhibitory

cells (I_{stay} and I_{switch}). For each pair of connected populations (or for self-connected excitatory

populations) pairs of cells were connected probabilistically with a probability,

$P(connection) = .5$. The strength of connections was symmetric across "stay" and "switch"

populations but depended on whether presynaptic or postsynaptic cells were excitatory or

inhibitory, as indicated in Table 1.

Code Availability

The code used to simulate our model is freely available online at

https://github.com/johnksander/naturalistic-decision-making

Table 1. Model neuron parameters.

Name	Description	value

E_{rev}	Reversal potential	Excitatory cells: $0\ mV$ Inhibitory cells: $-70\ mV$
E_l	Leak potential	$-70\ mV$
E_K	Potassium potential	$-80\ mV$
R_m	Membrane resistance	$100\ M\Omega$
C_m	Membrane capacity	$100\ pF$
τ_s	Synaptic gating timescale	Excitatory cells: $50\ ms$ Inhibitory cells: $10\ ms$
V_{reset}	Reset membrane potential	$-80\ mV$
V_{spike}	Spike threshold	$20\ mV$
τ_{ext}	noisy-background conductance timescale	Excitatory cells: $3.5\ ms$ Inhibitory cells: $2\ ms$
G_{syn}	Synaptic max conductance	$10\ nS$
τ_{fast}	Fast depression timescale	$300\ ms$
τ_{slow}	Slow depression timescale	$7\ s$
p_R	Vesicle release probability	.1
f_D	Ratio of max docked vesicles to max reserve vesicles	.05
D_{fast}	Ratio of docked vesicles out of max possible	$\dfrac{N_{docked}}{N_{sites}}$
D_{slow}	Ratio of reserve- vesicles out of max possible	$\dfrac{N_{pool}}{N_{max}}$
ΔG_{ext}	Conductance step-increase to external input spike	$1\ nS$
V_{th}	exponential spiking-term threshold	$-50\ mV$

Δ_{th}	spiking range	2 mV
τ_{ref}	Refractory conductance timescale	25 ms
ΔG_{ref}	Step change in refractory conductance	12.5 nS
dt	Simulation timestep	.1 ms

A Model summary	
Populations	Stay: 1 excitatory, 1 inhibitory Leave: 1 excitatory, 1 inhibitory
Connectivity	Within-pool (stay or leave): I-to-E and recurrent E-to-E Cross-pool (stay-to-leave or leave-to-stay): E-to-I
Neuron model	Exponential Leaky Integrate and Fire (ELIF) with dynamic refractory conductance
Synapse model	Conductance based, step increase followed by exponential decay
Plasticity	Depression with two timescales
Input	Noisy background input: fixed-rate Poisson spike trains to all cells Stimuli: Poisson spike trains to E-stay and E-leave cells
Measurements	Spike trains, activity state-durations, connection strengths

B Populations		
Name	**Elements**	**Size**
E-stay	ELIF neurons	100
I-stay	ELIF neurons	25

E-leave	ELIF neurons	100
I-leave	ELIF neurons	25
Noisy background input	Poisson trains	500
Aversive stimulus	Poisson trains	100
Hedonic stimulus	Poisson trains	100

C Connectivity			
Name	Source	Target	Pattern
E-to-I	E-stay E-leave	I-leave I-stay	Random, $p = .5$, model-dependent fixed weight: 'Entice' network $\quad$ 0.0909 $\qquad$'Repel' network $\quad$ 0.4242
I-to-E	I-stay I-leave	E-stay E-leave	Random, $p = .5$, model-dependent weight: $\qquad$'Entice' network $\quad$ 9.6192 $\qquad$'Repel' network $\quad$ 9.4939
E-to-E	E-stay, E-leave	E-stay, E- leave	Random, $p = .5$, fixed weight, $W^{EE} = 0.0405$

D Neuron and Synapse Model	
Name	LIF neuron
Type	Dynamic leaky integrate-and-fire with dynamic refractory conductance

Subthreshold dynamics	$$C_m \frac{dV_m}{dt} = \frac{E_L - V_m + \Delta_{th}\exp\left(\frac{V_m - V_{th}}{\Delta_{th}}\right)}{R_m} + G_{syn} \cdot S_I \left(E_{rev_I} - V_m\right)$$ $$+ G_{syn} \cdot S_E \left(E_{rev_E} - V_m\right) + G_{ref}(E_K - V_m) + G_{ext_I}\left(E_{rev_I} - V_m\right)$$ $$+ G_{ext_E}\left(E_{rev_E} - V_m\right)$$ $$\frac{dG_{ref}}{dt} = -\frac{G_{ref}}{\tau_{ref}}$$ $$\frac{dG_{ext}}{dt} = -\frac{G_{ext}}{\tau_{ext}}$$
Spiking	If $V_m > V_{spike}$: 1. Emit spike with timestamp t 2. $G_{ref} \mapsto G_{ref} + \Delta G_{ref}$ 3. $V_m \mapsto V_{reset}$
Synapse	$$S_i = \sum_j W_{ij} s_j$$ following a spike by neuron i: $$s_i \mapsto s_i + p_R D_{fast}(1 - s_i)$$

$$D_{fast,i} \mapsto D_{fast,i}(1 - p_R)$$

Between spikes:

$$\frac{ds_i}{dt} = -\frac{s_i}{\tau_S}$$

$$\frac{dD_{fast,i}}{dt} = \frac{(D_{slow,i} - D_{fast,i})}{\tau_{fast}}$$

$$\frac{dD_{slow,i}}{dt} = \frac{(1 - D_{slow,i})}{\tau_{slow}} - f_D \frac{(D_{slow,i} - D_{fast,i})}{\tau_{fast}}$$

F Input	
Type	**Description**
All external spiking input	Input spikes increase conductance: $G_{ext} \mapsto G_{ext} + B \cdot \Delta G_{ext}$ Conductance G_{ext} decays: $$\frac{dG_{ext}}{dt} = -\frac{G_{ext}}{\tau_{ext}}$$
Background noisy input	One excitatory spike-train per neuron, and one inhibitory spike-train per neuron (all 1540 Hz Poisson spike-trains).
Stimulus	One excitatory spike-train per neuron in the E-Stay pool ("Entice" network) or the E-Leave pool ("Repel" network). In any simulated

preference test two distinct stimulus strengths were used from the following sets of three: "Entice" network stimulus strengths of increasing palatability were 94.35Hz, 377.4Hz, 660.45Hz. "Repel" network stimulus strengths of increasing palatability were 198.62Hz, 113.5Hz, 28.35Hz.

G Measurements
Active state: when mean difference between E-stay and E-leave excitatory synaptic gating exceeds .02 for 50ms (consecutively).
State duration/sampling duration: time between state transitions (i.e. transitioning from E-stay to E-leave active state).

Network states and stimuli

A network's active state was evaluated by comparing the mean values of synaptic output, s_E,

averaged across all excitatory cells in each of the two excitatory populations. Specifically, when

the difference between the mean output of the previously less active excitatory population

exceeded that of the previously more active excitatory population by a threshold of 0.02

consistently for 50ms, we recorded a state change.

We did not simulate the animal's behavior in between bouts of sampling a stimulus. Once the

excitatory neurons in the "switch" population (E-switch cells) were recorded as more active

than those in the "stay" population, using the threshold mentioned above, we removed the

stimulus input to the network. 100 ms later, we induced a subsequent transition back to the

"stay" state to represent the animal initiating a new bout of stimulus sampling. The transition

back to sampling was accomplished by halving the noisy background input to E-switch cells until

the network transitioned again to the "stay" state. At all other times in simulations, the noisy

background input remained constant. Once a transition to the "stay" state was recorded (by

excitatory cells in the "stay" population being more active than those in the "switch"

population) input stimulus was applied to indicate the next bout of sampling. The choice of

subsequent next stimulus was probabilistic, with 50% probability of each of the pair being

compared in the simulated preference test. Individual taste preference task simulations lasted

1500 seconds total. Each simulation compared sampling bout durations in response to two

stimuli each with a fixed value across the session. For a given network the stimulus inputs

targeted the same population for all sessions.

To produce linear regression coefficients in Figure 10, we regressed the log of the state

duration as a function of the stimulus strengths used, because state durations depend

exponentially on stimulus strengths in our model, which is based on noise-induced transitions

between attractor states (Kramers, 1940; Miller & Wang, 2006).

References

Abbott, L. F., Varela, J., Sen, K., & Nelson, S. (1997). Synaptic depression and cortical gain control. *Science*, *275*(5297), 221-224.
Barack, D. L., Chang, S. W. C., & Platt, M. L. (2017). Posterior Cingulate Neurons Dynamically Signal Decisions to Disengage
during Foraging. *Neuron*, *96*(2), 339-347 e335. https://doi.org/10.1016/j.neuron.2017.09.048
Bateson, M., & Kacelnik, A. (1996). Rate currencies and the foraging starling: the fallacy of the averages revisited. *Behavioral*
Ecology, *7*(3), 341-352. https://doi.org/10.1093/beheco/7.3.341
Blanchard, T. C., & Hayden, B. Y. (2014). Neurons in dorsal anterior cingulate cortex signal postdecisional variables in a foraging
task. *J Neurosci*, *34*(2), 646-655. https://doi.org/10.1523/JNEUROSCI.3151-13.2014

Blanchard, T. C., Pearson, J. M., & Hayden, B. Y. (2013). Postreward delays and systematic biases in measures of animal
temporal discounting. *Proc Natl Acad Sci U S A*, *110*(38), 15491-15496. https://doi.org/10.1073/pnas.1310446110
Charnov, E. L. (1976). Optimal foraging, the marginal value theorem. *Theor Popul Biol*, *9*(2), 129-136.
https://doi.org/10.1016/0040-5809(76)90040-x
Constantino, S. M., & Daw, N. D. (2015). Learning the opportunity cost of time in a patch-foraging task. *Cogn Affect Behav*
Neurosci, *15*(4), 837-853. https://doi.org/10.3758/s13415-015-0350-y
Davis, J. D. (1996). Deterministic and probabilistic control of the behavior of rats ingesting liquid diets. *Am J Physiol*, *270*(4 Pt 2),
R793-800. https://doi.org/10.1152/ajpregu.1996.270.4.R793
Davis, J. D., & Smith, G. P. (1992). Analysis of the microstructure of the rhythmic tongue movements of rats ingesting maltose
and sucrose solutions. *Behav Neurosci*, *106*(1), 217-228.
Fourcaud-Trocmé, N., Hansel, D., van Vreeswijk, C., & Brunel, N. (2003). How spike generation mechanisms determine the
neuronal response to fluctuating inputs. *J Neurosci*, *23*(37), 11628-11640. https://doi.org/10.1523/jneurosci.23-37-
11628.2003
Hayar, A., Bryant, J. L., Boughter, J. D., & Heck, D. H. (2006). A low-cost solution to measure mouse licking in an
electrophysiological setup with a standard analog-to-digital converter. *J Neurosci Methods*, *153*(2), 203-207.
https://doi.org/10.1016/j.jneumeth.2005.10.023
Hayden, B. Y., Pearson, J. M., & Platt, M. L. (2011). Neuronal basis of sequential foraging decisions in a patchy environment. *Nat*
Neurosci, *14*(7), 933-939. https://doi.org/10.1038/nn.2856
Kramers, H. A. (1940). Brownian motion in a field of force and the diffusion model of chemical reactions. *Physica*, *7*(4), 284-304.
https://doi.org/https://doi.org/10.1016/S0031-8914(40)90098-2
Ksander, J., Katz, D. B., & Miller, P. (2021). A model of naturalistic decision making in preference tests. *PLoS Comput Biol*, *17*(9),
e1009012. https://doi.org/10.1371/journal.pcbi.1009012
Kusick, G. F., Chin, M., Raychaudhuri, S., Lippmann, K., Adula, K. P., Hujber, E. J., Vu, T., Davis, M. W., Jorgensen, E. M., &
Watanabe, S. (2020). Synaptic vesicles transiently dock to refill release sites. *Nat Neurosci*, *23*(11), 1329-1338.
https://doi.org/10.1038/s41593-020-00716-1
Miller, P., & Wang, X.-J. (2006). Stability of discrete memory states to stochastic fluctuations in neuronal systems. *Chaos*
(Woodbury, N.Y.), *16*(2), 026109-026109. https://doi.org/10.1063/1.2208923
Pearson, J. M., Hayden, B. Y., & Platt, M. L. (2010). Explicit information reduces discounting behavior in monkeys. *Front Psychol*,
1, 237. https://doi.org/10.3389/fpsyg.2010.00237
Pearson, J. M., Watson, K. K., & Platt, M. L. (2014). Decision making: the neuroethological turn. *Neuron*, *82*(5), 950-965.
https://doi.org/10.1016/j.neuron.2014.04.037
Sadacca, B. F., Rothwax, J. T., & Katz, D. B. (2012). Sodium concentration coding gives way to evaluative coding in cortex and
amygdala. *J Neurosci*, *32*(29), 9999-10011. https://doi.org/10.1523/JNEUROSCI.6059-11.2012
Stephens, D. W. (2002). Discrimination, discounting and impulsivity: a role for an informational constraint. *Philos Trans R Soc*
Lond B Biol Sci, *357*(1427), 1527-1537. https://doi.org/10.1098/rstb.2002.1062
Stephens, D. W., & Anderson, D. (2001). The adaptive value of preference for immediacy: when shortsighted rules have
farsighted consequences. *Behavioral Ecology*, *12*(3), 330-339. https://doi.org/10.1093/beheco/12.3.330
Tabak, J., Senn, W., O'Donovan, M. J., & Rinzel, J. (2000). Modeling of spontaneous activity in developing spinal cord using
activity-dependent depression in an excitatory network. *J Neurosci*, *20*(8), 3041-3056.
https://www.ncbi.nlm.nih.gov/pubmed/10751456
Varela, J. A., Sen, K., Gibson, J., Fost, J., Abbott, L., & Nelson, S. B. (1997). A quantitative description of short-term plasticity at
excitatory synapses in layer 2/3 of rat primary visual cortex. *Journal of Neuroscience*, *17*(20), 7926-7940.
822