1

Cell Adhesion-Dependent Biphasic Axon Outgrowth Elucidated by Femtosecond Laser Impulse

Sohei Yamada[a,b,†,*], Kentarou Baba[c,†], Naoyuki Inagaki[c], Yoichiroh Hosokawa[a,d,*]

[a] Division of Materials Science, Nara Institute of Science and Technology, 8916-5 Takayama, Ikoma, Nara 630-0192, Japan

[b] Graduate School of Science and Technology, Hirosaki University, 3 Bunkyo-Cho, Hirosaki, Aomori, 036-8561, Japan

[c] Division of Biological Science, Nara Institute of Science and Technology, 8916-5 Takayama, Ikoma, Nara 630-0192, Japan

[d] Medilux Research Center, Nara Institute of Science and Technology, 8916-5 Takayama, Ikoma, Nara 630-0192, Japan

[†] These authors contributed equally

*Corresponding authors: Sohei Yamada, so-yamada@hirosaki-u.ac.jp and Yoichiroh Hosokawa, hosokawa@ms.naist.jp

Abstract

Axon outgrowth is promoted by the mechanical coupling between F-actin and adhesive
substrates via clutch and adhesion molecules in an axonal growth cone. In this study, we utilized
a femtosecond laser-induced impulse to break the coupling between the growth cone and the
substrate, enabling us to evaluate the strength of the binding between the growth cone and a
laminin on the substrate, and also determine the contribution of adhesion strength to neurite
outgrowth and traction force for the outgrowth. We found that the adhesion strength of axonal L1
cell adhesion molecule (L1CAM)-laminin binding increased with the laminin density on the
substrate. In addition, fluorescent speckle microscopy revealed that the retrograde flow of F-actin
in the growth cone was dependent on the laminin density such that the flow speed reduced with
increasing L1CAM-laminin binding. However, neurite outgrowth and the traction force did not
increase monotonically with increased L1CAM-laminin binding but rather exhibited biphasic
behavior, in which the outgrowth was suppressed by excessive L1CAM-laminin binding. Our
quantitative evaluations suggest that the biphasic outgrowth is regulated by the balance between
traction force and adhesion strength. These results imply that adhesion modulation is key to the
regulation of neurite outgrowth.

Introduction

During neuronal development, axons elongate and form functional connections with other neurons and relevant cells. The axonal migration is an essential process to regulate formation and regeneration of neuronal network. The previous studies proposed that extracellular guidance cue (chemical ligands) induces the axon outgrowth[1,2]. The growth cone located at the tip of an elongating axon senses chemical ligands in the external environment and undergoes directional migration[1-3]. In addition, growth cone senses a rich source of the extracellular matrix (ECM) to facilitate directional migration. An adhesion coupling between the growth cone and its substrate induces rapid migration and axon outgrowth.

The traction force underlying growth cone migration is regulated by modulation of the coupling efficiency between actin filament (F-actin) retrograde flow and adhesive substrates via clutch and cell adhesion molecules[4,5]. Thus, the traction force transmitted to the substrate through the F-actin-adhesion coupling promotes axon outgrowth[1,6].

We previously identified shootin1a and cortactin as clutch molecules for growth cone migration[7,8]. These molecules mediate the linkage between F-actin retrograde flow and the cytoplasmic domain of L1 cell adhesion molecule (L1CAM)[9]. The extracellular domain of L1CAM interacts with adhesive ligands such as laminins in the extracellular matrix[10-12]. L1CAM linked to the F-actin flow undergoes gripping (stop) and slipping (retrograde flow) on the substrate. It is the balance between these grip and slip states that regulates growth cone migration[12].

Our investigation focuses on the mechanism by which the interaction between L1CAM and laminin generates force. The challenge is to quantify key processes of growth cone migration, for which the adhesion strength via L1CAM-laminin binding is fundamental. However, it is difficult to quantify adhesion strength by conventional methods. For instance, the shear flow assay[13,14] is not suitable for evaluating local adhesion at the interface between a growth cone and the substrate, whereas it is difficult to apply single-cell force spectroscopy[15,16] to evaluate force without disturbing the adhesion required for axon outgrowth. The adhesion strength of migrating growth cone was previously evaluated form a detachment by micromanipulation with calibrated glass needles[17]. However, these methods are not suitable for the evaluation of adhesion strength at the growth cone level. Therefore, the relationship between adhesion strength and axon outgrowth has not been comprehensively elucidated. We thus developed a method utilizing a femtosecond laser-induced impulsive force, which we used to quantify adhesions between leukocytes and endothelial cells, among epithelial cells, and between neurons and mast cells[18,19].

A near-infrared femtosecond laser focused through an objective lens into a water solution generates stress and shock waves at the laser focal point. Vogel and his co-workers have observed these generation processes and the propagation dynamics, and confirmed that the femtosecond laser pulse generates stress waves with the smallest photon energy and confines

the propagation in the smallest area compared to other pulsed lasers[20,21]. Referring the

experimental results, we construct a simplified model to explain the action of these waves to

small particles near the laser focal point and confirmed the reliability with experiments. Based on

these results, the action is approximated that these waves propagate out spherically and act as

an impulsive force on nearby cells. As the force is localized to a diameter of 1–10 µm and breaks

intercellular adhesions at a single-cell resolution, we considered the impulse could be used to

assess adhesion of the growth cone.

We also developed an original method to quantify the magnitude of the impulsive force by

using atomic force microscopy (AFM)[22], enabling us to quantify the strength of intercellular

adhesions on the basis of the force needed to break the connection[18,19,23]. In this method, the

force is estimated from vibration of the AFM cantilever which placed instead of the cell samples.

The vibration was analyzed with taking propagations of shock and stress waves into account, and

then the total force generated at the laser focal point was estimated as a function of the laser

pulse energy. We can calibrate the force to break adhesion of the growth cone by applying this

estimation to the geometrical relation between the laser focal point and the growth cone to break

the adhesion.

In this work, we applied our previously established methods for generating laser-induced

impulsive force[7,12] to investigate the contribution of L1CAM-laminin binding to axon outgrowth.

The specific interaction between laminin and L1CAM was confirmed by L1CAM knockdown in

neurons. The strength of L1CAM-mediated adhesion was confirmed to be dependent on the

density of laminin on the substrate. In addition, we used fluorescent speckle microscopy to

observe the motions of F-actin and L1CAM in the axonal growth cone and then further assessed

the contribution of L1CAM-laminin binding to F-actin-substrate coupling. These results were

compared with axon outgrowth and traction force for the outgrowth as a dependence of laminin

density on the substrate. We propose a mechanism of neurite outgrowth that depends on the

density of laminin on the substrate, identifying L1CAM–laminin binding as a key regulator of this

process.

Results

Adhesion breaking by a femtosecond laser-induced impulsive force

Hippocampal neurons cultured for 3 days on a glass-bottom dish coated with 10 µg/ml

laminin were placed on an inverted microscope equipped with a femtosecond laser irradiation

system. The single-shot femtosecond laser pulses were focused in the vicinity of axonal growth

cones to assess the adhesion breaking threshold (Fig. 1A). The force was estimated by

measuring the distance from the growth cone at which the laser pulse broke the adhesion to the

substrate. For example, the laser with a pulse energy of 700 nJ was initially focused at a position

20 μm from a targeted growth cone. After the first pulse irradiation, the focal point was moved

closer to the target in 5 μm steps via an electrical microscope stage until adhesion was broken;

the distance between the growth cone and the final laser focal position was recorded.

111   Representative result of the adhesion breaking of the growth cone is shown in Fig. 1B

and Video S1. The growth cone moves on μm scale in the differential interference contrast (DIC)

images before and after the laser irradiation at the final laser focal point. Such displacement was

only observed at the final laser irradiation point. The distance between the growth cone and the

final point became greater with increasing the pulse energy (Fig. 2A). The detachment of the

growth cone due to force loading may result from the dissociation of laminin from the substrate.

To confirm this possibility, fluorescent dye-conjugated laminins were coated onto the grass-

bottom dish, and their fluorescent intensity was monitored following impulsive force loading. As

shown in Fig. S1, no significant change in fluorescence intensity was observed around laser focal

point. In addition, we have confirmed that the effects of cavitation and temperature are negligible

for the cell detachment by the femtosecond laser impulse. These findings suggest that the axonal

detachment results from the dissociation of L1 from laminin rather than the detachment of laminin

from the substrate. This observation indicates that growth cones can be selectively detached from

the substrate by the laser-induced impulsive force.

Quantification of the adhesion breaking force

127   We evaluated the threshold distance (R) to break growth cone adhesion to the glass

surface coated with 10 μg/ml laminin at different laser pulse energies (L) (Fig. 2A). As the

impulsive force near the growth cone increases with L, the positive correlation between R and L

indicates that R increases with an increasing impulsive force. We quantified the threshold for

breaking the adhesion by using our previously established AFM method[24] in which an AFM

cantilever replaces the tip of the growth cone and the impulsive force loaded on the cantilever is

estimated from its bending movement[18] as a function of L. From the estimation, the impulsive

force F_0 generated at the laser focal point is related to L as follows:

$F_0 = -0.003573L^2 + 0.644L - 1.5758.$ [1]

136   Because the water medium absorbs the pulse with nonlinear processes such as multi-

photon and multi-cyclic absorption, the force increases nonlinearly with the pulse energy.

Therefore, although the curve in Fig. 2A looks linear in energy, it does not mean that the

impulsive force F_0 increases linearly with R. Assuming that F_0 propagates spherically in the

vicinity of the laser focal point, the impulsive force as a unit of pressure (P) is expressed by the

following equation:

$P \text{ (kPa)} = \dfrac{F_0 \text{ (μN)} \times 10^3}{4 \pi R^2 \text{ (μm}^2)}.$ [2]

Fig. 2B is a plot of pressure to break the growth cone adhesion calculated with Eq. [2] for
each data point in Fig. 2A. The mean value of the minimum pressure to break growth cone
adhesion to a 10 µg/ml laminin substrate was 4.35 $\pm$ 1.08 kPa, comparable with the breaking
threshold reported in our previous study[23,24].

Adhesion strength of the growth cone depends on L1CAM-laminin binding

Fluorescent dye-conjugated laminin was used to assess the density of laminin on the
glass substrate. The bright spots in images in Fig. 2C are aggregates of laminin less than 1 µm in
size. The number of aggregates increased with concentration (*C*) of laminin used to coat the
glass and saturated when *C* > 50 µg/ml. The fluorescence was observed not only from the
aggregates but also from the substrate. The intensity of fluorescence (*I*) increased with *C* until
reaching saturation (Fig. 2D). This result implies that the number of laminin molecules attached to
substrate *N* is not proportional to *C* but saturates when *C* > 50 µg/ml. Here we use *I* as an index
of coverage of laminin on the substrate. The saturation of *I* is simply expressed by an exponential
plateau curve:

$I = I_{max}\left(1 - e^{-C/k}\right)$ [3]

We assumed that (i) *I* is proportional to the number of laminin molecules attached to
substrate *N*; (ii) *N* has a maximum that determines I_{max}. The constant *k* presumably reflects
attachment of laminin molecules on the substrate, which are induced by collision of the laminin
molecules to the substrate on the coating period (12 h in this experiment). In addition, we
neglected the dissociation of laminin from the substrate because *I* was not significantly different
after replacing the medium to one without laminin for the laser irradiation experiment. We defined
the coverage (*A*) of laminin as an index of laminin density on the substrate using the following
equation:

$A = 1 - e^{-C/k}$ [4]

An *A* of 1 means that the laminin attached maximumly on the substrate. The constant *k*
was estimated by least-squares fitting with Eq. [3] from the data in Fig. 2D to obtain *A* on the
substrate coated with laminin solution at concentration *C*.

The breaking threshold was evaluated according to *A* (left plots in Fig. 3A) and compared
with that determined at L1CAM-knockdown samples (right plots in Fig. 3A). The threshold with
respect to the coverage of laminin increased with *A* (Fig. 3B). On the other hand, the L1CAM-
knockdown samples maintained thresholds that matched those shown by controls at the low
coverage (*A* = 0.01). As the expression of L1CAM is almost suppressed by RNAi[12] (see also
Materials and Methods), the major adhesion of the control sample is not due to L1CAM-laminin
binding. These findings indicate that the adhesion strength between the growth cone and

substrate is strengthened by L1CAM-laminin binding. The offset threshold (~2.5 kPa) likely
reflects laminin-independent adhesive interactions.
The increase in adhesion strength with increasing A in the control samples (Fig. 3B)
suggests that the adhesion strength due to the L1CAM-laminin binding increases with the laminin
density on the substrate, despite the variability as a result of individual differences among the
cells. As the adhesion strength is integral to the individual binding strength between L1CAM and
laminin, the adhesion strength reflects the number of L1CAM-laminin interactions. Thus, the
increase in the breaking threshold with A may reflect the increase in the number of the L1CAM-
laminin interactions that occur with increased laminin density. This relationship is presumably
satisfied until the L1CAM sites available for binding are saturated.
188

L1CAM-laminin binding promotes F-actin-adhesion coupling

The contribution of the laminin coverage A to F-actin-adhesion coupling in the retrograde
flow was investigated next by visualizing F-actin and L1CAM molecules in filopodia at the growth
cone. F-actin dynamics were observed by the motion of fluorescent actin speckles tagged with
HaloTag (Fig. 4A; Video S2; Video S3; Video S4) which were observed moving along filaments
toward the leading edge of the growth cone. The speed at which they moved decreased linearly
with increasing A, slowing from 3.70 μm/min at an A of 0.06 to 2.29 μm/min at an A of 0.99 (Fig.
4B). These data suggest that A promotes the cytoskeletal-adhesion coupling.
The dynamics of L1CAM-laminin binding were evaluated as grip and slip motions of
L1CAM-HaloTag as shown in Fig. 4C (Video S5; Video S; Video S7). The ratios of grip and slip
states increased and decreased, respectively, with increasing A (Fig. 4D). Consistent with this,
the speed at which HaloTag traveled (i.e., flow speed) decreased (Fig. 4E) while the duration
spent in the grip phase increased (Fig. 4F) with increasing A. The differences between slip and
grip states were proportionate to A in the range of 0.06 to 0.99.
L1CAM-laminin binding promotes the L1CAM grip state, transmitting the traction force to
the substrate[12]. With increased cytoskeletal-adhesion coupling, F-actin flow slows and the traction
force transmitted to the substrate for growth cone migration increases[4,25]. Therefore, the
associations described above support the result from the adhesion breaking test, i.e., the number
of the L1CAM-laminin interactions, reflective of the adhesion strength of the growth cone, are
nearly proportional to A. Conversely, the dissociation of L1CAM-laminin interactions disrupts
cytoskeletal-adhesion coupling such that the force of retrograde flow is no longer transmitted to
the substrate[12]. The grip and slip motions observed in this study indicate that L1CAM-laminin
binding is not static but changes dynamically. Thus, when the average number of L1CAM-laminin
interactions between the growth cone and substrate is increased, the grip state is prolonged.
213

L1CAM-laminin binding results in biphasic axonal migration

To examine the effect of laminin coverage on growth cone migration, hippocampal

neurons were transfected with either control RNA or L1CAM miRNA, and their migration was

monitored over a 7 h in different concentrations of *A* (Fig. 5A left panels; Video S8). The migration

speed of the growth cone did not increase linearly but instead exhibited a biphasic response

dependent on *A*: within the range of 0.06 to 0.45, the migration speed exhibited a significantly

increased, whereas within the range of 0.45 to 0.99, it was markedly reduced (Fig. 5B). In

neurons with L1CAM RNAi, the migration speed significantly decreased compared to Control

RNAi. Notably, the biphasic migration pattern of the growth cone was abolished in the L1CAM

knockdown neurons (Fig. 5A, right panels; Video S8).

L1CAM-laminin binding results in biphasic axon outgrowth

Axon outgrowth also depends on the laminin coverage (*A*), as shown in Fig. 6. Axon

lengths were the longest (>130 μm) when neurons were cultured under conditions where *A* was

between 0.78 and 0.95. However, the lengths of axons from L1CAM knockdown neurons were

not affected by *A* (blue box in Fig. 5B), with shorter axons overall. These data suggest that axon

outgrowth is regulated by the L1CAM-laminin binding to the laminin substrate and that the

regulation is biphasic.

Traction force for the axon outgrowth exhibits biphasic behavior

Traction force microscopy was utilized for monitoring the force loaded on the substrate in

the axon outgrowth. In this method, the traction force is observed as displacement of the

fluoresce microbeads enclosed in the substrate and quantified from the bead shift from original

position to displaced position in the axon outgrowth (Fig. 7A). At the region of the growth cone,

the mean force for $A = 0.45$ (35 pN/nm^2) was larger than not only that for $A = 0.01$ (25 pN/nm^2)

but also that for $A = 0.99$ (25 pN/nm^2) (Fig. 7B). The dependence, which is not monophasic but

biphasic, is similar to that for axon outgrowth (Fig. 5), suggesting that the traction force is directly

related with the axon outgrowth.

Discussion

In this study, we quantified adhesion strength of axonal growth cone (Fig. 3B), axon

elongation (Fig. 5, 6), and traction force of growth cone in the elongation (Fig. 7B) as a function of

laminin density on the substrate. In addition, we evaluated molecular dynamics of F-actin and

L1CAM in the growth cone (Fig. 4). Although the adhesion strength increased with the laminin

density, the laminin dependency of the axon elongation and traction force indicated biphasic

behavior. These differences revealed in the measurements allow us to argue the mechanism of
axonal outgrowth promoted by the growth cone.

The adhesion strength measured by femtosecond laser impulses is not only for the
adhesion between L1CAM and laminin but for total amount of the adhesion including other
specific and nonspecific adhesions. The adhesion strength estimated as the adhesion breaking
force is on the order of kilopascals (3-5 kPa). On the other hand, traction force detects the force
to promote axon outgrowth. The force is generated by force transmission of F-actin retrograde
flow; transmission is effective when the binding forces of the clutch and adhesion molecules
between F-actin and substrate are higher than the traction force[12]. The force is estimated in the
range 20 – 50 pN/μm^2. Since the unit of force for traction (pN/μm^2) corresponds to pascals (Pa =
N/m^2), the adhesion strength is about 100 times larger than the traction force (0.02 – 0.05 kPa).
As shown in Fig. S1, the laser impulse does not directly break the laminin on the substrate. This
means the adhesion breaking force in Fig. 3 reflects the adhesion between the growth cone and
the substrate. The traction force in Fig. 7 is the force transmitted by the F-actin retrograde flow.
Conclusively, our results suggest that the adhesion between the growth cone and the substrate is
strong enough to transmit the force of F-actin retrograde flow to the substrate.

The L1CAM-laminin interactions promote the axon outgrowth transmitting the force of F-
actin retrograde flow to the substrate[26]. In this study, we confirmed that axon outgrowth depends
on laminin concentration, incorporating adhesion strength and traction force. When the laminin
density is low, F-actin is not coupled to the adhesive substrate through clutch molecules (e.g.,
shootin1a and cortactin) and L1CAM. As a result, the force of retrograde flow is not effectively
transmitted to the substrate to produce sufficient traction for axon outgrowth. This was
demonstrated by the short axon lengths of neurons cultured on the substrate with low laminin
density. By contrast, the rate of retrograde flow slows as the laminin density increases, indicating
greater cytoskeletal-adhesion coupling that promotes transmission of the force from the flow to
the traction force for growth cone migration. Our results indicate that laminin densities resulting in
an *A* between 0.45 and 0.78 are optimal for providing suitable traction force. Interestingly, we
observed a decrease in axon outgrowth at a high laminin density, suggesting that excessive
L1CAM-laminin binding suppresses axon outgrowth.

Similar biphasic behavior has been reported depending for concentration of guidance
molecules[17,27] and integrin-ligand binding[28-30]. Notably, neurite outgrowth exhibits a biphasic
dependence on ligand concentration, with intermediate concentrations resulting in maximal
neurite outgrowth[31.] Our results consistent with previous studies, reinforcing the validity of our
method. By integrating genetic manipulation, we extend mechanical measurement to provide the
specific contribution of L1-laminin interactions to adhesion strength. Moreover, our study extends
these findings by demonstrating that traction force generation also exhibits a biphasic

dependence with laminin concentration. This integration of force measurement and genetical
approaches helps fill the gap between classical adhesion studies and contemporary molecular
insights into growth cone migration.

Cell adhesion strength was also evaluated by shear-stress flow assay and compared with
the migration speed. With strong adhesion, the cells spread and extended lamellae, but the cell
body did not move. The suppression of migration was attributed to the inability of cells to
overcome adhesion to the substrate[30,32-34]. For axons, outgrowth is promoted not only by the
traction force at the forward side but also by detachment of the back side. Thus, the decrease in
growth we observed may be attributable to a lack of detachment as a result of excessive L1CAM-
laminin interactions which does not couple with the retrograde flow. On this assumption, the
traction force is also suppressed by the excessive interaction as well as the axon outgrowth.

Notably, axon outgrowth was suppressed when A increased from 0.78 to 0.99, an
estimated proportional increase in adhesion strength of ~20%. This result indicates that the
modification of outgrowth is on the order of 10% of the modification in the adhesion strength as a
result of the number of L1CAM-laminin interactions. Our data therefore suggest that these
interactions are responsible for adhesion to the substrate and thus for the regulation of axon
outgrowth This regulation is key for growth cone migration and axon outgrowth through the
extracellular matrix in brain, thereby contributing to the formation of network connections with
other neurons and relevant cells.

This investigation utilized femtosecond laser impulses to quantitatively evaluate the
adhesion strength between axonal growth cones and a laminin substrate. The data show relative
contribution of the L1CAM-laminin interactions in adhesion of the growth cone i.e., the strength of
the interaction increases with the laminin density. Notably, axon outgrowth does not increase
monotonically with increased L1CAM-laminin binding but instead exhibits biphasic behavior, in
which outgrowth is suppressed in the presence of high amounts of L1CAM-laminin binding. This
biphasic outgrowth is regulated by altered adhesion caused by changes in the number of binding
interactions on the order of 10%. These results suggest that the balance between the traction
force from the cytoskeletal-adhesion coupling and growth cone adhesion is one of the keys to
regulating axon outgrowth. Future studies on the biphasic regulation of axon outgrowth should
seek to further elucidate the guidance mechanism.

Materials and Methods

Preparation of cell culture substrate

For each experiment, a 35 mm glass-bottom dish (Matsunami, Osaka, Japan) was
coated with 100 µg/ml poly-lysine (FUJIFILM WAKO Pure Chemical Corporation) at 37°C for 12
h. After washing, the plate was coated with laminin (laminin 1; FUJIFILM WAKO Pure Chemical
Corporation) in phosphate-buffered saline (PBS) at 37°C for 12 h. The surfaces were washed
three times with PBS. The laminin density on the dish was modified by altering the concentration
of the deposited laminin solution (0.01–100 µg/ml). The laminin density was evaluated by coating
the dish with laminin conjugated to a fluorescent dye (green fluorescent HiLyte 488;
Cytoskeleton), which was observed under a confocal laser scanning microscope (Zeiss LSM710;
excitation, 488 nm; emission, 510 nm). The fluorescence intensity was estimated as an area
integration (15×15 µm) of the substrate fluorescence.

Cell culture

332   Hippocampal neurons were prepared from embryonic day 18 rats and seeded on glass-
333   bottom dishes. To induce axon outgrowth, neurons were cultured in neurobasal medium (Thermo
334   Fisher Scientific) containing B-27 supplement (Thermo Fisher Scientific) and 1 mM glutamine for
335   3 days. All relevant aspects of the experimental procedures were approved by the Institutional
336   Animal Care and Use Committee of Nara Institute of Science and Technology.

Femtosecond laser irradiation system

The cultures were imaged on an inverted microscope (IX71; Olympus) utilizing
femtosecond laser pulses from a regeneratively amplified Ti:Sapphire femtosecond laser (800 ± 5
nm, 100 fs, <1 mJ/pulse, 32 Hz) (Solstice Ace; Spectra-Physics). The pulse was focused near the
growth cone (Fig. 1A) through a 100× lens objective (UMPlanFl, numerical aperture [NA], 1.25;
Olympus). The irradiation was controlled by a mechanical optical shutter (Σ-65GR; Sigma Koki).
The laser pulse energy was tuned by a half-wave ($\lambda/2$) plate and dual polarizers. A single
femtosecond pulse (50–1,000 nJ/pulse) was applied near the growth cone, and adhesion
breaking was monitored by a charge-coupled-device (CCD) camera.

Impulsive force measurement system using AFM

AFM was used to quantify the force needed to break the adhesion, as described
previously[18]. An AFM cantilever (thickness, 4.0 µm; length, 125 µm; width, 30 µm; force constant,
42 N/m; resonance frequency, 330 Hz in air) (TL-NCH-10; Nano World, Neuchatel, Switzerland)

was attached to the AFM head (Nano Wizard 4 BioScience; JPK Instruments, Berlin, Germany)

and placed in pure water on the microscope stage. The laser pulse was focused 10 μm away

from the top of the cantilever. The transient oscillation of the cantilever induced by the laser pulse

irradiation was detected by an oscilloscope. The magnitude of the cantilever movement was

estimated from the oscillation.

Evaluation of laminin detachment on the substrate by impulsive force loading

The system for the laser irradiation and the condition of fluorescence laminin-coat are the

same as described above. The change in fluorescence intensity before and after laser irradiation

was analyzed in estimated shock wave generation area.

L1CAM knockdown experiment

L1CAM knockdown neurons were prepared by using a Block-iT Pol II miR RNAi

expression kit (Invitrogen). The targeting sequence of L1CAM miRNA and its effectiveness were

reported previously[12]. Hippocampal neurons were transfected with the miRNA expression vector

and incubated for 20 h. The cells were then collected and cultured on the laminin-coated glass-

bottom dishes. In this system, GFP is expressed with the L1CAM miRNA, enabling the growth

cones of transfected cells to be visualized and monitored.

Fluorescent speckle microscopy

The retrograde flow of F-actin and slip and grip motions of L1CAM were investigated by

fluorescent speckle microscopy. HaloTag-actin and L1CAM-HaloTag were expressed in

hippocampal neurons. To introduce HaloTag tetramethylrhodamine (TMR) to L1CAM-HaloTag

and HaloTag-actin, hippocampal neurons were incubated with HaloTag TMR ligand (Promega) at

a 1:1,500 dilution in L15 medium containing B27 supplement and 1 mM glutamine for 1 h at 37°C.

The medium was then replaced with fresh L15 medium. The preparation method of HaloTag-actin

is specified in the literature[12].

HaloTag-actin speckles were observed at 37°C using a fluorescence inverted microscope

(Axio Observer A1; Carl Zeiss) equipped with a C-apochromat 63× NA 1.20 lens objective (Carl

Zeiss), an illumination laser (561 nm), and an EM-CCD camera (Ixon3; Andor). Fluorescent

L1CAM-HaloTag speckles in growth cones were observed using total internal reflection

fluorescence (TIRF) on an inverted microscope (IX81; Olympus) equipped with a TIRF lens

objective (UAPON 100×OTIRF NA 1.49; Olympus), an illumination laser (488 nm), and a

scientific complementary metal-oxide semiconductor (sCMOS) camera (ORCA Flash4.0LT;

HAMAMATSU). The flow speed of F-actin and slip speed of L1CAM were analyzed by monitoring

the fluorescence signals of the HaloTags at 5 s intervals. L1CAM puncta that were visible for at

least 10 s (two intervals) were analyzed; immobile ones were defined as L1CAM in stop (grip)
phase, while those that flowed retrogradely were defined as in flow (slip) phase.

Evaluation of growth cone migration by confocal microscopy

In the neuron, EGFP is expressed to track the growth. The growth cone was
monitored by the EGEP fluorescence and DIC images. The surface of polyacrylamide gels was
coated with laminin using the same procedure as that used to coat glass-bottom dishes (see
above). EGFP and DIC signals were observed at 5 min intervals for 7 h using a fluorescence
microscope (IX81; Olympus) equipped with an EM-CCD camera (Ixon DU888; Andor), using a
20×/0.8 objective lens. Speed of growth cone migration was measured using ImageJ (Fiji
version).

Evaluation of axon length by immunofluorescence staining

Axon length was evaluated by immunofluorescence imaging. Neurons were cultured on
polyacrylamide gels coated with laminin for 3 days, fixed with 3.7% formaldehyde in PBS
overnight at 4°C, treated for 15 min with 0.05% Triton X-100 in PBS at 4°C, and then incubated
with 10% fetal bovine serum in PBS overnight at 4°C. The cells were then incubated with anti-
GFP antibody (MBL, 598) and anti-Tubulin β3 (Biolegend, 801201), as described by Toriyama et
al[25]. We used secondary antibodies conjugated with Alexa Fluor 488 (Invitrogen, A11008) or
Alexa Fluor 594 (Invitrogen A11032), and observed with a confocal laser scanning microscope
(LSM 710). The lengths of all axons of 50 cells were measured for each coverage.

Traction force microscopy

Traction force microscopy was performed as described in literatures[12,25]. Briefly, neurons
were cultured on polyacrylamide gels with embedded fluorescent microspheres (200 nm
diameter; Invitrogen). Time-lapse imaging of florescent microsphere and growth cones was
performed at 37℃ using the confocal microscope (LSM710; Carl Zeiss) with a C-Apochromat
63x/1.2 W Corr objective (Carl Zeiss). In the neuron, EGFP is expressed to track the growth. The
growth cone was monitored by the EGEP fluorescence and DIC images. The surface of
polyacrylamide gels was coated with laminin using the same procedure as that used to coat
glass-bottom dishes (see above).

Traction forces under the growth cones were detected by displacement of the beads from
their original positions. Force vectors were estimated from the displacement. To compare the
forces under different laminin densities, the magnitude of the force vectors of individual beads
under the growth cones were statistically analyzed and expressed separately as means ± SEM.
The quantification of traction force was performed using MATLAB code TFM2021 and software[35].

424
425
426 ***Statistical analysis***
427   Graphs were generated and statistical analyses were performed by using Microsoft Excel
428 and R. For nonparametric datasets, we used Mann–Whitney test to compare two groups, and
429 Kruskal–Wallis test with Dunn's multiple comparison to compare more than two groups.
430

431 **Acknowledgments**
432   We thank Dr. Ryohei Yasukuni and Dr. Kazunori Okano (Nara Institute of Science and
433 Technology, Japan) for fruitful discussions.
434

435 **Funding**
436   This research was supported in part by AMED under grant number 21gm0810011h0005,
437 ACT-X under grant number JPMJAX191K, and the Foundation of Nara Institute of Science and
438 Technology.
439

440 **Author Contributions**
441 S.Y. and Y.H. designed the research. N.I. supervised the research and contributed to
442 presentation of the mechanism of axon outgrowth. S.Y. performed almost all experiments and
443 data analysis. K.B. prepared primary cultured neurons. S.Y. and Y.H. wrote the article.

References

1.   A. L. Kolodkin, M. Tessier-Lavigne, Mechanisms and molecules of neuronal wiring: a primer. Cold Spring Harb Perspect Biol 3 (2011).

2.   L. A. Lowery, D. Van Vactor, The trip of the tip: understanding the growth cone machinery. Nat Rev Mol Cell Biol 10, 332-343 (2009).

3.   K. Franze, Integrating Chemistry and Mechanics: The Forces Driving Axon Growth. Annu Rev Cell Dev Biol 36, 61-83 (2020).

4.   D. M. Suter, P. Forscher, Substrate-cytoskeletal coupling as a mechanism for the regulation of growth cone motility and guidance. J Neurobiol 44, 97-113 (2000).

5.   N. A. Medeiros, D. T. Burnette, P. Forscher, Myosin II functions in actin-bundle turnover in neuronal growth cones. Nat Cell Biol 8, 215-226 (2006).

6.   C. E. Chan, D. J. Odde, Traction dynamics of filopodia on compliant substrates. Science 322, 1687-1691 (2008).

7.   T. Shimada et al., Shootin1 interacts with actin retrograde flow and L1-CAM to promote axon outgrowth. J Cell Biol 181, 817-829 (2008).

8.   Y. Kubo et al., Shootin1-cortactin interaction mediates signal-force transduction for axon outgrowth. J Cell Biol 210, 663-676 (2015).

9.   K. Baba et al., Gradient-reading and mechano-effector machinery for netrin-1-induced axon guidance. Elife 7 (2018).

10.   H. Hall, S. Carbonetto, M. Schachner, L1/HNK-1 carbohydrate- and beta 1 integrin-dependent neural cell adhesion to laminin-1. J Neurochem 68, 544-553 (1997).

11.   T. Esch, V. Lemmon, G. Banker, Local presentation of substrate molecules directs axon specification by cultured hippocampal neurons. Journal of Neuroscience 19, 6417-6426 (1999).

478   12.   K. Abe et al., Grip and slip of L1-CAM on adhesive substrates direct growth cone
479   haptotaxis. Proceedings of the National Academy of Sciences of the United States of America
480   115, 2764-2769 (2018).
481

482   13.   S. Gupton, C. Waterman-Storer, Spatiotemporal feedback between actomyosin and
483   focal-adhesion systems optimizes rapid cell migration. Cell 125, 1361-1374 (2006).
484

485   14.   J. de Rooij, A. Kerstens, G. Danuser, M. Schwartz, C. Waterman-Storer, Integrin-
486   dependent actomyosin contraction regulates epithelial cell scattering. Journal of Cell Biology 171,
487   153-164 (2005).
488

489   15.   M. Benoit, D. Gabriel, G. Gerisch, H. Gaub, Discrete interactions in cell adhesion
490   measured by single-molecule force spectroscopy. Nature Cell Biology 2, 313-317 (2000).
491

492   16.   M. Krieg et al., Tensile forces govern germ-layer organization in zebrafish. Nature Cell
493   Biology 10, 429-U122 (2008).
494

495   17.   J Zheng, R E Buxbaum, S R Heidemann, Measurements of growth cone adhesion to
496   culture surfaces by micromanipulation. Journal of Cell Biology 15, 2049-2060 (1994).
497

498   18.   Y. Hosokawa, Applications of the femtosecond laser-induced impulse to cell research.
499   Japanese Journal of Applied Physics 58 (2019).
500

501   19.   Y. Hosokawa, M. Hagiyama, T. Iino, Y. Murakami, A. Ito, Noncontact estimation of
502   intercellular breaking force using a femtosecond laser impulse quantified by atomic force
503   microscopy. Proceedings of the National Academy of Sciences of the United States of America
504   108, 1777-1782 (2011).
505

506   20.   A. Vogel, V. Venugopalan, Mechanisms of pulsed laser ablation of biological tissues.
507   Chemical Reviews 103, 577-644 (2003).
508

509   21.   A. Vogel, J. Noack, G. Huttman, G. Paltauf, Mechanisms of femtosecond laser
510   nanosurgery of cells and tissues. Applied Physics B-Lasers and Optics 81, 1015-1047 (2005).
511

512 22.   T. Iino, M. Hagiyama, T. Furuno, A. Ito, Y. Hosokawa, Time-Course Statistical Evaluation
513 of Intercellular Adhesion Maturation by Femtosecond Laser Impulse. Biophysical Journal 111,
514 2255-2262 (2016).
515
516 23.   K. Oikawa et al., Physical interaction between peroxisomes and chloroplasts elucidated
517 by in situ laser analysis. Nat Plants 1, 15035 (2015).
518
519 24.   S. Yamada, T. Iino, Y. Bessho, Y. Hosokawa, T. Matsui, Quantitative analysis of
520 mechanical force required for cell extrusion in zebrafish embryonic epithelia. Biology Open 6,
521 1575-1580 (2017).
522
523 25.   M. Toriyama et al., Shootin1: a protein involved in the organization of an asymmetric
524 signal for neuronal polarization. Journal of Cell Biology 175, 147-157 (2006).
525
526 26.   RH. Nichol IV, KM. Hagen, DC. Lumbard, EW. Dent, TM. Gómez, Guidance of Axons by
527 Local Coupling of Retrograde Flow to Point Contact Adhesions. Journal of Neuroscience 17,
528 2267-2282 (2016).
529
530 27.   D. Koch, WJ. Rosoff, J. Jiang, HM. Geller, JS. Urbach, Strength in the Periphery: Growth
531 Cone Biomechanics and Substrate Rigidity Response in Peripheral and Central Nervous System
532 Neurons. Biophysical Journal 102, 452-460 (2012).
533
534 28.   S. Palecek, J. Loftus, M. Ginsberg, D. Lauffenburger, A. Horwitz, Integrin-ligand binding
535 properties govern cell migration speed through cell-substratum adhesiveness. Nature 385, 537-
536 540 (1997).
537
538 29.   DiMilla PA, Barbee K, Lauffenburger DA. Mathematical model for the effects of adhesion
539 and mechanics on cell migration speed. Biophysical Journal 60, 15-37 (1991).
540
541 30.   A. Huttenlocher, M. H. Ginsberg, A. F. Horwitz, Modulation of cell migration by integrin-
542 mediated cytoskeletal linkages and ligand-binding affinity. Journal of Cell Biology 134, 1551-1562
543 (1996).
544
545 31.   JC. Schense, JA. Hubbell, Three-dimensional Migration of Neurites Is Mediated by
546 Adhesion Site Density and Affinity. Journal of Biological Chemistry 275, 6813-6818 (2000)
547

548 32. M. P. Sheetz, D. P. Felsenfeld, C. G. Galbraith, Cell migration: regulation of force on
549 extracellular-matrix-integrin complexes. Trends Cell Biol 8, 51-54 (1998).
550
551 33. T. Minegishi, N. Inagaki, Forces to Drive Neuronal Migration Steps. Front Cell Dev Biol 8,
552 863 (2020).
553
554 34. K. Hennig et al., Stick-slip dynamics of cell adhesion triggers spontaneous symmetry
555 breaking and directional migration of mesenchymal cells on one-dimensional lines. Sci Adv 6,
556 eaau5670 (2020).
557
558 35. T. Minegishi, R. Fujikawa, RF. Kastian, Y. Sakumura, N. Inagaki, Analyses of Actin
559 Dynamics, Clutch Coupling and Traction Force for Growth Cone Advance. Journal of visualized
560 experiments : JoVE 176 (2021).
561

Figures

Figure 1. Observation of adhesion breaking of an axonal growth cone by femtosecond laser-induced impulsive force. (A) Schematics of the spatial relation between the femtosecond laser pulse and targeted axonal growth cone of a neuron on a glass substrate. The laser focal point was sequentially moved closer to the growth cone, as indicated by an arrow in the upper cartoon. The effect of the impulsive force to the growth cone was indicated in the lower cartoon, which is enlarged view of the tip of the axon in the upper cartoon. (B) A representative result of adhesion breaking of a growth cone observed by a differential interference contrast (DIC) imaging. The left panel is DIC image before the laser irradiation at the final point. The middle panel indicates the laser irradiation process, in which the laser focal point (cross marks) was sequentially moved along an arrow in the image. The right panel is a merged images of the growth cones before and after the final laser irradiation, which are colorized in green and magenta, respectively. Cross marks are laser focal points. Scale bar, 10 µm.

Figure 2. Quantitative evaluation of breaking force for growth cone adhesion by using

femtosecond laser impulsive force. (A) Pulse energy L dependence of threshold distance R to

break the growth cone adhesion on a glass substrate coated with a 10 µg/ml laminin solution,

corresponding to an A of 0.45 (see Eq. [4]). $n = 36$. (B) Box-and-whisker plot of adhesion

breaking threshold. Dots in the graph is the threshold calculated independently by substituting

data from panel A into Eq. [2]. The mean value and standard deviation are 4.35 and 1.08 kPa,

respectively. (C) Images of fluorescent dye-conjugated laminin on the substrate. Concentrations

of the laminin solution used for the coating are indicated at the top. Scale bar, 10 µm. (D)

Fluorescence intensity as a function of the laminin concentration. The fluorescence intensities

were measured on substrates coated with laminin solutions with concentrations of 0.01, 0.1, 1,

10, 50, and 100 µg/ml. The fitting curve was calculated by Eq. [3], where $I_{max} = 25.5$ and $k = 16.6$.

$n = 50$ for each concentration. Data are means ± SDs from three independent experiments.

594
595

Figure 3. Adhesion breaking of growth cone on a laminin-coated substrate.

(A) Means and SDs of the adhesion breaking threshold. Dots in the graph is the threshold

calculated independently by substituting data from panel A into Eq. [2]. The laminin concentration

is indicated at the top. Red and blue dots indicate control neurons and L1CAM knockdown

neurons, respectively. (B) Box-and-whisker plots of adhesion breaking threshold. Control

neurons: n = 41, 35, 36, 12, and 25 signals for A = 0.01, 0.06, 0.45, 0.78, and 0.99, respectively.

L1CAM knockdown neurons: n = 18, 44, 34, 20, and 19 signals for A = 0.01, 0.06, 0.45, 0.78, and

0.99, respectively. **p < 0.01 (two-tailed Student's t test), n.s., not significant.

Figure 4. Molecular dynamics of F-actin and L1CAM in the axonal growth cone detected by fluorescence speckle microscopy.

(A) Fluorescence speckle images of the HaloTag-actin in a filopodium extended from an axonal growth cone. The coverage of laminin (*A*) is indicated at the top. Kymographs (right) depict HaloTag-actin behavior in boxed region in the image on the left. Slope of the yellow dashed line corresponds to retrograde flow speed of the F-actin. Time interval between frames, 5 s. Scale bar, 5 µm. (B) Retrograde flow speed of F-actin. *n* = 125, 160, and 130 signals for *A* = 0.06, 0.45, and 0.99, respectively. (C) Fluorescence speckle images of L1CAM-HaloTag in a filopodium. Kymographs (right) depict L1CAM-HaloTag behavior in a boxed region in the image on the left. Dashed pink and blue lines connect L1CAM in grip and slip states, respectively. Time interval between frames, 5 s. Scale bar, 5 µm. (D) Ratios of the grip and slip states of L1CAM-HaloTag in filopodia obtained from the kymograph analyses. *n* = 261, 450, and 197 signals for *A* = 0.06, 0.45, and 0.99, respectively. (E) Flow speed of L1CAM-HaloTag in the slip state. The speed corresponds to slopes of dashed blue lines in panel C. (F) Duration time of L1-HaloTag grip. White, red, and blue bars represent data for *A* = 0.06, 0.45, and 0.99, respectively. Data are means ± SDs; **p < 0.01.

624
Figure 5. Growth cone migration on laminin-coated substrate.

(A) Time-lapse phase-contrast/fluorescence images of hippocampal neurons in control conditions

(left panels) or L1 KD conditions (Right panels). Yellow arrowheads indicate growth cone at the

first and last time-points, respectively. See also Video – and -. Scale bar: 30 µm. (B)

Quantification of axon outgrowth velocity expressing control RNA (red) and L1-CAM miRNA

(blue) for A = 0.06, 0.45, and 0.99, respectively. Control: n = 8, 9, and 8 for A = 0.06, 0.45, 0.78,

and 0.99, respectively; L1 KD: n = 8, 8, and 8for A = 0.06, 0.45, and 0.99, respectively. Data are

means ± SDs. Kruskal-Walls test. *p < 0.05, **p < 0.01. n.s., not significant.

Figure 6. **Axon elongation on laminin-coated substrate.**

(A) Confocal images of neurons visualized with a GFP (green) and Tuj1 (Tubulin β3, magenta) antibody. Neurons were cultured on polyacrylamide gels coated with laminin. The coverage of laminin (*A*) is indicated on the left. Scale bar; 50 μm. (B) Box-and-whisker plots of axon length. Control neurons (red): n = 91, 28, 79, 60, and 71 signals for *A* = 0.01, 0.06, 0.45, 0.78, and 0.99, respectively; L1CAM knockdown neurons (blue): n = 43, 49, 73, 103, and 48 signals for *A* = 0.01, 0.06, 0.45, 0.78, and 0.99, respectively. Kruskal-Walls test. *p < 0.05, **p < 0.01. n.s., not significant.

Figure 7. Measurement of traction force on a laminin-coated substrate.

(A) DIC (left panels) and fluorescence images (right panels) of an axonal growth cone cultured on polyacrylamide gels with embedded 200 nm fluorescent beads. Dashed lines indicate the boundary of the growth cone. The blue color in the dashed line is due to EGFP which is expressed in the neuron to track the growth cone position. The original and displaced bead positions in the gel are indicated by green and red colors in fluorescence images, respectively. The yellow color overlaps the green and red colors, i.e. the bead is rarely displaced. Laminin coverage *A* is shown at the top of the panel column. Scale bar; 10 μm. (B) Magnitude of the traction forces under axonal growth cones. Data represent mean ± SEM (error bars). Kruskal-Walls test. **p < 0.01.

Figure 1 Yamada et al.

A

Initial shoot

Final shoot

Axon

Neuron

Femtosecond laser

Stress and shock wave

Growth cone

Femtosecond laser

B

Before irradiation

Irradiation process

Merge

DIC

Initial point

Final point

Figure 2 Yamada et al.

A

R : Distance (μm) vs L : Laser pulse energy (nJ/pulse)

B

Pressure (kPa) — 10 μg/ml

C

1 μg/ml   10 μg/ml   50 μg/ml   100 μg/ml

D

Laminin intensity a.u. vs Concentration (μg/ml)

Figure 3 Yamada et al.

Figure 4 Yamada et al.

Figure 5 Yamada et al.

Figure 6 Yamada et al.

Figure 7 Yamada et al.