Cell Adhesion-Dependent Biphasic Axon Outgrowth Elucidated by Femtosecond Laser Impulse

Sohei Yamada[a,b,†,*], Kentarou Baba[c,†], Naoyuki Inagaki[c], Yoichiroh Hosokawa[a,d,*]

[a] Division of Materials Science, Nara Institute of Science and Technology, 8916-5 Takayama, Ikoma, Nara 630-0192, Japan

[b] Graduate School of Science and Technology, Hirosaki University, 3 Bunkyo-Cho, Hirosaki, Aomori, 036-8561, Japan

[c] Division of Biological Science, Nara Institute of Science and Technology, 8916-5 Takayama, Ikoma, Nara 630-0192, Japan

[d] Medilux Research Center, Nara Institute of Science and Technology, 8916-5 Takayama, Ikoma, Nara 630-0192, Japan

[†] These authors contributed equally

*Corresponding authors: Sohei Yamada, so-yamada@hirosaki-u.ac.jp and Yoichiroh Hosokawa, hosokawa@ms.naist.jp

Abstract

Axon outgrowth is promoted by the mechanical coupling between F-actin and adhesive
substrates via clutch and adhesion molecules in an axonal growth cone. In this study, we utilized a
femtosecond laser-induced impulse to break the coupling between the growth cone and the
substrate, enabling us to evaluate the strength of the binding between the growth cone and a
laminin on the substrate, and also determine the contribution of adhesion strength to neurite
outgrowth and traction force for the outgrowth. We found that the adhesion strength of axonal L1
cell adhesion molecule (L1CAM)-laminin binding increased with the laminin density on the
substrate. In addition, fluorescent speckle microscopy revealed that the retrograde flow of F-actin
in the growth cone was dependent on the laminin density such that the flow speed reduced with
increasing L1CAM-laminin binding. However, neurite outgrowth and the traction force did not
increase monotonically with increased L1CAM-laminin binding but rather exhibited biphasic
behavior, in which the outgrowth was suppressed by excessive L1CAM-laminin binding. Our
quantitative evaluations suggest that the biphasic outgrowth is regulated by the balance between
traction force and adhesion strength. These results imply that adhesion modulation is key to the
regulation of neurite outgrowth.

Introduction

During neuronal development, axons elongate and form functional connections with other neurons and relevant cells. The axonal migration is an essential process to regulate formation and regeneration of neuronal network. The previous studies proposed that extracellular guidance cue (chemical ligands) induces the axon outgrowth[1,2]. The growth cone located at the tip of an elongating axon senses chemical ligands in the external environment and undergoes directional migration[1-3]. In addition, growth cone senses a rich source of the extracellular matrix (ECM) to facilitate directional migration. An adhesion coupling between the growth cone and its substrate induces rapid migration and axon outgrowth.

The traction force underlying growth cone migration is regulated by modulation of the coupling efficiency between actin filament (F-actin) retrograde flow and adhesive substrates via clutch and cell adhesion molecules[4,5]. Thus, the traction force transmitted to the substrate through the F-actin-adhesion coupling promotes axon outgrowth[1,6].

We previously identified shootin1a and cortactin as clutch molecules for growth cone migration[7,8]. These molecules mediate the linkage between F-actin retrograde flow and the cytoplasmic domain of L1 cell adhesion molecule (L1CAM)[9]. The extracellular domain of L1CAM interacts with adhesive ligands such as laminins in the extracellular matrix[10-12]. L1CAM linked to the F-actin flow undergoes gripping (stop) and slipping (retrograde flow) on the substrate. It is the balance between these grip and slip states that regulates growth cone migration[12].

Our investigation focuses on the mechanism by which the interaction between L1CAM and laminin generates force. The challenge is to quantify key processes of growth cone migration, for which the adhesion strength via L1CAM-laminin binding is fundamental. However, it is difficult to quantify adhesion strength by conventional methods. For instance, the shear flow assay[13,14] is not suitable for evaluating local adhesion at the interface between a growth cone and the substrate, whereas it is difficult to apply single-cell force spectroscopy[15,16] to evaluate force without disturbing the adhesion required for axon outgrowth. The adhesion strength of migrating growth cone was previously evaluated form a detachment by micromanipulation with calibrated glass needles[17]. However, these methods are not suitable for the evaluation of adhesion strength at the growth cone level. Therefore, the relationship between adhesion strength and axon outgrowth has not been comprehensively elucidated. We thus developed a method utilizing a femtosecond laser-induced impulsive force, which we used to quantify adhesions between leukocytes and endothelial cells, among epithelial cells, and between neurons and mast cells[18,19].

A near-infrared femtosecond laser focused through an objective lens into a water solution generates stress and shock waves at the laser focal point. Vogel and his co-workers have observed these generation processes and the propagation dynamics, and confirmed that the femtosecond laser pulse generates stress waves with the smallest photon energy and confines the propagation

in the smallest area compared to other pulsed lasers[20,21]. Referring the experimental results, we
construct a simplified model to explain the action of these waves to small particles near the laser
focal point and confirmed the reliability with experiments. Based on these results, the action is
approximated that these waves propagate out spherically and act as an impulsive force on nearby
cells. As the force is localized to a diameter of 1–10 μm and breaks intercellular adhesions at a
single-cell resolution, we considered the impulse could be used to assess adhesion of the growth
cone.

79   We also developed an original method to quantify the magnitude of the impulsive force by
using atomic force microscopy (AFM)[22], enabling us to quantify the strength of intercellular
adhesions on the basis of the force needed to break the connection[18,19,23]. In this method, the force
is estimated from vibration of the AFM cantilever which placed instead of the cell samples. The
vibration was analyzed with taking propagations of shock and stress waves into account, and then
the total force generated at the laser focal point was estimated as a function of the laser pulse
energy. We can calibrate the force to break adhesion of the growth cone by applying this estimation
to the geometrical relation between the laser focal point and the growth cone to break the adhesion.

87   In this work, we applied our previously established methods for generating laser-induced
impulsive force[7,12] to investigate the contribution of L1CAM-laminin binding to axon outgrowth. The
specific interaction between laminin and L1CAM was confirmed by L1CAM knockdown in neurons.
The strength of L1CAM-mediated adhesion was confirmed to be dependent on the density of
laminin on the substrate. In addition, we used fluorescent speckle microscopy to observe the
motions of F-actin and L1CAM in the axonal growth cone and then further assessed the contribution
of L1CAM-laminin binding to F-actin-substrate coupling. These results were compared with axon
outgrowth and traction force for the outgrowth as a dependence of laminin density on the substrate.
We propose a mechanism of neurite outgrowth that depends on the density of laminin on the
substrate, identifying L1CAM–laminin binding as a key regulator of this process.

Results

Adhesion breaking by a femtosecond laser-induced impulsive force

100   Hippocampal neurons cultured for 3 days on a glass-bottom dish coated with 10 μg/ml
laminin were placed on an inverted microscope equipped with a femtosecond laser irradiation
system. The single-shot femtosecond laser pulses were focused in the vicinity of axonal growth
cones to assess the adhesion breaking threshold (Fig. 1A). The force was estimated by measuring
the distance from the growth cone at which the laser pulse broke the adhesion to the substrate. For
example, the laser with a pulse energy of 700 nJ was initially focused at a position 20 μm from a
targeted growth cone. After the first pulse irradiation, the focal point was moved closer to the target

in 5 μm steps via an electrical microscope stage until adhesion was broken; the distance between
the growth cone and the final laser focal position was recorded.

Representative result of the adhesion breaking of the growth cone is shown in Fig. 1B and
Video S1. The growth cone moves on μm scale in the differential interference contrast (DIC) images
before and after the laser irradiation at the final laser focal point. Such displacement was only
observed at the final laser irradiation point. The distance between the growth cone and the final
point became greater with increasing the pulse energy (Fig. 2A). The detachment of the growth
cone due to force loading may result from the dissociation of laminin from the substrate. To confirm
this possibility, fluorescent dye-conjugated laminins were coated onto the grass-bottom dish, and
their fluorescent intensity was monitored following impulsive force loading. As shown in Fig. S1, no
significant change in fluorescence intensity was observed around laser focal point. In addition, we
have confirmed that the effects of cavitation and temperature are negligible for the cell detachment
by the femtosecond laser impulse. These findings suggest that the axonal detachment results from
the dissociation of L1 from laminin rather than the detachment of laminin from the substrate. This
observation indicates that growth cones can be selectively detached from the substrate by the
laser-induced impulsive force.

Quantification of the adhesion breaking force

We evaluated the threshold distance (R) to break growth cone adhesion to the glass
surface coated with 10 μg/ml laminin at different laser pulse energies (L) (Fig. 2A). As the impulsive
force near the growth cone increases with L, the positive correlation between R and L indicates
that R increases with an increasing impulsive force. We quantified the threshold for breaking the
adhesion by using our previously established AFM method[24] in which an AFM cantilever replaces
the tip of the growth cone and the impulsive force loaded on the cantilever is estimated from its
bending movement[18] as a function of L. From the estimation, the impulsive force F_0 generated at
the laser focal point is related to L as follows:

$F_0 = -0.003573L^2 + 0.644L - 1.5758.$ [1]

Because the water medium absorbs the pulse with nonlinear processes such as multi-
photon and multi-cyclic absorption, the force increases nonlinearly with the pulse energy.
Therefore, although the curve in Fig. 2A looks linear in energy, it does not mean that the impulsive
force F_0 increases linearly with R. Assuming that F_0 propagates spherically in the vicinity of the
laser focal point, the impulsive force as a unit of pressure (P) is expressed by the following equation:

$P \text{ (kPa)} = \frac{F_0 \text{ (μN)} \times 10^3}{4 \ \pi R^2 \text{ (μm}^2)}.$ [2]

Fig. 2B is a plot of pressure to break the growth cone adhesion calculated with Eq. [2] for
each data point in Fig. 2A. The mean value of the minimum pressure to break growth cone adhesion

to a 10 μg/ml laminin substrate was 4.35 $\pm$ 1.08 kPa, comparable with the breaking threshold

reported in our previous study[23,24].

Adhesion strength of the growth cone depends on L1CAM-laminin binding

Fluorescent dye-conjugated laminin was used to assess the density of laminin on the glass

substrate. The bright spots in images in Fig. 2C are aggregates of laminin less than 1 μm in size.

The number of aggregates increased with concentration (C) of laminin used to coat the glass and

saturated when $C > 50$ μg/ml. The fluorescence was observed not only from the aggregates but

also from the substrate. The intensity of fluorescence (I) increased with C until reaching saturation

(Fig. 2D). This result implies that the number of laminin molecules attached to substrate N is not

proportional to C but saturates when $C > 50$ μg/ml. Here we use I as an index of coverage of

laminin on the substrate. The saturation of I is simply expressed by an exponential plateau curve:

$I = I_{max}\left(1 - e^{-C/k}\right)$ [3]

We assumed that (i) I is proportional to the number of laminin molecules attached to

substrate N; (ii) N has a maximum that determines I_{max}. The constant k presumably reflects

attachment of laminin molecules on the substrate, which are induced by collision of the laminin

molecules to the substrate on the coating period (12 h in this experiment). In addition, we neglected

the dissociation of laminin from the substrate because I was not significantly different after replacing

the medium to one without laminin for the laser irradiation experiment. We defined the coverage

(A) of laminin as an index of laminin density on the substrate using the following equation:

$A = 1 - e^{-C/k}$ [4]

An A of 1 means that the laminin attached maximumly on the substrate. The constant k

was estimated by least-squares fitting with Eq. [3] from the data in Fig. 2D to obtain A on the

substrate coated with laminin solution at concentration C.

The breaking threshold was evaluated according to A (left plots in Fig. 3A) and compared

with that determined at L1CAM-knockdown samples (right plots in Fig. 3A). The threshold with

respect to the coverage of laminin increased with A (Fig. 3B). On the other hand, the L1CAM-

knockdown samples maintained thresholds that matched those shown by controls at the low

coverage ($A = 0.01$). As the expression of L1CAM is almost suppressed by RNAi[12] (see also

Materials and Methods), the major adhesion of the control sample is not due to L1CAM-laminin

binding. These findings indicate that the adhesion strength between the growth cone and substrate

is strengthened by L1CAM-laminin binding. The offset threshold (~2.5 kPa) likely reflects laminin-

independent adhesive interactions.

The increase in adhesion strength with increasing A in the control samples (Fig. 3B)

suggests that the adhesion strength due to the L1CAM-laminin binding increases with the laminin

density on the substrate, despite the variability as a result of individual differences among the cells.

As the adhesion strength is integral to the individual binding strength between L1CAM and laminin,

the adhesion strength reflects the number of L1CAM-laminin interactions. Thus, the increase in the

breaking threshold with A may reflect the increase in the number of the L1CAM-laminin interactions

that occur with increased laminin density. This relationship is presumably satisfied until the L1CAM

sites available for binding are saturated.

L1CAM-laminin binding promotes F-actin-adhesion coupling

The contribution of the laminin coverage A to F-actin-adhesion coupling in the retrograde

flow was investigated next by visualizing F-actin and L1CAM molecules in filopodia at the growth

cone. F-actin dynamics were observed by the motion of fluorescent actin speckles tagged with

HaloTag (Fig. 4A; Video S2; Video S3; Video S4) which were observed moving along filaments

toward the leading edge of the growth cone. The speed at which they moved decreased linearly

with increasing A, slowing from 3.70 μm/min at an A of 0.06 to 2.29 μm/min at an A of 0.99 (Fig.

4B). These data suggest that A promotes the cytoskeletal-adhesion coupling.

The dynamics of L1CAM-laminin binding were evaluated as grip and slip motions of

L1CAM-HaloTag as shown in Fig. 4C (Video S5; Video S; Video S7). The ratios of grip and slip

states increased and decreased, respectively, with increasing A (Fig. 4D). Consistent with this, the

speed at which HaloTag traveled (i.e., flow speed) decreased (Fig. 4E) while the duration spent in

the grip phase increased (Fig. 4F) with increasing A. The differences between slip and grip states

were proportionate to A in the range of 0.06 to 0.99.

L1CAM-laminin binding promotes the L1CAM grip state, transmitting the traction force to

the substrate[12]. With increased cytoskeletal-adhesion coupling, F-actin flow slows and the traction

force transmitted to the substrate for growth cone migration increases[4,25]. Therefore, the

associations described above support the result from the adhesion breaking test, i.e., the number

of the L1CAM-laminin interactions, reflective of the adhesion strength of the growth cone, are nearly

proportional to A. Conversely, the dissociation of L1CAM-laminin interactions disrupts cytoskeletal-

adhesion coupling such that the force of retrograde flow is no longer transmitted to the substrate[12].

The grip and slip motions observed in this study indicate that L1CAM-laminin binding is not static

but changes dynamically. Thus, when the average number of L1CAM-laminin interactions between

the growth cone and substrate is increased, the grip state is prolonged.

L1CAM-laminin binding results in biphasic axonal migration

To examine the effect of laminin coverage on growth cone migration, hippocampal neurons

were transfected with either control RNA or L1CAM miRNA, and their migration was monitored over

a 7 h in different concentrations of A (Fig. 5A left panels; Video S8). The migration speed of the

growth cone did not increase linearly but instead exhibited a biphasic response dependent on A:

within the range of 0.06 to 0.45, the migration speed exhibited a significantly increased, whereas
within the range of 0.45 to 0.99, it was markedly reduced (Fig. 5B). In neurons with L1CAM RNAi,
the migration speed significantly decreased compared to Control RNAi. Notably, the biphasic
migration pattern of the growth cone was abolished in the L1CAM knockdown neurons (Fig. 5A,
right panels; Video S8).
219
220 *L1CAM-laminin binding results in biphasic axon outgrowth*

221 Axon outgrowth also depends on the laminin coverage (A), as shown in Fig. 6. Axon lengths
222 were the longest (>130 μm) when neurons were cultured under conditions where A was between
223 0.78 and 0.95. However, the lengths of axons from L1CAM knockdown neurons were not affected
224 by A (blue box in Fig. 5B), with shorter axons overall. These data suggest that axon outgrowth is
225 regulated by the L1CAM-laminin binding to the laminin substrate and that the regulation is biphasic.
226
227 *Traction force for the axon outgrowth exhibits biphasic behavior*

Traction force microscopy was utilized for monitoring the force loaded on the substrate in
the axon outgrowth. In this method, the traction force is observed as displacement of the fluoresce
microbeads enclosed in the substrate and quantified from the bead shift from original position to
displaced position in the axon outgrowth (Fig. 7A). At the region of the growth cone, the mean force
for A = 0.45 (35 pN/nm^2) was larger than not only that for A = 0.01 (25 pN/nm^2) but also that for A
= 0.99 (25 pN/nm^2) (Fig. 7B). The dependence, which is not monophasic but biphasic, is similar to
that for axon outgrowth (Fig. 5), suggesting that the traction force is directly related with the axon
outgrowth.
236
237 Discussion

In this study, we quantified adhesion strength of axonal growth cone (Fig. 3B), axon
elongation (Fig. 5, 6), and traction force of growth cone in the elongation (Fig. 7B) as a function of
laminin density on the substrate. In addition, we evaluated molecular dynamics of F-actin and
L1CAM in the growth cone (Fig. 4). Although the adhesion strength increased with the laminin
density, the laminin dependency of the axon elongation and traction force indicated biphasic
behavior. These differences revealed in the measurements allow us to argue the mechanism of
axonal outgrowth promoted by the growth cone.
The adhesion strength measured by femtosecond laser impulses is not only for the
adhesion between L1CAM and laminin but for total amount of the adhesion including other specific
and nonspecific adhesions. The adhesion strength estimated as the adhesion breaking force is on
the order of kilopascals (3-5 kPa). On the other hand, traction force detects the force to promote
axon outgrowth. The force is generated by force transmission of F-actin retrograde flow;

transmission is effective when the binding forces of the clutch and adhesion molecules between F-
actin and substrate are higher than the traction force[12]. The force is estimated in the range $20 - 50$
$pN/\mu m^2$. Since the unit of force for traction ($pN/\mu m^2$) corresponds to pascals ($Pa = N/m^2$), the
adhesion strength is about 100 times larger than the traction force ($0.02 - 0.05$ kPa). As shown in
Fig. S1, the laser impulse does not directly break the laminin on the substrate. This means the
adhesion breaking force in Fig. 3 reflects the adhesion between the growth cone and the substrate.
The traction force in Fig. 7 is the force transmitted by the F-actin retrograde flow. Conclusively, our
results suggest that the adhesion between the growth cone and the substrate is strong enough to
transmit the force of F-actin retrograde flow to the substrate.

The L1CAM-laminin interactions promote the axon outgrowth transmitting the force of F-
actin retrograde flow to the substrate[26]. In this study, we confirmed that axon outgrowth depends
on laminin concentration, incorporating adhesion strength and traction force. When the laminin
density is low, F-actin is not coupled to the adhesive substrate through clutch molecules (e.g.,
shootin1a and cortactin) and L1CAM. As a result, the force of retrograde flow is not effectively
transmitted to the substrate to produce sufficient traction for axon outgrowth. This was
demonstrated by the short axon lengths of neurons cultured on the substrate with low laminin
density. By contrast, the rate of retrograde flow slows as the laminin density increases, indicating
greater cytoskeletal-adhesion coupling that promotes transmission of the force from the flow to the
traction force for growth cone migration. Our results indicate that laminin densities resulting in an
A between 0.45 and 0.78 are optimal for providing suitable traction force. Interestingly, we observed
a decrease in axon outgrowth at a high laminin density, suggesting that excessive L1CAM-laminin
binding suppresses axon outgrowth.

Similar biphasic behavior has been reported depending for concentration of guidance
molecules[17,27] and integrin-ligand binding[28-30]. Notably, neurite outgrowth exhibits a biphasic
dependence on ligand concentration, with intermediate concentrations resulting in maximal neurite
outgrowth[31.] Our results consistent with previous studies, reinforcing the validity of our method. By
integrating genetic manipulation, we extend mechanical measurement to provide the specific
contribution of L1-laminin interactions to adhesion strength. Moreover, our study extends these
findings by demonstrating that traction force generation also exhibits a biphasic dependence with
laminin concentration. This integration of force measurement and genetical approaches helps fill
the gap between classical adhesion studies and contemporary molecular insights into growth cone
migration.

Cell adhesion strength was also evaluated by shear-stress flow assay and compared with
the migration speed. With strong adhesion, the cells spread and extended lamellae, but the cell
body did not move. The suppression of migration was attributed to the inability of cells to overcome
adhesion to the substrate[30,32-34]. For axons, outgrowth is promoted not only by the traction force at

the forward side but also by detachment of the back side. Thus, the decrease in growth we
observed may be attributable to a lack of detachment as a result of excessive L1CAM-laminin
interactions which does not couple with the retrograde flow. On this assumption, the traction force
is also suppressed by the excessive interaction as well as the axon outgrowth.

Notably, axon outgrowth was suppressed when A increased from 0.78 to 0.99, an
estimated proportional increase in adhesion strength of ~20%. This result indicates that the
modification of outgrowth is on the order of 10% of the modification in the adhesion strength as a
result of the number of L1CAM-laminin interactions. Our data therefore suggest that these
interactions are responsible for adhesion to the substrate and thus for the regulation of axon
outgrowth This regulation is key for growth cone migration and axon outgrowth through the
extracellular matrix in brain, thereby contributing to the formation of network connections with other
neurons and relevant cells.

This investigation utilized femtosecond laser impulses to quantitatively evaluate the
adhesion strength between axonal growth cones and a laminin substrate. The data show relative
contribution of the L1CAM-laminin interactions in adhesion of the growth cone i.e., the strength of
the interaction increases with the laminin density. Notably, axon outgrowth does not increase
monotonically with increased L1CAM-laminin binding but instead exhibits biphasic behavior, in
which outgrowth is suppressed in the presence of high amounts of L1CAM-laminin binding. This
biphasic outgrowth is regulated by altered adhesion caused by changes in the number of binding
interactions on the order of 10%. These results suggest that the balance between the traction force
from the cytoskeletal-adhesion coupling and growth cone adhesion is one of the keys to regulating
axon outgrowth. Future studies on the biphasic regulation of axon outgrowth should seek to further
elucidate the guidance mechanism.

Materials and Methods

Preparation of cell culture substrate

For each experiment, a 35 mm glass-bottom dish (Matsunami, Osaka, Japan) was coated with 100 µg/ml poly-lysine (FUJIFILM WAKO Pure Chemical Corporation) at 37°C for 12 h. After washing, the plate was coated with laminin (laminin 1; FUJIFILM WAKO Pure Chemical Corporation) in phosphate-buffered saline (PBS) at 37°C for 12 h. The surfaces were washed three times with PBS. The laminin density on the dish was modified by altering the concentration of the deposited laminin solution (0.01–100 µg/ml). The laminin density was evaluated by coating the dish with laminin conjugated to a fluorescent dye (green fluorescent HiLyte 488; Cytoskeleton), which was observed under a confocal laser scanning microscope (Zeiss LSM710; excitation, 488 nm; emission, 510 nm). The fluorescence intensity was estimated as an area integration (15 × 15 µm) of the substrate fluorescence.

Cell culture

Hippocampal neurons were prepared from embryonic day 18 rats and seeded on glass-bottom dishes. To induce axon outgrowth, neurons were cultured in neurobasal medium (Thermo Fisher Scientific) containing B-27 supplement (Thermo Fisher Scientific) and 1 mM glutamine for 3 days. All relevant aspects of the experimental procedures were approved by the Institutional Animal Care and Use Committee of Nara Institute of Science and Technology.

Femtosecond laser irradiation system

The cultures were imaged on an inverted microscope (IX71; Olympus) utilizing femtosecond laser pulses from a regeneratively amplified Ti:Sapphire femtosecond laser (800 ± 5 nm, 100 fs, <1 mJ/pulse, 32 Hz) (Solstice Ace; Spectra-Physics). The pulse was focused near the growth cone (Fig. 1A) through a 100× lens objective (UMPlanFl, numerical aperture [NA], 1.25; Olympus). The irradiation was controlled by a mechanical optical shutter (Σ-65GR; Sigma Koki). The laser pulse energy was tuned by a half-wave (λ/2) plate and dual polarizers. A single femtosecond pulse (50–1,000 nJ/pulse) was applied near the growth cone, and adhesion breaking was monitored by a charge-coupled-device (CCD) camera.

Impulsive force measurement system using AFM

AFM was used to quantify the force needed to break the adhesion, as described previously[18]. An AFM cantilever (thickness, 4.0 µm; length, 125 µm; width, 30 µm; force constant, 42 N/m; resonance frequency, 330 Hz in air) (TL-NCH-10; Nano World, Neuchatel, Switzerland)

was attached to the AFM head (Nano Wizard 4 BioScience; JPK Instruments, Berlin, Germany)

and placed in pure water on the microscope stage. The laser pulse was focused 10 μm away from

the top of the cantilever. The transient oscillation of the cantilever induced by the laser pulse

irradiation was detected by an oscilloscope. The magnitude of the cantilever movement was

estimated from the oscillation.

Evaluation of laminin detachment on the substrate by impulsive force loading

The system for the laser irradiation and the condition of fluorescence laminin-coat are the

same as described above. The change in fluorescence intensity before and after laser irradiation

was analyzed in estimated shock wave generation area.

L1CAM knockdown experiment

L1CAM knockdown neurons were prepared by using a Block-iT Pol II miR RNAi expression

kit (Invitrogen). The targeting sequence of L1CAM miRNA and its effectiveness were reported

previously[12]. Hippocampal neurons were transfected with the miRNA expression vector and

incubated for 20 h. The cells were then collected and cultured on the laminin-coated glass-bottom

dishes. In this system, GFP is expressed with the L1CAM miRNA, enabling the growth cones of

transfected cells to be visualized and monitored.

Fluorescent speckle microscopy

The retrograde flow of F-actin and slip and grip motions of L1CAM were investigated by

fluorescent speckle microscopy. HaloTag-actin and L1CAM-HaloTag were expressed in

hippocampal neurons. To introduce HaloTag tetramethylrhodamine (TMR) to L1CAM-HaloTag and

HaloTag-actin, hippocampal neurons were incubated with HaloTag TMR ligand (Promega) at a

1:1,500 dilution in L15 medium containing B27 supplement and 1 mM glutamine for 1 h at 37°C.

The medium was then replaced with fresh L15 medium. The preparation method of HaloTag-actin

is specified in the literature[12].

HaloTag-actin speckles were observed at 37°C using a fluorescence inverted microscope

(Axio Observer A1; Carl Zeiss) equipped with a C-apochromat 63× NA 1.20 lens objective (Carl

Zeiss), an illumination laser (561 nm), and an EM-CCD camera (Ixon3; Andor). Fluorescent

L1CAM-HaloTag speckles in growth cones were observed using total internal reflection

fluorescence (TIRF) on an inverted microscope (IX81; Olympus) equipped with a TIRF lens

objective (UAPON 100×OTIRF NA 1.49; Olympus), an illumination laser (488 nm), and a scientific

complementary metal-oxide semiconductor (sCMOS) camera (ORCA Flash4.0LT; HAMAMATSU).

The flow speed of F-actin and slip speed of L1CAM were analyzed by monitoring the fluorescence

signals of the HaloTags at 5 s intervals. L1CAM puncta that were visible for at least 10 s (two

intervals) were analyzed; immobile ones were defined as L1CAM in stop (grip) phase, while those

that flowed retrogradely were defined as in flow (slip) phase.

Evaluation of growth cone migration by confocal microscopy

In the neuron, EGFP is expressed to track the growth. The growth cone was

monitored by the EGEP fluorescence and DIC images. The surface of polyacrylamide gels was

coated with laminin using the same procedure as that used to coat glass-bottom dishes (see

above). EGFP and DIC signals were observed at 5 min intervals for 7 h using a fluorescence

microscope (IX81; Olympus) equipped with an EM-CCD camera (Ixon DU888; Andor), using a

20×/0.8 objective lens. Speed of growth cone migration was measured using ImageJ (Fiji version).

Evaluation of axon length by immunofluorescence staining

Axon length was evaluated by immunofluorescence imaging. Neurons were cultured on

polyacrylamide gels coated with laminin for 3 days, fixed with 3.7% formaldehyde in PBS overnight

at 4°C, treated for 15 min with 0.05% Triton X-100 in PBS at 4°C, and then incubated with 10%

fetal bovine serum in PBS overnight at 4°C. The cells were then incubated with anti-GFP antibody

(MBL, 598) and anti-Tubulin β3 (Biolegend, 801201), as described by Toriyama et al[25]. We used

secondary antibodies conjugated with Alexa Fluor 488 (Invitrogen, A11008) or Alexa Fluor 594

(Invitrogen A11032), and observed with a confocal laser scanning microscope (LSM 710). The

lengths of all axons of 50 cells were measured for each coverage.

Traction force microscopy

Traction force microscopy was performed as described in literatures[12,25]. Briefly, neurons

were cultured on polyacrylamide gels with embedded fluorescent microspheres (200 nm diameter;

Invitrogen). Time-lapse imaging of florescent microsphere and growth cones was performed at

37℃ using the confocal microscope (LSM710; Carl Zeiss) with a C-Apochromat 63x/1.2 W Corr

objective (Carl Zeiss). In the neuron, EGFP is expressed to track the growth. The growth cone was

monitored by the EGEP fluorescence and DIC images. The surface of polyacrylamide gels was

coated with laminin using the same procedure as that used to coat glass-bottom dishes (see

above).

Traction forces under the growth cones were detected by displacement of the beads from

their original positions. Force vectors were estimated from the displacement. To compare the forces

under different laminin densities, the magnitude of the force vectors of individual beads under the

growth cones were statistically analyzed and expressed separately as means ± SEM. The

quantification of traction force was performed using MATLAB code TFM2021 and software[35].

Statistical analysis

Graphs were generated and statistical analyses were performed by using Microsoft Excel and R. For nonparametric datasets, we used Mann–Whitney test to compare two groups, and Kruskal–Wallis test with Dunn's multiple comparison to compare more than two groups.

Acknowledgments

We thank Dr. Ryohei Yasukuni and Dr. Kazunori Okano (Nara Institute of Science and Technology, Japan) for fruitful discussions.

Funding

This research was supported in part by AMED under grant number 21gm0810011h0005, ACT-X under grant number JPMJAX191K, and the Foundation of Nara Institute of Science and Technology.

Author Contributions

S.Y. and Y.H. designed the research. N.I. supervised the research and contributed to presentation of the mechanism of axon outgrowth. S.Y. performed almost all experiments and data analysis. K.B. prepared primary cultured neurons. S.Y. and Y.H. wrote the article.

References

1. A. L. Kolodkin, M. Tessier-Lavigne, Mechanisms and molecules of neuronal wiring: a

primer. Cold Spring Harb Perspect Biol 3 (2011).

2. L. A. Lowery, D. Van Vactor, The trip of the tip: understanding the growth cone machinery.

Nat Rev Mol Cell Biol 10, 332-343 (2009).

3. K. Franze, Integrating Chemistry and Mechanics: The Forces Driving Axon Growth. Annu

Rev Cell Dev Biol 36, 61-83 (2020).

4. D. M. Suter, P. Forscher, Substrate-cytoskeletal coupling as a mechanism for the

regulation of growth cone motility and guidance. J Neurobiol 44, 97-113 (2000).

5. N. A. Medeiros, D. T. Burnette, P. Forscher, Myosin II functions in actin-bundle turnover in

neuronal growth cones. Nat Cell Biol 8, 215-226 (2006).

6. C. E. Chan, D. J. Odde, Traction dynamics of filopodia on compliant substrates. Science

322, 1687-1691 (2008).

7. T. Shimada et al., Shootin1 interacts with actin retrograde flow and L1-CAM to promote

axon outgrowth. J Cell Biol 181, 817-829 (2008).

8. Y. Kubo et al., Shootin1-cortactin interaction mediates signal-force transduction for axon

outgrowth. J Cell Biol 210, 663-676 (2015).

9. K. Baba et al., Gradient-reading and mechano-effector machinery for netrin-1-induced

axon guidance. Elife 7 (2018).

10. H. Hall, S. Carbonetto, M. Schachner, L1/HNK-1 carbohydrate- and beta 1 integrin-

dependent neural cell adhesion to laminin-1. J Neurochem 68, 544-553 (1997).

11. T. Esch, V. Lemmon, G. Banker, Local presentation of substrate molecules directs axon

specification by cultured hippocampal neurons. Journal of Neuroscience 19, 6417-6426 (1999).

12.   K. Abe et al., Grip and slip of L1-CAM on adhesive substrates direct growth cone
haptotaxis. Proceedings of the National Academy of Sciences of the United States of America 115,
2764-2769 (2018).

13.   S. Gupton, C. Waterman-Storer, Spatiotemporal feedback between actomyosin and focal-
adhesion systems optimizes rapid cell migration. Cell 125, 1361-1374 (2006).

14.   J. de Rooij, A. Kerstens, G. Danuser, M. Schwartz, C. Waterman-Storer, Integrin-
dependent actomyosin contraction regulates epithelial cell scattering. Journal of Cell Biology 171,
153-164 (2005).

15.   M. Benoit, D. Gabriel, G. Gerisch, H. Gaub, Discrete interactions in cell adhesion measured
by single-molecule force spectroscopy. Nature Cell Biology 2, 313-317 (2000).

16.   M. Krieg et al., Tensile forces govern germ-layer organization in zebrafish. Nature Cell
Biology 10, 429-U122 (2008).

17.   J Zheng, R E Buxbaum, S R Heidemann, Measurements of growth cone adhesion to
culture surfaces by micromanipulation. Journal of Cell Biology 15, 2049-2060 (1994).

18.   Y. Hosokawa, Applications of the femtosecond laser-induced impulse to cell research.
Japanese Journal of Applied Physics 58 (2019).

19.   Y. Hosokawa, M. Hagiyama, T. Iino, Y. Murakami, A. Ito, Noncontact estimation of
intercellular breaking force using a femtosecond laser impulse quantified by atomic force
microscopy. Proceedings of the National Academy of Sciences of the United States of America
108, 1777-1782 (2011).

20.   A. Vogel, V. Venugopalan, Mechanisms of pulsed laser ablation of biological tissues.
Chemical Reviews 103, 577-644 (2003).

21.   A. Vogel, J. Noack, G. Huttman, G. Paltauf, Mechanisms of femtosecond laser
nanosurgery of cells and tissues. Applied Physics B-Lasers and Optics 81, 1015-1047 (2005).

2122.T. Iino, M. Hagiyama, T. Furuno, A. Ito, Y. Hosokawa, Time-Course Statistical Evaluation of
Intercellular Adhesion Maturation by Femtosecond Laser Impulse. Biophysical Journal 111, 2255-
2262 (2016).

2223.   K. Oikawa et al., Physical interaction between peroxisomes and chloroplasts elucidated by
in situ laser analysis. Nat Plants 1, 15035 (2015).

2324.   S. Yamada, T. Iino, Y. Bessho, Y. Hosokawa, T. Matsui, Quantitative analysis of
mechanical force required for cell extrusion in zebrafish embryonic epithelia. Biology Open 6, 1575-
1580 (2017).

2425.   M. Toriyama et al., Shootin1: a protein involved in the organization of an asymmetric signal
for neuronal polarization. Journal of Cell Biology 175, 147-157 (2006).

26.   RH. Nichol IV, KM. Hagen, DC. Lumbard, EW. Dent, TM. Gómez, Guidance of Axons by
Local Coupling of Retrograde Flow to Point Contact Adhesions. Journal of Neuroscience 17, 2267-
2282 (2016).

27.   D. Koch, WJ. Rosoff, J. Jiang, HM. Geller, JS. Urbach, Strength in the Periphery: Growth
Cone Biomechanics and Substrate Rigidity Response in Peripheral and Central Nervous System
Neurons. Biophysical Journal 102, 452-460 (2012).

2528.   S. Palecek, J. Loftus, M. Ginsberg, D. Lauffenburger, A. Horwitz, Integrin-ligand binding
properties govern cell migration speed through cell-substratum adhesiveness. Nature 385, 537-
540 (1997).

2629.   DiMilla PA, Barbee K, Lauffenburger DA. Mathematical model for the effects of adhesion
and mechanics on cell migration speed. Biophysical Journal 60, 15-37 (1991).

2730.   A. Huttenlocher, M. H. Ginsberg, A. F. Horwitz, Modulation of cell migration by integrin-
mediated cytoskeletal linkages and ligand-binding affinity. Journal of Cell Biology 134, 1551-1562
(1996).

31.   JC. Schense, JA. Hubbell, Three-dimensional Migration of Neurites Is Mediated by
Adhesion Site Density and Affinity. Journal of Biological Chemistry 275, 6813-6818 (2000)

~~28~~32.   M. P. Sheetz, D. P. Felsenfeld, C. G. Galbraith, Cell migration: regulation of force on
extracellular-matrix-integrin complexes. Trends Cell Biol 8, 51-54 (1998).

~~29~~33.   T. Minegishi, N. Inagaki, Forces to Drive Neuronal Migration Steps. Front Cell Dev Biol 8,
863 (2020).

~~30~~34.   K. Hennig et al., Stick-slip dynamics of cell adhesion triggers spontaneous symmetry
breaking and directional migration of mesenchymal cells on one-dimensional lines. Sci Adv 6,
eaau5670 (2020).

35.   T. Minegishi, R. Fujikawa, RF. Kastian, Y. Sakumura, N. Inagaki, Analyses of Actin
Dynamics, Clutch Coupling and Traction Force for Growth Cone Advance. Journal of visualized
experiments : JoVE 176 (2021).

Figures

A

B

Figure 1. Observation of adhesion breaking of an axonal growth cone by femtosecond laser-

induced impulsive force. (A) Schematics of the spatial relation between the femtosecond laser

pulse and targeted axonal growth cone of a neuron on a glass substrate. The laser focal point was

sequentially moved closer to the growth cone, as indicated by an arrow in the upper cartoon. The

effect of the impulsive force to the growth cone was indicated in the lower cartoon, which is enlarged

view of the tip of the axon in the upper cartoon. (B) A representative result of adhesion breaking of

a growth cone observed by a differential interference contrast (DIC) imaging. The left panel is DIC

image before the laser irradiation at the final point. The middle panel indicates the laser irradiation

process, in which the laser focal point (cross marks) was sequentially moved along an arrow in the

image. The right panel is a merged images of the growth cones before and after the final laser

irradiation, which are colorized in green and magenta, respectively. Cross marks are laser focal

points. Scale bar, 10 µm.

Figure 2. Quantitative evaluation of breaking force for growth cone adhesion by using

femtosecond laser impulsive force. (A) Pulse energy L dependence of threshold distance R to

break the growth cone adhesion on a glass substrate coated with a 10 μg/ml laminin solution,

corresponding to an A of 0.45 (see Eq. [4]). n = 36. (B) Box-and-whisker plot of adhesion breaking

threshold. Dots in the graph is the threshold calculated independently by substituting data from

panel A into Eq. [2]. The mean value and standard deviation are 4.35 and 1.08 kPa, respectively.

(C) Images of fluorescent dye-conjugated laminin on the substrate. Concentrations of the laminin

solution used for the coating are indicated at the top. Scale bar, 10 μm. (D) Fluorescence intensity

as a function of the laminin concentration. The fluorescence intensities were measured on

substrates coated with laminin solutions with concentrations of 0.01, 0.1, 1, 10, 50, and 100 μg/ml.

The fitting curve was calculated by Eq. [3], where I_{max} = 25.5 and k = 16.6. n = 50 for each

concentration. Data are means ± SDs from three independent experiments.

Figure 3. Adhesion breaking of growth cone on a laminin-coated substrate.

(A) Means and SDs of the adhesion breaking threshold. Dots in the graph is the threshold

calculated independently by substituting data from panel A into Eq. [2]. The laminin concentration

is indicated at the top. Red and blue dots indicate control neurons and L1CAM knockdown neurons,

respectively. (B) Box-and-whisker plots of adhesion breaking threshold. Control neurons: n = 41,

35, 36, 12, and 25 signals for A = 0.01, 0.06, 0.45, 0.78, and 0.99, respectively. L1CAM knockdown

neurons: n = 18, 44, 34, 20, and 19 signals for A = 0.01, 0.06, 0.45, 0.78, and 0.99, respectively.

**p < 0.01 (two-tailed Student's t test), n.s., not significant.

Figure 4. Molecular dynamics of F-actin and L1CAM in the axonal growth cone detected by

fluorescence speckle microscopy. (A) Fluorescence speckle images of the HaloTag-actin in a

filopodium extended from an axonal growth cone. The coverage of laminin (*A*) is indicated at the

top. Kymographs (right) depict HaloTag-actin behavior in boxed region in the image on the left.

Slope of the yellow dashed line corresponds to retrograde flow speed of the F-actin. Time interval

between frames, 5 s. Scale bar, 5 µm. (B) Retrograde flow speed of F-actin. *n* = 125, 160, and

130 signals for *A* = 0.06, 0.45, and 0.99, respectively. (C) Fluorescence speckle images of

L1CAM-HaloTag in a filopodium. Kymographs (right) depict L1CAM-HaloTag behavior in a boxed

region in the image on the left. Dashed pink and blue lines connect L1CAM in grip and slip states,

respectively. Time interval between frames, 5 s. Scale bar, 5 µm. (D) Ratios of the grip and slip

states of L1CAM-HaloTag in filopodia obtained from the kymograph analyses. *n* = 261, 450, and

197 signals for *A* = 0.06, 0.45, and 0.99, respectively. (E) Flow speed of L1CAM-HaloTag in the

slip state. The speed corresponds to slopes of dashed blue lines in panel C. (F) Duration time of

L1-HaloTag grip. White, red, and blue bars represent data for *A* = 0.06, 0.45, and 0.99,

respectively. Data are means ± SDs; **$p < 0.01$.

Figure 5. Growth cone migration on laminin-coated substrate.

(A) Time-lapse phase-contrast/fluorescence images of hippocampal neurons in control conditions

(left panels) or L1 KD conditions (Right panels). Yellow arrowheads indicate growth cone at the first

and last time-points, respectively. See also Video – and -. Scale bar: 30 μm. (B) Quantification of

axon outgrowth velocity expressing control RNA (red) and L1-CAM miRNA (blue) for A = 0.06, 0.45,

and 0.99, respectively. Control: n = 8, 9, and 8 for A = 0.06, 0.45, 0.78, and 0.99, respectively; L1

KD: n = 8, 8, and 8for A = 0.06, 0.45, and 0.99, respectively. Data are means ± SDs. Kruskal-Walls

test. *p < 0.05, **p < 0.01. n.s., not significant.

Figure 6. **Axon elongation on laminin-coated substrate.**

(A) Confocal images of neurons visualized with a GFP (green) and Tuj1 (Tubulin β 3, magenta) antibody. Neurons were cultured on polyacrylamide gels coated with laminin. The coverage of laminin (*A*) is indicated on the left. Scale bar; 50 μm. (B) Box-and-whisker plots of axon length. Control neurons (red): *n* = 91, 28, 79, 60, and 71 signals for *A* = 0.01, 0.06, 0.45, 0.78, and 0.99, respectively; L1CAM knockdown neurons (blue): *n* = 43, 49, 73, 103, and 48 signals for *A* = 0.01, 0.06, 0.45, 0.78, and 0.99, respectively. Kruskal-Walls test. *p < 0.05, **p < 0.01. n.s., not significant.

Figure 7. Measurement of traction force on a laminin-coated substrate.

(A) DIC (left panels) and fluorescence images (right panels) of an axonal growth cone cultured on polyacrylamide gels with embedded 200 nm fluorescent beads. Dashed lines indicate the boundary of the growth cone. The blue color in the dashed line is due to EGFP which is expressed in the neuron to track the growth cone position. The original and displaced bead positions in the gel are indicated by green and red colors in fluorescence images, respectively. The yellow color overlaps the green and red colors, i.e. the bead is rarely displaced. Laminin coverage A is shown at the top of the panel column. Scale bar; 10 μm. (B) Magnitude of the traction forces under axonal growth cones. Data represent mean ± SEM (error bars).