Somatic mutation detection: a critical evaluation through simulations and reanalyses in oaks

Sylvain Schmitt (iD), Thibault Leroy (iD), Myriam Heuertz (iD), Niklas Tysklind (iD)

CNRS, UMR EcoFoG (Agroparistech, Cirad, INRAE, Université des Antilles, Université de la Guyane), Campus Agronomique, 97310 Kourou, French Guiana;   Department of Botany and Biodiversity Research, University of Vienna, Rennweg 14, A-1030 Vienna, Austria;   IRHS-UMR1345, Université d'Angers, INRAE, Institut Agro, SFR 4207 QuaSaV, 49071 Beaucouzé, France;   Université Bordeaux, INRAE, BIOGECO, 69 route d'Arcachon, CS 80227, 33612 Cestas Cedex, France;   INRAE, UMR EcoFoG (Agroparistech, CNRS, Cirad, Université des Antilles, Université de la Guyane), Campus Agronomique, 97310 Kourou, French Guiana

- **Corresponding author**: Sylvain Schmitt, +33 6 49 19 32 63, sylvain.m.schmitt@gmail.com
- **Running title:** Somatic mutation detection in plants

Abstract

1. Mutation, the source of genetic diversity, is the raw material of evolution; however, the mutation process remains understudied, especially in plants. Using both a simulation and reanalysis framework, we set out to test the performance of two types of variant callers, generic ones and those developed for cancer research, to detect *de novo* somatic mutations.

2. In an *in silico* experiment, we generated Illumina-like sequence reads spiked with simulated mutations at different allele frequencies to compare the performance of seven commonly-used variant callers to recall them. More empirically, we then reanalyzed two of the largest datasets available for plants, both developed for identifying within-individual variation in long-lived pedunculate oaks.

3. Even in plants, variant callers developed for cancer research outperform generic callers regarding mutation recall and precision, especially at low allele frequency. Such variants at low allele frequency are typically expected for within-individual *de novo* plant mutations. Reanalysis of published oak data with the best-performing caller based on our simulations identified up to 7x more somatic mutations than initially reported.

4. Our results advocate the use of cancer research callers to boost *de novo* mutation research in plants, and to reconcile empirical reports with theoretical expectations.

Introduction

DNA sequence mutation is the raw material for evolutionary change, but, despite its crucial role, many fundamental questions around the mutation process are still open. The understanding of mutation processes is one of the most common conceptual difficulties in biology (Smith & Knight, 2012; Prevost *et al.*, 2013). Mutations are often assumed to occur at a relatively constant pace (i.e. following the hypothesis of a 'perfect' molecular clock). Despite the extremely low number of direct mutation rates estimates, mutation rates are however known to be highly variable along the tree of life, differing by several orders of magnitude among species and kingdoms, and are considered as an evolvable trait *per se* Lynch *et al.,* (2016). Mutations are assumed to be random, but the rate at which different nucleotides mutate strongly depends on the genomic context, in particular the surrounding nucleotides (Martincorena & Campbell, 2015), hereafter referred to as a mutation spectrum. The mutation spectra themselves are now believed to evolve over time (Milholland *et al.,* 2017), even at relatively short evolutionary timescales (Harris & Pritchard, 2017). The drivers of new mutations, previously thought to be simply due to DNA replication errors, are now also debated (Gao *et al.,* 2019).

Unlike most animals that transmit to the next generation only mutations present in their germ cells (*i.e.* sperm and eggs), plants are expected to produce heritable somatic mutations as they grow throughout their lives, departing from the so-called Weismann's germ plasm theory (Weismann, 1893; but see also Lanfear, 2018). As a consequence, long-lived species, such as trees, are generally assumed to accumulate more heritable mutations than herbs

per generation (Hanlon *et al.,* 2019). To generate new knowledge on plant mutation

processes[9], several studies examined within-individual variation in long-lived trees, whose

individuals can live for more than a thousand years (Schöngart *et al.,* 2017). Two studies

used the pedunculate oak (*Quercus robur*), a long-lived European tree species, as a plant

model to identify somatic mutations. Schmid-Siegert *et al.,* (2017) identified 17 mutations

by comparing sequencing data from two branches of a 234-year-old individual. The authors

therefore argued that their results are consistent with a low mutation rate in pedunculate

oak. Plomion *et al.,* (2018) identified 46 mutations using three branches of a younger

(century-old) individual, which is an almost 10-fold higher rate after taking the tree age

difference into account. Plomion *et al.,* (2018) also recovered these new mutations on acorn

embryos collected on the same branches as those used for the *de novo* mutation

identification, therefore producing empirical support for departure from Weismann's germ

plasm theory in oaks. A shared limitation of both studies is that the authors have selected a

single variant caller, without having investigated beforehand the robustness of the results

from the selected method. The absence of a simulation work to identify the best suited

detection method prior to the empirical investigations therefore represents a major limit

with regards to the accuracy and completeness of the previously reported *de novo*

mutations.

The development of tools to detect mutations in humans is rapidly expanding in cancer

research (Kim *et al.,* 2018; Alioto *et al.,* 2015). Detecting mutations in cancers is

conceptually similar to detecting somatic mutations in plants, *i.e.*, the aim is to detect

mutations that potentially affect only a small fraction of the sequenced tissue. This specific

challenge is poorly addressed in plants, where mutation detection remains based on generic

variant callers, which were initially designed to detect heterozygous sites, which have an

expected frequency of 0.5 (Schmid-Siegert *et al.,* 2017; Watson *et al.,* 2016; Hanlon *et al.,*

2019; Orr *et al.,* 2019). Generic variant callers primarily detect candidate mutations per

sample against the reference genome and validate mutation robustness by comparing

results between sample pairs, while cancer callers identify mutations by comparing two

samples, one mutated and one normal sample, against the reference genome (Fig. 1). The

per-sample strategy used in generic variant callers carries the risk of overlooking low-

frequency mutated reads in one or more samples that should invalidate the mutation in the

other sample, whereas the consideration of paired samples in cancer variant callers instead

better addresses low allelic frequency mutations in one or both samples. Transferring

mutation detection tools from cancers to plants requires evaluating their performance in a

plant research context. Cancer research frequently uses very high sequencing depths (100X

- 1000X), while the depth available for plants is often considerably lower (e.g., 34X for

Hanlon *et al.,* 2019; 40X for Wang *et al.,* 2018; or 70X for Schmid-Siegert *et al.,* 2017), bar a

few exceptions (240X for Orr *et al.,* 2019; 250X for Plomion *et al.,* 2018; or 1000X for

Watson *et al.,* 2016). To improve the detection of mutations for basic and applied plant

research, a deep evaluation of the performance of variant callers is needed in relation to the

biological features and quality of data typical of plant studies.

Here, we performed both an *in silico* and an empirical data-based evaluation of the

performance of variant callers to detect somatic mutations, using two large published

datasets on the same species (pedunculate oak, *Quercus robur*) that applied different

strategies for sequencing depth and mutation detection (Schmid-Siegert *et al.,* 2017;

Plomion *et al.,* 2018; see Fig. S1). We particularly explored the recall and precision rates

depending on the sequencing depth and allelic frequency of the somatic mutation in tissues

to answer the following questions: (1) Can cancer research methods, both in terms of

protocols (*i.e.* sequencing depth) and tools (*i.e.* callers), improve the detection of somatic

mutations?; and (2) Can reanalyses of within-individual sequencing data provide new

insights regarding plant mutation processes?

Figure 1. Generic variant callers (top rows) detect candidate mutations per tissue sample

(dark green and light green) against the reference genome (blue) and validate the robustness

of mutations by comparing results between sample pairs, while cancer callers identify

mutations by comparing two samples, one mutated (tissue A, dark green) and one normal

*(tissue B, light green), against the reference genome (blue). At low sequencing depth (**A** and*

***E**), neither the generic nor the cancer variant callers detect a low (**A**) or high (**E**) frequency*

*mutation. At intermediate sequencing depths (**B** and **F**), both generic and cancer variant*

*callers detect high-frequency mutations (**F**), but cancer variant callers are expected to be*

*better at detecting low-frequency mutations than generic variant callers (**B**), which were*

originally designed to detect the expected high-frequency heterozygous sites. At high

*sequencing depths (**C** and **G**), both the generic and cancer variant callers detect high*

*frequency (**C**) and low frequency (**G**) mutations. However, with intermediate sequencing depth*

*(**D** and **H**), a poorly represented heterozygous site in one tissue may remain undetected in that*

tissue by the generic caller while it may be detected in the second tissu and thus be considered

*a mutation, resulting in a false positive (**D**). By comparing the two samples together, cancer*

*callers will avoid this error (**H**).*

Material and methods

Study design

We developed two workflows: 1) to generate Illumina-like sequencing reads including mutations with varying biological and sequencing parameters; and 2) to detect mutations with multiple variant callers (Fig. S1). We used both *singularity* containers (Kurtzer *et al.,* 2017) and the *snakemake* workflow engines (Köster *et al.,* 2012) to build automated, highly reproducible (FAIR), and scalable workflows. We then used both workflows to test the best performing variant caller for mutation detection *in silico* based on biological and sequencing parameters. We finally used the identified variant caller to detect mutations in pedunculate oak, *Quercus robur* L., by re-analysing data from two somatic mutation projects on oaks led by INRA Bordeaux, France (Plomion *et al.,* 2018) and the University of Lausanne, Switzerland (Schmid-Siegert *et al.,* 2017).

Generation of mutations

To ensure the feasibility of the project and to limit the computational load, a first step is to subsample one or several sequences of user-defined length in the reference genome. The first workflow named *generateMutations* therefore uses a bespoke R script named *sample_genome* to generate these subsets. The workflow then takes advantage of the two scripts included in *simuG* (Yue & Liti, 2019), *vcf2model.pl,* and *simuG.pl,* respectively, 1) to build a model of heterozygous sites distribution for an haploid reference genome based on

a user-defined set of known heterozygous sites in *vcf* format and 2) to build the second

reference haploid genome comprising a user-defined number of heterozygous sites to

accurately represent diploidy. Typically, the user can define a number of heterozygous sites

based on the product of nucleotide diversity (π) and genome length (L). The workflow uses

a homemade R script named *generate_mutations* to spike randomly the reference genome

with a user-defined number of mutations which are drawn in a binomial distribution using

a user-defined transition/transversion ratio (**R**). Finally, the workflow takes advantage of

InSilicoSeq (Gourlé *et al.,* 2019) defined with the model option *hiseq* to generate datasets of

mutated and non-mutated *in silico* Illumina-like sequencing reads using (1) the original

reference haploid genome; (2) the reference haploid genome with heterozygous sites, as the

workflow was developed for a diploid species; and (3) the reference genome spiked with

mutations following user-defined allelic fraction (**AF**) and depth of sequencing depth (**C**).

154   Detection of mutations

The second workflow named *detectMutations* aims to detect somatic mutations from

mapped sequencing reads on a genome reference. Pair-end sequencing reads of every

library are quality checked using *FastQC* before trimming using *Trimmomatic* (Bolger *et al.,*

2014) keeping only paired-end reads without adaptors and a phred score above 15 in a

sliding window of 4 bases. Reads are aligned against the reference per chromosome using

BWA *mem* with the option to mark shorter splits (Li & Durbin, 2009). Alignments are then

compressed using *Samtools view* in CRAM format, sorted by coordinates using *Samtools*

sort, and indexed using *Samtools index* (Li *et al.,* 2009). Duplicated reads in alignments are

marked using *GATK MarkDuplicates* (Auwera *et al.,* 2013). Finally, the workflow uses seven

variant callers to detect mutations, including generic variant callers to detect variants and

dedicated variant callers for mutation detection. Generic variant callers to detect variants

include *GATK HaplotypeCaller* with *GATK GenotypeGVCFs* (Auwera *et al.,* 2013) and

freebayes (Garrison & Marth, 2012) using and reporting genotype qualities, without priors

on allele balance, with a minimum alternate allele fraction of 0.03, a minimum repeated

entropy of 1 and a minimum alternate allele count of 2. Cancer variant callers developed for

mutation detection include *VarScan* (Koboldt *et al.,* 2009), *Strelka2* (Kim *et al.,* 2018), *MuSE*

(Fan *et al.,* 2016), *Mutect2* (using a panel of normal and without soft clipped bases;

Benjamin *et al.,* 2019), and *Somatic Sniper* (filtering reads with mapping quality less than

25, filtering mutations with quality less than 15 with prior probability of a mutation of

0.0001; Larson *et al.,* 2012). Then we only focused on the simulated mutations, and

therefore excluded from the analyses the known heterozygous sites provided by the user

thanks to the vcf file for *GATK*, *freebayes*, *Somatic Sniper*, and *Strelka2* using *BEDTools*

subtract (Quinlan & Hall, 2010) or directly within the variant caller for *Mutect2* and

VarScan.

In silico experiment

We used the *generateMutations* workflow to generate 1000 mutations in the oak genome

with varying biological and sequencing parameters. To ensure consistency between the *in*

silico experiment and the reanalysis of empirical data, we used the reference genome

"Qrob_PM1N" of *Quercus robur* 3P from Bordeaux, ENA accession number PRJEB8388

(Plomion *et al.,* 2018), thus assessing the behaviour of variant callers in the same genomic

context as used for the empirical work. To reduce the computational load, we only

generated mutations on the first megabase of the first chromosome of the oak assembly

("Qrob_Chr01") in order to later focus the detection on this region. To check that the

conclusions regarding the callers are independent of the considered genomic region, we

ran five independent investigations based on randomly selected genome areas of a

megabase in length. Our results were highly congruent over all our investigations

(Pearson's correlations, recall: 0.999, precision: 0.947). We used known heterozygous sites

from the reference genome (Plomion *et al.,* 2018) to simulate back one thousand

heterozygous sites (π = 0.01 , L = 1 Mb , N = π x L = 10^4). We used varying values of

transition/transversion ratio (**R** = [2, 2.5, 3]), allelic fraction (**AF** = [0.05, 0.1, 0.25, 0.5]), and

sequencing depth (**C** = [25, 50, 100, 150, 200]), resulting in 60 simulated datasets of

mutated and associated base reads (3**R** x 4**AF** x 5**C**). We then used the *detectMutations*

workflow to detect (recall) spiked mutations with every variant caller (*Mutect2, freebayes,*

GATK, Strelka2, VarScan, Somatic Sniper, and *MuSe*). Using known spiked mutations, we

assessed the number of true positive (TP), false positive (FP), and false negative (FN) for

each variant caller to detect mutations and each combination of biological and sequencing

parameters. We used the resulting confusion matrix to calculate the recall ($\frac{TP}{TP+FN}$) and the

precision rates ($\frac{TP}{TP+FP}$). The recall rate represents the ability of the variant caller to detect

all mutations, while the precision rate represents the ability of the variant caller to not

confound other sites with mutations. We finally assessed each variant caller to detect

mutations using the recall and the precision rates with varying transition/transversion

ratio (**R**), allelic fraction (**AF**), and sequencing depth (**C**) to identify the best performing

variant caller based on biological and sequencing parameters.

Oak data reanalyses

We re-analyzed publicly available oak data from two projects led by Bordeaux, France

(Plomion *et al.,* 2018) and Lausanne, Switzerland (Schmid-Siegert *et al.,* 2017) (SRA

PRJNA327502 and ENA PRJEB8388, respectively). We then used the best-performing

variant caller based on our *in silico* investigation, *Strelka2*, and the variant caller for

mutation detection from the original publication to compare the results, *i.e.*, *GATK* with *Best*

Practices for Swiss data (Schmid-Siegert *et al.,* 2017) and *Mutect2* for French data (Plomion

et al., 2018). The Swiss data comprised 2 libraries of medium sequencing depth (60X)

representing one lower and one upper branch. The French data comprised 3 libraries of

high sequencing depth (160X) representing 3 branches (lower, mid, and upper). For both

Swiss and French data, we compared each pair of sample points sequentially as the

reference library and the potentially mutated library to distinguish mutations among

branches from heterozygous sites and sequencing errors. For the French data, we further

filtered out candidate somatic mutations by using a cross-validation procedure to keep a

coherent temporal pattern among mutations following the original publication (Plomion *et*

al., 2018). Contrary to a general expectation and a common view in the field (Schmid-

Siegert *et al.*, 2017, Orr *et al.*, 2019), detected mutations do not always accumulate following

the developing plant architecture (Zahradníková *et al.*, 2020). As a consequence, our cross-

validation represents a conservative strategy for the mutation detection, but it should be

noted that this strategy could have removed some true somatic mutations. We used these

raw datasets to identify the mutations from the original studies after realigning the

megabase containing the mutation on the 3P genome using *BLAT* (Kent, 2002). For both

datasets, we finally kept candidate mutations with (1) a read depth for both the normal and

mutated samples between half and two times the mean sequencing depth (30-120X and 80-

320X for Swiss and French datasets, respectively), (2) an absence of the mutated allele in

the normal sample, (3) a minimum of 10 copies of the mutated allele in the mutated sample

and (4) an allelic frequency <0.5. In addition, *Strelka2* calculates an empirical variant score

(EVS) based on a supervised random forest classifier trained on data from sequencing runs

under various conditions, which provides an overall quality score for each variant (Kim *et*

al., 2018). We took advantage of the EVS to define a conservative set of candidate mutations

for both datasets, hereafter referred to as the EVS datasets. Given that the proportion of the

genome falling within the sequencing depth boundaries used for the detection (i.e. between

50 and 200% of the mean sequencing depth) varies depending on the dataset, we weighted

the observed number of mutations by the proportion of the genome satisfying the

sequencing depth criteria to provide a more accurate and comparable estimate of the real

total number of mutations. Across both empirical studies, the proportion of the genome

with 50-200% sequencing depth varies between 71 and 87%, therefore the impact of the

weighting in the estimation of the real total number of mutations is low.

Results

To compare the performance of different variant callers to detect mutations, we simulated sequencing data containing new mutations at a given allele frequency (fraction of simulated reads per genomic position carrying the mutated allele), and using varying depths of sequencing (for variable transition/transversion ratios, see Supplementary Note S1). We then evaluated the performance of variant callers as a function of allele frequency and sequencing depth. We found marked differences in: (1) the recall, the ability to recover the simulated mutations; and (2) the precision, the proportion of truly simulated mutations among all variants detected. For allele frequencies equal to, or lower than, 0.25, cancer-specific variant callers (*Strelka2, Mutect2, MuSE,* but not *Somatic Sniper*) outperform generic variant callers such as *GATK*, *freebayes*, and *VarScan* (Fig. 2). For allele frequencies over 0.25, all variant callers perform similarly well, except for *freebayes*, which identified many false positives. Over the 80 tested parameter combinations, *Strelka2* was the best performing variant caller for various allelic frequencies and sequencing depths (in 57/80 simulated datasets , with an average recall of 0.95 for a precision of 0.98, Fig. S2-4 and Table S1 and S2) and the second fastest caller (Fig. S9).

Figure 2: Variant caller performances to identify simulated mutations for varying allelic frequencies and sequencing depths (see Fig. S5 for all parameter combinations). The recall is the ability to detect (recover) the simulated mutations. The precision is the proportion of simulated mutations among all variants detected (i.e. including false positives). Each point represents the averaged mutation recall or precision (10 simulations) for increasing allelic frequency and sequencing depth. The shaded area represents the variation of recall and precision rates over the 10 replicates computed for all callers, but only visible for the precision of Muse, Mutect2, and VarScan. Linetype opposes generic callers (dashed) against cancer variant callers developed for cancer research (solid).

We further investigated the performance of the best performing variant caller, *Strelka2*, on

two empirical datasets on pedunculate oak (Schmid-Siegert *et al.,* 2017; Plomion *et al.,*

2018) in comparison to the variant callers used in the original publications, *i.e.*, *GATK* and

Mutect2, respectively (see Supplementary Note S2). Mapping the raw data of Schmid-

Siegert *et al.,* (2017) and Plomion *et al.,* (2018) on the oak genome that we used as a

mapping reference for our empirical study, we successfully mapped 14 and 60 of the

mutations detected in the original articles, respectively. Across variant callers, we

recovered 12 (86%) and 60 (100%) of these original mutations in our total list of candidate

somatic mutations (Fig. 3A), strongly supporting the results shown by the two previous

studies. However, our analyses were able to detect far more robust candidate mutations

than initially reported. Applying filtering based on sequencing depth and mutated allele

copies (see Supplementary Note S2), *Strelka2* produced a smaller set of candidate

mutations than *GATK* but similar to *Mutect2*, with an estimated number of mutation

candidates 10- to 25-fold higher than that of the original studies (Fig. 3A). Adding *Strelka2*

recommended filtering based on empirical variant scores yielded the most conservative

dataset with a similar number of mutations between both studies and a 2 to 7-fold increase

compared to the original number of mutations. Due to lack of access to biological material

from the original studies, conclusions were drawn from this list of conservative candidate

somatic mutations (but see supplementary note S4 for a discussion regarding validation of

mutations). The distribution of allelic frequencies of detected mutations partly explains

differences among detection methods (Fig. 3A), with *Strelka2* and *Mutect2* detecting

mutations with lower allelic frequencies than the candidate mutations presented in the

original publications, especially for the Plomion *et al.,* (2018) study that used higher

sequencing depths.

Based on the set of conservative mutations detected by *Strelka2* (EVS), we then explored

annotations and mutation spectra in both datasets (Fig. 3B-C), which have rarely been

explored in model plant species (but see first evidence based on mutation accumulation

lines in Arabidopsis in Weng *et al.,* 2019) and never in the wild. The proportions of

mutations found in different genomic regions (e.g. genic, intergenic) were highly correlated

between both original studies and proportional to the representation of the genomic

regions, supporting a random distribution of mutations throughout the genome (Fig. 3B).

Mutation spectra of the two studies are significantly correlated (Pearson'r=0.49, p<7.4 10-

5), with an enrichment in C>T transitions, particularly in some specific genomic contexts

(Fig. 3C).

Figure 3: Candidate mutation spectra depending on variant callers and filtering in Schmid-Siegert et al., (2017) and Plomion et al., (2018). A. Allelic frequency distribution for every dataset, including the candidate mutations from the original article present in the raw data from the reanalysis (red), the results of GATK with Best Practices (blue), Mutect2 after filtering (green), and Strelka2 after filtering (purple), and the results of Strelka2 using the filtering based on empirical variant scores named EVS (orange). The labels indicate the number of candidate mutations in each dataset. Per caller comparisons are available in Fig. S7. B. Annotation of the mutations detected by Strelka2 across chromosomes using the filtering based on empirical variant scores named EVS for Schmid-Siegert et al., (2017, green) and Plomion et al., (2018, orange) compared to the genomic expectation (grey, see Supplementary Note S3). Error bars represent the standard deviation (SD) of the observed

percentages across chromosomes, and the annotation above the columns indicates the

significance of the Student's t-test two-sided comparing the mean percentage of mutations to

the mean genomic expectation, with ns, **, and *** corresponding to non-significant, p<0.01,

and p<0.001 differences, respectively. **C.** *Context-dependent mutation spectra depending on*

mutation types for the results of Strelka2 using the filtering based on empirical variant scores

named EVS. Mutation types have been summarised into six main classes with thicker lines for

transversion compared to transition, and then differentiated depending on their 5' and 3'

genomic contexts, see Fig. S8-9. Pearson's correlation r measures the two-sided correlation of

the mutation spectra between Schmid-Siegert et al., (2017) and Plomion et al., (2018).

Discussion

Mutation research in plants still primarily uses generic variant callers and methodologies

that are not developed for the specificity and complexity of within-individual *de novo*

mutation detection. We examined if plant mutation research could benefit from the

development of tools and protocols initially designed for human cancer research, which is a

rapidly expanding field (Kim *et al.,* 2018). We found marked differences in the performance

of variant callers for mutation detection based on sequencing depth and allelic frequency.

We found that cancer variant callers performed better than generic variant callers for

mutation detection at low or intermediate allelic frequency or with low sequencing depth,

and similarly well for high allelic frequency. Low allelic frequency mutations, potentially

due to the chimeric nature of plant shoot apical meristems structures (Burian, 2021), might

be very important due to their great abundance that may balance out their low chance of

transmissions. Therefore, plant mutation studies should make greater use of cancer variant

callers such as *Strelka2* rather than generic variant callers such as *GATK* to detect somatic

mutations, in agreement with previous studies on germline mutations detection (Chen *et al.,*

2019), especially for detecting low frequency mutations and when using low sequencing

depth. The importance of allele frequency-dependency in variant detection is not restricted

to somatic mutations, but also concerns for instance polyploid species, which includes many

agriculturally important autopolyploid plant species (e.g. potato, sugarcane). Our

simulation framework therefore provides general insights regarding the impact of allelic

dosage in mutation detection which go beyond somatic mutation detection.

One problem that may arise when analysing sample pairs with cancer variant callers

is the rapid increase in pairwise comparisons when using a larger sample size than

previous studies (e.g., N=3 in Plomion *et al.* 2018). A simple solution is the use of a single

reference sample such as a cambium sample from the base of the tree, which is therefore

considered as the closest genome to the seed, to compare it to all samples from branches

(Hanlon *et al.,* 2019). By reanalyzing the raw oak data (Schmid-Siegert *et al.,* 2017; Plomion

et al., 2018), we found that the marked differences in the performance of variant callers

could account for the discrepancies in genome-wide plant somatic mutation rate estimates.

Our reanalysis shows robust evidence for an up to 7-fold higher number of mutations than

previously reported, a value closer to the expectations based on the theory (Schoen &

Schultz, 2019; Burian, 2021). We argue that knowledge and methodological transfers from

cancer to plant mutation detection are expected to contribute strongly to the upward trend

of this field and to reconcile empirical reports with theoretical expectations.

Acknowledgments

The manuscript benefited from the comments of three anonymous reviewers.

Funding

This study was funded through an Investissement d'Avenir grant of the ANR: CEBA (ANR-10-LABEX-0025).

Conflict of interest disclosure

The authors declare they have no conflict of interest relating to the content of this article.

MH is a recommender for PCI Evol Biol.

Data, script and code availability

Reanalyzed reads and corresponding genomes were extracted from GenBank under accession BioProject PRJNA327502 and from European Nucleotide Archive under project accession code PRJEB8388. generateMutations and detectMutations pipelines are available on GitHub (https://github.com/sylvainschmitt/generateMutations and https://github.com/sylvainschmitt/detectMutations).

Authors' contributions

All authors conceived the ideas; SS developed the pipelines, conducted the virtual

experiment and the data reanalyses; SS analysed outputs and led the writing of the

manuscript. All authors contributed critically to the drafts and gave final approval for

publication.

References

Alioto, T.S., Buchhalter, I., Derdak, S., Hutter, B., Eldridge, M.D., Hovig, E., Heisler, L.E., Beck,

T.A., Simpson, J.T., Tonon, L., Sertier, A.S., Patch, A.M., Jäger, N., Ginsbach, P., Drews, R.,

Paramasivam, N., Kabbe, R., Chotewutmontri, S., Diessl, N., Previti, C., Schmidt, S., Brors, B.,

Feuerbach, L., Heinold, M., Gröbner, S., Korshunov, A., Tarpey, P.S., Butler, A.P., Hinton, J.,

Jones, D., Menzies, A., Raine, K., Shepherd, R., Stebbings, L., Teague, J.W., Ribeca, P., Giner,

F.C., Beltran, S., Raineri, E., Dabad, M., Heath, S.C., Gut, M., Denroche, R.E., Harding, N.J.,

Yamaguchi, T.N., Fujimoto, A., Nakagawa, H., Quesada, V., Valdés-Mas, R., Nakken, S., Vodák,

D., Bower, L., Lynch, A.G., Anderson, C.L., Waddell, N., Pearson, J.V., Grimmond, S.M., Peto, M.,

Spellman, P., He, M., Kandoth, C., Lee, S., Zhang, J., Létourneau, L., Ma, S., Seth, S., Torrents, D.,

Xi, L., Wheeler, D.A., López-Otín, C., Campo, E., Campbell, P.J., Boutros, P.C., Puente, X.S.,

Gerhard, D.S., Pfister, S.M., McPherson, J.D., Hudson, T.J., Schlesner, M., Lichter, P., Eils, R.,

Jones, D.T.W. & Gut, I.G. (2015). A comprehensive assessment of somatic mutation detection

in cancer using whole-genome sequencing. *Nature Communications*, **6**.

Auwera, G.A., Carneiro, M.O., Hartl, C., Poplin, R., Angel, G. del, Levy-Moonshine, A., Jordan,

T., Shakir, K., Roazen, D., Thibault, J., Banks, E., Garimella, K.V., Altshuler, D., Gabriel, S. &

DePristo, M.A. (2013). From FastQ Data to High-Confidence Variant Calls: The Genome

Analysis Toolkit Best Practices Pipeline. *Current Protocols in Bioinformatics*, **43**, 483–492.

Retrieved from https://onlinelibrary.wiley.com/doi/abs/10.1002/0471250953.bi1110s43

Benjamin, D., Sato, T., Cibulskis, K., Getz, G., Stewart, C. & Lichtenstein, L. (2019). Calling

Somatic SNVs and Indels with Mutect2.

Bolger, A.M., Lohse, M. & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina

sequence data. *Bioinformatics*, **30**, 2114–2120.

Burian, A. (2021). Does Shoot Apical Meristem Function as the Germline in Safeguarding

Against Excess of Mutations? *Frontiers in Plant Science*, **12**, 1–9.

Chen, Z.L., Meng, J.M., Cao, Y., Yin, J.L., Fang, R.Q., Fan, S.B., Liu, C., Zeng, W.F., Ding, Y.H., Tan,

D., Wu, L., Zhou, W.J., Chi, H., Sun, R.X., Dong, M.Q. & He, S.M. (2019). A high-speed search

engine pLink 2 with systematic evaluation for proteome-scale identification of cross-linked

peptides. *Nature Communications*, **10**. Retrieved from http://dx.doi.org/10.1038/s41467-

019-11337-z

Fan, Y., Xi, L., Hughes, D.S.T., Zhang, J., Zhang, J., Futreal, P.A., Wheeler, D.A. & Wang, W.

(2016). MuSE: accounting for tumor heterogeneity using a sample-specific error model

improves sensitivity and specificity in mutation calling from sequencing data. *Genome*

biology, **17**, 178. Retrieved from http://dx.doi.org/10.1186/s13059-016-1029-6

Gao, Z., Moorjani, P., Sasani, T.A., Pedersen, B.S., Quinlan, A.R., Jorde, L.B., Amster, G. &

Przeworski, M. (2019). Overlooked roles of DNA damage and maternal age in generating

human germline mutations. *Proceedings of the National Academy of Sciences of the United*

States of America, **116**, 9491–9500.

Garrison, E. & Marth, G. (2012). Haplotype-based variant detection from short-read

sequencing. 1–9. Retrieved from http://arxiv.org/abs/1207.3907

Gourlé, H., Karlsson-Lindsjö, O., Hayer, J. & Bongcam-Rudloff, E. (2019). Simulating Illumina

metagenomic data with InSilicoSeq. *Bioinformatics*, **35**, 521–522.

Hanlon, V.C.T., Otto, S.P. & Aitken, S.N. (2019). Somatic mutations substantially increase the

per-generation mutation rate in the conifer Picea sitchensis. *Evolution Letters*, **3**, 348–358.

Harris, K. & Pritchard, J.K. (2017). Rapid evolution of the human mutation spectrum. *eLife*,

6, 1–17.

Kent, W.J. (2002). BLAT—The BLAST-Like Alignment Tool. *Genome Research*, **12**, 656–664.

Kim, S., Scheffler, K., Halpern, A.L., Bekritsky, M.A., Noh, E., Källberg, M., Chen, X., Kim, Y.,

Beyter, D., Krusche, P. & Saunders, C.T. (2018). Strelka2: fast and accurate calling of

germline and somatic variants. *Nature Methods*, **15**, 591–594.

Koboldt, D.C., Chen, K., Wylie, T., Larson, D.E., McLellan, M.D., Mardis, E.R., Weinstock, G.M.,

Wilson, R.K. & Ding, L. (2009). VarScan: Variant detection in massively parallel sequencing

of individual and pooled samples. *Bioinformatics*, **25**, 2283–2285.

Köster, J. & Rahmann, S. (2012). Snakemake-a scalable bioinformatics workflow engine.

Bioinformatics, **28**, 2520–2522.

Kurtzer, G.M., Sochat, V. & Bauer, M.W. (2017). Singularity: Scientific containers for mobility

of compute. *PLoS ONE*, **12**, 1–20.

Lanfear, R. (2018). Do plants have a segregated germline? *PLoS Biology*, **16**, 1–13.

Larson, D.E., Harris, C.C., Chen, K., Koboldt, D.C., Abbott, T.E., Dooling, D.J., Ley, T.J., Mardis,

E.R., Wilson, R.K. & Ding, L. (2012). Somaticsniper: Identification of somatic point mutations

in whole genome sequencing data. *Bioinformatics*, **28**, 311–317.

Li, H. & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics*, **25**, 1754–1760.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G. &

Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, **25**,

2078–2079.

Lynch, M., Ackerman, M.S., Gout, J.F., Long, H., Sung, W., Thomas, W.K. & Foster, P.L. (2016).

Genetic drift, selection and the evolution of the mutation rate. *Nature Reviews Genetics*, **17**,

704–714. Retrieved from http://dx.doi.org/10.1038/nrg.2016.104

Martincorena, I. & Campbell, P.J. (2015). Somatic mutation in cancer and normal cells.

Science, **349**, 1483–1489.

Milholland, B., Dong, X., Zhang, L., Hao, X., Suh, Y. & Vijg, J. (2017). Differences between

germline and somatic mutation rates in humans and mice. *Nature Communications*, **8**, 1–8.

Retrieved from http://dx.doi.org/10.1038/ncomms15183

Orr, A.J., Padovan, A., Kainer, D., Külheim, C., Bromham, L., Bustos-Segura, C., Foley, W., Haff,

T., Hsieh, J.F., Morales-Suarez, A., Cartwright, R.A. & Lanfear, R. (2019). A phylogenomic

approach reveals a low somatic mutation rate in a long-lived plant. *bioRxiv*.

Plomion, C., Aury, J.M., Amselem, J., Leroy, T., Murat, F., Duplessis, S., Faye, S., Francillonne,

N., Labadie, K., Le Provost, G., Lesur, I., Bartholomé, J., Faivre-Rampant, P., Kohler, A., Leplé,

J.C., Chantret, N., Chen, J., Diévart, A., Alaeitabar, T., Barbe, V., Belser, C., Bergès, H., Bodénès,

C., Bogeat-Triboulot, M.B., Bouffaud, M.L., Brachi, B., Chancerel, E., Cohen, D., Couloux, A., Da

Silva, C., Dossat, C., Ehrenmann, F., Gaspin, C., Grima-Pettenati, J., Guichoux, E., Hecker, A.,

Herrmann, S., Hugueney, P., Hummel, I., Klopp, C., Lalanne, C., Lascoux, M., Lasserre, E.,

Lemainque, A., Desprez-Loustau, M.L., Luyten, I., Madoui, M.A., Mangenot, S., Marchal, C.,

Maumus, F., Mercier, J., Michotey, C., Panaud, O., Picault, N., Rouhier, N., Rué, O., Rustenholz,

C., Salin, F., Soler, M., Tarkka, M., Velt, A., Zanne, A.E., Martin, F., Wincker, P., Quesneville, H.,

Kremer, A. & Salse, J. (2018). Oak genome reveals facets of long lifespan. *Nature Plants*, **4**,

440–452. Retrieved from http://dx.doi.org/10.1038/s41477-018-0172-3

Prevost, L., Knight, J., Smith, M., & Lurain, U. M. (2013). Student writing reveals their

heterogeneous thinking about the origin of genetic variation in populations.

Quinlan, A.R. & Hall, I.M. (2010). BEDTools: A flexible suite of utilities for comparing

genomic features. *Bioinformatics*, **26**, 841–842.

Schmid-Siegert, E., Sarkar, N., Iseli, C., Calderon, S., Gouhier-Darimont, C., Chrast, J., Cattaneo,

P., Schütz, F., Farinelli, L., Pagni, M., Schneider, M., Voumard, J., Jaboyedoff, M., Fankhauser,

C., Hardtke, C.S., Keller, L., Pannell, J.R., Reymond, A., Robinson-Rechavi, M., Xenarios, I. &

Reymond, P. (2017). Low number of fixed somatic mutations in a long-lived oak tree.

Nature Plants, **3**, 926–929. Retrieved from http://dx.doi.org/10.1038/s41477-017-0066-9

Schoen, D.J. & Schultz, S.T. (2019). Somatic Mutation and Evolution in Plants. *Annual Review

of Ecology, Evolution, and Systematics*, **50**, 49–73.

Schöngart, J., Bräuning, A., Barbosa, A.C.M.C., Lisi, C.S. & Oliveira, J.M. de. (2017).

Dendroecology Tree-Ring Analyses Applied to Ecological Studies.

Smith, M. K., & Knight, J. K. (2012). Using the Genetics Concept Assessment to document

persistent conceptual difficulties in undergraduate genetics courses. Genetics, 191(1), 21–

32. https://doi.org/10.1534/genetics.111.137810

Wang, L., Ji, Y., Hu, Y., Hu, H., Jia, X., Jiang, M., Zhang, X., Zhao, L., Zhang, Y., Jia, Y., Qin, C., Yu,

L., Huang, J., Yang, S., Hurst, L.D. & Tian, D. (2019). The architecture of intra-organism

mutation rate variation in plants. *PLoS Biology*, **17**, 1–29.

Watson, J.M., Platzer, A., Kazda, A., Akimcheva, S., Valuchova, S., Nizhynska, V., Nordborg, M.

& Riha, K. (2016). Germline replications and somatic mutation accumulation are

independent of vegetative life span in Arabidopsis. *Proceedings of the National Academy of*

Sciences of the United States of America, **113**, 12226–12231.

Weismann, A. (1893). *The germ-plasm: a theory of heredity*. Scribner's.

Weng, M.L., Becker, C., Hildebrandt, J., Neumann, M., Rutter, M.T., Shaw, R.G., Weigel, D. &

Fenster, C.B. (2019). Fine-grained analysis of spontaneous mutation spectrum and

frequency in arabidopsis thaliana. *Genetics*, **211**, 703–714.

Yue, J.X. & Liti, G. (2019). SimuG: A general-purpose genome simulator. *Bioinformatics*, **35**,

4442–4444.

Zahradníková, E., Ficek, A., Brejová, B., Vinař, T., & Mičieta, K. (2020). Mosaicism in old trees

and its patterns. *Trees - Structure and Function,* **34**, 357–370.

https://doi.org/10.1007/s00468-019-01921-7

Supplementary material

The following Supporting Information is available for this article:

Note S1. Simulation results

Note S2. Reanalyses results

Note S3. Genomic expectations for the annotation of the mutations

Note S4. 'validation' of mutations

Fig. S1. Study scheme

Fig. S2. Variation in the performance of variant callers for mutation detection with varying
biological and sequencing parameters

Fig. S3. Variation in the performance of variant callers for mutation detection with varying
biological and sequencing parameters

Fig. S4. Best performing variant callers for mutation detection depending on allelic fraction
(allelic frequency) and coverage (sequencing depth)

Fig. S5. Mutation recall and precision rates for generic and mutation-specific variant callers
by allelic fraction and sequencing depth

Fig. S6. Observed allelic frequencies of candidate mutations depending on variant callers
and filtering in Schmid-Siegert *et al.,* (2017) and Plomion *et al.,* (2018)

Fig. S7. Percentage of nucleotide change types of candidate mutations depending on variant

callers and filtering in Schmid-Siegert *et al.,* (2017) and Plomion *et al.,* (2018)

Fig. S8. Context-dependent mutation spectrum depending on variant callers and filtering in

Schmid-Siegert *et al.,* (2017) and Plomion *et al.,* (2018)

Tab. S1. Mean and standard deviation in performance of variant callers for mutation

detection across all simulations

Tab S2. Mean and standard deviation in performance of variant callers for mutation

detection with varying allelic frequency and sequencing depth

526