Somatic mutation detection: a critical evaluation through simulations and reanalyses in oaks

Sylvain Schmitt [iD], Thibault Leroy [iD], Myriam Heuertz [iD], Niklas Tysklind [iD]

CNRS, UMR EcoFoG (Agroparistech, Cirad, INRAE, Université des Antilles, Université de la Guyane), Campus Agronomique, 97310 Kourou, French Guiana;   Department of Botany and Biodiversity Research, University of Vienna, Rennweg 14, A-1030 Vienna, Austria;   IRHS-UMR1345, Université d'Angers, INRAE, Institut Agro, SFR 4207 QuaSaV, 49071 Beaucouzé, France;   Université Bordeaux, INRAE, BIOGECO, 69 route d'Arcachon, CS 80227, 33612 Cestas Cedex, France;   INRAE, UMR EcoFoG (Agroparistech, CNRS, Cirad, Université des Antilles, Université de la Guyane), Campus Agronomique, 97310 Kourou, French Guiana

- **Corresponding author**: Sylvain Schmitt, +33 6 49 19 32 63, sylvain.m.schmitt@gmail.com
- **Running title:** Somatic mutation detection in plants

Abstract

1. Mutation, the source of genetic diversity, is the raw material of evolution; however, the mutation process remains understudied, especially in plants. Using both a simulation and reanalysis framework, we set out to explore and demonstrate the improved performance of variant callers developed for cancer research compared to single nucleotide polymorphism (SNP) callers in detecting *de novo* somatic mutations.

2. In an *in silico* experiment, we generated Illumina-like sequence reads spiked with simulated mutations at different allelic fractions to compare the performance of seven commonly-used variant callers to recall them. More empirically, we then reanalyzed two of the largest datasets available for plants, both developed for identifying within-individual variation in long-lived pedunculate oaks.

3. Based on the *in silico* experiment, variant callers developed for cancer research outperform SNP callers regarding plant mutation recall and precision, especially at low allele frequency. Such variants at low allelic fractions are typically expected for within-individual *de novo* plant mutations, which initially appear in single cells. Reanalysis of published oak data with *Strelka2*, the best-performing caller based on our simulations, identified up to 3.4x more candidate somatic mutations than reported in the original studies.

4. Our results advocate the use of cancer research callers to boost *de novo* mutation research in plants, and to reconcile empirical reports with theoretical expectations.

Introduction

DNA sequence mutation is the raw material for evolutionary change, but, despite its crucial role, many fundamental questions around the mutation process are still open. Notwithstanding its apparent simplicity, the understanding of mutation processes is one of the most common conceptual difficulties for biologists (Smith & Knight, 2012; Prevost *et al.*, 2013). Mutations are often assumed to occur at a relatively constant pace (i.e. following the hypothesis of a 'perfect' molecular clock). Despite the extremely low number of direct mutation rates estimates currently available in the literature, mutation rates are, however, known to be highly variable across the tree of life, differing by several orders of magnitude among species and kingdoms, and are considered as an evolvable trait *per se* Lynch *et al.*, (2016). In mammals, the somatic mutation rate varies directly with life span (Cagan *et al.*, 2022). Mutations are assumed to be random, but the rate at which different nucleotides mutate strongly depends on the genomic context, in particular the surrounding nucleotides (Martincorena & Campbell, 2015), hereafter referred to as a mutation spectrum. The mutation spectra themselves are now believed to evolve over time (Milholland *et al.*, 2017), even at relatively short evolutionary timescales (Harris & Pritchard, 2017). The relative contribution of DNA replication and DNA repair errors to mutation rates represents another timely evolutionary question in the field (Gao *et al.*, 2019).

Unlike most animals that transmit to the next generation only mutations present in their germ cells (*i.e.* sperm and eggs), plants are expected to produce heritable somatic mutations as they grow throughout their lives, departing from the so-called Weismann's germ plasm

theory (Weismann, 1893; but see also Lanfear, 2018). As a consequence, long-lived species,

such as trees, are generally assumed to accumulate more heritable mutations than short-

lived species per generation (Hanlon *et al.,* 2019). To generate new knowledge on plant

mutation processes (Schoen & Schultz, 2019), several studies examined within-individual

variation in long-lived trees, whose individuals can live for more than a thousand years

(Schöngart *et al.,* 2017). Two studies used the pedunculate oak (*Quercus robur*), a long-lived

diploid and highly heterozygous European tree species, as a plant model to identify somatic

mutations. Schmid-Siegert *et al.,* (2017) identified 17 mutations by comparing sequencing

data from two branches of a 234-year-old individual. The authors therefore argued that

their results are consistent with a low mutation rate in pedunculate oak. Plomion *et al.,*

(2018) identified 46 mutations using three branches of a younger century-old individual,

which is an almost 10-fold higher rate after taking the tree age difference into account.

Plomion *et al.,* (2018) also recovered these new mutations on acorn embryos collected on

the same branches as those used for the *de novo* mutation identification, therefore

producing empirical support for departure from Weismann's germ plasm theory in oaks. A

shared limitation of both studies is that the authors used a single variant caller for *de novo*

mutation detection, without investigating beforehand the robustness of the results from the

selected variant caller. The absence of a simulation work to identify the best suited

detection method prior to the empirical investigations therefore represents a major limit

with regards to the accuracy and completeness of the previously reported *de novo*

mutations.

Variant callers are designed each for a specific purpose, with choices made by the developer

on sequence read filtering, and models and thresholds of sensitivity for the output of

variants. Broadly, we can distinguish 1. Single nucleotide polymorphism (SNP) callers

designed to detect heterozygous sites, i.e. sites with an expected allelic fractions of 0.5 in

the analysed sample, and 2. Cancer callers, designed to detect sites that are mutated in a

fraction of mutated cells in a sample, i.e. with expected allelic fraction $\leq$0.5. When used to

detect mutations, SNP callers primarily detect candidate mutations per sample against the

reference genome and validate mutation robustness by comparing results between sample

pairs (Fig. 1). The per-sample strategy used in SNP callers carries the risk of overlooking

low-frequency mutated reads in one or more samples, which risks invalidating the mutation

in the other sample. To better address the specificities of detecting low frequency

mutations, the development of tools to detect mutations in humans is rapidly expanding in

cancer research (Kim et al., 2018; Alioto et al., 2015), where cancer callers identify

mutations by comparing two samples, one mutated and one normal sample, against the

reference genome (Fig. 1). Detecting mutations in cancers is conceptually similar to

detecting somatic mutations in plants, i.e., the aim is to detect mutations that potentially

affect only a small fraction of the sequenced tissue. However, cancer research frequently

uses very high sequencing depths (100X - 1000X), while the depth available for plants is

often considerably lower (e.g., 34X for Hanlon et al., 2019; 40X for Wang et al., 2018; or 70X

for Schmid-Siegert et al., 2017), bar a few exceptions (240X for Orr et al., 2019; 250X for

Plomion et al., 2018; or 1000X for Watson et al., 2016). Despite the advantage of cancer

callers to identify mutations, the specific challenge of detecting low frequency mutations is

so far poorly addressed in plants, where SNP callers have been the most frequently used

method to detect somatic mutations (Schmid-Siegert et al., 2017; Watson et al., 2016;

Hanlon et al., 2019; Orr et al., 2019). To demonstrate the advantages of using cancer callers

in somatic mutation detection for basic and applied plant research, we evaluated the

performance of the cancer callers compared to SNP callers using biological characteristics

and sequencing depth typical of plant studies.

Here, we performed both an *in silico* and an empirical data-based evaluation of the

performance of variant callers to detect somatic mutations, using simulated reads with

known mutations and two large published datasets on the same species (pedunculate oak,

Quercus robur) that applied different strategies for sequencing depth and mutation

detection (Schmid-Siegert *et al.,* 2017; Plomion *et al.,* 2018; see Fig. S1). We particularly

explored the cancer callers in a plant research context to answer the following questions:

(1) Can cancer research methods, both in terms of protocols (*i.e.* sequencing depth) and

tools (*i.e.* callers), improve the detection of plant somatic mutations?; and (2) Can

reanalyses of within-individual sequencing data provide new insights regarding plant

mutation processes?

Figure 1. SNP callers (top rows) detect candidate mutations per tissue sample (dark green and

light green) against the reference genome (blue) and validate the robustness of mutations by

comparing results between sample pairs, while cancer callers identify mutations by comparing

two samples, one mutated (tissue A, dark green) and one normal (tissue B, light green),

*against the reference genome (blue). At low sequencing depth (**A** and **E**), neither the SNP nor*

*the cancer variant callers detect a low (**A**) or high (**E**) frequency mutation. At intermediate*

*sequencing depths (**B** and **F**), both SNP and cancer callers detect high-frequency mutations*

*(**F**), but cancer callers are expected to be better at detecting low-frequency mutations than*

*SNP callers (**B**), which were originally designed to detect the expected high-frequency*

*heterozygous sites. At high sequencing depths (**C** and **G**), both the SNP and cancer callers*

*detect high frequency (**C**) and low frequency (**G**) mutations. However, with intermediate*

*sequencing depth (**D** and **H**), a poorly represented heterozygous site in one tissue may remain*

undetected in that tissue by the SNP caller while it may be detected in the second tissu and

*thus be considered a mutation, resulting in a false positive (**D**). By comparing the two samples*

*together, cancer callers will avoid this error (**H**).*

Material and methods

Study design

We developed two workflows: 1) to generate Illumina-like sequencing reads including mutations with varying biological and sequencing parameters; and 2) to detect mutations with multiple variant callers (Fig. S1). We used *singularity* containers (Kurtzer *et al.,* 2017) and the *snakemake* workflow engines (Köster *et al.,* 2012) to build automated, highly reproducible (FAIR), and scalable workflows. We then used both workflows to test for the best performing variant caller for mutation detection *in silico* based on biological and sequencing parameters. We finally used the identified variant caller to detect mutations in pedunculate oak, *Quercus robur* L., by re-analysing data from two somatic mutation projects on oaks led by INRAE Bordeaux, France (Plomion *et al.,* 2018) and the University of Lausanne, Switzerland (Schmid-Siegert *et al.,* 2017).

Generation of mutations

To ensure the feasibility of the project and to limit the computational load, a first step is to subsample one or several sequences of user-defined length in the reference genome. The first workflow named *generateMutations* therefore uses a bespoke R script named *sample_genome* to generate these subsets. The workflow then takes advantage of the two scripts included in *simuG* (v1.0.1, Yue & Liti, 2019), *vcf2model.pl,* and *simuG.pl,* respectively, 1) to build a model of heterozygous sites distribution for an haploid reference genome

based on a user-defined set of known heterozygous sites in *vcf* format and 2) to build the

second reference haploid genome comprising a user-defined number of heterozygous sites

to accurately represent diploidy. Typically, the user can define a number of heterozygous

sites based on the product of nucleotide diversity (π) and genome length (L). The workflow

then uses a homemade R script named *generate_mutations* to spike randomly the reference

genome with a user-defined number of *de novo* mutations which are drawn in a binomial

distribution using a user-defined transition/transversion ratio (**R**). Finally, the workflow

takes advantage of *InSilicoSeq* (v1.5.3, Gourlé *et al.,* 2019) defined with the model option

hiseq to generate datasets of mutated and non-mutated *in silico* Illumina-like sequencing

reads with a user-defined sequencing depth (**C**), sequentially using (1) the reference

haploid genome; (2) considering reference and alternate allele versions at heterozygous

sites, as the workflow was developed for a diploid species; and (3) considering, or ignoring,

the simulated de novo mutations which all feature the same user-defined allelic fraction

(**AF**).

Detection of mutations

The second workflow named *detectMutations* aims to detect somatic mutations from

mapped sequencing reads on a genome reference. Paired-end sequencing reads of every

library are quality checked using *FastQC* (v0.11.9) before trimming using *Trimmomatic*

(v0.39, Bolger *et al.,* 2014) keeping only paired-end reads without adaptors and a phred

score above 15 in a sliding window of 4 bases. Reads are aligned against the reference per

chromosome using BWA *mem* with the option to mark shorter splits (v0.7.17, Li & Durbin,

2009). Alignments are then compressed using *Samtools view* in CRAM format, sorted by

coordinates using *Samtools sort,* and indexed using *Samtools index* (v1.10, Li *et al.,* 2009).

Duplicated reads in alignments are marked using *GATK MarkDuplicates* (v4.2.6.1, Auwera *et*

al., 2013). Finally, the workflow uses seven variant callers to detect mutations, including

variant callers developed originally for SNP calling and variant callers developed initially

for cancer research. SNP callers to detect variants included *GATK HaplotypeCaller* with

GATK GenotypeGVCFs (Auwera *et al.,* 2013) and *freebayes* (v1.3.2, Garrison & Marth, 2012)

using and reporting genotype qualities, without priors on allele balance, with a minimum

alternate allelic fraction of 0.03, a minimum repeated entropy of 1 and a minimum alternate

allele count of 2. Cancer callers developed for mutation detection included *VarScan* (v2.4.3,

Koboldt *et al.,* 2009), *Strelka2* (v2.9.10, Kim *et al.,* 2018), *MuSE* (v0.1.1, Fan *et al.,* 2016),

Mutect2 (using a panel of normal and without soft clipped bases; within v4.2.6.1; Benjamin

et al., 2019), and *Somatic Sniper* (v1.0.5.0, filtering reads with mapping quality less than 25,

filtering mutations with quality less than 15 with prior probability of a mutation of 0.0001;

Larson *et al.,* 2012). Then we only focused on the simulated mutations, and therefore

excluded from the analyses the known heterozygous sites provided by the user thanks to

the vcf file for *GATK, freebayes, Somatic Sniper,* and *Strelka2* using *BEDTools subtract*

(v2.29.2, Quinlan & Hall, 2010) or directly within the variant caller for *Mutect2* and *VarScan*.

In silico experiment

We used the *generateMutations* workflow to generate 1,000 mutations in the oak genome

with varying biological and sequencing parameters. To ensure consistency between the *in*

silico experiment and the reanalysis of empirical data, we used the reference genome

"Qrob_PM1N'' of Quercus robur 3P from Bordeaux, ENA accession number PRJEB8388

(Plomion *et al.,* 2018), thus assessing the behaviour of variant callers in the same genomic

context as used for the empirical work. To reduce the computational load, we only

generated mutations on the first megabase of the first chromosome of the oak assembly

("Qrob_Chr01") in order to later focus the detection on this region. To check that the

conclusions regarding the callers are independent of the considered genomic region, we

ran five independent investigations based on randomly selected genome areas of a

megabase in length. Our results were highly congruent over all our investigations

(Pearson's correlations across all callers and all simulations, recall: 0.999, precision: 0.947,

but see Table S1 for differences among callers). We used known heterozygous sites from

the reference genome (Plomion *et al.,* 2018) to simulate back ten thousand heterozygous

sites (N = π x L = 10^4, assuming π = 0.01 (Plomion *et al.,* 2018) and L = 1 Mb). We used

varying values of transition/transversion ratio (**R** = [2, 2.5, 3]), allelic fraction (**AF** = [0.05,

0.1, 0.25, 0.5]), and sequencing depth (**C** = [25, 50, 100, 150, 200]), resulting in 60 simulated

datasets of mutated and associated base reads (3**R** x 4**AF** x 5**C**). We then used the

detectMutations workflow to detect (recall) spiked mutations with every variant caller

(*Mutect2, freebayes, GATK, Strelka2, VarScan, Somatic Sniper,* and *MuSe*). Using known

spiked mutations, we assessed the number of true positive (TP), false positive (FP), and

false negative (FN) for each variant caller to detect mutations and each combination of

biological and sequencing parameters. We used the resulting confusion matrix to calculate

the recall ($\frac{TP}{TP+FN}$) and the precision rates ($\frac{TP}{TP+FP}$). The recall rate represents the ability of

the variant caller to detect all mutations, while the precision rate represents the ability of

the variant caller to not confound other sites with mutations. We finally assessed each

variant caller to detect mutations using the recall and the precision rates with varying

transition/transversion ratio (**R**), allelic fraction (**AF**), and sequencing depth (**C**) to identify

the best performing variant caller based on biological and sequencing parameters.

Oak data reanalyses

We re-analyzed publicly available data on pedunculate oak from two projects conducted in

Lausanne, Switzerland (Schmid-Siegert *et al.,* 2017) and Bordeaux, France (Plomion *et al.,*

2018) (SRA PRJNA327502 and ENA PRJEB8388, respectively). Sequenced reads of every

library were quality checked, trimmed, and mapped with the same strategy than previously

described for the simulation work, i.e. using *FastQC, Trimmomatic,* and *BWA mem*. We then

used the best-performing variant caller at low coverage and allelic fraction based on our *in*

silico investigation, *Strelka2*, and the variant caller for mutation detection from the original

publication to compare the results, *i.e.*, *GATK* with *Best Practices* for the data from Schmid-

Siegert *et al.* (2017) and *Mutect2* for the data from Plomion *et al.* (2018). The former

comprised 2 libraries of medium sequencing depth (60X) representing one lower and one

upper branch. The latter comprised 3 libraries of high sequencing depth (160X)

representing 3 branches (lower, mid, and upper). For both data sets, we compared each

pair of sample points sequentially as the reference library and the potentially mutated

library to distinguish mutations among branches from heterozygous sites and sequencing

errors. For the data from Plomion *et al.* 2018, we further filtered out candidate somatic

mutations by using a cross-validation procedure to keep a coherent temporal pattern

among mutations following the original publication (Plomion *et al.,* 2018). Contrary to a

general expectation and a common view in the field (Schmid-Siegert *et al.*, 2017, Orr *et al.*,

2019), detected mutations do not always accumulate following the developing plant

architecture (Zahradníková *et al.*, 2020; Ren *et al.*, 2021). As a consequence, our cross-

validation represents a conservative strategy for the mutation detection, but it should be

noted that this strategy could have removed some true somatic mutations. We used these

raw datasets to identify the mutations from the original studies after realigning the

megabase containing the mutation on the 3P genome using *BLAT* (Kent, 2002). For both

datasets, we finally kept candidate mutations with (1) a read depth for both the normal and

mutated samples between half and two times the mean sequencing depth (30-120X and 80-

320X for Schmid-Siegert *et al.,* (2017) and Plomion *et al.,* (2018) datasets, respectively), (2)

an absence of the mutated allele in the normal sample, and (3) a minimum of 10 copies of

the mutated allele in the mutated sample. In addition, *Strelka2* calculates an empirical

variant score (EVS) based on a supervised random forest classifier trained on data from

sequencing runs under various conditions, which provides an overall quality score for each

variant (Kim *et al.*, 2018). We took advantage of the EVS to define a conservative set of

candidate mutations for both datasets, hereafter referred to as the EVS datasets. Given that

the proportion of the genome falling within the sequencing depth boundaries used for the

detection (i.e. between 50 and 200% of the mean sequencing depth) varies depending on

the dataset, we weighted the observed number of mutations by the proportion of the

genome satisfying the sequencing depth criteria to provide a more accurate and

comparable estimate of the real total number of mutations. Across both empirical studies,

the proportion of the genome with 50-200% sequencing depth varies between 71 and 87%.

Results

To advocate the use of cancer callers in somatic mutation detection for plant research, we

simulated sequencing data containing new mutations at a given allelic fraction (i.e. fraction

of simulated reads per genomic position carrying the mutated allele), and using varying

depths of sequencing (for variable transition/transversion ratios, see Fig S2). We then

evaluated the performance of variant callers as a function of allelic fraction and sequencing

depth. We found marked differences in: (1) the recall, the ability to recover the simulated

mutations; and (2) the precision, the proportion of true simulated mutations among all

variants detected. For allelic fractions equal to, or lower than, 0.25, cancer variant callers

(*Strelka2, Mutect2, MuSE,* but not *Somatic Sniper*) outperform SNP callers such as *GATK*,

freebayes, and *VarScan* (Fig. 2 and S3), mainly based on the recall. For allelic fractions over

0.25, all variant callers perform similarly well, except for *freebayes*, which identified many

false positives. Over the 80 tested parameter combinations, *Strelka2* was the best

performing variant caller for various allelic frequencies and sequencing depths (in 57/80

simulated datasets , with an average recall of 0.95 for a precision of 0.98, Fig. S2-4 and

Table S2 and S3) and an excellent computational efficiency (second fastest caller, Fig. S9).

Figure 2: Variant caller performances to identify simulated mutations for varying allelic

fractions and sequencing depths (see Fig. S5 for all parameter combinations). The recall is the

ability to detect (recover) the simulated mutations. The precision is the proportion of

simulated mutations among all variants detected (i.e. including false positives). Each point

represents the averaged mutation recall or precision (10 simulations) for increasing allelic

fraction and sequencing depth. The shaded area represents the variation of recall and

precision rates over the 10 replicates computed for all callers, but only visible for the precision

of Muse, Mutect2, and VarScan. Linetype distinguishes SNP callers (dashed) and cancer callers

(solid).

We further investigated the performance of the best performing variant caller based on our

in silico experiment, *Strelka2*, on two empirical datasets on pedunculate oak (Schmid-

Siegert *et al.,* 2017; Plomion *et al.,* 2018) in comparison to the variant callers used in the

original publications, *i.e.*, *GATK* and *Mutect2*, respectively. Mapping the raw data of Schmid-

Siegert *et al.,* (2017) and Plomion *et al.,* (2018) on the oak genome that we used as a

mapping reference for our empirical study, we successfully mapped 14 and 60 of the

mutations detected in the original articles, respectively. Across variant callers, we

recovered 12 (86%) and 60 (100%) of these original mutations in our total list of candidate

somatic mutations (Fig. 3A), strongly supporting the results shown by the two previous

studies. However, our analyses were able to detect far more candidate mutations than

initially reported. The filtering based on sequencing depth reduced the proportion of the

genome covered with adequate sequencing depth to 72% (assuming 30-120X) for Schmid-

Siegert *et al.,* (2017) and to 84% (assuming 80-320X) for Plomion *et al.,* (2018). Similarly,

the cross-validation between branches following the plant development (see Materials and

Methods for details) filtered out 27% of mutations detected for the dataset of Plomion *et al.,*

(2018). Using filtering based on sequencing depth and mutated allele copies with cross-

validation, *Strelka2* produced a smaller set of candidate mutations than *GATK* but similar to

Mutect2, with an estimated number of mutation candidates 2- to 4.3-fold higher than that

of the original studies (Fig. 3A). Adding *Strelka2* recommended filtering based on empirical

variant scores (EVS) yielded the most conservative dataset with a 1.5 to 3.4-fold increase

compared to the original number of mutations. Due to lack of access to biological material

from the original studies, conclusions were drawn from this list of conservative candidate

somatic mutations (but see discussion regarding validation of mutations). The distribution

of allelic fractions of detected mutations partly explains differences among detection

methods (Fig. 3A), with *Strelka2* and *Mutect2* detecting mutations with lower allelic

fractions than the candidate mutations presented in the original publications, especially for

the Plomion *et al.,* (2018) study that used higher sequencing depths allowing the

identification of these numerous low frequency mutations.

Based on the set of conservative mutations detected by *Strelka2* (EVS), we then explored

annotations and mutation spectra in both datasets (Fig. 3B-C), which have rarely been

explored in model plant species (but see first evidence based on mutation accumulation

lines in *Arabidopsis thaliana* in Weng *et al.,* 2019) and never in the wild. The proportions of

mutations found in different genomic regions (e.g. genic, intergenic) were highly correlated

between both original studies and proportional to the representation of the genomic

regions, supporting a random distribution of mutations throughout the genome (Fig. 3B).

Mutation spectra of the two studies are significantly correlated (Pearson'r=0.49, p<10-5),

with an enrichment in C>T transitions, particularly in some specific genomic contexts (Fig.

3C).

Figure 3: Candidate mutation spectra depending on variant callers and filtering in Schmid-Siegert et al., (2017) and Plomion et al., (2018). **A.** *Allelic* *fraction* *distribution for every dataset, including the candidate mutations from the original article present in the raw data from the reanalysis (red), the results of GATK with Best Practices (blue), Mutect2 after filtering (green), and Strelka2 after filtering (purple), and the results of Strelka2 using the filtering based on empirical variant scores named EVS (orange). The labels indicate the number of candidate mutations in each dataset. Per caller comparisons are available in Fig. S7.* **B.** *Annotation of the mutations detected by Strelka2 across chromosomes using the filtering based on empirical variant scores named EVS for Schmid-Siegert et al., (2017, green) and Plomion et al., (2018, orange) compared to the genomic expectation (grey, see Supplementary Note S1). Error bars represent the standard deviation (SD) of the observed*

percentages across chromosomes, and the annotation above the columns indicates the significance of the Student's t-test two-sided comparing the mean percentage of mutations to the mean genomic expectation, with ns, **, and *** corresponding to non-significant, p<0.01, and p<0.001 differences, respectively. **C.** *Context-dependent mutation spectra depending on mutation types for the results of Strelka2 using the filtering based on empirical variant scores named EVS. Mutation types have been summarised into six main classes with thicker lines for transversion compared to transition, and then differentiated depending on their 5' and 3' genomic contexts, see Fig. S8-9. Pearson's correlation r measures the two-sided correlation of the mutation spectra between Schmid-Siegert et al., (2017) and Plomion et al., (2018). All figures compare the reanalysed data and not the original results.*

Discussion

Mutation research in plants still primarily uses SNP callers and methodologies that are not developed for the specificity and complexity of within-individual *de novo* mutation detection. We demonstrated that plant mutation research could benefit from the development of tools and protocols initially designed for human cancer research, which is a rapidly expanding field (Kim *et al.,* 2018). We quantified expected marked differences in the performance of variant callers for mutation detection based on sequencing depth and allelic fraction. We demonstrated that cancer callers performed better than SNP callers for mutation detection at low or intermediate allelic fraction or with low sequencing depth, and similarly well for high allelic fraction. The overall higher detection efficiency demonstrates the benefit of using tools and protocols initially designed for human cancer research. Low

allelic fraction mutations, potentially due to the chimeric nature of plant shoot apical

meristems structures (Burian, 2021), might be very important due to their great abundance

that may balance out their low chance of transmissions. Therefore, plant mutation studies

should make greater use of cancer variant callers such as *Strelka2* rather than SNP callers

such as *GATK* to detect somatic mutations, in agreement with previous studies on germline

mutations detection (Chen *et al.,* 2019), especially for detecting low frequency mutations

and when using low sequencing depth. The importance of allele fraction-dependency in

variant detection is not restricted to somatic mutations, but also concerns for instance

polyploid species, which includes many agriculturally important autopolyploid plant

species (e.g. potato, sugarcane). Our simulation framework therefore provides general

insights regarding the impact of allele fraction in mutation detection which go beyond

somatic mutation detection.

Validation of mutations

Intuitively, after identifying candidate mutations, one expects their "validation". In our

study, it was impossible to validate mutations using genotyping or resequencing because

we performed a reanalysis of already available sequence data, without having access to the

biological material. Yet, we argue that validation is more complex than generally thought.

We specifically discuss (i) allele fraction, (ii) fraction-aware approach, and (iii) offspring

validation. High frequency mutations (e.g. AF=0.5) will be easily confirmed by genotyping

or Sanger sequencing. But if one considers that low frequency mutations are only present in

a fraction of the tissue, SNP genotyping technologies are hindered by the complexity of

defining the genotype clustering while Sanger sequencing is hindered by the difficulty of

isolating the allelic assay from the background when reading the chromatogram. Therefore,

such validation strategies are conservative and favour high frequency mutations. To

unambiguously validate low-frequency mutations using intra-individual samples, an allelic

fraction-aware approach can be used, such as a hybridisation capture-based sequencing

strategy targeting regions of candidate mutations with high-coverage sequencing. However,

this would mean that the validation is based on another round of sequencing, bringing back

all the biases of the bioinformatics steps. Another strategy could be genotyping of inherited

mutations in offspring, working at the family rather than the individual level, which also has

its limitations (see Note S2). From our point of view, the current literature on somatic

mutations in plants does not mention the limits of validation, focusing on "fully validated"

mutations, thus probably favouring high frequency mutations. However, high-frequency

mutations are expected to be relatively rare compared to low-frequency mutations, which

affect conclusions about the total number of somatic mutations. In the relatively near

future, the use of single cell sequencing approaches to reveal intra-individual genetic

variation is likely to eliminate the problem at its source, avoiding the use of mixed samples

of mutated and non-mutated cells, leading to simple detection and validation.

Limits and future directions

396 Our study provided a first simple case of evaluating the performance of callers to detect

397 somatic mutations in plants. Our study was limited to variant callers in the cases of mutated

398 and normal sample pairs using the recommended default parameters. Future studies

 should consider and quantify the performance of other types of callers, such as subclonal or

mosaic variant callers (e.g. *deepSNV* and *MosaicForecast*). Future studies could also consider

exploring tuning parameters for the specific task of detecting plant somatic mutations. We

hope that the open-source automated, highly reproducible, and scalable workflows

constructed for our study, *generateMutations* and *detectMutations*, will allow further

exploration of these questions. Our study also overlooked the importance of genome size, as

plants can have larger genomes than humans. Trees have a very wide range of genome

sizes, from small genomes (0.35 GB) to very large genomes (>20 GB), and our study focused

on the relatively small oak genome (0.75 GB). A large genome will obviously be a barrier to

high sequencing depth with a fixed budget, but with regard to mutation detection, our study

argues for the use of cancer callers in large genomes as they also outperform SNP callers in

terms of computational time (Fig. S9). Similarly, our study did not take into account the role

of polyploidy as we only focused on a diploid species. We assume that for ploidy greater

than two, the performance of cancer callers relative to SNP callers will increase since the

cancer caller will be even more efficient at low frequencies. But future studies could also

take advantage of our tools to explore the role of polyploidy in caller performance. Another

problem that may arise when analysing sample pairs with cancer callers is the rapid

increase in pairwise comparisons when using a larger sample size than previous studies

(e.g., N=3 in Plomion *et al.* 2018). A simple solution is the use of a single reference sample

such as a cambium sample from the base of the tree, which is therefore considered as the

closest genome to the seed, to compare it to all samples from branches (Hanlon *et al.,* 2019).

Conclusion

By reanalyzing the raw oak data (Schmid-Siegert *et al.,* 2017; Plomion *et al.,* 2018), we

found that the marked differences in the performance of variant callers could account for

the discrepancies in genome-wide plant somatic mutation rate estimates. Our reanalysis

using the best-performing caller based on our simulations, *Strelka2*, suggests an up to 3.4-

fold higher number of mutations than previously reported, a value closer to the

expectations based on the theory (Schoen & Schultz, 2019; Burian, 2021). We argue that

knowledge and methodological transfers from cancer to plant mutation detection are

expected to contribute strongly to the upward trend of this field and to reconcile empirical

reports with theoretical expectations.

Acknowledgments

431   The manuscript benefited from the comments of Nicolas Bierne and five anonymous

432   reviewers.

Funding

434   This study was funded through an Investissement d'Avenir grant of the ANR: CEBA (ANR-

435   10-LABEX-0025).

Conflict of interest disclosure

The authors declare they have no conflict of interest relating to the content of this article.

MH is a recommender for PCI Evol Biol.

Data, script and code availability

Reanalyzed reads and corresponding genomes were extracted from GenBank under accession BioProject PRJNA327502 and from European Nucleotide Archive under project accession code PRJEB8388. `generateMutations` and `detectMutations` pipelines are available on GitHub (https://github.com/sylvainschmitt/generateMutations and https://github.com/sylvainschmitt/detectMutations).

Authors' contributions

References

Alioto, T.S., Buchhalter, I., Derdak, S., Hutter, B., Eldridge, M.D., Hovig, E., Heisler, L.E., Beck, T.A., Simpson, J.T., Tonon, L., Sertier, A.S., Patch, A.M., Jäger, N., Ginsbach, P., Drews, R.,

Paramasivam, N., Kabbe, R., Chotewutmontri, S., Diessl, N., Previti, C., Schmidt, S., Brors, B.,

Feuerbach, L., Heinold, M., Gröbner, S., Korshunov, A., Tarpey, P.S., Butler, A.P., Hinton, J.,

Jones, D., Menzies, A., Raine, K., Shepherd, R., Stebbings, L., Teague, J.W., Ribeca, P., Giner,

F.C., Beltran, S., Raineri, E., Dabad, M., Heath, S.C., Gut, M., Denroche, R.E., Harding, N.J.,

Yamaguchi, T.N., Fujimoto, A., Nakagawa, H., Quesada, V., Valdés-Mas, R., Nakken, S., Vodák,

D., Bower, L., Lynch, A.G., Anderson, C.L., Waddell, N., Pearson, J.V., Grimmond, S.M., Peto, M.,

Spellman, P., He, M., Kandoth, C., Lee, S., Zhang, J., Létourneau, L., Ma, S., Seth, S., Torrents, D.,

Xi, L., Wheeler, D.A., López-Otín, C., Campo, E., Campbell, P.J., Boutros, P.C., Puente, X.S.,

Gerhard, D.S., Pfister, S.M., McPherson, J.D., Hudson, T.J., Schlesner, M., Lichter, P., Eils, R.,

Jones, D.T.W. & Gut, I.G. (2015). A comprehensive assessment of somatic mutation detection

in cancer using whole-genome sequencing. *Nature Communications*, **6**.

Auwera, G.A., Carneiro, M.O., Hartl, C., Poplin, R., Angel, G. del, Levy-Moonshine, A., Jordan,

T., Shakir, K., Roazen, D., Thibault, J., Banks, E., Garimella, K.V., Altshuler, D., Gabriel, S. &

DePristo, M.A. (2013). From FastQ Data to High-Confidence Variant Calls: The Genome

Analysis Toolkit Best Practices Pipeline. *Current Protocols in Bioinformatics*, **43**, 483–492.

Retrieved from https://onlinelibrary.wiley.com/doi/abs/10.1002/0471250953.bi1110s43

Benjamin, D., Sato, T., Cibulskis, K., Getz, G., Stewart, C. & Lichtenstein, L. (2019). Calling

Somatic SNVs and Indels with Mutect2.

Bolger, A.M., Lohse, M. & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina

sequence data. *Bioinformatics*, **30**, 2114–2120.

Burian, A. (2021). Does Shoot Apical Meristem Function as the Germline in Safeguarding

Against Excess of Mutations? *Frontiers in Plant Science*, **12**, 1–9.

Cagan, A., Baez-Ortega, A., Brzozowska, N., Abascal, F., Coorens, T. H. H., Sanders, M. A., …

Martincorena, I. (2022). Somatic mutation rates scale with lifespan across mammals.

Nature, 604(7906), 517–524. https://doi.org/10.1038/s41586-022-04618-z

Chen, Z.L., Meng, J.M., Cao, Y., Yin, J.L., Fang, R.Q., Fan, S.B., Liu, C., Zeng, W.F., Ding, Y.H., Tan,

D., Wu, L., Zhou, W.J., Chi, H., Sun, R.X., Dong, M.Q. & He, S.M. (2019). A high-speed search

engine pLink 2 with systematic evaluation for proteome-scale identification of cross-linked

peptides. *Nature Communications*, **10**. Retrieved from http://dx.doi.org/10.1038/s41467-

019-11337-z

Fan, Y., Xi, L., Hughes, D.S.T., Zhang, J., Zhang, J., Futreal, P.A., Wheeler, D.A. & Wang, W.

(2016). MuSE: accounting for tumor heterogeneity using a sample-specific error model

improves sensitivity and specificity in mutation calling from sequencing data. *Genome*

biology, **17**, 178. Retrieved from http://dx.doi.org/10.1186/s13059-016-1029-6

Gao, Z., Moorjani, P., Sasani, T.A., Pedersen, B.S., Quinlan, A.R., Jorde, L.B., Amster, G. &

Przeworski, M. (2019). Overlooked roles of DNA damage and maternal age in generating

human germline mutations. *Proceedings of the National Academy of Sciences of the United*

States of America, **116**, 9491–9500.

Garrison, E. & Marth, G. (2012). Haplotype-based variant detection from short-read

sequencing. 1–9. Retrieved from http://arxiv.org/abs/1207.3907

Gourlé, H., Karlsson-Lindsjö, O., Hayer, J. & Bongcam-Rudloff, E. (2019). Simulating Illumina

metagenomic data with InSilicoSeq. *Bioinformatics*, **35**, 521–522.

Hanlon, V.C.T., Otto, S.P. & Aitken, S.N. (2019). Somatic mutations substantially increase the

per-generation mutation rate in the conifer Picea sitchensis. *Evolution Letters*, **3**, 348–358.

Harris, K. & Pritchard, J.K. (2017). Rapid evolution of the human mutation spectrum. *eLife*,

6, 1–17.

Kent, W.J. (2002). BLAT—The BLAST-Like Alignment Tool. *Genome Research*, **12**, 656–664.

Kim, S., Scheffler, K., Halpern, A.L., Bekritsky, M.A., Noh, E., Källberg, M., Chen, X., Kim, Y.,

Beyter, D., Krusche, P. & Saunders, C.T. (2018). Strelka2: fast and accurate calling of

germline and somatic variants. *Nature Methods*, **15**, 591–594.

Koboldt, D.C., Chen, K., Wylie, T., Larson, D.E., McLellan, M.D., Mardis, E.R., Weinstock, G.M.,

Wilson, R.K. & Ding, L. (2009). VarScan: Variant detection in massively parallel sequencing

of individual and pooled samples. *Bioinformatics*, **25**, 2283–2285.

Köster, J. & Rahmann, S. (2012). Snakemake-a scalable bioinformatics workflow engine.

Bioinformatics, **28**, 2520–2522.

Kurtzer, G.M., Sochat, V. & Bauer, M.W. (2017). Singularity: Scientific containers for mobility

of compute. *PLoS ONE*, **12**, 1–20.

Lanfear, R. (2018). Do plants have a segregated germline? *PLoS Biology*, **16**, 1–13.

Larson, D.E., Harris, C.C., Chen, K., Koboldt, D.C., Abbott, T.E., Dooling, D.J., Ley, T.J., Mardis,

E.R., Wilson, R.K. & Ding, L. (2012). Somaticsniper: Identification of somatic point mutations

in whole genome sequencing data. *Bioinformatics*, **28**, 311–317.

Li, H. & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics*, **25**, 1754–1760.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G. &

Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, **25**,

2078–2079.

Lynch, M., Ackerman, M.S., Gout, J.F., Long, H., Sung, W., Thomas, W.K. & Foster, P.L. (2016).

Genetic drift, selection and the evolution of the mutation rate. *Nature Reviews Genetics*, **17**,

704–714. Retrieved from http://dx.doi.org/10.1038/nrg.2016.104

Martincorena, I. & Campbell, P.J. (2015). Somatic mutation in cancer and normal cells.

Science, **349**, 1483–1489.

Milholland, B., Dong, X., Zhang, L., Hao, X., Suh, Y. & Vijg, J. (2017). Differences between

germline and somatic mutation rates in humans and mice. *Nature Communications*, **8**, 1–8.

Retrieved from http://dx.doi.org/10.1038/ncomms15183

Orr, A.J., Padovan, A., Kainer, D., Külheim, C., Bromham, L., Bustos-Segura, C., Foley, W., Haff,

T., Hsieh, J.F., Morales-Suarez, A., Cartwright, R.A. & Lanfear, R. (2019). A phylogenomic

approach reveals a low somatic mutation rate in a long-lived plant. *bioRxiv*.

Plomion, C., Aury, J.M., Amselem, J., Leroy, T., Murat, F., Duplessis, S., Faye, S., Francillonne,

N., Labadie, K., Le Provost, G., Lesur, I., Bartholomé, J., Faivre-Rampant, P., Kohler, A., Leplé,

J.C., Chantret, N., Chen, J., Diévart, A., Alaeitabar, T., Barbe, V., Belser, C., Bergès, H., Bodénès,

C., Bogeat-Triboulot, M.B., Bouffaud, M.L., Brachi, B., Chancerel, E., Cohen, D., Couloux, A., Da

Silva, C., Dossat, C., Ehrenmann, F., Gaspin, C., Grima-Pettenati, J., Guichoux, E., Hecker, A.,

Herrmann, S., Hugueney, P., Hummel, I., Klopp, C., Lalanne, C., Lascoux, M., Lasserre, E.,

Lemainque, A., Desprez-Loustau, M.L., Luyten, I., Madoui, M.A., Mangenot, S., Marchal, C.,

Maumus, F., Mercier, J., Michotey, C., Panaud, O., Picault, N., Rouhier, N., Rué, O., Rustenholz,

C., Salin, F., Soler, M., Tarkka, M., Velt, A., Zanne, A.E., Martin, F., Wincker, P., Quesneville, H.,

Kremer, A. & Salse, J. (2018). Oak genome reveals facets of long lifespan. *Nature Plants*, **4**,

440–452. Retrieved from http://dx.doi.org/10.1038/s41477-018-0172-3

Prevost, L., Knight, J., Smith, M., & Lurain, U. M. (2013). Student writing reveals their

heterogeneous thinking about the origin of genetic variation in populations.

Quinlan, A.R. & Hall, I.M. (2010). BEDTools: A flexible suite of utilities for comparing

genomic features. *Bioinformatics*, **26**, 841–842.

Ren, Y., He, Z., Liu, P., Traw, B., Sun, S., Tian, D., ... Wang, L. (2021). Somatic Mutation

Analysis in Salix suchowensis Reveals Early-Segregated Cell Lineages. *Molecular Biology and*

Evolution, **38**, 5292–5308. https://doi.org/10.1093/molbev/msab286

Schmid-Siegert, E., Sarkar, N., Iseli, C., Calderon, S., Gouhier-Darimont, C., Chrast, J., Cattaneo,

P., Schütz, F., Farinelli, L., Pagni, M., Schneider, M., Voumard, J., Jaboyedoff, M., Fankhauser,

C., Hardtke, C.S., Keller, L., Pannell, J.R., Reymond, A., Robinson-Rechavi, M., Xenarios, I. &

Reymond, P. (2017). Low number of fixed somatic mutations in a long-lived oak tree.

Nature Plants, **3**, 926–929. Retrieved from http://dx.doi.org/10.1038/s41477-017-0066-9

Schoen, D.J. & Schultz, S.T. (2019). Somatic Mutation and Evolution in Plants. *Annual Review*

of Ecology, Evolution, and Systematics, **50**, 49–73.

Schöngart, J., Bräuning, A., Barbosa, A.C.M.C., Lisi, C.S. & Oliveira, J.M. de. (2017).

Dendroecology Tree-Ring Analyses Applied to Ecological Studies.

Smith, M. K., & Knight, J. K. (2012). Using the Genetics Concept Assessment to document

persistent conceptual difficulties in undergraduate genetics courses. Genetics, 191(1), 21–

32. https://doi.org/10.1534/genetics.111.137810

Wang, L., Ji, Y., Hu, Y., Hu, H., Jia, X., Jiang, M., Zhang, X., Zhao, L., Zhang, Y., Jia, Y., Qin, C., Yu,

L., Huang, J., Yang, S., Hurst, L.D. & Tian, D. (2019). The architecture of intra-organism

mutation rate variation in plants. *PLoS Biology*, **17**, 1–29.

Watson, J.M., Platzer, A., Kazda, A., Akimcheva, S., Valuchova, S., Nizhynska, V., Nordborg, M.

& Riha, K. (2016). Germline replications and somatic mutation accumulation are

independent of vegetative life span in Arabidopsis. *Proceedings of the National Academy of*

Sciences of the United States of America, **113**, 12226–12231.

Weismann, A. (1893). *The germ-plasm: a theory of heredity*. Scribner's.

Weng, M.L., Becker, C., Hildebrandt, J., Neumann, M., Rutter, M.T., Shaw, R.G., Weigel, D. &

Fenster, C.B. (2019). Fine-grained analysis of spontaneous mutation spectrum and

frequency in arabidopsis thaliana. *Genetics*, **211**, 703–714.

Yue, J.X. & Liti, G. (2019). SimuG: A general-purpose genome simulator. *Bioinformatics*, **35**,

4442–4444.

Zahradníková, E., Ficek, A., Brejová, B., Vinař, T., & Mičieta, K. (2020). Mosaicism in old trees

and its patterns. *Trees - Structure and Function,* **34**, 357–370.

https://doi.org/10.1007/s00468-019-01921-7

Supplementary material

The following Supporting Information is available for this article:

Note S1. Genomic expectations for the annotation of the mutations

Note S2. 'Validation' of mutations in offsprings

Fig. S1. Study scheme

Fig. S2. Variation in the performance of variant callers for mutation detection with varying biological and sequencing parameters

Fig. S3. Variation in the performance of variant callers for mutation detection with varying biological and sequencing parameters

Fig. S4. Best performing variant callers for mutation detection depending on allelic fraction (allelic fraction) and coverage (sequencing depth)

Fig. S5. Mutation recall and precision rates for SNP and cancer variant callers by allelic fraction and sequencing depth

Fig. S6. Observed allelic frequencies of candidate mutations depending on variant callers and filtering in Schmid-Siegert *et al.,* (2017) and Plomion *et al.,* (2018)

Fig. S7. Percentage of nucleotide change types of candidate mutations depending on variant callers and filtering in Schmid-Siegert *et al.,* (2017) and Plomion *et al.,* (2018)

Fig. S8. Context-dependent mutation spectrum depending on variant callers and filtering in

Schmid-Siegert *et al.,* (2017) and Plomion *et al.,* (2018)

Tab. S1. Mean and standard deviation in performance of variant callers for mutation

detection across all simulations

Tab S2. Mean and standard deviation in performance of variant callers for mutation

detection with varying allelic fraction and sequencing depth

600