RSV immunizes bystander cells to IAV infection using interferons β and λ

Maciej Czerkies[1,=], Marek Kochańczyk[1,=], Zbigniew Korwek[1], Wiktor Prus[1], and Tomasz Lipniacki[1*]

[1] Institute of Fundamental Technological Research, Polish Academy of Sciences, 02-106 Warsaw, Poland.

[=]These authors contributed equally to this work.

[*]Corresponding author: tlipnia@ippt.pan.pl (T.L.)

Abstract

We observed the interference between two prevalent respiratory viruses, respiratory syncytial virus (RSV) and influenza A virus (IAV, H1N1), and characterized its molecular underpinnings in alveolar epithelial cells (A549). We found that RSV induces higher interferon (IFN) β production than IAV and that IFNβ priming confers higher protection against infection with IAV than with RSV. Consequently, we focused on the sequential infection scheme: RSV-then-IAV. Using the A549 WT, IFNAR1 KO, IFNLR1 KO, and IFNAR1–IFNLR1 double KO cell lines we found that both IFNβ and IFNλ are necessary for maximum protection against subsequent infection. Immunostaining revealed that preinfection with RSV partitions the cell population into a subpopulation susceptible to subsequent infection with IAV and an IAV-proof subpopulation. Strikingly, the susceptible cells turned out to be those already compromised and efficiently expressing RSV, whereas the bystander, interferon-primed cells are resistant to IAV infection. Thus, the virus–virus exclusion at the cell population level is not realized through a direct competition for a shared ecological niche (single cell) but rather achieved with the involvement of specific cytokines induced within the host innate immune response.

Author summary

After being challenged with a pathogen, infected cells warn bystander cells about the potential threat by synthesizing and secreting cytokines, especially interferons. In this way, the not yet infected but warned cells are given an opportunity to reinforce their defense mechanisms before they encounter the pathogen replicating in the infected cells whose defense mechanisms have been overcome. Consequently, cell population is partitioned into the cells that may be more prone to infection with another pathogen and the cells that may be expected to be resistant to another pathogen. By studying at the single-cell level the sequential infection with two prevalent respiratory viruses, respiratory syncytial virus (RSV) and influenza A virus (IAV), we showed that the interferon-mediated resistance induced by infection with RSV protects non-infected cells from infection with IAV. The cells compromised by RSV turned out however to be susceptible to invasion of IAV. This demonstrates that, even in the absence of within-cell viral competition, the innate immune response can induct viral interference between primary infection with RSV and secondary infection with IAV.

Introduction

Human respiratory tract infections are predominantly caused by viruses. Out of a plethora of respiratory viruses, recently expanded by the novel beta coronavirus SARS-CoV-2 [1], the three most common recurrent infectants, originating from different viral families, are rhinovirus (human RV, HRV), influenza A virus (IAV), and the respiratory syncytial virus (RSV, human orthopneumovirus) [2], with the latest being the most common pathogen in severe disease in young children and the elderly [3–5]. In temperate areas of the northern hemisphere, the incidence of respiratory infections surges in the winter season and subsides during summer. Detailed examination of temporal patterns of disease outbreaks reveals not only seasonal coincidence but also apparent avoidance of some types of cocirculating respiratory viruses [6–9]. It is unclear whether the epidemiological observations are population-level consequences of an interplay of multiple viral cooperation and competition mechanisms that operate at a single-host level [10]. Intriguingly, the epidemiological avoidance patterns are visible also for taxonomically different groups of viruses, suggesting their potential reliance on mechanisms that are not related to specific antibody-mediated

cross-protection. Although viral interference can be readily observed *in vitro* [11], its underlying mechanisms are necessarily inextricably intertwined with the response of the host organism and await elucidation from a systemic perspective.

Non-antigenic competitive interactions among respiratory viruses have been suggested based on clinical data analysis [12,13] and were recapitulated within animal models [14,15]. Two very recent studies [16,17] demonstrated that an innate immunity mechanism, specifically antiviral cytokine signaling, is implicated in the interference between RV and IAV: blocking the type-I interferon (IFN) response restores IAV replication following RV infection. Whereas type-I IFNs (IFNα and IFNβ) were originally discovered specifically because of their activity against the influenza virus [18], their roles in RSV infection and disease have been controversial [19–25]. A recent animal model study [14] suggested that the RSV–IAV viral interference is antigen-independent, however direct involvement of IFNs has not been demonstrated.

Type I IFNs, together with the more recently identified type III IFNs (IFNλ1–λ4) [26,27], are produced and secreted by infected cells to trigger transcription of an array of IFN-stimulated genes (ISGs) in both the infected and the bystander cells [28–30]. Elevated ISGs restrict viral replication until virus-specific host defense mechanisms develop [31]. IFNα and IFNβ additionally attract specific immune system cells to the site of inflammation [22,32]. IFNλ acts more locally, preferentially at epithelial barriers [33], and differentially affects tissue homeostasis: while IFNλ confers longer lasting protection than IFNα [34], chronic IFNλ exposure inhibits proliferation of respiratory epithelial cells, impairing post-infection lung regeneration [35,36]. A significant relative potency of IFNλ in induction of antiviral state in response to IAV has been demonstrated [34,37–40]. In the absence of IFNβ, IFNλ has been proposed to be the primary factor preventing RSV infection [25].

In this study, we sought to determine molecular underpinnings of the interactions between RSV and IAV. We used a human alveolar epithelial cell line (A549) and characterized direct inter-viral relations and interferon-mediated interference *in vitro* at the single-cell level. We found that RSV induces higher IFNβ production than IAV and that IFNβ is a more potent inducer of STAT1/2 signaling than IFNλ1. Moreover, IFNβ-stimulated cells are more resistant to infection with IAV than to infection with RSV. In light of these findings, we focused on the sequential coinfection scheme, in which cells are preinfected with RSV

and then infected with IAV. We developed A549-derived cell lines with knockout (KO) of type I IFN receptor (IFNAR1 KO), type III IFN receptor subunit (IFNLR1 KO), and double KO (IFNAR1–IFNLR1 dKO). They enabled us to find that IFNβ is the main inducer of the antiviral state in RSV-infected A549 cells and both interferons, IFNβ and IFNλ, are simultaneously necessary for building maximum protection against subsequent infection with IAV. Immunostaining revealed that preinfection with RSV partitions the cell population into a subpopulation susceptible to subsequent infection with IAV and an IAV-proof subpopulation. Strikingly, the susceptible cells turned out to be those already compromised and efficiently expressing the priming virus. Overall, we conclude that in the sequential RSV-then-IAV infection scheme, the virus–virus exclusion at the cell-population level is not realized through a direct competition for a shared ecological niche (single cell) but rather achieved with the involvement of specific cytokines induced within the host innate immune response.

Results

IFNβ and IFNλ mediate STAT1/2 activation upon RSV and IAV infection

We first characterized induction of the antiviral state by interferon signaling. We found that RSV is a more potent inducer of IFNβ than IAV at comparable multiplicities of infection (MOIs) during first two days post-infection (*cf*. Fig 1A, Fig 1B).

We then analyzed interferon-mediated activation of STAT1/2 during RSV or IAV infection and consecutive induction of ISGs that are the primary correlates of the antiviral state: RIG-I (viral RNA sensor [41,42]), PKR (indirect translation inhibitor [43]), and OAS1 (direct activator of RNase L, the RNA eradicator [44]). We juxtaposed results for A549 WT cells with those for the A549-derived cell lines in which we knocked out IFNβ receptor IFNAR1, heterodimeric IFNλ receptor subunit IFNLR1, and both these genes (Fig 1C, Fig 1D). We observed that infection with RSV or IAV leads to robust activation (phosphorylation) of STAT1 and STAT2 in A549 WT cells as well as in IFNLR1 KO cells. In IFNAR1–IFNLR1 dKO cells, there is no observable STAT1/2 phosphorylation, whereas in IFNAR1 KO cells, secreted IFNλ leads to relatively weak STAT1/2 phosphorylation. The observed patterns of STAT activity translate directly to the patterns of accumulation of RIG-I, PKR, and OAS1, as well as STAT1 and STAT2. Cells devoid of IFNβ receptor, but not cells devoid of IFNλ receptor, are more permissive to the spread of RSV infection. Of note, the

lack of either IFNβ receptor or IFNλ receptor facilitates IAV replication, despite IFNLR1 KO having no impact on STAT1/2 signaling nor accumulation of ISGs. Altogether, we conclude that despite divergent efficacy of IFNβ induction by RSV and IAV, upon infection with any of these viruses, IFNβ is the main activator of STAT1 and STAT2 signaling. Both IFNβ and IFNλ secreted by infected cells are responsible for confinement of the spread of IAV, however only IFNβ appears to be capable of attenuating the propagation of RSV.

Prestimulation with IFNβ inhibits propagation of RSV and IAV

To estimate the impact of priming with IFNβ and IFNλ1 on the infectiousness and propagation of RSV and IAV, we quantified viral RNA molecules per cell and analyzed the abundance of viral proteins. We performed infection experiments without or with a preceding, day-long stimulation with IFNβ (1000 U/ml) or with IFNλ1 (50 ng/ml). In Fig 2, we juxtaposed results for these two stimulation protocols. We found that 24 hours post-infection at MOI 0.1, prestimulation with IFNβ limits RSV progeny by nearly one order of magnitude, whereas IAV progeny is reduced by at least two orders of magnitude (*cf.* Fig 2A, Fig 2B). At MOI 1, when nearly all cells are infected, prestimulation with IFNβ limits RSV proliferation about five-fold, while the relative reduction of IAV proliferation is as significant as in the case of the lower MOI (*cf.* Fig 2A, Fig 2B). Prestimulation with IFNβ triggers an antiviral state manifested by accumulation of ISGs: RIG-I, PKR, and OAS1 and STAT1/2. These proteins maintain their elevated levels 48 hours post-infection (Fig 2C, Fig 2D). Consequently, viral proteins (RSV fusion glycoprotein, IAV nucleoprotein) are less abundant in IFNβ-primed cells than in the naïve cells.

Impact of RSV or IAV infection on STAT1 activation

To analyze at the single-cell level how infected cells induce the antiviral state in bystander cells, we performed infections at MOI 0.1, fixed the cells 24 hours post-infection, and immunostained them for active IRF3 (as an evidence of detection of viral RNA by cell), p-STAT1 (used as an indication of the antiviral state), and viral proteins (RSV fusion glycoprotein or IAV nucleoprotein). In two-channel overlays shown in Fig 3A, Fig 3B, it can be noticed that p-STAT1 is present in most bystander cells (at various levels), whereas the infected cells exhibit p-STAT1 only sporadically. This effect indicates the capability of the viruses to suppress the response to IFN stimulation [45–49]. We quantified this effect by

looking directly at the presence of viral proteins as the definite manifestation of productive infection, and at nuclear p-STAT1 as an indication of antiviral alertness. Based on the expression of RSV fusion glycoprotein or IAV nucleoprotein, we classified cells as either virus-positive or virus-negative and collected histograms of nuclear p-STAT1 intensity in each cell within the two groups. In Fig 3C, Fig 3D we can see that the two distributions are indeed different for virus-positive and virus-negative cells. To express the extent to which the two distributions are disjoint, we calculated the Kolmogorov–Smirnov (KS) statistic. Across all experimental replicates, KS lies in the range 0.42–0.58 (95% CrI: 0.39–0.67) in the case of RSV infection and in the range 0.57–0.59 (95% CrI: 0.51–0.68) in the case of IAV infection.

Effect of prestimulation with IFNβ or IFNλ1 on IAV spread in A549 WT cells

Thus far, we have used cell-population techniques to show that IFNβ is a more potent inducer of the antiviral state than IFNλ1 (Fig 1) and, as a consequence, in A549 cells viral proliferation is more suppressed after prestimulation with IFNβ than with IFNλ1 (Fig 2A). Since RSV turned out to elicit stronger IFNβ response than IAV, to characterize the potential interferon-mediated cross-protection between the two viruses we decided to investigate specifically the dual infection scheme in which the cells are first preinfected with RSV and after 24 hours infected with IAV.

To gain insight into the interferon-mediated interaction at the single-cell level, we quantified the protective effect of IFNβ or IFNλ1 in the context of IAV infection based on imaging data (Fig 4A). In Fig 4B we show proportions of cells expressing viral nucleoprotein 24 hours after treatment with IAV at MOI = 1 in each field of view in the case of naïve (that is, devoid of any prior IFN stimulation) A549 WT cells and A549 WT cells that were prestimulated with either IFNβ or IFNλ1. We estimated that more than 40% non-prestimulated cells express nucleoprotein 24 hours after infection. This proportion drops to about 3% after prestimulation with IFNβ and about 10% after prestimulation with IFNλ1. Relations between these proportions provide additional evidence for strong IAV spread attenuation in A549 cells conferred by IFNs.

RSV preinfection inhibits IAV infection in an interferon-dependent manner

In the course of viral infection there are cells that are initially infected by the primary virus and bystander cells that, in response to IFN secreted by the initially infected cells, may develop the antiviral state and become partially protected from secondary infections. Our prior experiments with IFN prestimulation and high IAV MOI allowed us to characterize the response of the cells that had uniformly developed the antiviral state and were then challenged with the virus, which mimics only one subpopulation of the cells that appears during ongoing infection. Since in our dual infection scheme we wanted to apply the second virus to a cell population that consist of the subpopulation of the preinfected and IFN-producing cells and the subpopulation of not (yet) infected but IFN-stimulated cells, we primed cells with the first virus, RSV, at MOI = 0.1 and after 30 hours applied the second virus, IAV, at MOI = 1. To corroborate and discern the relative importance of IFNβ or IFNλ in their native RSV-elicited mixture, we used additionally A549 IFNAR1 KO, IFNLR1 KO, and IFNAR1–IFNLR1 dKO cell lines.

In Fig 5A and Fig 5B, we show imaging-based single-cell level analysis of the cell population after dual infection. After preinfection with RSV, proportion of WT A549 cells infected with the second virus, IAV, is reduced by an order of magnitude with respect to non-preinfected cells (*cf*. naïve WT cells in Fig 4B and RSV-primed WT cells in Fig 5B, which gives a tenfold reduction of IAV nucleoprotein-expressing cells, from ~40% to ~4%). The protective effect conferred by RSV-priming is significantly weaker in IFNAR1 KO cells incapable of responding to IFNβ (reduction from ~40% to ~20%) and only slightly weaker in IFNLR1 cells incapable of responding to IFNλ (reduction from ~40% to ~10%). The priming effect is virtually non-existent in cells devoid of receptors for both IFNβ and IFNλ. In Fig 5C we show dPCR results of corresponding experiments that are in qualitative agreement with results of single-cell analysis in Fig 5B. Overall, this demonstrates that preinfection with RSV protects from infection with IAV in an interferon-dependent manner in A549 cells. The protective effect may be attributed nearly entirely to the paracrine stimulation with both IFNβ and IFNλ, with a major contribution from IFNβ.

IAV infects preferentially the cells preinfected by and expressing RSV

Visual inspection of WT cells suggested that IAV nucleoprotein is preferentially expressed in the cells with apparent expression of RSV fusion glycoprotein (see Fig 6A). To quantitatively verify this hypothesis, as in Fig 3D, we classified cells as virus-positive and virus-negative. We decided to use the measured level of IAV NP as the binary criterion because of its mostly nuclear location, which facilitates image analysis. We then collected histograms of intensity of RSV fusion glycoprotein in the cells belonging to these two groups. Clearly, in A549 WT cells, the IAV+ cells are associated with higher abundance of RSV fusion glycoprotein than IAV− cells (KS of 0.62 indicates good discernibility). This effect is much weaker in IFNAR1–IFNLR1 dKO cells, which cannot respond to neither IFNβ nor IFNλ (KS of 0.15 indicates that the two distributions are nearly overlapping). This clearly shows that the RSV–IAV exclusion is not realized through a direct competition for a shared ecological niche, a single cell, but rather achieved with the involvement of interferon signaling.

Discussion

To replicate, an invading pathogen has to evade or compromise the innate immunity. When challenged by the infectant, the innate immune system acts to impede pathogen replication before a more specific adaptive response is evoked. In the context of sequential viral coinfections, the priming virus both breaks through the innate protection in infected cells and, indirectly, via interferon secreted by these cells, triggers the protective antiviral state in bystander cells. These two antagonistic processes determine the fate of both the primary and the secondary infection at the cell-population level.

To obtain insight into this interplay at the single-cell level, we studied how upon a priming infection with RSV, individual cells become either cross-protected or, conversely, more vulnerable to the second virus, IAV. The choice of these two pathogens has been motivated by the fact that RSV and influenza viruses show relatively high coincidence [50]. The particular order of infections that we focused on results from our observations that (1) RSV induces higher IFNβ production than IAV, (2) IFNβ-stimulated cells are more resistant to infection with IAV than to infection with RSV, and (3) IAV is sensitive to IFNλ and this sensitivity does not rely solely on STAT1/2 signaling. We thus inferred that the potential

cross-protection may be more pronounced in the RSV-then-IAV protocol, which is consistent with a comparison of RSV-then-IAV and IAV-then-RSV coinfections in a mouse model, showing that the IAV-then-RSV coinfection is associated with higher IAV loads and mouse mortality [51] (see also [52]).

We found that IFNβ is the main inducer of the STAT1/2-associated antiviral state in RSV-infected A549 cells and both interferons, IFNβ and IFNλ, are simultaneously necessary for building maximum protection against a subsequent infection with IAV. A synergistic effect of type I and type III interferon, consistent with our findings, has been previously reported in mice by Mordstein *et al.* [53]. In our coinfection experiments, we infected the epithelial alveolar cells with RSV at MOI of 0.1 to readily observe their stratification into the compromised infected cells and the reinforced bystander cells. Immunostaining revealed that preinfection with RSV partitions the cell population into a subpopulation susceptible to a subsequent infection with IAV and an IAV-proof subpopulation. This dynamic functional differentiation of cells may be amplified *in vivo* by differential propensity of specific cell types to permit or restrict viral replication, produce interferons, and build up the antiviral state [54]. We found that the susceptible cells turned out to be predominantly those already compromised and efficiently expressing the priming virus. This means that after preinfection the population of cells susceptible to a secondary infectant is largely limited to those yielding preinfectant progeny, proportion of which depends on the multiplicity of the preinfection and time interval between infections. The optimal cross protection effect may thus ensue when the proportion of cells preinfected with RSV is relatively low yet sufficient to trigger a pervasive antiviral state in bystander cells. In the case of massive infection with RSV, when most cells are simultaneously compromised by the virus, the interferon-mediated protection likely does not come into effect.

The natural question is to what extent the described cross-protection mechanism reduces the incidence of coinfections or severity of the disease featured by a secondary infection *in vivo*. Answering the question is especially important, but appears complex, in the context of secondary SARS-CoV-2 infections. An early study by Ziegler *et al.* [55] showed that the receptor for SARS-CoV-2, ACE2, is a protein coded by ISG, but later Onabajo *et al.* [56] identified the transcriptionally independent truncated isoform of *ACE2, deltaACE2,* not *ACE2,* as an ISG. Since deltaACE2 may not bind SARS-CoV-2 spike protein, the latter

study suggests that interferon signaling is not hijacked by this virus to enhance its proliferation. Several studies suggested beneficial roles of type I and type III interferon at the early stage of infection [57]. It is thus possible that interferons secreted in response to a primary infection exhibit a protective role against a secondary infection with SARS-CoV-2 (see also [58]).

At the clinical level, one may expect that the occurrence of coinfections may result from defective work of the immune system (which aggravates the course of the disease and also imaginably cannot mediate antagonistic viral interactions). Although this effect cannot be excluded and coinfection data should be treated with caution, impact of coinfections on disease severity is unclear. Most respiratory viral coinfections, detected usually in a small proportion of patients, appear not to increase severity of clinical outcomes [59–62], with two noteworthy exceptions of a coinfection by IAV and IBV [50] and coinfections with RSV in paediatric patients [63]. It should be noted that interactions of two respiratory viruses within the host can hardly be translated to their coincidence or avoidance in human populations. This is because at the population level there are two additional and antagonistic effects: on one hand, viral coincidence of contagious diseases can be seasonal (the effective contact rate is modulated by weather and behavioral patterns); on the other hand, infected and ill individuals change their behavior to avoid contact. A significant behavioral reduction of the contact rate may result from an outbreak of an epidemic of a threatening virus, decreasing the risk of propagation of other viruses, as observed during the COVID-19 pandemic in 2020 [64,65].

In conclusion, our study demonstrates that the infection with RSV cross-protects alveolar epithelial cells against infection with IAV. Whereas RSV-infected cells are more vulnerable to infection with IAV, priming with RSV indirectly protects bystander non-infected cells from IAV. The cross-protection mechanism relies on induction of and paracrine stimulation with both IFNβ and IFNλ.

Materials and Methods

Cell lines and cultures

The A549 and MDCK cell lines were purchased from ATTC. A549 cells were cultured in F12K basal medium supplemented with 10% fetal bovine serum (FBS) and penicillin/streptomycin antibiotic solution. MDCK cells and HEK-293T cells were maintained in DMEM medium supplemented with 10% FBS and penicillin/streptomycin antibiotic solution. All cell lines were cultured in standard conditions (37^0C, 5% CO_2) and kept in monolayers up to 90% confluency. The IFNAR1, IFNLR1, and IFNAR1–IFNLR1 knockouts based on A549 were engendered using the CRISPR lentivirus system (Abm). To obtain A549-based knockouts, HEK-293T cells were co-transfected with commercially available plasmids encoding Cas9 nuclease (Abm, cat. K002), lentivirus packaging particles (Abm, cat. LV053) and sgRNA targeted against IFNAR1 (Abm, cat. 2426411) or IFNLR1 (Abm, cat. 2487611). The double IFNAR1/IFNLR1 mutant was created based on IFNAR1 KO cells. After 2 days post-transfection, the media with lentiviral particles were collected, enriched with 8 µg/ml polybrene and filtered through 0.45 µm syringe filters. Then each individual lentiviral supernatant was used to transduce A549 wild type cells, which were subcultured at low confluency (seeding density 5×10^4 cells per 30 mm dish). After another 2 days, A549 cells were subjected to selection with 800 µg/ml of G418 for 10 days and seeded into a 96-well plate to obtain single cell colonies. Clones were validated by Western blot or immunostaining analysis of pStat1 activation in response to IFNβ or IFNλ. Additionally, the selected KO clones were verified by sequencing.

Virus amplification and isolation

Respiratory Syncytial Virus A2 strain and Influenza A virus H1N1, strain A/PR/8/34, were purchased from ATCC and amplified in HeLa or MDCK cells, respectively. Cells were seeded on a 225 cm^2 Tissue Culture Flasks (Falcon) and cultured as described above for 2-3 days until reaching 90% confluency. On the day of infection, virus growth medium was prepared: DMEM with 2% FBS for RSV or MEM basal medium with 0.3% BSA and 1 ug/ml of TPCK-Trypsin for IAV. The dilutions of virus were prepared in an appropriate media, with target MOI around 0.05. Culture media were removed, cells were washed once with PBS,

and overlaid with 10 ml of inoculum. Virus was allowed to adsorb to cells for 2 hours at 37°C with occasional stirring. Then, additional virus growth medium was added to a total volume of 40 ml per flask. Infected cells were cultured at 37°C until the development of cytopathic effects could be observed in at least 80% of cells (typically around 3 days for IAV and 5 days for RSV). Virus-containing culture fluid was then collected and clarified by centrifugation at 3000 g, 4°C, for 20 min. Then, virus particles were precipitated by adding 50% (w/v) PEG6000 (Sigma-Aldrich) in NT buffer (150 mM NaCl, 50 mM Tris-HCl, pH 7.5) to a final concentration of 10% and stirring gently at 4°C for 90 min. Virus was centrifuged at 3250 g, 4°C, for 20 min and re-centrifuged after removing supernatant to remove the remaining fluid. Pellet was suspended in 1 ml of NT buffer or in 20% sucrose in NT buffer in case of RSV, aliquoted and stored at −80°C.

Virus quantification

Virus concentration in collected samples was quantified using immunofluorescence protocol. HeLa or MDCK cells were seeded on microscopic cover slips and cultured upon reaching 90-100% confluency. Serial dilutions of virus samples were made in virus growth medium in a 10^{-3} to 10^{-6} range. After washing with PBS, cells were overlaid in duplicates with diluted virus, which was allowed to adhere for 2 hours with occasional stirring. Afterwards, virus-containing medium was removed, cells were overlaid with fresh virus growth medium and cultured for 16 (for IAV) or 24 hours (for RSV). Then, cells were washed with PBS and fixed with 4% formaldehyde for 20 min, RT. Cells were stained using standard immunofluorescence protocol with anti-RSV fusion glycoprotein antibody (Abcam, cat. ab43812) or anti-influenza A virus nucleoprotein antibody [C43] (Abcam, cat. ab128193). Cells containing stained viral proteins were counted using Leica SP5 confocal microscope. Virus concentration was calculated using the following formula: (avg. number of infected cells)/(dilution factor × volume containing virus added) = infectious particles/ml.

Compounds and stimulation protocols

Human interferon β1A and interferon λ1 were purchased from Thermo Fisher Scientific (cat. numbers PHC4244 and 34-8298-64, respectively) and prepared according to manufacturer's instructions. For cell stimulation, interferons were further diluted to desired concentrations in

the F12K medium supplemented with 2% FBS. Lowered FBS content was used to prevent inhibition of viral attachment and entry at the second stage of experiments.

For interferon + virus experiments, the cell culture media were exchanged for interferon-containing or control media at time = 0 and were not removed afterwards till the end of experiment. Appropriately diluted virus was added in small volumes (less than 10 µl) directly into the wells. Even distribution of virus across cell population was aided by intermittent rocking of the plate for 2 hours.

Similarly, for coinfection experiments, the F12K 2% FBS medium containing the first virus (RSV) were added to the cells at the start of experiment, while the dilutions of second virus (IAV) were added directly into the wells at a later time point and distributed by rocking.

Antibodies

Antibodies for Western blotting

Primary: anti-phospho-STAT1 (Tyr 701) (58D6) (Cell Signaling Technologies, cat. 9167, 1:1000); anti-phospho-Stat2 (Tyr 690) (D3P2P) (Cell Signaling Technologies, cat. 88410, 1:1000); anti-RIG-I (D14G6) (Cell Signaling Technologies, cat. 3743, 1:1000); anti-STAT1 (BD Biosciences, cat. 610116, 1:1000), anti-STAT2 Antibody (R&D Systems, cat. PAF-ST2, 1:1000), anti-PKR (B-10) (Santa Cruz Biotechnology, cat. sc-6282, 1:1000), anti-OAS1 Antibody (F-3) (Santa Cruz Biotechnology, cat. sc-374656, 1:1000), anti-Influenza A Virus Nucleoprotein antibody [C43] (Abcam, cat. ab128193, 1:1000), anti-Respiratory Syncytial Virus antibody [2F7] (Abcam, cat. ab43812, 1:1000); hFAB Rhodamine anti-GAPDH primary antibody (Bio-Rad, cat. 12004168, 1:10,000).

Secondary: goat anti-Rabbit IgG (H+L) Secondary Antibody, DyLight 800 4X PEG (Thermo Fisher Scientific, cat. SA5-35571, 1:10,000); goat anti-Mouse IgG (H+L) Secondary Antibody, DyLight 800 4X PEG (Thermo Fisher Scientific, cat. SA5-35521, 1:10,000); StarBright Blue 700 Goat Anti-Mouse IgG (Bio-Rad, cat. 12004159, 1:10,000), rabbit anti-goat immunoglobulins/HRP (Agilent, cat. P0449, 1:10,000).

Antibodies for immunostaining

Primary: anti-phospho-STAT1 Tyr 701 (58D6) (Cell Signaling Technologies, cat. 9167, 1:1000); anti-IRF-3 (D-3) (Santa Cruz Biotechnology, cat. sc-376455, 1:500), anti-Respiratory Syncytial Virus antibody (Abcam, cat. ab20745, 1:1000), anti-Influenza A virus (Abcam, cat. ab20841, 1:1000), anti-Influenza A Virus Nucleoprotein antibody [C43] (Abcam, cat. ab128193, 1:1000).

Secondary: donkey anti-rabbit IgG (H+L), Alexa Fluor 488 conjugate (Thermo Fisher Scientific, cat. A-21206, 1:1000), donkey anti-mouse IgG (H+L), Alexa Fluor 555 conjugate (Thermo Fisher Scientific, cat. A-31570, 1:1000), donkey anti-Goat IgG (H+L) Alexa Fluor 633 conjugate (Thermo Fisher Scientific, cat. A-21082 1:1000).

ELISA

For measuring IFNβ production, cells were seeded on 96-well plates (Falcon) at a density of 20,000 cells per well and infected with RSV or IAV as described above. Culture medium (a total of 200 µl) from infected cells was collected at designated time points and stored at −20°C until further analysis. IFNβ levels were estimated using the VeriKine Mouse IFNβ ELISA kit (PBL Assay Science). Standards and diluted samples in duplicates or triplicates were added to a precoated plate included in the kit and incubated for 1 h. After subsequent washing, antibody solution was prepared and added to wells for another 1 h, followed by another washing and 1 h incubation with HRP solution. Finally, TMB Substrate Solution was added to wells and developing color reaction was stopped after 15 min with the addition of Stop Solution. Optical densities of samples after resulting color development were determined using Multiskan GO plate reader (Thermo Fisher Scientific) set to 450 nm, with wavelength correction at 570 nm. IFNβ concentrations were obtained based on the standard curve (4-parameter logistic function fitted to 7 data points from standards).

Digital PCR (dPCR)

For gene expression experiments, cells were seeded on 24-well plates at a density of 100 000 cells per well. RNA from virus-infected cells was isolated using PureLink RNA Mini Kit (Thermo Fisher Scientific), following manufacturer's instructions: cells were harvested and vortexed in Lysis Buffer with 2-mercaptoethanol and then vortexed again with one volume of

70% ethanol. Upon transferring to the spin cartridge, cellular RNA content was bound to the column, washed with appropriate buffers and eluted, all by centrifugation at 12,000g. Eluted RNA in RNase-free water was used immediately for reverse transcription or stored for later use at −80°C. Concentration and quality of isolated total RNA was determined by measuring UV absorbance at 260 nm and at 280 nm using Multiskan GO spectrophotometer (Thermo Fisher Scientific). Around 0.5 µg of RNA was used as a template for reverse transcription, performed using High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher Scientific): diluted RNA samples were mixed 1:1 with freshly prepared Master Mix containing RT Buffer, RT Random Primers, dNTP Mix, MultiScribe Reverse Transcriptase and RNase Inhibitor. Reaction was performed in Mastercycle Gradient thermal cycler (Eppendorf) under following conditions: 10 min at 25°C, 120 min at 37°C, and 5 min at 85°C.

Measurements of the viral RNA copy number was then performed using QuantStudio 3D system (Thermo Fisher Scientific) and TaqMan Gene Expression Assays Vi99990011_po and Vi99990014_po for quantification of, respectively, IAV and RSV RNA. Appropriately diluted samples were mixed with the reaction master mix and the chosen assay, loaded on the digital PCR chips in duplicates and thermocycled using ProFlex Flat Block Thermal Cycler. Subsequently, chips were analyzed with QuantStudio 3D Digital PCR Instrument. Chip quality check and final data analysis were performed using QuantStudio 3D AnalysisSuite software.

Western blotting

Cells at the indicated time points were washed twice with PBS, lysed in a Laemmli sample buffer containing DTT and boiled at 95°C for 10 min. Even amounts of each protein sample were separated on a 4–16% Mini-PROTEAN TGX Stain-Free Precast Gels using Mini-PROTEAN Tetra Cell electrophoresis system (Bio-Rad). Upon completion of electrophoresis, proteins were transferred to the nitrocellulose membrane using wet electrotransfer in the Mini-PROTEAN apparatus (400 mA, 50 min). Membrane was rinsed with TBST (TBS buffer containing 0.1% Tween-20, Sigma-Aldrich) and blocked for 1h with 5% BSA/TBS or 5% non-fat dry milk. Subsequently, membranes were incubated at 4°C overnight with one of the primary antibodies diluted in a 5% BSA/TBS buffer. After thorough washing with TBST, membranes were incubated with secondary antibodies conjugated with

specific fluorochrome or horseradish peroxidase diluted in 5% non-fat dry milk for 1 h, RT. Chemiluminescent reaction was developed with the Clarity Western ECL system (Bio-Rad). For GAPDH detection, hFAB Rhodamine Anti-GAPDH Primary Antibody (Bio-Rad) was used. Specific protein bands were detected using the ChemiDoc MP Imaging System (Bio-Rad).

Immunostaining

For staining of intracellular proteins, cells were seeded on 12 mm round glass coverslips, which were previously washed in 60% ethanol/40% HCl, thoroughly rinsed with water and sterilized. After stimulation with interferon or viral infection at desired time points, cells on coverslips were washed with PBS and immediately fixed with 4% formaldehyde (20 min, RT). Cells were then washed three times with PBS and incubated for 20 min at −20°C with 100% cold methanol. After washing with PBS, coverslips with cells were blocked and permeabilized for 1.5 hour with 5% BSA (Sigma-Aldrich) with 0.3% Triton X-100 (Sigma-Aldrich) in PBS at room temperature. After removing the blocking solution, coverslips with cells were incubated overnight at 4°C with primary antibodies diluted in PBS containing 1% BSA and 0.3% Triton X-100. After washing cells five times with PBS, appropriate secondary antibodies conjugated with fluorescent dyes were added and incubated for 1 hour at room temperature. Subsequently, cells were washed and their nuclei were stained for 10 min with 200 ng/ml DAPI (Sigma-Aldrich). After final washing in miliQ water, coverslips with stained cells were mounted on microscope slides with a drop of Mowiol (Sigma-Aldrich). Cellular sublocalization of stained proteins was observed using Leica SP5 confocal microscope and Leica Application Suite AF software.

Data analysis

Image analysis

Confocal images obtained from immunostaining were analyzed using our in-house software (https://pmbm.ippt.pan.pl/software/shuttletracker). Nuclear outlines were detected based on DAPI staining. Outlines of nuclei that were partially out of frame or mitotic were excluded from analysis; outlines of overlapping nuclei were split based on geometric convexity defects when possible. Cells were classified as either RSV-negative or RSV-positive based on the

combination of the sum of intensities of pixels inside nuclear outlines (weight 5) and the perinuclear region (weight 1, decreasing sigmoidally with the distance from the nuclear contour). Cells were classified as either IAV-negative or IAV-positive based on the sum of intensities of pixels inside nuclear outlines.

Credible interval

Credible interval (CrI) of the Kolmogorov–Smirnov statistics (Fig 3C, Fig 3D, Fig 6B) has been estimated by resampling data points 10^5 times. In each resampling, data points were drawn at random with probabilities proportional to the number of data points in each category (virus-positive, virus-negative) and the number of resampled data points was equal to the total number of data points in both categories.

References

1. Hu B, Guo H, Zhou P, Shi Z-L. Characteristics of SARS-CoV-2 and COVID-19. Nat Rev Microbiol. 2021;19: 141–154. doi:10.1038/s41579-020-00459-7

2. Crowe JE. Human Respiratory Viruses. Reference Module in Biomedical Sciences. 2014; B978-0-12-801238-3.02600-3. doi:10.1016/B978-0-12-801238-3.02600-3

3. Bruyndonckx R, Coenen S, Butler C, Verheij T, Little P, Hens N, et al. Respiratory syncytial virus and influenza virus infection in adult primary care patients: Association of age with prevalence, diagnostic features and illness course. Int J Infect Dis. 2020;95: 384–390. doi:10.1016/j.ijid.2020.04.020

4. Falsey AR, Hennessey PA, Formica MA, Cox C, Walsh EE. Respiratory syncytial virus infection in elderly and high-risk adults. N Engl J Med. 2005;352: 1749–1759. doi:10.1056/NEJMoa043951

5. Scheltema NM, Gentile A, Lucion F, Nokes DJ, Munywoki PK, Madhi SA, et al. Global respiratory syncytial virus-associated mortality in young children (RSV GOLD): a retrospective case series. Lancet Glob Health. 2017;5: e984–e991. doi:10.1016/S2214-109X(17)30344-3

6. Casalegno JS, Ottmann M, Bouscambert-Duchamp M, Valette M, Morfin F, Lina B. Impact of the 2009 influenza A(H1N1) pandemic wave on the pattern of hibernal respiratory virus epidemics, France, 2009. Euro Surveill. 2010;15: 19485.

7. Shrestha S, King AA, Rohani P. Statistical inference for multi-pathogen systems. PLoS Comput Biol. 2011;7: e1002135. doi:10.1371/journal.pcbi.1002135

8. Míguez A, Iftimi A, Montes F. Temporal association between the influenza virus and respiratory syncytial virus (RSV): RSV as a predictor of seasonal influenza. Epidemiol Infect. 2016;144: 2621–2632. doi:10.1017/S095026881600090X

9. Nickbakhsh S, Mair C, Matthews L, Reeve R, Johnson PCD, Thorburn F, et al. Virus-virus interactions impact the population dynamics of influenza and the common cold. Proc Natl Acad Sci U S A. 2019; 201911083. doi:10.1073/pnas.1911083116

10. DaPalma T, Doonan BP, Trager NM, Kasman LM. A systematic approach to virus-virus interactions. Virus Res. 2010;149: 1–9. doi:10.1016/j.virusres.2010.01.002

11. Andrewes CH. Interference by One Virus with the Growth of Another in Tissue-Culture. Br J Exp Pathol. 1942;23: 214–220.

12. Greer RM, McErlean P, Arden KE, Faux CE, Nitsche A, Lambert SB, et al. Do rhinoviruses reduce the probability of viral co-detection during acute respiratory tract infections? J Clin Virol. 2009;45: 10–15. doi:10.1016/j.jcv.2009.03.008

13. Tao KP, Chong M, Pun JC, Tsun JG, Chow SM, Ng CS, et al. Suppression of influenza virus infection by rhinovirus interference at the population, individual and cellular levels. 2021 Aug p. 2021.08.09.21256656. doi:10.1101/2021.08.09.21256656

14. Chan KF, Carolan LA, Korenkov D, Druce J, McCaw J, Reading PC, et al. Investigating Viral Interference Between Influenza A Virus and Human Respiratory Syncytial Virus in a Ferret Model of Infection. J Infect Dis. 2018;218: 406–417. doi:10.1093/infdis/jiy184

15. Gonzalez AJ, Ijezie EC, Balemba OB, Miura TA. Attenuation of Influenza A Virus Disease Severity by Viral Coinfection in a Mouse Model. J Virol. 2018;92: e00881-18. doi:10.1128/JVI.00881-18

16. Wu A, Mihaylova VT, Landry ML, Foxman EF. Interference between rhinovirus and influenza A virus: a clinical data analysis and experimental infection study. Lancet Microbe. 2020;1: e254–e262. doi:10.1016/s2666-5247(20)30114-2

17. Van Leuven JT, Gonzalez AJ, Ijezie EC, Wixom AQ, Clary JL, Naranjo MN, et al. Rhinovirus Reduces the Severity of Subsequent Respiratory Viral Infections by Interferon-Dependent and -Independent Mechanisms. mSphere. 2021;6: e0047921. doi:10.1128/mSphere.00479-21

18. Lindenmann J. From interference to interferon: a brief historical introduction. Philos Trans R Soc Lond B Biol Sci. 1982;299: 3–6. doi:10.1098/rstb.1982.0101

19. Gardner PS, McGuckin R, Beale A, Fernandes R. Interferon and respiratory syncytial virus. Lancet. 1970;1: 574–575. doi:10.1016/s0140-6736(70)90821-4

20. Hall CB, Douglas RG, Simons RL, Geiman JM. Interferon production in children with respiratory syncytial, influenza, and parainfluenza virus infections. J Pediatr. 1978;93: 28–32. doi:10.1016/s0022-3476(78)80594-0

21. Roberts NJ, Hiscott J, Signs DJ. The limited role of the human interferon system response to respiratory syncytial virus challenge: analysis and comparison to influenza virus challenge. Microb Pathog. 1992;12: 409–414. doi:10.1016/0882-4010(92)90003-7

22. Goritzka M, Makris S, Kausar F, Durant LR, Pereira C, Kumagai Y, et al. Alveolar macrophage-derived type I interferons orchestrate innate immunity to RSV through recruitment of antiviral monocytes. J Exp Med. 2015;212: 699–714. doi:10.1084/jem.20140825

23. Chu C-Y, Qiu X, McCall MN, Wang L, Corbett A, Holden-Wiltse J, et al. Insufficiency in airway interferon activation defines clinical severity to infant RSV infection. 2019 May p. 641795. doi:10.1101/641795

24. Hijano DR, Vu LD, Kauvar LM, Tripp RA, Polack FP, Cormier SA. Role of Type I Interferon (IFN) in the Respiratory Syncytial Virus (RSV) Immune Response and Disease Severity. Front Immunol. 2019;10: 566. doi:10.3389/fimmu.2019.00566

25. Villenave R, Broadbent L, Douglas I, Lyons JD, Coyle PV, Teng MN, et al. Induction and Antagonism of Antiviral Responses in Respiratory Syncytial Virus-Infected Pediatric Airway Epithelium. J Virol. 2015;89: 12309–12318. doi:10.1128/JVI.02119-15

26. Kotenko SV, Gallagher G, Baurin VV, Lewis-Antes A, Shen M, Shah NK, et al. IFN-lambdas mediate antiviral protection through a distinct class II cytokine receptor complex. Nat Immunol. 2003;4: 69–77. doi:10.1038/ni875

27. Sheppard P, Kindsvogel W, Xu W, Henderson K, Schlutsmeyer S, Whitmore TE, et al. IL-28, IL-29 and their class II cytokine receptor IL-28R. Nat Immunol. 2003;4: 63–68. doi:10.1038/ni873

28. Czerkies M, Korwek Z, Prus W, Kochańczyk M, Jaruszewicz-Błońska J, Tudelska K, et al. Cell fate in antiviral response arises in the crosstalk of IRF, NF-κB and JAK/STAT pathways. Nat Commun. 2018;9: 493. doi:10.1038/s41467-017-02640-8

29. Lazear HM, Schoggins JW, Diamond MS. Shared and Distinct Functions of Type I and Type III Interferons. Immunity. 2019;50: 907–923. doi:10.1016/j.immuni.2019.03.025

30. Forero A, Ozarkar S, Li H, Lee CH, Hemann EA, Nadjsombati MS, et al. Differential Activation of the Transcription Factor IRF1 Underlies the Distinct Immune Responses

Elicited by Type I and Type III Interferons. Immunity. 2019;51: 451-464.e6. doi:10.1016/j.immuni.2019.07.007

31. Schoggins JW. Interferon-Stimulated Genes: What Do They All Do? Annu Rev Virol. 2019;6: 567–584. doi:10.1146/annurev-virology-092818-015756

32. Seo S-U, Kwon H-J, Ko H-J, Byun Y-H, Seong BL, Uematsu S, et al. Type I interferon signaling regulates Ly6C(hi) monocytes and neutrophils during acute viral pneumonia in mice. PLoS Pathog. 2011;7: e1001304. doi:10.1371/journal.ppat.1001304

33. Kotenko SV, Durbin JE. Contribution of type III interferons to antiviral immunity: location, location, location. J Biol Chem. 2017;292: 7295–7303. doi:10.1074/jbc.R117.777102

34. Klinkhammer J, Schnepf D, Ye L, Schwaderlapp M, Gad HH, Hartmann R, et al. IFN-λ prevents influenza virus spread from the upper airways to the lungs and limits virus transmission. Elife. 2018;7: e33354. doi:10.7554/eLife.33354

35. Broggi A, Ghosh S, Sposito B, Spreafico R, Balzarini F, Lo Cascio A, et al. Type III interferons disrupt the lung epithelial barrier upon viral recognition. Science. 2020;369: 706–712. doi:10.1126/science.abc3545

36. Major J, Crotta S, Llorian M, McCabe TM, Gad HH, Priestnall SL, et al. Type I and III interferons disrupt lung epithelial repair during recovery from viral infection. Science. 2020;369: 712–717. doi:10.1126/science.abc2061

37. Mordstein M, Kochs G, Dumoutier L, Renauld J-C, Paludan SR, Klucher K, et al. Interferon-lambda contributes to innate immunity of mice against influenza A virus but not against hepatotropic viruses. PLoS Pathog. 2008;4: e1000151. doi:10.1371/journal.ppat.1000151

38. Essaidi-Laziosi M, Geiser J, Huang S, Constant S, Kaiser L, Tapparel C. Interferon-Dependent and Respiratory Virus-Specific Interference in Dual Infections of Airway Epithelia. Sci Rep. 2020;10: 10246. doi:10.1038/s41598-020-66748-6

39. Davidson S, McCabe TM, Crotta S, Gad HH, Hessel EM, Beinke S, et al. IFNλ is a potent anti-influenza therapeutic without the inflammatory side effects of IFNα treatment. EMBO Mol Med. 2016;8: 1099–1112. doi:10.15252/emmm.201606413

40. Lozhkov AA, Klotchenko SA, Ramsay ES, Moshkoff HD, Moshkoff DA, Vasin AV, et al. The Key Roles of Interferon Lambda in Human Molecular Defense against Respiratory Viral Infections. Pathogens. 2020;9: E989. doi:10.3390/pathogens9120989

41. Liu P, Jamaluddin M, Li K, Garofalo RP, Casola A, Brasier AR. Retinoic acid-inducible gene I mediates early antiviral response and Toll-like receptor 3 expression in

respiratory syncytial virus-infected airway epithelial cells. J Virol. 2007;81: 1401–1411. doi:10.1128/JVI.01740-06

42. Liu G, Park H-S, Pyo H-M, Liu Q, Zhou Y. Influenza A Virus Panhandle Structure Is Directly Involved in RIG-I Activation and Interferon Induction. J Virol. 2015;89: 6067–6079. doi:10.1128/JVI.00232-15

43. Meurs E, Chong K, Galabru J, Thomas NS, Kerr IM, Williams BR, et al. Molecular cloning and characterization of the human double-stranded RNA-activated protein kinase induced by interferon. Cell. 1990;62: 379–390. doi:10.1016/0092-8674(90)90374-n

44. Silverman RH. Viral encounters with 2',5'-oligoadenylate synthetase and RNase L during the interferon antiviral response. J Virol. 2007;81: 12720–12729. doi:10.1128/JVI.01471-07

45. Ramaswamy M, Shi L, Varga SM, Barik S, Behlke MA, Look DC. Respiratory syncytial virus nonstructural protein 2 specifically inhibits type I interferon signal transduction. Virology. 2006;344: 328–339. doi:10.1016/j.virol.2005.09.009

46. Oshansky CM, Krunkosky TM, Barber J, Jones LP, Tripp RA. Respiratory syncytial virus proteins modulate suppressors of cytokine signaling 1 and 3 and the type I interferon response to infection by a toll-like receptor pathway. Viral Immunol. 2009;22: 147–161. doi:10.1089/vim.2008.0098

47. Wei H, Wang S, Chen Q, Chen Y, Chi X, Zhang L, et al. Suppression of interferon lambda signaling by SOCS-1 results in their excessive production during influenza virus infection. PLoS Pathog. 2014;10: e1003845. doi:10.1371/journal.ppat.1003845

48. Wei H, Wang S, Chen Q, Chen Y, Chi X, Zhang L, et al. Correction: Suppression of Interferon Lambda Signaling by SOCS-1 Results in Their Excessive Production during Influenza Virus Infection. PLoS Pathog. 2016;12: e1005402. doi:10.1371/journal.ppat.1005402

49. Du Y, Yang F, Wang Q, Xu N, Xie Y, Chen S, et al. Influenza a virus antagonizes type I and type II interferon responses via SOCS1-dependent ubiquitination and degradation of JAK1. Virol J. 2020;17: 74. doi:10.1186/s12985-020-01348-4

50. Goka E, Vallely P, Mutton K, Klapper P. Influenza A viruses dual and multiple infections with other respiratory viruses and risk of hospitalisation and mortality. Influenza Other Respir Viruses. 2013;7: 1079–1087. doi:10.1111/irv.12020

51.	George JA, AlShamsi SH, Alhammadi MH, Alsuwaidi AR. Exacerbation of Influenza A Virus Disease Severity by Respiratory Syncytial Virus Co-Infection in a Mouse Model. Viruses. 2021;13: 1630. doi:10.3390/v13081630

52.	Chen HD, Fraire AE, Joris I, Welsh RM, Selin LK. Specific history of heterologous virus infections determines anti-viral immunity and immunopathology in the lung. Am J Pathol. 2003;163: 1341–1355. doi:10.1016/S0002-9440(10)63493-1

53.	Mordstein M, Neugebauer E, Ditt V, Jessen B, Rieger T, Falcone V, et al. Lambda interferon renders epithelial cells of the respiratory and gastrointestinal tracts resistant to viral infections. J Virol. 2010;84: 5670–5677. doi:10.1128/JVI.00272-10

54.	Hillyer P, Shepard R, Uehling M, Krenz M, Sheikh F, Thayer KR, et al. Differential Responses by Human Respiratory Epithelial Cell Lines to Respiratory Syncytial Virus Reflect Distinct Patterns of Infection Control. J Virol. 2018;92: e02202-17. doi:10.1128/JVI.02202-17

55.	Ziegler CGK, Allon SJ, Nyquist SK, Mbano IM, Miao VN, Tzouanas CN, et al. SARS-CoV-2 Receptor ACE2 Is an Interferon-Stimulated Gene in Human Airway Epithelial Cells and Is Detected in Specific Cell Subsets across Tissues. Cell. 2020;181: 1016-1035.e19. doi:10.1016/j.cell.2020.04.035

56.	Onabajo OO, Banday AR, Stanifer ML, Yan W, Obajemu A, Santer DM, et al. Interferons and viruses induce a novel truncated ACE2 isoform and not the full-length SARS-CoV-2 receptor. Nat Genet. 2020;52: 1283–1293. doi:10.1038/s41588-020-00731-9

57.	Sa Ribero M, Jouvenet N, Dreux M, Nisole S. Interplay between SARS-CoV-2 and the type I interferon response. PLoS Pathog. 2020;16: e1008737. doi:10.1371/journal.ppat.1008737

58.	Dee K, Goldfarb DM, Haney J, Amat JAR, Herder V, Stewart M, et al. Human Rhinovirus Infection Blocks Severe Acute Respiratory Syndrome Coronavirus 2 Replication Within the Respiratory Epithelium: Implications for COVID-19 Epidemiology. J Infect Dis. 2021;224: 31–38. doi:10.1093/infdis/jiab147

59.	Nishimura N, Nishio H, Lee MJ, Uemura K. The clinical features of respiratory syncytial virus: lower respiratory tract infection after upper respiratory tract infection due to influenza virus. Pediatr Int. 2005;47: 412–416. doi:10.1111/j.1442-200x.2005.02099.x

60.	Martin ET, Kuypers J, Wald A, Englund JA. Multiple versus single virus respiratory infections: viral load and clinical disease severity in hospitalized children. Influenza Other Respir Viruses. 2012;6: 71–77. doi:10.1111/j.1750-2659.2011.00265.x

61. Scotta MC, Chakr VCBG, de Moura A, Becker RG, de Souza APD, Jones MH, et al. Respiratory viral coinfection and disease severity in children: A systematic review and meta-analysis. J Clin Virol. 2016;80: 45–56. doi:10.1016/j.jcv.2016.04.019

62. Nolan VG, Arnold SR, Bramley AM, Ampofo K, Williams DJ, Grijalva CG, et al. Etiology and Impact of Coinfections in Children Hospitalized With Community-Acquired Pneumonia. J Infect Dis. 2018;218: 179–188. doi:10.1093/infdis/jix641

63. Aberle JH, Aberle SW, Pracher E, Hutter H-P, Kundi M, Popow-Kraupp T. Single versus dual respiratory virus infections in hospitalized infants: impact on clinical course of disease and interferon-gamma response. Pediatr Infect Dis J. 2005;24: 605–610. doi:10.1097/01.inf.0000168741.59747.2d

64. Jones N. How COVID-19 is changing the cold and flu season. Nature. 2020;588: 388–390. doi:10.1038/d41586-020-03519-3

65. Chan CP, Wong NS, Leung CC, Lee SS. Positive impact of measures against COVID-19 on reducing influenza in the Northern Hemisphere. J Travel Med. 2020;27: taaa087. doi:10.1093/jtm/taaa087

Figures

Figure 1. IFNβ and IFNλ mediate STAT1 and STAT2 activation upon RSV and IAV infection in A549 cells.

(**A**, **B**) ELISA for IFNβ following (A) RSV and (B) IAV infection, both at MOI = 0.1 and MOI = 1. Open circles mark data points with cytokine concentration below detection range. Error bars are s.d. of technical replicates.

(**C**, **D**) Activation of STAT1 and STAT2 and interferon-stimulated genes (RIG-I, PKR, OAS1) upon (C) RSV and (D) IAV infection, both at MOI 0.1, at 24 and 48 hours post-infection (h p.i.), in A549 WT, IFNAR1 KO, IFNLR1 KO, and IFNAR1–IFNLR1 dKO cells. Non-treated cells are labeled n.t. Activated STATs, p-STAT1 and p-STAT2, are STAT1 and STAT2 phosphorylated at Tyr 701 and at Tyr 690, respectively. RSV F, RSV fusion glycoprotein; IAV NP, IAV nucleoprotein.

Figure 2. Prestimulation with IFNβ or IFNλ1 inhibits propagation of RSV and IAV in A549 WT cells.

(**A**, **B**) Effect of prestimulation with IFNβ (1000 U/ml) or 50 ng/ml IFNλ1 24 h prior to infection on the relative abundance of viral RNA 24 hours post (A) RSV and (B) IAV infection, both at MOI = 0.1 and MOI = 1. Each filled circle corresponds to a single experimental replicate. Error bars show the smallest and the largest ratio of technical replicates for the naive and IFN-prestimulated cells.

(**C**, **D**) Effect of prestimulation with IFNβ (1000 U/ml) 24 h prior to infection on the abundance and activation of STAT1 and STAT2 and abundance of interferon-stimulated genes (RIG-I, PKR, OAS1) and (C) RSV fusion glycoprotein (RSV F) and (D) IAV nucleoprotein (IAV NP), 24 and 48 hours post-infection (h p.i.), both at MOI = 0.1 and MOI = 1. Non-treated cells are labeled n.t. Activated STATs, p-STAT1 and p-STAT2, are STAT1 and STAT2 phosphorylated at Tyr 701 and at Tyr 690, respectively.

Figure 3. Impact of RSV or IAV infection on STAT1 activation.

(**A**, **B**) A549 WT cells 24 hours post-infection with (A) RSV at MOI = 0.1 or (B) IAV at MOI = 0.1. Channels with IRF3 (red), p-STAT1 (yellow), RSV (fusion glycoprotein in magenta, only in panel B) or IAV (nucleoprotein in cyan, only in panel C) are shown as incremental overlays. White dotted lines are nuclear outlines determined based on DAPI counterstaining (channel not shown). Scale bar, 50 μm.

(**C**, **D**) Histograms showing the empirical probability density of p-STAT1 24 hours post-infection with (C) RSV at MOI = 0.1 or (D) IAV at MOI = 0.1 in cells classified as either expressing (+) or not expressing (−) the virus. In each pair of overlaid histograms, a Kolmogorov–Smirnov statistic is reported (with its credible interval, Crl) and a corresponding maximally discriminating threshold is marked with a vertical dash-dotted line. In each subpanel of (C) and (D), three experimental replicates are shown separately and the numbers of cells that were classified as either virus-positive or virus-negative are given on the left- and the right-hand side of the plus sign, respectively.

Figure 4. Effect of prestimulation with IFNβ or IFNλ1 on IAV spread in A549 WT cells.
Naïve A549 WT cells or A549 WT cells prestimulated with either IFNβ (100 U/ml) or IFNλ1
(50 ng/ml) for 24 hours were infected with IAV at MOI = 1 and fixed 24 hours post-infection
(i.e., 48 hours since the beginning of prestimulation). Interferon-containing cell culture media
were not displaced upon infection.

(**A**) Representative fields of view showing naïve cells and cells prestimulated with either
IFNβ or IFNλ1 after infection with IAV (cyan, IAV nucleoprotein). White dotted lines are
nuclear outlines determined based on DAPI counterstaining (channel not shown). For each
pretreatment condition, sample images from two experimental replicates are shown. Scale
bar, 50 μm.

(**B**) Proportions of productively IAV-infected cells among naïve cells and cells prestimulated
with either IFNβ or IFNλ1, based on nucleoprotein expression. Data from two independent
experiments are shown separately; each circle corresponds to a single field of view. The
fields of view do not overlap. Black diamonds and featuring error bars denote mean
proportions ± s.e.m. The number of cells analyzed in all fields of view from each experiment
is given above respective data series.

Figure 5. RSV preinfection inhibits IAV infection in an interferon-dependent manner in A549 cells. Analyzed are infections in four A549 cell lines: WT, IFNAR1 KO, IFNLR1 KO, and IFNAR1–IFNLR1 double KO (dKO). Naïve cells and cells preinfected with RSV at MOI = 0.1 for 30 hours were infected with IAV at MOI = 1 and cultured subsequently for 24 hours. Cell culture media was not displaced upon IAV infection.

(**A**) Representative fields of view showing WT and IFNAR1–IFNLR1 dKO cells preinfected with RSV (magenta, RSV fusion glycoprotein) after infection with IAV (cyan, IAV nucleoprotein). White dotted lines are nuclear outlines determined based on DAPI counterstaining (channel not shown). Scale bar, 50 µm.

(**B**) Proportions of IAV-infected cells in naïve cells and in cells preinfected with RSV. Data from two independent experiments are shown separately; each circle corresponds to a single FOV. Black diamonds and featuring error bars denote mean proportions ± s.e.m. Two encircled gray diamonds above the data points for WT cells are mean proportions of the IAV-infected WT cells that were not preinfected with RSV (naïve WT cells in Fig 4B), shown here for comparison.

(**C**) Impact of preinfection with RSV on the number of IAV RNA molecules per cell in all four considered cell lines. Cells were infected with RSV and, after 30 hours, with IAV. dPCR has been performed for cells collected 24 hours post IAV (i.e., 54 hours post RSV) infection. Error bars are s.d. of technical replicates.

Figure 6. IAV infects preferentially A549 WT cells preinfected by and expressing RSV, but this effect is not observed in A549 IFNAR1–IFNLR1 dKO cells. A549 WT and A549 IFNAR1–IFNLR1 dKO cells were preinfected with RSV (MOI = 0.1) 30 hours before infection with IAV (MOI = 1) and fixed 24 hours after infection with IAV.

(**A**) Representative fields of view showing WT cells and IFNAR1–IFNLR1 dKO cells preinfected with RSV (magenta, RSV fusion glycoprotein), after subsequent IAV (cyan, IAV nucleoprotein) infection (overlays). White dotted lines are nuclear outlines determined based on DAPI counterstaining (channel not shown). Scale bar, 50 μm.

(**B**) Histograms of RSV intensity in cells classified as either infected (cyan) or non-infected (sepia) with IAV in A549 WT cells and A549 IFNAR1–IFNLR1 dKO cells. Histograms show aggregated data from two experimental replicates. The numbers of cells that were classified as either virus-positive or virus-negative are given on the left- and the right-hand side of the plus sign, respectively.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6