1

Hydrophobicity in clearwing Lepidoptera: impact of scale micro and nanostructure, and trade-off with optical transparency

Doris Gomez[1*], Jonathan Pairraire[2], Charline Pinna[3], Monica Arias[1], Céline Houssin[3], Jérôme Barbut[3],

Serge Berthier[2], Christine Andraud[4], Thierry Ondarçuhu[5], Marianne Elias[3]

[1] CEFE, CNRS, Univ Montpellier, EPHE, IRD, Montpellier, France, 34090

[2] INSP, Sorbonne University, CNRS, Paris, France, 75005

[3] Institut de Systématique, Evolution, Biodiversité (ISYEB), CNRS, Muséum national d'Histoire

naturelle, Sorbonne Université, EPHE, Université des Antilles, France, 75005

[4] CRC, MNHN, Paris, France, 75005

[5] IMFT, Univ. Toulouse, CNRS, Toulouse, France, 31400

Corresponding author: Doris Gomez

Email: doris.gomez@cefe.cnrs.fr

Abstract

While scales ensure hydrophobicity in opaque Lepidoptera, their reduction in clearwing species question hydrophobicity and its dependency on wing microstructures (scale presence, morphology, insertion, coloration) and nanostructures (scale ridge features). By measuring static contact angle (CA) of water droplets at different stages of evaporation in opaque and transparent patches of 23 clearwing species with varied microstructure, we establish for the first time that transparency is costly for hydrophobicity, a cost depending on microstructure: transparent patches lose more hydrophobicity with water evaporation than opaque patches, a loss attenuated for higher scale densities, coloured scales (for erect scales), and when combining two types of scales (piliform and lamellar). Nude membranes show lowest hydrophobicity. Models best predicting CA include microstructures for all scales, and nanostructures only for flat scales. We reveal for the first time that wing hydrophobicity negatively relates to optical transparency. Finally, tropical species have more hydrophobic transparent patches but similarly hydrophobic opaque patches compared to temperate species. All findings are consistent with the physics of hydrophobicity, especially on multiscale roughness, and underline that wings are evolutionary compromises.

Keywords: Lepidoptera; transparency; trade-off; multiscale roughness; hydrophobicity

INTRODUCTION

Hydrophobicity is essential for terrestrial organisms. As predicted by physics (1) and illustrated in plants (2), a key parameter for hydrophobicity is surface texture or roughness. A water droplet sitting on a textured hydrophobic surface can exhibit two different wetting states. First, in the Cassie-Baxter state (Figure 1, series a), the water droplet sits on top of the texture, with trapped air underneath and cavities filled with air (composite state: under the drop, water can be in contact with solid or air), and hydrophobicity is at a maximum. If this state is thermodynamically unstable, the water droplet may undergo the so-called Cassie-Baxter to Wenzel transition, in which water penetrates the air-filled cavities by capillarity. In the Wenzel wetting state (Figure 1 series b), the water droplet fully fills all the cavities of the textured surface and adheres to the surface (non-composite state: under the drop, water is in contact with solid only and no longer with air), decreasing the surface energy; hydrophobicity is then lost (3). Compared to the Wenzel state, the Cassie-Baxter state is of high biological interest as it offers an incomplete water-surface contact and a weak water adhesion. Maintaining a stable Cassie-Baxter state is crucial to maintain high hydrophobicity under harsh environmental conditions, rainfall for instance. Roughness at nanoscale – the parameter most studied to date in animals and plants - increases hydrophobicity, as shown in cicadids and dragonflies (4–6). Yet, multiscale roughness –at nano and micro scale – is even more efficient: it increases hydrophobicity and its thermodynamic stability (in modelling studies 7, 8, illustrated in the Lotus (so-called 'Lotus effect') and other plants in 9) and reduces water adhesion (7). Increasing thermodynamic stability allows maintaining hydrophobicity with water droplets of various sizes (dew, fog, rain) and increases anti-fogging properties, i. e. the resistance to tiny water droplets condensing on the surface. While the role of nanostructures in hydrophobicity has been extensively documented (e.g. 10, 11), the role of microstructures shape in determining hydrophobic properties has been limited to simple geometries (cones in 12, cylinders in 13, 14) and remains poorly investigated from an empirical perspective. The only existing empirical studies with such an approach either focus on one type of micro-architecture (6, 15) thereby excluding microstructural influence, or they describe variation in hydrophobicity between various micro-architectures but without invoking explanations (16).

Lepidoptera (from the ancient Greek λεπίς: scale and πτερόν: wing) – butterflies and moths – offer an outstanding group to investigate this question. They are characterized by wings entirely covered with flat and coloured lamellar scales (17). Through their pigmentation and structure, scales are involved in multiple functions such as antipredator defences (e. g. camouflage, deflection, mimicry in 18, 19), communication (20), thermoregulation (17, 21–23), or flight enhancement (24, 25). They also confer superhydrophobic properties to the wing, resulting in water repellency and self-cleaning (26, 27). Superhydrophobicity *sensu lato* is defined by water droplets making high contact angles (>150°) with a surface. Self-cleaning – superhydrophobicity *stricto sensu* (definition not taken here) –

adds to this condition a weak water adhesion, estimated by a minimal tilt from the horizontal plane

needed for water droplets to roll-off (roll-off angle of a few degrees) or a minimal hysteresis (difference

between advancing and receding contact angles). Superhydrophobicity is thus a *sine qua non* condition

for water repellency and self-cleaning. Opaque butterflies and moths typically have self-cleaning

wings, as attested by small roll-off angles (15, 28). Scarce relevant studies suggest that wing

hydrophobicity may depend on wing microstructure (presence and type of scale in 26, scale type and

insertion angle in 29), and on wing macrostructure: species with longer wings (4), or larger ratio of

wing area to body mass (26) show higher hydrophobicity and wing shape was invoked to explain

natural variations in hydrophobicity (4).

While the vast majority of Lepidoptera species has opaque wings, some species from various

lineages show transparent or translucent wings (30), which reduces their detectability from visually-

hunting predators (31–33). Transparency shows a broad microstructural diversity (i. e. scale diversity,

see examples in Figure 2), the membrane being nude or covered with scales varying in type (piliform,

i. e., hair-like, and/or lamellar), insertion on the membrane (flat or erect), and colouration (coloured

or transparent) (30). All combinations of scale type, insertion, and colouration (i. e., structural

strategies 30) can be found in nature (Figure 2), and they differ in their efficiency at transmitting light

: the nude membrane are most efficient while flat coloured scales (lamellar alone or in combination

with piliform) are least efficient (30). Microstructures are complemented by nanostructures:

longitudinal ridges on scales, and nanostructures on the wing membrane, of various shape and density.

Membrane nanostructures reduce reflection levels and increase light transmission (34–37).

Because transparency often entails profound modifications of scale dimensions and density

(30), we can suppose that transparency may be potentially costly for hydrophobicity, for both water

repellency and self-cleaning. These functions are vital for butterflies and moths: water repellency is

crucial for flight and for preventing wings from sticking together, especially in tropical rainforest

species with daily rain and high humidity. Likewise, self-cleaning helps removing dust contamination

that impairs flight (26). Among the lepidopteran species investigated so far for hydrophobicity (15, 26,

27, 29, 38), only three clearwing butterfly species have been included: *Parantica sita* (with lamellar

titled scales) and *Parnassius glacialis* (with flat lamellar scales), with high or moderate hydrophobicity

respectively (15, 29), and *Greta oto* (with piliform scales) with one of the lowest hydrophobicity values

found in butterflies (27). Scarce data suggest that a greater reduction in scale dimensions or coverage

on the wing membrane may entail higher costs in terms of hydrophobicity. However, large-scale

comparative studies are currently lacking.

To fill that gap in our knowledge, here we explore to what extent anti-wetting ability is influenced by

micro and nanostructure in species that largely differ in their wing microstructure, by selecting a subset

of 23 species (Figure 3) from a broad study of 123 clearwing Lepidoptera species (30). In these species,

we explored the links between structure, hydrophobicity and optics while controlling for phylogenetic relatedness between species. First, we explored the relationships between hydrophobicity and wing structure, at macro-, micro- and nano-structural level: (i) we measured the contact angle (CA) made by water droplets of various sizes on the wing (ii) We then quantified wing macro-, micro-, and nano-structures, and related them to hydrophobicity. (iii) Using a modelling perspective, we ran various Cassie-Baxter and Wenzel models differing by their assumptions and compared the predicted to the observed CA values to assess the relative importance of microstructure and nanostructures in determining the observed hydrophobicity. Second, to identify the selective pressures acting on hydrophobicity, we tested whether hydrophobicity and light transmission showed potential trade-off or synergy. If microstructures play a dominant role in conferring hydrophobicity, species most efficient at transmitting light – which lack scales or have highly modified scales in low coverage on the wing surface – are expected be less efficient at repelling water. Third, to identify whether hydrophobicity is influenced by environmental conditions, we tested the links between habitat latitude and hydrophobicity: if repelling water is more important in the tropics where rain and humidity are inescapable, tropical species are expected to show higher hydrophobicity than temperate species.

RESULTS

Variation in hydrophobicity and relation to wing macrostructure

We measured the contact angle of water droplets and wing surface in the transparent and opaque zones of the forewing of three museum specimens per species, and we monitored contact angle at three times, as water evaporated and droplet size decreased (Figures 2 and S1). We performed classic mixed models and Bayesian models, the latter controlling for species phylogenetic relatedness, to explore to what extent hydrophobicity depends on general wing macrostructure (opaque and transparent zone, wing area, shape). Comparing models controlling or not for phylogeny helped us purge our results from spurious correlations arising from phylogenetic ancestry.

We observed a general decrease in hydrophobicity with water evaporation in the opaque zone and in the transparent zone (Figure 4A). Transparency appeared costly: (i) the transparent zone showed lower hydrophobicity than the opaque zone of the same wing, whatever the size of the water droplet considered (Figure 4A, zone effect in Table S1, see Figure S2 for distribution of hydrophobicity levels), (ii) and we observed a stronger decrease in hydrophobicity with water evaporation in the transparent than in the opaque zone of the same wing (time x zone effect in mixed models in Table S1, Figure 4A). CA did not correlate to wing length but negatively correlated to the Wing Area / Body Volume ratio (Table S1, effect only important when correcting for phylogeny, Figure 4C). Contrary to our expectation, species with more elongated wings (higher FW ratio) had a slightly lower hydrophobicity (Figure 4C, effect marginally important when corrected for phylogeny in Table S1).

Finally, species with more elongated wings or with shorter wings exhibited a greater loss of

hydrophobicity with evaporation (LWratio x time effect in Table S1).

Variation in hydrophobicity and relation to wing microstructure

We used the same approach (classic mixed and Bayesian models) to explore to what extent

hydrophobicity depends on wing microstructure – namely scale presence, type (piliform and/or

lamellar), insertion (erect or flat on the wing membrane), coloration (transparent or coloured), and

density in both the transparent and the opaque zones – and wing scale nanostructures – namely the

width and spacing of longitudinal ridges that were present on all scales.

The influence of wing microstructure on hydrophobicity was pervasive in our results (Figure

5): (i) we found a higher interspecific variance in contact angle values in the transparent than in the

opaque zone (Figure 4A, Fligner-Killeen tests with all times together χ^2=79.48, p<0.001 or separated

at T1: χ^2=49.57, p<0.001; T2: χ^2=29.24, p<0.001; T3 χ^2=26.47, p<0.001), maybe in relation to the higher

interspecific microstructural diversity of the transparent zone. (ii) The nude membrane (N) yielded a

lower hydrophobicity (Scale presence effect shown in classic models in Table S3a, Scale Nb (0>1) effect

in classic and Bayesian models in S3b) but a similar decrease in hydrophobicity compared to the

structural strategies that involved scales (Table S2, scale presence effect in Table S3a, Figure 5). (iii)

Combining two types of scales (piliform and lamellar) yielded comparable levels of hydrophobicity, but

a lower decrease in hydrophobicity with evaporation than having only one type of scales only (Table

S2, Scale Nb (2>1) x Time interaction in Table S3b, Figure 5). (iv) The decrease in hydrophobicity with

evaporation was lower for erect coloured scales than for erect transparent scales (Table S2, Colour

EC>ET x Time interaction in Table S3c, Figure 5). (v) The decrease in hydrophobicity with water

evaporation was attenuated for denser scales (Table S2, time x density interaction in Table S3c, Figure

4B). This attenuation by density was stronger in the transparent zone than in the opaque zone (Table

S2, time x density x zone interaction in Table S3a, S3b, S3d, Figure 4B), and for erect scales (Table S2,

time x density interaction in Table S3c). (vi) For flat lamellar scales, the decrease in hydrophobicity with

evaporation was attenuated when scales were arranged in a higher number of layers, be they in the

transparent or in the opaque zone (Table S2, time x nb layers interaction in Table S3d). (vii) Finally,

scale nanostructures did not explain an important part of the variation in hydrophobicity (scale ridge

ratio never retained in the best models).

Because erect geometries (involving piliform and/or lamellar scales: PLE, PE, LE) appear to

behave differently in reaction to water than flat geometries, we analysed scale dimensions and spacing

by performing classic mixed models and Bayesian analyses on the broad dataset of 123 species used

by Gomez et al. (30) to get more representative trends. Compared to flat scales, erect scales were

shortened when one type of scale was involved, especially in lamellar scales (Figure S3, Table S4).

When both present, piliform and lamellar scales were in similar densities, close in space, and piliform

scales were 2.6 times longer than lamellar scales, creating a multi-hierarchical roughness at
microscopic scales (see SI for supplementary results, Table S4, Figure S4).

For modelling, we first assigned water droplets at T1 to a Cassie-Baxter regime when droplets
showed a high contact angle after evaporation (CA≥120° at T3; results remained the same when
shifting the threshold to 110°); otherwise, water droplets at T1 were assigned to a Wenzel regime
(CA<120° at T3). We tested Cassie-Baxter models with microstructures alone (scale dimensions,
insertion, and density) or with scale microstructures and nanostructures (ridge-ratio) to assess their
relative importance in determining hydrophobicity observed at T1, selecting as best the model that
minimized the difference between predictions and observations, for each structural strategy
separately (SI, Figure S5 for model details, Figure S6 for best model selection).

The best model included only microstructures for all erect strategies (with piliform and/or
lamellar scales PE, LE, PLE, with a mean difference between observations and predictions of 7°, 14°,
and 7° respectively) and for flat lamellar scales in low densities (not fully covering the wing membrane,
with a mean difference between observations and predictions of 22° and 25° for the transparent and
the opaque zone respectively). This was consistent with the fact that in these cases water contacted
scale contour/edge, which did not exhibit any nanostructures. For strategies involving erect piliform
scales (PE, PLE), the model with bending piliform scales outperformed the model with fully erect
piliform scales. For flat piliform scales (PF) and for flat lamellar scales (LF) in high densities (fully
covering the membrane), the best model included microstructures and scale nanostructures (with a
mean difference between observations and predictions of 9° for PF, 22° and 25° for LF for the
transparent and the opaque zone respectively). In this case, the scale upper side is in contact with
water and both types of structures come into play. In general, predictions were rather close to
observations with no systematic bias towards under or over-estimation of Φ_S, the fraction of the
droplet contact area where water is in contact with the solid (Figure 6).

For the water droplets not categorized as in a Cassie-Baxter state, we tested whether a Wenzel
model with microstructures only could predict the observed values at T1 (Figure S7). We could not test
a Wenzel model with micro and nanostructures, as it required to precisely characterize scale
nanogroove height and membrane nanostructures, which was not allowed on museum specimens.
Predictions of Wenzel models were far below observations (Figure S8).

DISCUSSION
We here considered hydrophobicity as a proxy for self-cleaning ability. Indeed, although we could not
quantify water droplet roll-off angles precisely with our set-up, we observed that water droplets rolled

off extremely fast from the wings when not perfectly horizontal (D Gomez & M Elias personal observation July 2016; J Pairraire personal observation May 2018; C Houssin personal observation April 2021), which made our measurements particularly time-consuming. This suggested small roll-off angles and weak water adhesion. Small roll-off angles are commonly found in opaque butterflies (15, 28) and in *Parantica sita* and *Parnassius glacialis*, the two clearwing butterfly species studied to date (29).

The observed decrease in hydrophobicity with water evaporation is commonly observed in hydrophobic human-made surfaces (14, 39, 40) and in natural surfaces, as in the transparent-winged damselfly *Ischnura heterosticta* (3). It is interpreted as a loss of self-cleaning ability when contact angle values get below the hydrophilicity threshold (3). Beyond this general trend, the lower hydrophobicity and the stronger loss in hydrophobicity with water evaporation in the transparent than in the opaque zone undoubtedly show that transparency entails important costs in terms of hydrophobicity.

Variation in hydrophobicity and relation to wing macrostructure

Relationships between hydrophobicity and wing macrostructure were surprising, probably due to the taxonomic level investigated. Our absence of relationship between CA and wing length contrasted with Byun et al.'s (4) marginal positive correlation between these variables (study on 24 species, 10 insect orders, wing length effect on CA, estimate=10.58±5.33, t=1.99, p=0.067), Wing shape was much more diverse in Byun et al.'s dataset (LWratio: min=1.2, mean=3.6, max=9.3) than in ours (23 species, 1 insect order), LWratio: min=1.8, mean=2.2, max=3.2) When restricting Byun's dataset to our maximal value for LWratio, the effect disappeared (n=13 species, wing length estimate=2.42 ± 5.09, t=-0.47, p=0.652). Our finding a negative correlation between CA and the ratio of wing area to body volume contrasted with Wagner et al.'s (26) positive correlation between CA and the ratio of wing area to body mass (study on 38 species from 14 insect orders, wing area/body mass ratio effect on CA: estimate=11.09 ± 3.75, t=2.96, p=0.007), When restricting Wagner et al's dataset to insects without elytra but with a microstructured membrane (Odonata, Ephemeroptera, Lepidoptera, some Planipennia), the effect disappeared (n=21 species (4 insect orders), estimate=0.15 ± 1.12, t=-0.14, p=0.895). Finally, the negative correlation between forewing elongation and CA (Figure 4C) is surprising. Water droplets exert higher moment force when further away from the butterfly body and should select for higher hydrophobicity; yet, more elongated wings may already show higher elasticity that may already ensure rapid droplet roll-off through movement, and offset the need for a higher hydrophobicity.

Variation in hydrophobicity and relation to wing microstructure

Going back to the physical theory behind hydrophobicity, several studies have shown that a
single-level structure does not necessarily guarantee a low water adhesion, even in the Cassie-Baxter
state (see references in 7). Introducing higher levels of hierarchy increases the robustness of a surface
hydrophobicity (8): it stabilizes the Cassie-Baxter state by dramatically decreasing the contact area
fraction (ratio of contact area to the total surface area of the structure) and thus the adhesion force of
water droplets, and by enlarging the energy difference between the Cassie-Baxter and the Wenzel
states. Hierarchical structures can be frequently found in plants and in animals. For instance, in the
water strider *Gerris remigis*, leg water resistance is due to the hierarchical structures of nano-grooved
microsetae, which prevents striders from being drowned under heavy rainfall (42). This likely explains
why, in our study, the combination of erect piliform scales and lamellar scales yields a lower loss of
hydrophobicity with evaporation than piliform or lamellar scales alone. Such geometries have a 3-level
roughness: (1) erect piliform scales bending over lamellar scales (piliform scales are 2.6 times longer
than lamellar scales and first in contact with water), (2) erect lamellar scales tightly associated in space
to piliform scales (similar density and close spacing), and (3) nanostructures on scales and on the wing
membrane. Hydrophobicity likely results from the combination of the complex geometry of erect
microstructures (which considerably reduces the proportion of the total surface in contact with water),
and the gain in mechanical resistance (gain in elasticity and resistance against breakage) of piliform
scales when bending against lamellar scales. The importance of elasticity of bending hair-like
microstructures has been found in several studies. In the Lady's mantle plant (*Alchemilla vulgaris*),
hairs are hydrophilic when measured individually, but they bend and coalesce into bundles when in
contact with water droplets; their elasticity results in a repulsive interaction between the droplet and
the plant surface, which maintains hydrophobicity (CA above 90°) (43). Likewise, in *Nasutitermes*
termits, large bending hairs and small micrasters (micraster wavelength was around 11,7 µm according
to our measurements taken on Figure 3C from 41) enable hydrophobicity (CA above 90°) in both rain
and mist conditions (44).

Increasing the density of microstructures does not significantly change hydrophobicity, but
leads to a lower loss of hydrophobicity with water evaporation, i. e., to a higher resistance to tiny water
droplets. This can be seen for all structural strategies (effect time x density) and for flat scales organized
in layers (effect time x number of layers). This is consistent with the fact that, during droplet
evaporation, the Cassie-Baxter regime is more robust for large microstructure density. Not only scale
architecture but also coloration can contribute to hydrophobicity. Erect scales show a lower loss of
hydrophobicity when pigmented than when transparent. In the transparent zone, coloured scales
exhibit colours ranging from pale yellow to brown and black. They are likely impregnated by melanin
pigments, which are known to be involved – for some biochemical forms – in cuticle sclerotization
(hardening) (45). Hence, the additional hardening conferred by pigments may increase their

mechanical resistance to deformation and may contribute to maintaining hydrophobicity, even when

evaporation occurs.

We found that scale nanostructures did not contribute significantly to wing hydrophobicity for

most structural strategies, except for flat lamellar scales fully covering the wing membrane and

organized in layers, in the transparent or in the opaque zone, and for flat piliform scales in the

transparent zone (Figure S6). Our results bring novel evidence for a major role of microstructures in

explaining large variations in hydrophobicity when diverse microstructures are considered. The rare

existing studies on the subject suggest a synergetic effect of scale nanostructures and microstructures

on enhancing surface hydrophobicity (experiments on one type of microstructure, namely flat lamellar

scales in opaque butterflies, in   (15, 46) or hairs in the wing of the housefly *Musca domestica* (47);

theoretical modelling on one type of microstructure in (48)), or even a major role of nanostructures in

the overall variation (15, 47). Yet, these two analyses only examine one type of microstructure, thereby

potentially underestimating the importance of microstructures when more types of microstructures

are considered. Overlap in scales is assumed to help anisotropy in hydrophobicity (49). Here, we find

that it attenuates the loss of hydrophobicity with water evaporation, thereby maintaining self-cleaning

ability more efficiently.

Wing mechanical resistance is crucial for flight and geometries that limit protrusion height are

more resistant to breakage and less hydrophobic (50). Several of our results suggest scale height may

be limited: (i) erect piliform scales are likely bending over the membrane as shown by modelling. (ii)

When piliform scales are alone, they have similar height, be they flat or erect, maybe because they

bend easily, which may limit their sensitivity to breakage. (iii) Erect lamellar scales are shortened

compared to flat lamellar scales, which likely increases their resistance to breakage. (iv) Erect

transparent lamellar scales are densely packed, as shown in Gomez et al. (30), which can also increase

their mechanical resistance. Further experiments are needed to elucidate these aspects, and clarify

the role of nanostructures, as not only their presence, but their topography and its randomness have

been recently suggested to play a role in determining antiwetting properties (51).

Finally, the poor predictability of Wenzel model is not surprising since the validity of Wenzel

model – as currently formulated – is widely questioned in the literature (41 and references therein).

This is especially true for our dataset given that most microstructures only weakly increase the

roughness parameter r (see materials and methods) while providing strong pinning sites for the contact

line and thus significantly increasing the contact angle measured at T1.

Trade-off between hydrophobicity and optical transparency

Using spectrometric measurements of wing direct transmittance, we found a negative relationship

between contact angle and mean transmittance over 300-700 nm (Table S5, Figure 7). A 10% increase in transmittance resulted in a 4° loss in CA. While this relationship was marginally significant without controlling for phylogenetic relatedness, whatever the level of analysis (all CA values, mean CA values per individual, mean CA values per species), it was statistically important (i. e., the 95% credibility interval did not contain 0) in phylogeny-controlled analyses when considering all measurements or mean individual values, but less important at species level (90% credibility interval), likely because of weaker statistical power. In agreement with our prediction that microstructures play a major role in hydrophobicity, we find a negative relationship between hydrophobicity and transparency, a condition associated with major modifications in scale shape and density. This trade-off can be seen from the literature: the nymphalid butterfly *Greta oto* has been shown to exhibit a high transparency resulting from poorly dense erect piliform scales and efficient antireflective nanostructures (35, 37) but a weak hydrophobicity (27). Likewise, the trade-off can be seen in the dragonfly *Gynacantha dravida* (which has micro and nanospikes), in which distal wing parts show higher hydrophobicity but lower transmittance compared to proximal wing parts (52). Finding this trade-off is fully compatible with the fact that some species with nude membrane show high hydrophobicity: membrane nanostructures are at full play in species with nude membrane and can potentially efficiently reduce both reflection and water adhesion, like in the cicada *Aleeta curvicosta* (53). In species with erect scales, our models show a major role of microstructures and a negligible role of nanostructures in hydrophobicity. If selection for hydrophobicity is relaxed on membrane nanostructures in species with erect scales, this may explain why nanostructures are so diverse in their architecture (type, density), as recently shown in clearwing Lepidoptera (36).

Hydrophobicity and latitude

Compared to their temperate counterparts, species living in the tropics had a higher hydrophobicity in their transparent zone – loss of 10° CA for 10° increase in latitude – but a similar hydrophobicity in their opaque zone (Table S6, Figure 8) All species showed superhydrophobic opaque patches (intercept above 150° in Table S6, Figure 8B) and there was no relationship between the proportion of wing area occupied by transparency and latitude that could have explained the observed variations in CA (model with proportion of transparency as dependent variable, latitude effect = 0.08 ± 0.22, t=-0.31, p=0.76, the best model was the null model). This result is consistent with the prediction that in tropical climates where species face more humid conditions, and where rainfall can happen daily, there is a stronger selective pressure for increased hydrophobicity. While the opaque zone allows maximizing hydrophobicity in all environmental conditions, the differential in environmental conditions reveals the costs of transparency. To our knowledge, this is the first evidence for a higher hydrophobicity in more humid conditions. Scarce relevant studies have explored the link between habitat humidity and

species hydrophobicity: at local geographical scale, all four cicada species studied by (6) show superhydrophobicity regardless of whether they live in dry or more humid habitats, but annual species are more hydrophobic than the species that emerges in large swarms every 17 years. Likewise, Goodwyn et al. (29) suggest that in transparent butterflies hydrophobicity may depend on lifespan and migration ability. Further studies are needed to elucidate the links between hydrophobicity and species ecology.

This study is the first to mix many architectures at microscale and give access to the relative role of micro and nanostructures at transmitting light and repelling water. It shows that selection likely acts on multiple features of scales (shape, orientation, coloration, density) in relation to climatic conditions. Considering more natural geometric complexity in experimental and theoretical studies on hydrophobicity should open new venues for applied physics and answer open questions like the role of randomness in nanostructures, shown to improve optical transparency (35) but suggested to impair hydrophobicity (54, but see 51).

MATERIALS AND METHODS

Species selection

Scale type and scale insertion have been suggested to influence hydrophobicity(29); scale coloration, often involving melanin deposition which increases cuticle hardening in insects (45), could increase scale stiffness and ability to repel water droplets. Hence, we selected a set of species varying in structural strategies – scale type (N=nude membrane, P=piliform bifid or monofid scales, L=shape different than piliform, hereafter called lamellar, or PL=association of piliform and lamellar scales), insertion (E=erect or F=flat), and colouration (C=coloured or T=transparent) – from the study of 123 species of clearwing Lepidoptera (30). We minimized the phylogenetic relatedness between species harbouring the same type of structural strategies to increase the power of comparative analyses. We selected a total of 23 species from 10 families (Figure 1 & 2, list in Table S7), comprising 3 species for the structural strategies (N, PFC, PEC, LFC, LFT, LEC, LET), 2 species for PLEC and LEC, and 1 species for PLET, as for some species only a limited number of specimens were present in the collections. For each species, we selected three specimens in good condition either from Paris MNHN collections or from our own private collections. 54/69 specimens (all species but *Eutresis hypereia*) had labels with exact collect location that could be tracked down to GPS coordinates.

Hydrophobicity measurements

For each specimen, we used a purpose-built water-droplet dispenser (a graduated pipette on a holder)

and a Keyence VHX-5000 microscope (equipped with Z20 zoom) to image water droplets on butterfly

wings. As a general procedure, we dropped a series of three 1µl water droplets (volume usually taken

to assess hydrophobicity (3, 29)) at three locations of the transparent and opaque zones of the dorsal

side of a wing. After the water droplet was dropped (time T1), we allowed its volume to be

approximately divided by two (time T2) and by four (time T3) compared to its original volume. Since

evaporation kinetics depended on droplet shape, time intervals elapsed between consecutive photos

were not identical from one species to another. At each time, we took a photo (Figure 1) in which we

measured the static contact angle (Figure S1). Contact angle measurements were first measured on

both wings and found highly repeatable (see detailed methods and results in SI, Table S8). We thus

kept the same protocol, but we measured only the forewing.

Measurements of wing macro, micro and nanostructure

To characterize wing macrostructure, we took photos of the three specimens of each species using a

camera (D800E Nikon, 60mm lens, annular light). We analysed photos using ImageJ (55). Given the role

of wing length (4), maybe wing shape (4), and ratio of total wing area to body mass (53) on

hydrophobicity and self-cleaning ability, we computed wing length, length-to-width LW ratio and the

ratio of total wing area to body volume, taking the volume as a proxy for mass for dry specimens, and

assuming the body to be a cylinder, for which we measured length (thorax+abdomen) and width. Using

the 'rptR' package (56), we found that all wing macrostructural measurements were repeatable, i.e.

that a specimen was representative of its species for all wing macrostructural variables (Table S8).

To characterize wing microstructure (i.e. scale characteristics, presence, type, insertion,

coloration, density), we imaged the dorsal side of forewing transparent and opaque zones using

microscopes (Zeiss Stereo Discovery V20 and Keyence VHX-5000). We did that in one specimen per

species because scale dimensions and density had already been found repeatable at zone by species

level in Gomez et al. (30). Using ImageJ or the Keyence built-in tool, we measured scale density (per

mm²), length and width (µm), scale surface (in µm²) as the product of length by width, and scale

coverage as the product of scale surface (expressed in mm²) by scale density. We counted the number

of different scale types: 0=nude membrane, 1= lamellar scales or piliform scales, 2= combination of

piliform scales and lamellar scales. For flat lamellar scales, we also computed the density of scale top

layer and computed the number of layers as the ratio between density and top layer density.

We quantified scale nanostructures – width and spacing (in µm) of longitudinal ridges

present on all scales – on the detailed top-view microscopical images. We computed the ridge-ratio as

ridge width to spacing (SI, Fig S2). Since museum specimens are patrimonial material, we could not

damage or destroy specimens to image membrane nanostructures or scale ridge height.

Optical measurements

For one specimen per species, we measured specular transmittance from 300 to 700 nm as in Gomez et al. (30), using a deuterium-halogen lamp (Avalight DHS), direct optic fibres (FC-UV200-2-1.5 x 100) and a spectrometer (Avaspec-2048 L, Avantes). Wing samples were placed perpendicular at equal distance between fibres aligned 5 mm apart (1 mm diameter spot). We took five measurements of the forewing in various points of the transparent zone. Using Avicol v6 (57), we computed the mean transmittance over [300-700] nm, which described the level of optical transparency. Optical measurements had been found highly repeatable at species level in Gomez et al. (30).

Comparative analyses

All analyses were conducted using the R environment (58). We conducted mixed models not controlling for phylogeny using the 'nlme' R package (59). We selected the best mixed model using AICc minimization. We used the formulated model for Bayesian mixed models with MCMC analyses controlling for phylogeny, using the 'mulTree' R package (60). Bayesian analyses used the maximum clade credibility (MCC) phylogeny obtained in Gomez et al. (30) and pruned to targeted species. Comparing classic and Bayesian mixed models allowed us to assess the influence, if any, of phylogeny on the observed relationships. For Bayesian analyses, we used uninformative priors, an Inverse-Gamma distribution (shape = 0.001, scale = 0.001) for both random effect and residual variances (61), 2 chains of 500,000 iterations, burn-in of 10,000, and thinning interval of 300. Fixed effects were considered statistically important when associated with 95% credibility intervals excluding zero, and less important (marginally important) when associated with 90% credibility interval excluding zero.

Considering all contact angle measurements, we analysed the variation in contact angle with (i) wing macrostructure descriptors – time, zone, forewing size, surface, LW ratio, the ratio of total wing area divided by body volume, and relevant two and three-way interactions –; (ii) with wing microstructure descriptors – time, zone, wing length (to correct for variation in scale dimensions), scale length, width, density, scale type, number of different types, scale insertion, scale colouration, number of layers, ridge-ratio, and biologically relevant two and three-way interactions. (iii) To characterize spatial geometries, we analysed scale length against scale insertion for structural strategies with one scale type. For structural strategies based on both scale types (piliform and lamellar), we analyzed length ratio, density ratio and spatial association between the two scale types in relation to scale insertion (see SI for details). (iv) We analyzed the difference between contact angles predicted by various Cassie-Baxter and Wenzel models (see below) and measured contact angles at T1 to select the best models. (v) We tested for a potential trade-off between optical transparency and wing hydrophobicity, considering all measurements of contact angle, individual mean values, or species

mean values at T1. (vi) Finally, we tested whether tropical species were more hydrophobic than

temperate species. To do so, we related for each specimen its average CA value to its latitude to the

equator, the proportion of wing area occupied by transparency and wing length, while taking species

as random effect, for the opaque and transparent zone separately. We also tested whether variation

in the proportion of transparency could be explained by the latitude to the equator.

Hydrophobicity modelling

To assess to what extent scale micro- and nanostructure could explain the observed variation in

hydrophobicity, we elaborated Cassie-Baxter model (62), whereby water only wets the top of the

surface and the apparent contact angle observed at equilibrium is θ^*.

$$\cos\theta^* = \Phi_S \cos\theta_S + \Phi_A \cos\theta_A \ldots\ldots\ldots\ldots\ldots\ldots\ldots(\text{Eq 1})$$

where Φ_S is the fraction of the droplet contact area where water is in contact with the solid, and Φ_A

is the fraction of droplet contact area in contact with air, with $\Phi_S + \Phi_A = 1$. $\theta_A = 180°$ (contact angle

of water with air) and the angle θ_S (also called Young angle) is the contact angle of water on an ideal

smooth surface of the same material (wing membrane), set to 95°, as in Fang et al (15), a plausible

value given the presence of epicuticular wax on the wing membrane (63), even in clearwing

Lepidoptera (37). This equation is valid for water droplet size largely exceeding surface roughness

wavelength (64), which is the case for 1µl water droplets. Considering flat-topped geometries

simplifies Eq1 to:

$$\cos\theta^* = \Phi_S \cos\theta_S - (1 - \Phi_S)\ldots\ldots\ldots\ldots\ldots\ldots\ldots(\text{Eq 2})$$

While $\Phi_S = 1$ for a nude membrane, $\Phi_S < 1$ for all the other microstructures (Figure S5 for

calculations). We explored several hypotheses: piliform scales fully erect or bending over a fraction pp

= 3/5 of their length (the value 3/5 comes from P/L length ratio = 2.6 found for PLE strategies in Figure

S4), an organization of flat lamellar scales in one layer if in low density, or several layers (NL) when in

high density, models with microstructures only, or with micro and nanostructures. In the latter case,

we considered water droplets contacted scale nanostructures whether flat scales (PF or LF) covered

the wing membrane entirely or not, or only when scales were in high coverage, building a

homogeneous horizontal surface.

We also elaborated a Wenzel model (1), whereby water fills all the pores of the surface and

the wetting contact angle is primarily determined by surface roughness r. The apparent contact angle

θ^* is obtained by:

$$\cos\theta^* = r\cos\theta_S\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots(\text{Eq 3})$$

where r is the ratio between the total solid surface and the projected surface, and the angle θ_S is the

same as previously. We only included microstructures in the model since we had no measurement of

scale ridge height or membrane nanostructures. While r=1 for a nude membrane, r exceeds 1 for all

the other microstructures (see Figure S7 for calculations).

ACKNOWLEDGEMENTS

We warmly thank Jacques Pierre and Rodolphe Rougerie for help with species choice, identification,

and data on species ecology, Edgar Attivissimo for contributing to Keyence imaging, and Thibaud

Decaëns, Daniel Herbin, and Claude Tautel for species selection and identification.

COMPETING INTERESTS

This work was funded by Clearwing ANR project (ANR-16-CE02-0012), HFSP project on transparency

(RGP0014/2016) and a France-Berkeley fund grant (FBF #2015-58). No competing interest. The authors

declare no competing interests.

AUTHOR CONTRIBUTIONS

Doris Gomez study and protocol design, funding acquisition, specimen selection, macrostructure and

microstructure measurements, optical measurements, data curation and analysis, hydrophobicity

modelling, writing original draft. Jonathan Pairraire: protocol design, structure and CA measurements,

CA computation. Charline Pinna: protocol design. Monica Arias: hydrophobicity modelling. Celine

Houssin: protocol design, structure and CA measurements, CA computation. Jerome Barbut: species

selection. Serge Berthier: study design, funding acquisition, writing review and editing. Christine

Andraud: study and protocol design, writing, funding acquisition, writing review and editing. Thierry

Ondarçuhu: hydrophobicity modelling, writing review and editing, Conceptualization. Marianne Elias:

study and protocol design, funding acquisition, specimen selection, data curation and analysis, writing

review and editing.

DATA AND CODE AVAILABILITY

Data are available at: https://doi.org/10.17605/OSF.IO/GB49E

Code availability: upon manuscript acceptance in a journal

ORCIDS

Doris Gomez http://orcid.org/0000-0002-9144-3426

Charline Sophie Pinna http://orcid.org/0000-0002-4947-0893

Monica Arias: http://orcid.org/0000-0003-1331-2604

Christine Andraud http://orcid.org/0000-0002-3112-9363

Serge Berthier http://orcid.org/0000-0002-9255-059X

Thierry Ondarçuhu: http://orcid.org/0000-0001-7657-5271

Marianne Elias http://orcid.org/0000-0002-1250-2353

REFERENCES

1. R. N. Wenzel, Resistance of solid surfaces to wetting by water. *Ind. Eng. Chem.* **28**, 988–994 (1936).

2. W. Barthlott, C. Neinhuis, Purity of the sacred lotus, or escape from contamination in biological surfaces. *Planta* **202**, 1–8 (1997).

3. J. Hasan, *et al.*, Spatial Variations and Temporal Metastability of the Self-Cleaning and Superhydrophobic Properties of Damselfly Wings. *Langmuir* **28**, 17404–17409 (2012).

4. D. Byun, *et al.*, Wetting Characteristics of Insect Wing Surfaces. *J. Bionic Eng.* **6**, 63–70 (2009).

5. M. Sun, G. S. Watson, Y. Zheng, J. A. Watson, A. Liang, Wetting properties on nanostructured surfaces of cicada wings. *J Exp Biol* **212**, 3148–3155 (2009).

6. J. Oh, *et al.*, Exploring the Role of Habitat on the Wettability of Cicada Wings. *ACS Appl. Mater. Interfaces* **9**, 27173–27184 (2017).

7. Y. Su, *et al.*, Nano to Micro Structural Hierarchy Is Crucial for Stable Superhydrophobic and Water-Repellent Surfaces. *Langmuir* **26**, 4984–4989 (2010).

8. M. S. Bell, A. Shahraz, K. A. Fichthorn, A. Borhan, Effects of Hierarchical Surface Roughness on Droplet Contact Angle. *Langmuir* **31**, 6752–6762 (2015).

9. W. Barthlott, M. Mail, C. Neinhuis, Superhydrophobic hierarchically structured surfaces in biology: evolution, structural principles and biomimetic applications. *Philos. Trans. R. Soc. A-Math. Phys. Eng. Sci.* **374**, 20160191 (2016).

10. N. A. Patankar, Mimicking the lotus effect: Influence of double roughness structures and slender pillars. *Langmuir* **20**, 8209–8213 (2004).

11. F. Porcheron, P. A. Monson, Mean-Field Theory of Liquid Droplets on Roughened Solid Surfaces: Application to Superhydrophobicity. *Langmuir* **22**, 1595–1601 (2006).

12. W. Ding, M. Fernandino, C. A. Dorao, Conical micro-structures as a route for achieving super-repellency in surfaces with intrinsic hydrophobic properties. *Appl. Phys. Lett.* **115**, 053703 (2019).

13. C. E. Cansoy, H. Y. Erbil, O. Akar, T. Akin, Effect of pattern size and geometry on the use of Cassie–Baxter equation for superhydrophobic surfaces. *Colloids and Surfaces A: Physicochemical and Engineering Aspects* **386**, 116–124 (2011).

14. P. Tsai, R. G. H. Lammertink, M. Wessling, D. Lohse, Evaporation-Triggered Wetting Transition for Water Droplets upon Hydrophobic Microstructures. *Phys. Rev. Lett.* **104**, 116102 (2010).

15. Y. Fang, G. Sun, Y. H. Bi, H. Zhi, Multiple-dimensional micro/nano structural models for hydrophobicity of butterfly wing surfaces and coupling mechanism. *Sci. Bull.* **60**, 256–263 (2015).

568   16.   A. Sanchez-Monge, J. Rodriguez Arrieta, M. Jimenez-Chavarria, A. Retana-Salazar, Observations
569   on the Ultrastructure and Hydrophobicity of the Wings of Thirteen Neotropical Families of
570   Diptera (insecta) with Comments on Their Flight. *Acta Microsc.* **24**, 111–117 (2015).

571   17.   I. N. Miaoulis, B. D. Heilman, Butterfly thin films serve as solar collectors. *Ann. Entomol. Soc. Am.*
572   **91**, 122–127 (1998).

573   18.   I. C. Cuthill, *et al.*, Disruptive coloration and background pattern matching. *Nature* **434**, 72–74
574   (2005).

575   19.   M. Stevens, C. L. Stubbins, C. J. Hardman, The anti-predator function of "eyespots" on
576   camouflaged and conspicuous prey. *Behav. Ecol. Sociobiol.* **62**, 1787–1793 (2008).

577   20.   D. J. Kemp, Female butterflies prefer males bearing bright iridescent ornamentation. *Proc Royal*
578   *Soc B* **274**, 1043–1047 (2007).

579   21.   S. Berthier, Thermoregulation and spectral selectivity of the tropical butterfly *Prepona meander*:
580   a remarkable example of temperature auto-regulation. *Appl Phys A-Mater* **80**, 1397–1400 (2005).

581   22.   A. Krishna, *et al.*, Infrared optical and thermal properties of microstructures in butterfly wings.
582   *Proc Natl Acad Sci USA* **117**, 1566–1572 (2020).

583   23.   C.-C. Tsai, *et al.*, Physical and behavioral adaptations to prevent overheating of the living wings
584   of butterflies. *Nat Commun* **11**, 551 (2020).

585   24.   W. Nachtigall, Aerodynamische Messungen am Tragfluegelsystem segeinder Schmetterlinge.
586   *Journal of Comparative Physiology A* **54**, 210–231 (1967).

587   25.   N. Slegers, *et al.*, Beneficial aerodynamic effect of wing scales on the climbing flight of butterflies.
588   *Bioinspir. Biomim.* **12**, 016013 (2017).

589   26.   T. Wagner, C. Neinhuis, W. Barthlott, Wettability and contaminability of insect wings as a function
590   of their surface sculptures. *Acta Zoologica* **77**, 213–225 (1996).

591   27.   N. D. Wanasekara, V. B. Chalivendra, Role of surface roughness on wettability and coefficient of
592   restitution in butterfly wings. *Soft Matter* **7**, 373–379 (2011).

593   28.   Y. Fang, *et al.*, "Micro-morphological Models for the Special Wettability of Locust and Moth
594   Wing" in *Proceedings of the 2017 5th International Conference on Mechatronics, Materials,*
595   *Chemistry and Computer Engineering (Icmmcce 2017)*, M. Wang, X. Zhou, Eds. (Atlantis Press,
596   2017), pp. 330–333.

597   29.   P. Perez Goodwyn, Y. Maezono, N. Hosoda, K. Fujisaki, Waterproof and translucent wings at the
598   same time: problems and solutions in butterflies. *Naturwissenschaften* **96**, 781–787 (2009).

599   30.   D. Gomez, *et al.*, Transparency in butterflies and moths: structural diversity, optical properties
600   and   ecological   relevance.   *Ecological   Monographs*   (2021)
601   https:/doi.org/10.1101/2020.05.14.093450 (May 19, 2020).

31.   M. McClure, *et al.*, Why has transparency evolved in aposematic butterflies? Insights from the

largest radiation of aposematic butterflies, the Ithomiini. *Proc Royal Soc B* **286**, 20182769 (2019).

32.   M. Arias, *et al.*, Transparency reduces predator detection in chemically protected clearwing
butterflies. *Funct Ecol* (2019) https://doi.org/10.1111/1365-2435.13315.

33.   M. Arias, M. Elias, C. Andraud, S. Berthier, D. Gomez, Transparency improves concealment in
cryptically coloured moths. *J Evol Biol* **33**, 247–252 (2020).

34.   A. Yoshida, M. Motoyama, A. Kosaku, K. Miyamoto, Antireflective nanoprotuberance array in the
transparent wing of a hawkmoth, *Cephonodes hylas*. *Zoological Science* **14**, 737–741 (1997).

35.   R. H. Siddique, G. Gomard, H. Holscher, The role of random nanostructures for the
omnidirectional anti-reflection properties of the glasswing butterfly. *Nature Communications* **6**
(2015).

36.   C. Pinna, *et al.*, Convergence in light transmission properties of transparent wing areas in
clearwing mimetic butterflies. *bioRxiv*, 2020.06.30.180612 (2020).

37.   A. F. Pomerantz, *et al.*, Developmental, cellular, and biochemical basis of transparency in the
glasswing butterfly Greta oto. *Journal of Experimental Biology* **accepted**, eb237917 (2021).

38.   Y. M. Zheng, X. F. Gao, L. Jiang, Directional adhesion of superhydrophobic butterfly wings. *Soft*
Matter **3**, 178–182 (2007).

39.   G. McHale, S. Aqil, N. J. Shirtcliffe, M. I. Newton, H. Y. Erbil, Analysis of Droplet Evaporation on a
Superhydrophobic Surface. *Langmuir* **21**, 11053–11060 (2005).

40.   M. Reyssat, J. M. Yeomans, D. Quéré, Impalement of fakir drops. *EPL* **81**, 26006 (2007).

41.   D. Quéré, Wetting and Roughness. *Annual Review of Materials Research* **38**, 71–99 (2008).

42.   X. Gao, L. Jiang, Water-repellent legs of water striders: Biophysics. *Nature* **432**, 36–36 (2004).

43.   A. Otten, S. Herminghaus, How Plants Keep Dry: A Physicist's Point of View. *Langmuir* **20**, 2405–
2408 (2004).

44.   G. S. Watson, B. W. Cribb, J. A. Watson, Contrasting Micro/Nano Architecture on Termite Wings:
Two Divergent Strategies for Optimising Success of Colonisation Flights. *PLOS ONE* **6**, e24368
(2011).

45.   M. Sugumaran, Complexities of cuticular pigmentation in insects. *Pigment Cell & Melanoma*
Research **22**, 523–525 (2009).

46.   S. N. Aideo, D. Mohanta, Unusually diverse surface-wettability features found in the wings of
butterflies across Lepidoptera order and evaluation of generic and vertical gibbosity-based
models. *Phys. Scr.* **96**, 085004 (2021).

47.   Q. Wan, *et al.*, Combination of active behaviors and passive structures contributes to the
cleanliness of housefly wing surfaces: A new insight for the design of cleaning materials. *Colloid*
Surf. B-Biointerfaces **180**, 473–480 (2019).

48.   S. H. Sajadinia, F. Sharif, Thermodynamic analysis of the wetting behavior of dual scale patterned

hydrophobic surfaces. *Journal of Colloid and Interface Science* **344**, 575–583 (2010).

49.   D. Bixler, B. Bhushan, Rice- and butterfly-wing effect inspired self-cleaning and low drag

micro/nanopatterned surfaces in water, oil, and air flow. *Nanoscale* **6**, 76–96 (2014).

50.   E. Bittoun, A. Marmur, The Role of Multiscale Roughness in the Lotus Effect: Is It Essential for

Super-Hydrophobicity? *Langmuir* **28**, 13933–13942 (2012).

51.   Y. Li, *et al.*, Biomimetic Random Arrays of Nanopillars and Nanocones with Robust Antiwetting

Characteristics. *J. Phys. Chem. C* **124**, 17095–17102 (2020).

52.   S. N. Aideo, D. Mohanta, Limiting hydrophobic behavior and reflectance response of dragonfly

and damselfly wings. *Applied Surface Science* **387**, 609–616 (2016).

53.   G. S. Watson, S. Myhra, B. W. Cribb, J. A. Watson, Putative functions and functional efficiency of

ordered cuticular nanoarrays on insect wings. *Biophys. J.* **94**, 3352–3360 (2008).

54.   M. Sun, *et al.*, Influence of Cuticle Nanostructuring on the Wetting Behaviour/States on Cicada

Wings. *PLOS ONE* **7**, e35056 (2012).

55.   C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image analysis.

Nat. Methods **9**, 671–675 (2012).

56.   M. A. Stoffel, S. Nakagawa, H. Schielzeth, rptR: repeatability estimation and variance

decomposition by generalized linear mixed-effects models. *Methods Ecol Evol* **8**, 1639–1644

(2017).

57.   D. Gomez, *AVICOL v6. a program to analyse spectrometric data. Free program available from the*

author   upon   request   at   dodogomez@yahoo.fr   or   by   download   from

http://sites.google.com/site/avicolprogram/ (2011).

58.   R Development Core Team, *R: a language and environment for statistical computing* (R

Foundation for Statistical Computing, 2013).

59.   J. Pinheiro, D. Bates, D. DebRoy, D. R. Sarkar, nlme: linear and nonlinear mixed effects models. R

package version 3.1-145 (2020).

60.   T. Guillerme, K. Healy, mulTree: performs MCMCglmm on multiple phylogenetic trees. R package

version 1.3.6. (2019).

61.   J. D. Hadfield, MCMC methods for multi-response generalized linear mixed models: The

MCMCglmm *R* Package. *J Stat Soft* **33** (2010).

62.   A. B. D. Cassie, S. Baxter, Wettability of porous surfaces. *Transactions of the Faraday Society* **40**,

546 (1944).

63.   E. P. Ivanova, *et al.*, Molecular Organization of the Nanoscale Surface Structures of the Dragonfly

Hemianax papuensis Wing Epicuticle. *PLOS ONE* **8**, e67893 (2013).

64.   A. J. B. Milne, A. Amirfazli, The Cassie equation: How it is meant to be used. *Advances in Colloid*

and Interface Science **170**, 48–55 (2012).

Figure 1. Examples of water droplets dropped in the transparent zone: Cassie-Baxter regime (series a) for *Eutresis hypereia* combining erected coloured piliform and lamellar scales and Wenzel regime (series b) for *Neorcarnegia basirei* with a nude membrane. Water droplet evolution is shown at different times: T1 (a', b'), T2 (a'', b''), and at T3 (a''', b''').

Figure 2. Examples of structural strategies in clearwing butterflies. Structural strategy is a combination of scale type (N: no scales, P: piliform scales, L: lamellar scales, PL: combination of piliform scales and lamellar scales), scale insertion (E: erected, and F: flat), and scale colour (C: coloured, and T: transparent). Notice that *Macrosoma albipannosa* has transparent lamellar scales but coloured piliform scales.

Figure 3. Phylogeny and distribution of trait values in the study species, for the forewing. Scale type (column 1), scale insertion on the membrane (column 2), scale colouration (column 3), and structural strategy (columns 4 and 5). For scale type: N=no scales, P=piliform scales, L= lamellar scales PL=combination of piliform scales and lamellar scales. For scale insertion on the membrane: E=erected, F=flat, U=undefined (for absent scales). For scale colouration: C=coloured, T=transparent, U=undefined (for absent scales). The strategy NUU was simplified into N.

712
713
714

Figure 4. Variation in contact angle with evaporation time(A) and with scale density (B) in the transparent and opaque zones. Variation in contact angle with forewing length-to-width ratio (C) and total area (D) in the transparent zone. Structural strategy is a combination of scale type (N: no scales, P: piliform scales, L: lamellar scales, PL: combination of piliform scales and lamellar scales), scale insertion (E: erected, and F: flat), and scale colour (C: coloured, and T: transparent). Superhydrophobic: >150° (above the red line), hydrophobic: <150° and >90°; hydrophilic: <90° (below the blue line). (A) all measurements considered, (BCD) mean CA values for each species, zone, and time. Results are presented in Tables S1 to S3d.

Figure 5. Variations of contact angle with wing zone, microstructure and time, i.e. water droplet size. Structural strategy is a combination of scale type (N: no scales, P: piliform scales, L: lamellar scales, PL: combination of piliform scales and lamellar scales), scale insertion (E: erected, and F: flat), and scale colour (C: coloured, and T: transparent). Superhydrophobic: >150° (above the red line), hydrophobic: <150° and >90°; hydrophilic: <90° (below the blue line). All individuals and droplets were considered. Results are presented in Tables S2 to S3d.

Figure 6. Comparison between contact angle values predicted by the best Cassie-Baxter model for the transparent zone (A) and the opaque zone (B). For the measured values, we included all the values from water droplets characterized as following a Cassie-Baxter regime, i.e. showing a CA>120° at T3. The best model considered the effect of microstructures for all structural strategies (with bending P scales when erect) and an effect of nanostructures, but only for flat lamellar scales in high coverage. Structural strategy is a combination of scale type (N: no scales, P: piliform scales, L: lamellar scales, PL: combination of piliform scales and lamellar scales), scale insertion (E: erect, and F: flat), and scale colour (C: coloured, and T: transparent). The dashed black line represents perfect agreement between prediction and measurement. We considered all the measured water droplets, and for time T1.

Figure 7. Variations of contact angle with wing transmittance for the different structural strategies. Structural
strategy is a combination of scale type (N: no scales, P: piliform scales, L: lamellar scales, PL: combination of
piliform scales and lamellar scales), scale insertion (E: erected, and F: flat), and scale colour (C: coloured, and
T: transparent). Superhydrophobic: >150° (above the red line), hydrophobic: <150° and >90°; hydrophilic:
<90° (below the blue line). We considered only the mean of CA for each species, for time T1, and for the
transparent zone. The black plain line indicates the significant fitted regression line based on the Bayesian
model. Results are presented in Table S5.

Figure 8. Relationship between contact angle in the transparent (A) and in the opaque (B) zone and the distance in latitude to the equator. Structural strategy is a combination of scale type (N: no scales, P: piliform scales, L: lamellar scales, PL: combination of piliform scales and lamellar scales), scale insertion (E: erected, and F: flat), and scale colour (C: coloured, and T: transparent). Superhydrophobic: >150° (above the red line), hydrophobic: <150° and >90°; hydrophilic: <90° (below the blue line). The black plain line in A indicates the significant fitted regression line based on the Bayesian model. Results are presented in Table S6.