Nipah virus Bangladesh infection elicits organ-specific innate and inflammatory responses in

the marmoset model

Christian S. Stevens[1], Jake Lowry[2], Terry Juelich[3], Colm Atkins[3,#], Kendra Johnson[4], Jennifer K Smith[3],

Maryline Panis[1], Tetsuro Ikegami[3], Benjamin tenOever[1], Alexander N. Freiberg[3], Benhur Lee[1]

[1]Icahn School of Medicine at Mount Sinai, Department of Microbiology

[2]University of Texas Medical Branch, Animal Resource Center

[3]University of Texas Medical Branch, Department of Pathology

[4]University of Texas Medical Branch, Department of Microbiology and Immunology

[#]Present address: Rutgers University, Department of Cell Biology and Neuroscience

Abstract. The common marmoset (*Callithrix jacchus*) has, in recent years, received more recognition as

an ideal non-human primate (NHP) model not only for studies related to neuroscience, but also at high-

biocontainment due to its smaller size and relative ease of handling. Here, we evaluated the susceptibility

and pathogenesis of Nipah virus Bangladesh strain (NiV_B) infection in marmosets at biosafety level 4.

Four marmosets were infected via the intranasal and intratracheal route and monitored for disease

development. We obtained histopathological data, quantified genome copies on >25 tissue-types, and

performed RNA-seq on six different organs from all infected and control marmosets. All four subjects

developed fatal disease between days 8 and 11 post infection, with three animals developing pulmonary

edema and hemorrhage as well as multi-focal hemorrhagic lymphadenopathy. One subject recapitulated

neurologic clinical symptoms and cardiomyopathy on gross pathology. In all animals, syncytia were

evident in endothelial cells in pulmonary vessels, cells in alveolar septum, and in splenic follicles or red

pulp. To define the organ-specific innate and inflammatory responses, we performed RNA-seq on six

different organs from all infected and compared to naïve non-infected marmoset tissues. RNA-seq gave

17.4 million reads per sample on average, with the most highly infected tissue sample, the lung of one

marmoset, containing >180,000 virus-specific reads. NiV V and W transcripts comprised ~40% of all

phosphoprotein-derived transcripts with V and W being very close to each other proportionally. Principal

component analysis showed that in brain stem, the marmoset exhibiting neurological symptoms

displayed a unique RNA transcriptome relative even to other infected marmosets. Upregulated genes in

the various tissues belonged to distinct GO pathways. Additionally, one male and female animal had

detectable viral reads in their ovaries (27,120) and testes (858). Our results provide a more

comprehensive understanding of NiV pathogenesis in an accessible and novel NHP model, closely

reflecting clinical disease as observed in NiV patients.

Introduction

Nipah virus (NiV), a zoonotic, bat-borne negative-sense paramyxovirus of the genus *Henipavirus*, first emerged in Malaysian 1998/1999 during an outbreak in pigs [1]. Since then, outbreaks have occurred almost every year, resulting in over 600 cases of respiratory and/or neurological disease in patients, with 59% percent of which resulted in death [2]. Two major clades, NiV-Malaysia (NiV_M) and NiV-Bangladesh (NiV_B), have since emerged. All but two outbreaks have been NiV_B, which has had a higher case fatality rate (CFR) of 76% relative to NiV_M which has had a CFR of 39% [2]. In addition to a higher observed CFR, NiV_B also has a shorter average incubation period, a higher rate of respiratory symptoms, including patients developing an acute respiratory distress syndrome, and a smaller proportion of patients exhibiting segmental myoclonus relative to NiV_M [3–5]. Another important difference has been the extent of human-to-human transmission, common in NiV_B outbreaks but extremely rare in NiV_M outbreaks [6]. Given these significant differences, and that NiV_B has made up all but two outbreaks to date, animal models successfully recapitulating NiV_B pathology in humans are especially critical.

Many animal models have been used to investigate NiV pathogenesis and vaccine research including cats (*Felis catus*), ferrets (*Mustela putorius furo*), golden hamsters (*Mesocricetus auratus*), guinea pigs (*Cavia porcellus*), pigs (*Sus scrofa domesticus*), and mice (*Mus musculus*) [7–19]. Additionally, several non-human primate (NHP) models have been explored: African green monkeys (*Chlorocebus aethiops*; AGM), cynomolgus macaques (*Macaca fascicularis*), squirrel monkeys (*Saimiri sciureus*), and grivets (*Chlorocebus aethiops*) [20–34]. Among these, only AGMs have been able to faithfully recapitulate the high lethality of NiV_B exposure as well as most other clinical signs. However, overt clinical signs of neurological disease are seen very rarely in AGMs despite neurological disease being commonly seen via MRI and post-mortem histopathology. Additionally, myocarditis has been consistently reported among patients infected with NiV_B but to date there have been no reports of myocarditis in any animal models [35].

NiV infection in NHP models consistently replicate much of the hallmark clinical pathology and histopathology of NiV infection. Systemic vasculitis, hypoalbuminemia, pneumonitis, coagulopathy, and congestion in the liver and spleen are commonly seen in AGMs [21,22], although models such as the cynomolgus macaque are unable to replicate many of these signs [25]. Transcriptional profiling in AGMs has also shown significant upregulation in innate immune signaling genes (*MX2, OASL, OAS2*), particularly among non-survivors of infection relative to survivors [25]. Others have shown that NHP models show an especially robust expression of interferon stimulated genes in the lung, including *ISG15* and *OAS1*, [36]. Additionally, the success of the AGM model of NiV infection has given insight into the peripheral immune response to infection. For example, while CD8+ T cell activation and proliferation markers are drastically increased during infection, increased natural killer cell response was correlated with improved survival [30].

Although AGMs are the current standard NHP model of NiV infection, they suffer from a few shortcomings. Thus far there are few reports of overt clinical signs of neurological disease and no reports of myocarditis in NiV_B-infected AGMs. Additionally, given the biosafety level 4 (BSL4) requirements of working with NiV, AGMs large size and associated costs are historically a significant barrier. In order to diversify the preclinical testing portfolio of NHP models, we show here that NiV_B infection of the common marmoset (*Callithrix jacchus*) can faithfully recapitulate pathology seen in humans. Marmosets were chosen because they have several distinct advantages. First, they are an ideal NHP model for studying high-biocontainment human respiratory pathogens because of their small size—approximately 20-30 centimeters in body length and 250-400 grams in body weight—and relative ease of handling. Second, they are the first among New World monkeys to have their whole genome sequenced and assembled allowing for techniques like RNA-sequencing (RNA-seq) to elucidate RNA expression changes in infected and uninfected tissues [37]. Lastly, marmosets provide other logistical advantages such as the ability to perform transmission experiments given that they can be co-housed and a high reproductive efficiency

91   with a short gestation period and delivery interval allows for easier development of transgenic animals

92   [38]. With these advantages in mind, results of this study identified marmosets as a highly susceptible

93   novel new-world NHP model of NiV_B infection and pathogenesis.

Material and Methods

Ethics statement. Experiments were approved by the Institutional Animal Care and Use Committee of the University of Texas Medical Branch (UTMB) and performed following the guidelines of the Association for Assessment and Accreditation of Laboratory Animal Care International (AAALAC) by certified staff in an AAALAC-approved facility. Animals were co-housed in primate cages allowing social interactions, under controlled conditions of humidity, temperature and light (12-hour light/12-hour dark cycles). Animals were monitored pre- and post-infection and fed commercial monkey chow, treats and fruit twice daily. Food and water were available ad libitum and environmental enrichment consisted of commercial toys. All procedures were conducted by BSL4-trained personnel under the oversight of an attending BSL4-trained veterinarian. All invasive procedures described here were performed on animals under isoflurane anesthesia. Animals that reached euthanasia criteria were euthanized using a pentobarbital-based euthanasia solution.

Virus and cells. Nipah virus (Bangladesh strain, NiV_B; 200401066 isolate) was kindly provided by Dr. Thomas Ksiazek (UTMB). The virus was propagated on Vero E6 cells (ATCC, CRL1586) in Dulbecco's minimal essential medium (DMEM) supplemented with 5% fetal bovine serum (HyClone, Logan, UT), penicillin, and streptomycin at 37°C in a humidified CO_2 incubator (5%). Virus titers were determined by plaque assay on Vero cells (ATCC, CCL-81) as previously described [39]. All work with infectious virus was performed at BSL4 biocontainment at the Galveston National Laboratory (GNL) at UTMB.

Animal studies. Four healthy marmosets (2 females, 2 males; 348 to 453 gram) were inoculated with 6.33×10^4 plaque forming units (PFU) of NiV_B with the dose being equally divided between the intranasal (IN) and intratracheal (IT) routes for each animal. After infection, animals were closely monitored for signs of clinical illness, changes in body temperature (BMDS transponders) and body weight. A scoring system was utilized monitoring respiratory, food intake, responsiveness, and neurological parameters to

determine study endpoint. In addition, changes in blood chemistry and hematology were monitored. On days -7, 1, 3, 5, 7, and 10 days post infection, blood was collected via the femoral vein and swabs (oral, nasal, rectal).

Hematology and serum biochemistry. Blood was collected in tubes containing EDTA and complete blood counts (CBC) were evaluated with a VetScan HM5. (Abaxis, Union City, CA). Hematological analysis included total white blood cell counts, white blood cell differentials, red blood cell counts, platelet counts, hematocrit values, total hemoglobin concentrations, mean cell volumes, mean corpuscular volumes and mean corpuscular hemoglobin concentrations. Clinical chemistry analysis was performed using a VetScan2 Chemistry Analyzer (Abaxis, Union City, CA). Serum samples were tested for concentrations of albumin (ALB), amylase (AMY), alanine aminotransferase (ALT), alkaline phosphatase (ALP), glucose (GLU), total protein (TP), total bilirubin (TBIL), plasma electrolytes (calcium, phosphorus, sodium, and potassium, blood urea nitrogen (BUN), creatinine (CRE) and globulin (GLOB)).

X-ray images. X-ray images were obtained using the MinXray model HF 100/30 portable x-ray unit (Northbrook, IL) operated at 78 kVp 12 3.20 mAs at a distance of 40 inches from the subject. A Fuji DR 14x17 inch panel served as the exposure plate and the images were processed in the SoundVet application (Carlsbad, CA).

Sample collection and RNA isolation. Nasal, oral, and rectal swabs were collected in 1 ml of Dulbecco's minimal essential medium (DMEM) (Sigma-Aldrich, St. Louis, MO) and vortexed for 30 s. Whole blood was collected in EDTA Vacutainers (Beckman Dickinson, Franklin Lakes, NJ) for hematology and serum biochemistry, while another part was mixed with TRIzol LS for RNA extraction. At time of euthanasia, full necropsies were performed, and tissues were homogenized in TRIzol reagent (Qiagen, Inc.) using Qiagen TissueLyser and stainless-steel beads. The liquid phase was transferred into a new cryovial and stored at -80°C. The tissues sampled include bronchial lymph node, mandibular lymph node, mediastinal

lymph node, inguinal lymph node, salivary gland lymph node, axillary lymph node, mesenteric lymph node, femoral bone marrow, tonsil, oro/nasopharynx, nasal mucosa, trachea, lung (left upper lobe, left lower lobe, right upper lobe, right lower lobe), right bronchos, left bronchos, heart, liver, spleen, kidney, adrenal gland, pancreas, jejunum, colon transversum, brachial plexus, brain (frontal, cerebellum, stem), cervical spinal cord, pituitary gland, urinary bladder, and testes/ovaries. All blood samples and swabs were inactivated in TRIzol LS and TRIzol reagent, respectively, prior to removal from the BSL4 laboratory. Subsequently, RNA was isolated from blood and swabs using the QIAamp viral RNA kit (Qiagen, Inc.) and from tissues using the RNeasy minikit (Qiagen, Inc.) according to the manufacturer's instructions supplied with each kit.

RT-qPCR. RT-qPCR was used to quantitate viral loads in tissues, swabs and blood. RNA was extracted from tissues homogenized in TRIzol reagent or from blood and swabs mixed with TRIzol LS reagent using Direct-zol RNA Miniprep kits (Zymo Research, Irvine, CA). RT-qPCR assays were then run using QuantiFast RT-PCR mix (Qiagen, Hilden, Germany), probes targeting NiV_M P gene (5′-ACATACAACTGGACCCARTGGTT-3′ and 5′-CACCCTCTCTCAGGGCTTGA-3′) (IDT, Coralville, IA), and fluorescent probe (5′-6FAM-ACAGACGTTGTATA + C + CAT + G-TMR) (TIB MOLBIOL, Adelphia, NJ). RT-qPCR was performed using the following cycle: 10 minutes at 50 °C, 5 minutes at 95 °C, and 40 cycles of 10 seconds at 95 °C and 30 seconds at 60 °C using a BioRad CFX96 real time system. Assays were run in parallel with uninfected hamster tissues and a NiV_M stock standard curve.

Virus titration. Virus titration was performed by plaque assay on Vero cells as previously described [39].

Histopathology and immunohistochemistry. Necropsies were performed on euthanized animals, and tissues collected for histopathology and immunohistochemistry. Tissues were immersion-fixed in 10% neutral buffered formalin for at least 21 days under BSL-4 conditions with regular formalin changes. Specimens were then processed under BSL-2 conditions. Briefly, tissues were dehydrated through a

series of graded ethanol baths, infiltrated by and embedded with paraffin wax, sectioned at 5 µm

thickness, and stained with hematoxylin and eosin. Immunohistochemistry for NiV nucleoprotein

detection was performed using a rabbit anti-NiV-nucleoprotein antibody incubated overnight (kindly

provided by Dr C. Broder, Uniformed Services University, Bethesda, Maryland) and a secondary

horseradish peroxidase (HRP)-conjugated goat antirabbit antibody (Abcam) incubated for 2 hours (both

1:1000). DAB substrate (ThermoScientific) was then added for 2 minutes to enable chromogenic

detection (brown precipitate) of HRP activity.

RNA sequencing (RNA-seq). Polyadenylated RNA libraries were prepared using the TruSeq RNA

Library Prep Kit v2 (Illumina) or TruSeq Stranded mRNA Library Prep Kit (Illumina) according to the

manufacturer's instructions. Libraries were sequenced on an Illumina NextSeq 500 platform using a

paired end 2 x 150-base pair, dual index format. Across all five marmosets (#193, #300, #340, #425, and

naïve) six tissue samples were sequence (lung, inguinal lymph node, kidney, gonads, brain stem, and

spinal cord) except gonads in the naïve marmoset. For the other five tissues in the naïve marmoset, all

samples were sequenced in technical duplicate. Tissues from naïve marmosets were kindly provided by

Dr. Ricardo Carrion Jr (Texas Biomedical Research Institute).

RNA-seq data processing. Trimming, alignment, and quantification were all performed in the Partek

Flow v10.0 [40]. Raw sequencing reads were trimmed on both edges based on quality score. Paired

reads were then aligned to the *Callithrix jacchus* genome (Ensemble release 91) [41] using Bowtie2

v2.2.5 with default parameters. Counts were generated using the Partek Quantify to Annotation Model

(Partek E/M) *Callithrix jacchus* annotation v105. Data normalization was performed by trimmed mean of

M values (TMM) adding 1.0E-4 (default parameters). Samples were filtered if the number of aligned reads

was less than 1 million mapped reads to genes or if they were determined to be outliers due to PCA. In

total, six samples were removed for the final analysis. Differential gene expression analysis was

performed using Partek's gene specific analysis (GSA) differential expression model.

Reads that did not map to the *Callithrix jacchus* genome were then aligned to the Nipah virus Bangladesh

genome (GenBank: AY988601.1) using the BWA-MEM algorithm within BWA v0.7.17 with all default

parameters. Transcriptional gradient and single nucleotide polymorphism analyses of the NiV_B

sequences was performed using IGVtools with a minimum read coverage of 100 per position and a

minimum variant frequency of 5%. Insert counts at the P editing site were calculated using mPileup in

Samtools v1.8 [42], selecting only for inserts of Gs and a minimal coverage of 100 reads per sample.

Statistical analysis, gene set analysis, and visualization. PCA was performed using the default

settings within Partek Flow v10.0. Functional enrichment analysis was performed using PantherDB's

Biological Processes Gene Ontology (GO) annotation set [43]. Submitted gene lists were acquired via

Partek's gene specific analysis (GSA) differential expression model. Lists were filtered using an FDR

step-up value of <0.05 and an absolute value of fold change >2. Gene lists were analyzed via the

PANTHER Overrepresentation Test, annotation version release data 2021-08-18. Because there are no

gene ontology sets for *Callithrix jacchus*, the *Homo sapiens* GO biological process complete annotation

data set was utilized. Only GO terms where FDR < 0.01 were used. GO terms along with accompanying

FDR were clustered and visualized using REVIGO [44].

Results

NiV$_B$ infection and pathogenesis in marmosets. In the studies described here, we evaluated the susceptibility and suitability of the common marmoset to infection with NiV$_B$. Four marmosets were infected with NiV$_B$ by the IN/IT routes and throughout the course of the study, survival, body weights, body temperature, blood chemistry and complete blood count (CBC) were determined (**Supplementary Figures S1, S3, and S4**). All four animals developed clinical signs of disease and reached euthanasia criteria between days 8 and 11 post infection (**Table 1**). Throughout the first seven days of the study, all subjects were normal, and only two animals (#'s 193 and 300) had loose stool on day 6 post-infection (PI). By day 8 PI, subjects #300 and #425 developed hyperventilation and did not consume their food. Subject #425 became lethargic at the end of day 8 and reached euthanasia criteria. Subject #300 started to deteriorate on day 9 PI, when found with open mouth breathing and reached euthanasia criteria the same day. Subject #193 had a reduced food consumption on days 9 and 10 PI, and on day 11 PI was found to be hyperventilating and lethargic and reached euthanasia criteria. Subject #340 was normal through day 10 PI, then developed a hunched posture, and on day 11 PI was hyperventilating and reached euthanasia criteria. Furthermore, this subject presented left hindlimb tremors during euthanasia, reflecting signs of neurological impairment. Over the course of the study, no significant changes in body weight were detected, as well as body temperature (**Supplementary Figure S1**).

X-ray autoradiography and gross pathology. To monitor development of respiratory disease, we used X-ray radiographic imaging (**Figure 1A and Supplementary Figure S2**). Over the course of the study, a decrease in normal lung tissue was observed. Prior to day 7, no radiographic changes were observed in the lungs and comparable to day 0. At day 7 PI, lungs were either still clinically normal (#193, #340), or some decreasing opacity was observed along with an appearance of an interstitial pattern (#300) and some inflammation around the smaller bronchi with a mild interstitial infiltrate (#425). At the time of euthanasia, virtually no normal lung was observed for animals #425 (euthanized day 8 PI), #300

(euthanized day 9 PI) and #193 (euthanized day 11 PI), as the lungs developed a nodular alveolar

pattern, suggesting the alveoli were filled with fluid. In animal #340 (euthanized day 11 PI), a mild

bronchial pattern emerged at the time of euthanasia, showing some inflammation around the smaller

bronchi.

The primary gross pathology finding identified in marmosets that reached euthanasia criteria after

infection with NiV$_B$ was the presence of minimal normal lung tissues (< 5-10%) in the cranial portion of

the lung (**Figure 1B and Table S1**). The remaining of the lung was fluid filled with a hemorrhagic/serous

fluid. In one animal (#340), the right lung was normal but the bronchus fluid filled. The livers had multi-

focal to coalescing regions of pale circles and were very friable (**Figure 1C**). Lymph nodes (mediastinal,

bronchial, inguinal, and/or retropharyngeal) had gross enlargement and hemorrhage. In one animal

(#425), the spleen was friable and congested. Interestingly, animal #340, which presented with a less

severe infection of the lungs, neurological involvement was detected. An infarct was found, as well as

malacia near the transition zone between grey and white matter in the occipital lobe of the left cerebral

hemisphere. In addition, edema was present in the cerebrum. According to these observations,

marmosets developed diffuse pulmonary edema and hemorrhage (3 out of 4 animals), diffuse

hemorrhagic hepatitis (4 out of 4 animals), multi-focal hemorrhagic lymphadenopathy. One animal (#340),

had left ventricular cardiomyopathy and neurological complications.

Histopathological changes in NiV$_B$ infected marmosets. Histopathological and immunohistochemical

examination of the lungs showed that all animals had mild to moderate infiltrations of mononuclear cells

and neutrophils in pulmonary vessels (**Figure 2A**). Subject #300 additionally showed pulmonary edema

with scattered presence of alveolar macrophages (data not shown). Syncytia formations were evident in

infected lungs (endothelial cells in pulmonary vessels or cells in alveolar septum) in all four animals. NiV

antigen was present in endothelial cells or smooth muscle cells of pulmonary arterioles or venules, as

well as in cells composing the pulmonary septa, most likely capillary endothelial cells (**Figure 2B**). In the

liver, all animals showed moderate to severe multifocal coagulation necroses affecting the midzonal area.

Mild to moderate portal inflammation with mononuclear cells and neutrophils were also evident (**Figure 2C**). Viral antigen was detected in necrotic hepatocytes and endothelial cells of the portal vein or sinusoid (**Figure 2D**). In the spleens, the red pulp was partly necrotic in subject #340, while all animals retained follicles without notable necrosis in white pulps. Syncytia formation was evident in mononuclear cells in follicles or in red pulps (**Figure 2E**). NiV antigen was detectable in mononuclear or multinuclear syncytial cells inside or surrounding follicles (**Figure 2F**). In the kidney, loss of endothelial and mesangial cells in affected glomeruli was observed, along with syncytia formations of endothelial cells in the medullary peritubular capillaries, cells constituting glomeruli, and the parietal epithelial cells of the Bowman's capsules (**Figure 2G**). NiV nucleoprotein was detectable in glomerular endothelial cells, and peritubular capillary endothelial cells (**Figure 2H**). Although there were no detectable lesions in the histopathological slides, viral antigen was found in the glomerular or peritubular capillary endothelial cells in subjects #300 and #340. Furthermore, histopathological lesions in subject #425 suggests myocarditis associated with the infiltration of mononuclear cells and neutrophils (**Figure 2I**). Viral antigen was found in the heart smooth muscle cells and capillary or arteriole endothelial cells (**Figure 2J**). While no evident lesions were found in the histopathological slides, viral antigen was also detected in the in heart smooth muscle cells and capillary or arteriole endothelial cells of subject #300. No detectable lesions or viral antigen were found in the testes (subjects #425, #193), bronchus or CNS of all subjects.

Blood chemistry and hematology. Over the course of the study, changes in CBC and blood chemistry were monitored at days 1, 3, 5, 7, and 10 PI (**Supplementary Figures S2 and S3**). Changes over time for specific parameters were compared to values obtained at day 1 PI, due to the fact that no uninfected control animal was included in this study. While no change was observed in white blood cell counts for subjects #300, 340, and 425, an increase at day 10 PI was measured for subject #193 one day before euthanasia, indicative for leukocytosis (**Supplementary Figure S2**). Neutrophilic and eosinophilic granulocytosis was noted for subjects #193, 300 and 340, and subjects #193, 300, and 425, respectively. These changes were observed imminent before euthanasia. Additionally, basophilia was noted for

subjects #193 and 340 starting 1 DPI. No obvious changes were noted in hematocrit and platelet counts, except for subject #425, for which an increase was observed at day 7 PI, one day prior to euthanasia, possibly indicating reactive thrombocytosis.

A decreased trend in albumin levels were detected between days 1 and 5 PI for all four subjects, followed by a sudden increase for animals #193 and #425 at day 7 PI, while levels for the other two subjects stayed leveled. At 10 DPI a continued downward trend was observed again for the remaining two animals, #193 and #340, just before euthanasia. Elevations in alkaline phosphatase at day 10 PI for subjects #193 and 340 were suggestive of hepatocyte damage. Similarly, a drastic increase in alanine aminotransferase was noted at days 7 and 10 PI for animals #425 and 193, respectively. Amylase levels showed a decreased trend over the course of the study for all four subjects, indicative for a damage in the pancreas. An increase in creatinine levels was noted for subject #300 at day 7 PI and is suggestive for impaired kidney function. For subject #193, a sudden decrease in creatinine levels was noted at day 10 PI, just before euthanasia and could be indicative for liver damage. A temporary sudden increase in potassium levels were measured for subjects #193 and 425 at day 7 PI. For animal #425, a sudden drop was detected in MCH and MCHC, and an increase in RBC, and PCT at 7 DPI potentially indicating acute hemorrhage. For animal #193, a temporary increase in monocytes and a drop in MCV was noted at 7 DPI.

Viral titers in tissues. Tissues were collected at time of euthanasia and viral titers determined by plaque assay (**Figure 3A**). Virus was found in 19 out of 23 tissues evaluated but not consistently for the four marmosets. Virus was found in the kidney and adrenals, as well as lower respiratory tract for three marmosets, followed by the spleen, liver, and genitourinary system. Other tissues where virus was detected in marmosets included the tonsil, upper respiratory tract, heart, gastrointestinal tract, and CNS. When comparing tissues that were positive for virus between the four animals, #425 had detectable NiV_B in 7 out of 10, and #300 in 6 out of the 10 tested tissue groups, while four tissues were positive for #193, and one for #340. Infectious virus was found in both the brain stem and the cerebellum of #300.

Viral RT-qPCR in tissues. RNA was extracted from 23 tissues and RT-qPCR was performed (**Figure 3B**). Viral genome was detected in all tissues evaluated, however not for all four marmosets. Viral RNA is consistently lowest for #193 but the two tissues where #193 shows the lowest Ct correspond to the tissues where #193 showed the highest viral titers (liver and spleen). As expected, viral load is highest across the board in tissues of the respiratory tract. Despite only one subject being positive for infectious virus in the CNS, viral RNA was found in at least one of these tissues for all marmosets except #193.

RNA-seq of NiV$_B$ in infected marmoset tissues. RNA was extracted from six different tissues across all infected and uninfected marmosets: spinal cord, brain stem, inguinal lymph node, gonads, kidney, and lung. The average tissue sample had 17.4 million reads aligned to *Callithrix jacchus*. The most highly infected tissue (#425 lung) contained >180,000 virus-specific reads and 15 samples had >100 reads aligned to NiV$_B$ (**Figure 4A**). Total reads aligned to NiV$_B$ was highly driven by overall sequencing depth, however lung tissue showed the highest number of viral read counts for all animals whose lung tissue sample passed quality controls.

Across all samples, reads were counted per gene, and we see that NiV in the marmoset model shows a transcriptional gradient where the fraction of total reads aligned to a particular gene does not drop off significantly until after N, P, and M (**Figure 4B**). In spite of the steep transcriptional gradient, three samples achieved >95% coverage of L: #425 lung, #300 lung, and #300 gonads. Additionally, we were able to filter all samples with >100 read coverage at the RNA editing site in P and calculate the number of G inserts, with the largest number found in any individual read being 12 Gs (**Figure 4C**). Approximately 40% of all P-derived reads are V and W transcripts (**Figure 4D**), with proportions relatively equal but varying somewhat between tissues and animals (not shown). Given the high depth of coverage we were also able to identify several singe nucleotide polymorphisms (SNPs) across the NiV$_B$ genome at rates >5% of all reads when filtering for positions with a read depth > 100 reads (not shown).

RNA-seq samples aligned to *Callithrix jacchus*. Across all samples, an average number of 17.4 million

reads per sample aligned to the marmoset genome. Across all samples, we first performed differential

expression analysis comparing samples from NiV_B infected and uninfected animals. Only genes with an

FDR step up < 0.05 and an absolute value of the fold change > 2 were chose for gene ontology gene set

enrichment analysis. The resulting GO biological process terms were then clustered in semantic space

based on shared and related genes (**Figure 5A**). We see that a few clusters form with the two most

notable and expected dealing primarily with regulation of viral processes and interferon responses.

However, other clusters also emerge such as those dealing with blood circulation and angiogenesis.

We performed the same analysis on specific tissues as well, first looking at the lungs. From this we

identified the top 25 most significant differentially expressed genes, meeting the same filtering thresholds

described previously. We then visualized the differential expression using a hierarchically clustered

heatmap (**Figure 5B**). Here we see expected interferon stimulated genes such as ISG15, ISG20, OAS3,

OASL, and IFNB1 upregulated in infected tissues. However, we see only a single gene downregulated

relative to the naïve samples make the top 25 cutoff: SCGB3A2. Secretoglobin (SCGB) 3A2 is known to

have anti-inflammatory properties, especially in allergic airway inflammation models [45] although its role

in viral infections has not been previously shown.

We also performed principal component analysis of brain stem samples from infected and uninfected

animals (**Figure 5C**). One sample stands as a clear outlier, the brain stem from marmoset #340, the only

marmoset to have shown overt neurologic clinical symptoms (blue sphere). Due to the unique clinical

symptoms and the results of the PCA we performed differential expression analysis comparing the brain

stem sample from #340 to the other three infected brain stem samples (#193, #300, and #425). Filtering

genes using the same thresholds as above, then extracting enriched GO biological process terms, and

mapping the interconnectedness of the resulting GO terms, we found a large, interconnected graph of

371   unique and related terms (**Figure 5D**). The largest interconnected tree graph shows a family of GO terms

372   relating to myelination, oligodendrocyte development, and glial cell development. These results are

373   noteworthy given previous findings of significant myelin degradation and axonal NiV aggregation in

374   hamsters infected with NiV_M [46].

Discussion

In this study we identify marmosets as a highly susceptible novel new-world NHP model of NiV$_B$ infection. Four marmosets, two male and two female, were infected with NiV$_B$ via the IN/IT routes and all four animals succumbed to infection by day 11 PI. The marmosets recapitulated many of the same clinical symptoms seen in patients and other NHP models infected with NiV, such as anorexia, lethargy, and hyperventilation. Of special note, one marmoset, subject #340, presented with left hindlimb tremors on day 11 PI, displaying overt neurological clinical symptoms. In the setting of acute infection with NiV$_B$, neurological clinical symptoms are rare [34] although neurological disease seen on MRI, in histopathology, or in gross pathology is relatively common [22,24,27,31,33]. However, in infections with NiV$_M$, neurological clinical symptoms are more common, especially among NHPs surviving longer [21,28,29]. This may give some insight into the development of neurological clinical symptoms in the only marmoset (#340) that presented a less severe infection of the lungs, with all other animals having <5-10% normal lung tissue following euthanasia. #340 was found to have a cerebral infarct, malacia in the occipital lobe of the left cerebral hemisphere, and cerebral edema. This animal, uniquely, also displayed left ventricular cardiomyopathy.

On histopathological examination, all marmosets were found to have significant multifocal coagulation in the liver, syncytia formation of endothelial cells in the kidney, and infiltration in the lung. Immunohistochemistry showed NiV antigen in syncytial cells across multiple tissues, notably in the smooth muscle cells and capillary endothelial cells of the heart. No detectable viral antigen was found in the CNS of all subjects. Blood chemistry and hematology showed consistent hypoalbuminemia across all animals, reactive thrombocytosis in marmoset #425, decreases in amylase indicating pancreatic damage across all marmosets, and increased creatinine in subject #300 indicating impaired kidney function. These findings replicate the clinical and pathophysiological picture of NiV infection in NHPs.

Especially importantly is the finding not only of neurological disease, but a marmoset with neurological

clinical symptoms following NiV$_B$ infection.

Viral titers were performed showing titerable virus from all tissue sets evaluated, but some variance was

seen across subjects. Infectious virus was also tittered from both the brain stem and cerebellum of #300.

Performing RT-qPCR showed that viral RNA was detectable in all tissues tested but not in all marmosets.

Viral RNA was most prevalent in the lower respiratory tract, as expected, but was also found in at least

one tissue each within the CNS for #300, 340, and 425.

RNA-seq was performed providing deeper insight into both the NiV and marmoset transcriptional patterns

across multiple tissues. The number of reads aligned to the *Callithrix* jacchus genome was high across

all tissues, but the number of reads aligning to NiV$_B$ was highly variable. More highly infected tissues,

especially the lungs, tended to have the highest number and proportion of NiV reads. Additionally, RNA-

seq of NiV$_B$ showed a transcriptional gradient that did not drop significantly until after matrix, a pattern

not typically seen among other viruses of the family *Paramyxoviridae* [47,48]. We also showed that RNA

editing of P results in very similar amounts of W and V, in contrast to the steep drop from P often seen in

other paramyxoviruses [49,50]. However, others have shown this same pattern in NiV$_B$ with

approximately 40-50% of reads being P while the remaining reads are nearly evenly split between V and

W [51]. Similarly, we replicated *in vivo* the findings that the number of G additions is nearly flat from +1

to +5 as seen in 293 cells [51].

By looking at the marmoset transcriptome, across all tissues we observed a consistent up-regulation of

gene ontologies relating to regulation of viral processes and interferon stimulated genes. We found

results mirroring those found in AGMs showing especially significant upregulation of genes such as *MX2,*

OASL, CCL2, and *ISG15* in the most highly infected tissue, the lung. When evaluating the tissue-specific

transcriptomic responses to infection, the principal component analysis of brain stem samples showed

that #340 was a significant outlier relative to all other samples. When exploring the differential expression

in the sample from #340 we found a particular enrichment in genes relating to myelination, axon

ensheathment, and oligodendrocytes relative to other infected marmosets. Previous findings of myelin

degradation and axonal NiV antigen aggregation in hamsters may be related findings to what we observe

here in marmosets [46].

Marmosets may provide a new NHP model to study severe respiratory disease caused by NiV. These

findings are of particular note given many of the logistical advantages afforded by working with

marmosets in the context of high-biocontainment human respiratory pathogens like NiV. Marmosets offer

distinct advantages to the current standard NHP model of NiV infection, AGMs, due to their small size

and ease of handling, typically lower costs, and ease of co-housing allowing for a potential model of

transmission studies. Given the unique transcriptional profile in the brain stem of the only marmoset that

showed neurologic clinical symptoms, there is a need for additional follow up with a larger cohort number

of both infected and uninfected marmosets. More detailed pathogenicity studies are required to monitor

levels of viremia during the course of disease development, as well as evaluate different routes of

infection to potentially allow in depth studying of neuroinvasion. Overall, marmosets were highly

susceptible to infection via the IN/IT route resulting in a 100% mortality in non-protected animals.

References

1. Looi L-M, Chua K-B. Lessons from the Nipah virus outbreak in Malaysia. Malays J Pathol
    [Internet]. **2007**; 29(2):63–7. Available from: http://www.ncbi.nlm.nih.gov/pubmed/19108397

2. Howley PM, Knipe DM (David M, Whelan SPJ, Fields BN. Fields virology. Volume 1, Emerging
    viruses . Seventh ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2021.

3. Goh KJ, Tan CT, Chew NK, et al. Clinical Features of Nipah Virus Encephalitis among Pig
    Farmers in Malaysia. N Engl J Med [Internet]. Massachusetts Medical Society; **2000**;
    342(17):1229–1235. Available from: https://doi.org/10.1056/NEJM200004273421701

4. Hossain MJ, Gurley ES, Montgomery JM, et al. Clinical presentation of nipah virus infection in
    Bangladesh. Clin Infect Dis an Off Publ Infect Dis Soc Am. United States; **2008**; 46(7):977–984.

5. Chadha MS, Comer JA, Lowe L, et al. Nipah Virus-associated Encephalitis Outbreak, Siliguri,
    India. Emerg Infect Dis [Internet]. **2006**; 12(2):235–240. Available from:
    http://wwwnc.cdc.gov/eid/article/12/2/05-1247_article.htm

6. Homaira N, Rahman M, Hossain MJ, et al. Nipah virus outbreak with person-to-person
    transmission in a district of Bangladesh, 2007. Epidemiol Infect [Internet]. **2010**; 138(11):1630–
    1636. Available from:
    https://www.cambridge.org/core/product/identifier/S0950268810000695/type/journal_article

7. Mungall BA, Middleton D, Crameri G, et al. Vertical Transmission and Fetal Replication of Nipah
    Virus in an Experimentally Infected Cat. J Infect Dis [Internet]. **2007**; 196(6):812–816. Available

from: https://academic.oup.com/jid/article-lookup/doi/10.1086/520818

8.   McEachern JA, Bingham J, Crameri G, et al. A recombinant subunit vaccine formulation protects

against lethal Nipah virus challenge in cats. Vaccine [Internet]. **2008**; 26(31):3842–3852.

Available from: https://linkinghub.elsevier.com/retrieve/pii/S0264410X08005720

9.   Dups J, Middleton D, Long F, Arkinstall R, Marsh GA, Wang L-F. Subclinical infection without

encephalitis in mice following intranasal exposure to Nipah virus-Malaysia and Nipah virus-

Bangladesh. Virol J [Internet]. **2014**; 11(1):102. Available from:

https://virologyj.biomedcentral.com/articles/10.1186/1743-422X-11-102

10.   Valbuena G, Halliday H, Borisevich V, Goez Y, Rockx B. A Human Lung Xenograft Mouse Model

of Nipah Virus Infection. Pekosz A, editor. PLoS Pathog [Internet]. **2014**; 10(4):e1004063.

Available from: https://dx.plos.org/10.1371/journal.ppat.1004063

11.   Kasloff SB, Leung A, Pickering BS, et al. Pathogenicity of Nipah henipavirus Bangladesh in a

swine host. Sci Rep [Internet]. **2019**; 9(1):5230. Available from:

http://www.nature.com/articles/s41598-019-40476-y

12.   Torres-Velez FJ, Shieh W-J, Rollin PE, et al. Histopathologic and Immunohistochemical

Characterization of Nipah Virus Infection in the Guinea Pig. Vet Pathol [Internet]. **2008**;

45(4):576–585. Available from: http://journals.sagepub.com/doi/10.1354/vp.45-4-576

13.   Clayton BA, Middleton D, Bergfeld J, et al. Transmission Routes for Nipah Virus from Malaysia

and Bangladesh. Emerg Infect Dis [Internet]. **2012**; 18(12):1983–1993. Available from:

http://wwwnc.cdc.gov/eid/article/18/12/12-0875_article.htm

484 14. Satterfield BA, Cross RW, Fenton KA, et al. The immunomodulating V and W proteins of Nipah

485   virus determine disease course. Nat Commun [Internet]. **2015**; 6(1):7483. Available from:

486   http://www.nature.com/articles/ncomms8483

487 15. Bossart KN, Zhu Z, Middleton D, et al. A Neutralizing Human Monoclonal Antibody Protects

488   against Lethal Disease in a New Ferret Model of Acute Nipah Virus Infection. Basler CF, editor.

489   PLoS Pathog [Internet]. **2009**; 5(10):e1000642. Available from:

490   https://dx.plos.org/10.1371/journal.ppat.1000642

491 16. Wong KT, Grosjean I, Brisson C, et al. A Golden Hamster Model for Human Acute Nipah Virus

492   Infection. Am J Pathol [Internet]. **2003**; 163(5):2127–2137. Available from:

493   https://linkinghub.elsevier.com/retrieve/pii/S0002944010635699

494 17. Weingartl H, Czub S, Copps J, et al. Invasion of the Central Nervous System in a Porcine Host

495   by Nipah Virus. J Virol [Internet]. **2005**; 79(12):7528–7534. Available from:

496   https://journals.asm.org/doi/10.1128/JVI.79.12.7528-7534.2005

497 18. Middleton DJ, Westbury HA, Morrissy CJ, et al. Experimental Nipah Virus Infection in Pigs and

498   Cats. J Comp Pathol [Internet]. **2002**; 126(2–3):124–136. Available from:

499   https://linkinghub.elsevier.com/retrieve/pii/S0021997501905328

500 19. Dhondt KP, Mathieu C, Chalons M, et al. Type I Interferon Signaling Protects Mice From Lethal

501   Henipavirus Infection. J Infect Dis [Internet]. **2013**; 207(1):142–151. Available from:

502   https://academic.oup.com/jid/article-lookup/doi/10.1093/infdis/jis653

503 20. Geisbert TW, Daddario-DiCaprio KM, Hickey AC, et al. Development of an Acute and Highly

504   Pathogenic Nonhuman Primate Model of Nipah Virus Infection. Jacobson S, editor. PLoS One

[Internet]. **2010**; 5(5):e10690. Available from: https://dx.plos.org/10.1371/journal.pone.0010690

21.   Johnston SC, Briese T, Bell TM, et al. Detailed Analysis of the African Green Monkey Model of

Nipah Virus Disease. Elankumaran S, editor. PLoS One [Internet]. **2015**; 10(2):e0117817.

Available from: https://dx.plos.org/10.1371/journal.pone.0117817

22.   Prasad AN, Agans KN, Sivasubramani SK, et al. A Lethal Aerosol Exposure Model of Nipah

Virus Strain Bangladesh in African Green Monkeys. J Infect Dis [Internet]. **2020**;

221(Supplement_4):S431–S435. Available from:

https://academic.oup.com/jid/article/221/Supplement_4/S431/5610171

23.   Lo MK, Feldmann F, Gary JM, et al. Remdesivir (GS-5734) protects African green monkeys from

Nipah virus challenge. Sci Transl Med [Internet]. **2019**; 11(494):eaau9242. Available from:

https://stm.sciencemag.org/lookup/doi/10.1126/scitranslmed.aau9242

24.   Mire CE, Geisbert JB, Agans KN, et al. Use of Single-Injection Recombinant Vesicular Stomatitis

Virus Vaccine to Protect Nonhuman Primates Against Lethal Nipah Virus Disease. Emerg Infect

Dis [Internet]. **2019**; 25(6):1144–1152. Available from: http://wwwnc.cdc.gov/eid/article/25/6/18-

1620_article.htm

25.   Prasad AN, Woolsey C, Geisbert JB, et al. Resistance of Cynomolgus Monkeys to Nipah and

Hendra Virus Disease Is Associated With Cell-Mediated and Humoral Immunity. J Infect Dis

[Internet]. **2020**; 221(Supplement_4):S436–S447. Available from:

https://academic.oup.com/jid/article/221/Supplement_4/S436/5727782

26.   Geisbert TW, Mire CE, Geisbert JB, et al. Therapeutic Treatment of Nipah Virus Infection in

Nonhuman Primates with a Neutralizing Human Monoclonal Antibody. Sci Transl Med [Internet].

2014; 6(242):242ra82-242ra82. Available from:

https://stm.sciencemag.org/lookup/doi/10.1126/scitranslmed.3008929

27.   Geisbert JB, Borisevich V, Prasad AN, et al. An Intranasal Exposure Model of Lethal Nipah Virus

Infection in African Green Monkeys. J Infect Dis [Internet]. **2020**; 221(Supplement_4):S414–

S418. Available from: https://academic.oup.com/jid/article/221/Supplement_4/S414/5610161

28.   Liu J, Coffin KM, Johnston SC, et al. Nipah virus persists in the brains of nonhuman primate

survivors. JCI Insight [Internet]. **2019**; 4(14). Available from:

https://insight.jci.org/articles/view/129629

29.   Lee JH, Hammoud DA, Cong Y, et al. The Use of Large-Particle Aerosol Exposure to Nipah

Virus to Mimic Human Neurological Disease Manifestations in the African Green Monkey. J

Infect Dis [Internet]. **2020**; 221(Supplement_4):S419–S430. Available from:

https://academic.oup.com/jid/article/221/Supplement_4/S419/5613084

30.   Lara A, Cong Y, Jahrling PB, et al. Peripheral immune response in the African green monkey

model following Nipah-Malaysia virus exposure by intermediate-size particle aerosol. Geisbert T,

editor. PLoS Negl Trop Dis [Internet]. **2019**; 13(6):e0007454. Available from:

https://dx.plos.org/10.1371/journal.pntd.0007454

31.   Cong Y, Lentz MR, Lara A, et al. Loss in lung volume and changes in the immune response

demonstrate disease progression in African green monkeys infected by small-particle aerosol

and intratracheal exposure to Nipah virus. Horby P, editor. PLoS Negl Trop Dis [Internet]. **2017**;

11(4):e0005532. Available from: https://dx.plos.org/10.1371/journal.pntd.0005532

32.   Marianneau P, Guillaume V, Wong KT, et al. Experimental Infection of Squirrel Monkeys with

Nipah Virus. Emerg Infect Dis [Internet]. **2010**; 16(3):507–510. Available from:

http://wwwnc.cdc.gov/eid/article/16/3/09-1346_article.htm

33. Hammoud DA, Lentz MR, Lara A, et al. Aerosol exposure to intermediate size Nipah virus

particles induces neurological disease in African green monkeys. Bente DA, editor. PLoS Negl

Trop Dis [Internet]. **2018**; 12(11):e0006978. Available from:

https://dx.plos.org/10.1371/journal.pntd.0006978

34. Mire CE, Satterfield BA, Geisbert JB, et al. Pathogenic Differences between Nipah Virus

Bangladesh and Malaysia Strains in Primates: Implications for Antibody Therapy. Sci Rep

[Internet]. **2016**; 6(1):30916. Available from: http://www.nature.com/articles/srep30916

35. Gómez Román R, Wang L-F, Lee B, et al. Nipah@20: Lessons Learned from Another Virus with

Pandemic Potential. Imperiale MJ, editor. mSphere [Internet]. **2020**; 5(4). Available from:

https://journals.asm.org/doi/10.1128/mSphere.00602-20

36. Pelissier R, Iampietro M, Horvat B. Recent advances in the understanding of Nipah virus

immunopathogenesis and anti-viral approaches. F1000Research [Internet]. **2019**; 8:1763.

Available from: https://f1000research.com/articles/8-1763/v1

37. Worley KC, Warren WC, Rogers J, et al. The common marmoset genome provides insight into

primate biology and evolution. Nat Genet [Internet]. **2014**; 46(8):850–857. Available from:

https://doi.org/10.1038/ng.3042

38. Okano H, Hikishima K, Iriki A, Sasaki E. The common marmoset as a novel animal model

system for biomedical and neuroscience research applications. Semin Fetal Neonatal Med

[Internet]. **2012**; 17(6):336–340. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S1744165X12000844

39.   Yun T, Park A, Hill TE, et al. Efficient Reverse Genetics Reveals Genetic Determinants of

Budding and Fusogenic Differences between Nipah and Hendra Viruses and Enables Real-Time

Monitoring of Viral Spread in Small Animal Models of Henipavirus Infection. J Virol [Internet].

American Society for Microbiology; **2021**; 89(2):1242–1253. Available from:

https://doi.org/10.1128/JVI.02583-14

40.   Partek Inc. Partek® Flow® (Version 10.0). 2020.

41.   Howe KL, Achuthan P, Allen J, et al. Ensembl 2021. Nucleic Acids Res [Internet]. **2021**;

49(D1):D884–D891. Available from: https://academic.oup.com/nar/article/49/D1/D884/5952199

42.   Danecek P, Bonfield JK, Liddle J, et al. Twelve years of SAMtools and BCFtools. Gigascience

[Internet]. **2021**; 10(2). Available from:

https://academic.oup.com/gigascience/article/doi/10.1093/gigascience/giab008/6137722

43.   Thomas PD, Kejariwal A, Guo N, et al. Applications for protein sequence-function evolution data:

mRNA/protein expression analysis and coding SNP scoring tools. Nucleic Acids Res [Internet].

2006; 34(Web Server):W645–W650. Available from: https://academic.oup.com/nar/article-

lookup/doi/10.1093/nar/gkl229

44.   Supek F, Bošnjak M, Škunca N, Šmuc T. REVIGO Summarizes and Visualizes Long Lists of

Gene Ontology Terms. Gibas C, editor. PLoS One [Internet]. **2011**; 6(7):e21800. Available from:

https://dx.plos.org/10.1371/journal.pone.0021800

45.   Yoneda M, Xu L, Kajiyama H, et al. Secretoglobin Superfamily Protein SCGB3A2 Alleviates

House Dust Mite-Induced Allergic Airway Inflammation in Mice. Int Arch Allergy Immunol

[Internet]. **2016**; 171(1):36–44. Available from: https://www.karger.com/Article/FullText/450788

46. Munster VJ, Prescott JB, Bushmaker T, et al. Rapid Nipah virus entry into the central nervous

system of hamsters via the olfactory route. Sci Rep [Internet]. **2012**; 2(1):736. Available from:

http://www.nature.com/articles/srep00736

47. Wignall-Fleming EB, Hughes DJ, Vattipally S, et al. Analysis of Paramyxovirus Transcription and

Replication by High-Throughput Sequencing. Dutch RE, editor. J Virol [Internet]. **2019**; 93(17).

Available from: https://journals.asm.org/doi/10.1128/JVI.00571-19

48. Cox RM, Krumm SA, Thakkar VD, Sohn M, Plemper RK. The structurally disordered

paramyxovirus nucleocapsid protein tail domain is a regulator of the mRNA transcription

gradient. Sci Adv [Internet]. **2017**; 3(2). Available from:

https://www.science.org/doi/10.1126/sciadv.1602350

49. Cox R, Plemper RK. The paramyxovirus polymerase complex as a target for next-generation

anti-paramyxovirus therapeutics. Front Microbiol [Internet]. **2015**; 6. Available from:

http://www.frontiersin.org/Virology/10.3389/fmicb.2015.00459/abstract

50. Rao PL, Gandham RK, Subbiah M. Molecular evolution and genetic variations of V and W

proteins derived by RNA editing in Avian Paramyxoviruses. Sci Rep [Internet]. **2020**; 10(1):9532.

Available from: http://www.nature.com/articles/s41598-020-66252-x

51. Lo MK, Harcourt BH, Mungall BA, et al. Determination of the henipavirus phosphoprotein gene

mRNA editing frequencies and detection of the C, V and W proteins of Nipah virus in virus-

infected cells. J Gen Virol [Internet]. **2009**; 90(2):398–404. Available from:

https://www.microbiologyresearch.org/content/journal/jgv/10.1099/vir.0.007294-0

<u>**Figure Legends**</u>

Table 1. Clinical description and diagnosis in NiV$_B$ infected marmosets.

Graphical representation of clinical descriptions and diagnoses of all four marmosets infected with NiV$_B$.

Further detail is available in **Table S1**.

Figure 1: Progressive pathogenesis in marmosets after NiV$_B$ infection.

(A) Subject #300 was positioned in a ventral dorsal position and all images are in a R -> L orientation.

X-ray images of the chest were taken on days 0, 1, 3, 5, 7, and at euthanasia. Prior to day 7, lungs

appeared normal. Progressive lung pathology was detected from day 7 onward, with a decrease in

normal lung tissue over the experimental infection course. By day 7, some decreasing opacity was

observed along with an appearance of an interstitial pattern. At the time of Euthanasia (day 9 post-

infection), virtually no normal lung was observed as the lungs develop a nodular alveolar pattern,

suggesting the alveoli are filled with fluid. **(B)** Lung of subject #425 at time of euthanasia at day 8 post-

infection. Acute diffuse pulmonary hemorrhage, with hemorrhage involving all lobes and <10% of

normal tissue remaining. **(C)** Liver of subject #425. Diffuse massive necrosis of the liver, with necrosis

involving all lobes and normal structure visible.

Figure 2: Histopathological changes in marmoset tissues after NiV$_B$ infection.

Panels A, C, E, G, I: H&E staining. **Panels B, D, F, H, J:** Immunohistochemical staining for NiV

nucleoprotein.

(A) Lung (subject #340): Mild to moderate infiltration of mononuclear cells (macrophages, monocytes)

and neutrophils in perivascular space. **(B)** Lung (subject #425): NiV antigen present in endothelial cells

and smooth muscle cells of pulmonary vessels (arteriole, venule). **(C)** Liver (subject 340): Focal

coagulation necrosis in midzonal hepatocytes, surrounded by inflammatory cells (neutrophils,

mononuclear cells). **(D)** Liver (subject #425): NiV antigen present in hepatocytes, sinusoidal endothelial

cells, and portal vein endothelial cells. **(E)** Spleen (subject #340): Necrosis of red pulp and formation of

syncytial cells. **(F)** Spleen (subject #425): NiV antigen present in mononuclear or multinuclear cells in

peripheral zone of follicles, or in red pulp. **(G)** Kidney (subject #300): Loss of endothelial and mesangial

cells in affected glomeruli. **(H)** Kidney (subject #425): NiV antigen present in glomerular cells and

peritubular capillary endothelial cells. **(I)** Heart (subject #425): Mild myocarditis with necrosis of

myocytes and infiltration of mononuclear cells and neutrophils. **(J)** Heart (subject #425): NiV antigen

present in capillary and arteriole endothelial cells, as well as heart muscle cells.

Figure 3. Viral titers and viral RT-qPCR data across various tissues.

(A) At time of euthanasia, tissues were collected. Viral titers were determined by plaque assay in 23

tissues, combined into 10 tissues systems seen here. Values equal to zero were set to 10^0 for

visualization on a log-scale. Tissue systems with more than one tissue type are upper respiratory tract

(nasal mucosa and trachea), lower respiratory tract (left upper lobe, left lower lobe, right upper lobe,

and right lower lobe of the lung as well as each of the left and right bronchi), kidney and adrenals

(kidney and adrenal gland), gastrointestinal tract (pancreas, jujunum, and colon transversum), central

nervous system (brain-frontal, brain-cerebellum, brain stem, and cervical spinal cord), and genitourinary

system (urinary bladder and gonads). **(B)** RT-qPCR was performed to determine relative amounts of

viral RNA across 23 tissues. Average cycle threshold for uninfected samples is shown (dotted line).

Alternating white and gray fill for box and whisker plots separate similar tissue types (e.g. all lymph

node (LN) samples are adjacent and grey). Both right (R) and left (L) lower and upper lung samples are

also shown adjacent and with the same fill color (grey).

Figure 4. RNA-seq of NiV$_B$ in infected marmoset tissues.

(A) Total number of reads aligned to *Callithrix jacchus* (cyan) and NiV$_B$ (dark blue) are shown across all

tissues and samples. The dotted line indicates 100 aligned reads, the threshold for inclusion into 4B.

(B) The fraction of total NiV$_B$ aligned reads corresponding to each Nipah virus gene across all samples

meeting the 100 read threshold. **(C)** The fraction of reads with the indicated number of inserted

guanines (Gs) at the RNA editing site in P for all samples meeting the threshold of 100 reads aligned to

the RNA editing site. **(D)** The fraction reads at the P editing site resulting in P, V, or W based on the

number of inserted Gs.

Figure 5. RNA-seq samples aligned to *Callithrix jacchus*.

For all subfigures, differential expressed genes must meet the threshold of FDR step up < 0.0 and,

absolute value of fold change > 2. **(A)** Across all animals and all tissues, differential expression analysis

was performed comparing infected and uninfected samples. Filtered differentially expressed genes

were used to obtain enriched gene ontology (GO) gene sets, and similar terms were clustered.

Log_{10}(FDR) of the gene set is indicated by the color and log_{10} of the number of terms in the gene set is

indicated by the size of the circle. **(B)** Differential expression analysis only in the lung samples across

all animals were used to identify the top 25 differentially expressed genes and then hierarchical

clustering was performed to obtain a heatmap. Genes in red are more expressed relative to the other

samples and genes in blue are less expressed. **(C)** Principal component analysis was performed on the

brain stem samples across all animals. The top three principal components are shown. Tissues from

infected animals (blue) and tissues from uninfected animals (red) are further broken down into tissues

from animals showing clinical neurological signs (sphere). This was only seen in marmoset #340. **(D)**

The top differentially expressed genes between #340 and the other animals were used to ascertain

enriched GO terms. The largest interconnected family of GO terms is shown with darker red indicating

a lower FDR and a larger circle indicating more genes in a particular GO term.

 Supplementary Figure Legends

 Table S1. Clinical description and diagnosis in NiV$_B$ infected marmosets.

 Detailed descriptions of clinical and gross pathology as well as clinical diagnoses for all four marmosets

 infected with NiV$_B$.

 Figure S1: Changes in body weight and temperature in marmosets after NiV$_B$ infection.

 (A) Changes in body weight are shown as percent change compared to the individual starting weight at

 day 0 of the study. (B) Changes in body temperature are shown as changes in ºF compared to the first

 measurement made on day 1 post-infection. Subject #193 is shown in orange circles, subject #300 in

 red circles, subject #340 in green circles, and subject #425 in blue circles.

 Figure S2: Progressive lung pathogenesis in marmosets after NiV$_B$ infection.

 Subject were positioned in a left lateral position and all images are in a Cranial to Caudal orientation. X-

 ray images were taken on days 0, 1, 3, 5, 7, and at euthanasia. Prior to day 7, lungs appeared normal.

 Progressive lung pathology was detected from day 7 onward. **(A)** Subject #340: A mild bronchial

 pattern emerged at the time of euthanasia (day 9 post-infection) and showed some inflammation

 around the smaller bronchi, compared to day 7. **(B)** Subject #425: A severe interstitial pattern with

 nodules emerged at the time of euthanasia (day 11 post-infection), compared to day 7 where the

 animal was showing some inflammation around the smaller bronchi with a mild interstitial infiltrate. **(C)**

 Subject #193: A severe interstitial pattern emerged at the time of euthanasia (day 11 post-infection),

 compared to day 7 where the lung was completely normal

 Supplementary Figure S3. Changes in complete blood count.

 Taken at days 1, 3, 5, 7, and 10, hematological analysis included total white blood cell counts, white

 blood cell differential, red blood cell counts, platelet counts, hematocrit values, total hemoglobin

concentrations, mean cell volumes, mean corpuscular volumes and mean corpuscular hemoglobin concentrations. White blood cell count (WBC), neutrophils (Neu), eosinophils (Eos), basophils (Bas), hematocrit (HCT), and platelets (PLT.

Supplementary Figure S4. Changes in blood chemistry.

Clinical chemistry analysis was performed at days 1, 3, 5, 7, and 10. Serum samples were tested for concentrations of albumin, alanine aminotransferase (ALT), alkaline phosphatase (ALP), glucose, plasma electrolytes (blood urea nitrogen (BUN) and creatinine (CRE)).

Supplementary Figure S5. Additional changes in complete blood count.

Taken at days 1, 3, 5, 7, and 10, hematological analysis included total white blood cell counts, white blood cell differential, red blood cell counts, platelet counts, hematocrit values, total hemoglobin concentrations, mean cell volumes, mean corpuscular volumes and mean corpuscular hemoglobin concentrations. Red blood cell count (RBC), lymphocytes (LYM), monocytes (Mon), plateletcrit (PCT), platelet distribution width (PDWs), platelet distribution width % (PDWc), hemoglobin (HGB), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC), red cell distribution width (RDWs), red cell distribution width % (RDWc), and mean platelet volume (MPV).

Supplementary Figure S6. Additional changes in blood chemistry.

Clinical chemistry analysis was performed at days 1, 3, 5, 7, and 10. Serum samples were tested for concentrations of amylase (AMY), total protein (TP), total bilirubin (TBIL), plasma electrolytes (calcium (CA), phosphorus (Phos), sodium (Na+), and potassium (K+), and globulin (GLOB)).

Nipah virus Bangladesh infection elicits organ-specific innate and inflammatory responses in the marmoset model

Christian S. Stevens[1], Jake Lowry[2], Terry Juelich[3], Colm Atkins[3,#], Kendra Johnson[4], Jennifer K Smith[3], Maryline Panis[1], Tetsuro Ikegami[3], Benjamin tenOever[1], Alexander N. Freiberg[3], Benhur Lee[1]

[1]Icahn School of Medicine at Mount Sinai, Department of Microbiology

[2]University of Texas Medical Branch, Animal Resource Center

[3]University of Texas Medical Branch, Department of Pathology

[4]University of Texas Medical Branch, Department of Microbiology and Immunology

[#]Present address: Rutgers University, Department of Cell Biology and Neuroscience

Table 1. Clinical description and diagnosis in NiV$_B$ infected marmosets

	Clincal Diagnosis	193 Male 11 DPI	300 Female 9 DPI	340 Female 11 DPI	425 Male 8 DPI
(1)	Diffuse Pulmonary Edema and Hemorrhage	✅	✅	✅	✅
(2)	Diffuse hemorrhagic hepatitis	✅	✅	✅	✅
(3)	Multi-focal hemorrhagic Lymphadenopathy	✅	✅		✅
(4)	Hemorrhagic Tonsilitis		✅		
(5)	Splenic Congestion				✅
(6)	Left Ventricular Cardiomyopathy / Right Atrial Dilation			✅	
(7)	Cerebral Infarct and malacia in the occipital lobe			✅	
(8)	Cerebral Edema			✅	
	(Left Hindlimb tremors at euthansia)			✅	

Figure 1. X-ray autoradiography and gross pathology.

Day 0

Day 7

Euthanasia Day 9

Figure 2. Histopathological changes in NiV$_B$ infected marmosets.

Figure 3. Viral titers and viral RT-qPCR data.

Figure 4. RNA-seq of NiV$_B$ in infected marmoset tissues.

Figure 5. RNA-seq samples aligned to *Callithrix jacchus*.

Supplementary Figures

Table S1. Clinical description and diagnosis in NiV$_B$ infected marmosets

Subject No.	Sex	Euthanized (DPI)	Clinical and gross pathology	Clinical diagnosis
193	Male	11	(1) Minimal (<10%) normal lung tissue detected in the cranial portion of the lung. The rest of lung was fluid filled with hemorrhagic/serous fluid. The left and right bronchus had clear fluid present and the trachea had minimal fluid in it. (2) Liver had multi-focal to coalescing regions of pale circles of varying sizes throughout all fields and was very friable. (3) Mediastinal and bronchial lymph nodes had gross enlargement and hemorrhage.	(1) Diffuse Pulmonary Edema and Hemorrhage (2) Diffuse hemorrhagic hepatitis (3) Multi-focal hemorrhagic Lymphadenopathy
300	Female	9	(1) Minimal (<5%) normal lung tissue detected in the cranial portion of the lung. The rest of lung was fluid filled with hemorrhagic/serous fluid. The left and right bronchus had serosanguineous fluid present and the trachea had minimal fluid in it. (2) Liver had multi-focal to coalescing regions of pale circles of varying sizes throughout all fields and was very friable. (3) Inguinal and retropharygneal lymph nodes had gross enlargement but minimal hemorrhage associated with them. (4) Tonsils had pin point hemorrhages associated throughout the tissue. The tissue however was not enlarged. (5) Diffuse subcutaneous liquefactive necrosis of fat. (6) Thoracic cavity contained approximately 1mL of serosanguineous fluid. (7) Peritoneal cavity contained approximately 2mL of serosanguineous fluid.	(1) Diffuse Pulmonary Edema and Hemorrhage (2) Diffuse hemorrhagic hepatitis (3) Multi-focal Lymphadenopathy (4) Hemorrhagic Tonsilitis
340	Female	11	(1) Right lung was normal, though the bronchus was fluid filled. Left lung was congested as it was the dependent lung at the time of death and is a post-mortem change. (2) Liver had multi-focal to coalescing regions of pale circles of varying sizes throughout all fields and was very friable. (3) Inguinal lymph node had gross enlargement but minimal hemorrhage associated with it. (4) Significant hypertrophy of the left ventricular (LV) wall, approximately 10:1 left vs. right. Dilation of the right atrium, though milder than the LV hypertrophy. (5) Infarct was found and malacia near the transition zone between grey and white matter in the occipital lobe of the left cerebral hemisphere. Edema was appreciated in the cerebrum. (6) Trachea was filled with serous fluid at necropsy. (7) Multiple pinpoint areas of hemorrhage were appreciated in the right renal pelvis. (8) Peritoneal cavity contained approximately 0.5 mL of serosanguineous fluid.	(1) Diffuse hemorrhagic hepatitis (2) Left Ventricular Cardiomyopathy / Right Atrial Dilation (3) Cerebral Infarct and malacia in the occipital lobe (4) Cerebral Edema
425	Male	8	(1) Minimal (<5%) normal lung tissue detected in the cranial portion of the lung. The rest of the lung was fluid filled with hemorrhagic/serous fluid. The left and right bronchus had clear fluid present and the trachea had minimal fluid in it. (2) Liver had multi-focal to coalescing regions of pale circles of varying sizes throughout all fields and was very friable. (3) Inguinal, mediastinal and retropharygneal lymph nodes had gross enlargement and hemorrhage. (4) Spleen was extremely friable and congested, no normal tissue was observed on cut sections.	(1) Diffuse Pulmonary Edema and Hemorrhage (2) Diffuse hemorrhagic hepatitis (3) Multi-focal hemorrhagic Lymphadenopathy (4) Splenic Congestion

Supplementary Figure S1. Weight and Temperature.

Supplementary Figure S2. X-ray autoradiography.

Supplementary Figure S3. Changes in complete blood count.

Supplementary Figure S4. Changes in blood chemistry.

Supplementary Figure S5. Additional changes in complete blood count.

Supplementary Figure S6. Additional changes in blood chemistry.