1   **RNAi screening of RING/U-box domain ubiquitin ligases identifies**

2   **critical regulators of stem cells and tissue regeneration in planarians**

John M. Allen[1,2], Madison Balagtas[1], Elizabeth Barajas[1], Carolina Cano Macip[1], Sarai

Alvarez Zepeda[1], Ionit Iberkleid[1], Elizabeth M. Duncan[2], and Ricardo M. Zayas[1]

[1]Department of Biology, San Diego State University, 5500 Campanile Dr., San Diego,

CA 92182-4614, USA.

[2]Deparment of Biology, University of Kentucky, 101 T.H. Morgan Building, Lexington

KY, 40506-022, USA.

*Correspondence should be addressed to:

Dr. Ricardo M. Zayas

E-mail: rzayas@sdsu.edu

Short Title: "Planarian RING/U-box domain E3 ligases."

Keywords: Planarian, Stem Cells, Regeneration, E3 ubiquitin ligases, PRC1, Prpf19,

RNF2

Abstract

Regenerative processes depend on the interpretation of signals to coordinate cell

behaviors. The role of ubiquitin-mediated signaling in regeneration is not well

understood. To investigate how ubiquitylation might specifically impact tissue

regeneration, we are studying planarians that are capable of regenerating from nearly

any injury using a population of stem cells. Here we used RNAi to screen RING/U-box

E3 ubiquitin ligases that are highly expressed in planarian stem cells and stem cell

progeny. RNAi screening identified nine genes with functions in regeneration, including

the spliceosomal factor *prpf19* and epigenetic regulator *rnf2*; based on roles in

developmental processes, we further explored these two genes. We found that *prpf19*

was required for survival but not for stem cell maintenance, suggesting a role in

promoting cell differentiation. Because RNF2 is the catalytic subunit of the Polycomb

Repressive Complex 1 (PRC1), we also examined other cofactors of *rnf2* and observed

a striking phenotype of regional tissue misspecification in *cbx* and *phc* RNAi planarians.

To identify genes regulated by PRC1, we performed RNA-seq after knocking down *rnf2*

or *phc* and found that the set of genes differentially expressed were largely non-

overlapping despite being predicted to function in the same complex. Using *in situ*

hybridization, we showed that *rnf2* regulates gene expression levels within a tissue type,

whereas *phc* is necessary for the spatial restriction of gene expression. This work

uncovered roles for RING/U-box E3 ligases in stem cell regulation and regeneration and

identified differential gene targets for PRC1 factors required for maintaining cell-type-

specific gene expression in planarians.

Introduction

A deep understanding of the networks and signaling pathways that direct the

maintenance and differentiation of adult stem cells is essential for regenerative

therapies. The freshwater planarian, *Schmidtea mediterranea*, is an important model for

studying the molecular mechanisms that underpin stem cell-based regeneration[1,2].

These worms maintain a large population of adult stem cells, a subset of which have

been demonstrated to be pluripotent[3,4]. This population of stem cells continuously

renews planarian tissues during homeostasis and is also mobilized in response to injury

to regenerate tissues[5,6]. As such, they offer an amenable model to study stem cell

biology in a whole-organism *in vivo* context.

Extensive work has been performed to understand the molecular basis of

planarian regeneration[7], yet most studies have primarily examined transcriptional

changes[8-12]. Comparatively, fewer studies have focused on proteomic regulation in

planarian stem cells[13,14] or the post-translational regulation of proteins vital for stem cell

function[15]. An essential post-translation regulator of proteins is the addition of the small,

highly conserved polypeptide ubiquitin, which modifies protein function in myriad cellular

contexts, including transcription, cell cycle regulation, translational fidelity, protein

turnover, and degradation[16-19].

Ubiquitin-dependent signaling events have emerged as essential regulators of

stem cell functions, including self-renewal and differentiation[20]. The transfer of free

ubiquitin onto a target substrate typically occurs through a tripartite enzymatic cascade

that terminates with the E3 ubiquitin ligases. The E3 ligases can be grouped into two

major classes, the HECT (Homologous to the E6-AP Carboxyl Terminus) and the more

prevalent RING (Really Interesting New Gene) class. Of the approximately 617 genes

encoding putative E3 ligases identified in the human genome, 309 were predicted to

contain a RING finger (RNF) or the related U-box domain; a further 270 E3 genes that

function in complexes associated with RINGs[21]. The RNFs are defined by a zinc-finger

domain with an evolutionarily conserved arrangement of cysteine and histidine residues

that coordinate two zinc ions and bind an E2-ubiquitin conjugate[22]. The U-box domain

forms a similar structure to the RING domain and can bind conjugated E2 but does not

coordinate zinc[23]. Substrate recognition and binding are achieved by additional domains

within the RNF protein or association with other proteins as part of a multi-protein

complex. Previous work on E3 ligase function in planarians has implicated HECT E3

and Cullin-Ring complex member ligases as essential regulators of regeneration and

stem cells[15,24].

Here we performed functional analysis on a subset of RING and U-box domain-

containing genes that are expressed in the planarian stem cells or progeny. We found

several to be essential for homeostatic maintenance, regeneration, and tissue

patterning. These genes included spliceosomal factor *prpf19* and epigenetic factors *rnf2*

and *bre1*, known to ubiquitylate histones. We found that *prpf19* was required for worm

survival but not for stem cell maintenance, suggesting a role in promoting cell

differentiation. Polycomb Repressive Complex 1 (PRC1) component gene *rnf2* was

required for global monoubiquitylation of histone H2A (H2Aub1) and promoting proper

regeneration. In addition, when we disrupted PRC1genes *phc* and *cbx*, we did not

observe a global reduction in H2Aub1 levels but did observe defects in the organization

of tissue near the base of the planarian pharynx. Taken together, analysis of RING/U-

box E3 ligases identified stem cell biology and regeneration regulators and led to

discovering differential phenotypes and transcriptional targets for PRC1 factors.

Results

Identification of RING and U-Box E3 ubiquitin ligase genes in *S. mediterranea*

The RING and U-Box protein domains have been identified as having a pivotal role in mediating the ubiquitylation of a target substrate. To identify genes in *S. mediterranea* that are predicted to encode a RING/U-box domain, we filtered a reference planarian transcriptome[25] using InterPro domain annotations and generated a list of 393 transcripts. Next, we used the predicted RING and U-box domain-containing gene transcripts to perform BLAST analysis using a curated list of human E3 ubiquitin ligases[21]. We found 376 planarian genes that were predicted to have homology with a human RING/U-box gene (Supplemental Table S1) and 17 planarian transcripts that did not have predicted significant homology to a human RING/U-box gene.

A functional screen reveals genes with roles in planarian stem cell regulation and regeneration

To identify RING/U-box genes that regulate planarian stem cells and regeneration, we assessed the function of 93 genes from our list, including 72 genes most highly expressed in stem cells or stem cell progeny[8] (Supplemental Table S2). RNAi treatments were performed over the course of four weeks while the worms were observed for defects in homeostasis. After 28 days, the worms were amputated to assess the effect of RNAi treatment on regeneration (Figure 1A). We found that RNAi of nine genes produced phenotypes related to stem cell function in homeostasis during regeneration (Table 1). Phenotypes observed during homeostasis included head

regression, epidermal lesions, ventral curling, and lysis (Figure 1B); other genes

displayed abnormalities and delays during regeneration when disrupted (Figure 1C).

During homeostasis, head regression was observed after RNAi-mediated disruption of

prpf19, *march5*, *traf2-like*, *not4*, and *bre1*; lesions were observed after disruption of

march5, *ran*, and *rnf8-like*; and ventral curling was observed after disruption of *prpf19*

and *not4*. The genes *prpf19*, *march5*, and *ran* were essential for worm survival, and

depletion of these genes caused worm lysis. Knockdown of *rnf8-like*, *bre1*, *rnf2*, and

ring1 caused defective regeneration, typically manifested as a delayed appearance of

visible eyespots compared to *control(RNAi)* treatments. We chose to characterize a

subset of these phenotypes further based on their strong expression in stem cells

(Supplemental Figure S1B) or known roles as epigenetic regulators of stem cells during

developmental processes.

Spliceosomal factor *prpf19* is required for worm survival and stem cell function

The U-box gene *prpf19* had enriched expression in planarian stem cells, and our

initial screen revealed that its expression was required for worm survival. Other aspects

of the RNAi phenotype, including head regression and ventral curling, are typically

associated with depletion or loss of stem cells in planarians. These phenotypes are

consistent with an earlier report for *prpf19* as being up-regulated during, and necessary

for, head regeneration in planarians[26]. *prpf19* is a member of NineTeen Complex (NTC),

with a well-described role in regulating mRNA splicing that is conserved in

Saccharomyces cerevisiae. Consistent with a role in an essential cellular process, we

found broad expression of this gene using WISH (Supplemental Figure S1A).

Furthermore, we found that at least a subset of this expression is in the stem cells or

stem cell progeny by performing WISH of irradiated worms (Supplemental Figure S1B).

We confirmed this observation using double fluorescent in situ hybridization (FISH) to

observe co-expression of *prpf19* with stem cell markers *piwi-1* and *h2b*, and stem cell

progeny markers *prog-1* and *agat*-1 (Supplemental Figure S2A-B). As *prpf19* is known

to function as an E3 ubiquitin ligase, we assayed the effect of *prpf19* inhibition on

ubiquitylated proteins by western blotting and found no noticeable difference with

control samples (Supplemental Figure S2C), suggesting that *prpf19(RNAi)* does not

have an appreciable effect on general proteasome function but rather or has only a

minor effect on global ubiquitylation or is targeting specific proteins that are not

resolvable on a total ubiquitin blot.

To investigate if the *prpf19(RNAi)* phenotypes observed were the result of stem

cell depletion, we performed WISH to planarian stem cell marker genes *tgs-1*, *piwi-1,*

and *h2b*, on *prpf19(RNAi)* and control worms. Surprisingly, all marker genes analyzed

showed robust expression, even in worms where the phenotype had significantly

progressed (Figure 2A). Because *prpf19* was found to be expressed in additional cell

types besides stem cells (Supplemental Figure S2B), we also examined the effect of

prpf19 inhibition on epidermal progeny by performing WISH with markers for early and

late epidermal progeny, *prog-1* and *agat-1*, respectively[10,27]. Consistent with the

epidermal lesions observed during the progression of the *prpf19* phenotype, staining for

epidermal lineage markers was reduced in *prpf19(RNAi)* worms. These results suggest

that *prpf19* function is not required for the maintenance and survival of planarian stem

cells but may affect the differentiation of stem cells into progenitors or the maintenance

of progenitor populations.

Inhibition of *prpf19* causes defects in stem cell proliferation and increases

apoptosis

Despite being dispensable for stem cell maintenance, the strong expression of

prpf19 in stem cells and robust phenotypes that resulted from *prpf19* inhibition

suggested a role for *prpf19* in regulating stem cell dynamics. To examine the effect of

prpf19 RNAi on cell proliferation, we stained *control(RNAi)* and *prpf19(RNAi)* treated

worms with anti-phospho-histone H3 (pH3) to mark mitotic cells across several time

points of treatment. We found that at late time points (day 18), when the phenotype is

beginning to manifest, there was a significant decrease in the number of mitotic cells in

prpf19(RNAi) worms (Figure 2B and 2C). This decrease in the number of mitotic cells

was not correlated with a reduction in the expression of stem cell marker genes (Figure

2A, Supplemental Figure S2A), suggesting that *prpf19(RNAi)* treatment blocks or alters

the rate of stem cell differentiation.

To better understand the severe phenotypes observed in *prpf19(RNAi)* worms, including

epidermal lesioning and worm lysis, we assayed the worms for apoptotic cells. Not

surprisingly, we found an increase in apoptotic cells in *prpf19(RNAi)*-treated worms

compared to control worms at the time point before observing phenotypes, and a

marked increase was observed as the *prp19f(RNAi)* phenotype progressed (Figure 2D

and 2E). This result is congruous with reports of *prpf19* having anti-apoptotic effects in

human cell lines[28]. Together with the observed loss of epidermal progenitor markers

(Figure 2A), these data suggest that the phenotypes observed after *prpf19* depletion are

not caused by a loss of stem cells but rather by dysfunction in the proper homeostatic

replacement of differentiated tissues. After *prpf19* disruption, stem cells are unable to

replenish tissues, either through impaired differentiation or reduced proliferation, and

epidermal integrity becomes compromised. In addition, there could also be a role for

prpf19 as an antiapoptotic factor in differentiated tissues, resulting in increased

apoptosis in *prpf19(RNAi)* worms.

NTC components and targets are necessary for tissue renewal and regeneration

NTC is a large protein complex with various cellular roles but has its best-

described role in regulating pre-mRNA splicing. Named after its founding member,

prpf19, the complex is conserved between humans and yeast. In the spliceosome, NTC

functions as an E3 ligase through its PRPF19 subunit (Figure 3A). To examine if the

effects of *prpf19* RNAi were being mediated through disruption of spliceosomal

assembly and function, we knocked down three homologs of core NTC component

members, *cdc5l*, *prlg1*, and *spf27* (Supplemental Table S3). We found that these genes

were also necessary for worm survival and regeneration (Figure 3B and 3C). *cdc5l* and

prlg1 are essential for NTC function in yeast and presented very severe phenotypes of

head regression, ventral curling, and lysis that mirrored the observations for

prpf19(RNAi). *Spf27(RNAi)* worms displayed a milder phenotype than other NTC genes

examined and showed delayed or absent regeneration in 28/37 head fragments and

33/37 trunk fragments; an additional three trunk fragments showed a more severe

phenotype of ventral curling and lysis. Also, we postulated that if the *prpf19* RNAi

phenotype resulted from its ubiquityl ligase activity within the NTC complex, inhibiting

prpf3 or *prpf8* should result in a similar phenotype. Indeed, we found that *prpf3(RNAi)*

and *prpf8(RNAi)* worms exhibited severe phenotypes like *prpf19(RNAi)*, including head

regression, ventral curling, epidermal lesions, and lysis (Figure 3D). WISH analysis of

NTC genes, *prpf3* and *prpf8*, demonstrated broad parenchymal expression patterns like

prpf19, with *prpf8* showing a noticeable stem cell expression pattern (Supplemental

Figure S3A). The similar phenotypes and expression patterns observed for NTC genes

and factors downstream to *prpf19* suggested that the *prpf19(RNAi)* phenotype is

mediated through its role in NTC and that the NTC and spliceosome function is critical

for stem cell regulation during homeostasis and regeneration.

Histone-modifying ubiquitin ligases are essential for regeneration and

homeostasis

Ubiquitylation of histone H2B is associated with transcriptional activation and is

mediated by the E3 ligase complex RNF20/40 (Bre1 in yeast)[29,30]. We found that

planarians have a single homolog for this complex named *Smed-bre1* (referred to as

bre1 hereon), and that knockdown of *bre1* caused the worms to exhibit head regression

and lesions prior to day 28 of treatment in 33/53 worms assayed (Figure 1B).

Furthermore, most *bre1(RNAi)* worms failed to regenerate when amputated, and many

lysed with 31/53 head fragments and 21/53 trunk fragments lysing by the end of the

observation period (day 14 post-amputation). To investigate if *bre1(RNAi)* affected

global levels of monoubiquityl-histone H2B (H2Bu1), we performed a protein blot using

an H2Bub1-specific antibody. We found reduced levels of H2Bu1 in whole worm

homogenates as soon as 14 days after beginning RNAi treatment (Supplemental Figure

S4A).

In contrast to histone H2B ubiquitylation, monoubiquitylation of histone H2A is

associated with transcriptional repression. It occurs in various cellular contexts,

including developmental processes, stem cell regulation, and the DNA damage

response. Histone H2A is targeted for ubiquitylation by RING1 and RNF2, which act as

RING E3 ligases within PRC1. PRC1 is a repressive epigenetic complex active during

development to monoubiquitylate histone H2A[31] and stably silence genes[32] (Figure 4A).

We identified two candidate homologs of RING1 and RNF2 and found that depletion of

each caused delayed or absent regeneration compared to controls. These phenotypes

were most evident in the trunk fragments where 37/58 *rnf2(RNAi)* and 19/29

ring1(RNAi) worms exhibited a delayed regeneration phenotype (measured by the

appearance of dark eyespots) compared to 7/54 and 2/30 *control(RNAi)* worms assayed

at the same regeneration time point (7 days post-amputation). Of the 37/58 *rnf2(RNAi)*

trunks and 19/29 *ring1(RNAi)* trunks with regeneration defects, 13/37 and 4/19 failed to

form regeneration blastemas, respectively, whereas all *control(RNAi)* worms formed

normal-sized blastemas (Figure 1C). No obvious phenotypes were observed during

homeostatic maintenance, even during long-term RNAi treatment (>16 feeds over eight

weeks). To assess if the ubiquityl ligase activity of *rnf2* towards histone H2A was

conserved in planarians, we examined bulk levels of H2Aub1 by protein blot analysis

using an H2Aub1 specific antibody and observed markedly reduced levels of H2Aub1

after *rnf2* inhibition (Figure 4B). In contrast, we found that *ring1* RNAi did not

appreciably affect global H2Aub1 levels (Supplemental Figure S4B), consistent with *rnf2*

being the E3 ligase subunit within PRC1 that is responsible for H2A ubiquitylation[31,33].

Disruption of canonical PRC1 subunit *phc* affects the patterning of the planarian

pharyngeal body region

To gain insight into the role of PRC1 in planarians, we examined the functional

requirements for PRC1 subunits. In vertebrates, the composition of PRC1 is variable[34];

the complex is defined by which of the six mammalian paralogs of PGCF is present.

PCGF2 and PCGF4 define the mammalian canonical PRC1 complex (cPRC1), which

also includes one each of several chromobox (CBX) and Polyhomeotic (PHC) paralogs

(Figure 4A). We identified planarian homologs for these other core PRC1 genes and

found a single homolog for *cbx* and *phc* and two homologs for *pcgf* (Supplemental Table

S3). To investigate if the phenotypes for *rnf2(RNAi)* and *ring1(RNAi)* were mediated

through their function in canonical PRC1, we used RNAi to deplete *cbx*, *phc*, *pcgf2*, and

pcgf3.

In contrast to the impaired regeneration observed after *rnf2* or *ring1* knockdown,

RNAi for *phc* or *cbx* demonstrated a complex homeostasis phenotype that included the

abnormal appearance of a dorsal lesion anterior to the pharynx (Figure 4C). In some

cases, we observed the pharynx protruding from the lesioned region and extending

ectopically from the dorsal surface of the worm. As the phenotypes progressed, these

RNAi worms began to exhibit defects along the body axis, showing crimped tails unable

to affix to the dish and epidermal lesions. We also assayed the effect of inhibition of the

canonical PRC1 genes on H2Aub1 levels. We found that inhibition of *phc* or *cbx* did not

impact bulk H2Aub1 levels (Supplemental Figure S4B), suggesting that cPRC1 is not

responsible for most H2Aub1 deposition, like findings in vertebrate models[35]. Both *phc*

and *cbx* had similar mRNA expression patterns, suggesting that they can function in the

same complex (Supplemental Figure S4C). This expression pattern overlapped with the

diffuse parenchymal expression pattern for *rnf2* and *ring1* (Supplemental Figure S1A)

but had more robust expression near the planarian brain and intestinal branches, the

latter of which are areas known to be enriched in stem cells.

While similar, the penetrance of the *phc(RNAi)* phenotype was higher than for

cbx(RNAi), and we chose to examine the *phc(RNAi)* phenotype further using known

markers of tissue patterning. The appearance of a dorsal lesion and mislocalization of

the pharynx to the dorsal surface in *phc(RNAi)* animals suggests that planarian PRC1

may be involved in the specification of specific tissues related to the pharynx or in

regulating genes that provide axial positioning cues to stem cell progeny during

homeostatic tissue turnover. We examined dorsal-ventral patterning factor *bmp-4*[36] and

anterior-posterior factor *ndl-3*[37] expression after *phc(RNAi)* and did not observe a

noticeable change in the expression pattern of these factors relative to controls

(Supplemental Figure S4D). We then further examined genes that mark specific tissues

related to the pharynx, including the pharynx marker *laminin*[38] and the gene *NB.22.1E*[39],

which labels marginal adhesive gland cells, the ventral mouth opening, and a population

of cells near the base of the pharynx. Following *phc* inhibition, we observed that *laminin*

expression was reduced to a single condensed spot of expression near the location of

the dorsal lesion and a few scattered cells near the midline (Figure 4D). Likewise, we

observed the specific disappearance of the *NB.22.1E⁺* population of cells at the base of

the pharynx following *phc(RNAi)*. In contrast, expression along the body margin and

ventral mouth opening was unaffected (Figure 4D). These data establish a role for

PRC1 factors in maintaining tissue identity in a non-embryological context.

RNA-seq analysis of gene expression after PRC1 subunit inhibition reveals

candidate transcriptional targets of PRC1

To investigate which genes are differentially expressed after PRC1 inhibition and

to understand the transcriptional basis for the *rnf2* and *phc* RNAi phenotypes, we

performed RNA-seq. We chose time points based on the phenotypic progression,

quantitative PCR analysis to confirm a robust reduction in target RNAi transcript levels

(not shown), and, for *rnf2(RNAi)*, protein blot analysis to ensure the RNAi treatment was

reducing levels of H2Aub1. Based on these parameters, we extracted RNA after 11

days of *phc* RNAi treatment and 14 and 28 days after *rnf2* RNAi (Supplemental Figure

S5A). For *rnf2(RNAi)*, we identified 264 differentially expressed genes (126 down-

regulated and 138 up-regulated) (Figure 5A, Supplemental Figure S5B). Not

surprisingly, an extended *rnf2* RNAi treatment increased the number of differentially

expressed genes: 247 genes were differentially expressed after 28 days compared to

29 after 14 days of treatment (Table 2). Also, there was substantial overlap between the

rnf2(RNAi) data sets, with 12 of 29 genes in the day 14 data set represented in the day

28 data set (Supplemental Figure S5C). After 11 days of *phc(RNAi)*, 49 genes were

differentially expressed: 20 were down-regulated and 29 up-regulated (Figure 5C).

Consistent with a repressive role in transcriptional regulation, more genes were

significantly up-regulated when either *phc* or *rnf2* was inhibited. Importantly, *rnf2* and

phc were each significantly down-regulated when targeted for RNAi.

Surprisingly, despite being predicted to function in a complex together, only a

single gene was found in common between the *phc(RNAi)* and *rnf2(RNAi)* data sets.

Intriguingly, this gene was *cbx*, which encodes a chromatin-binding element within

PRC1, and was the most significantly up-regulated gene after *phc* knockdown. This

overall lack of overlap between the data sets suggests that *phc* and *rnf2* regulate

different processes and pathways *in vivo,* and this difference explains the disparate

phenotypes observed after RNAi treatment.

To gain insight into the RNA-seq data set, we performed Gene Ontology (GO)

analysis on the differentially expressed gene set from the day 28 *rnf2(RNAi)* gene set.

The down-regulated genes were significantly enriched for GO biological process terms

related to metabolic and catabolic processes (Supplemental Figure S5D). Conversely,

among GO terms enriched in up-regulated genes were cellular stress (Supplemental

Figure S5E), especially low oxygen conditions, including "response to hypoxia"

(GO:0001666), "cellular response to decreased oxygen levels" (GO:0036294), "ATF6-

mediated unfolded protein response" (GO:0036500), "regulation of transcription from

RNA polymerase II promoter in response to stress" (GO:0043618), "chaperone

cofactor-dependent protein refolding" (GO:0051085), "protein folding in endoplasmic

reticulum" (GO:0034975), and "protein refolding" (GO:0042026). These GO annotations

suggest that *rnf2* activity represses cellular responses to stress during normal

homeostatic conditions and that epigenetic mechanisms facilitate the switch between

homeostasis and cellular stress responses.

To investigate the spatial expression changes of differentially expressed genes

from our RNA-seq data sets, we selected a subset to examine using WISH after *phc* or

rnf2 RNAi (Supplemental Table S4). For *rnf2(RNAi)*, we selected 33 differentially

expressed genes that were predicted to be involved in the extracellular matrix, stress

response factors, cell signaling, and chromatin regulation or transcription and assayed

their expression after *rnf2* depletion. In general, *rnf2(RNAi)* caused a subtle effect on

gene expression levels. However, in some instances, a robust change in expression

occurred in *rnf2(RNAi)* worms, as seen clearly for *Smed-colec10* and *Smed-colec11*;

expression of these genes is nearly undetectable in control worms as compared to

rnf2(RNAi) worms (Figure 5B). Taken together, the GO and *in situ* analyses indicate

that *rnf2* functions in broad cellular processes and that it maintains gene expression in

differentiated tissues at appropriate levels.

In contrast to the mild effect on gene expression observed in *rnf2(RNAi)* animals,

assaying mRNA expression of putative PHC target genes revealed striking changes in

expression levels and spatial patterning in *phc(RNAi)* worms. We examined 11 genes

using *in situ* hybridization, including genes involved in cell adhesion, cell signaling,

transcription, and chromatin regulation. For 7 of these 11 genes, strong ectopic

expression was observed in *phc(RNAi)* worms in the region of the worm where the

dorsal lesion forms (Figure 5D). Genes ectopically expressed in this region included the

cell adhesion factor *icam5*, the Cut homeobox transcription factor *onecut1*, and *roar*,

which encodes an orphan nuclear receptor. We also found several chromatin regulators

that were misexpressed in the region near the pharynx, including *cbx*, *pc-like*, *smc4*,

and *kat6a*. Additionally, we found that the extracellular matrix protein *egflam*, which is

normally expressed in the nervous system and pharynx tip, was significantly down-

regulated throughout the worm.

The ectopic expression of specific factors and disruption of *NB.22.1E* and

laminin expression at the site of tissue defects in *phc(RNAi)* worms indicates that *phc*

function is required to maintain the proper specification and integrity of tissues in this

region.

Discussion

To address the role of ubiquitin signaling in stem cell regulation and regeneration

in a whole organism *in vivo* context, we performed a functional screen of the RING/U-

box class of E3 ubiquitin ligases that are expressed in stem cells and progeny in *S.*

mediterranea. The screen identified nine genes that demonstrated phenotypes related

to stem cell function or regeneration, building on previous studies from our lab on the

HECT[24] and Cullin[15] classes of E3 ligases.

A U-box gene that we chose to investigate in further detail was *prpf19*, the

founding member of the large protein complex NTC. First characterized in yeast, the

best described role for NTC is in the spliceosome, where the E3 ligase function of

Prpf19 is essential for forming snRNP conformations[40]. We found that depletion of

prpf19 caused a strong homeostasis phenotype that included head regression,

lesioning, ventral curling, and lysis, all of which are morphological effects often caused

by stem cell depletion. We depleted other NTC member genes in this study and

observed similar phenotypes to *prpf19(RNAi)*, suggesting that the *prpf19(RNAi)*

phenotype is mediated through its role in NTC. In addition, we found that the stem cell

population was maintained in *prpf19(RNAi)* worms, suggesting an alternative

mechanism of dysregulation. Our results are consistent with a previous study that

observed a similar phenotype upon depletion of *prp19,* which showed an effect on head

regeneration without disrupting the stem cells[26].

Post-transcriptional processing of RNAs is emerging as a major regulator of

planarian stem cells and differentiation. The PIWI homolog *smedwi-2* was identified as

nonessential for stem cell maintenance but necessary for proper differentiation[41], and

smedwi-3 was shown to regulate stem cell mRNAs through two distinct activities[42]. A

screen of planarian ribonucleoprotein granule component homolog genes demonstrated

that most were expressed in planarian stem cells and that depletion of several genes,

including mRNA turnover factors, exoribonucleases, and DEAD-box RNA helicases,

inhibited regeneration without affecting proliferation or stem cell maintenance[43].

Similarly, the CCR4-NOT complex regulates the post-translational degradation of

mRNAs and has been shown to have a critical role in planarian stem cell biology[44]. The

phenotype of CCR4-NOT complex member gene *Smed-not1* was reported to have a

similar phenotype to *prpf19*, where the animals maintained proliferative stem cells

despite presenting a phenotypic that suggests loss of tissue renewal[44]. This study found

that an additional CCR4-NOT subunit, *not4*, is critical for worm homeostasis and causes

head regression and ventral curling upon inhibition (Figure 1B). This phenotype is

consistent with that of *not1(RNAi)*; however, in the future, it will be necessary to

examine the stem cell population using marker genes in *not4(RNAi)* worms to resolve if

the phenotype is mediated through a similar mechanism. Regulation of mRNAs in

planarian stem cells by several pathways, including piRNAs, deadenylation, or splicing,

is crucial for homeostasis and regeneration while being dispensable for stem cell

maintenance. These studies implicate post-transcriptional regulation of mRNAs in

planarian stem cells as a critical process for regulating differentiation.

Epigenetic regulation of gene expression is essential during development and

throughout organismal life. Our RNAi screen uncovered planarian homologs of histone-

targeting RING E3 ubiquitin ligases that affected worm homeostasis and regeneration

and confirmed that inhibition of *bre1* and *rnf2* reduced levels of monoubiquityl-histone

H2B and H2A, respectively. This work demonstrates that activating and repressive

signals provided through histone modifiers are essential for the proper specification of

stem cells and establishing cellular identity during regenerative events.

PRC1 is a major repressive complex that works during development to

ubiquitylate histone H2A, compact chromatin, and silence target gene expression[31,45-49].

PRC1 function was first discovered and was best characterized as a repressor of the

HOX genes during development[50]. The core PRC1 complex is defined by a RING and

PCGF protein that forms canonical (cPRC1) or variant PRC1 (vPRC1) depending on the

presence of additional factors[51]. The RING subunit acts as an E3 ligase that targets

histone H2A, and in vertebrates is either RING1 or RNF2[52]. In contrast to *Drosophila*,

we found that planarians have two homologs of RING1 and RNF2. While these are

likely to be lineage-specific paralogs instead of direct homologs of each vertebrate

gene, we find that, as in vertebrates, the *S. mediterranea rnf2* paralog acts as the major

ligase and is responsible for the bulk of histone H2A ubiquitylation. We did not observe

a noticeable difference in global H2Aub1 levels after *Smed-ring1* inhibition

(Supplemental Figure S4B). However, as both genes demonstrated similar

regeneration-specific phenotypes, they may share common targets or pathways.

In contrast to *rnf2(RNAi)*, when we perturbed other cPRC1 core elements *phc*

and *cbx*, we did not see a reduction in bulk H2Aub1 levels by western blotting. Work in

mammalian cell lines has determined that vPRC1 activity is responsible for most H2A

ubiquitylation with a minimal contribution from cPRC1 complexes[35,53]. Invertebrates

were not thought to contain vPRC1, but more recent phylogenic analysis that included a

greater variety of invertebrate model organisms indicates that vPRC1 likely evolved as

early as cnidarians[54]. Our protein blot results show a minimal contribution of cPRC1

genes *cbx* and *phc* to overall H2Aub1 levels *in vivo,* and the presence of two *S.*

mediterranea pcgf genes support the existence of vPRC1 in *S. mediterranea*.

To gain insight into which genes are regulated by *rnf2* and *phc,* we performed

RNA-seq following RNAi. Consistent with predicted roles in transcriptional repression,

inhibiting the function of either gene led to more up-regulated than down-regulated

differentially expressed genes. The only gene shared between the data sets was the

significantly up-regulated factor *cbx*. Thus, it is possible that cPRC1 autoregulates its

expression in planarians and that the disruption to chromatin homeostasis induced by

cPRC1 disruption causes a compensatory response involving other chromatin factors.

Using GO analysis, we found that *rnf2* regulates genes related to the cellular

stress response, and when we examined the expression of candidate genes from our

RNA-seq data set using WISH, we observed that gene expression changes in

rnf2(RNAi) animals occurred mainly within the endogenous expression pattern. These

data support a role for RNF2 and potentially H2A ubiquitylation, in tuning transcription

levels within a particular cell type, especially for pathways that are adaptive and

responsive to stressful stimuli.

In contrast to *rnf2(RNAi)*, we saw dramatic shifts in the spatial expression of

specific genes after *phc(RNAi)*, including several genes that showed ectopic expression

near the base of the pharynx where lesions formed in *phc(RNAi)* planarians. In this

affected region, we observed both up- (*intercellular adhesion molecule 5*) and down-

regulation (*pikachurin*) of genes that encode extracellular matrix and intercellular

adhesion molecules, suggesting that their dysregulation is likely linked to the formation

of the dorsal lesion seen after *phc* RNAi. The ectopically expressed genes also included

regulators of cellular specification, including nuclear receptors, transcription factors, and

chromatin modifiers. One gene we identified as being misexpressed after *phc* depletion

was that encoding the nuclear factor *onecut1*, a CUT, and homeobox domain-containing

transcription factor that promotes hepatocyte proliferation remodels chromatin

accessibility and promotes tumor growth in colorectal cancers[55-57]. Based on its role in

regulating transcription and tissue identity in other animal models, we suspect it may be

contributing to the change in patterning near the pharynx. Future investigation into this

factor will elucidate if *onecut1* misexpression drives regional tissue misspecification and

if inhibition of *onecut1* suppresses the *phc(RNAi)* phenotype.

PRC1 regulates gene expression through the post-translational ubiquitylation of

histone H2A and by compacting chromatin. A deeper understanding of the role of this

complex in regulating regeneration will require uncovering which genomic elements it

targets. In the future, assays like ChIP-seq[58] or Cut&Run[59] could be used to identify the

localization of H2Aub1 in the planarian genome to inform how genes are regulated

epigenetically to promote a robust regenerative response during injury and tissue re-

specification and remodeling.

Methods

Planarian care

484 A clonal line of asexual *S. mediterranea* (CIW4) was used in all experiments and kept in

485 1X Montjuïc salts (1.6 mM NaCl, 1.0 mM $CaCl_2$, 1.0 mM $MgSO_4$, 0.1 mM $MgCl_2$, 0.1 mM

486 KCl, 1.2 mM $NaHCO_3$, pH 7.0) in food-grade plastic containers at 20°C[60]. Animals

487 selected for experiments were 3-6 mm in length and starved for one week before

488 experimentation.

Gene identification and Cloning

To find RING and U-box domain-containing genes in *S. mediterranea,* we filtered the

Dresden transcriptome[25,61] using InterPro Domain IDs[62], IPR001841 (Zinc finger, RING-

type), and IPR003613 (U box domain). This list was filtered to include only the longest

gene contig for each hit and was used as query sequences for a BLAST search to a

curated list of human RING and U-box genes[21] at an expected value cut-off of 10^{-3}. We

additionally filtered the Dresden transcriptome for contigs annotated with IPR013083

(Zinc finger, RING/FYVE/PHD-type). This list was filtered to remove duplicate entries,

and a BLAST search was performed against our list of human RING and U-box genes.

As the IPR013083 family contains non-RING and U-box genes, only genes that had

predicted homology to a human gene at an expected cut-off of 10^{-3} were appended to

our initial list. The sequences of interest were obtained from either an EST library[63] or

cloned using gene-specific primers into pPR-T4P using ligation independent cloning[64,65].

EST clone accession numbers and the primer sequences used are listed in

Supplemental Tables S2-S4.

RNA interference

During the initial screening, animals were fed double-stranded RNA (dsRNA) mixed with

a ≈3:1 mixture of liver-water paste twice per week for eight feeds and were amputated

pre-pharyngeally on day 28 of treatment to observe regeneration. *In vitro* transcribed

and dsRNA expressed in bacteria were used to perform RNAi during the initial

screening of RING and U-box genes; all subsequent RNAi knockdowns were performed

using bacterially-expressed dsRNA. *In vitro* dsRNA was synthesized as previously

described[66]; the entire reaction mixture was separated into eight aliquots, mixed with

liver paste, and stored until feeding. Bacterially expressed dsRNA was prepared by

growing cultures of *E. coli* strain HT115 transformed with the pPR-T4P plasmid[64,65]

containing the gene of interest and inducing dsRNA expression using IPTG. Bacteria

pellets were purified using centrifugation and mixed with liver paste for administration to

animals[67].

In situ hybridization

Antisense probes for *in situ* hybridization were synthesized as previously described[68]

from DNA templates amplified from pBS II SK(+) (Stratagene) or pPR-T4P[64,65] plasmid

vectors incorporating either digoxigenin or FITC labeled UTPs. Animals for whole-mount

in situ hybridization (WISH) were processed and hybridized as outlined previously[69].

Briefly, samples were sacrificed in 5% *n*-acetyl cysteine, fixed in 4% formaldehyde, and

bleached in a formamide/hydrogen peroxide bleaching solution. Samples were pre-

hybridized for two hours and then hybridized with probe overnight at 56°C. Next,

samples were incubated with an appropriate antibody, depending on the probe label

and subsequent development strategy. For chromogenic development, samples were

incubated with an anti-digoxigenin-AP antibody (Roche, 1:2000) and developed with

NBT/BCIP in AP buffer. Fluorescent *in situ* development was performed using Fast Blue

salts[70] or Tyramide Signal Amplification (TSA) after incubation with anti-digoxigenin-AP

or anti-FITC-POD (Roche, 1:300) antibodies, respectively, following previously

described protocols[69,71]. For irradiation experiments to eliminate dividing cells, worms

were exposed to 60 Gy of X-ray irradiation in a Precision CellRad Irradiation System

and processed for WISH 7 days post-irradiation.

Anti-phosphohistone H3 immunohistochemistry

Animals were incubated in ice-cold 2% hydrochloric acid for 5 minutes and fixed for 2

hours in Carnoy's solution at 4°C. Samples were washed in methanol for 1 hour at 4°C

and bleached overnight in 6% H_2O_2 diluted in methanol at room temperature. Animals

were washed out of methanol and into PBS-Tx and blocked in 1% Bovine serum

albumin (BSA) diluted in PBS-Tx for 4 hours at room temperature. Samples were

incubated with anti-phosphohistone H3 (Ser 10) (Cell Signaling 3377, 1:1000) diluted in

1% BSA/PBS-Tx overnight at 4°C. Antibody was developed using TSA as previously

described[69].

TUNEL staining

The Terminal Deoxynucleotidyl Transferase-mediated deoxyuridine triphosphate Nick

End-labeling (TUNEL) assay was performed to quantify apoptotic cells. Animals were

incubated in 5% *n*-acetyl cysteine (diluted in PBS) for 5 minutes and fixed in 4%

formaldehyde (diluted in PBS-Tx [PBS with 0.3% Triton X-100]) for 15 minutes.

Samples were then permeabilized in 1% SDS (diluted in PBS) and bleached overnight

in 6% H_2O_2 (diluted in PBS-Tx). Samples were then rinsed and stained using the

Apoptag Kit (Millipore-Sigma) as previously described[72].

Protein extraction and western blotting

Protein samples from RNAi worms were homogenized in Trizol and extracted from the

organic phase following the manufacturer-provided protocol with a modified

solubilization buffer (4M Urea, 0.5% SDS) and added sonication step of 10 one-second

pulses that were performed to increase protein recovery[73,74]. Samples were loaded onto

AnyKD TGX gels (BioRad), transferred using the semidry method to a 0.45m PVDF

membrane, and blocked in 5% nonfat milk/TBS-Tw (Tris-buffered saline with 0.1%

Tween-20). Antibodies to monoubiquityl-Histone H2A (Cell Signaling 8240) and

monoubiquityl-Histone H2B (Cell Signaling 5546) were diluted in 5% bovine serum

albumin in TBS-Tw at 1:2000 and 1:1000 respectively and incubated overnight at 4°C.

Washes were performed with TBS-Tw and anti-rabbit-HRP (Cell Signaling 7074) was

diluted in 5% nonfat milk/TBS-Tw at 1:2500 and incubated for 1 hour at room

temperature. Signal was developed using BioRad Clarity Western ECL Substrate

(BioRad 1705061).

RNA sequencing

Worms from three independent control and experimental RNAi groups per time

point were homogenized in Trizol, and RNA was extracted and purified following

manufacturer protocol. RNA was treated with the Turbo DNA-free kit and column

purified using the Qiagen RNeasy MinElute Cleanup kit. Three independent biological

groups were collected at each time point assayed for both control and experimental

(*rnf2* or *phc*) RNAi treatments. Samples were sequenced on an Illumina HiSeq 4000 to

a read depth of at least 15 million 150 bp paired-end reads. The sequenced reads were

submitted to the NCBI BioProject PRJNA768725. Reads were pseudoaligned to the

Dresden (dd_Smed_v6) transcriptome using kallisto[75], and differential gene expression

analysis was performed using the R Bioconductor package[76] and DESeq2[77] with an

FDR cut-off value of ≤ 0.1 applied. To perform Gene Ontology (GO) analysis,

differentially expressed transcripts from the day 28 *rnf2(RNAi)* data set were compared

to the human proteome using BLASTX (cut-off e-value < $1e^{-3}$). Human UniProt IDs were

used as input for annotation and overrepresentation analysis (http://geneontology.org/)

using Fisher's Exact test with an FDR multiple comparisons correction cut-off of ≤ 0.05

applied.

Acknowledgments

We thank Celeste Romero for help with RNAi experiments, Dr. Kelly Ross for

helpful discussions and feedback on the manuscript, and Catherine May from Boston

Children's Hospital for helping to set up the RNA-seq analysis and providing template R

scripts. J.M.A was supported by the ARCS Fellowship from the ARCS Foundation. This

work was supported by NIH R35GM142679 to E.M.D. and NSF IOS-1350302 and NIH

R01GM135657 to R.M.Z.

598   Author Contributions

J.M.A. and R.M.Z. conceptualized the study and designed experiments. J.M.A., M.B.,

E.B., C.C.M., S.A.Z., and I.I. conducted experiments. J.M.A. and R.M.Z. wrote the

original draft of the manuscript. The manuscript was reviewed and edited by J.M.A.,

E.M.D., and R.M.Z. Funding for the project was acquired by R.M.Z.

Figure Legends

Figure 1. RNAi screen of RING/U-Box E3 ubiquitin ligases identifies regulators of stem

cells and regeneration. (A) Feeding and amputation schedule of RNAi screen. Worms

were fed twice per week for a total of eight feeds and amputated pre-pharyngeally on

day 28. (B) Knockdown of the indicated genes displayed phenotypes including ventral

curling and lesions (white arrow). Animals are shown after the conclusion of the RNAi

feedings and before amputation. (C) Knockdown of indicated genes that demonstrated

phenotypes of delayed or absent regeneration after amputation shown by smaller than

normal or absent blastemas (white arrow) and missing or faint eyespots (white

arrowhead) when compared to *control(RNAi)* worm at the same regeneration time point.

Scale bars = 200 μm.

Figure 2. Inhibition of *prpf19* disrupts stem cell function but is not required for stem cell

maintenance. (A) WISH to stem cell markers *tgs-1*, *piwi-1* and *h2b*, and early and late

epidermal progeny markers *prog-1* and *agat-1*, respectively, in *control(RNAi)* (upper

panels) and *prpf19(RNAi)* animals at 14 (middle panels) and 18 (bottom panels) days

after first RNAi feeding. (B) Representative image of animals fixed 14 days after first

RNAi feeding for *control(RNAi)* (left) or *prpf19(RNAi)* (right) and immunostained for

mitotic marker phospho-histone H3. (C) Quantification of phospho-histone H3$^+$ cells per

mm^2 of worms fixed at 11, 14, and 18 days after first RNAi feed (N = 17 – 24 per time

point). (D) Representative image of animals fixed 14 days after first RNAi feeding for

control(RNAi) (left) or *prpf19(RNAi)* (right) and processed for TUNEL staining. (E)

Quantification of TUNEL$^+$ cells per mm^2 of worms fixed at 11, 14, and 18 days after first

RNAi feed (N = 13 – 16 per time point) All data are represented as mean ± SD. *p-value

< 0.05, **p-value < 0.001, ***p-value < 0.0001,Student's t-test with Holm-Sidak

correction for multiple comparisons. Scale bars = 200 μm.

Figure 3. *prpf19*-associated factors and downstream targets recapitulate *prpf19(RNAi)*

phenotypes. (A) Prpf19 acts as an E3 ligase in NTC, interacting with core complex

members PLRG1, CDC5l, and SPF27 to modify U4/U6 snRNP subunit PRPF3 with

nonproteolytic K63-linked ubiquitin chains. This ubiquityl mark stabilizes the interaction

of PRPF3 with U5 snRNP subunit PRPF8 to allow the stable formation of the U4/U6.U5

tri-snRNP and the catalytic activity of the spliceosome. (B) Knockdown of indicated NTC

core components displaying head regression. (C) Knockdown of NTC core component

spf27 caused a reduced and delayed regenerative response in amputated worms. At 6

days post-amputation, *spf27(RNAi)* worms have smaller blastemas compared to

control(RNAi) worms at the same time point. At day 11 post-amputation, the

regenerative response in *control(RNAi)* worms is largely concluded with large blastemas

and visible reformed eyespots present in trunk fragments. In comparison, *spf27(RNAi)*

worms have smaller blastemas and have not regenerated eyespots. (D) Inhibition of

Prpf19 target *prpf3* and ubiquityl-Prpf3 binding factor *prpf8* demonstrate phenotypes like

prpf19(RNAi) and includes head regression, lesions, and ventral curling. Scale bars =

200 μm.

Figure 4. RNAi-mediated inhibition of canonical PRC1 function disrupts pharyngeal

patterning and histone ubiquitylation. (A) Composition and function of PRC1. PRC1

functions to ubiquitylate histone H2A and compact chromatin to repress gene

expression. (B) Western blot analysis showing a reduction in H2Aub1 levels following

rnf2 inhibition across three biological replicates and two experimental time points. (C)

RNAi of PRC1 genes *phc* and *cbx* causes phenotypes of a dorsal lesion anterior to the

pharynx and mislocalization of the pharynx on the dorsal surface of the worm. (D) WISH

to *NB.22.1E* marks marginal adhesive gland cells, mouth opening, and a population of

cells at the base of the pharynx and for *laminin,* which marks the pharynx feeding organ

in *control(RNAi)* animals (upper panels) and *phc(RNAi)* animals at days 21 (middle

panels) and 28 (bottom panels) after the first RNAi treatment. Scale bars = 200 μm.

Figure 5. Loss of PRC1 function causes changes to gene expression levels and spatial

patterning. (A) Volcano plot of differentially expressed genes after 28 days of *rnf2* RNAi

treatment. (B) WISH analysis of select genes that were differentially expressed by RNA-

seq of *rnf2* RNAi treatment. (C) Volcano plot of differentially expressed genes after 28

days of *phc* RNAi. (D) WISH analysis of selected genes indicated to be differentially

expressed after *phc* RNAi knockdown. The length of time each probe was developed

was equal between target RNAi and control RNAi samples. Arrows indicate up- or

down-regulated expression measured by RNA-seq. The red arrows highlight regions

with changed expression after RNAi in the worm's brain (B) and mouth (D) regions.

Arrowheads indicate regions of ectopic gene expression after RNAi treatment. Scale

bars = 200 μm.

Supplemental Figure Legends

Supplemental Figure S1. (A) WISH expression patterns for genes showing

phenotypes in the RNAi screen. All the genes examined were expressed throughout the

parenchyma; a subset of genes displayed enriched expression near the cephalic

ganglia or the intestine. (B) WISH analysis showing expression patterns of *prpf19*, *rnf2*,

and *phc* in untreated controls (right) and irradiated worms (left). Arrows show

expression in regions enriched in stem cells in untreated worms that are undetectable in

irradiated worms. Scale bars = 200 μm.

Supplemental Figure S2. (A-B) Double fluorescence *in situ* hybridization showing co-

expression of *prpf19* (magenta) with marker genes (green) for stem cells (*piwi-1*, *h2b*),

early epidermal progeny marker (*prog-1*), and late epidermal progeny marker (*agat-1*).

(C) Western blot probed using an anti-ubiquitin antibody for whole worm homogenate

protein extracts from *control(RNAi)* or *prp19f(RNAi)* treated animals collected at days

11, 14, and 18 of treatment. Scale bars in A-B = 20 μm.

Supplemental Figure S3. (A) WISH to NTC core elements *cdc5l*, *pflg1*, *spf27*, and

spliceosomal RNP members *prpf3* and *prpf8*. Scale bars = 200 μm.

Supplemental Figure S4. (A) Western blot analysis shows a reduction in H2Bub1

levels following inhibition of *bre1* at days 14, 21, and 28 of RNAi treatment. (B) Western

blot analysis shows no reduction in H2Aub1 levels following inhibition of canonical

PRC1 genes *ring1*, *cbx*, and *phc* after 21 days of RNAi treatment. (C) WISH analysis

showing expression patterns for PRC1 genes *phc* and *cbx*. (D) WISH marker gene

analysis for dorsoventral marker *bmp4* and anterior-posterior marker *ndl-3* showing no

change in expression domains following *phc* inhibition. Scale bars = 200 μm.

Supplemental Figure S5. (A) Feeding and sampling schedule for RNA-seq

experiments. Worms were fed twice a week and sampled for RNA extraction at the days

indicated by the red arrowheads. (B) Volcano plot of differentially expressed genes after

28 days of *rnf2* RNAi treatment. (C) Venn diagram showing the overlap between day 28

and day 14 sample sets for *rnf2(RNAi)*. (D) GO analysis of down-regulated genes in

rnf2(RNAi) worms after 28 days of treatment. (E) GO analysis of up-regulated genes in

rnf2(RNAi) worms after 28 days of treatment.

Table 1: RING/U-box E3 ubiquitin ligases showing phenotypes following RNAi.

Gene Name	Dresden Contig ID	Human RING or U-box Homolog	E-value	Phenotypes Observed
Smed-bre1	dd_Smed_v6_4070_0_1	RNF40	1e-88	HR, DR
Smed-march5	dd_Smed_v6_4602_0_1	MARCH5	8e-93	HR, Lesions, Lysis
Smed-*not4*	dd_Smed_v6_4767_0_1	CNOT4	5e-87	HR, VC
Smed-prpf19	dd_Smed_v6_1276_0_1	PRPF19	0.0	HR, VC, Lysis
Smed-ran	dd_Smed_v6_330_0_1	CBLB*	3e-94	Lesions, Lysis
Smed-ring1	dd_Smed_v6_12141_0_1	RING1	6e-33	DR
Smed-rnf2	dd_Smed_v6_8989_0_1	RNF2	6e-46	DR
Smed-rnf8-like	dd_Smed_v6_1137_0_5	RNF8	4e-05	DR, Lesions
Smed-traf2-like	dd_Smed_v6_3837_0_1	TRAF2	2e-69	HR

706 HR: Head regression. VC: Ventral Curling. DR: Delayed regeneration; *Top Human BLAST hit:
707 RAN, e-value: 9e-116

References

1   Elliott, S. A. & Sánchez Alvarado, A. The history and enduring contributions of
planarians to the study of animal regeneration. *Wiley Interdiscip Rev Dev Biol* **2**, 301-
326, doi:10.1002/wdev.82 (2013).

2   Ivankovic, M. *et al.* Model systems for regeneration: planarians. *Development* **146**,
dev167684, doi:10.1242/dev.167684 (2019).

3   Baguñà, J., Saló, E. & Auladell, C. Regeneration and pattern formation in planarians. III.
Evidence that neoblasts are totipotent stem cells and the source of blastema cells.
Development **107**, 77- 86 (1989).

4   Wagner, D. E., Wang, I. E. & Reddien, P. W. Clonogenic neoblasts are pluripotent adult
stem cells that underlie planarian regeneration. *Science* **332**, 811-816,
doi:10.1126/science.1203983 (2011).

5   Saló, E. & Baguñà, J. Cell movement in intact and regenerating planarians. Quantitation
using chromosomal, nuclear and cytoplasmic markers. *Development* **89**, 57-70,
doi:10.1242/dev.89.1.57 (1985).

6   Abnave, P. *et al.* Epithelial-mesenchymal transition transcription factors control
pluripotent adult stem cell migration in vivo in planarians. *Development* **144**, 3440-3453,
doi:10.1242/dev.154971 (2017).

7   Reddien, P. W. The Cellular and Molecular Basis for Planarian Regeneration. *Cell* **175**,
327-345, doi:10.1016/j.cell.2018.09.021 (2018).

8   Labbe, R. M. *et al.* A comparative transcriptomic analysis reveals conserved features of
stem cell pluripotency in planarians and mammals. *Stem Cells* **30**, 1734-1745,
doi:10.1002/stem.1144 (2012).

9   Onal, P. *et al.* Gene expression of pluripotency determinants is conserved between
mammalian and planarian stem cells. *EMBO J* **31**, 2755-2769,
doi:10.1038/emboj.2012.110 (2012).

10   van Wolfswinkel, J. C., Wagner, D. E. & Reddien, P. W. Single-cell analysis reveals
functionally distinct classes within the planarian stem cell compartment. *Cell Stem Cell*
15, 326-339, doi:10.1016/j.stem.2014.06.007 (2014).

11   Fincher, C. T., Wurtzel, O., de Hoog, T., Kravarik, K. M. & Reddien, P. W. Cell type
transcriptome atlas for the planarian Schmidtea mediterranea. *Science* **360**,
doi:10.1126/science.aaq1736 (2018).

12   Plass, M. *et al.* Cell type atlas and lineage tree of a whole complex animal by single-cell
transcriptomics. *Science*, doi:10.1126/science.aaq1723 (2018) (2018).

13   Boser, A. *et al.* SILAC proteomics of planarians identifies Ncoa5 as a conserved
component of pluripotent stem cells. *Cell Rep* **5**, 1142-1155,
doi:10.1016/j.celrep.2013.10.035 (2013).

14   Fernandez-Taboada, E., Rodriguez-Esteban, G., Salo, E. & Abril, J. F. A proteomics
approach to decipher the molecular nature of planarian stem cells. *BMC Genomics* **12**,
133, doi:10.1186/1471-2164-12-133 (2011).

15   Strand, N. S. *et al.* Dissecting the function of Cullin-RING ubiquitin ligase complex genes
in planarian regeneration. *Dev Biol* **433**, 210-217, doi:10.1016/j.ydbio.2017.10.011
(2018).

16   Ciechanover, A., Finley, D. & Varshavsky, A. Ubiquitin dependence of selective protein
degradation demonstrated in the mammalian cell cycle mutant ts85. *Cell* **37**, 57-66,
doi:10.1016/0092-8674(84)90300-3 (1984).

17   Endoh, M. *et al.* Histone H2A mono-ubiquitination is a crucial step to mediate PRC1-
dependent repression of developmental genes to maintain ES cell identity. *PLoS Genet*
8, e1002774, doi:10.1371/journal.pgen.1002774 (2012).

18   Higgins, R. *et al.* The Unfolded Protein Response Triggers Site-Specific Regulatory
Ubiquitylation of 40S Ribosomal Proteins. *Mol Cell* **59**, 35-49,
doi:10.1016/j.molcel.2015.04.026 (2015).

19   Nakayama, K. I. & Nakayama, K. Regulation of the cell cycle by SCF-type ubiquitin
ligases. *Semin Cell Dev Biol* **16**, 323-333, doi:10.1016/j.semcdb.2005.02.010 (2005).

20   Werner, A., Manford, A. G. & Rape, M. Ubiquitin-Dependent Regulation of Stem Cell
Biology. *Trends Cell Biol* **27**, 568-579, doi:10.1016/j.tcb.2017.04.002 (2017).

21   Li, W. *et al.* Genome-wide and functional annotation of human E3 ubiquitin ligases
identifies MULAN, a mitochondrial E3 that regulates the organelle's dynamics and
signaling. *PLoS One* **3**, e1487, doi:10.1371/journal.pone.0001487 (2008).

22   Lorick, K. L. *et al.* RING fingers mediate ubiquitin-conjugating enzyme (E2)-dependent
ubiquitination. *Proc Natl Acad Sci U S A* **96**, 11364-11369,
doi:10.1073/pnas.96.20.11364 (1999).

23   Aravind, L. & Koonin, E. V. The U box is a modified RING finger - a common domain in
ubiquitination. *Curr Biol* **10**, 132-134, doi:http://dx.doi.org/10.1016/S0960-
9822(00)00398-5 (2000).

24   Henderson, J. M. *et al.* Identification of HECT E3 ubiquitin ligase family genes involved
in stem cell regulation and regeneration in planarians. *Dev Biol* **404**, 21-34,
doi:10.1016/j.ydbio.2015.04.021 (2015).

25   Brandl, H. *et al.* PlanMine--a mineable resource of planarian biology and biodiversity.
Nucleic Acids Res **44**, D764-773, doi:10.1093/nar/gkv1148 (2016).

26   Roberts-Galbraith, R. H., Brubacher, J. L. & Newmark, P. A. A functional genomics
screen in planarians reveals regulators of whole-brain regeneration. *Elife* **5**,
doi:10.7554/eLife.17002 (2016).

27 Eisenhoffer, G. T., Kang, H. & Sánchez Alvarado, A. Molecular analysis of stem cells
and their descendants during cell turnover and regeneration in the planarian Schmidtea
mediterranea. *Cell Stem Cell* **3**, 327-339, doi:10.1016/j.stem.2008.07.002 (2008).

28 Lu, X. & Legerski, R. J. The Prp19/Pso4 core complex undergoes ubiquitylation and
structural alterations in response to DNA damage. *Biochem Biophys Res Commun* **354**,
968-974, doi:10.1016/j.bbrc.2007.01.097 (2007).

29 Hwang, W. W. *et al.* A Conserved RING Finger Protein Required for Histone H2B
Monoubiquitination and Cell Size Control. *Molecular Cell* **11**, 261-266,
doi:10.1016/s1097-2765(02)00826-2 (2003).

30 Henry, K. W. *et al.* Transcriptional activation via sequential histone H2B ubiquitylation
and deubiquitylation, mediated by SAGA-associated Ubp8. *Genes Dev* **17**, 2648-2663,
doi:10.1101/gad.1144003 (2003).

31 Wang, H. *et al.* Role of histone H2A ubiquitination in Polycomb silencing. *Nature* **431**,
873-878, doi:10.1038/nature02985 (2004).

32 Bunker, C. A. & Kingston, R. E. Transcriptional repression by Drosophila and
mammalian Polycomb group proteins in transfected mammalian cells. *Mol Cell Biol* **14**,
1721-1732, doi:10.1128/mcb.14.3.1721-1732.1994 (1994).

33 de Napoles, M. *et al.* Polycomb group proteins Ring1A/B link ubiquitylation of histone
H2A to heritable gene silencing and X inactivation. *Dev Cell* **7**, 663-676,
doi:10.1016/j.devcel.2004.10.005 (2004).

34 Gao, Z. *et al.* PCGF homologs, CBX proteins, and RYBP define functionally distinct
PRC1 family complexes. *Mol Cell* **45**, 344-356, doi:10.1016/j.molcel.2012.01.002 (2012).

35 Fursova, N. A. *et al.* Synergy between Variant PRC1 Complexes Defines Polycomb-
Mediated Gene Repression. *Mol Cell* **74**, 1020-1036 e1028,
doi:10.1016/j.molcel.2019.03.024 (2019).

36 Gavino, M. A. & Reddien, P. W. A Bmp/Admp regulatory circuit controls maintenance
and regeneration of dorsal-ventral polarity in planarians. *Curr Biol* **21**, 294-299,
doi:10.1016/j.cub.2011.01.017 (2011).

37 Rink, J. C., Gurley, K. A., Elliott, S. A. & Sánchez Alvarado, A. Planarian Hh signaling
regulates regeneration polarity and links Hh pathway evolution to cilia. *Science* **326**,
1406-1410, doi:10.1126/science.1178712 (2009).

38 Adler, C. E., Seidel, C. W., McKinney, S. A. & Sánchez Alvarado, A. Selective
amputation of the pharynx identifies a FoxA-dependent regeneration program in
planaria. *Elife* **3**, e02238, doi:10.7554/eLife.02238 (2014).

39 Tu, K. C. *et al.* Egr-5 is a post-mitotic regulator of planarian epidermal differentiation.
Elife **4**, e10501, doi:10.7554/eLife.10501 (2015).

40   Song, E. J. *et al.* The Prp19 complex and the Usp4Sart3 deubiquitinating enzyme control
reversible ubiquitination at the spliceosome. *Genes Dev* **24**, 1434-1447,
doi:10.1101/gad.1925010 (2010).

41   Reddien, P. W., Oviedo, N. J., Jennings, J. R., Jenkin, J. C. & Sánchez Alvarado, A.
SMEDWI-2 is a PIWI-like protein that regulates planarian stem cells. *Science* **310**, 1327-
1330, doi:10.1126/science.1116110 (2005).

42   Kim, I. V. *et al.* Planarians recruit piRNAs for mRNA turnover in adult stem cells. *Genes*
Dev **33**, 1575-1590, doi:10.1101/gad.322776.118 (2019).

43   Rouhana, L., Shibata, N., Nishimura, O. & Agata, K. Different requirements for
conserved post-transcriptional regulators in planarian regeneration and stem cell
maintenance. *Dev Biol* **341**, 429-443, doi:10.1016/j.ydbio.2010.02.037 (2010).

44   Solana, J. *et al.* The CCR4-NOT complex mediates deadenylation and degradation of
stem cell mRNAs and promotes planarian stem cell differentiation. *PLoS Genet* **9**,
e1004003, doi:10.1371/journal.pgen.1004003 (2013).

45   Francis, N. J., Kingston, R. E. & Woodcock, C. L. Chromatin compaction by a polycomb
group protein complex. *Science* **306**, 1574-1577, doi:10.1126/science.1100576 (2004).

46   Shao, Z. *et al.* Stabilization of Chromatin Structure by PRC1, a Polycomb Complex. *Cell*
98, 37-46, doi:10.1016/s0092-8674(00)80604-2 (1999).

47   Tamburri, S. *et al.* Histone H2AK119 Mono-Ubiquitination Is Essential for Polycomb-
Mediated Transcriptional Repression. *Mol Cell*, doi:10.1016/j.molcel.2019.11.021 (2019).

48   Blackledge, N. P. *et al.* PRC1 Catalytic Activity Is Central to Polycomb System Function.
Mol Cell **77**, 857-874 e859, doi:10.1016/j.molcel.2019.12.001 (2020).

49   Pengelly, A. R., Kalb, R., Finkl, K. & Muller, J. Transcriptional repression by PRC1 in the
absence of H2A monoubiquitylation. *Genes Dev* **29**, 1487-1492,
doi:10.1101/gad.265439.115 (2015).

50   Lewis, E. B. A gene complex controlling segmentation in Drosophila. *Nature* **276**, 565-
570, doi:10.1038/276565a0 (1978).

51   Conway, E. M. & Bracken, A. P. in *Polycomb Group Proteins*   57-80 (2017).

52   Cao, R., Tsukada, Y. & Zhang, Y. Role of Bmi-1 and Ring1A in H2A ubiquitylation and
Hox gene silencing. *Mol Cell* **20**, 845-854, doi:10.1016/j.molcel.2005.12.002 (2005).

53   Blackledge, N. P. *et al.* Variant PRC1 complex-dependent H2A ubiquitylation drives
PRC2 recruitment and polycomb domain formation. *Cell* **157**, 1445-1459,
doi:10.1016/j.cell.2014.05.004 (2014).

54   Gahan, J. M., Rentzsch, F. & Schnitzler, C. E. The genetic basis for PRC1 complex
diversity emerged early in animal evolution. *Proc Natl Acad Sci U S A,*
doi:10.1073/pnas.2005136117 (2020).

55   Peng, Y. C., Lv, T. H., Du, Z. K., Cun, X. N. & Yang, K. M. Liver Macrophages Stimulate
the Expression of Hepatocyte Nuclear Factor-6 and Promote Hepatocyte Proliferation at
the Early Stage of Liver Regeneration. *Bull Exp Biol Med* **170**, 40-45,
doi:10.1007/s10517-020-05000-7 (2020).

56   van der Raadt, J., van Gestel, S. H. C., Nadif Kasri, N. & Albers, C. A. ONECUT
transcription factors induce neuronal characteristics and remodel chromatin accessibility.
Nucleic Acids Res **47**, 5587-5602, doi:10.1093/nar/gkz273 (2019).

57   Jiang, K. *et al.* HNF6 promotes tumor growth in colorectal cancer and enhances liver
metastasis in mouse model. *J Cell Physiol* **234**, 3675-3684, doi:10.1002/jcp.27140
(2019).

58   Lee, T. I., Johnstone, S. E. & Young, R. A. Chromatin immunoprecipitation and
microarray-based analysis of protein location. *Nat Protoc* **1**, 729-748,
doi:10.1038/nprot.2006.98 (2006).

59   Skene, P. J. & Henikoff, S. An efficient targeted nuclease strategy for high-resolution
mapping of DNA binding sites. *Elife* **6**, doi:10.7554/eLife.21856 (2017).

60   Merryman, M. S., Sánchez Alvarado, A. & Jenkin, J. C. Culturing Planarians in the
Laboratory. *Methods Mol Biol* **1774**, 241-258, doi:10.1007/978-1-4939-7802-1_5 (2018).

61   Rozanski, A. *et al.* PlanMine 3.0-improvements to a mineable resource of flatworm
biology and biodiversity. *Nucleic Acids Res* **47**, D812-D820, doi:10.1093/nar/gky1070
(2019).

62   Blum, M. *et al.* The InterPro protein families and domains database: 20 years on. *Nucleic
Acids Res* **49**, D344-D354, doi:10.1093/nar/gkaa977 (2021).

63   Zayas, R. M. *et al.* The planarian Schmidtea mediterranea as a model for epigenetic
germ cell specification: analysis of ESTs from the hermaphroditic strain. *Proc Natl Acad
Sci U S A* **102**, 18491-18496, doi:10.1073/pnas.0509507102 (2005).

64   Liu, S. Y. *et al.* Reactivating head regrowth in a regeneration-deficient planarian species.
Nature **500**, 81-84, doi:10.1038/nature12414 (2013).

65   Adler, C. E. & Sánchez Alvarado, A. Systemic RNA Interference in Planarians by
Feeding of dsRNA Containing Bacteria. *Methods Mol Biol* **1774**, 445-454,
doi:10.1007/978-1-4939-7802-1_17 (2018).

66   Rouhana, L. *et al.* RNA interference by feeding in vitro-synthesized double-stranded
RNA to planarians: methodology and dynamics. *Dev Dyn* **242**, 718-730,
doi:10.1002/dvdy.23950 (2013).

67   Gurley, K. A., Rink, J. C. & Sánchez Alvarado, A. β-Catenin Defines Head Versus Tail
Identity During Planarian Regeneration and Homeostasis. *Science* **319**, 323-327 (2008).

68   Pearson, B. J. *et al.* Formaldehyde-based whole-mount in situ hybridization method for
planarians. *Dev Dyn* **238**, 443-450, doi:10.1002/dvdy.21849 (2009).

69   King, R. S. & Newmark, P. A. *In situ* hybridization protocol for enhanced detection of
gene expression in the planarian *Schmidtea mediterranea*. *BMC Developmental Biology*
13 (2013).

70   Lauter, G., Soll, I. & Hauptmann, G. Two-color fluorescent in situ hybridization in the
embryonic zebrafish brain using differential detection systems. *BMC Dev Biol* **11**, 43,
doi:10.1186/1471-213X-11-43 (2011).

71   Brown, D. D. R. & Pearson, B. J. in *In Situ Hybridization Methods Neuromethods* Ch.
Chapter 7, 127-150 (2015).

72   Pellettieri, J. *et al.* Cell death and tissue remodeling in planarian regeneration. *Dev Biol*
338, 76-85, doi:10.1016/j.ydbio.2009.09.015 (2010).

73   Simoes, A. E. *et al.* Efficient recovery of proteins from multiple source samples after
TRIzol((R)) or TRIzol((R))LS RNA extraction and long-term storage. *BMC Genomics* **14**,
181, doi:10.1186/1471-2164-14-181 (2013).

74   Duncan, E. M., Chitsazan, A. D., Seidel, C. W. & Sánchez Alvarado, A. Set1 and MLL1/2
Target Distinct Sets of Functionally Different Genomic Loci In Vivo. *Cell Rep* **13**, 2741-
2755, doi:10.1016/j.celrep.2015.11.059 (2015).

75   Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq
quantification. *Nat Biotechnol* **34**, 525-527, doi:10.1038/nbt.3519 (2016).

76   Huber, W. *et al.* Orchestrating high-throughput genomic analysis with Bioconductor. *Nat*
Methods **12**, 115-121, doi:10.1038/nmeth.3252 (2015).

77   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15**, 550, doi:10.1186/s13059-014-0550-8
(2014).

Figure 1

A

B

control(RNAi)	prpf19(RNAi)	not4(RNAi)	ran(RNAi)	march5(RNAi)	bre1(RNAi)
d20	d22	d28	d15	d28	d21

C

	control(RNAi)	rnf2(RNAi)	ring1(RNAi)	traf2-like(RNAi)	rnf8-like(RNAi)
7 dpa head					
7 dpa tail					

Supplemental Figure S1

A

cephalic ganglia	intestinal

prpf19 *not4* *march5* *bre1* *rnf2* *traf2-like* *rnf8-like* *ring1* *ran*

B

	Untreated	Irradiated

prpf19

rnf2

Figure 2

Supplemental Figure S2

Figure 3

A NineTeen Complex (NTC)

PRLG1 CDC5L

SPF27 PRPF19

E3 Ligase Activity

Ub Ub Ub

U5 snRNP PRPF8   PRPF3

Stabilized Association

U4/U6 snRNP

B

cdc5l(RNAi)   *prlg1(RNAi)*

d14   d11

C

control (RNAi) *spf27 (RNAi)* *control (RNAi)* *spf27 (RNAi)*

Head

Tail

D

prpf3(RNAi)   *prpf8(RNAi)*

d19   d18

Supplemental Figure S3

Figure 4

A

B

C

D

Supplemental Figure S4

A day 14   day 21   day 28

RNAi: control   bre1   control   bre1   control   bre1

ub-H2B

β-tubulin

B day 21

RNAi: control   rnf2   ring1   cbx   phc

ub-H2A

C *phc*   *cbx*

D *bmp-4*   *ndl-3*

control

phc(RNAi)   d28   d28

Figure 5

A

Total number of differentially expressed genes identified by RNA-seq at FDR value < 0.1.

Gene name	Days of RNAi	Down-regulated genes	Up-regulated genes	Total
Smed-phc	11	20	29	49
Smed-rnf2	14	16	13	29
Smed-rnf2	28	113	134	247

Supplemental Figure S5

A

No. of feeds: 1   2   3   4   5   6   7   8

Days after first feed: 1   7   11   14   21   28

Sampling days and genes: *phc*   *rnf2*   *rnf2*

B

C

Day 14   17   12   Day 28   235

D

Down-regulated genes GO term

- cellular catabolic process
- primary metabolic process
- organic substance catabolic process
- organic substance metabolic process

$-Log_{10}$ (*P*-value)

E

Up-regulated genes GO term

- response to hypoxia
- alpha-amino acid catabolic process
- aromatic amino acid family catabolic process
- proteasomal ubiquitin-independent protein catabolic process
- cellular amino acid catabolic process
- neutrophil degranulation
- regulation of cellular amino acid metabolic process
- regulation of transcription from RNA polymerase II promoter in response to stress
- ATF6-mediated unfolded protein response
- chaperone cofactor-dependent protein refolding
- protein-containing complex assembly
- protein folding in endoplasmic reticulum
- antigen processing and presentation of exogenous peptide antigen via MHC class I, TAP-dependent
- cellular response to decreased oxygen levels
- glycosyl compound catabolic process
- protein refolding
- cellular biosynthetic process
- regulation of stem cell differentiation

$-Log_{10}$ (*P*-value)