Genetic diversity loss in the Anthropocene

Moises Exposito-Alonso[1,2,3*], Tom R. Booker[4,5,&], Lucas Czech[1,&], Tadashi Fukami[2,&], Lauren Gillespie[1,6,&], Shannon Hateley[1,&], Christopher C. Kyriazis[7,&], Patricia L. M. Lang[2,&], Laura Leventhal[1,2,&], David Nogues-Bravo[8,&], Veronica Pagowski[2,&], Megan Ruffley[1,&], Jeffrey P. Spence[9,&], Sebastian E. Toro Arana[1,2,&], Clemens L. Weiß[9,&], Erin Zess[1,&].

[1]Department of Plant Biology, Carnegie Institution for Science, Stanford, CA 94305, USA.
[2]Department of Biology, Stanford University, Stanford, CA 94305, USA.
[3]Department of Global Ecology, Carnegie Institution for Science, Stanford, CA 94305, USA.
[4]Department of Zoology, University of British Columbia, Vancouver, Canada.
[5]Biodiversity Research Centre, University of British Columbia, Vancouver, Canada.
[6]Department of Computer Science, Stanford University, Stanford, CA 94305, USA.
[7]Department of Ecology and Evolutionary Biology, University of California, Los Angeles, CA 90095, USA.
[8]Center for Macroecology, Evolution and Climate, GLOBE Inst., Univ. of Copenhagen, Copenhagen, Denmark.
[9]Department of Genetics, Stanford University, Stanford, CA 94305, USA.

[&]Authors are listed alphabetically
[*]To whom correspondence should be addressed: moisesexpositoalonso@gmail.com

Keywords: extinction, genetic diversity, climate change, habitat loss, Anthropocene

More species than ever before are at risk of extinction due to anthropogenic habitat loss and climate change. But even species that are not threatened have seen reductions in their populations and geographic ranges, likely impacting their genetic diversity. Although preserving genetic diversity is key to maintaining adaptability of species, we lack predictive tools and global estimates of genetic diversity loss across ecosystems. By bridging theories of biodiversity and population genetics, we introduce a mathematical framework to understand the loss of naturally occurring DNA mutations within decreasing habitat within a species. Analysing genome-wide variation data of 10,095 geo-referenced individuals from 20 plant and animal species, we show that genome-wide diversity follows a power law with geographic area (the mutations-area relationship), which can predict genetic diversity loss in spatial computer simulations of local population extinctions. Given pre-21st century values of ecosystem transformations, we estimate that over 10% of genetic diversity may already be lost, surpassing the United Nations targets for genetic preservation. These estimated losses could rapidly accelerate with advancing climate change and habitat destruction, highlighting the need for forecasting tools that facilitate implementation of policies to protect genetic resources globally.

Anthropogenic habitat loss and climate change (*1*, *2*) have led to the extinction of hundreds of species over the last centuries (*1*, *2*) and approximately one million more species (25% of all known species) are at risk of extinction (*3*). It has been estimated that an even larger fraction—at least 47%—of plant and animal species have lost part of their geographic range in response to the last centuries of anthropogenic activities (*4*, *5*). Though this loss might seem inconsequential compared to losing an entire species, this range contraction reduces genetic diversity, which dictates species' ability to adapt to new environmental conditions (*6–8*). The loss of geographic range can spiral into a feedback loop where diversity loss further increases the risk of species extinction (*9*, *10*).

Although genetic diversity is a key dimension of biodiversity (*11*), it has been overlooked in international conservation initiatives. Only in 2021 did the United Nations' Convention of Biological Diversity propose to preserve at least 90% of all species' genetic diversity (*12*, *13*). Although analyses of genetic markers in animal populations sampled over time with the aim of quantifying recent genetic change are emerging (*14*, *15*) and simulation studies with species distribution models or sensitivity analyses suggest within-species range variation may be strongly impacted (*5*, *16*, *17*), theory and scalable approaches to estimate genome-wide diversity loss across species do not yet exist, impairing prioritization and evaluation of conservation targets. Here, we introduce a framework to estimate global genetic diversity loss by bridging biodiversity theory with population genetics, and by combining data on global ecosystem transformations with newly available genomic datasets.

The first studies that predicted biodiversity reductions in response to habitat loss and climate change in the 1990s and the 2000s projected species extinctions using the relationship of biodiversity with geographic area—termed the species-area relationship (SAR) (*18*) (see **Supplementary Materials [SM] I** for a comparison of mathematical models for predicting biodiversity). In this framework, ecosystems with a larger area (*A*) harbour a larger number of species (*S*) resulting from a balance of limited dispersal, habitat heterogeneity, and colonisation-extinction-speciation dynamics. The more a study area is extended, the more species are found. The SAR has been empirically shown to follow a power law, $S = A^z$. It scales consistently across continents and ecosystems (*19*), with a higher z characterising more speciose and spatially structured ecosystems. Given estimates of decreasing ecosystem areas over time ($A_{t-1} > A_t$), Thomas et al. (*20*) proposed rough estimates of the percentage of species extinctions in the 21[st] century ranging from 15 to 37% (**SM I.3**). Though this may be an oversimplification, SAR has become a common tool for policy groups including the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES) (*3*).

As species richness is for to ecosystems' biodiversity, within-species variation can be quantitatively described by the richness of genetic mutations within a species, defined here as DNA nucleotide variants appearing in individuals of a species. Although population genetics theory has long established that larger populations have higher genetic diversity (*21*), and it is known that geographic isolation between populations within the same species results in geographically separated accumulation of different mutations, there have been no attempts to describe the extent of genetic diversity loss driven by species' geographic range reduction using an analogous "mutations-area relationship" (MAR).

We suspected that such a mutations-area relationship must exist given that another general assumption is shared with species studies, namely that when mutations appear they are first in only one individual, and they typically remain at low frequency in a population, though a few prevail to high frequency through stochastic genetic drift and natural selection (*22*). This principle of "commonness of rarity" is well-known for species (i.e. most species in an ecosystem are rare while only a few are

92   common) and, together with limited spatial dispersal of species and communities, is a key statistical
93   condition that led to the power-law SAR.
94
To examine the expectation of a power-law MAR, we begin quantifying the rarity of mutations
using millions of biallelic genetic variants of the *Arabidopsis thaliana* 1001 genomes dataset (**Fig. 1A**)
(*23*) by fitting several common models of species abundances (*24*) to the distribution of mutation
frequencies (*q*), termed the Site Frequency Spectrum in population genetics (**Fig. 1B**, **SM II.1**). The
canonical L-shaped probability distribution (*1/q*) of this spectrum—which is expected under
population-equilibrium and the absence of natural selection processes—fit this data well (Fig. 1B),
although the more parameter rich Preston's species abundance log-normal model achieved the best AIC
value (**Fig. 1B**, **SM III.1, Table S3, Table S10**). Despite the small differences in fit, these models all
showcase the similarities of abundance distributions of mutations within species and species within
ecosystems, suggesting that they may behave similarly in their relationship to geographic area (*22*, *24*).
105
106

107
Fig. 1 | Mutations across populations follow a log-normal abundance distribution and a power
law with species range area. (**A**) Density of individuals projected in a 1 x 1 degree latitude/longitude
map of Europe and exemplary subsample areas of different sizes. (**B**) Distribution of mutation (SNPs)
frequencies in 1,001 *Arabidopsis thaliana* plants using a site frequency spectrum histogram (grey inset)
and a Whittaker's rank abundance curve plot, and the fitted models of common species abundance
functions in *A. thaliana* using a dataset random sample of 10,000 mutations also used in (C). The AIC
fit of the three models is indicated with respect to the top model, log-normal. (**C**) The mutations-area

relationship (MAR) in log-log space built from 10 random subsamples of different areas of increasing
size within *A. thaliana*'s geographic range along with the number of mutations discovered for each area
subset.
118
To quantify how genetic diversity within a species increases with geographic area, we
constructed the MAR by subsampling different regions of different sizes of *Arabidopsis thaliana*'s
native range using over one thousand geo-referenced genomes (**Fig. 1A, C**). As a metric of genetic
diversity, we modelled the number of mutations (*M*) in space (number of segregating sites) consistent
with the species-centric approach of SAR, which uses species richness as the metric of biodiversity
(**SM II.2**). The MAR also followed the power law relationship $M = cA^z$ with a scaling value $z_{MAR} =$
0.324 (CI95% = 0.238–0.41) (**Fig. 1C**). Naturally, subsamples of larger areas may also contain more
individuals, and therefore should also have more mutations. But the observed power law relationship
goes beyond what is expected from the increase of number of samples in an area (which only accounts
for increases of $M \approx log(A)$, see theoretical derivation **SM II.3**). The remainder may be attributed to
population genetic drift and spatial natural selection causing structuring of genetic diversity across
populations. The discovered power law scaling appears robust to different methods of area
quantification, the effects of non-random spatial patterns, random area sampling, fully nested outward
or inward sampling (*19*), raster area calculations, raster grid resolution (~10–1,000 km side cell size),
and is adjusted for limited sample sizes (**SM II.3.2, III.3, Fig. S14-18, Tables S7-9**).
134
We then wondered whether MAR can predict the loss of genetic diversity due to species' range
contractions. We explored several scenarios of range contraction in *A. thaliana* by removing *in silico*
grid cells in a map representing populations that are lost (**Fig. 2B**). Our simulations included random
local population extinction as if deforestation was scattered across large continents, radial expansion of
an extinction front due to intense localised mortality, or local extinction in the warmest regions within
a species range (*4, 25*), among others (**SM III.4**). The MAR-based predictions of genetic loss, using *1-*
$(1-A_t / A_{t-1})^z$ and assuming $z = 0.3$, conservatively followed the simulated local loss in *A. thaliana*
(pseudo-R^2 = 0.87, taking all simulations together) (**SM II.4, III.4**).
143
Since genetic diversity is ultimately created by spontaneous DNA errors passed onto offspring
every generation, the loss of genetic diversity seems reversible, as these mutations could happen again.
However, the recovery of genetic diversity through natural mutagenesis is extremely slow (*57*),
especially for mutations affecting adaptation. Simulating a species undergoing only a 5–10% in area
reduction, it would take at least ≈140–520 generations to recover its original genetic diversity (2,100–
7,800 years for a fast-growing tree or medium-lifespan mammal of 15 year generation length), although
for most simulations, recovery virtually never happened over millennia (see **SM II.4-5, Fig. S11, SM**
III.6).
152
153
154

A Genetic diversity loss by geographic area loss using the mutations-area relationship

Fig. 2 | The power law of genetic diversity loss with range area loss. (A) Percentage of loss of total genetic diversity in *Arabidopsis thaliana* from several stochastic simulations (red) of local extinction in (B), and theoretical model projections of genetic diversity loss using the MAR (dotted lines). The expectation for genetic diversity loss based only on individuals is in grey (using starting populations of $N=10^4$-10^9) (**SM II.4**). **(B)** Cartoon of several possible range contractions simulated by progressively removing grid cells across the map of Eurasia (red/grey boxes) following different hypothesised spatial extinction patterns. (**C**) A metric of adaptive capacity loss during warm edge extinction in (B). Using Genome Wide Associations (GWA) to estimate effects of mutation on fitness in different rainfall conditions, water use efficiency [wue], flowering time, seed dormancy, plant growth rate, and plant size. Plotted are the fraction loss of the summed squared effects ($\sum a^2$) of 10,000 mutations from the top 1% tails of effects. We also plot (yellow) the fraction of protein-coding alleles lost (nonsynonymous, stop codon loss/gain, and frameshift mutations).

To test the generality of the MAR, we searched in public nucleotide repositories for datasets of hundreds to thousands of whole-genome sequenced individuals for the same species sampled across geographic areas within their native ranges (**Table 1, SM IV**). In total, we identified 20 wild plant and animal species with such published resources and assembled a dataset amassing a total of 10,095 individuals of these species, with 1,522 to 88,332,015 naturally occurring mutations per species, covering a geographic area ranging from 0.03 to 115 million km^2. Fitting MAR for these diverse species, we recovered z_{MAR} values similar to *A. thaliana*, with many species overlapping in confidence intervals, with the exception of some outliers (mean (SE) $z_{MAR} = 0.31$ (± 0.038), median = 0.26, IQR = ± 0.15, range=0.10–0.82, mean (SE) z^*_{MAR} scaled = 0.26 (± 0.048). See **Table 1, SM IV, Fig. S22, Table S10).** Theoretical derivations show that z_{MAR} is a consequence of fundamental evolutionary and ecological forces (mutation, dispersal, selection) and should range from 0 to 1, depending on the strength of population structure (**SM II.3,** see **Fig. S10** for its relationship with isolation-by-distance). These predictions were further confirmed by spatial population genetics coalescent and individual-based

183 simulations in 2D and continuous space (**SM II.3**), as well as with mainland-island community
184 assembly simulations according to the Unified Neutral Theory of Biodiversity (UNTB) (**SM V.3**).
185
186
Table 1 |The mutations-area relationship across diverse species. Summary statistics of individuals
sampled broadly across species distributions, sequencing method and mutations studied, and convex
hull area extent of all samples within a species. The mutations-area relationship (MAR) parameter z,
which captures how spatially restricted mutations are, including a scaled correction z^* for low sampling
genomic effort. Percent area that needs to be kept for a species to maintain 90% of its genetic diversity,
using the per-species MAR value estimates. Area predictions are not provided for threatened species,
as these have likely already lost substantial genetic diversity and require protection of their full
geographic range (Fig. 3).

Species	N	M_{tot} Method	A_{tot} Km²x10⁶	MAR z [CI95%]	MAR scaled z* [CI95%]	Min area 90% %
Arabidopsis thaliana	1,135 (1,001)[#]	11,769,920 W	27.34	0.324 (0.238–0.41)	0.312 (0.305 - 0.32)	71–78
Arabidopsis lyrata	108	17,813,817 W	2.79	0.236 (0.218–0.254)	0.151 (0.137–0.165)	50–66
Amaranthus tuberculatus	162 (155)	1,033,443 W	0.80	0.109 (0.081–0.136)	0.142 (0.136–0.149)	48–65
Eucalyptus melliodora[VU]	275 (36)[*]	9,378 GBS	0.95	0.466 (0.394–0.538)	0.403 (0.398–0.407)	77–82
Yucca brevifolia[CA]	290	10,695 GBS	NA	?0.128 (0.109–0.147)	0.049 (0.037–0.062)	-
Mimulus guttatus	521 (286)[#*]	1,522 GBS	25.14	0.274 (0.259–0.29)	0.231 (0.221–0.241)	63–73
Panicum virgatum	732 (576)[†]	33,905,044 W	6.29	0.232 (0.211–0.252)	0.126 (0.116–0.136)	43–63
Panicum hallii	591	45,589 W	2.19	0.824 (0.719 - 0.928)	0.814 (0.745 - 0.883)	88–90
Pinus contorta	929	32,449 GC	0.89	?0.015 (0.014–0.016)	-0.061(-0.062-0.060)	-
Pinus torreyana[CR]	242	478,238 GBS	NA	?0.236 (0.19–0.282)	0.105 (0.099–0.11)	-
Populus trichocarpa	882	28,342,826 W	1.12	0.275 (0.218–0.332)	0.165 (0.155–0.176)	53–67
Anopheles gambiae	1142 (29)[*]	52,525,957 W	19.96	0.214 (0.164–0.264)	0.122 (0.111–0.132)	42–62
Acropora millepora[NT]	253 (12)[*]	17,931,448 W	0.03	0.246 (0.209–0.283)	0.287 (0.28–0.294)	69–77
Drosophila melanogaster	271[%]	5,019 W	115.21	0.437 (0.397–0.477)	0.325 (0.314–0.336)	72–79
Empidonax traillii Decline	219 (199)[&]	349,014 GBS/GC	7.03	0.214 (0.174–0.254)	0.074 (0.047–0.102)	24–54
Setophaga petechia[Decline]	199	104,711 GBS	15.17	0.251 (0.236 - 0.267)	0.149 (0.135 - 0.163)	49--66
Peromyscus maniculatus	80 (78)[&]	14,076 GBS	22.61	0.488 (0.264–0.713)	0.683 (0.615–0.751)	86--88
Dicerorhinus sumatrensis[CR]	16	8,870,513 W	NA	?0.412 (0.369–0.456)	0.127 (0.11–0.144)	-
Canis lupus	349 (230)[†]	1,517,226 W	19.10	0.256 (0.232–0.28)	0.184 (0.175–0.193)	56–70
Homo sapiens	2504 (24)[*]	88,332,015 W	80.76	?0.431 (0.347–0.514)	0.281 (0.23–0.332)	NA

[#]*Only individuals in the native range were used for the analyses.*
[&]*Only individuals with available coordinates or matching IDs were used for analyses.*
[%]*Numbers indicate pools of flies used for Pool-Sequencing.*
[*]*Number of geographically separated populations, as multiple individuals were collected per population.*
[†]*Only natural populations were used, excluding breeds, landraces, and cultivars.*
Area was not reported for species with unknown locations or where less than 2 populations were sampled.
?*Values excluded from global averages used for conservation applications due to uncertain estimates, suboptimal genomic data type, or*
because estimates should not be applied for conservation (i.e. humans or nearly extinct Sumatran rhinoceros).
Acronyms: W = whole-genome re-sequencing or discovery SNP calling. GBS = genotyping by sequencing of biallelic SNP markers. GC =
genotyping chip; CR = Red List Critically Endangered. VU= Red List Vulnerable. CA = included in the California Endangered Species Act.
Decline = population decline reported in the Red List.
206
207

Although we expect species-specific traits related to dispersibility or gene flow to affect z_{MAR}
(e.g. migration rate and environmental selection in population genetic simulations significantly
influences z_{MAR}, **Table S2**), no significant association was found between z_{MAR} and different
ecologically-relevant traits, mating systems, home continents, etc., for the 20 species analysed. Perhaps
this is simply that there are still too few species that have large population genomic data to find such a
signal (**Table 1**, **Table S12-13**). Nevertheless, the relative consistency of z_{MAR} across largely different
species may be promising for conservation purposes, as an average z_{MAR} ~0.3 (IQR ±0.15 , **Table 1,**
Table S11) could be predictive of large-scale trends of genetic diversity loss in many range-reduced
species that lack genomic information. Further, although species will naturally have different starting

levels of total genetic diversity prior to range reductions, for instance, due to genome size, structure, or mating system differences (*26*), the application of z_{MAR} provides relative estimates of genetic diversity loss. For instance, assuming z_{MAR} ~0.3, we would predict that an area reduction of ~50% creates an approximate loss of ~20% of genetic diversity relative to the total genetic diversity of a given species.

Finally, we used MAR to estimate the average global genetic diversity loss caused by pre-21st century land transformations. Although accurate species-specific geographic area reduction data in the last centuries are scarce, we leveraged global land cover transformations from primary ecosystems to urban or cropland systems (*3*, *27*) (**Table S14-15**). Using the average scaled z^{*}_{MAR} (**Table S18**) and several global averages of Earth's land and coastal transformations for present day (38% global area transformation from (*27*), 34% from (*28*), and 43-50% from (*29*)), we estimate a 10-16% global genetic diversity loss on average across species (**Fig. 3A**). While these estimates may correctly approximate central values across species in an ecosystem, we expect a substantial variation in the extent of loss across species, ranging theoretically from 0 to 100% (**Fig. 3, Fig. S26**). One cause of this variation is the heterogeneity in land cover transformations across ecosystems; for example, more pristine high-altitude systems have only lost 0.3% of their area, while highly managed temperate forests and woodlands have lost 67% (**Fig. 3B**, **Table S14-15)**.

Another cause for the variability in genetic loss among species (even within the same ecosystem) may be their differential geographic ranges and abundances, life histories, or species-specific threats. We gathered data from species red-listed by the International Union for Conservation of Nature (IUCN) (*1*), which evaluates recent population or geographic range area reduction over ±10 years / ±3 generations to place assessed species in different threat categories using several thresholds (guidelines for assessments and thresholds available at www.iucn.org). Again, assuming that with the average z_{MAR} ~0.3 we can capture general patterns, we translate these category thresholds into genetic diversity loss (**Fig. 3C,** see **SM V, Table S17)**. *Vulnerable* species, having lost at least 30% of their geographic distribution, may have experienced >9% of genetic diversity loss, *endangered* species, which have lost over 50% of their geographic distribution, should have incurred >16% of genetic diversity loss, and *critically endangered* species, with over 80% area reduction, likely suffered >33% of genetic diversity loss (**Fig. 3B**). This clearly showcases that even species in no imminent risk of extinction (e.g. least concern, near threatened, vulnerable), such as the majority of species for which population genomic data exists, may already be losing substantial genetic diversity (**Fig. 3A**).

Fig. 3 | The parameter space of genetic diversity loss mapping pre-21st century ecosystem transformations and species threat categories against possible values of the mutations-area relationship. (A) Possible values of two key parameters, the mutations-area relationship scaling parameter (MAR) and % of area reduction of a species geographic range (as a proxy of entire ecosystem transformation). The theoretical % of genetic diversity loss is represented as filled grey colour, with isolines in white. Estimates of scaled z^*_{MAR} from Table 1 per species are in orange with their 95% confidence intervals (for unscaled z_{MAR} see Fig. S23). Although exact area losses per species are unknown, species are plotted based on their IUCN Red list (C) status, using the broad ranges of minimum and maximum recent population or area decline per category. The global average is calculated with the average z_{MAR} across species and % of the Earth transformed from IPBES. **(B)** Percentage of transformed ecosystem area from the Millennium Ecosystem Assessment (MEA) (*27*) are represented by light blue arrows, from the Intergovernmental Science-Policy Panel for Biodiversity and Ecosystem Services (IPBES) (*28*) for 2010 and 2050 are dark blue arrows, and from the Land Use Harmonization 2 (LUH2) dataset (*29*) are in dark purple. **(C)** The minimum criterion value of population or geographic area loss to be classified in each category of the IUCN Red List are indicated with pink arrows (the near threatened category does not have a range of values, instead we used 30% ±10%). The number of plant species (for which population abundance loss approximates area loss) included in each category is shown as box sizes (*1*). The IUCN ranges were used to place ranges of estimates in (A) per species.

The ultimate challenge is to understand how genetic diversity loss relates to loss of adaptive capacity of a species. To this end, we leveraged the extensive knowledge of the effect of mutations in ecologically relevant traits in *A. thaliana* from Genome-Wide Associations (GWA) (**Fig. 2C, SM III**). We again conducted spatial warm edge extinction simulations, this time tracking metrics of adaptive capacity, including the total sum of effects estimated from GWA of remaining mutations ($\sum_i a_i$ for i=1…10,000 variants of putative a_i effect), the additive genetic variance ($Va = \sum_i p_i(1-p_i)a_i^2$, which accounts for each variant's population frequency p_i), and the loss of nonsynonymous mutations (**SM III.5**). Although determining the effect of mutations through GWA is technically challenging even in model species (*30, 31*), and variants may even be either deleterious or advantageous depending on genomic backgrounds (*32*) or environments (*33*), our simulations suggest putatively functional

mutations may be lost more slowly ($z<0.3$, **Fig. 2C**) than neutral genetic diversity (**Fig. 2A)**. In fact,
the additive variance Va parameter, often equated to the rate of adaptation, appears rather stable (*34*)
until just before the extinction event when it sharply collapses (**Fig. S21**; see also **Fig. 2C,** and **SM**
II.3.4 for simulations that replicate this pattern). This is analogous to the famous "rivet popper"
metaphor where ecosystem structure and function may suddenly collapse as species are inadvertently
lost (*35*). Projections of the MAR using genome-wide variation may crucially serve as early
conservation tool in non-threatened species (*36*, *37*), before species reach accelerating collapsing
extinction dynamics—an acceleration that we expect to be even more dramatic due to elevated drift and
accumulation of deleterious mutations of small critically-endangered populations (*38*, *39*).

To achieve the recently published United Nations target to protect "at least 90% of genetic
diversity within all species"(*13*), it will be necessary to aggressively protect as many populations as
possible for each species. Here, we have discovered the existence of a mutations-area relationship
(MAR) and provided a mathematical framework to forecast genetic diversity loss with shrinking
geographic species ranges. The MAR contrasts with existing studies on the risk of losing entire species
by focusing on quantifying the magnitude and dynamics of genetic diversity loss likely ongoing in most
species. This framework demonstrates that even with conservative estimates, substantial area protection
will be needed to meet the UN Sustainable Development Goals. For vulnerable or endangered species,
we may have likely already failed.

ADDITIONAL INFORMATION

Author contribution M.E.-A. conceived and led the project. M.E.-A., J.P.S., M.R., S.H., L.G., L.C.,
L.L., S.T.A., V.P., E.Z., P.L.M.L., C.C.K., T.B., C.W. conducted research, all authors interpreted the
results and wrote the manuscript.

Data availability. The analysed datasets are publicly available or were shared by authors upon request
(see Supplementary Materials for details). Code is available at Github
(https://github.com/moiexpositoalonsolab/mar) and Zenodo (https://doi.org/10.5281/zenodo.6408624).

Acknowledgements. We are grateful for many researchers who made their genomic data publicly
available and thus made this research possible: the 1001 Arabidopsis Genomes Consortium, the
Anopheles gambiae 1000 Genomes Consortium, the 1000 (Human) Genome Consortium, the European
"Drosophila Evolution over Space and Time" (DEST) Consortium, MacLachlan et al., Fuller et al.,
Ruegg et al., Kingsley et al., Schweizer et al., and Royer et al.. In addition, we are grateful to all the
authors who directly shared intermediate genome variant files: Sujan Mamidi, John Lovell, Dan
Jacobson, Manesh Shah, Julia Kreiner, Kay Lucek, Yvonne Willi, Juan D. Palacio Mejia, Justin
Borevitz, Megan Supple, Mario Vallejo-Marin, Nicolas Dussex, Lionel Di Santo, and Jill Hamilton.
We thank John Wiens, Sue Rhee, Detlef Weigel, Ellie Armstrong, Marty Kardos, Dmitri Petrov, Rob
Colwell, Rasmus Nielsen, Anna Michalak, Sean Hoban, Jonathan Pritchard, and members of the Moi
and Mordecai Labs for comments, discussion, or references. M.E.-A. is supported by the Office of the
Director of the National Institutes of Health's Early Investigator Award with award number:
1DP5OD029506-01; by the U.S. Department of Energy, Office of Biological and Environmental
Research, grant number: DE-SC0021286; and by the Carnegie Institution for Science. J.P.S. is
supported by an NIH training grant (5T32HG000044-23), and S.T.A by the NIGMS Center of the NIH
under award number T32GM007276. S.H. and C.L.W. are supported by Stanford's Center for
Computational, Evolutionary, and Human Genomics. P.L.M.L. is supported by a Human Frontier
Science Program Long-Term Fellowship (LT000330/2019-L). M. R. is supported by the NSF's Plant

329   Genome Postdoctoral Research Fellowship in Biology. L. L. and L.G. are supported by NSF's GRFP.
330   Computational analyses were done on the High-Performance Computing clusters *Memex* and *Calc*
331   supported by the Carnegie Institution for Science.

REFERENCES

334   1.   IUCN, The IUCN Red List of Threatened Species. *https://www.iucnredlist.org*.

335   2.   S. Díaz, J. Settele, E. S. Brondízio, H. T. Ngo, J. Agard, A. Arneth, P. Balvanera, K. A.
336   Brauman, S. H. M. Butchart, K. M. A. Chan, L. A. Garibaldi, K. Ichii, J. Liu, S. M.
337   Subramanian, G. F. Midgley, P. Miloslavich, Z. Molnár, D. Obura, A. Pfaff, S. Polasky, A.
338   Purvis, J. Razzaque, B. Reyers, R. R. Chowdhury, Y.-J. Shin, I. Visseren-Hamakers, K. J. Willis,
339   C. N. Zayas, Pervasive human-driven decline of life on Earth points to the need for
340   transformative change. *Science*. **366**, eaax3100 (2019).

341   3.   IPBES, *Global Assessment Report on Biodiversity and Ecosystem Services* (IPBES Secretariat,
342   Bonn, Germany, 2019; https://www.ipbes.net/global-assessment-report-biodiversity-ecosystem-
343   services).

344   4.   J. J. Wiens, Climate-Related Local Extinctions Are Already Widespread among Plant and
345   Animal Species. *PLoS Biol.* **14**, e2001104 (2016).

346   5.   W. Thuiller, S. Lavorel, M. B. Araújo, M. T. Sykes, I. C. Prentice, Climate change threats to
347   plant diversity in Europe. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 8245–8250 (2005).

348   6.   M. Exposito-Alonso, F. Vasseur, W. Ding, G. Wang, H. A. Burbano, D. Weigel, Genomic basis
349   and evolutionary potential for extreme drought adaptation in Arabidopsis thaliana. *Nat Ecol*
350   *Evol*. **2**, 352–358 (2018).

351   7.   C. Parmesan, Ecological and Evolutionary Responses to Recent Climate Change. *Annu. Rev.*
352   *Ecol. Evol. Syst.* **37**, 637–669 (2006).

353   8.   T. Capblancq, M. C. Fitzpatrick, R. A. Bay, M. Exposito-Alonso, S. R. Keller, Genomic
354   Prediction of (Mal)Adaptation Across Current and Future Climatic Landscapes. *Annu. Rev. Ecol.*
355   *Evol. Syst.* **51**, 245–269 (2020).

356   9.   M. Lynch, J. Conery, R. Burger, Mutation Accumulation and the Extinction of Small
357   Populations. *Am. Nat.* **146**, 489–518 (1995).

358   10.   D. Spielman, B. W. Brook, R. Frankham, Most species are not driven to extinction before
359   genetic factors impact them. *Proceedings of the National Academy of Sciences*. **101** (2004), pp.
360   15261–15264.

361   11.   W. Steffen, K. Richardson, J. Rockström, S. E. Cornell, I. Fetzer, E. M. Bennett, R. Biggs, S. R.
362   Carpenter, W. de Vries, C. A. de Wit, C. Folke, D. Gerten, J. Heinke, G. M. Mace, L. M.
363   Persson, V. Ramanathan, B. Reyers, S. Sörlin, Planetary boundaries: Guiding human
364   development on a changing planet. *Science*. **347** (2015), doi:10.1126/science.1259855.

365   12.   S. Díaz, N. Zafra-Calvo, A. Purvis, P. H. Verburg, D. Obura, P. Leadley, R. Chaplin-Kramer, L.
366   De Meester, E. Dulloo, B. Martín-López, M. R. Shaw, P. Visconti, W. Broadgate, M. W.
367   Bruford, N. D. Burgess, J. Cavender-Bares, F. DeClerck, J. M. Fernández-Palacios, L. A.
368   Garibaldi, S. L. L. Hill, F. Isbell, C. K. Khoury, C. B. Krug, J. Liu, M. Maron, P. J. K.
369   McGowan, H. M. Pereira, V. Reyes-García, J. Rocha, C. Rondinini, L. Shannon, Y.-J. Shin, P.
370   V. R. Snelgrove, E. M. Spehn, B. Strassburg, S. M. Subramanian, J. J. Tewksbury, J. E. M.
371   Watson, A. E. Zanne, Set ambitious goals for biodiversity and sustainability. *Science*. **370**, 411–

413 (2020).

13. CBD, 1st Draft of The Post-2020 Global Biodiversity Framework (2021), (available at
https://www.cbd.int/doc/c/abb5/591f/2e46096d3f0330b08ce87a45/wg2020-03-03-en.pdf).

14. D. M. Leigh, A. P. Hendry, E. Vázquez-Domínguez, V. L. Friesen, Estimated six per cent loss of
genetic variation in wild populations since the industrial revolution. *Evol. Appl.* **12**, 1505–1512
(2019).

15. K. L. Millette, V. Fugère, C. Debyser, A. Greiner, F. J. J. Chain, A. Gonzalez, No consistent
effects of humans on animal genetic diversity worldwide. *Ecol. Lett.* **23**, 55–67 (2020).

16. I. G. Alsos, D. Ehrich, W. Thuiller, P. B. Eidesen, A. Tribsch, P. Schönswetter, C. Lagaye, P.
Taberlet, C. Brochmann, Genetic consequences of climate change for northern plants. *Proc. Biol.*
Sci. **279**, 2042–2051 (2012).

17. S. Theodoridis, C. Rahbek, D. Nogues-Bravo, Exposure of mammal genetic diversity to mid-21st
century global change. *Ecography (Cop.)* (2021), doi:10.1111/ecog.05588.

18. O. Arrhenius, Species and Area. *J. Ecol.* **9**, 95–99 (1921).

19. D. Storch, P. Keil, W. Jetz, Universal species-area and endemics-area relationships at continental
scales. *Nature*. **488**, 78–81 (2012).

20. C. D. Thomas, A. Cameron, R. E. Green, M. Bakkenes, L. J. Beaumont, Y. C. Collingham, B. F.
N. Erasmus, M. F. de Siqueira, A. Grainger, L. Hannah, L. Hughes, B. Huntley, A. S. van
Jaarsveld, G. F. Midgley, L. Miles, M. A. Ortega-Huerta, A. Townsend Peterson, O. L. Phillips,
S. E. Williams, Extinction risk from climate change. *Nature*. **427**, 145–148 (2004).

21. V. Buffalo, Quantifying the relationship between genetic diversity and population size suggests
natural selection cannot explain Lewontin's paradox. *Elife*. **10** (2021), doi:10.7554/eLife.67509.

22. R. A. Fisher, XVII.—The Distribution of Gene Ratios for Rare Mutations. *Proceedings of the*
Royal Society of Edinburgh. **50**, 204–219 (1931).

23. 1001 Genomes Consortium, 1,135 Genomes Reveal the Global Pattern of Polymorphism in
Arabidopsis thaliana. *Cell*. **166**, 481–491 (2016).

24. F. W. Preston, The canonical distribution of commonness and rarity: Part I. *Ecology*. **43**, 185
(1962).

25. A. Hampe, R. J. Petit, Conserving biodiversity under climate change: the rear edge matters. *Ecol.*
Lett. **8**, 461–467 (2005).

26. V. Buffalo, Why do species get a thin slice of π? Revisiting Lewontin's Paradox of Variation.
bioRxiv (2021), p. 2021.02.03.429633.

27. Millennium Ecosystem Assessment, *Millennium ecosystem assessment* (Millennium Ecosystem
Assessment, 2005; http://chapter.ser.org/europe/files/2012/08/Harris.pdf).

28. Ipbes, *Global assessment report of the Intergovernmental Science-Policy Platform on*
Biodiversity and Ecosystem Services (IPBES Secretariat, Bonn, Germany, 2019;
https://www.ipbes.net/news/ipbes-global-assessment-summary-policymakers-pdf).

29. G. C. Hurtt, L. Chini, R. Sahajpal, S. Frolking, B. L. Bodirsky, K. Calvin, J. C. Doelman, J. Fisk,
S. Fujimori, K. Klein Goldewijk, T. Hasegawa, P. Havlik, A. Heinimann, F. Humpenöder, J.
Jungclaus, J. O. Kaplan, J. Kennedy, T. Krisztin, D. Lawrence, P. Lawrence, L. Ma, O. Mertz, J.

Pongratz, A. Popp, B. Poulter, K. Riahi, E. Shevliakova, E. Stehfest, P. Thornton, F. N. Tubiello, D. P. van Vuuren, X. Zhang, Harmonization of global land use change and management for the period 850–2100 (LUH2) for CMIP6. *Geosci. Model Dev.* **13**, 5425–5464 (2020).

30. G. Gibson, Rare and common variants: twenty arguments. *Nat. Rev. Genet.* **13**, 135–145 (2011).

31. M. V. Rockman, The QTN program and the alleles that matter for evolution: all that's gold does not glitter. *Evolution*. **66**, 1–17 (2012).

32. M. J. Harms, J. W. Thornton, Evolutionary biochemistry: revealing the historical and physical causes of protein properties. *Nat. Rev. Genet.* **14**, 559–571 (2013).

33. M. Exposito-Alonso, 500 Genomes Field Experiment Team, H. A. Burbano, O. Bossdorf, R. Nielsen, D. Weigel, Natural selection in the *Arabidopsis thaliana* genome in present and future climates. *Nature*. **573**, 126–129 (2019).

34. H. R. Taft, D. A. Roff, Do bottlenecks increase additive genetic variance? *Conserv. Genet.* **13**, 333–342 (2012).

35. P. Ehrlich, B. Walker, Rivets and redundancy. *Bioscience*. **48**, 387 (1998).

36. C. C. Kyriazis, R. K. Wayne, K. E. Lohmueller, Strongly deleterious mutations are a primary determinant of extinction risk due to inbreeding depression. *Evolution Letters*. **n/a** (2020), doi:10.1002/evl3.209.

37. M. Kardos, E. Armstrong, S. W. Fitzpatrick, S. Hauser, P. Hedrick, J. Miller, D. Tallmon, W. Chris Funk, The crucial role of genome-wide genetic variation in conservation. *bioRxiv* (2021), p. 2021.07.05.451163.

38. R. Lande, Risks of Population Extinction from Demographic and Environmental Stochasticity and Random Catastrophes. *Am. Nat.* **142**, 911–927 (1993).

39. M. Lynch, R. Lande, The critical effective size for a genetically secure population. *Anim. Conserv.* **1**, 70–72 (1998).

SUPPLEMENTAL REFERENCES

1. S. P. Hubbell, *The Unified Neutral Theory of Biodiversity and Biogeography* (Monographs in Population Biology, 2001).

2. M. Tokeshi, Niche Apportionment or Random Assortment: Species Abundance Patterns Revisited. *J. Anim. Ecol.* **59**, 1129–1146 (1990).

3. M. Tokeshi, in *Advances in Ecological Research*, M. Begon, A. H. Fitter, Eds. (Academic Press, 1993; https://www.sciencedirect.com/science/article/pii/S0065250408600422), vol. 24, pp. 111–186.

4. MOTOMURA, I, A statistical treatment of ecological communities. *Zoological Magazine*. **44**, 379–383 (1932).

5. R. H. MacArthur, On the relative abundance of bird species. *Proc. Natl. Acad. Sci. U. S. A.* **43**, 293–295 (1957).

6.   R. A. Fisher, A. S. Corbet, C. B. Williams, The Relation Between the Number of Species and the
Number of Individuals in a Random Sample of an Animal Population. *J. Anim. Ecol.* **12**, 42–58
(1943).

7.   F. W. Preston, The canonical distribution of commonness and rarity: Part I. *Ecology*. **43**, 185
(1962).

8.   F. W. Preston, The commonness, and rarity, of species. *Ecology*. **29**, 254–283 (1948).

9.   D. Alonso, A. J. McKane, Sampling Hubbell's neutral theory of biodiversity. *Ecol. Lett.* **7**, 901–
910 (2004).

10. D. Storch, P. Keil, W. Jetz, Universal species-area and endemics-area relationships at continental
scales. *Nature*. **488**, 78–81 (2012).

11. S. L. Pimm, G. J. Russell, J. L. Gittleman, T. M. Brooks, The future of biodiversity. *Science*.
269, 347–350 (1995).

12. C. D. Thomas, A. Cameron, R. E. Green, M. Bakkenes, L. J. Beaumont, Y. C. Collingham, B. F.
N. Erasmus, M. F. de Siqueira, A. Grainger, L. Hannah, L. Hughes, B. Huntley, A. S. van
Jaarsveld, G. F. Midgley, L. Miles, M. A. Ortega-Huerta, A. Townsend Peterson, O. L. Phillips,
S. E. Williams, Extinction risk from climate change. *Nature*. **427**, 145–148 (2004).

13. J. F. C. Kingman, The coalescent. *Stochastic Process. Appl.* **13**, 235–248 (1982).

14. R. C. Griffiths, S. Tavaré, The age of a mutation in a general coalescent tree. *Communications in
Statistics. Stochastic Models*. **14**, 273–295 (1998).

15. R. A. Fisher, XVII.—The Distribution of Gene Ratios for Rare Mutations. *Proceedings of the
Royal Society of Edinburgh*. **50**, 204–219 (1931).

16. M. W. Hahn, *Molecular Population Genetics* (Oxford University Press, 2018;
https://play.google.com/store/books/details?id=3BDkswEACAAJ).

17. V. Buffalo, Quantifying the relationship between genetic diversity and population size suggests
natural selection cannot explain Lewontin's paradox. *Elife*. **10** (2021), doi:10.7554/eLife.67509.

18. D. R. Marshal, A. D. H. Brown, in *Crop genetic resources for today and tomorrow*, O. H.
Frankel, J. G. Hawkes, Eds. (Cambridge University Press, 1975;
https://www.researchgate.net/publication/280057199), vol. 2.

19. S. Wright, Isolation by Distance. *Genetics*. **28**, 114–138 (1943).

20. J. Novembre, M. Stephens, Interpreting principal component analyses of spatial population
genetic variation. *Nat. Genet.* **40**, 646–649 (2008).

21. H. Fan 樊海英, Q. Zhang 张清臣, J. Rao 饶娟娟, J. Cao 曹静文, X. Lu 卢欣, Genetic Diversity-
Area Relationships across Bird Species. *Am. Nat.* **194**, 736–740 (2019).

22. P. R. A. Campos, V. M. de Oliveira, A. Rosas, Epistasis and environmental heterogeneity in the
speciation process. *Ecol. Modell.* **221**, 2546–2554 (2010).

23.   J. Kelleher, A. M. Etheridge, G. McVean, Efficient Coalescent Simulation and Genealogical
485   Analysis for Large Sample Sizes. *PLoS Comput. Biol.* **12**, e1004842 (2016).

486   24.   B. C. Haller, P. W. Messer, SLiM 3: Forward Genetic Simulations Beyond the Wright–Fisher
487   Model. *Mol. Biol. Evol.* **36**, 632–637 (2019).

488   25.   T. R. Booker, S. Yeaman, M. C. Whitlock, The WZA: A window-based method for
489   characterizing genotype-environment association. *bioRxiv* (2021), p. 2021.06.25.449972.

490   26.   J. Kelleher, K. R. Thornton, J. Ashander, P. L. Ralph, Efficient pedigree recording for fast
491   population genetics simulation. *PLoS Comput. Biol.* (2018), p. e1006581.

492   27.   B. C. Haller, J. Galloway, J. Kelleher, P. W. Messer, P. L. Ralph, Tree-sequence recording in
493   SLiM opens new horizons for forward-time simulation of whole genomes. *Mol. Ecol. Resour.*
494   **19**, 552–566 (2019).

495   28.   R. R. Hudson, M. Slatkin, W. P. Maddison, Estimation of levels of gene flow from DNA
496   sequence data. *Genetics*. **132**, 583–589 (1992).

497   29.   1001 Genomes Consortium, 1,135 Genomes Reveal the Global Pattern of Polymorphism in
498   Arabidopsis thaliana. *Cell*. **166**, 481–491 (2016).

499   30.   B. J. McGill, B. J. Enquist, E. Weiher, M. Westoby, Rebuilding community ecology from
500   functional traits. *Trends Ecol. Evol.* **21**, 178–185 (2006).

501   31.   P. I. Prado, M. Miranda, A. Chalom, *sads: R package for fitting species abundance distributions*
502   (Github, 2018; https://github.com/piLaboratory/sads).

503   32.   T. J. Matthews, K. A. Triantis, R. J. Whittaker, F. Guilhaumon, sars: an R package for fitting,
504   evaluating and comparing species–area relationship models. *Ecography*. **42**, 1446–1455 (2019).

505   33.   F. He, S. P. Hubbell, Species-area relationships always overestimate extinction rates from habitat
506   loss. *Nature*. **473**, 368–371 (2011).

507   34.   R. J. H. &. J. van Etten, raster: Geographic analysis and modeling with raster data (2012),
508   (available at http://CRAN.R-project.org/package=raster).

509   35.   M. Exposito-Alonso, 500 Genomes Field Experiment Team, H. A. Burbano, O. Bossdorf, R.
510   Nielsen, D. Weigel, Natural selection in the *Arabidopsis thaliana* genome in present and future
511   climates. *Nature*. **573**, 126–129 (2019).

512   36.   Y. B. Simons, K. Bullaughey, R. R. Hudson, G. Sella, A population genetic interpretation of
513   GWAS findings for human quantitative traits. *PLoS Biol.* **16**, e2002985 (2018).

514   37.   M. Exposito-Alonso, C. Becker, V. J. Schuenemann, E. Reiter, C. Setzer, R. Slovak, B. Brachi, J.
515   Hagmann, D. G. Grimm, J. Chen, W. Busch, J. Bergelson, R. W. Ness, J. Krause, H. A. Burbano,
516   D. Weigel, The rate and potential relevance of new mutations in a colonizing plant lineage. *PLoS*
517   *Genet.* **14**, e1007155 (2018).

518   38.   H. Seebens, T. M. Blackburn, E. E. Dyer, P. Genovesi, P. E. Hulme, J. M. Jeschke, S. Pagad, P.
519   Pyšek, M. Winter, M. Arianoutsou, S. Bacher, B. Blasius, G. Brundu, C. Capinha, L. Celesti-

Grapow, W. Dawson, S. Dullinger, N. Fuentes, H. Jäger, J. Kartesz, M. Kenis, H. Kreft, I. Kühn,
B. Lenzner, A. Liebhold, A. Mosena, D. Moser, M. Nishino, D. Pearman, J. Pergl, W. Rabitsch,
J. Rojas-Sandoval, A. Roques, S. Rorke, S. Rossinelli, H. E. Roy, R. Scalera, S. Schindler, K.
Štajerová, B. Tokarska-Guzik, M. van Kleunen, K. Walker, P. Weigelt, T. Yamanaka, F. Essl,
No saturation in the accumulation of alien species worldwide. *Nat. Commun.* **8**, 14435 (2017).

39. H. Seebens, F. Essl, W. Dawson, N. Fuentes, D. Moser, J. Pergl, P. Pyšek, M. van Kleunen, E.
Weber, M. Winter, B. Blasius, Global trade will accelerate plant invasions in emerging
economies under climate change. *Glob. Chang. Biol.* **21**, 4128–4140 (2015).

40. S. Purcell, B. Neale, K. Todd-Brown, L. Thomas, M. A. R. Ferreira, D. Bender, J. Maller, P.
Sklar, P. I. W. de Bakker, M. J. Daly, P. C. Sham, PLINK: a tool set for whole-genome
association and population-based linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

41. C. Mérot, R. A. Oomen, A. Tigano, M. Wellenreuther, A Roadmap for Understanding the
Evolutionary Significance of Structural Genomic Variation. *Trends Ecol. Evol.* **35**, 561–572
(2020).

42. K. Lucek, Y. Willi, Drivers of linkage disequilibrium across a species' geographic range. *PLoS
Genet.* **17**, e1009477 (2021).

43. J. M. Kreiner, D. A. Giacomini, F. Bemm, B. Waithaka, J. Regalado, C. Lanz, J. Hildebrandt, P.
H. Sikkema, P. J. Tranel, D. Weigel, J. R. Stinchcombe, S. I. Wright, Multiple modes of
convergent adaptation in the spread of glyphosate-resistant Amaranthus tuberculatus. *Proc. Natl.
Acad. Sci. U. S. A.* **116**, 21076–21084 (2019).

44. M. A. Supple, J. G. Bragg, L. M. Broadhurst, A. B. Nicotra, M. Byrne, R. L. Andrew, A.
Widdup, N. C. Aitken, J. O. Borevitz, Landscape genomic prediction for restoration of a
Eucalyptus foundation species under climate change. *Elife.* **7** (2018), doi:10.7554/eLife.31835.

45. M. Vallejo-Marín, J. Friedman, A. D. Twyford, O. Lepais, S. M. Ickert-Bond, M. A. Streisfeld,
L. Yant, M. van Kleunen, M. C. Rotter, J. R. Puzey, Population genomic and historical analysis
suggests a global invasion by bridgehead processes in Mimulus guttatus. *Commun Biol.* **4**, 327
(2021).

46. J. T. Lovell, A. H. MacQueen, S. Mamidi, J. Bonnette, J. Jenkins, J. D. Napier, A. Sreedasyam,
A. Healey, A. Session, S. Shu, K. Barry, S. Bonos, L. Boston, C. Daum, S. Deshpande, A.
Ewing, P. P. Grabowski, T. Haque, M. Harrison, J. Jiang, D. Kudrna, A. Lipzen, T. H.
Pendergast 4th, C. Plott, P. Qi, C. A. Saski, E. V. Shakirov, D. Sims, M. Sharma, R. Sharma, A.
Stewart, V. R. Singan, Y. Tang, S. Thibivillier, J. Webber, X. Weng, M. Williams, G. A. Wu, Y.
Yoshinaga, M. Zane, L. Zhang, J. Zhang, K. D. Behrman, A. R. Boe, P. A. Fay, F. B. Fritschi, J.
D. Jastrow, J. Lloyd-Reilley, J. M. Martínez-Reyna, R. Matamala, R. B. Mitchell, F. M.
Rouquette Jr, P. Ronald, M. Saha, C. M. Tobias, M. Udvardi, R. A. Wing, Y. Wu, L. E. Bartley,
M. Casler, K. M. Devos, D. B. Lowry, D. S. Rokhsar, J. Grimwood, T. E. Juenger, J. Schmutz,
Genomic mechanisms of climate adaptation in polyploid bioenergy switchgrass. *Nature* (2021),
doi:10.1038/s41586-020-03127-1.

47. I. R. MacLachlan, T. K. McDonald, B. M. Lind, L. H. Rieseberg, S. Yeaman, S. N. Aitken,
Genome-wide shifts in climate-related variation underpin responses to selective breeding in a
widespread conifer. *Proc. Natl. Acad. Sci. U. S. A.* **118** (2021), doi:10.1073/pnas.2016900118.

48. G. Tuskan, W. Muchero, J.-G. Chen, D. Jacobson, T. Tschaplinski, D. Rokhsar, W. Schackwitz, J. Schmutz, S. DiFazio, Populus Trichocarpa Genome-Wide Association Study (GWAS) Population SNP Dataset Released, (available at https://doi.ccs.ornl.gov/ui/doi/55).

49. Anopheles gambiae 1000 Genomes Consortium, Data analysis group, Partner working group, Sample collections—Angola:, Burkina Faso:, Cameroon:, Gabon:, Guinea:, Guinea-Bissau:, Kenya:, Uganda:, Crosses:, Sequencing and data production, Web application development, Project coordination, Genetic diversity of the African malaria vector Anopheles gambiae. *Nature*. **552**, 96–100 (2017).

50. Z. L. Fuller, V. J. L. Mocellin, L. A. Morris, N. Cantin, J. Shepherd, L. Sarre, J. Peng, Y. Liao, J. Pickrell, P. Andolfatto, M. Matz, L. K. Bay, M. Przeworski, Population genetics of the coral Acropora millepora: Toward genomic prediction of bleaching. *Science*. **369** (2020), doi:10.1126/science.aba4674.

51. K. Ruegg, E. C. Anderson, M. Somveille, R. A. Bay, M. Whitfield, E. H. Paxton, T. B. Smith, Linking climate niches across seasons to assess population vulnerability in a migratory bird. *Glob. Chang. Biol.* **27**, 3519–3531 (2021).

52. R. A. Bay, R. J. Harrigan, V. Le Underwood, H. Lisle Gibbs, T. B. Smith, K. Ruegg, Genomic signals of selection predict climate-driven population declines in a migratory bird. *Science*. **359**, 83–86 (2018).

53. E. P. Kingsley, K. M. Kozak, S. P. Pfeifer, D.-S. Yang, H. E. Hoekstra, The ultimate and proximate mechanisms driving the evolution of long tails in forest deer mice. *Evolution*. **71**, 261–273 (2017).

54. L. Smeds, J. Aspi, J. Berglund, I. Kojola, K. Tirronen, H. Ellegren, Whole-genome analyses provide no evidence for dog introgression in Fennoscandian wolf populations. *Evol. Appl.* **14**, 721–734 (2021).

55. R. M. Schweizer, J. Robinson, R. Harrigan, P. Silva, M. Galverni, M. Musiani, R. E. Green, J. Novembre, R. K. Wayne, Targeted capture and resequencing of 1040 genes reveal environmentally driven functional variation in grey wolves. *Mol. Ecol.* **25**, 357–379 (2016).

56. 1000 Genomes Project Consortium, A. Auton, L. D. Brooks, R. M. Durbin, E. P. Garrison, H. M. Kang, J. O. Korbel, J. L. Marchini, S. McCarthy, G. A. McVean, G. R. Abecasis, A global reference for human genetic variation. *Nature*. **526**, 68–74 (2015).

57. J. D. Palacio-Mejía, P. P. Grabowski, E. M. Ortiz, G. A. Silva-Arias, T. Haque, D. L. Des Marais, J. Bonnette, D. B. Lowry, T. E. Juenger, Geographic patterns of genomic diversity and structure in the C4 grass Panicum hallii across its natural distribution. *AoB Plants*. **13**, lab002 (2021).

58. A. M. Royer, M. A. Streisfeld, C. I. Smith, Population genomics of divergence within an obligate pollination mutualism: Selection maintains differences between Joshua tree species. *Am. J. Bot.* **103**, 1730–1741 (2016).

59. M. Kapun, J. C. B. Nunez, M. Bogaerts-Márquez, J. Murga-Moreno, M. Paris, J. Outten, M. Coronado-Zamora, C. Tern, O. Rota-Stabelli, M. P. García Guerreiro, S. Casillas, D. J. Orengo,

E. Puerma, M. Kankare, L. Ometto, V. Loeschcke, B. S. Onder, J. K. Abbott, S. W. Schaeffer, S. Rajpurohit, E. L. Behrman, M. F. Schou, T. J. S. Merritt, B. P. Lazzaro, A. Glaser-Schmitt, E. Argyridou, F. Staubach, Y. Wang, E. Tauber, S. V. Serga, D. K. Fabian, K. A. Dyer, C. W. Wheat, J. Parsch, S. Grath, M. S. Veselinovic, M. Stamenkovic-Radak, M. Jelic, A. J. Buendía-Ruíz, M. Josefa Gómez-Julián, M. Luisa Espinosa-Jimenez, F. D. Gallardo-Jiménez, A. Patenkovic, K. Eric, M. Tanaskovic, A. Ullastres, L. Guio, M. Merenciano, S. Guirao-Rico, V. Horváth, D. J. Obbard, E. Pasyukova, V. E. Alatortsev, C. P. Vieira, J. Vieira, J. Roberto Torres, I. Kozeretska, O. M. Maistrenko, C. Montchamp-Moreau, D. V. Mukha, H. E. Machado, A. Barbadilla, D. Petrov, P. Schmidt, J. Gonzalez, T. Flatt, A. O. Bergland, Drosophila Evolution over Space and Time (DEST) - A New Population Genomics Resource. *bioRxiv* (2021), p. 2021.02.01.428994.

60. L. N. Di Santo, S. Hoban, T. L. Parchman, J. W. Wright, J. A. Hamilton, Reduced representation sequencing to understand the evolutionary history of Torrey pine (Pinus torreyana Parry) with implications for rare species conservation. *bioRxiv* (2021), p. 2021.07.02.450939.

61. J. von Seth, N. Dussex, D. Díez-Del-Molino, T. van der Valk, V. E. Kutschera, M. Kierczak, C. C. Steiner, S. Liu, M. T. P. Gilbert, M.-H. S. Sinding, S. Prost, K. Guschanski, S. K. S. S. Nathan, S. Brace, Y. L. Chan, C. W. Wheat, P. Skoglund, O. A. Ryder, B. Goossens, A. Götherström, L. Dalén, Genomic insights into the conservation status of the world's last remaining Sumatran rhinoceros populations. *Nat. Commun.* **12**, 2393 (2021).

62. Millennium Ecosystem Assessment, *Millennium ecosystem assessment* (Millennium Ecosystem Assessment, 2005; http://chapter.ser.org/europe/files/2012/08/Harris.pdf).

63. G. C. Hurtt, L. Chini, R. Sahajpal, S. Frolking, B. L. Bodirsky, K. Calvin, J. C. Doelman, J. Fisk, S. Fujimori, K. Klein Goldewijk, T. Hasegawa, P. Havlik, A. Heinimann, F. Humpenöder, J. Jungclaus, J. O. Kaplan, J. Kennedy, T. Krisztin, D. Lawrence, P. Lawrence, L. Ma, O. Mertz, J. Pongratz, A. Popp, B. Poulter, K. Riahi, E. Shevliakova, E. Stehfest, P. Thornton, F. N. Tubiello, D. P. van Vuuren, X. Zhang, Harmonization of global land use change and management for the period 850–2100 (LUH2) for CMIP6. *Geosci. Model Dev.* **13**, 5425–5464 (2020).

64. S. Theodoridis, C. Rahbek, D. Nogues-Bravo, Exposure of mammal genetic diversity to mid-21st century global change. *Ecography (Cop.)* (2021), doi:10.1111/ecog.05588.

65. I. Overcast, M. Ruffley, J. Rosindell, L. Harmon, P. A. V. Borges, B. C. Emerson, R. S. Etienne, R. Gillespie, H. Krehenwinkel, D. Luke Mahler, F. Massol, C. E. Parent, J. Patiño, B. Peter, B. Week, C. Wagner, M. J. Hickerson, A. Rominger, A unified model of species abundance, genetic diversity, and functional diversity reveals the mechanisms structuring ecological communities. *bioRxiv* (2020), p. 2020.01.30.927236.

66. S. Díaz, N. Zafra-Calvo, A. Purvis, P. H. Verburg, D. Obura, P. Leadley, R. Chaplin-Kramer, L. De Meester, E. Dulloo, B. Martín-López, M. R. Shaw, P. Visconti, W. Broadgate, M. W. Bruford, N. D. Burgess, J. Cavender-Bares, F. DeClerck, J. M. Fernández-Palacios, L. A. Garibaldi, S. L. L. Hill, F. Isbell, C. K. Khoury, C. B. Krug, J. Liu, M. Maron, P. J. K. McGowan, H. M. Pereira, V. Reyes-García, J. Rocha, C. Rondinini, L. Shannon, Y.-J. Shin, P. V. R. Snelgrove, E. M. Spehn, B. Strassburg, S. M. Subramanian, J. J. Tewksbury, J. E. M. Watson, A. E. Zanne, Set ambitious goals for biodiversity and sustainability. *Science*. **370**, 411–413 (2020).

642   67.   C. Rahbek, R. K. Colwell, Biodiversity: Species loss revisited. *Nature*. **473** (2011), pp. 288–289.

643   68.   H. M. Pereira, G. C. Daily, Modeling biodiversity dynamics in countryside landscapes. *Ecology*.
644   **87**, 1877–1885 (2006).

645   69.   J. Harte, A. B. Smith, D. Storch, Biodiversity scales from plots to biomes with a universal
646   species-area curve. *Ecol. Lett.* **12**, 789–797 (2009).

647   70.   D. R. Lockwood, C. M. Richards, G. M. Volk, Probabilistic models for collecting genetic
648   diversity: Comparisons, caveats, and limitations. *Crop Sci.* **47**, 861–866 (2007).

649   71.   L. Czech, M. Exposito-Alonso, grenepipe: A flexible, scalable, and reproducible pipeline to
650   automate variant and frequency calling from sequence reads. *arXiv* (2021), (available at
651   http://arxiv.org/abs/2103.15167).

652   72.   A. Miraldo, S. Li, M. K. Borregaard, A. Flórez-Rodríguez, S. Gopalakrishnan, M. Rizvanovic, Z.
653   Wang, C. Rahbek, K. A. Marske, D. Nogués-Bravo, An Anthropocene map of genetic diversity.
654   *Science*. **353**, 1532–1535 (2016).

655   73.   D. H. Parks, T. Mankowski, S. Zangooei, M. S. Porter, D. G. Armanini, D. J. Baird, M. G. I.
656   Langille, R. G. Beiko, GenGIS 2: Geospatial Analysis of Traditional and Genetic Biodiversity,
657   with New Gradient Algorithms and an Extensible Plugin Framework. *PLoS One*. **8**, e69885
658   (2013).

659   74.   C. Li, D. Tian, B. Tang, X. Liu, X. Teng, W. Zhao, Z. Zhang, S. Song, Genome Variation Map: a
660   worldwide collection of genome variations across multiple species. *Nucleic Acids Res.* **49**,
661   D1186–D1191 (2021).

662   75.   M. Kardos, E. Armstrong, S. W. Fitzpatrick, S. Hauser, P. Hedrick, J. Miller, D. Tallmon, W.
663   Chris Funk, The crucial role of genome-wide genetic variation in conservation. *bioRxiv* (2021),
664   p. 2021.07.05.451163.

665   76.   D. Spielman, B. W. Brook, R. Frankham, Most species are not driven to extinction before
666   genetic factors impact them. *Proceedings of the National Academy of Sciences*. **101** (2004), pp.
667   15261–15264.

668   77.   Ipbes, *Global assessment report of the Intergovernmental Science-Policy Platform on
669   Biodiversity and Ecosystem Services* (IPBES Secretariat, Bonn, Germany, 2019;
670   https://www.ipbes.net/news/ipbes-global-assessment-summary-policymakers-pdf).

Genetic diversity loss in the Anthropocene

Moises Exposito-Alonso, Tom R. Booker, Lucas Czech, Tadashi Fukami, Lauren Gillespie, Shannon Hateley, Christopher C. Kyriazis, Patricia L. M. Lang, Laura Leventhal, David Nogues-Bravo, Veronica Pagowski, Megan Ruffley, Jeffrey P. Spence, Sebastian E. Toro Arana, Clemens L. Weiß, Erin Zess.

Correspondence to: moisesexpositoalonso@gmail.com

This PDF file includes:

Supplementary Methods
Supplementary Results
Figs. S1 to S27
Tables S1 to S18

Table of contents

120 ## SUPPLEMENTAL METHODS
121
122 ## I. Background on species biodiversity and biogeography
123 ### I.1 Theoretical models of biodiversity
124
125 Studies in biogeography have modelled the species-area relationship with several functions.
126 Below we summarise the different approaches using an example of richness of S = 100
127 species, with variable abundance or area, A.
128
We may visualise the different areas or abundances of species as a frequency
histogram (Fig. S1, Preston plot), with x-axis: logarithm of abundance bins (historically log2
as a rough approximation to the natural logarithm), and y-axis: number of species at given
abundance. Alternatively, as a rank-abundance diagram (Fig. S1, Whittaker plot): x-axis:
species list, ranked in order of descending abundance (i.e. from common to rare), and y-axis:
logarithm of % relative abundance.
135
136

138 ***Fig. S1 | Example of typical plots used for species abundance curve studies***
139 *Due to their strong skew, Species Abundance Curves are often plotted using the Preston plot (left) where the x axis*
140 *represents bins of log2 abundances (also referred to as octaves), or using the Whittaker plot (right) where the x axis is the*
141 *rank of each species in a dataset and y axis the species' relative abundance.*
142
143
144
145 ### I.1.2. Niche apportionment approaches
146
147 A series of theoretical deterministic and stochastic "niche apportionment models" have been
148 put forward (summarised in (*1*) or (*2, 3*)).
149
150 The Motomura (*4*) geometric series suggests that each species that arrives takes half
151 the area. The first would take 50%, the second 50% of 50%, and so forth, which can be
152 expressed as:
153
154 $P_i = 0.5^i$.
155
156 Similarly, one can imagine that as a species colonises a habitat, it takes up a fraction different
157 than 50%. This gives a geometric series with parameters k which can be written as
158

$P_i = k(1 - k)^{i-1}$.

Other geometric series-related models include stochasticity, where k instead of being a fixed
parameter is a random uniform variable and there is a k_i each time i a new species arrives to
the ecosystem. The "dominance preemption" model draws from 50-100% at any new arrival
of a species, the random fraction model draws from 0-100%. Then the abundance of a species
depends on the stochastic process of previous $f = 1...i-1$ species arriving first:

$$E[P_i | P_1, ..., P_{i-1}, k_i] = k_i \times \left(1 - \sum_{f=1}^{i-1} P_f \right)$$

167
168

169   Another approach is the broken stick by MacArthur (*5*), which theorised a habitat is broken
170   into *S-1* places at random, which creates S fractions of an area. Then the relative area of a
171   species is:

$$E[P_i] = (\frac{1}{S}) \sum_{w=i}^{S} \frac{1}{w}.$$

173
174
175
176   **I.1.2. Niche statistical approaches of species sampling**

178   Differently from niche partitioning functions, statistical approaches such as the log-series
179   from Fisher (*6*) and log-normal from Preston (*7*) are probability distributions, and approach
180   modelling in a conceptually different way: they model the sampling process of species
181   collections given an underlying relative abundance (see below).

182
183

Statistical-based derivations probably began with Fisher (*6*), with the log-series
distribution. It assumes that species abundances in the community are independent identically
distributed variables, sampling is a Poisson process, sampling is done with replacement, or
the fraction sampled is small enough to approximate a sample with replacement. Here,

$S_n = \frac{\alpha x^n}{n}$,

where x is a constant $x \in [0, 1]$ related to the sample dataset (typically close to 1),
$x = \frac{N}{\alpha + N}$, and α is a new constant term (ecosystem-specific) that is used as a measure of
biodiversity. Fisher proposed the number of species could be estimated as:

$S = \alpha \times log\left(1 + \frac{N}{\alpha}\right)$.

Finally, Preston (*8*) posed that the skewness of previous proposals is due to lack of
sampling. With little data, common species are collected sooner, but with more abundant
sampling, the rarest species are also well-sampled and have abundances well above 0. Preston
then proposed that the octaves (bins of doubling abundance) follow a normal distribution,
making the raw abundance log-normal distributed. Given S_0 is the number of species in the
model octave of abundance and a variance composite of the log-Normal σ^2, the number of
species per abundance (octave) bin R (=*log(n)*) is:

$S_R = S_0 e^{-R^2/2\sigma^2}.$
206
The Unified Neutral Theory of Biodiversity (UNTB) by Hubbell (*1*) takes a stochastic
approach of a community with immigrants, extinctions, and speciation in continuous
dynamics. Interestingly, the UNTB's key parameter, *θ*, coincides with Fisher's *α*, as the log-
series is a limiting case of UNTB. Hubbell's discovery was that *α=2J$_m$v*, where J_m is the size
of the external metacommunity that provides migrants of species to the focal community, and
v is the speciation rate. Alonso and McKane (*9*) derived the so-called Metacommunity Zero-
Sum Multinomial (MZSM) distribution from the UNTB. In practice, both distributions have
almost-identical fits (lines completely overlapping in Fig. S2 below).
215
216

217
218   *Fig. S2 | **Summary of theoretical models of Species Abundance Curves.***
219   *Five niche partitioning or statistical models shown in a Whittaker plot. The different models expect different levels of*
220   *evenness in abundance across the species in the community, from the lowest (geometric series) to the highest (log-normal).*
221
222

223   **I.2 Metric of species diversity**
224
225   Although a number of metrics exist to measure species diversity, such as the Shannon index,
226     $H' = -\sum_{i=1}^{S} P_i log P_i$ (with *Pi* the relative proportions of species abundances) or Fisher's non-
227   dimensional *α* parameter, the study of species abundances and area relationships has focused
228   on species richness *S*, that is, the total number of species in a given location or area. Below
229   we therefore focus on species richness.
230
231
232   **I.3 Biogeography of species and extinction.**
233
234   **SAD and SAR connection**
235
236   Due to many species being rare, it is expected that as researchers sample an area, the most
237   common species will be sampled first, and as the area studied increases, more and more
238   species will be discovered. This is thought to happen following a power law relationship,

239 where the number of species in that area S_A increases with the sampled area A, with scaling z
240 (slope in a log-log plot), and with a constant c:
241
242 $SAR(A) = cA^z$.
243
244   Preston (*7*) derived theoretically that from a log-normal series, one would expect
245 $z=0.27$, under a number of assumptions (Fig. S3). This has been empirically shown to be
246 close to reality (*7*, *10*), although there is some variation across ecosystems and spatial scales.
247

248
249 ***Fig. S3 | Example of a Species-Area Relationship in Galapagos Islands***
250 *Classic species richness dataset from the Galapagos Islands (Preston, 1962). It depicts species richness as a function of*
251 *island area in a log-log plot.*
252
253

254 **I.4 Estimating extinction of species from the species area relationship**
255
256 The first estimates of species extinction used the SAR relationship. Given a reduction of
257 ecosystem area, A, by an area of a (*11*, *12*). If these areas, as well as the SAR scaling, z, are
258 known, then one can predict the number of species in the future as:
259
260 $S_{\mathrm{now}} - S_{\mathrm{fut}} = cA^z_{\mathrm{now}} - cA^z_{\mathrm{fut}}$,
261
262 However, we are normally interested in the fraction of species that will go extinct X_s so we
263 can take the ratio:
264
265 $X_s = \frac{S_{\mathrm{now}} - S_{\mathrm{fut}}}{S_{\mathrm{now}}} = 1 - \frac{cA^z_{\mathrm{fut}}}{cA^z_{\mathrm{now}}} = 1 - \left(\frac{A_{\mathrm{fut}}}{A_{\mathrm{now}}}\right)^z$.
266
267

268 **II. Population genetics models and the site frequency spectrum.**
269 **II.1 The Wright-Fisher model and the site frequency spectrum**
270
Statisticians and population geneticists from the 20[th] century, Wright and Fisher, built a
simple statistical model of evolution of a population. It assumes that each generation a
population of N monoecious (hermaphrodite) individuals mate randomly to create a new
generation of N individuals and then immediately die so that only N individuals remain in the
population at any given time. This random sampling process causes the frequency of a variant

in one generation to possibly differ from its frequency in the previous generation—a process
known as genetic drift.
278
When a nucleotide mutation or variant (e.g. ACG**A**A → ACG**T**A) emerges by a
random process of, for instance, DNA replication error, it will first be in $1/N$ individuals (if
we consider these diploid, 1/2N chromosomes). Through random sampling that **T** mutation
may be lost, stay at the same frequency, or randomly move to higher frequency. Although
rarely, just by chance, the mutation may reach 100% frequency. This results in a
"commonness of rarity" when looking at mutations in a population, as we have seen in
previous sections for species. Since these genetic drift dynamics affect all mutations genome-
wide, we therefore expect the majority of mutations to be absent, or rare, and only a much
smaller proportion of variants to be at moderate or high frequencies.
288
The site frequency spectrum (SFS) refers to the distribution of frequencies of variants
in a population. This is the number of sites at which we observe a variant at frequency q in a
sample of n individuals. To derive the expected SFS distribution, we turn to Kingman's
Coalescent (*13*). Both models describe the same ideal population of random mating, constant
population size, and mutations emerging at a low rate and drifting in frequency. But while the
Wright-Fisher model describes the dynamics of a whole population forward-in-time, the
Kingman's Coalescent describes the genealogy of a sample of individuals from a population,
going backward in time. By building a model around the individuals that are sampled or that
survived, rather than of an entire population, the Coalescent provides a simpler way to derive
expectations in small populations or in cases, for example here, where a limited sample of
genomes are sequenced. Using the Coalescent (see (*14*) for details), one obtains that the
expected number of mutations of a given abundance, n, is inversely related to their frequency,
q:
302
303 $M_n = c\frac{1}{q}$
304
305 for some constant c that depends on the mutation rate and the population size. This SFS from
306 population genetics theory is remarkably similar to the Species Abundance Relationship. In
307 fact, Fisher himself (*15*) derived an expression similar to the above.
308
309   Rearranging terms, one can see this is a constrained version of the log-series
310 Probability Mass Function (PMF), which Fisher also proposed for the distribution of species
311 abundances (*6*). Below, one can graphically see the similarities (Fig. S4):
312

314 ***Fig. S4 | Similarity between the Species Abundance Distribution and the Site Frequency Spectrum***
315 *Left is the Probability Mass Function of the log-series (p=0.999), center is the SFS (N=100, c=1), and right is*
316 *the log-series-based abundance of species (alpha=100, N=10000).*
317
318   Keeping the abundance, n, constant (and low), when the number of individuals
319 $N \to \infty$, we know that the constant x from Fisher's SAD approaches 1, $x = \frac{N}{N+\alpha} \to 1$. Then,
320 we can rewrite the number of species at any given abundance (S_n) as:
321
322 $$S_n = \alpha \frac{(\frac{N}{\alpha+N})^n}{n} = \alpha \frac{1^n}{n} = c\frac{1}{n} = M_n$$
323
324   So both have the same form as the log series PMF: $f(k) = \frac{-1}{ln(1-p)} \frac{p^k}{k}$ when $p \to 1$. In
325 the next section we will see that the constants of the SAD and the SFS are proportional to
326 species and mutation diversity, although the Site Frequency Spectrum (SFS) is a specific case
327 of SAD. One can also see that because the constant in the SFS is the population scaled
328 mutation rate, $c = \theta = Ne\mu$, and Fisher's $\alpha \approx \theta$ for large N.
329
330 **II.2 Metrics of genetic diversity**
331
In population genetics, multiple measurements of genetic diversity have been put forward.
The most straightforward is the allelic richness, also number of mutations, or also called the
number of segregating sites. Segregating sites, *M*, is the direct equivalent of the species
richness*, S*, and it depends on the number of samples used and length of DNA sequence
explored (Note: we use the non-standard notation, *M*, as the standard in population genetics is
S [for segregating sites] but this is already in use for species richness. We then use *M* for
mutations and *S* for species). This metric can also be thought of as the area under the curve of
the SFS. Two other metrics that describe the SFS but that aim to be sequence-length- and
individual independent are Watterson's Theta, θ_W, and Nucleotide diversity, π, (also called θ_π
). These two metrics of diversity are identical at population equilibrium and are estimates of
4N$_e$μ (when the SFS follows a 1/q relationship), with effective population size N_e and per-
generation mutation rate *μ*, whereas they differ in non-equilibrium demographics, under
natural selection, or under other behaviors not considered in the Wright-Fisher neutral model,
such as different mating systems (*16*).
346
347 First, π is described as:
348
349 $$\pi = \frac{\sum_{i=1}^{n-1} i(n-i)M_i}{n(n-1)/2},$$
350
351 and θ_W as:
352
353 $$\theta_W = \frac{\sum_{i=1}^{n-1} M_i}{\sum_{i=1}^{n-1} 1/i},$$
354
355   where $\sum_{i=1}^{n-1} 1/i$ is the *n-1th* Harmonic number, which serves to scale the segregating
356 sites based on the assumption that the abundance of mutations follows a 1/q SFS. The
357 diversity metrics π and θ_W are both functions of the SFS, as opposed to Fisher's α from the
358 Species Abundance Distribution, which is a parameter that changes the shape of the
359 distribution.
360

361   Although often nucleotide diversity π is reported as a typical measure of genetic
362   diversity of a species, since it can be calculated for a single genome and it captures the
363   process of inbreeding of a population (*17*), classic literature relating germplasm management
364   for conservation and breeding has advocated for allelic richness (*18*).
365
366
367   **II.3 Spatial genetics and the mutations-area relationship (MAR)**
368
Since its inception, a number of concepts in population genetics have dealt with genetic
variation in populations of different sizes, or populations separated in space. For instance, one
classic result in population genetics is the relationship of $\pi \approx 4N_c\mu,$ which relates genetic
diversity π with the effective population size N_e and the mutation rate of the species μ. A
relationship which is still studied nowadays in an effort to reconcile data with theory (*17*).
In 1943, Sewall Wright turned to study the genetics of multiple populations within a
species. He proposed that populations sampled further apart geographically must differ more
in allele frequency due to more independent drift (*19*), leading to the commonly used
correlation between geographic distance and the metric of differentiation F_{ST}. Most
prominently, the use of correlation in the accumulation of mutations of populations that are
geographically close or share evolutionary history has been uncovered using dimensionality
reduction approaches such as PCA (*20*).
Despite these enormous advances in understanding spatial genetic structures,
surprisingly little quantitative work has been done to parametrize the loss of genetic diversity
by direct loss of habitat.
Because of the abundance of rare mutations in populations, it is straightforward to
think that the more area and individuals sampled, the more segregating sites will be found.
Analogous to the Species Area Relationship (SAR), $S=cA^z$, we should thus be able to
estimate the equivalent scaling for a mutations-area relationship (MAR):
388
389        $M=cA^z,$
390
391   with a scaling $z = z_{MAR}$, which corresponds to the slope of best fit in a log-log-plot of
392   *A* and *M* for a given species. (Other functions are often fit empirically for SAR datasets,
393   which we explore later in section III.3. We work with the power law because of its historical
394   use, mathematical convenience, and because other more complicated functions only
395   improved fitting marginally, see Table S4).
396
397   This differs from other efforts to understand the number of segregating sites or
398   heterozygosity differences across species that differ in their total census size or geographic
399   distribution (*21, 22*). The MAR instead is built within a species, as its ultimate aim is to relate
400   the number of mutations left in a species as it loses spatial populations.
401
402   Below we derive what are the expectations of MAR taking two opposite scenarios of
403   neutral population evolution, and study how many segregating sites or mutations *M* are
404   discovered with increasing area in the simulations. We further test the scenario of meta-
405   populations in space with varying migration rates and neutral or natural selection processes.
406
407   **II.3.1 Panmictic population**
408
409   The expected number of mutations, *M*, is a constant that depends on the mutation rate, μ, and
410   the expected total branch length of the population genealogy, *L*, with $M=\mu L$. Under the

411   coalescent, the total branch length is equal to the number of lineages or individuals sampled
412   from the population, *n*, times the time of the genealogy during which there are such lineages,
413   T_n, plus *n-1* times the time in the genealogy with such number of lineages, and so forth:
414
415   $$L = nT_n + (n-1)T_{n-1} + ... + 2T_2.$$
416
417   Under the coalescent,
418
419   $$E[T_n] = \frac{2Ne}{n(n-1)},$$
420
421   and thus:
422
423   $$E[L] = n\frac{2N_e}{n(n-1)} + (n-1)\frac{2N_e}{(n-1)(n-2)} + ...,$$
424
425   which simplifies to
426
427   $$E[L] = 2N_e(\frac{1}{n-1} + \frac{1}{n-2} + ... + 1) = 2N_e H_{n-1},$$
428
where H_{n-1} is the (n-1)th harmonic number. This is of course related to one of the
diversity metrics (section II.2), where Watterson's Θ_W scales the number of segregating sites
(*M*) by the harmonic number of sampled individuals. This is based on the expectation that as
more individuals are sampled, we expect to discover more mutations proportional to the
above harmonic number. Because such number is not so easy to work with to create an
expectation for z_{MAR}, we further simplify this expectation following the Taylor expansion
approximation of the harmonic number:
436
437   $$H_n = \gamma + \log(n) + \frac{1}{2n} + O(\frac{1}{n^2}) \simeq \gamma + \log(n) + \frac{1}{2n},$$
438
439   which we can further approximate as:
440
441   $$E[L] \approx 2N_e \log(n-1) + c.$$
442
Therefore, assuming a constant mutation rate and effective population size (N_e) under
panmixia, *M* grows following *log(n)*. In such a case, a log-log plot (typical power law plot)
does not display a linear relationship, and the slope is asymptotic to *z → 0* for *N → ∞*. On the
other hand, with low values of *x* (area or individuals sampled close to 0), the slope z_{MAR} will
be incorrectly high. We can show this effect trivially by studying the local derivative of the
function $log_{10}(M) = log_{10}(log(N))$. The local slope of that function is an approximation of our
z_{MAR} parameter. This can be locally estimated at any given point *N* by taking the derivative:
450
451   $$\frac{d \log_{10}(\log(N))}{d(\log_{10}(N))} = \frac{1}{\log_{10}(N)\log(10)}.$$
452
The implication of this nonlinear function is that if we sampled only few individuals
or areas of a species (e.g., n=100), even if this species was completely panmictic we would
expect a non-zero z_{MAR}, a value that will change with sampling effort. We can roughly approximate
z_{MAR} by the local slope of the number in the midpoint of the graph, e.g., for n=100 we look at
the slope at *n=50*, and obtain *1/(log_{10}(50) x log(10))≅0.256*. Therefore, with small sample
sizes, this parameter will not be helpful to understand whether a species behaves

panmictically or is limited by migration, which may be problematic for estimates of genetic
diversity loss later. We can visualise our expectation of the z_{MAR} under panmixia plotting the
first derivative above (Fig. S5). Because—as we will show below—we do expect a power
law relationship under a migration-limited scenario, z_{MAR} should theoretically not change with
sample size. The graphical study of the (non-)linearity of the log-log plots between the
number of mutations and area sampled should be diagnostic to this problem (We see for
instance that *Pinus contorta* has a highly nonlinear relationship, likely due to the use of
ascertained intermediate frequency markers instead of genome-wide data, Fig. S22).
467
Finally, we used msprime (*23*) to corroborate this finding (z_{MAR} being constant with
respect to sample size) with simulations, simulating 1600 demes in a 40x40 grid of demes or
populations of $N=N_e=1000$ that are completely panmictic (universal gene flow or dispersal,
so this is equivalent to a single panmictic deme). We observed the z_{MAR} for $t=100...10,000$
generations in log_{10} increments. After this time, we sample $n=1...100$ individuals in
increasingly large groups of adjacent demes. The range of estimates of z_{MAR} in these
simulations was 0.07-0.15.
475
476   Fig. S5 indicates that the minimum average z_{MAR} even under panmixia would
477 continuously increase with lower numbers of individuals of a species sampled. This is due to
478 the fact that the site frequency spectrum is not fully sampled with small numbers of
479 individuals. Therefore, we devised an approach to rescale z_{MAR}.
480
481
482

484   **_Fig. S5 | Expected ranges of z_{MAR} given sample sizes._**
485   *For increasing numbers of individuals sampled, we plot the expected mean z_{MAR} under two theoretical trends of a migration-*
486   *limited (green) and a panmictic (purple) species (Purple dots indicate averages from SLiM simulations under panmixia to*
487   *confirm the theoretical trend based on the derivative approach above). In black, z_{MAR} and 95% Confidence Interval of*
488   *species analyzed in section IV are plotted (see section for details).*
489
490
491   **II.3.2 Scaling z_{MAR} for low sampling and low census size**
492
493   Let $z_{pan-n} = E[z_{MAR} \mid n, panmixia]$, be the expected value of z_{MAR} of a panmictic
494   species given that we only have small sampling of n. Although theoretically z_{MAR} should
495   approach 0, with small samples it can be upwardly biased. In order to force the possible
496   values of z_{MAR} to range 0-1 despite small sample sizes, we can scale it as:
497
498           $z_{naive\ scaled} = (z_{MAR} - z_{pan-n}) / (1-z_{pan-n})$.
499
500   In words, this moves the purple line in Fig. S5 to zero, stretching the space above it
501   accordingly.
502
503   Most species have census sizes so large that z_{MAR} should indeed approach 0 under
panmixia, so we should correct the sample estimate z_{MAR} to range 0-1. However, some
species have such low census size N that even if we sample all individuals of a species, the
sample size will still be small. In those cases, we should not scale z_{MAR} to range 0-1, but
rather scale it from $z_{pan-N} - 1$, where $z_{pan-N} = E[z_{MAR} \mid N, panmixia]$ is the expected value of
z_{MAR} given a census size N (plants or animals living in the wild). The updated scaling
approach for both census and sample size would then be:
510
511           $z^*_{scaled} = (1-z_{pan-N}) (z_{MAR} - z_{pan-n}) / (1-z_{pan-n}) + z_{pan-N}$.
512
513
514   Note that this scaled estimate must be conservative because while we adjust the
minimum z for the average value expected for low sample sizes, we do not adjust for the
maximum possible z, which only under very extraordinary theoretical conditions can be $z=1$,
namely under an unrealistic complete disconnection of populations by gene flow (see below).
Because deriving the maximum z would require more biological knowledge of the species'
demography, landscape connectivity, genome structure, etc., and because we rather create
conservative estimates, we do not create further scaling approaches.
521
522   **II.3.3 Meta-populations in space**
523
A more realistic simulation than a panmictic population is that of the same 40x40 deme grid
where migration can happen between adjacent demes. This migration rate can be changed to
understand the effect of population structure and migration on z_{MAR}. Under no migration (or
very low migration), we expect the mutations in two distinct populations (and thus their SFS)
to be (almost) completely independent. Hence, when explored demes are doubled (N_e
doubles), we discover twice as many mutations. In this case, the number of mutations should
scale linearly with the area, so we expect the following to be true: $M=A$, $log(M) = log(A)$,
and $z_{MAR}=1$. Our analyses under different sampling schemes, and with different numbers of
"burn-in generations" (generations since a single deme colonised the full 40x40 space)
confirm that z_{MAR} approaches 1 in the limit of low migration (see Table S1 and Fig. S6).
Different from the panmictic situation, as we increase the sampled area, we not only increase
n, which would lead to a $log(A)$ in mutations, but also increase N_e.

536
537
538

539
540   ***Fig. S6 | msprime 2D deme simulations and the mutations-area relationship***
541   *Simulations with different burn-in and migration rates under neutrality, and their corresponding zmar.*
542

543   ***Table S1 | msprime population genetic simulations in 2D***
544   *Simulations summarised by grouping ranges of the resulting z_{MAR} parameters. The average parameters of the simulations*
545   *with similar z_{MAR} EW provided. (Acronyms: Nemt = product of effective population size, migration rate, and simulated*
546   *generations).*
547

z_{MAR}	Samples/deme	Generations	Migration rate	$N_e$mt
0.2 +/- 0.05	2.4	50001.7	0.0271675	5000044.23
0.3 +/- 0.05	20.25	70003	0.0561655	7000075.77
0.4 +/- 0.05	26.5714286	13057.4286	0.04450857	1305497.96
0.5 +/- 0.05	12.9230769	121759.462	0.04017769	752221.743
0.6 +/- 0.05	15.6111111	3218.77778	0.045735	321174.768
0.7 +/- 0.05	35.6842105	35034.8421	0.03395895	143791.614
0.8 +/- 0.05	35.030303	15655.1212	0.03055818	58023.5539
0.9 +/- 0.05	36.5806452	3057.12903	0.0253029	15290.4081
1 +/- 0.05	42.0140845	13625.4085	0.00861178	1798.36141

548
549
These simulations corroborated that we can recover z_{MAR} values ranging between 0-1
just varying migration and burn-in generation parameters. We found that it was both the time
of the system to reach an equilibrium as well as the migration rate that determined z_{MAR}. In
the future, it will be interesting to study different non-equilibrium scenarios to better
understand how genetic drift, gene flow, and different landscape structures may shape the
z_{MAR} .
556
557   **II.3.4 Metapopulations in space with local adaptation**
558
In order to simulate local adaptation, we use the individual-based simulation software SLiM
(*24*) following the approach of (*25*). These simulations were set up for 196 demes arranged in
a 14 x 14 grid. Each grid cell contains a population of *N=1000* and has an environment
attribute, *e*, which varied spatially from the lower-left to the upper-right corners (approx. *-7 <*
e < 7). 12 locations in the genome were allowed to be under directional natural selection. The
selection coefficient was fixed for a simulation, and grid runs were conducted with
0<s<0.05, but this selection would vary based on the environmental selection value of a grid
cell, according to *e × s*. Therefore, these alleles are antagonistic pleiotropic. Selected
mutations across the 12 loci in the genome behaved additively (e.g. if an individual in grid

cell i had two of the selected mutations, fitness would be $w=1+2s \times e_i$). The migration rate
varied from one individual in a billion (1×10^{-9}), to one individual every ten (1×10^{-1}). Finally,
the mutation rate was set to 10^{-8} mutations/bp/generation and the recombination rate to 10^{-7}
crossovers/bp/generation.

572
573
574

Fig. S7 | SLiM population genetic simulations in 2D with selection and local adaptation
576   *Simulations were carried out with different combinations of migration rates and strength of antagonistic*
577   *pleiotropic selection at 12 QTLs. (A) Marginal relationship between z_{MAR} with the strength of spatially-varying*
578   *selection s. (B) Marginal relationship between z_{MAR} with the migration rate m.*
579

580
These results, together with individual-based simulations, corroborate what we had
observed with coalescent simulations, i.e. that z_{MAR} is lowest with a high migration rate. The
simulations also appear to show a negative effect of selection on z_{MAR} . Generating a linear
model fitting migration rate and selection and their interaction to understand what factors
explain the scaling coefficient: $z_{MAR} \sim log_{10}(m) + s + log_{10}(m) s$; we confirm that both had a
significant effect, and that selection significantly reduces z_{MAR} (Fig. S7, see below summary
Table S2). This may seem counterintuitive, as one may expect that locally-adaptive mutations
are rare and will be localised only to where they are adaptive. More work is necessary to
understand the signatures that spatially-varying natural selection (and its different types)
create on z_{MAR}, but we can think that under migration limited scenarios (where z approaches
1) adaptive alleles and their linked mutations permeate faster to similar neighbour
environments than neutral alleles.
593

594   **Table S2 | Linear model explaining z_{MAR} by migration rate and natural selection**
595   *Summary table of the linear model $z_{MAR} \sim mig + s + mig:s$*

	Estimate	**SE**	**t-value**	**P-value**
intercept	0.3385022	0.0469174	7.214859	0.0000001
mig	-0.0419733	0.0085804	-4.891792	0.0000407
s	-4.693492	1.6290184	-2.881178	0.0076725
mig : s	-0.4998393	0.2426463	-2.059950	0.0491621

596
597
598   **II.3.5. Metapopulations in space with purifying selection**
599
To understand the effect of purifying selection on z_{MAR} we also ran 2D simulations with a
fraction of the genome allowed to be globally-deleterious (i.e. independent of the spatially-
varying environment). We simulated an increasingly strong purifying selection ($|s|$ range
from 0.0 to 0.1), simulating roughly that 29% of the genome of Arabidopsis is coding
(arabidopsis.org) and mutations can be deleterious. We also varied the degree of
recombination. Following our expectation, with stronger purifying selection deleterious

mutations are pushed to lower allele frequencies, stopping their geographic spread, which
increases z_{MAR}. Recombination rate appears to have a minor role on z_{MAR} (Fig. S8).
608
609

610
611 ***Fig. S8 | SLiM population genetic simulations in 2D with purifying selection***
612 *Simulations were carried out with varying strengths of purifying selection (|s| range from 0.0 to 0.1) at coding positions,*
613 *representing about 29% of the genome. Different values of recombination rate were also used in all pairwise combinations*
614 *with |s|.*
615
616 **II.3.6 Continuous-space non-Wright-Fisher models**
617
In order to confirm z_{MAR} generality in highly realistic conditions and its behavior through the
population extinction process (II.4), we set up SLiM simulations in continuous space using
non-Wright-Fisher dynamics (*24*). Spatial population structure in these simulations was
established through individual dispersal, local mate choice and spatial competition, which we
chose to lead to realistic values of F_{ST} across space. Spatial competition also acted as
population control, by keeping the total population size below a target carrying capacity
through direct effects on individual fitness. In addition to competition, fitness was also
affected by individual age as well as by a polygenic trait under stabilising selection. A subset
of variants (final proportion ~10%) directly affected this trait with effect sizes drawn from a
Gaussian distribution with mean = 0.0 and standard deviation = 0.1, and a fitness penalty was
incurred by deviating from the optimal trait value using a Gaussian fitness function centered
at the optimum and with a standard deviation = 5.0. We initialised functional variation for
SLiM using neutral coalescent simulations with msprime (*23*) to reduce the computational
burden of burn-in, and loaded the resulting tree sequences into SLiM (*26, 27*). We drew
functional effect sizes for these variants, placed individuals into continuous space, and ran
simulations forward-in-time for 5,000 generations. After that, the geographic distribution of
the species experienced impacts as expected during global change: every generation, 0.001 of
one edge of the species distribution got its carrying capacity reduced to 0. This meant that
over 1,000 generations the whole species would disappear (note that this is a reasonable
fraction of area reduction given the estimates of yearly deforestation and habitat change in
section V). We subsequently overlayed neutral mutations on the tree sequence using
msprime, and analysed genomes sampled throughout the extinction process (by tracking them
in the tree sequence output) and extracted using tskit.
641
642

II.3.7 Connection of z_{MAR} with the isolation-by-distance pattern

Ultimately, z_{MAR} is a complex integrator of evolutionary forces acting in space (mutation, migration, drift, selection) and captures how structured the distribution of a species' mutations is. Although the isolation-by-distance pattern conceptually resembles z_{MAR}, we have found no obvious analytical expression that relates both. Note that F_{ST} is defined based on heterozygosity or π, instead of the number of segregating sites (i.e., mutations M). For instance, using Hudson's estimator (*28*) to compute F_{ST} across a set of populations we calculate $F_{ST} = 1- (\pi_w / \pi_b)$, where π_w is the diversity or heterozygosity within a population and π_b is the same parameter calculated for the meta-population. Plotting F_{ST} of a metapopulation by the distance of the farthest demes shows the typical non-linear trend of isolation-by-distance, which shows that very close populations have similar allele frequencies whereas populations further away drift apart. A challenge of F_{ST} is that it requires pre-defining discrete populations, which is straightforward in stepping-stone simulations but hard in real data. Comparing average F_{ST} of our 14x14 spatial demes and z_{MAR}, we see that the two parameters correlate (Fig. S10C). However, it appears that for low values of F_{ST}, z_{MAR} captures more variation across the simulations (Fig. S10). These patterns were also confirmed in continuous space simulations (not shown).

Fig. S10 | SLiM population genetic simulations in 2D comparing F_{ST} and z_{MAR}

Neutral SLiM simulations with different degrees of migration. (A) Hudson's F_{ST} across populations with different area subsamples. Following the expectation of the isolation-by-distance pattern, as the distance between the farthest demes in the subsample increases, F_{ST} becomes larger and saturates at large distances. (B) The mutations-area relationship. (C) Comparison between F_{ST} and z_{MAR}.

II.4 The loss of mutations (genetic diversity) in space

The aim is to predict the fraction of genetic diversity loss, x_M, from shrinking of an ecosystem by an area a. To define all terms, we then have a past area A_{t-1} and a present reduced area $A_t = A_{t-1} - a$, and a fraction of area extinct $x = a/A_{t-1}$

       We first think of the loss of genetic diversity x_M through the basic process of losing individuals. From the population genetics's coalescent theory derivation of the number of mutations or segregating sites from individuals we got the approximation $M \sim log(N)$. Assuming the loss of area is simply the loss of individuals ($A=N$), we can derive the fraction of genetic diversity loss as:

$$x_M = 1 - \frac{M_t}{M_{t-1}} = 1 - \frac{log(N_t)}{log(N_{t-1})} = 1 - \frac{log(N_{t-1}(1-x))}{log(N_{t-1})} = 1 - \frac{log(N_{t-1}) + log(1-x)}{log(N_{t-1})}$$

$$= -\frac{log(1-x)}{log(N_{t-1})}$$

       The loss of mutations is then in the scale of: *log(1-x)*; which is very slow, as we expected from having derived the trend that under panmixia $z_{MAR} \approx 0$. A substantial loss of genetic diversity in this case only happens when population extinction is almost complete.

698
699   Species do not typically behave perfectly panmictic given different z_{MAR} values. Under
700   population structure, we can use our relationship to project the number of mutations (genetic
701   diversity) lost as the geographic distribution due to habitat loss or climate change following
702   equation:
703

704
$$x_M = 1 - \frac{M_t}{M_{t-1}} = 1 - \frac{MAR(A_t)}{MAR(A_{t-1})} = 1 - \frac{A_t^z}{A_{t-1}^z} = 1 - \left(\frac{A_t}{A_{t-1}}\right)^z = 1 - (1-x)^z.$$

705
706   In the most extreme scenario of $z_{MAR} \approx 1$, the fraction loss of geographic area directly
707   translates to the same fraction loss of genetic diversity.
708
709   Reality should be in between the panmictic and fully-migration-limited cases. With
710   combinations of environmental selection, non-equilibrium demography, and long-range
711   dispersal, we may get intermediate z_{MAR} values, and it will be empirical estimates that can
712   inform us how much may be lost (Section III).
713
714   **II.5 Recovery of genetic diversity after a bottleneck or local extinction**
715

716
717   *Fig. S11 | 2D stepping-stone msprime simulations with extinction and recovery*
718   *(A) Recovery of genetic diversity (number mutations) after loss of a fraction of the population. (B) Recovery of genetic*
719   *diversity after instantaneous loss of a fraction of the population and consecutive repopulation.*
720   **Simulations with number of generations until recovery that are exceedingly large are assigned a value of 1,500, as none*
721   *are realistic for current conservation timelines.*
722
723
The intuition that rapid recovery of genetic diversity may be possible is likely flawed.
While genetic recovery may be faster than speciation rates, which are on the order of millions
of years, the time for a set of populations that went through a simulation burn-in of 1,000
generations (not yet in diversity equilibrium), and that suffer an instantaneous 5% reduction
of area and an instantaneous recovery (e.g., through reforestation) would range from 20-90
generations. This number of generations for long-lived species would translate into centuries
or millennia of recovery without further impacts. About 49% of simulations – including every
simulation that reached equilibrium (burn-in generations >10,000) – have a recovery time of
more than a thousand generations (Fig. S11).
733
734

735   SUPPLEMENTAL RESULTS
736
737   **III. The mutations-area relationship with the 1001 Arabidopsis Genomes**
738
739   We begin testing the idea of a general mutations-area relationship using the extensive
740   sampling of the model plant species *Arabidopsis thaliana* and the 1001 Arabidopsis Genomes
741   Project (*29*). This section will serve as a case study to explore different approaches and biases
742   when building MAR to then apply the learned lessons across species (section IV).
743
744
745   **III.1 The Site Frequency Spectrum of the 1001 Arabidopsis Genomes**
746
747   We began analyzing the frequency distribution of 11,769,920 biallelic genetic variants (i.e.,
748   mutations), which is typically called the Site Frequency Spectrum (SFS) in population
749   genetics.
750
751

753   ***Fig. S12 | Mutation abundance study in A. thaliana***
754   *(A) Site Frequency Spectrum (SFS). (B) Preston plot of mutation abundances. (C) Whittaker plot of mutation rank*
755   *abundances.*
756
757   To showcase the similarities to the Species Abundance Distributions (SAD), we use
758   the Whittaker plot of mutation rank abundance (Fig. S12) that suggests a log-normal of S-
759   shape may be the best fitting model (Table S3). For a review listing many popular models,
760   see (*30*), and for implementation details of 13 SAD models see the thorough manual of R
761   package SADS (*31*). As we shall see later, the log-normal distribution seems to be the best fit
762   across species.
763
764

Fig. S13 | Fit of mutation abundance study in A. thaliana with different SAD models
Representative models from Table S3 are plotted along with the observed frequency of 11,769,920 mutations.

Although model AIC captures best the fit of a curve accounting for the difference in
parameter complexity of each model and the statistical distributions behind, we often are
interested in the variance explained. We then calculated a proxy of predictive accuracy using
a pseudo-R^2 approach of the difference between the model fit and the observed data as:
$R^2 = 1 - \frac{SS_{res}}{SS_{tot}}$. For *A. thaliana*, we used 10,000 SNPs sampled at random to an accuracy of
over $R^2 > 0.999$ for both the top log-Normal model and the bottom log-Series model,
indicating that all "commonness of rarity" models must have a pretty good fit of mutation
frequency data.

Table S3 | AIC values for model fit of common species distribution curves.
For each SAD model, the degrees of freedom and the delta AIC compared to the top model are reported.

Model	dAIC	df
log-Normal	0	2
Poisson	7204.37509	2
Geometric	44267.5475	1
Weibull	45872.3678	2
Gamma	48805.6065	2
Broken Stick	49076.4368	0
UNTB (MTZSM)	168434.181	1
log-Series	168434.726	1

The typical SFS from population genetics is of course not implemented in current packages for Species
Abundance Distributions like R sads. For comparison, in the main text we also calculate the log
likelihood and AIC of this following the standard population genetics likelihood:

$$logL = \sum_i log(\frac{1}{Nq_i}) - log(H_n(N-1))$$
,

where N represents the number of individuals in a sample, and q_i is the minor allele frequency of a
SNP in the sample, in the main text calculated for i=1…10000 random SNPs (see main text). As
before, H_n is the harmonic number function.

794
795 **III.2 Building the Mutations-Area Relationship**
796
In the following, we explain how the area was estimated that was used to compute z_{MAR} on
real world data. In short, we used a grid on the world map, with samples placed on the map
based on their geo-coordinates of origin (Fig. 1). We first create square spatial subsamples of
the *Arabidopsis thaliana* geographic distribution (Fig. 1, Fig. S15) and quantify diversity M
as the total segregating sites. Excluding zeros, these two variables are fed to the sars_power
function from the R SARS package (*32*).
803
Although the power law mutations-area relationship was already theoretically
motivated (II.3), here we also fit different types of functions typically applied to the Species-
Area Relationship. Doing this, we reach the conclusion that multiple models perform very
similarly, and the classic power law is among the top models, see Table S4. Although small
marginal fitting accuracy could be achieved with other models, for mathematical convenience
and historical continuity, we use the power law for later sections and the study of MAR
across species (Sections IV and V).
811
812
813 *Table S4 | Different SAR curves fit to mutations.*
814 We fit 20 different functions and calculated the variance explained (R2), Pearson's r, and Spearman's rho.
815

Model	R2	r	rho
Asymptotic regression	0.21825683	0.46717965	0.53510077
Beta-P cumulative	0.22012799	0.46917799	0.53374757
Chapman Richards	0	NA	NA
Cumulative Weibull 3 par.	0.21929646	0.468291	0.53374757
Cumulative Weibull 4 par.	0.21930145	0.46829633	0.53374757
Extended Power model 1	0.21833611	0.46726449	0.53026812
Extended Power model 2	0.21682584	0.46564561	0.53462775
Gompertz	0.16393078	0.40488366	0.45964364
Heleg(Logistic)	0.21929721	0.4682918	0.53531975
Kobayashi	0.22228406	0.47147011	0.53526975
Linear model	0.19579007	0.44248171	0.53510077
Logarithmic	0.20280401	0.45033767	0.53430311
Logistic(Standard)	0.22536996	0.47473146	0.53549765
Monod	0.22500999	0.47435217	0.53579276
Negative exponential	0.22801633	0.47751055	0.53447179
Persistence function 1	0.21929612	0.46829063	0.53501182
Persistence function 2	0.21760028	0.46647645	0.53409266
Power	**0.21929556**	**0.46829004**	**0.53543785**
PowerR	0.21753225	0.46640353	0.53493321
Rational function	0.22072491	0.46981369	0.53451874

816
817
Because in the species literature it is recommended to only quantify richness of
endemic species (*33*), we also count segregating sites that are private to the area subsample,
creating the equivalent endemic-mutations-area relationship (EMAR) (*33*). The MAR slope
and 95% Confidence Interval was $z = 0.324$ (0.238 - 0.41) (Table S5, Fig. S14 A), while the
EMAR was $z = 1.241$ (1.208 - 1.274) (Table S6, Fig. S14 B). Interestingly, the endemics-area
relationship of $z \approx 1$ resembles that of endemic species, whereas the total mutation
relationship with area is above that of species relationships, which typically follows the
canonical $z \approx 0.2 - 0.4$.
826

827   We must note that EMAR, the genetic analogy of the Endemic-(species)-Area
828 Relationship (EAR) may not be that meaningful when analyzing genomic data (we did not
829 find a way to theoretically motivate it in section II), and later we see it overestimates loss in
830 our simulations (Fig. S18)
831

832 *Table S5 | The mutations-area relationship (MAR).*
833 Fitted values in a log-log power function between area sampled and mutations discovered.

	Estimate	Std. Error	t value	P	2.5%	97.5%	nls.Est.	nls.2.5%	nls.97.5%
c	494.565432	135.6314588	3.646392	0.0003138	223.3025141	765.8283493	494.5531270	278.1107276	822.829918
z	0.323727	0.0430277	7.523681	0.0000000	0.2376715	0.4097824	0.3237367	0.2430303	0.413162

834
835

836 *Table S6 | The endemic-mutations-area relationship (EMAR).*
837 Fitted values in a log-log power function between area sampled and endemic mutations discovered.

	Estimate	Std. Error	t value	P	2.5%	97.5%	nls.Est.	nls.2.5%	nls.97.5%
c	0.0001001	0.0000231	4.337758	1.98e-05	0.0000539	0.0001463	0.0001001	0.0000635	0.0001555
z	1.2411831	0.0165268	75.101442	0.00e+00	1.2081296	1.2742366	1.2412125	1.2096087	1.2737927

838
839
840

842 **Fig. S14 | The mutations-area and endemic-mutations-area relationships in A. thaliana.**
843 *Dividing* A. thaliana *native geographic distribution into a 1 degree lat/long grid, square areas with 1 degree side-length to*
844 *36 degrees side-length were randomly placed (n=100 for each size) across the distribution, and genetic diversity metrics*
845 *were computed to produce the (A) Mutations-Area Relationship and (B) Endemic-Mutations Area relationship.*
846
847

848 **III.3 Testing for potential numerical artefacts**
849

We wondered whether MAR estimates may be affected by some numerical artefacts in our
software pipeline (available at https://github.com/moiexpositoalonsolab/mar). For instance,
real world data may have uneven sampling in space, the spatial resolution of georeferenced
samples may vary, projection of samples into gridded maps may have limited resolution,
software pipelines may produce biased estimates, etc. To test this, we conducted several
experiments:
856

Lower bound of the method for z_{MAR}. Our first experiment when building the MAR
aimed to make sure that spatial sampling, or some unknown bias in genome sequencing, or
the number of samples used, are not creating artificially large z_{MAR}. We then simulated a mock
dataset of *A. thaliana* with the same number of mutations, samples, and using the original
geographic locations. The number of SNPs were also sampled in a way that we created a
canonical 1/q SFS for the whole species. Under no biases, we then expect the MAR to follow
the theoretical derivation under panmixia with a $z\sim0$. This exercise confirmed we get a value
approaching zero: $z=0.033$, (-0.095 - 0.162).
865

866   *Table S7 | MAR built with different area calculations and grid sizes*
867

Grid resolution (deg.)	z_{MAR} [CI95%] (cell area)	z_{MAR}[CI95%] (total area)
A=N	0.431 (0.423 - 0.439)	NA
0.1	0.435 (0.424 - 0.446)	0.367 (0.281 - 0.454)
0.25	0.454 (0.449 - 0.459)	0.422 (0.376 - 0.467)
0.5	0.488 (0.465 - 0.511)	0.352 (0.152 - 0.551)
1	0.543 (0.529 - 0.558)	0.389 (0.295 - 0.483)
2.5	0.644 (0.6 - 0.688)	0.388 (0.251 - 0.526)
5	0.617 (0.205 - 1.029)	0.403 (-0.204 - 1.011)

868
869
870
Grid sizes, area calculations, and non-random spatial sampling. In order to
streamline geospatial operations, we implemented the MAR relationship calculations in this
project using R raster objects (*34*). This required projecting the collected samples of a species
and the observations of any given mutation into a world map (i.e., each mutation's geographic
distribution). Necessarily, in order to be able to assign areas to sets of samples or mutations
on the map, the projection requires the choice of a grid size. The larger the grid size (e.g.,
lower spatial resolution), the faster the spatial operations can be performed. Further, for larger
grid sizes, we expect the slope of MAR to be more influenced by larger-scale patterns, while
for smaller grid sizes, the MAR will be influenced by smaller-scale patterns. To test this, we
repeated the subsampling of *A. thaliana* distribution with grid sizes ranging 0.1 degrees
latitude/longitude (roughly 10km side-length in temperate regions) to 10 degrees (roughly
1,000 km side-length). The estimates were roughly consistent between 0.4-0.6, but increases
in value at larger grid sizes (row in Table S7 for large grid size values), a scale-dependent
pattern that resembles results of SAR of species in ecosystems fitted at different scales (*10*).
885
Because we often have sparse samples of individuals in space, we devised two
strategies to calculate areas during the subsampling of MAR (see cartoon in Fig. S15): (A)
the total square area of the minimum and maximum latitude/longitude values of all the
samples analyzed. That is, simply the area of the red box in the figure. (B) the sum of areas of
grid cells that contain at least one sample. That is, the sum of the grey squares within the red
box in the figure. In addition, we also calculated the MAR relationship assuming the total
area is equal to the number of individuals (*A=N*) (which should be theoretically equivalent to
a grid of very high resolution where we end up with a maximum of one individual sampled at
any grid cell).
895
Table S7 values suggest there is a dependency of z_{MAR} with the grid size when areas
are calculated as the sum of grid cells with at least one sample. Our intuition for this pattern
is that lower resolution grids (e.g., 5 degrees side) lead to some grid cells having many
samples, which would increase the number of mutations discovered when discovering the

area. On the other hand, the calculation of z_{MAR} using the total area does not seem to affect the
z_{MAR} estimate; however, because large areas often do not have samples (limiting the potential
to find new mutations), it creates a higher variance in the estimate of z_{MAR} (see confidence
intervals in Table S7 and Fig. S16). Here, we favored consistency of z at the expense of
broader, more conservative confidence intervals. All the estimates reported below and in the
main text therefore use the total area approach.
906
907
908

910   ***Fig. S15 | Cartoon of raster sampling to build the MAR***
911   *Map of mock samples of a species projected into a raster. Grey scale indicates the number of samples per grid cell. Red*
912   *boxes exemplify the process of spatial subsampling of increasing area to build the MAR relationship. Two example grid sizes*
913   *were created for illustrative purposes: (A) Small grid size or high spatial resolution. (B) Large grid size or low spatial*
914   *resolution.*
915
916

918   ***Fig. S16 | MAR comparison with different area calculations.***
919   *(A) Using total area, (B) using grid cell sum with at least one sample. For 1 degree latitude/longitude grid cell.*
920
921   **Geographic subsampling strategy (inwards, outwards, random)**. It has been
indicated that the way the Species-Area Relationship (SAR) and Endemics-Area Relationship
(EAR) are created may create differences in the scaling parameter z. The plots and estimates
above were produced by randomly placing boxes of different size or area across the
distribution of the species. Often, however, either discovery of species or extinction happen

in certain patterns. For instance, we often imagine sampling an ecosystem concentrically
outwards from a focal point, whereas we may think of the extinction process of species area
reductions being concentrically inwards (*33*). Because these patterns seem of importance, we
also calculated the MAR and EMAR outwards from the latitude and longitude median of all
the samples in the map, moving outwardly until the map is filled (Fig. S17, Table S8).
Likewise, the inward pattern is conducted in an inverse manner.
932
933
934

936   ***Fig. S17 | MAR and EMAR in Arabidopsis thaliana using outward and inward sampling.***
937   *Dividing A. thaliana native distribution in 1 degree lat/long grid, a square area of 1 degree was placed at the median of the*
938   *sampling range and was expanded iteratively by 1 degree lat/long until all the area of the distribution was covered. (A-B)*
939   *MAR and EMAR using a typical outward sampling. (C-D) MAR and EMAR using an inward sampling. The latter may not be*
940   *a common process of sample collection, but it is common for extinction progress.*
941

942   ***Table S8 | Outward and inward MAR and EMAR***
943   *The MAR and EMAR relationship computed with inward or outward nested subsampling, calculating area only as those*
944   *cells with samples.*
945

Relationship	z
MAR outwards	0.444 (0.412 - 0.476)
EMAR outwards	1.086 (0.982 - 1.189)
MAR inwards	0.561 (0.524 - 0.597)
EMAR inwards	1.295 (1.192 - 1.399)

946
947

Incomplete sampling of the species. To check whether the relationship holds with
few individuals of a species or limited geographic distributions, we compared the species-
wide MAR with that of subset populations. Downsampling the native distribution of *A.*
thaliana to a region within North-East Spain (-2.00–4.25 degrees East, 36.52–42.97 degrees
North), or to a region within Germany (2.69–13.73 degrees East, 50.0–52.0 degrees North),
and using only 1,000 SNPs, we recovered z_{MAR}= 0.423(0.233-0.614) for Spain and
0.525(0.242-0.807) for Germany, which were close to the estimate based on the whole
distribution (Table 1). This result is reassuring in that if migratory patterns are relatively
homogeneous, one may be able to estimate this parameter from a subset of the species
distribution. For heterogeneous population structure cases, we expect incomplete sampling to
produce unreliable estimates.
959
960   **Number of genome-wide SNPs used.** To check whether different numbers of SNPs
961   used for the analyses would lead to different z_{MAR}, we conducted analyses with random
962   subsets consisting of 100, 1,000, and 10,000 SNPs, replicated 3 times. Estimates had a
963   coefficient of variation of 4.7%, which is way below the standard error of typical estimates
964   (Table 1).
965
Locally-adaptive variants. We then aimed to understand the effect of utilizing SNPs
that appear to be related to adaptation. To study this, we utilized an outdoor climate-
manipulated experiment that recorded fitness data (survivorship and reproduction output of
seeds) for 515 *Arabidopsis thaliana* ecotypes part of the 1001 Genomes set in 8 environments
(Exposito-Alonso, 2019). We devised two sets of alleles: 10,000 that were negatively
correlated with fitness in a Genome-Wide Association across 8 different environments, and
10,000 alleles that were associated positively with fitness in one environment but negatively
in another (antagonistic pleiotropic). The MAR relationship was computed as before and
compared to the original random (putatively neutral) set of alleles from the previous sections
(Table S9). Although we see a trend that locally-adaptive alleles have a slightly higher *z*,
estimates overlap. The effects seen here of having smaller *z* for adaptive alleles than neutral
variation could, however, be due to top GWA SNPs often being ascertained to higher
frequency than background SNPS.
979
980

981   *Table S9 | MAR for putatively neutral, deleterious, and locally adaptive alleles in Arabidopsis thaliana*
982

SNP set	*z*
neutral	0.324 (0.238 - 0.41)
globally deleterious	0.209 (0.13 - 0.288)
locally adaptive	0.291 (0.217 - 0.365)
globally positive	0.234 (0.137 - 0.332)

983
984
985   **III.4 Local population extinction in Arabidopsis**
986
Using the MAR framework, we can make projections of loss of mutations (or its inverse, the
remaining genetic diversity. By doing this, the known intuition is that with *z >1* (as from
EMAR) the decrease of diversity is much faster than the decrease of habitat, but with *z < 1*
(as from MAR), there is a (desirable) slower dynamics of genetic loss. In the latter, despite
habitats disappearing, reservoirs of mutations distributed across different locations enable
conservation of certain variation. To study which one is more likely and to observe the

stochastic nature of genetic diversity loss, we simulated in silico population extinctions of
map cells from the Arabidopsis map (Fig. 1) and directly estimated from the genome matrix
of remaining individuals the remaining genetic diversity. These simulations were
implemented to capture different hypothesised patterns of extinction (see main text). All,
however, agree with the more hopeful estimate of $z_{MAR} \approx 0.3$.
998
999
1000

1001
1002   ***Fig. S18 | Loss of mutations with habitat loss in A. thaliana.***
1003   *Predictions based on MAR and EMAR functions and in silico extinction stochastic simulations in A. thaliana.*
1004
To study the fit of the genetic loss predictions based on MAR relationships and the
results from computer simulations, we calculated a pseudo-R^2 based on the squared
differences between the predicted line and the "observed" genetic loss as: $R^2 = 1 - \frac{SS_{res}}{SS_{tot}}$.
This results in a high fit R^2=0.872 of the MAR, built from random samples of distribution
areas, while the EMAR had a poor fit due to overestimation of genetic loss: R^2=-0.710
(negative values indicate predictions are worse than the mean of the data).
1011
1012
1013   **III.5 Potential impacts of genetic loss in adaptability**
1014
Although likely imperfect, Genome-Wide Associations could help to understand the
relevance of mutations in different frequency classes in model organisms such as *Arabidopsis*
thaliana. Fig. S19 shows the site frequency spectrum and a metric of the "total accumulated
effect in fitness" of the alleles in every bin. Effect sizes were retrieved from GWA on lifetime
fitness of 515 ecotypes in outdoor experiments (*35*). The average effect size across 8 fitness
GWA from 8 experimental combinations were used: high/low precipitation, high/low latitude
of outdoor stations, and high/low plant density. This exercise showcases the phenomenon that
low frequency variants often have strong effect sizes, which is expected under a stabilising
selection quantitative model (*36*). Because low frequency alleles will be the first to be lost
during a bottleneck (as would happen with the rapid extinction of populations of a species),
we may expect to lose variants that are related to fitness and thus potentially lose diversity
that could be advantageous in some environments. Alternatively, deleterious mutations are
also expected to be at low frequency, in which case would also make them more easily lost.

1028
1029

1030

1031 ***Fig. S19 | Bias of low frequency mutations and effect size for fitness traits in A. thaliana.***
1032 *Grey bars represent the site frequency spectrum (scaled for visualisation purposes). The black dots represent the mean*
1033 *absolute effects of alleles as estimated from GWAs with 515 accessions scored for fitness traits in 8 outdoor experiments.*

1034
1035

To further build intuition on the progress of extinction in relation to loss of genetic
diversity that is not neutral, we repeated warm edge extinction simulations with several
subsets of alleles: randomly selected SNPs, SNPs that were associated positively in 2
environments (low precipitation Spain and high precipitation Germany) (labelled globally
positive), and SNPs that were associated positively in one environment and negatively in the
other (labelled antagonistic pleiotropic or putatively locally-adaptive). This (Fig. S20)
supports our intuition that although putatively functional alleles (or alleles tightly linked to
such functional ones) may have slower loss dynamics than neutral variants due to a high
frequency and z_{MAR} , certain population extinction patterns may actually lead to rapid loss of
potentially-adaptive genetic diversity. The complexity of these patterns, together with the
evolutionary feedback created by lowering genetic standing variation that affects fitness,
make the inference of adaptive capacity loss even more difficult than just inferring the loss of
genetic diversity itself.
1049

1050

Fig. S20 | Simulations illustrating the potential loss of locally-adaptive mutations in A. thaliana.
Simulations of extinction using multiple patterns of population losses with different subsets of alleles ascertained to show positive associations in fitness GWA in two outdoor experiments (green), positive associations in one environment (e.g. low precipitation) but negative in a second environment (e.g. high precipitation) or vice versa (green). These were compared to a random set (grey).

1058

Fig. S21| Extinction simulations showing proxies of adaptive capacity of A. thaliana.
Using estimated allele effect sizes from 10,000 SNPs in the 1% P-value tails of several Genome-Wide Associations, we show (A) Percentage of change of Va as a proxy of adaptive potential and (B) raw square sum of allele effects to showcase the inflating effect of intermediate frequency alleles. Grey background shape indicates the minimum and maximum boundaries of trajectories created by replicated frequency-matched non-effect sets of SNPs (one per GWA). The trajectories of some effect alleles appear to show faster loss than the non-effect background trajectories.

III.6 Case study of a massive natural bottleneck

1067
1068

A recent colonisation of North America by *Arabidopsis thaliana* can help us understand the recovery of genetic variation. Whole-genome sequencing of 100 specimens of North American *A. thaliana* indicates that it migrated from its native range of Europe to North

1069
1070
1071

America in the 17th century, and began spreading across the continent from a genetically-
homogeneous population (*37*). Despite ideal conditions to re-gain genetic diversity—a
continental population expansion aided by human travel (*38*, *39*)—only ~8,000 new
mutations were detected through spontaneous accumulation, equivalent to only ~0.067% of
the species-wide native genetic diversity. Because most of these mutations are at very low
frequency, as expected during population expansion, the scaling of genetic diversity with area
is approximately 1 (z_{MAR} = 1.025 [CI95%: 0.878 - 1.173]).
1079
1080
1081

IV. The mutations-area relationship in diverse species

Every dataset was retrieved online either from the published article in the form of VCF or fastq files, or provided by the study authors upon request. All datasets were first transformed into PLINK files using PLINK v1.9 (*40*). For computational efficiency, and since we showed random subsampling does not appear to affect calculations of z_{MAR} (Section III.3), we conducted all analyses with up to 10,000 randomly selected SNPs for each species sampled genome-wide, or in the largest chromosome for those species with large genomes. We aim to use mostly unfiltered SNP datasets to avoid ascertainment biased toward intermediate frequency SNPs, and therefore we did not apply a MAF filter for any analyses. By default, PLINK transforms SNP matrices into biallelic (if multiallelic, it takes the two most common alleles). Although the preservation of structural genetic variation may also be relevant and may have important consequences in adaptation (*41*), we do not expect dramatic differences in their scaling relationship compared to biallelic SNPs, as their SFS are relatively similar (Structural variants may show a skew to lower frequency, resulting in steeper z_{MAR}. By excluding those, our analyses may be conservative). In order to properly characterise the geographic distribution of a mutation using all available geo-tagged individuals, we filtered for genotyping rate (plink --geno), and the final value is reported per dataset.

Details for dataset processing or homogenization are described below.

- The 1001 Arabidopsis Genomes Consortium (*29*) generated a WGS Illumina sequencing dataset of *Arabidopsis thaliana* comprising 1,135 individuals and 11,769,920 SNPs. The VCF with the data is available at: https://1001genomes.org. The raw sequencing data is available at https://www.ncbi.nlm.nih.gov/bioproject/PRJNA273563. These included recently colonised regions such as North America or Japan. Analyses of z_{MAR} were calculated only for the native range, which comprises most of the species diversity (>99%) and 1001 individuals. For computational efficiency, we conducted analyses using randomly sampled SNPs from chromosome 1, as we did not observe any difference when sampling from other chromosomes. A number of MAR approaches were tested in this species (section III). For homogeneity, the final reported estimate (Table 1) was conducted following the same procedures as other species with a random sample of 10,000 SNPs.

- Lucek & Willi (*42*) recently published a dataset of WGS Illumina sequencing 108 *Arabidopsis lyrata* individuals from North America, which the authors directly shared as a VCF. The raw data is available at https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJEB30473. We retrieved the latitude/longitude data from the supplemental material. We applied a genotyping rate filter ending with a dataset of 0.955431 genotyping rate. 10,000 SNPs were subsetted at random from the genome-wide data.

- Kreiner et al. (*43*) WGS Illumina sequenced 165 individuals of *Amaranthus tuberculatus*. The raw data is available in the link https://www.ebi.ac.uk/ena/browser/view/PRJEB31711. The authors provided a VCF. Overall, 155 individuals contained latitude and longitude information and were kept for the analyses. The genotyping rate was 0.98162 and we subsetted randomly 10,000 SNPs.

1132   -   Supple et al. (*44*) generated a dataset of *Eucalyptus melliodora* of 275 individuals
1133   from 36 broadly distributed populations. The dataset was produced by Illumina
1134   sequence Genotyping-by-Sequencing (GBS) libraries digested with ApeKI as in
1135   Elshire et al. (2011). The raw data is available at
1136   https://www.ncbi.nlm.nih.gov/bioproject/PRJNA413429/. The authors provided the
1137   dataset in PLINK format. Genotyping rate was 0.769807 but we did not apply a
1138   further filter to avoid reducing the total number of variants. We conducted analyses
1139   with all 9378 SNPs. The genotyping rate in this dataset is likely not problematic as the
1140   total number of GPS locations is 36, with multiple individuals sampled closely. This
1141   sampling scheme probably allows to characterise an allele's distribution correctly
1142   despite the lower genotyping rate.
1143

1144   -   Vallejo-Marin et al. (*45*) generated a GBS dataset of 521 Mimulus plants, with 286
1145   samples being *Mimulus guttatus* from its native distribution. Libraries for
1146   Genotyping-By-Sequencing were prepared with PstI enzyme as described in Twyford
1147   & Friedman (2015) and sequenced using Illumina. The VCF of this dataset is
1148   available at http://hdl.handle.net/11667/168 and was also directly shared by the
1149   authors. After applying a filtering for missingness, we ended up with a genotyping
1150   rate of 0.904192 and 1,498 SNPs, which were used for the analyses.
1151

1152   -   Lovell & MacQueen (*46*) generated a WGS Illumina sequencing dataset of
1153   Switchgrass, *Panicum virgatum*, of a collection of 732 individuals and 33,905,044
1154   variants. The raw data is available at:
1155   https://www.ncbi.nlm.nih.gov/bioproject/PRJNA622568. The authors provided a
1156   VCF file and latitude/longitude tables. 576 individuals were from natural collections.
1157   The dataset contains also other collections such as cultivars, which were not used to
1158   build the MAR. The genotyping rate was 0.976393 and analyses were conducted with
1159   10,000 SNPs drawn from the largest chromosome.
1160

1161   -   MacLachlan et al. (*47*) generated a SNP chip dataset of *Pinus contorta* comprising
1162   929 trees with latitude and longitude information and 32,449 SNPs. Genotyping was
1163   conducted with the AdapTree lodgepole pine Affymetrix Axiom 50,298 SNP array
1164   and data was provided in the supplemental material of the paper along with custom
1165   scripts to parse the data. The database is available at
1166   https://datadryad.org/stash/dataset/doi:10.5061/dryad.ncjsxkstp. The genome matrix
1167   was transformed into PLINK. The genotyping rate was 0.959146, and analyses were
1168   conducted with 10,000 randomly drawn SNPs. The fact that this dataset was created
1169   with ascertained SNPs likely generates a frequency bias. In Fig. S22, one can see that
1170   this may be a problem to calculate z_{MAR}, as the mutations~area graph appears
1171   nonlinear and rapidly saturates. This confirms the expectation that SNPs are
1172   ascertained to be common, as they are discovered immediately with very few samples.
1173

1174   -   Tuskan et al. (*48*) WGS Illumina sequenced 882 *Populus trichocarpa* trees. The
1175   dataset includes 28,342,826 SNPs. The data is available under this DOI
1176   https://doi.ccs.ornl.gov/ui/doi/55 which redirects to a globus data sharing platform.
1177   The authors provided the dataset as a VCF along with latitude/longitude coordinates.
1178   This dataset was downsampled to the first chromosome. The genotyping rate was
1179   0.921191, and 10,000 SNPs were randomly sampled for analyses.
1180

1181   -   The Anopheles gambiae 1000 Genomes Consortium (*49*) (Phase 2) produced Whole-
1182   Genome Illumina sequencing data for 1142 wild-caught mosquitoes of *Anopheles*
1183   *gambiae*. All raw and processed data are available through
1184   https://www.malariagen.net/data. We downloaded a VCF and latitude/longitude
1185   coordinate files. The VCF was filtered for genotyping rate ending up at a 0.998895
1186   rate. For efficiency, 10,000 randomly-selected SNPs from the VCF of the largest
1187   chromosome 2L were used for analyses downstream.
1188

1189   -   Fuller et al. (*50*) WGS Illumina sequenced 253 coral individuals of *Acropora*
1190   *millepora* in 12 reefs. The dataset was downloaded as fastq files from the published
1191   online material from https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA593014,
1192   and SNPs were called as described in the supplemental material ending with
1193   17,931,448, which were filtered to achieve a genotyping rate of 0.935709 for a total of
1194   2,512 SNPs, which were used in the analyses.
1195

1196   -   Ruegg et al. (*51*) generated a dataset of 219 birds *Empidonax traillii*, for which 199
1197   could be matched with geographic coordinates. SNPs were ascertained from several
1198   publications using RAD seq and Fluidigm 96.96 IFC described and available in their
1199   repository https://github.com/eriqande/ruegg-et-al-wifl-genoscape. A total of 349,014
1200   SNPs were parsed using their custom scripts and we transformed them into PLINK
1201   files. A genotyping rate filter was applied ending with a 0.96061 rate and 195,700
1202   SNPs. 10,000 SNPs were selected at random for downstream analyses. Similarly, as
1203   with the *Pinus contorta*, the incorporation of some ascertained SNPs in the dataset
1204   based on Fluidigm technology could lead to quick saturation of the MAR curve (Fig.
1205   S22).
1206

1207   -   Bay et al. (*52*) generated a dataset of 199 *Setophaga petechia* birds using a Restriction
1208   site–associated DNA sequencing (RAD-Seq). The raw data is available at
1209   https://www.ncbi.nlm.nih.gov/bioproject/421926. The authors shared a VCF file, with
1210   a genotyping rate of 0.962419 and a total of 104,711 SNPs. 10,000 SNPs were
1211   selected at random for downstream analyses.
1212

1213   -   Kingsley et al. (*53*) produced a dataset of 80 *Peromyscus maniculatus* deermice, for
1214   which 78 could be matched with geographic locations. The SNP dataset was produced
1215   using MY-select capture followed by Illumina sequencing. The VCF and PLINK files
1216   are available via Figshare at https://doi.org/10.6084/m9.figshare.1541235. The dataset
1217   included a total of 14,076 variants which were filtered to achieve a genotyping rate of
1218   0.940411 for 2,946 SNPs, which were used in subsequent analyses.
1219

1220   -   We identified two published datasets for wolves. Smeds et al. (*54*) produced a WGS
1221   Illumina sequencing dataset and combined it with pre-existing datasets for a total of
1222   349 local dog breeds and wolves, of which 230 were *Canis lupus* from natural
1223   populations. However, these samples did not have GPS locations assigned. The
1224   second dataset we identified was from Schweizer et al. (*55*), which contained 107
1225   geo-tagged grey wolves from North America using a capture and resequencing
1226   approach for 1040 genes. The raw data is available at
1227   https://trace.ncbi.nlm.nih.gov/Traces/sra/?study=SRP065570, and meta-data along
1228   with a VCF area available at https://doi.org/10.1111/mec.13467. This data contained
1229   13,092 SNPs at 0.993061 calling rate, and a better geographic resolution. We report
1230   data for the second dataset.

1231
-    The 1000 Genome Consortium (*56*) created WGS Illumina sequencing for over 2,504
humans and 24 unique geographic locations. We downloaded chromosome 1 from
http://ftp.1000genomes.ebi.ac.uk/vol1/ftp/datacollections/
1000G2504highcoverage/working/20190425NYGCGATK/ and gathered the
population locations from https://www.internationalgenome.org/data-
portal/population. To conduct analyses, we subsampled 10,000 SNPs at genotping
rate 0.991069.
1239
-    Palacio-Mejia (*57*) used WGS for 591 *Panicum hallii* individuals to sequence at low
coverage. The raw data is available at
https://www.ncbi.nlm.nih.gov/bioproject/PRJNA390994. The authors shared an
unfiltered VCF of 45,589 SNPs. Because of the low-coverage, stringent filters of
calling rates as used for other species would lead to removing all SNPs, and we settled
on a genotyping rate of 0.825824 for 242 variants, all of which were used for
downstream analyses.
1247
1248   -   Royer et al. (*58*) produced a SNP dataset using RAD-Seq based Genotyping-By-
1249   Sequencing of 290 *Yucca brevifolia* (Joshua Tree) individuals. A total of 10,695 SNPs
1250   with a genotyping rate of 0.897501 wre used for the analyses. The data was available
1251   at Dryad https://datadryad.org/stash/dataset/doi%253A10.5061%252Fdryad.7pj4t.
1252
1253   -   Kapun et al. (*59*) produced a WGS dataset of pooled *Drosophila melanogaster*,
1254   sequencing ~80 pooled individuals from each of 271 populations as part of the
1255   European "Drosophila Evolution over Space and Time" (DEST) project. A total of
1256   5,019 shared SNPs with a genotyping rate of 0.937697 were used for analyses. The
1257   dataset, both raw and processed, is available through https://dest.bio.
1258
1259   -   Di Santo et al. (*60*) studied the highly-threatened species *Pinus torreyana*. They used
1260   Genotyping-by-Sequencing of 242 individuals of the last remaining populations. The
1261   dataset is not yet available through NCBI but the authors kindly shared a VCF directly
1262   with us. From a total set of 166,564 SNPs with a genotyping rate of 0.964632, 10,000
1263   were randomly selected for our analyses.
1264
-    von Seth et al. (*61*) studied the highly-threatened species *Dicerorhinus sumatrensis*.
They used Illumina WGS of 16 individuals of the last remaining populations. The raw
data is available at https://www.ebi.ac.uk/ena/browser/view/PRJEB35511. The
authors shared a VCF. In total, this comprises a set of 8,870,513 SNPs, with a
genotyping rate of 0.854862, which we did not further filter due to the small number
of individuals. For computational efficiency we selected 10,000 SNPs from the largest
chromosome.
1272
1273   Information and results per species are gathered in Table 1 and its extended version, Table
1274   S10, and the average z_{MAR} across species are provided in Table S11.
1275
1276

1281

Table S10 | The mutations-area relationship across species. Extended Table 1
The Mutations-Area Relationship (MAR) fitted with Area = Individuals and the scaled version. In the main text areas to
protect 90% of genetic diversity per species are provided given the scaled z*. Here, we also provide the average estimated
area based on % of grid cells per species to be transformed from 2015 to 2050 using the LUH² dataset, the area where at
least 10% of grid cells will be transformed, and the genetic loss corresponding to those area transformations (see section
_V.2)._
1288

Species (study)	SFS mod [ΔAIC]	MAR (A=N) z_N [CI95%]	MAR scaled z* [CI95%]	LUH² change '50	LUH² >10% change '50	LUH² extinct '50	LUH² >10% extinct '50
Arabidopsis thaliana (29)	logN (85.8)	0.431 (0.423 - 0.439)	0.312 (0.305–0.32)	4.58	13.54	1.12	3.43
Arabidopsis lyrata (42)	logN (9592.4)	0.254 (0.238 - 0.27)	0.15 (0.136–0.165)	0.79	2.64	0.19	0.64
Amaranthus tuberculatus (43)	logN (7317.5)	0.244 (0.237 - 0.251)	0.142 (0.135–0.148)	4.86	11.13	1.19	2.79
Eucalyptus melliodora (44)	logN (157.5)	0.531 (0.526 - 0.536)	0.402 (0.397–0.406)	3.82	7.77	0.93	1.92
Yucca brevifolia (58)	logN(33300)	0.141 (0.128 - 0.155)	0.049 (0.037–0.062)	0.74	0	0.18	0
Mimulus guttatus (45)	logN (580.8)	0.342 (0.331 - 0.353)	0.231 (0.221–0.241)	3.78	NA	0.92	NA
Panicum virgatum (46)	logN (8345.2)	0.226 (0.215 - 0.237)	0.126 (0.116–0.136)	8.07	27.65	2	7.47
Panicum hallii (57)	logN (86)	0.983 (0.907 - 1.059)	0.814 (0.745 - 0.883)	3.78	11.36	0.92	2.85
Pinus contorta (47)	Wei (19413.7)	0.019 (0.018 - 0.02)	-	1.95	5.54	0.47	1.36
Pinus torreyana (60)	logN(766156)	0.239 (0.232 - 0.245)	0.105 (0.099–0.11)	25.4	NA	6.79	NA
Populus trichocarpa (48)	logS (0)	0.268 (0.257 - 0.28)	0.164 (0.154–0.175)	4.68	17.28	1.14	4.45
Anopheles gambiae (49)	logS (0)	0.221 (0.209 - 0.233)	0.121 (0.11–0.132)	9.95	21.96	2.48	5.78
Acropora millepora (50)	logN (452.3)	0.403 (0.395 - 0.41)	0.287 (0.28–0.293)	72.73	84.69	26.79	36.26
Drosophila melanogaster (59)	logN(33300)	0.445 (0.433 - 0.458)	0.324 (0.313–0.336)	0.95	NA	0.23	NA
Empidonax traillii (51)	Wei (640401.9)	0.169 (0.139 - 0.199)	0.074 (0.047–0.101)	5.55	15.14	1.36	3.86
Setophaga petechia (52)	ln (67138.5)	0.251 (0.236 - 0.267)	0.149 (0.135 - 0.163)	2.83	7.54	0.69	1.86
Peromyscus maniculatus (53)	logN (1449.7)	0.844 (0.769 - 0.919)	0.68 (0.613–0.748)	5.61	13.68	1.38	3.47
Dicerorhinus sumatrensis (61)	w (107864.2)	0.474 (0.449 - 0.498)	0.123 (0.106–0.14)	0.25	NA	0.06	NA
Canis lupus (55)	logN (85.8)	0.29 (0.28 - 0.301)	0.183 (0.174–0.193)	0.23	NA	0.06	NA
Homo sapiens (56)	logN (9592.4)	0.395 (0.339 - 0.451)	0.28 (0.229–0.331)	28.81	40.13	7.83	11.58

1289   _Extended acronyms:_
1290   _logN: log Normal distribution. logS: log Series distribution. Wei: Weibull distribution._
1291

1292   **IV.1 Exclusion of species from global averages**
1293

1294   To avoid contaminating across-species averages of z_{MAR} with estimates of species whose data
1295   we do not fully trust, we conducted global averages excluding species for which we are not
1296   confident z_{MAR} reflects the correct species diversity-area relationships.
1297
1298   *Pinus contorta* showed a lower z_{MAR} than what is expected in a theoretical baseline from
1299   individual sampling (section II). This is most likely due to this being the only species for
1300   which SNPs were previously ascertained to be intermediate frequency (i.e. the genome
1301   technology was a SNP chip). This alters SFS, so we are not confident the z_{MAR} is the true
1302   parameter of the species.
1303
1304   *Yucca brevifolia* was a dense sampling of several local populations within a constrained area
1305   that is a hybrid zone. Since this species was not sampled range-wide we do not feel confident
1306   to include it in downstream analyses. The species also has a lower z than expected (Fig. S5)
1307
1308   *Pinus torreyana* only has two wild populations left, and therefore the MAR is based on two
1309   area sizes (Fig. S22). Because this is such a threatened species with already most of its range
1310   loss, we do not have confidence in the z parameter.
1311
1312   *Dicerorhinus sumatrensis* has only ~30 estimated adult individuals in the wild. Again we do
1313   not have confidence in the z parameter in such extinction-edge cases.
1314
1315   *Homo sapiens*. We exclude our own species.
1316

1317   ***Table S11 | Mean z_{MAR} and other summary statistics across species.***
1318   *We selected those species that did not show artefacts in Fig. S22 or whose z_{MAR} overlapped with 0 to calculate a species-*
1319   *wide mean. See section IV.1.*
1320

	z_{MAR}	z_{MAR} *(A=N)*	$z*_{MAR}$ *scaled*
mean	0.31	0.39	0.27
mean se	0.038	0.053	0.048
median	0.25	0.29	0.18
IQR	0.15	0.19	0.17

1321
1322
Although we could not see any obvious patterns relating z_{MAR} with certain groups of
species (Table 1), we wondered whether any life history trait of the species analysed could
explain the variation we observed (see Table S12 of traits). An ANOVA did not show any
significant relationship. Because we know theoretically this parameter must be related to the
degree of dispersal ability of genotypes of a species relative to the whole species geographic
range, we expect traits involved in determining these to be good predictors. Future work will
be necessary to validate this, as the sample size (n=19) may not permit enough power to
detect these expected patterns.
1331

1332   ***Table S12 | Traits, life history, and other characteristics of the analyzed species.***

Species	RedList	Known Decline	Kingdom	Reproduction	Pollination	Mobility	AreaRange
Arabidopsis thaliana	NO	NO	Plantae	Selfing	Selfing	Sessile	27337467.4
Arabidopsis lyrata	NO	NO	Plantae	Outcrossing	Vector	Sessile	2791301.4
Amaranthus tuberculatus	LC	NO	Plantae	Outcrossing	Vector	Sessile	804124.8
Eucalyptus melliodora	VU	NO	Plantae	Outcrossing	Wind	Sessile	948699.3
Yucca brevifolia	LC	YES	Plantae	Outcrossing	Vector	Sessile	1213454.4
Mimulus guttatus	LC	NO	Plantae	Outcrossing	Vector	Sessile	25138310.6
Panicum virgatum	LC	NO	Plantae	Outcrossing	Wind	Sessile	6291400.2

Panicum hallii	NO	NO	Plantae	Outcrossing	Wind	Sessile	2188807.4
Pinus contorta	LC	NO	Plantae	Outcrossing	Wind	Sessile	886182.2
Pinus torreyana	CR	YES	Plantae	Outcrossing	Wind	Sessile	30781.95
Populus trichocarpa	LC	NO	Plantae	Outcrossing	Wind	Sessile	1119664.1
Drosophila melanogaster	NO	NO	Animalia	Outcrossing	Activemating	Fly	115208408
Anopheles gambiae	NO	NO	Animalia	Outcrossing	Activemating	Fly	19959809.9
Acropora millepora	NT	YES	Animalia	Outcrossing	Activemating	Fly	26725.9
Empidonax traillii	LC	YES	Animalia	Outcrossing	Activemating	Fly	7027395.2
Setophaga petechia	LC	NO	Animalia	Outcrossing	Activemating	Fly	15172431.15
Peromyscus maniculatus	LC	NO	Animalia	Outcrossing	Activemating	Mobile	22609152.6
Dicerorhinus sumatrensis	CR	YES	Animalia	Outcrossing	Activemating	Mobile	3335605.58
Canis lupus	LC	NO	Animalia	Outcrossing	Activemating	Mobile	19102403.5
Homo sapiens	NA	NA	NA	NA	NA	NA	80763121.8

1333

1334

1335 ***Table S13 | Association of traits, life history, and other characteristics with z_{MAR}.***
1336 Acronyms: NO=not assessed but likely non-threatened, LC=low concern, VU=vulnerable, CR=critically endangered

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
RedList	4	0.0952396	0.0238099	0.5580988	0.7040464
KnownDecline	1	0.0275537	0.0275537	0.6458527	0.4580865
Kingdom	1	0.0011684	0.0011684	0.0273876	0.8750400
Reproduction	1	0.0003238	0.0003238	0.0075890	0.9339612
Pollination	1	0.0375975	0.0375975	0.8812784	0.3909509
Mobility	1	0.1600627	0.1600627	3.7518370	0.1104995
AreaRange	1	0.0174745	0.0174745	0.4095989	0.5503439
Residuals	5	0.2133125	0.0426625	NA	NA

1337

1338

1339   While no association between life history and z_{MAR} was found (Table S13), this may
1340 be due to limited power, as the sample size of species analysed here is still small, n=20.
1341 Further studies expanding the numbers of species will be necessary to confirm or reject this
1342 expected association.

1343

1344

1345

1346

V. An estimate of global genetic diversity loss

Using the approach described in section II.4, we generated a number of estimates either per ecosystem or per species. All estimates below tried to be conservative, and thus we always used the scaled z_{MAR} values (section II.3.2.)

V.1 Estimates of ecosystem area losses

Table S14 | Millennium Ecosystem Assessment land cover transformation.
Changes of ecosystem area pre-21st century. Ecosystem names are repeated for ecosystem sub-classes.
Source: https://www.millenniumassessment.org

System	Area (km² x10⁶)	Earth % surface	Protected areas (%)	Area transformed (%)
MARINE	349.3	68.6	0.3	NA
COASTAL	17.2	4.1	7	NA
- TERRESTRIAL	6	4.1	4	11
- MARINE	11.2	2.2	9	NA
INLAND WATER	10.3	7	12	11
FOREST/WOODLAND	41.9	28.4	10	42
- TROPICAL	23.3	15.8	11	34
- TEMPERATE	6.2	4.2	16	67
- BOREAL	12.4	8.4	4	25
DRYLAND	59.9	40.6	7	18
- HYPERARID	9.6	6.5	11	1
- ARID	15.3	10.4	6	5
- SEMIARID	22.3	15.3	6	25
- SUBHUMID	12.7	8.6	7	35
ISLAND	7.1	4.8	17	17
- STATES	4.7	3.2	18	21
MOUNTAINS	35.8	24.3	14	12
- 300-1000	13	8.8	11	13
- 1000-2500	11.3	7.7	14	13
- 2500-4500	9.6	6.5	18	6
- 4500+	1.8	1.2	22	0.3
POLAR	23	15.6	42	0.38
CULTIVATED	35.3	23.9	6	47
- PASTURE	0.1	0.1	4	11
- CROPLAND	8.3	5.7	4	62
- MIXED	26.9	18.2	6	43
URBAN	3.6	2.4	0	100
GLOBAL	510	NA	4	38

Ecosystem transformation has been tracked over decades. We extracted ecosystem transformations from the Millennium Ecosystem Assessment (*62*), which estimated ecosystem transformations from presumably native systems to cultivated or urban areas by GLC2000 land cover dataset (Table S14). The forest/woodland is calculated as percentage change between potential vegetation from WWF ecoregions to the current actual forest/woodland areas from GLC2000. These provide bulk ecosystem reductions, not for a given species, but may be a good proxy for an average across species.

Table S15 | IPBES land cover transformation.
Source: https://ipbes.net

Region	Area(Mkm2)	MSA_2010	MSA_2050_SSP2	MSA_2050_SSP1	MSA_2050_SSP3
North America	20	65	56	NA	NA
Central and South America	18	65	53	NA	NA
Middle East and Northern Africa	11	81	77	NA	NA
Sub-Saharan Africa	24	70	56	NA	NA
Western and Central Europe	6	37	29	NA	NA
Russian region and Central Asia	21	73	65	NA	NA

South Asia	5	44	35	NA	NA
China region	11	56	49	NA	NA
Southeast Asia	7	55	43	NA	NA
Japan, Korea and Oceania	8	71	57	NA	NA
Polar	2	96	91	NA	NA
World	132	66	56	62	54

The Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES) recently used a PBL satellite product from the Netherlands Environmental Assessment Agency (https://www.pbl.nl/en/nature-and-biodiversity) to study the % of area ecosystem transformation in the world (Table S15). This provides an updated estimate to the Millennium Assessment as well as projections under several Shared Socioeconomic Pathways (1-3) for 2050. These were reported per region as of 2010, and for projections to 2050 (scenario SSP2). Instead of direct area, the metric is a composite of land use information to predict Mean Species Abundance (MSA), a measure of the size of populations of wild organisms as a percentage of their inferred abundance in their natural state (% MSA).

A global transformation metric can also be captured by the most updated land use transformation data, the Land Use Harmonization 2 (release v2e for 2015-2011 and release v2h for baseline 1850-2015) (*63*). Baseline transformation of primary ecosystems was calculated subtracting the total area covered by primary forest (primf) and primary non-forest (primn) variables between year 1850 layer (roughly pre-industrial baseline) and the present, 2015, as $1-A_{2015}/A_{1850}$ (Table S16). Analyses that use projections to mid-21st century were conducted similarly as in (*64*), summing over all transitions from primary forest (primf), primary non-forest (primn), secondary forest (secdf) and secondary non-forest (secdn) lands to any other category for all years within the 2015-2050 period (see Table S10).

Table S16 | Land Use Harmonization 2 from 1850 to 2015
Source: https://luh.umd.edu/data.shtml

	Area %
Primary forest transformed	43
Primary non-forest transformed	50

Finally, we searched for timely estimates of forest reduction (based on vegetation cover) reported in the Global Forest Watch website: https://www.globalforestwatch.org/dashboards/global/ (accessed June 2021). From 2002 to 2020, there has been a global tree cover loss of 10%, with an annual tree cover loss of 0.6-1.1%.

Although these are not direct area transformations, we also used the IUCN Red List resource (https://www.iucnredlist.org, Table S12 shows status of the species analysed here), which includes guides to categorise species as vulnerable, endangered, critically endangered, and extinct, and has conducted extensive assessments across thousands of species (Table S17).

Table S17 | IUCN Red List categories of extinction risk and number of species.
Source: www.iucnredlist.org, January 2021

IUCN Red List	**Description**	**Criterion of area or pop.**	**# plant species**

Category		reduction (>%)	
EX	Extinct	100	164
EW	Extinct in the Wild	100	
CR	Critically Endangered	80	4674
EN	Endangered	50	8593
VU	Vulnerable	30	8459
NC, LR, NT, DD,LC	No Concern, Low Risk, Near Threatened, Data Deficient, Least Concern, Other	0	32237

1417
1418

V.2 A global estimate of genetic loss

1419

1420

1421 Taking the estimates and standard error of z_{MAR} across species, and the world's reduction of
1422 ecosystems we can calculate the fraction of genetic diversity reduction following the MAR
1423 equation (section II.4), giving a range of estimates (Table S18).
1424

1425

Fig. S23 | The parameter space of genetic diversity loss, extended

(A) The theoretical space of genetic diversity loss. z_{MAR} values (using area, unscaled for samples, differently from Fig. 32) computed for species analyzed here are marked as orange vertical lines, with confidence intervals as orange shading. Blue horizontal lines correspond to ecosystem transformations from the Millennium Assessment (light blue) and IPBES Assessment (dark blue) (B) Density histogram of percentages of area transformed across ecosystems from the MA, with averages per ecosystem marked in the distribution as well as horizontal lines in (A). (C) The number of species of each of the IUCN categories and the most optimistic range of area or abundance reduction for each of the category brackets.

Table S18 | Estimates of average expected genetic loss for different ecosystems.

Assuming ecosystem transformation approximately translates into average species distribution reduction, and using the ranges of z_{MAR} from Table 1 of the main text, we project the average genetic loss using the Mutations Area Relationship.

System	Area transformed (%)	Genetic loss % (mean z based)	Genetic loss % (min z based)	Genetic loss % (max z based)

COASTAL TERRESTRIAL	*11*	*3.2*	*0.9*	*9*
INLAND WATER	*11*	*3.2*	*0.9*	*79.7*
FOREST/WOODLAND	*42*	*14.0*	*4*	*35.8*
FOREST/WOODLAND TROPICAL	*34*	*10.5*	*3*	*28.7*
FOREST/WOODLAND TEMPERATE	*67*	*26.5*	*7.9*	*59.4*
FOREST/WOODLAND BOREAL	*25*	*7.7*	*2.1*	*20.9*
DRYLAND	*18*	*5.4*	*1.5*	*14.9*
DRYLAND HYPERARID	*1*	*0.3*	*0.1*	*0.8*
DRYLAND ARID	*5*	*1.4*	*0.4*	*4.1*
DRYLAND SEMIARID	*25*	*7.7*	*2.1*	*20.9*
DRYLAND SUBHUMID	*35*	*11.3*	*3.2*	*29.6*
ISLAND	*17*	*5.0*	*1.4*	*14.1*
MOUNTAINS	*12*	*3.5*	*0.9*	*9.9*
MOUNTAINS 300-1000	*13*	*3.8*	*1*	*10.7*
MOUNTAINS 1000-2500	*13*	*3.8*	*1*	*10.7*
MOUNTAINS 2500-4500	*6*	*1.7*	*0.5*	*4.9*
MOUNTAINS 4500+	*0.3*	*0.1*	*0*	*0.2*
POLAR	*0.4*	*0.1*	*0*	*0.3*
GLOBAL	*38*	*12.4*	*3.5*	*32.2*

Assuming the average z_{MAR}, and utilising tree cover from the Global Forest Watch (https://www.globalforestwatch.org), which estimates 0.6-1.1% of transformation per year across Canada, United States and Australia, we extrapolated genetic diversity loss in the next 50 years for tree species to be 8-15% genetic diversity loss.

Assuming that the calculated z_{MAR} estimates (Table 1) are representative of plant species, we conducted an experiment to create a distribution of % of genetic diversity loss in threatened species. We used the number of species in each IUCN category (Table S17) for a total of 54,127 plant species. For plant species, one of the evaluation criteria of percentage of population loss likely translates faithfully to area reduction in the species. Thus, the proportion of species per category gives a discrete probability distribution of the ranges of percentage of area loss: P(0-29%)=0.596, P(30-49%)=0.156, P(50-79%)=0.159, P(80-99%)=0.086, P(99%-100%)=0.003. Using a simulation-based sampling approach, we drew 350,000 random area reductions A_t / A_{t-1} from the previous distribution and a z_{MAR} from the mean and variance of our estimates from Table 1 for plants. These were plugged into the MAR equation (Section II.4) to calculate the percentage of genetic diversity loss of these 350,000 random draws. The resulting distribution had a median and interquartile range of 17.53 % [7.51- 31.82]..

Using the Land Use Harmonization 2 dataset, we also create per-species predictions based on the % transformation of each of the sampled regions per species (Table S14). As before, the land use transformations that merit be considered area losses are all transitions from primary forest (primf), primary non-forest (primn), secondary forest (secdf) and secondary non-forest (secdn) lands to any other category. Taking all the locations where each species has been sampled, we extracted the predicted % of land use change per cell and summed over all cells where individuals had been sampled (we call this LUH[2] change '50, see column in Table S10). We also produced the alternative area loss estimate taking that at least 10% predicted habitat transformation for a grid cell renders the entire area of that grid cell as impacted or lost (we call this LUH[2] >10% change '50). These per-species area losses, in combination with the matched z_{MAR}, provided a range of potential loss estimates to 2050 ranging 0-36% depending on the species (Table S10).

1473

1474   **V.3 Community ecology simulations and MAR**

1475

To test whether intermediate levels of MAR would be expected across species in entire
ecosystems, we conducted community assembly simulations of ~100-500 species following
the Neutral Theory of Biodiversity (*1, 41*) and coalescent simulations (*23*) using the software
MESS (*65*). These simulations are computationally demanding and could not run in a
complete 2D spatial grid. Instead, they were simulated in a mainland-island system, with
islands of increasing areas. The community forms by species colonising an empty island
according to Hubbell's Unified Neutral Theory of Biodiversity and Biogeography (UNTB),
where all species are equally likely to colonise and persist in the local community. Continued
colonisation and migration to the local community continues to bring in new species that may
or may not survive, while also continuously bringing in individuals of species already in the
local community. The community assembly process ends when the community has reached
an equilibrium denoted as the balance between local extinction and new species dispersing
into the area (Hubbell 2001). Once the forward-time process has ended, we simulate the
coalescent history of each species backward in time. For this, MESS considers the population
size, divergence time, and migration rates of the meta and local communities. These
coalescent simulations provide us with genetic data and ultimately diversity estimates for
each species in the community.

1493

We simulated 100 MESS communities, and for each community the size of the local
community was varied from 1K to 100K. We varied the size of communities to emulate
variation in area occupied by a given community because we assume as the number of
individuals in a community increases from 1,000 to 100,000, so does the area occupied. All
other parameters were kept consistent across each of these community simulations, and most
remained at their default value. The parameters changed were the length of the sequences
simulated for the coalescent-based simulations, which was fixed at 10,000 bp, and the
migration rate, which was fixed at 0.01.

1502

1503   The simulation output was used to then compute a single z_{SAR} for the system as
1504 $S=cA^{z_{SAR}}$, and one z_{MAR} for each species in the same way, $M=cA^{z_{MAR}}$. This resulted in the
1505   distribution of z_{MAR} from Fig. S24. This confirmed that we can recover typical z_{SAR} and z_{MAR}
1506   values from completely stochastic neutral yet spatially structured systems such as species in
1507   communities and mutations in populations of a species.

1508

1509

Fig. S24 | z_{MAR} calculated from MESS eco-evolutionary simulations
Using the MESS framework of a mainland-island model with different island sizes, z_{MAR} per species is recovered. The
stochastic nature of the simulations results in each species having different abundances and migration histories that change

the scaling value. Values were typically around 0.3. Rarely some species had values above 1, which appear could be noisy
estimates from recently colonising species in the simulations.
.
1516
1517

V.4 The nested species extinction and genetic diversity loss processes

1518
1519
Finally, we worried that our estimates of V.2 would be mistaken as overestimates. In fact, we
believe these may be underestimated. Recent policy proposals for the United Nations'
Sustainability Goals emphasize that the target of protecting 90% of species genetic diversity
for all species cannot leave the already-extinct species behind (*66*) (That is, one cannot
protect 90% of species and leave 10% to become extinct to meet this goal). This clearly
exemplifies a problem in conservation biology that what researchers can study is (most of the
time) what has escaped extinction, and therefore if we do not account for extinct species in
our overall estimates of genetic diversity loss we may naively think ecosystems have not
suffered genetic diversity loss (i.e. in the extreme scenario, an ecosystem that has lost all but
one abundant species may not really appear genetically eroded if such species is in good
shape).

1531
1532 We then created spatial simulations in R where 1,000 species are distributed in
1533 100x100 grid cells following a UNTB abundance distribution and then proceeded with an
1534 edge extinction of the ecosystem (see Fig. S25 for a cartoon).

1535
1536
1537

1538
Fig. S25 | Cartoon of nested extinction of species and genetic diversity loss.
An ecosystem with multiple species within it (left), distributed in space, with few species broadly distributed and many
narrowly distributed. Moving one level of biological organization lower, mutations within species (right) are also spatially
distributed with many narrowly distributed. As extinction happens (red line moving bottom to top), all species below the red
line go extinct, but only the mutations within species 1 below the line are lost, while mutations above the line remain.
Species 3 has already become extinct, and therefore also all the mutations within it.
1545
1546
Two extreme types of distributions of species can be imagined: species are randomly
placed in space, or species are found mostly in perfectly contiguous ranges (We ended up
using as an example a simulation with 85% of the individuals of a species found in a core
square continuous distribution and 15% found outside that core in fragmented observations,
as this scenario produced the canonical SAR of *z*~0.3). Spatial structure interestingly creates
two extreme distributions of area reductions across species (Fig. S26): random placement of
cell habitats essentially show that the average area reduction per ecosystem is followed by
most species, while autocorrelated placement of cell habitats create a U distribution in area
reductions, where at the beginning of the extinction process most species have not

experienced any impact (Fig. S26B left) but at the end of ecosystem reduction virtually all
species are already extinct (Note we may be at the beginning of S26B process given the data
from IUCN, Fig 3C).
1559
1560

1561

Fig. S26 | The distribution of per-species area lost and total ecosystem extinction with 1000 species
Two ecosystems of 100x100 cells with 1000 species. Species are either randomly distributed in cells (A) or spatially
autocorrelated with occupying mostly contiguous cells (B). As the extinction process wipes out part of the ecosystem
(snapshots are provided at 5%, 50%, and 95%), the area loss per species (and hence genetic diversity lost) is tracked. In (A)
the average area lost per species is roughly the total reduction of the ecosystem, whereas in (B) the distribution is U shaped
(note the log-scaled y-axis). While in (B) the mean area lost in the distribution correctly captures the area loss of the
ecosystem, per species losses are highly uneven.
1569

To study the consequence of the above differential area loss and the effect of some
species going extinct on the total ecosystem genetic diversity, we conducted the next
analysis: For extant species, we assumed they would lose genetic diversity following the
MAR relationship (section II.4), with all species having $z_{MAR} = 0.3$ for simplicity (i.e. all
species lose genetic diversity at the same rate). For extinct species (100% of their area
reduced), we considered genetic diversity loss was 100%. The compound total genetic
diversity loss would then just be the sum of those $X_{Tot} = \sum_{i=1}^{1000} X_i$ (Of course, in reality species
may vary in their genome-wide diversity average, and we could for instance use Watterson's
Θ_W (see section II.2) to scale the total loss of genetic diversity in the ecosystem accounting
for different basal level of diversity per species: $\sum_{i=1}^{1000} \Theta_{Wi} X_i$). Interestingly, if we calculate the
z of the slope of compound genetic diversity across species in an ecosystem it is much larger
than MAR or SAR alone: $z_{compunded} = 0.6$ (Fig. S27).
1582

1583

Fig. S27 | Numeric simulation of nested species and genetic diversity loss.
(A) Simulating the extinction of an ecosystem with 1,000 species that follow a log-normal species abundance curve.
Extinction of the ecosystem creates a curve of species loss of z~0.3 (grey). Likewise, each species trajectory (light red, 15
species drawn randomly) follows a simulated genetic diversity loss of z_{MAR}~0.3 as they lose area. Because species'
geographic distributions are by construction smaller than the whole ecosystem area, those distributed closer to the start of
the extinction front lose area first, while those distributed farthest from the extinction front only lose area when the
ecosystem is almost completely destroyed. Because genetic diversity loss is both due to complete extinction of species as well
as area reduction of extant species, the compound genetic diversity loss curve (red) follows the faster loss dynamics. (B)
Holding z_{SAR}=0.3 constant, and varying z_{MAR} in independent simulations shows that the compound genetic diversity across
species is close to the sum of both z slopes (the SAR and the MAR), but it saturates at ca. 0.85 (grey dotted line shows z_{MAR}
+ z_{SAR}).

1595

1596

1597

VI. Limitations and outlook

In this last section we list some potential limitations of an inherently simple scaling law, and what approaches could be used to address those and improve genetic diversity loss projections.

VI.1 Reasons for overestimations

Many researchers have posited that SAR likely overestimates species extinction (*33*, *67*). For instance:

- Ignoring that a diversity-area relationship can be defined outwards, inwards, or focusing on endemisms can have an impact (*10*, *33*, *67*). To address this, we confirmed relative consistency between inward, outward, and random placement MAR, and proposed that the EMAR may not be that appropriate to study genetic diversity loss (or at least EMAR does not show predictability in our simulation).
- Species may persist in altered habitats, like some animals are known to do (*68*). We have focused some of the estimates in this study on plants, for which area loss should equate to population loss and vice versa, but further extensions could be applied in the future as described by Pereira and Daily (*68*).
- SAR is not a mechanistic model (*69*). We have derived its ranges of possible values and averages analytically and are beginning to understand how evolutionary forces shape MAR. Realistic simulations can help understand in a process-based framework how populations (and their MAR) react to partial population extinction (continuous space simulations with progressive area reductions appear to fit well with the MAR predictions calculated before the extinction process starts, section III.2.6).
- There is a scale dependence in the SAR slope, with slight increase in the slope at large scales (*10*). Since power laws are typically fit with large-scale datasets and used to predict local scale extinctions, predictions could be overestimated at local scales.

VI.2 Reasons for underestimations

While the simplicity of power laws to make predictions of species extinction may lead to overestimations, there are also important reasons to believe MAR would underestimate genetic loss.

- Perhaps even more so than in species list datasets and census, the discovery of low frequency genetic variants is highly underpowered (*70*). These are highly prevalent, but genomic pipelines, with the aim to be conservative, often filter out rare variants. This would underestimate z_{MAR} and therefore the degree of genetic diversity loss with area shrinkage. This is clear in the pre-selected-only marker dataset of *Pinus contorta*.
- Related to the previous: Although sequencing methods have an error rate that misreads true nucleotide sequences, this rate is typically extremely low (many sequencing projects described here used Illumina HiSeq series, which has a 0.112% error rate, or about 1 misread nucleotide in 1000). This could intuitively lead to overestimates in mutations in space but in fact, the mis-reading of DNA ends up causing an underestimation. This is because bioinformatic software that transforms raw data into SNP variant tables errs towards the conservative direction, often not calling mutations that have been observed very few times, and thus likely under-representing rare mutations (*71*).
- The use of scaled z_{MAR} proposed in section II.3.2. accounts for that the minimum z_{MAR} is rarely exactly 0, especially when sample sizes are limited. We use this correction

scaling down z_{MAR} to be conservative. However, z_{MAR} could only in very exceptional
circumstances be 1, but we do not correct for this, again, to have a conservatively low
z_{MAR}. Hence, our conservative approach would generally lead to underestimates of
genetic diversity loss.
-   When species shrink in area, the effective population size of the remaining population
decreases, increasing drift and moving towards a lower diversity equilibrium. This
reactive process is not captured by the phenomenological MAR relationship.
-   The nested extinction of species and genetic diversity loss (section V.3) would lead
us, by the right of "survival bias", to underestimate how much genetic diversity has
been lost cumulative in an ecosystem.
1658
1659 **VI.3 Final notes**
1660
Ultimately, to make accurate predictions of genetic diversity loss and increased extinction
risk of species, very detailed data and expert assessment per species will be required: census
sizes, genome size, migration in metapopulations, mating system, detailed maps of genetic
makeups, and finescale area transformations. This could enable mechanistic models projected
forward-in-time such as discussed in section II.3.6. The production of new genomic datasets
across entire ecosystems should further help create maps of genetic diversity at high
resolution to track losses (*72–74*).
1668
Our philosophy in this work has been to err on the conservative side when projecting
genetic diversity loss (e.g. using area calculations that produce lower z_{MAR} values, scaling
them for low sample bias, using lower estimates of ecosystem transformation, etc.). However,
this conservative approach can also lead us into under-estimating loss. As described in V.4.,
the phenomenon of survival bias likely leads us to underestimate what has been lost given we
do not observe it. A phenomenon also highlighted as a possible explanation for the relatively
shy difference in genetic diversity between threatened and non-threatened species (*75, 76*)
1676
Because to our knowledge, no other approaches exist to project genetic diversity, we
believe that MAR is a quantitative and scalable first-approximation of genetic diversity that
would just require accurate understanding of abundance or area reductions and minimal
information about population structure or mating/dispersal/range relationships. Given that
scaling relationships are already applied by conservation policy (*77*), and given that
assumptions and limitations are understood, we expect MAR to become a relevant tool to
project losses of a dimension of biodiversity so far mostly invisible or unaddressable in large
conservation projections.
1685
1686
1687

SUPPLEMENTAL REFERENCES

1. S. P. Hubbell, *The Unified Neutral Theory of Biodiversity and Biogeography* (Monographs in Population Biology, 2001).

2. M. Tokeshi, Niche Apportionment or Random Assortment: Species Abundance Patterns Revisited. *J. Anim. Ecol.* **59**, 1129–1146 (1990).

3. M. Tokeshi, in *Advances in Ecological Research*, M. Begon, A. H. Fitter, Eds. (Academic Press, 1993; https://www.sciencedirect.com/science/article/pii/S0065250408600422), vol. 24, pp. 111–186.

4. MOTOMURA, I, A statistical treatment of ecological communities. *Zoological Magazine*. **44**, 379–383 (1932).

5. R. H. MacArthur, On the relative abundance of bird species. *Proc. Natl. Acad. Sci. U. S. A.* **43**, 293–295 (1957).

6. R. A. Fisher, A. S. Corbet, C. B. Williams, The Relation Between the Number of Species and the Number of Individuals in a Random Sample of an Animal Population. *J. Anim. Ecol.* **12**, 42–58 (1943).

7. F. W. Preston, The canonical distribution of commonness and rarity: Part I. *Ecology*. **43**, 185 (1962).

8. F. W. Preston, The commonness, and rarity, of species. *Ecology*. **29**, 254–283 (1948).

9. D. Alonso, A. J. McKane, Sampling Hubbell's neutral theory of biodiversity. *Ecol. Lett.* **7**, 901–910 (2004).

10. D. Storch, P. Keil, W. Jetz, Universal species-area and endemics-area relationships at continental scales. *Nature*. **488**, 78–81 (2012).

11. S. L. Pimm, G. J. Russell, J. L. Gittleman, T. M. Brooks, The future of biodiversity. *Science*. **269**, 347–350 (1995).

12. C. D. Thomas, A. Cameron, R. E. Green, M. Bakkenes, L. J. Beaumont, Y. C. Collingham, B. F. N. Erasmus, M. F. de Siqueira, A. Grainger, L. Hannah, L. Hughes, B. Huntley, A. S. van Jaarsveld, G. F. Midgley, L. Miles, M. A. Ortega-Huerta, A. Townsend Peterson, O. L. Phillips, S. E. Williams, Extinction risk from climate change. *Nature*. **427**, 145–148 (2004).

13. J. F. C. Kingman, The coalescent. *Stochastic Process. Appl.* **13**, 235–248 (1982).

14. R. C. Griffiths, S. Tavaré, The age of a mutation in a general coalescent tree. *Communications in Statistics. Stochastic Models*. **14**, 273–295 (1998).

15. R. A. Fisher, XVII.—The Distribution of Gene Ratios for Rare Mutations. *Proceedings of the Royal Society of Edinburgh*. **50**, 204–219 (1931).

16. M. W. Hahn, *Molecular Population Genetics* (Oxford University Press, 2018; https://play.google.com/store/books/details?id=3BDkswEACAAJ).

17. V. Buffalo, Quantifying the relationship between genetic diversity and population size suggests natural selection cannot explain Lewontin's paradox. *Elife*. **10** (2021), doi:10.7554/eLife.67509.

1727   18.   D. R. Marshal, A. D. H. Brown, in *Crop genetic resources for today and tomorrow*, O. H.
1728   Frankel, J. G. Hawkes, Eds. (Cambridge University Press, 1975;
1729   https://www.researchgate.net/publication/280057199), vol. 2.

1730   19.   S. Wright, Isolation by Distance. *Genetics*. **28**, 114–138 (1943).

1731   20.   J. Novembre, M. Stephens, Interpreting principal component analyses of spatial population
1732   genetic variation. *Nat. Genet.* **40**, 646–649 (2008).

1733   21.   H. Fan 樊海英, Q. Zhang 张清臣, J. Rao 饶娟娟, J. Cao 曹静文, X. Lu 卢欣, Genetic Diversity-
1734   Area Relationships across Bird Species. *Am. Nat.* **194**, 736–740 (2019).

1735   22.   P. R. A. Campos, V. M. de Oliveira, A. Rosas, Epistasis and environmental heterogeneity in the
1736   speciation process. *Ecol. Modell.* **221**, 2546–2554 (2010).

1737   23.   J. Kelleher, A. M. Etheridge, G. McVean, Efficient Coalescent Simulation and Genealogical
1738   Analysis for Large Sample Sizes. *PLoS Comput. Biol.* **12**, e1004842 (2016).

1739   24.   B. C. Haller, P. W. Messer, SLiM 3: Forward Genetic Simulations Beyond the Wright–Fisher
1740   Model. *Mol. Biol. Evol.* **36**, 632–637 (2019).

1741   25.   T. R. Booker, S. Yeaman, M. C. Whitlock, The WZA: A window-based method for
1742   characterizing genotype-environment association. *bioRxiv* (2021), p. 2021.06.25.449972.

1743   26.   J. Kelleher, K. R. Thornton, J. Ashander, P. L. Ralph, Efficient pedigree recording for fast
1744   population genetics simulation. *PLoS Comput. Biol.* (2018), p. e1006581.

1745   27.   B. C. Haller, J. Galloway, J. Kelleher, P. W. Messer, P. L. Ralph, Tree-sequence recording in
1746   SLiM opens new horizons for forward-time simulation of whole genomes. *Mol. Ecol. Resour.*
1747   **19**, 552–566 (2019).

1748   28.   R. R. Hudson, M. Slatkin, W. P. Maddison, Estimation of levels of gene flow from DNA
1749   sequence data. *Genetics*. **132**, 583–589 (1992).

1750   29.   1001 Genomes Consortium, 1,135 Genomes Reveal the Global Pattern of Polymorphism in
1751   Arabidopsis thaliana. *Cell*. **166**, 481–491 (2016).

1752   30.   B. J. McGill, B. J. Enquist, E. Weiher, M. Westoby, Rebuilding community ecology from
1753   functional traits. *Trends Ecol. Evol.* **21**, 178–185 (2006).

1754   31.   P. I. Prado, M. Miranda, A. Chalom, *sads: R package for fitting species abundance distributions*
1755   (Github, 2018; https://github.com/piLaboratory/sads).

1756   32.   T. J. Matthews, K. A. Triantis, R. J. Whittaker, F. Guilhaumon, sars: an R package for fitting,
1757   evaluating and comparing species–area relationship models. *Ecography*. **42**, 1446–1455 (2019).

1758   33.   F. He, S. P. Hubbell, Species-area relationships always overestimate extinction rates from habitat
1759   loss. *Nature*. **473**, 368–371 (2011).

1760   34.   R. J. H. &. J. van Etten, raster: Geographic analysis and modeling with raster data (2012),
1761   (available at http://CRAN.R-project.org/package=raster).

1762   35.   M. Exposito-Alonso, 500 Genomes Field Experiment Team, H. A. Burbano, O. Bossdorf, R.
1763   Nielsen, D. Weigel, Natural selection in the *Arabidopsis thaliana* genome in present and future
1764   climates. *Nature*. **573**, 126–129 (2019).

36. Y. B. Simons, K. Bullaughey, R. R. Hudson, G. Sella, A population genetic interpretation of
GWAS findings for human quantitative traits. *PLoS Biol.* **16**, e2002985 (2018).

37. M. Exposito-Alonso, C. Becker, V. J. Schuenemann, E. Reiter, C. Setzer, R. Slovak, B. Brachi, J.
Hagmann, D. G. Grimm, J. Chen, W. Busch, J. Bergelson, R. W. Ness, J. Krause, H. A. Burbano,
D. Weigel, The rate and potential relevance of new mutations in a colonizing plant lineage. *PLoS*
Genet. **14**, e1007155 (2018).

38. H. Seebens, T. M. Blackburn, E. E. Dyer, P. Genovesi, P. E. Hulme, J. M. Jeschke, S. Pagad, P.
Pyšek, M. Winter, M. Arianoutsou, S. Bacher, B. Blasius, G. Brundu, C. Capinha, L. Celesti-
Grapow, W. Dawson, S. Dullinger, N. Fuentes, H. Jäger, J. Kartesz, M. Kenis, H. Kreft, I. Kühn,
B. Lenzner, A. Liebhold, A. Mosena, D. Moser, M. Nishino, D. Pearman, J. Pergl, W. Rabitsch,
J. Rojas-Sandoval, A. Roques, S. Rorke, S. Rossinelli, H. E. Roy, R. Scalera, S. Schindler, K.
Štajerová, B. Tokarska-Guzik, M. van Kleunen, K. Walker, P. Weigelt, T. Yamanaka, F. Essl,
No saturation in the accumulation of alien species worldwide. *Nat. Commun.* **8**, 14435 (2017).

39. H. Seebens, F. Essl, W. Dawson, N. Fuentes, D. Moser, J. Pergl, P. Pyšek, M. van Kleunen, E.
Weber, M. Winter, B. Blasius, Global trade will accelerate plant invasions in emerging
economies under climate change. *Glob. Chang. Biol.* **21**, 4128–4140 (2015).

40. S. Purcell, B. Neale, K. Todd-Brown, L. Thomas, M. A. R. Ferreira, D. Bender, J. Maller, P.
Sklar, P. I. W. de Bakker, M. J. Daly, P. C. Sham, PLINK: a tool set for whole-genome
association and population-based linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

41. C. Mérot, R. A. Oomen, A. Tigano, M. Wellenreuther, A Roadmap for Understanding the
Evolutionary Significance of Structural Genomic Variation. *Trends Ecol. Evol.* **35**, 561–572
(2020).

42. K. Lucek, Y. Willi, Drivers of linkage disequilibrium across a species' geographic range. *PLoS*
Genet. **17**, e1009477 (2021).

43. J. M. Kreiner, D. A. Giacomini, F. Bemm, B. Waithaka, J. Regalado, C. Lanz, J. Hildebrandt, P.
H. Sikkema, P. J. Tranel, D. Weigel, J. R. Stinchcombe, S. I. Wright, Multiple modes of
convergent adaptation in the spread of glyphosate-resistant Amaranthus tuberculatus. *Proc. Natl.*
Acad. Sci. U. S. A. **116**, 21076–21084 (2019).

44. M. A. Supple, J. G. Bragg, L. M. Broadhurst, A. B. Nicotra, M. Byrne, R. L. Andrew, A.
Widdup, N. C. Aitken, J. O. Borevitz, Landscape genomic prediction for restoration of a
Eucalyptus foundation species under climate change. *Elife.* **7** (2018), doi:10.7554/eLife.31835.

45. M. Vallejo-Marín, J. Friedman, A. D. Twyford, O. Lepais, S. M. Ickert-Bond, M. A. Streisfeld,
L. Yant, M. van Kleunen, M. C. Rotter, J. R. Puzey, Population genomic and historical analysis
suggests a global invasion by bridgehead processes in Mimulus guttatus. *Commun Biol.* **4**, 327
(2021).

46. J. T. Lovell, A. H. MacQueen, S. Mamidi, J. Bonnette, J. Jenkins, J. D. Napier, A. Sreedasyam,
A. Healey, A. Session, S. Shu, K. Barry, S. Bonos, L. Boston, C. Daum, S. Deshpande, A.
Ewing, P. P. Grabowski, T. Haque, M. Harrison, J. Jiang, D. Kudrna, A. Lipzen, T. H.
Pendergast 4th, C. Plott, P. Qi, C. A. Saski, E. V. Shakirov, D. Sims, M. Sharma, R. Sharma, A.
Stewart, V. R. Singan, Y. Tang, S. Thibivillier, J. Webber, X. Weng, M. Williams, G. A. Wu, Y.
Yoshinaga, M. Zane, L. Zhang, J. Zhang, K. D. Behrman, A. R. Boe, P. A. Fay, F. B. Fritschi, J.
D. Jastrow, J. Lloyd-Reilley, J. M. Martínez-Reyna, R. Matamala, R. B. Mitchell, F. M.
Rouquette Jr, P. Ronald, M. Saha, C. M. Tobias, M. Udvardi, R. A. Wing, Y. Wu, L. E. Bartley,
M. Casler, K. M. Devos, D. B. Lowry, D. S. Rokhsar, J. Grimwood, T. E. Juenger, J. Schmutz,
Genomic mechanisms of climate adaptation in polyploid bioenergy switchgrass. *Nature* (2021),
doi:10.1038/s41586-020-03127-1.

47.   I. R. MacLachlan, T. K. McDonald, B. M. Lind, L. H. Rieseberg, S. Yeaman, S. N. Aitken,
1812   Genome-wide shifts in climate-related variation underpin responses to selective breeding in a
1813   widespread conifer. *Proc. Natl. Acad. Sci. U. S. A.* **118** (2021), doi:10.1073/pnas.2016900118.

1814   48.   G. Tuskan, W. Muchero, J.-G. Chen, D. Jacobson, T. Tschaplinski, D. Rokhsar, W. Schackwitz,
1815   J. Schmutz, S. DiFazio, Populus Trichocarpa Genome-Wide Association Study (GWAS)
1816   Population SNP Dataset Released, (available at https://doi.ccs.ornl.gov/ui/doi/55).

1817   49.   Anopheles gambiae 1000 Genomes Consortium, Data analysis group, Partner working group,
1818   Sample collections—Angola:, Burkina Faso:, Cameroon:, Gabon:, Guinea:, Guinea-Bissau:,
1819   Kenya:, Uganda:, Crosses:, Sequencing and data production, Web application development,
1820   Project coordination, Genetic diversity of the African malaria vector Anopheles gambiae.
1821   *Nature*. **552**, 96–100 (2017).

1822   50.   Z. L. Fuller, V. J. L. Mocellin, L. A. Morris, N. Cantin, J. Shepherd, L. Sarre, J. Peng, Y. Liao, J.
1823   Pickrell, P. Andolfatto, M. Matz, L. K. Bay, M. Przeworski, Population genetics of the coral
1824   Acropora millepora: Toward genomic prediction of bleaching. *Science*. **369** (2020),
1825   doi:10.1126/science.aba4674.

1826   51.   K. Ruegg, E. C. Anderson, M. Somveille, R. A. Bay, M. Whitfield, E. H. Paxton, T. B. Smith,
1827   Linking climate niches across seasons to assess population vulnerability in a migratory bird.
1828   *Glob. Chang. Biol.* **27**, 3519–3531 (2021).

1829   52.   R. A. Bay, R. J. Harrigan, V. Le Underwood, H. Lisle Gibbs, T. B. Smith, K. Ruegg, Genomic
1830   signals of selection predict climate-driven population declines in a migratory bird. *Science*. **359**,
1831   83–86 (2018).

1832   53.   E. P. Kingsley, K. M. Kozak, S. P. Pfeifer, D.-S. Yang, H. E. Hoekstra, The ultimate and
1833   proximate mechanisms driving the evolution of long tails in forest deer mice. *Evolution*. **71**, 261–
1834   273 (2017).

1835   54.   L. Smeds, J. Aspi, J. Berglund, I. Kojola, K. Tirronen, H. Ellegren, Whole-genome analyses
1836   provide no evidence for dog introgression in Fennoscandian wolf populations. *Evol. Appl.* **14**,
1837   721–734 (2021).

1838   55.   R. M. Schweizer, J. Robinson, R. Harrigan, P. Silva, M. Galverni, M. Musiani, R. E. Green, J.
1839   Novembre, R. K. Wayne, Targeted capture and resequencing of 1040 genes reveal
1840   environmentally driven functional variation in grey wolves. *Mol. Ecol.* **25**, 357–379 (2016).

1841   56.   1000 Genomes Project Consortium, A. Auton, L. D. Brooks, R. M. Durbin, E. P. Garrison, H. M.
1842   Kang, J. O. Korbel, J. L. Marchini, S. McCarthy, G. A. McVean, G. R. Abecasis, A global
1843   reference for human genetic variation. *Nature*. **526**, 68–74 (2015).

1844   57.   J. D. Palacio-Mejía, P. P. Grabowski, E. M. Ortiz, G. A. Silva-Arias, T. Haque, D. L. Des
1845   Marais, J. Bonnette, D. B. Lowry, T. E. Juenger, Geographic patterns of genomic diversity and
1846   structure in the C4 grass Panicum hallii across its natural distribution. *AoB Plants*. **13**, lab002
1847   (2021).

1848   58.   A. M. Royer, M. A. Streisfeld, C. I. Smith, Population genomics of divergence within an obligate
1849   pollination mutualism: Selection maintains differences between Joshua tree species. *Am. J. Bot.*
1850   **103**, 1730–1741 (2016).

1851   59.   M. Kapun, J. C. B. Nunez, M. Bogaerts-Márquez, J. Murga-Moreno, M. Paris, J. Outten, M.
1852   Coronado-Zamora, C. Tern, O. Rota-Stabelli, M. P. García Guerreiro, S. Casillas, D. J. Orengo,
1853   E. Puerma, M. Kankare, L. Ometto, V. Loeschcke, B. S. Onder, J. K. Abbott, S. W. Schaeffer, S.
1854   Rajpurohit, E. L. Behrman, M. F. Schou, T. J. S. Merritt, B. P. Lazzaro, A. Glaser-Schmitt, E.

Argyridou, F. Staubach, Y. Wang, E. Tauber, S. V. Serga, D. K. Fabian, K. A. Dyer, C. W. Wheat, J. Parsch, S. Grath, M. S. Veselinovic, M. Stamenkovic-Radak, M. Jelic, A. J. Buendía-Ruíz, M. Josefa Gómez-Julián, M. Luisa Espinosa-Jimenez, F. D. Gallardo-Jiménez, A. Patenkovic, K. Eric, M. Tanaskovic, A. Ullastres, L. Guio, M. Merenciano, S. Guirao-Rico, V. Horváth, D. J. Obbard, E. Pasyukova, V. E. Alatortsev, C. P. Vieira, J. Vieira, J. Roberto Torres, I. Kozeretska, O. M. Maistrenko, C. Montchamp-Moreau, D. V. Mukha, H. E. Machado, A. Barbadilla, D. Petrov, P. Schmidt, J. Gonzalez, T. Flatt, A. O. Bergland, Drosophila Evolution over Space and Time (DEST) - A New Population Genomics Resource. *bioRxiv* (2021), p. 2021.02.01.428994.

60. L. N. Di Santo, S. Hoban, T. L. Parchman, J. W. Wright, J. A. Hamilton, Reduced representation sequencing to understand the evolutionary history of Torrey pine (Pinus torreyana Parry) with implications for rare species conservation. *bioRxiv* (2021), p. 2021.07.02.450939.

61. J. von Seth, N. Dussex, D. Díez-Del-Molino, T. van der Valk, V. E. Kutschera, M. Kierczak, C. C. Steiner, S. Liu, M. T. P. Gilbert, M.-H. S. Sinding, S. Prost, K. Guschanski, S. K. S. S. Nathan, S. Brace, Y. L. Chan, C. W. Wheat, P. Skoglund, O. A. Ryder, B. Goossens, A. Götherström, L. Dalén, Genomic insights into the conservation status of the world's last remaining Sumatran rhinoceros populations. *Nat. Commun.* **12**, 2393 (2021).

62. Millennium Ecosystem Assessment, *Millennium ecosystem assessment* (Millennium Ecosystem Assessment, 2005; http://chapter.ser.org/europe/files/2012/08/Harris.pdf).

63. G. C. Hurtt, L. Chini, R. Sahajpal, S. Frolking, B. L. Bodirsky, K. Calvin, J. C. Doelman, J. Fisk, S. Fujimori, K. Klein Goldewijk, T. Hasegawa, P. Havlik, A. Heinimann, F. Humpenöder, J. Jungclaus, J. O. Kaplan, J. Kennedy, T. Krisztin, D. Lawrence, P. Lawrence, L. Ma, O. Mertz, J. Pongratz, A. Popp, B. Poulter, K. Riahi, E. Shevliakova, E. Stehfest, P. Thornton, F. N. Tubiello, D. P. van Vuuren, X. Zhang, Harmonization of global land use change and management for the period 850–2100 (LUH2) for CMIP6. *Geosci. Model Dev.* **13**, 5425–5464 (2020).

64. S. Theodoridis, C. Rahbek, D. Nogues-Bravo, Exposure of mammal genetic diversity to mid-21st century global change. *Ecography (Cop.)* (2021), doi:10.1111/ecog.05588.

65. I. Overcast, M. Ruffley, J. Rosindell, L. Harmon, P. A. V. Borges, B. C. Emerson, R. S. Etienne, R. Gillespie, H. Krehenwinkel, D. Luke Mahler, F. Massol, C. E. Parent, J. Patiño, B. Peter, B. Week, C. Wagner, M. J. Hickerson, A. Rominger, A unified model of species abundance, genetic diversity, and functional diversity reveals the mechanisms structuring ecological communities. *bioRxiv* (2020), p. 2020.01.30.927236.

66. S. Díaz, N. Zafra-Calvo, A. Purvis, P. H. Verburg, D. Obura, P. Leadley, R. Chaplin-Kramer, L. De Meester, E. Dulloo, B. Martín-López, M. R. Shaw, P. Visconti, W. Broadgate, M. W. Bruford, N. D. Burgess, J. Cavender-Bares, F. DeClerck, J. M. Fernández-Palacios, L. A. Garibaldi, S. L. L. Hill, F. Isbell, C. K. Khoury, C. B. Krug, J. Liu, M. Maron, P. J. K. McGowan, H. M. Pereira, V. Reyes-García, J. Rocha, C. Rondinini, L. Shannon, Y.-J. Shin, P. V. R. Snelgrove, E. M. Spehn, B. Strassburg, S. M. Subramanian, J. J. Tewksbury, J. E. M. Watson, A. E. Zanne, Set ambitious goals for biodiversity and sustainability. *Science*. **370**, 411–413 (2020).

67. C. Rahbek, R. K. Colwell, Biodiversity: Species loss revisited. *Nature*. **473** (2011), pp. 288–289.

68. H. M. Pereira, G. C. Daily, Modeling biodiversity dynamics in countryside landscapes. *Ecology*. **87**, 1877–1885 (2006).

69. J. Harte, A. B. Smith, D. Storch, Biodiversity scales from plots to biomes with a universal species-area curve. *Ecol. Lett.* **12**, 789–797 (2009).

1900   70.   D. R. Lockwood, C. M. Richards, G. M. Volk, Probabilistic models for collecting genetic
1901   diversity: Comparisons, caveats, and limitations. *Crop Sci.* **47**, 861–866 (2007).

1902   71.   L. Czech, M. Exposito-Alonso, grenepipe: A flexible, scalable, and reproducible pipeline to
1903   automate variant and frequency calling from sequence reads. *arXiv* (2021), (available at
1904   http://arxiv.org/abs/2103.15167).

1905   72.   A. Miraldo, S. Li, M. K. Borregaard, A. Flórez-Rodríguez, S. Gopalakrishnan, M. Rizvanovic, Z.
1906   Wang, C. Rahbek, K. A. Marske, D. Nogués-Bravo, An Anthropocene map of genetic diversity.
1907   *Science.* **353**, 1532–1535 (2016).

1908   73.   D. H. Parks, T. Mankowski, S. Zangooei, M. S. Porter, D. G. Armanini, D. J. Baird, M. G. I.
1909   Langille, R. G. Beiko, GenGIS 2: Geospatial Analysis of Traditional and Genetic Biodiversity,
1910   with New Gradient Algorithms and an Extensible Plugin Framework. *PLoS One.* **8**, e69885
1911   (2013).

1912   74.   C. Li, D. Tian, B. Tang, X. Liu, X. Teng, W. Zhao, Z. Zhang, S. Song, Genome Variation Map: a
1913   worldwide collection of genome variations across multiple species. *Nucleic Acids Res.* **49**,
1914   D1186–D1191 (2021).

1915   75.   M. Kardos, E. Armstrong, S. W. Fitzpatrick, S. Hauser, P. Hedrick, J. Miller, D. Tallmon, W.
1916   Chris Funk, The crucial role of genome-wide genetic variation in conservation. *bioRxiv* (2021),
1917   p. 2021.07.05.451163.

1918   76.   D. Spielman, B. W. Brook, R. Frankham, Most species are not driven to extinction before
1919   genetic factors impact them. *Proceedings of the National Academy of Sciences.* **101** (2004), pp.
1920   15261–15264.

1921   77.   Ipbes, *Global assessment report of the Intergovernmental Science-Policy Platform on
1922   Biodiversity and Ecosystem Services* (IPBES Secretariat, Bonn, Germany, 2019;
1923   https://www.ipbes.net/news/ipbes-global-assessment-summary-policymakers-pdf).

1924