Visual space curves before eye movements

Ifedayo-EmmanuEL Adeyefa-Olasupo[1]*†

[1]Department of Psychiatry and Psychotherapy, Charité – Universitätsmedizin Berlin

*Berlin School of Mind and Brain, Humboldt-Universität zu Berlin

†Department of Neuroscience, Yale University

Despite the incessant retinal disruptions that necessarily accompany eye movements, our percept of the visual world remains continuous and stable – a phenomenon referred to as spatial constancy. How the visual system achieves spatial constancy remains unclear despite almost four centuries worth of experimentation (*1-4*). Here I measured visual sensitivity at geometrically symmetric locations, observing transient but marked sensitivity differences between them where none should be observed if cells that support spatial constancy indeed faithfully translate or converge (*5-10*). These differences, recapitulated by a novel neurobiological mechanical model, reflect an overriding influence of putative visually transient error signals that curve visual space. Intermediate eccentric locations likely to contain retinal disruptions are uniquely affected by curved visual space, suggesting that active visual processing at these locations is transiently turned off before an eye movement, and with the gating off of these error signals, turned back on after an eye-movement– a possible mechanism underlying spatial constancy.

We rapidly move our eyes, saccade, between two resting locations in space ~3-5 times per second (*11*). Saccades routinely displace our retinal image of the visual world. Yet we rarely ever notice these discrete displacements saccades accompany. To account for the visual continuity and stability, it is posited that cells in retinotopic brain areas predictively shift their spatial extent from their current locations of sensitivity to their future post-saccadic locations just before a saccade is made. However, it has been long suspected that these early studies do not sample enough spatial positions to thoroughly assess the reliability of these purely translational retinotopic shifts. While researchers have begun to speculate that there may be another mechanism underlying spatial constancy (*12*), experimental designs so far have been unable to implicate a possible mechanism.

Here, for the first time, faint probes were randomly flashed for 20ms at 2.5 dva from the fixation dot (Exp 1), from the midpoint (at eccentricity of 5°) between the fixation dot and the saccade target (Exp 2), or from the saccade target (Exp 3), at upper and lower tangential locations geometrically symmetric to one another with respect to the radial axis (Fig. 1A). To discourage any artificial restriction of attentional and oculomotor resources along the tangential axes, probes were also flashed along the radial axis (see Methods). Subjects reported with a button-press whether they were able to detect a flashed probe around the time of an eye movement towards a saccade target. With respect to target onset, flashed probes fell within the time window of 0 to ~1285ms after target onset (Fig. 1B).

The results in Fig. 2A, 1st panel, calculated across experiments from 0 to 209ms after target onset along the tangential axes show that gaze was averted away from the target in the direction of the foveal tangential clockwise fov_{cw} location (*13*). This shift in gaze is in agreement with transient increases in neural activity in the posterior parietal cortex during fixation at an eccentric foveal site (*14*). From 210ms to 820ms after target onset (2nd panel) gaze continued to remain eccentric while compressing downward towards the foveal tangential counterclockwise fov_{ccw} location. This compression in gaze, in anticipation of the saccade, is in part in agreement with the shrinkage of receptive fields in the extrastriate cortex (*8, 10*). The spatial distribution between these marked shifts in gaze along radial and tangential axes was statistically significant (rad_{shift}, p = 4.0058e-36; tan_{shift}, p = 0.0149). Fig. 2A, 3rd panel, shows that the saccade landing position, from 0 to 50ms after saccades land in the periphery, was dramatically mislocalized towards the peripheral tangential clockwise $peri_{cw}$ location. This observation is in agreement with the orthogonal mislocalization of flashed stimuli away from a stationary saccade target (*15, 16*). As expected, a larger variance was observed for the initial locus of gaze (6.84 deg²) when compared to the variance associated with the compressed locus of gaze (2.12 deg²). As for the variance associated with the saccade landing position, a larger variance (10 deg²) was observed when compared to the loci of the gaze in the fovea (Fig. 2A, 4th panel). These results suggest that the tangential bias in the locus of the saccade landing position (more eccentric region) remained mirroring the tangential bias in the initial shift in gaze within the fovea (less eccentric region), even as gaze actively compressed towards the fov_{ccw} location.

To account for this unexpected spatial difference, I predict the presence of "visually transient error signals" that delay visual processing at the more eccentric locations, in turn introducing an angular bearing (curvilinear) component within the representation of visual space. In line with this prediction, Fig. 2B, 1st panel, shows that when subjects began to compress their gaze downward towards the fov_{ccw} location, there was a transient but marked increase in peri-saccadic sensitivity at the fov_{ccw} location, while at the fov_{cw} location, there was a transient but marked decline in peri-saccadic sensitivity. From 820ms onwards after target onset there was a global decline in peri-saccadic sensitivity at both foveal locations. This global decline is in agreement with peri-saccadic corollary discharge commands originating from pre-motor brain areas (*4*). The 2nd and 3rd panels show the opposite pattern in the changes in peri-saccadic sensitivity. Specifically, from 210ms to 820ms after target onset, there are declines in peri-saccadic sensitivity at $para_{ccw}$ and $peri_{ccw}$ locations and increases in peri-saccadic sensitivity at $para_{cw}$ and $peri_{cw}$ locations. Equally revealing is the fact that at these eccentric locations, from 820ms onwards after target onset, there is a slow but clear peri-saccadic sensitivity "hand-off", as $para_{ccw}$ and $peri_{ccw}$ locations that once experienced declines in peri-saccadic sensitivity slowly begin to experience increases. Conversely, at $para_{cw}$ and $peri_{cw}$ locations that once experienced increases in peri-saccadic sensitivity slowly begin to experience declines. These hand-offs suggest that the visually transient error signals delaying visual processing at the more eccentric locations are eventually gated off around the time a saccade has been planned.

In line with the presumed curvilinear representation of visual space caused by transient error signals and the eventual gating off of these signals, Fig. 2C shows that saccade trajectories from the current position of gaze towards the periphery were curved across experiments, at a rate of 73% towards the $peri_{cw}$ location, while in 27% of all trials, saccades were curved towards the $peri_{ccw}$ location, reflecting a calibrated visual representation towards the actual location of the target. Specifically, when the saccade target was presented at azimuth angles of 0°, the trajectories were curved towards the per_{cw} location at a rate of 91% with an average curvature of 0.22 in area/amp (*17*). Conversely, for the remaining trials when saccades were curved towards the per_{ccw} location, an average curvature of -0.80 in area/amp was observed. Next, when the peripheral target was presented at an azimuth angle of 45°, the trajectories were curved towards the per_{cw} location at a rate of 65% with an average curvature of 0.56 in area/amp, while for the remaining trials (saccades curved towards the per_{ccw} location) an average curvature of -0.74 in area/amp was observed. Finally, when the saccade target was presented at an azimuth angle of 315°, trajectories were curved towards the per_{cw} location at a rate of 60% with an average curvature of 1.12 in area/amp, while for the remaining trials (saccades curved towards the per_{ccw}) location) an average curvature of -1.17 in area/amp was observed.

Fig. 3A, 1st panel, shows that a large part of the observed changes in peri-saccadic sensitivity at fov_{cw} and fov_{ccw} locations is actively meditated by rhythmicity in the delta frequency band at ~2 Hz and ~3 Hz respectively, which is known to support attentional shifts or, more often, their inhibitions during demanding visuomotor tasks (*18*). Fig. 3A, 2nd panel, shows that the spectral signature that mediated the fov_{cw} location was not location-specific when compared to shuffled data sets (p=0.2280). Conversely, the spectral signature that supported the fov_{ccw} location was specific to that retinotopic location (p=0.0230). The $para_{cw}$, $para_{cw}$, $peri_{ccw}$, and $peri_{ccw}$ locations also exhibited rhythmicity in the ~2 Hz delta frequency band (Fig. 3B–C, 1st panel). The spectral signature observed at eccentric clockwise and counterclockwise retinotopic locations were not location-specific (p=0.0915, p=0.9725, p=0.3179, p=0.9725) (Fig. 3B–C, 2nd panel). Furthermore, the phase of ~2 Hz within $para_{cw}$ and $para_{ccw}$ locations was consistently offset by approximately 201°, and even larger offset of 290° was observed at the more eccentric $peri_{cw}$ and $peri_{ccw}$ locations (Fig. 3B–C, 3rd panel). This antiphase periodicity at ~2 Hz suggests that the retinotopic brain areas that support these upper and lower eccentric tangential locations are synchronized by attentional dynamics at a common frequency but with location-specific phases (*19*).

To provide a mechanistic account of the neural computations and architecture that support this "curved visual space", a neurobiological mechanical model is proposed. At its core, the model assumes that underlying peri-saccadic attentional and oculomotor signals are actively modulated by visually transient error signals which give rise to an input and output forces exerted by corresponding time varying masses (Fig. 4B, Fig. 4E) that act upon a "retinotopic lever". Under an inverse force field ($\varepsilon = -0.55$), input and output forces transiently shift spring-loaded sensors – population retinotopic receptive fields (pRFs) that tile visual space and thereby shape normalized density

(putative population sensitivity) at a sampled retinotopic location in time (Fig. 4A, Fig. 4D, and Extended Data Fig. 1-2).

100   As far as the foveal simulations, location-specific visually transient error signals (Fig. 3A) give rise to an input force that is preparing to act on the input arm of a "retinotopic lever" which causes pRFs to shift upper ward in the direction of the fov_{cw} location and later compress towards the fov_{ccw} location (Fig. 2A, 1st and 2nd panel) (*14*). A compressed input force causes the lever to pivot on its "foveal fulcrum" thereby providing a transient neural advantage at the fov_{ccw} location. To always maintain retinotopic organization, pRFs that tile the more eccentric

105 regions in visual space are initially restricted from over-translating towards the foveal fov_{cw} location by their population elastic fields (pEFs, see Methods) which remains mirroring the onset location of the input force before a full compression occurs. For the sake of simplicity, foveal simulations began from the point in time when pRFs mirrored the initial shift in gaze (translational output force), followed by late compressive shift towards the fov_{ccw} location (compressed input force) (Fig. 4B). Under these input and output forces, simulation results in Fig. 4C show

110 that putative population sensitivity recapitulated signatures in peri-saccadic sensitivity in the foveal region of visual space (Fig. 2B, 1st panel). Assuming that retinotopic cells are not mediated by visually transient error signals (i.e., visual representation does not include a curved component), translational shifts anticipating retinal image disruptions should have yielded no difference in putative population sensitivity (Fig. 4C, dotted marker) in comparison to peri-saccadic sensitivity observed in Fig. 2B, 1st panel. Similarly, symmetrically compressed

115 retinotopic shifts towards a stationary peripheral site (Fig. 4C, thin markers) cannot explain the changes in peri-saccadic sensitivity observed in the fovea.

       As far as the eccentric simulations (Fig. 4D), after a fully compressed input force, pRFs moved further towards the fov_{ccw} location in proportion to the mislocalization of the target. Cells across retinotopic brain areas can receive both convergent and translational signals around the time of a saccade (*6, 9*). To this end, pRFs later

120 converged around an emerging output mass which exceeded the magnitude of the initial shift in gaze and approximated the location of the mislocalized target (Fig. 2A, 1st and 3rd panel) (*20*). This is immediately followed by the onset of a translational force proportional to the magnitude of a saccade orthogonal to the radial axis towards the $peri_{cw}$ location (Fig. 4E). Indeed, under three forces, especially when the forces that corresponds to convergent and translational signals emerge, a neural advantage at eccentric clockwise locations should be observed. As

125 predicted, Fig. 4F shows that putative population sensitivity was able to recapitulate peri-saccadic sensitivity signatures at parafoveal locations (Fig. 2B, 2nd panel). Once again, a purely translational or symmetrically compressed shifts cannot explain changes in peri-saccadic sensitivity.

       Overall, results suggest that peri-saccadic retinotopic shifts reflect an overriding influence of visually transient error signals that support both local shifts in gaze and erroneous perceived changes in the retinal location of

130 a distal target. A curvilinear representation of visual space, which underlies these observations, can also explain (i).

the frequent mislocalization of flashed stimuli away from a stationary target (*15, 16*), (ii). the neural computations and architecture that causes cells in the extrastriate cortex to shift orthogonal to the presumed saccade direction (*7, 8*) and (iii). the extent retinotopic cells will symmetrically compress towards a stationary target (*21*). These shifts in gaze remain a curious point since they are directed away from the distal target (*13*). It is possible that this reflects an evolutionarily conversed visuomotor avoidance strategy use to track a distal target (e.g., conspecific, predator) from a safe distance (e.g., corner of the eye), later followed by a triggered movement (e.g., saccade) towards or away from the target (*22*). Another point is the anticipatory compression of gaze, which could mark the enhancement of visual processing towards a selected peripheral site (*8, 10, 12*). However, it is more likely that this is the result of the visual system temporarily withholding the additional spreading of neural resources to the surrounding eccentric locations (regions likely to contain retinal disruptions introduced by the impending saccade) (*23*). This can explain why retinotopic representations at eccentric locations (parafoveal and peripheral), inferred from behavioral and spectral results highlighted above, remained unresponsive even as gaze actively compressed downward.

REFERENCES

1. Descartes, R. *Traité de l'homme* (Paris, France, 1644).

2. Helmholtz, H.V. *A treatise on physiological optics* (Dover, New York, NY, 1962).

3. von Holst, E., Mittelstaedt, H. The reafferent principle: reciprocal effects between central nervous system and periphery. *Naturwissenschaften*, 37, 464–476. (1950).

4. Sperry, R.W. Neural basis of the spontaneous optokinetic response produced by visual inversion. *J. Comp. Physiol. Psychol*, 43, 482–489. (1950).

5. Duhamel, J. R., Colby, C. L., Goldberg, M. E. The updating of the representation of visual space in parietal cortex by intended eye movements. *Science*, 255(5040), 90–92. (1992).

6. Umeno, M.M., Goldberg, M.E. Spatial processing in the monkey frontal eye field. I. Predictive visual responses. *J Neurophysiol*, 78:1373–138. (1997).

7. Connor, C. E., Gallant, J. L., Preddie, D. C., Van Essen, D. C. Responses in area V4 depend on the spatial relationship between stimulus and attention. *Journal of Neurophysiology*, 75(3), 1306–1308. (1996).

8. Tolias, A. S., Moore, T., Smirnakis, S. M., Tehovnik, E. J., Siapas, A. G., Schiller, P. H. (2001). Eye movements modulate visual receptive fields of V4 neurons. *Neuron,* 29(3), 757–767.

9. Zirnsak, M., Steinmetz, N. A., Noudoost, B., Xu, K. Z., Moore, T. Visual space is compressed in prefrontal cortex before eye movements. *Nature*, 507, 504–507. (2014).

10. Hartmann, T.S., Zirnsak, M., Marquis, M., Hamker, F.H., Moore, T. (2017). Two types of receptive field dynamics in area V4 at the time of eye movements? *Front. Syst. Neurosci*, 11, 1–7.

11. Rayner, K. Eye movements in reading and information processing: 20 years of research. *Psychol.*
175 *Bull*, 124, 372 –422. (1998).

12. Zirnsak, M., Moore, T. Saccades and shifting receptive fields: anticipating consequences or selecting targets? *Trends in cognitive sciences*, 18,12, 621-628. (2014)

180
13. Hafed, Z. M., Clark, J. J. Microsaccades as an overt measure of covert attention shifts. Vision Research, 180 42(22), 2533–2545. (2002).

14. Andersen, R. A., Mountcastle, V. B. The influence of the angle of gaze upon the excitability of the light-sensitive neurons of the posterior parietal cortex. *The Journal of Neuroscience*, 3(3), 532–548. (1983).

185
15. Kaiser, M., Lappe M. Perisaccadic mislocalization orthogonal to saccade direction. *Neuron*, 22;41(2), 293–300. (2004).

16. Zimmermann, E., Morrone, M. C., Burr, D. C. Visual mislocalization during saccades sequences. *Exp*
190 *Brain Res*, 233(2):577-585. (2014).

17. Walker, R., McSorley, E., Haggard, P. The control of saccade trajectories: direction of curvature depends on prior knowledge of target location and saccade latency. *Percept Psychophys*, 68(1), 129-138. (2006).

195
18. Harmony, T. The functional significance of delta oscillations in cognitive processing. *Frontiers in Integrative Neuroscience*, 7, 83. (2013).

19. Fries, P. A mechanism for cognitive dynamics: neuronal communication through neuronal coherence. *Trends in Cognitive Sciences,* 9(10), 474–480. (2005).

200
20. Dowiasch, S., Blohm, G., Bremmer, F. Neural correlate of spatial (mis-)localization during smooth eye movements. *The European Journal of Neuroscience*, 44(2), 1846–1855. (2016).

21. Ross, J., Morrone, M.C., and Burr, D.C. Compression of visual space before saccades. *Nature*, 386, 598–
205 601. (1997).

22. Porter, T. C. Contributions to the study of flicker II. *Proceedings of the Royal Society of London Series A*, 70, 313-329. (1902).

210   23. Womelsdorf, T., Anton-Erxleben, K., Treue, S. Receptive field shift and shrinkage in macaque middle temporal area through attentional gain modulation. *The Journal of Neuroscience*, 28, 8934–8944. (2008).

215   24. Rolfs, M., Jonikaitis, D., Deubel, H., Cavanagh, P. Predictive remapping of attention across eye movements. *Nat. Neurosci*, 14, 252–256. (2011).

25. Arkesteijn, K., Belopolsky, A. V., Smeets, J., Donk, M. The Limits of Predictive Remapping of Attention Across Eye Movements. *Frontiers in psychology*, 10, 1146. (2019).

220   26. Botev, Z. I., Grotowski, J. F., Kroese, D. P. Kernel density estimation via diffusion. *Annals of Statistics*, 38:2916–2957. (2010).

27. Crossland, M.D., Sims, M., Galbraith, R.F., Rubin, G.S. Evaluation of a new quantitative technique to assess the number and extent of preferred retinal loci in macular disease. *Vision Research*, 44(13), 1537-
225   1546. (2004).

28. Ludwig, C. J. H., Gilchrist, I. D. Measuring saccade curvature: A curve-fitting approach. *Behavior Research Methods, Instruments, & Computers*, 34, 618-624. (2002).

230   29. Nandy, A. S., Nassi, J. J., Reynolds, J. H. Laminar Organization of Attentional Modulation in Macaque Visual Area V4. *Neuron*, 4;93(1), 235-246. (2017).

30. Tass, P., Rosenblum, M.G., Weule, J., Kurths, J., Pikovsky, A., Volkmann, J., Schnitzler, A., Freund, H. J. Detection of n:m Phase Locking from Noisy Data: Application to Magnetoencephalography. *Physical*
235   *review letters,* 81, 3291-3294. (1998).

31. Adeyefa-Olasupo, I-E. Retinotopic mechanics derived using classical physics. *arXiv, 2109.11632*. (2021).

32. Bowman, A.W., Azzalini, A. *Applied Smoothing Techniques for Data Analysis: the Kernel Approach with*
240   *S-Plus Illustrations* (Oxford University Press, Oxford, 1997).

ACKNOWLEDGEMENTS

I would like to thank D. P. Melcher and S. Wirth for comments on the manuscript and B. Wichterlová for editing the

245   manuscript. I would also like to thank A. S. Nandy for his initial guidance on the experimental design.

COMPETING INTEREST

The author has no competing financial interest.

DATA AVAILABILITY

The data that supports this study is available upon reasonable request.

CODE AVAILABILITY

The custom Matlab code that supports the analyses reported in this study is available upon reasonable request.

SUPPLEMENTAL TEXT

Fig 1. Spatiotemporal profile of experiments. (A) Markers in the illustration of visual space show all retinotopic locations at which a probe was flashed before and after the deployment of attention, saccade planning, and, finally,

260   the execution of a saccade. Probes of interest, presented along the tangential axes, are conspicuously highlighted in open circle markers. **(B)** Temporal sequence of a valid trial across experiments with respect to target onset.

Fig 2. Gaze dynamics, saccade landing position, peri-saccadic sensitivity, and saccade deviation estimates. (A) Probability density map of foveal gaze and saccade landing was calculated using a bivariate gaussian kernel along

265   with their bivariate contour ellipse area estimates for each probability density map. Error estimates are standard error of means across observers. **(B)** Shifts in peri-saccadic sensitivity along tangential axes with respect to target onset; error estimates were calculated using a 20-fold jackknife procedure. **(C)** Mean saccade trajectory deviation for horizontal and oblique saccades across experiments. Positive values on the ordinate denote trajectory deviation towards a clockwise location (cw), while negative values denote trajectory deviation towards a counterclockwise

270   location (ccw) (*17*).

Fig 3. Frequency domain and cross-location phase coherence estimates. (A-C) Frequency domain representation of raw detrended peri-saccadic sensitivity at less and more eccentric locations in visual space. Shuffled (n=1000, dashed line) versus unshuffled (solid line) data set were used to compute location-specific spectral signatures at a

275    $p < 0.05$. For parafoveal and peripheral clockwise and counterclockwise locations a cross-location phase coherency analysis was performed.

Fig 4. Retinotopic lever of curved visual space. (A) For illustration purposes, a single eccentricity dependent population receptive field (pRF) within a retinotopic field is highlighted in pink. A pRF is defined by its center, size, and an eccentric dependent population elastic field (pEF). **(B)** MGα denotes gaze transitional signals modelled as a virtual mass at infinity located in the direction of an upper tangential location approximating the initial shift in gaze. MF denotes compressive signals modelled as a varying mass located at a lower tangential location in the fovea. **(C)** Simulation results in the foveal region when a pRF is under the influence of a resultant force exerted by (i) MF and MGα (curved space, thick marker) (ii) MC (convergent signals modelled as a varying mass at a stationary peripheral site; see Methods) and (iii) MT (saccade transitional signals modelled as a virtual mass at infinity located in the direction of a straight saccade; see Methods). **(D)** A pRF bounded by pEF is highlighted in pink for parafoveal simulations. **(E)** MGβ denotes gaze translational signals modelled as a virtual mass at infinity located in the direction of a downward shifts in gaze, MM denotes a varying mass at a mislocalized peripheral target, and MS denotes the curved transitional signals modelled as a virtual mass at infinity located in the direction of a curvilinear saccade. **(F)** Curved, convergent and translational simulation results.

Extended Data Fig. 1 Peri-saccadic spatiotemporal retinotopic shifts for foveal simulations. (A) hexagonal retinotopic field spanning the entire visual field. Retinotopic shifts are produced by the perturbation of pRFs under an inverse force field in discrete time increments of 1ms by the combination of external forces exerted by **(B)** corresponding masses ($M_{G\alpha} + M_F$).

Extended Data Fig. 2 Peri-saccadic spatiotemporal retinotopic shifts for parafoveal stimulations. (A) Retinotopic shifts when the combination of external forces is exerted by **(B)** corresponding masses ($M_{G\beta} + M_M + M_S$).

METHODS

Human subjects

Six consenting human subjects (5 females and 1 male; 21-26 years old) with corrected-to-normal visual acuity participated in the three experiments reported. Three of the six subjects participated in the foveal and parafoveal experiments. The remaining three subjects participated in the peripheral experiment. The Yale Ethics Review board approved the protocol of these experiments and the storage of the data that was collected.

Visual stimuli and experimental protocol

Fig. 1 summarizes the protocol used across experiments. In a dimly lit room, stimuli were presented on a gamma corrected display with a spatial resolution of 1400 x 1050 pixels and a mean luminance of 38cd/m^2. Eye movements were recorded using an infrared video-based eye-tracker sampled at 1kHz (I-Scan Inc., Woburn, MA). While tightly head fixed using a head rest and a customized bite bar, subjects at the onset of each trial acquired fixation for 300ms. Fixation within this temporal window was spatially aligned with the fixation point using a custom-developed 9-point eye-calibration procedure in Picto. Next, depending on the subject, a saccade target was presented at an eccentricity of 10° and at azimuth angles of 0° (for s_1), 45° (for s_2) and 315° (for s_3) (Fig. 1A). Following target onset, fixation was maintained for another 300-600ms within ~2dva fixation tolerance window, at the end of which the fixation spot was extinguished, which served as the central motor cue for subjects to make an eye movement towards the saccade target. For another 500ms after the central motor cue, the saccade target remained visible. Finally, a low contrast probe (at a contrast level which the subject could detect with 50% accuracy) were isotropically flashed, depending on the experiment, for 20ms either along the radial or tangential axis at a random time during the 300-600ms window between the target onset and the central motor cue (25% of trials) or at a random time between 0-340ms after the central motor cue (50% of the trials). On the remaining 25% (which yielded a false alarm rate of 1.2% along the tangential axes), no probes were presented. All three conditions were randomly interleaved across trials. With respect to target onset, the presentation of flashed probes fell within a time window of 0-1285ms after target onset (Fig. 1B). In addition to the execution of a saccade, each subject was asked to report, using a push button, whether they were able to detect the flashed probe. While the responses along the radial axis were not analyzed, the presentation of these probes was crucial in controlling for any response biases that may have emerged at upper or lower locations along tangential axes. To ensure that attentional resources was not being driven by an asymmetric stimuli configuration or insufficient control measurements that is to yield confusing results, no secondary salient spatial cue was presented (*24, 25*). Along the tangential axes, a total of ~6010 trials were collected across experiments, with a minimum of 650 trials per subject.

Statistical analysis

To investigate gaze dynamics, probability density maps and variance estimates were calculated using median filtered edge preserving fixational data, which was rotated to an angle of 0° across target locations. Next, I calculated a gaze

probability density map for the foveal and periphery regions using a bivariate Gaussian kernel (*26*). The estimated locus of gaze across subjects was defined as the peak of each probability density map (i.e., the yellow contour lines). To compare the distribution of gaze in the fovea during the 0 to 209ms window after target onset versus 210 to 820ms window after target onset, a Mann-Whitney U statistic was calculated at p value less than 0.05. Further, variance in gaze was calculated using a bivariate Contour Ellipse Area (BCEA) approach which included 68% of fixational data around the mean (*27*). Accompanying error bars were obtained which denote the standard error of the mean variance across subjects. To investigate the extent saccade trajectories were curved across experiments at each saccade location. Horizontal and vertical eye positions, with the time window starting point constituting the moment fixation was maintained to the moment the eye landed in the periphery, was isolated. Next to obtain a measurement of curvature, the curved area that emerged from the eye position during this time window was calculated. To then obtain a ratio measure of curvature per unit amplitude, this area was divided by the amplitude of the saccade (*28*). Accompanying calculated error bars denote the standard error of the mean at each target location across experiments. To investigate normalized changes in peri-saccadic sensitivity across subjects, I first binned each subject's percent correct responses for the foveal (Exp 1), parafoveal (Exp 2), and peripheral (Exp 3) experiments, using a 15ms sliding window that was moved in 45ms increments. Next, I obtained each subject's raw peri-saccadic sensitivity, using a 20-fold jackknife procedure by estimating sensitivity from 95% of the data. This was repeated 20 times, each time leaving out 5% of the data. This produced a mean and an error estimate of peri-saccadic sensitivity at each tangential location for each experiment across subjects (*29*). To control for any eccentric-dependent effect on raw peri-saccadic sensitivity, these traces were normalized by the average sensitivity over the initial 100ms in the data. To investigate frequency domain and cross-location phase coherence dynamics, a fast Fourier transform (FFT) was applied to the detrended raw peri-saccadic sensitivity data. Next, I computed the mean of the spectral signatures and corresponding standard error of the mean across subjects. To than assess whether the observed periodicities were location specific, I randomly shuffled (n=1000) the probe location identities across trials and calculated the sensitivity function and periodicity of the shuffled data using the same procedures noted above. A two tailed paired-sample t-test at p-values of 0.05 was calculated to assess the significance between shuffled and non-shuffled data. Finally, for the cross-location phase coherence analysis, I applied an approach to the shuffled and non-shuffled detrended data that is commonly used in assessing the underlying coherence in neural data (*30*). I then projected the difference between phase angles of each subject between counterclockwise (ccw) and clockwise (cw) locations and then calculated the resultant vector across subjects.

Retinotopic mechanics

The underlying architecture of this simple mechanical model is composed of a retinotopic field (ϕ_r) and a retinotopic lever which is inspired by a neurobiological force field. The retinotopic field is a two-dimensional overlapping hexagonal grid of eccentricity-dependent receptive fields - RF_i - that tiled the foveal, parafoveal and

370 peripheral regions in visual space (Extended Fig. 1-2). Each RF_i was defined by three parameters: (a) p_i, the spatial extent of the RF center (x_i, y_i) at t_0 (when there is no resultant force acting on the ϕ_r). p_i determines the eccentricity - e_i of RF_i, (b) s_i, the radius of RF_i is proportional to e_i, and (c) the movement extent (M_{max}) of RF_i is its EF_i which is also proportional to e_i. ϕ_r emerged from peri-saccadic attentional and oculomotor signals actively modulated by visually transient error signals that act as input and output forces of retinotopic lever in this field, and transiently

375 perturb a constellation of population receptive fields from equilibrium and thereby modulate their putative population sensitivity (density readouts) as a function of space and time. For simulations at the less eccentric ("foveal"), the resultant force ($\overrightarrow{F_{R1}}$) included an input and output force ($\overrightarrow{F_I}$, and $\overrightarrow{F_O}$) and an internal force ($\overrightarrow{F_E}$) which RF_i at t_{0+n} experiences:

380
$$\overrightarrow{F_O} = \overrightarrow{U_{M_{G\alpha}}} . M_{G\alpha} . |G\alpha|$$

$$\overrightarrow{F_I} = \overrightarrow{U_{M_F}} . M_F . R_\beta^\varepsilon$$

$$\overrightarrow{F_{R1}} = \overrightarrow{F_I} . k_I + \overrightarrow{F_O} + \overrightarrow{F_E}$$

where $M_{G\alpha}$ and M_F are the virtual mass at infinity in the upward direction which approximates an initial shift in gaze and a foveal mass located at a lower tangential foveal retinotopic site which approximates a compressed shift in gaze respectively. $\overrightarrow{U_{M_{G\alpha}}}$ and $\overrightarrow{U_{M_F}}$ are the unit vectors in the direction of the virtual mass and the foveal mass respectively. $|G\alpha|$ is the magnitude in gaze towards the upper tangential foveal retinotopic site, while R_β is the

390 spatial difference between RF_i and M_F raised to an epsilon scalar – ε. The value of ε, depending on the retinotopic mechanics that best explains the phenomenon in question, can either be consistent with a proportional- distance rule (i.e., >0) or an inverse-distance rule (i.e., <0) (*31*). Further, k_I was obtained by calculating the average of $R_{\beta0}^\varepsilon$ across RF_i's and then taking its reciprocal $\frac{1}{(R_{\beta0}^\varepsilon)}$, while $\overrightarrow{F_E}$ is proportional to the movement of RF_i. For the more eccentric

("parafoveal") simulations, the resultant force ($\overrightarrow{F_{R2}}$) included an input, output, an additional orthogonally biased

translational force ($\overrightarrow{F_I}$, $\overrightarrow{F_O}$ and $\overrightarrow{F_T}$) and an internal force ($\overrightarrow{F_E}$) which RF_i at t_{0+n} experiences:

$$\overrightarrow{F_I} = \overrightarrow{U_{M_\beta}} . M_{G\beta} . |G\beta|$$

$$\overrightarrow{F_O} = \overrightarrow{U_{M_M}} . M_M . R_\mu^\varepsilon$$

$$\overrightarrow{F_T} = \overrightarrow{U_{M_S}} . M_S . |S|$$

$$\overrightarrow{F_{R2}} = \overrightarrow{F_O} . k_O + \overrightarrow{F_I} + \overrightarrow{F_T} + \overrightarrow{F_E}$$

where $M_{G\beta}$ and M_M are the virtual mass at infinity in the direction towards lower foveal tangential retinotopic site which approximates a complete compression of gaze and the mislocalized mass located at a peripheral retinotopic site that approximates a mislocalized saccade target respectively. $\overrightarrow{U_{M_{G\beta}}}$ and $\overrightarrow{U_{M_M}}$ are the unit vectors in the direction of the virtual mass and the mislocalized peripheral mass respectively. $|G\beta|$ is the magnitude of a fully compressed gaze which is roughly proportional to the magnitude of a mislocalized target. R_μ is the spatial difference between RF_i and M_M raised to ε, while $|S|$ approximates the magnitude of an impending saccade. The same calculation used to obtain k_I was used to obtain k_O, however in this case, the average of $R_{\mu0}^\varepsilon$ across RF_i's was used. Depending on the stimulation, a time varying resultant force ($\overrightarrow{F_{R1}}$ or $\overrightarrow{F_{R2}}$) perturbed each RF_i, which produced spatiotemporal peri-saccadic retinotopic displacements (Extended Data Fig. 1-2). The population movements of the constellations of RF_i's manifested in time-varying modulation of raw density, modelled using a bivariate gaussian kernel function at sampled tangential locations (*32*). Density readouts were later normalized using the same approach used to normalize the raw peri-saccadic sensitivity traces.

Fig 1. Spatiotemporal profile of experiments. (A) Markers in the illustration of visual space show all retinotopic locations at which a probe was flashed before and after the deployment of attention, saccade planning, and, finally, the execution of a saccade. Probes of interest, presented along the tangential axes, are conspicuously highlighted in open circle markers. **(B)** Temporal sequence of a valid trial across experiments with respect to target onset.

Fig 2. Gaze dynamics, saccade landing position, peri-saccadic sensitivity, and saccade deviation estimates. (A)
Probability density map of foveal gaze and saccade landing was calculated using a bivariate gaussian kernel along with their bivariate contour ellipse area estimates for each probability density map. Error estimates are standard error of means across observers. **(B)** Shifts in peri-saccadic sensitivity along tangential axes with respect to target onset; error estimates were calculated using a 20-fold jackknife procedure. **(C)** Mean saccade trajectory deviation for horizontal and oblique saccades across experiments. Positive values on the ordinate denote trajectory deviation towards a clockwise location (cw), while negative values denote trajectory deviation towards a counterclockwise location (ccw) (17).

Fig 3. Frequency domain and cross-location phase coherence estimates. (A-C) Frequency domain representation of raw detrended peri-saccadic sensitivity at less and more eccentric locations in visual space. Shuffled (n=1000, dashed line) versus unshuffled (solid line) data set were used to compute location-specific spectral signatures at a p<0.05. For parafoveal and peripheral clockwise and counterclockwise locations a cross-location phase coherency analysis was performed.

Fig 4. Retinotopic lever of curved visual space. (A) For illustration purposes, a single eccentricity dependent population receptive field (pRF) within a retinotopic field is highlighted in pink. A pRF is defined by its center, size, and an eccentric dependent population elastic field (pEF). **(B)** $M_{G\alpha}$ denotes gaze transitional signals modelled as a virtual mass at infinity located in the direction of an upper tangential location approximating the initial shift in gaze. M_F denotes compressive signals modelled as a varying mass located at a lower tangential location in the fovea. **(C)** Simulation results in the foveal region when a pRF is under the influence of a resultant force exerted by (i) M_F and $M_{G\alpha}$ (curved space, thick marker) (ii) M_C (convergent signals modelled as a varying mass at a stationary peripheral site; see Methods) and (iii) M_T (saccade transitional signals modelled as a virtual mass at infinity located in the direction of a straight saccade; see Methods). **(D)** A pRF bounded by pEF is highlighted in pink for parafoveal simulations. **(E)** $M_{G\beta}$ denotes gaze translational signals modelled as a virtual mass at infinity located in the direction of a downward shifts in gaze, M_M denotes a varying mass at a mislocalized peripheral target, and M_S denotes the curved transitional signals modelled as a virtual mass at infinity located in the direction of a curvilinear saccade. **(F)** Curved, convergent and translational simulation results.