Visual space curves before eye movements

Ifedayo-EmmanuEL Adeyefa-Olasupo[1]* & Zixuan Xiao[4]

[1]Department of Psychiatry and Psychotherapy, Charité – Universitätsmedizin Berlin

*Berlin School of Mind and Brain, Humboldt-Universität zu Berlin

†Department of Neuroscience, Yale University

[4] Department of Psychology, Beijing Normal University

ABSTRACT

In most if not all experiments within cognitive and systems neuroscience, fixation is required before the onset of an experimental trial. However, the term "fixation" is rather misleading since our eyes are constantly moving. One type of transient miniature movements observed during fixation is commonly referred to as fixational eye movements. Traditionally, and even in more recent experiments, the inevitable effects of fixational eye movements on visual perception have been largely overlooked. To date, no study has been able to provide an account of the extent to which fixational eye movements actively mediate visual perception well before and around the time of a rapid eye movement (a saccade) at retinotopic locations that constitute distinct retinotopic regions of visual space. Oculomotor planning and performance effects associated with these transient shifts around the time of a saccade remain equally unknown. Here, we systematically measured visual sensitivity in human subjects to visual probes presented during fixation, well before and around the time of a saccade, at geometrically symmetric retinotopic locations in the foveal, parafoveal and peripheral regions of visual space. We observed transient local asymmetric visual sensitivity differences between these symmetric retinotopic locations where none should be observed. Surprisingly, we also observed that saccade trajectories, which are expected to travel along a linear path, routinely deviate along a curved path towards orthogonal eccentric retinotopic locations. As such, we proposed a novel neurobiologically inspired phenomenological force field model in which underlying attentional and oculomotor signals are modulated by transient eccentric error signals that manifest as temporally overlapping forces and act on retinotopic brain areas. These forces, which transiently shift putative population sensitivity towards retinotopic foveal sites to support transient orthogonal fixational eye movements, and, shortly after, along an axis orthogonal to the saccade direction towards a mislocalized peripheral site, succinctly capture the essence of our perceptual and oculomotor observations.

INTRODUCTION

The central 2° of the visual field is extensively represented in the retina, the superior colliculus

(SC), and the primary visual cortex (V1), and is thus well suited for tasks that necessitate detailed

inspections of everyday visual objects (e.g., fruits, human faces, traffic signals). Under these

conditions, transient shifts in gaze, which are marked by fixational eye movements, are ubiquitous

(*1*). These transient shifts in gaze, which, as previous studies have shown, actively mediate visual

perception in general well before they occur, precede rapid eye movements, saccades (*2-3*).

However, whether these transient shifts in gaze, once they occur, actively mediate visual

perception well before and around the time of a saccade at retinotopic locations that constitute

distinct retinotopic regions of visual space (i.e., the foveal, parafoveal or peripheral retinotopic

regions) remains unknown (*4-5*). The oculomotor consequences associated with the onset of these

transient shifts in gaze around the time of a saccade are also unknown.

Equally unclear is the extent to which early retinotopic representations that support local

transient shifts in gaze during fixation influence peri-saccadic spatiotemporal retinotopic shifts.

Specifically, recent behavioral and neurophysiological studies have suggested that around the time

of a saccade, cells across retinotopic brain areas do not purely shift (or translate) from their current

retinotopic locations of sensitivity to fixed future post-saccadic retinotopic locations of sensitivity

as they primarily converge towards the target of the saccade (*6, 7*). Oddly enough, some of these

studies have also reported convergent shifts orthogonal to the saccade direction, almost as if the

location of the fixed saccade target is actively being ignored by the visual system (*7-10*).

Unquestionably, these results, which report either purely translational (*11-12*) or symmetric and

orthogonal convergent effects emerge from experiments that require fixation, during which

transient shifts in gaze occur (*1-5*). Keeping in mind that impending fixational eye movements

cause human subjects to erroneously perceive eccentric stimuli as being shifted away from their

actual fixed locations (*2*), it is plausible that the extent to which retinotopic cells shift beyond the

spatial extent of their fixed future post-saccadic retinotopic locations or converge along an axis

that is orthogonal to the saccade direction could indeed be the result of retinotopic representations

which support transient shifts in gaze that occur well before a saccade – an account that has been

completely overlooked by previous translational and convergent studies.

In addition to these unknowns, the neural computations and architecture that support spatial

constancy – the visual system's ability to account for the incessant retinal disruptions that

necessarily accompany saccades (*13*) – remain poorly understood. Recent studies have shown that

retinal suppression signals triggered by fixational eye movements, which do not depend on

corollary discharge commands originating from pre-motor brain areas (*14*), temporarily reduce

visual sensitivity across visual space around the time of a saccade. It is therefore possible that

attentional and oculomotor signals that support the allocation of more neural resources towards the

foveal retina during fixation (*15-20*), at the expense of neural resources that would otherwise

support intermediate eccentric retinotopic locations likely to contain retinal disruptions, are

important for uncovering the neural computations and architecture underlying spatial constancy.

Assuming these attentional and oculomotor signals play an important role, purely translational

signals around the time of a saccade – which researchers once believed could explain spatial

constancy (*13, 21*) but have been exposed by recent behavioral and electrophysiological studies as

the result of coarse spatial sampling (*6, 22*) – would not be required for the visual system to achieve

spatial constancy (*21*).

RESULTS

To uncover how the onset and continual shifts in gaze during fixation actively influence (i) visual

sensitivity at retinotopic locations that constitute distinct retinotopic regions of visual space, (ii)

oculomotor planning and performance around the time of a saccade, (iii) peri-saccadic

spatiotemporal translational and convergent retinotopic shifts and (iv) the neural computations and

architecture that could support spatial constancy, we first assessed transient shifts in gaze after the

onset of a saccade target presented at a fixed retinotopic location during fixation and around the

time of a saccade. We later assessed saccade landing positions after subjects executed a saccade,

which informed us as to where subjects perceived the fixed saccade target to be located during

fixation. Following this, we assessed the transient consequences of visual sensitivity to visual

probes that were presented around the path of a saccade in the foveal, parafoveal, and peripheral

retinotopic regions of visual space during fixation and around the time of a saccade. In addition to

this, saccade trajectories were observed in order to assess whether the planning of a saccade and

its eventual execution along with observed transient changes in visual sensitivity share an

underlying cause. Next, we investigated the rhythmicity that actively supported these transient

changes in visual sensitivity in order to assess whether they resemble known rhythmic signatures

that mediate neural activity across retinotopic brain areas during fixation and around the time of a

saccade. Finally, we proposed a novel neurobiologically inspired phenomenological force field

model that succinctly recapitulated transient changes in visual sensitivity at less (foveal) and more

eccentric (parafoveal) retinotopic of visual space.

In a dimly lit room, during fixation, head-fixed human subjects were randomly flashed

low-contrast probes (at a contrast level which the subject could detect with 50% accuracy) for

20ms at 2.5 degrees of visual angle from the fixation dot (*Experiment 1*), from the midpoint (at an

eccentricity of 5°) between the fixation dot and the fixed saccade target (*Experiment 2*), or from

the fixed saccade target (*Experiment 3*), at tangential retinotopic locations (i.e., retinotopic

locations around the presumed path of a saccade, Fig. 1A), geometrically symmetric to one another

with respect to the radial axis (i.e., the presumed path along which subjects should plan their

impending saccade). Temporally, low contrast probes were flashed either during fixation or after

the extinguishment of the fixation dot, which served as the central oculomotor cue for subjects to

make a planned saccade towards the retinotopic location along which they perceived the fixed

saccade target to be positioned (See Methods, Fig. 1B). Subjects reported with a button-press

whether they were able to detect a flashed probe. To control for artificial restriction of attentional

and oculomotor resources along the tangential axes, probes were also isotropically flashed along

the radial axis. To then control for any false responses, in 25% of all experimental trials no probes

were flashed, which yielded a false alarm rate of 1.2% along the tangential axes. Indeed, because

of the aim of our study highlighted above, perceptual and oculomotor responses obtained in both

control conditions were discarded and are therefore not included in the main analyses reported

below. With respect to target onset, flashed probes fell within the time window of 0 to

1285ms after target onset (Fig. 1B).

During fixation, the retinotopic location of the fixed saccade target appeared orthogonally

mislocalized, which reflects initial transient orthogonal shifts in gaze but does not reflect

later transient orthogonally compressed shifts in gaze.

Across experiments, after the onset of the fixed saccade target during fixation, as expected, we

observed transient shifts in gaze (*1-5*). However, specifically from 0 to 209ms after the onset of

the fixed saccade target, these transient shifts in gaze were unexpectedly averted away from the

retinotopic location of the fixed saccade target in the direction of the foveal tangential clockwise

fov_{cw} retinotopic location (Fig. 2A, *5*, *23*). From 210ms to 820ms after the onset of the fixed

saccade target (Fig. 2B), we observed that these transient shifts in gaze continued to remain

eccentric but were compressed towards the foveal tangential counterclockwise fov_{ccw} retinotopic

location. This compression in gaze, in anticipation of a planned saccade, is in part in agreement

with previous reports of peri-saccadic compression of retinotopic receptive fields toward the

periphery, prior to the execution of a planned saccade (*7*, 24). The spatial distribution between

these marked transient shifts in gaze along the radial and tangential axes was statistically

significant (rad_{shift}, p = 4.0058e-36; tan_{shift}, p = 0.0149). In all, these transient shifts in gaze

are in alignment with previous studies which report increases in neural activity in the posterior

parietal cortex (PPC) during transient shifts in gaze that were biased towards eccentric retinotopic

locations in visual space (*25*). Not surprisingly, the PPC has direct and independent connections

to the SC which is known to support the maintenance of fixation and, at the same time, initiate

transient shifts in gaze during fixation (*26*, *27*). Across experiments, we observed that the saccade

landing position, from 0 to 50ms after saccade offset (i.e., the moment the eye lands in the

periphery) was dramatically mislocalized towards the peripheral tangential clockwise $peri_{cw}$

retinotopic location (Fig. 2C). Because the oculomotor system, as it plans an impending saccade,

must estimate the spatial extent of the eccentric saccade target that is concurrently being sampled

on the peripheral retina during fixation (*28*), this observation shows that subjects erroneously

perceived the fixed eccentric saccade target in the direction of the $peri_{cw}$ retinotopic location

before the onset of the central oculomotor cue. This observation is in part in agreement with

previous reports of increases in neural activity in the PPC as the organism attempts to sample a

visual target of interest that is inherently difficult to track given the elusiveness of the visual target

(e.g., its eccentric spatial extent) or during a cued oculomotor task (*29–30*).

As expected, a larger variance was observed for the initial locus of gaze (6.84 deg^2) when

compared to the variance associated with the compressed locus of gaze (2.12 deg^2). As for the

variance associated with the saccade landing position, a larger variance (10 deg^2) was observed

when compared to the loci of the gaze in the fovea (Fig. 2D). These results strongly suggest that

the orthogonal bias in the locus of the saccade landing position (more eccentric retinotopic regions)

erroneously continued to mirror the orthogonal bias we observed in the initial transient shifts in

gaze within the fovea, even as gaze actively compressed orthogonally towards the fov_{ccw}

retinotopic location.

Transient orthogonal shifts in gaze curve the representation of visual space and, on the

perceptual domain, correlate with local asymmetric differences in visual sensitivity at

geometrically symmetric retinotopic locations.

We hypothesize that observed transient orthogonally biased shifts in gaze during fixation and the

orthogonal mislocalization of the fixed saccade target during fixation is the result of a distorted

retinotopic representation of visual space in which the foveal retinotopic region is represented

along the radial axis and, inversely, eccentric retinotopic regions are erroneously represented along

an axis that is orthogonal to the radial axis as it has been reported in previous neurophysiological

studies $(7, 31)$. This interpretation of our results implies that well before a saccade and, to an extent,

before the execution of a saccade, the retinotopic representation of visual space erroneously

includes an angular component, that is, visual space curves. From a functional perspective, since

the retinotopic representation of eccentric retinotopic regions will not reflect active visual

processing, as it is the case for the less eccentric retinotopic region, measured eccentric retinotopic

locations (i.e., $para_{cw}$ and the $peri_{cw}$ retinotopic locations) tangentially positioned in the

direction which reflect initial transient orthogonal shifts in gaze should demonstrate transient

increases in visual sensitivity. Along the same lines, measured eccentric retinotopic locations (i.e.,

$para_{ccw}$ and the $peri_{ccw}$ retinotopic locations) tangentially positioned away from initial transient

orthogonal shifts in gaze should demonstrate transient declines in visual sensitivity. With respect

to the fovea, the measured retinotopic location (i.e., the fov_{ccw} retinotopic location) tangentially

positioned in the direction towards which gaze is actively being shifted should demonstrate

transient increases in visual sensitivity, while the retinotopic location (i.e., the fov_{cw} retinotopic

location) tangentially positioned away from the target retinotopic location of the impending

transient shift in gaze should demonstrate transient declines in visual sensitivity. Conversely, if

visual sensitivity does not demonstrate local transient asymmetric differences in visual sensitivity

as we expect, then observed transient differences, assuming they will be observed, cannot be

directly attributed to a curvilinear representation of visual space.

At geometrically symmetric retinotopic locations within distinct retinotopic regions of

visual space, we observed local transient asymmetric differences in visual sensitivity (Fig. 2E-G).

Specifically, in the foveal retinotopic region, when subjects began to orthogonally compress their

gaze towards the fov_{ccw} retinotopic location, we observed transient but marked increases in visual

sensitivity at the fov_{ccw} retinotopic location (*3*), while at the fov_{cw} retinotopic location, we

observed transient but marked declines in visual sensitivity (Fig. 2E). In the parafoveal retinotopic

region, from 210ms to 820ms after the onset of the fixed target, we observed transient but marked

increases in visual sensitivity at the $para_{cw}$ retinotopic location, while at the $para_{ccw}$ retinotopic

location we observed marked declines in visual sensitivity (Fig. 2F). Finally, in the peripheral

retinotopic region, from 210ms to 820ms after the onset of the fixed target, we observed transient

but marked increases in visual sensitivity at the $peri_{cw}$ retinotopic location, while at the $peri_{ccw}$

retinotopic location, we observed transient but marked declines in visual sensitivity (Fig. 2G).

Peri-saccadic sensitivity hand-offs in the more eccentric retinotopic regions of visual space

take place around the same time orthogonal convergent effects are known to occur.

In the parafoveal retinotopic region, we also observed peri-saccadic visual sensitivity "hand-offs"

between $para_{ccw}$ and $para_{cw}$ retinotopic locations. Specifically, 1014ms after the onset of the

fixed saccade target (i.e., 70ms before saccade onset), the $para_{ccw}$ retinotopic location that once

experienced transient declines in visual sensitivity begun to experience rapid increases in visual

sensitivity, while the $para_{cw}$ retinotopic location that once experienced transient increases in

visual sensitivity began to experience rapid declines in visual sensitivity (Fig. 2F). Along the same

lines, in the peripheral retinotopic region, we observed peri-saccadic hand-offs between $peri_{ccw}$

and $peri_{cw}$ retinotopic locations. Specifically, 920ms after the onset of the fixed saccade target

(i.e., 164ms before saccade onset), the $peri_{ccw}$ retinotopic location that once experienced transient

declines in visual sensitivity begun to experience rapid increases in visual sensitivity, while the

$peri_{cw}$ retinotopic location that once experienced transient increases in visual sensitivity began to

experience rapid declines in visual sensitivity (Fig. 2G). As expected, in the foveal retinotopic

region, from 820ms onwards after the onset of the fixed saccade target, we observed rapid global

declines in visual sensitivity at both foveal retinotopic locations (Fig. 2E). On one hand, these

global declines in visual sensitivity are in agreement with foveal suppression signals originating

from the retina to retinotopic brain areas before subjects execute a saccade (*32*). On the other hand,

observed peri-saccadic visual sensitivity hand-offs strongly suggest that, around the time subjects

began to prepare a saccade, eccentric retinotopic regions of visual space began to appear to subjects

more radially. This means that retinotopic locations (i.e., $para_{ccw}$ and $peri_{ccw}$ retinotopic
locations) tangentially positioned in the direction towards which eccentric retinotopic
representation of visual space is being calibrated, experienced rapid increases in visual sensitivity,
while retinotopic locations (i.e., $para_{cw}$ and $peri_{cw}$ retinotopic locations) tangentially positioned
away from the direction towards which eccentric retinotopic representation is being calibrated,
experienced rapid declines in visual sensitivity. These observed hand-offs in visual sensitivity,
along with the earlier observed fact that fixed saccade target can be orthogonally mislocalized
away from its actual retinotopic location during fixation strongly suggest that how quickly a
curvilinear visual space is calibrated by the visual system before saccade onset will fundamentally
determine (i) the extent to which cells in the extrastriate cortex will converge along an axis that is
orthogonal to the saccade direction (*7, 31*), and (ii) the extent to which human subjects
orthogonally mislocalize flash eccentric stimuli away from or towards the retinotopic location in
which the fixed saccade target is actually positioned (*32, 33*).

Saccades that are planned during fixation, expected to travel along the radial axis, deviate
along a curved path towards orthogonal eccentric retinotopic locations positioned beyond
the spatial extent of future post-saccadic retinotopic locations.

Following these observations on the perceptual domain, a natural question one might ask is
whether a curvilinear retinotopic representation of visual space accompanies any oculomotor
consequences, specifically around the time of a saccade. Keeping in mind that a curvilinear
representation of visual space predicts that eccentric retinotopic regions of visual space are
represented along an axis that is orthogonal to the radial axis, coupled with the fact that this
representation precedes and, in some cases, temporally overlaps with when subjects begin to plan
and prepare for an impending saccade (i.e., ~150 to 200ms before saccade onset) (*34, 35*), we
predict that saccades on average will deviate along a curved path towards the $peri_{cw}$ retinotopic
location – the retinotopic location towards which subjects on average mislocalized the fixed
saccade target (Fig. 2C). We further predict that because curved visual space is eventually
calibrated back towards $para_{ccw}$ and $peri_{ccw}$ retinotopic locations, as the hand-offs in peri-
saccadic visual sensitivity observed in our perceptual experiences suggest, some saccades should
also curve towards the $peri_{ccw}$ retinotopic location. Assuming our predictions are correct, this
would further strengthen the idea that the visual system is not exclusively spatially selective for

future post-saccadic retinotopic locations that are uniquely parallel to the radial axis (i.e., the path

saccades are expected to travel along, *36, 37, 38*), but can actively select retinotopic locations that

are positioned beyond these fixed post-saccadic retinotopic locations, for example, retinotopic

locations orthogonal to the radial axis that convey task-relevant information (*21, 22*).

255   Aligned with this prediction, we observed that trajectories towards the periphery

systemically deviated along a curvilinear path across experiments, at a rate of 73% towards the

$peri_{cw}$ retinotopic location. In 27% of all trials, saccades were curved towards the $peri_{ccw}$

retinotopic location. Specifically, when the saccade target was presented at a fixed azimuth angle

of 0°, the trajectories were curved towards the $peri_{ccw}$ retinotopic location at a rate of 91% with

an average curvature of 0.22 in area/amp. Conversely, for the remaining trials (saccades curved

towards the $peri_{ccw}$ retinotopic location), an average curvature of -0.80 in area/amp was observed.

When the saccade target was presented at a fixed azimuth angle of 45°, the trajectories were curved

towards the $peri_{cw}$ retinotopic location at a rate of 65% with an average curvature of 0.56 in

area/amp, while for the remaining trials (saccades curved towards the $peri_{ccw}$ retinotopic

location), an average curvature of -0.74 in area/amp was observed. Finally, when the saccade target

was presented at a fixed azimuth angle of 315°, trajectories were curved towards the retinotopic

$peri_{cw}$ retinotopic location at a rate of 60% with an average curvature of 1.12 in area/amp, while

for the remaining trials (saccades curved towards the retinotopic location $peri_{ccw}$ location), an

average curvature of -1.17 in area/amp was observed.

Delta-rhythmicity known to suspend neural resources and mediate neural activity in

retinotopic brain areas also supports local asymmetric differences in visual sensitivity at

geometrically symmetric retinotopic locations.

So far, we have provided an account of the functional correlates (i.e., the perceptual and

oculomotor consequences) of curved visual space, which arise from transient orthogonal shifts in

gaze during fixation. We posited that these transient orthogonal shifts occur because cells that

support the more eccentric retinotopic regions of visual space are devoid of neural resources they

need to correctly (or actively) process the visual world. This results in an erroneous representation

of eccentric retinotopic regions along an axis that is orthogonal to the radial axis which (i) gives

rise to asymmetric differences in visual sensitivity at geometrically symmetric retinotopic

locations at less and more eccentric retinotopic regions of visual space, and (ii) curves saccade

trajectories expected to travel along a linear path. Indeed, these effects can also account for
orthogonally biased translational and orthogonal convergent effects researchers have reported in
previous studies. Here, we hypothesize that the neural resources these eccentric retinotopic cells
once possessed are temporarily allocated toward orthogonal foveal sites (*15-20*), at and around the
fov_{cw} retinotopic location, for example. To test this hypothesis, we investigated whether changes
in visual sensitivity are uniquely supported by the same spectral signatures which suspend neural
resources during cognitively demanding tasks (*39*) and which are known to mediate neural activity
in brain areas that specifically support fixational eye movements and/or the planning and
preparation of a saccade (e.g., frontal and parietal cortex, *6, 12, 25, 29, 30, 40*). To this end, we
transformed our detrended raw time-series visual sensitivity data into the frequency domain. The
logic behind this investigation is as follows: if the changes in visual sensitivity we measured at
retinotopic locations are supported by rhythms which are known to suspend the spreading of neural
resources (with the exception of the fov_{cw} retinotopic location), then the spectral signature
associated with these changes in visual sensitivity should strongly resemble the known rhythmic
signatures (i.e., delta rhythms), especially at more eccentric retinotopic locations. Conversely, if
the spectral signatures we observe do not recapitulate these known rhythmic signatures, then these
transient changes cannot be directly attributed to the suspension of neural resources at the more
eccentric retinotopic regions of visual space.

At the fov_{ccw} retinotopic location, we observed that these transient changes in visual
sensitivity were actively mediated by rhythmicity in the delta frequency band at ~2 Hz. At the
fov_{cw} retinotopic location, we observed that these transient changes in visual sensitivity were
actively being mediated by rhythmicity at ~3 Hz. This higher rhythmicity is known to mediate the
allocation of attentional signals that support visual sensitivity at suprathreshold levels (*41–43*).
Furthermore, the spectral signature that mediated the fov_{ccw} retinotopic location was not location-
specific when compared to shuffled data sets (p=0.2280). Conversely, the spectral signature that
supported the fov_{cw} retinotopic location was specific to that retinotopic location (p=0.0230). The
$para_{cw}$, $para_{cw}$, $peri_{ccw}$, and $peri_{ccw}$ retinotopic locations also exhibited rhythmicity in the ~2
Hz delta frequency band (Fig. 3C & 3E). The spectral signature observed at these eccentric
clockwise and counterclockwise retinotopic locations were not location-specific (p=0.0915,
p=0.9725, p=0.3179, p=0.9725) (Fig. 3D & 3F). Furthermore, the phase of ~2 Hz within $para_{cw}$
and $para_{ccw}$ retinotopic locations was consistently offset by approximately 201°, and even larger
offset of 290° was observed at the more eccentric $peri_{cw}$ and $peri_{ccw}$ retinotopic locations (Fig.

3G & 3H). This antiphase periodicity at ~2 Hz further strengthens the idea of peri-saccadic visual

sensitivity hand-offs, suggesting that the retinotopic brain areas which support these geometrically

symmetric eccentric tangential retinotopic locations are synchronized by attentional dynamics at a

common frequency, but with location-specific phases (*44*).

Neurobiologically inspired retinotopic lever under an inverse force field.

Our spectral results suggest that observed transient changes in visual sensitivity may indeed be the

consequences of signals that suspend attentional resources at the more eccentric retinotopic

locations and directly enhance active foveal visual processing, namely at the fov_{cw} retinotopic

location. Remarkably, at this retinotopic location, rhythmic visual sensitivity patterns were

uniquely spatially selective. It is also this location towards which subjects initially shifted their

gaze after the onset of the fixed saccade target. Finally, it is from this very same retinotopic

location that attentional resources must be shifted towards the fov_{ccw} retinotopic location in order

to enhance visual sensitivity there (*3, 7, 24*).

With these insights from our spectral investigations, along with the fact that population

receptive fields (pRFs) across retinotopic brain areas can transiently shift beyond the spatial extent

of their classical receptive fields, we developed a simple but powerful novel neurobiologically

inspired phenomenological force field model (See Methods). From an architectural standpoint, our

model includes retinotopic pRFs that tile foveal, parafoveal, and peripheral retinotopic regions of

visual space. With the context of our model, these pRFs can be understood as spring-loaded

sensors. Another layer in our model is a force field, which is composed of forces, exerted by

corresponding time-varying masses that act on pRFs. Finally, the third layer of our model consists

of retinotopic population elastic fields (pEFs) which all pRFs possess in order to always maintain

retinotopic organization during transient retinotopic shifts in visual space (e.g., Fig 4A).

Functionally, at its core, our model assumes that underlying neural (i.e., attentional and

oculomotor) signals, are actively modulated by transient eccentric error signals (*25, 29, 30*). The

summation of these signals temporarily suspends neural resources at the more eccentric retinotopic

regions of visual space which causes the visual system to erroneously represent these regions along

an axis that is orthogonal to the radial axis while enhancing active visual processing at tangential

foveal retinotopic locations (*20*). This gives rise to temporally overlapping input and output forces

exerted by corresponding time varying masses positioned at fov_{ccw} and $peri_{cw}$ retinotopic

locations respectively (Fig. 4B & 4E), which then act upon a "retinotopic lever". Under an inverse
force field (ε = -0.55, see Methods), input and output forces transiently shift pRFs along an axis
that is orthogonal to the radial axis and thereby shape population density (putative population
sensitivity) at a sampled retinotopic location in time (Fig. 4A, Fig. 4D, and Extended Data Fig. 1–
2).
352
Spatio-temporal retinotopic shifts that curve visual space under an inverse-distance rule
explain local asymmetric transient differences in visual sensitivity at geometrically
symmetric locations in less eccentric and more eccentric retinotopic regions.

As far as the foveal simulations, attentional and oculomotor signals modulated by transient
eccentric error signals give rise to an input force that is preparing to act on the input arm of a
"retinotopic lever" which causes pRFs to orthogonally shift in the direction of the fov_{cw}
retinotopic location and later compress towards the fov_{ccw} retinotopic location. A compressed
input force causes the lever to pivot on its "foveal fulcrum" along a curvilinear path thereby
providing a transient neural advantage at the fov_{ccw} retinotopic location. To always maintain
retinotopic organization, pRFs that tile the more eccentric retinotopic regions in visual space are
initially restricted from over-translating towards the foveal fov_{cw} retinotopic location by their
pEFs, which remain mirroring the onset location of the input force before a full compression
occurs. For the sake of simplicity, foveal simulations began from the point in time when pRFs
mirrored the initial transient orthogonal shift in gaze (translational output force), followed by late
transient compressive shift in gaze towards the fov_{ccw} retinotopic location (compressed input
force) (Fig. 4B). Under these input and output forces, as expected, simulation results in Fig. 4C
show that putative population sensitivity recapitulated signatures in transient changes in visual
sensitivity in the less eccentric (foveal) retinotopic regions of visual space (Fig. 2E). Assuming
that retinotopic cells are not mediated by transient eccentric error signals (i.e., retinotopic
representation does not include a curvilinear component), translational shifts researchers once
thought anticipated retinal image disruptions saccades accompany (*11-13*) should have yielded no
difference in putative population sensitivity (Fig. 4C, dotted marker) in comparison to the transient
changes in visual sensitivity we observed in our behavioral experiments (Fig. 2E). Similarly,
symmetric convergent retinotopic shifts towards a fixed peripheral site (Fig. 4C, thin markers, *6*)
cannot explain observed transient changes in visual sensitivity.

As far as the eccentric simulations (Fig. 4D), after a fully compressed input force, pRFs

moved further towards the fov_{ccw} retinotopic location in proportion to the mislocalization of the

fixed saccade target. Cells across retinotopic brain areas can receive both convergent and

translational signals around the time of a saccade (*6, 12*). To this end, pRFs later converged around

an emerging output mass which exceeded the magnitude of the initial transient orthogonal shift in

gaze and approximated the retinotopic location of the mislocalized target (Fig. 2A & 2C). This is

immediately followed by the onset of a translational force proportional to the magnitude of a

saccade orthogonal to the radial axis towards the $peri_{cw}$ retinotopic location (Fig. 4E). Indeed,

under these three forces, especially when the forces that correspond to error modulated convergent

signals emerge, a neural advantage at eccentric clockwise retinotopic locations should occur. As

expected, Fig. 4F shows that putative population sensitivity was able to recapitulate transient visual

sensitivity signatures at parafoveal retinotopic locations (Fig. 2F). In all, as expected, a purely

translational or symmetric convergent retinotopic shifts cannot explain observed transient changes

in visual sensitivity.

DICUSSION

There is now a growing consensus that fixational eye movements actively mediate visual

perception well before they occur (*e.g., 2, 3*). For example, in 2011, Hafed found that before the

onset of fixational eye movements the retinotopic representation of visual space compresses

towards the site of the impending fixational eye movement. This peri-mircosaccadic compression

of visual space causes human subjects to erroneously perceive foveal stimuli as more eccentric

and conversely eccentric stimuli as more foveal. However, while some of the effects of fixational

eye movements on visual perception well before and even after they occur have been uncovered

(e.g., *5*), the extent to which transient shifts in gaze actively mediate visual sensitivity at retinotopic

locations that constitute distinct retinotopic regions of visual space remained unknown. Equally

unaccounted for is the influence of transient shifts in gaze on oculomotor planning and

performance around the time of a saccade. To provide an account from a functional perspective,

we systematically measured the consequences of transient shifts in gaze during fixation on visual

sensitivity at geometrically symmetric retinotopic locations in the foveal, parafoveal, and

peripheral regions of visual space. Next, we assessed how these transient shifts in gaze mediate

the planning and execution of a saccade trajectory. We show for the first time that these transient

shifts in gaze give rise to local asymmetric differences in visual sensitivity at geometrically

symmetric retinotopic locations, while saccade trajectories deviate along a curvilinear path

towards an orthogonal mislocalized peripheral site (Fig. 2E & 2H). We later constructed a

phenomenological force field model in which we proposed that transient shifts in gaze are

underlined by attentional and oculomotor signals that are actively modulated by transient eccentric

error signals. The summation of these signals manifests as temporally overlapping forces which

act on cells across retinotopic brain areas and thereby modulate their population sensitivity

equivalent to changes in visual sensitivity in our psychophysical experiments. Putative population

sensitivity signatures recapitulated local asymmetric differences in visual sensitivity we observed

at geometrically symmetric retinotopic locations in less and more eccentric retinotopic regions of

visual space (Fig. 2E & 2F, Fig. 4C, and Fig. 4F).

Apart from uncovering the effects of transient shifts in gaze on visual sensitivity and

saccade trajectory, our behavioral and simulation results provide three fundamental insights into

how early retinotopic representation actively modulates peri-saccadic representation. First, peri-

saccadic representation of visual space reflects an overriding influence of attentional and

oculomotor signals which are actively modulated by transient eccentric error signals and are not

the result of purely translational or symmetrically convergent signals, as suggested by previous

studies (*6, 11, 12, 21*). Indeed, the presence of or the gating off of these eccentric error signals

seems to determine (i) whether flashed stimuli will be orthogonally mislocaized away from or

towards a fixed saccade target (Fig. 2F & 2G, *32,33*), (ii.) whether cells in the extrastriate cortex

will align with or shift along an axis that is orthogonal to the radial axis (*7, 8*) and (iii) the extent

to which peri-saccadic retinotopic shifts will translate towards future post-saccadic retinotopic

locations or shift towards tangential retinotopic locations positioned beyond the spatial extent of

future post-saccadic sites (Fig. 2H, *22*). Second, keeping in mind that the purely translational (or

forward) account of spatial constancy which researchers once thought could explain this retinal

phenomenon (*13, 21*) has been challenged on methodological grounds (*6, 7, 24*), our results go

one step further and lead to a proposal of a novel mechanism that can explain spatial constancy.

Specifically, our results strongly suggest that well before a saccade and, to some extent, around

the time of a saccade, the visual system transiently allocates visual resources that would otherwise

support active visual processing at intermediate eccentric retinotopic locations towards the fovea,

resources, which are then then used to further enhance active visual processing at specific foveal

retinotopic locations. This results in a drastic diminishing of active visual processing at

intermediate eccentric retinotopic regions of visual space. Furthermore, because peri-saccadic
visual resources spread at a glacial pace (*36, 45*), we posit that resources which eventually revert
back towards intermediate eccentric retinotopic locations arrive well after the execution of a rapid
movement towards the periphery (*46*). As opposed to translating visual resources to fixed post-
saccadic future retinotopic locations, we posit that the visual system transiently shifts and
compresses resources backwards (i.e., in the opposite direction of the fixed saccade target, *5, 23*),
away from intermediate retinotopic locations likely to contain retinal disruptions. Because of this
difference in speed, that is, between eye movement during a saccade and the spread of eccentric
peri-saccadic visual resources to their equilibrium positions, our perception of the visual world can
escape the retinal disruptions that necessarily accompany rapid eye movements. Finally, the neural
computations that support this retinotopic representation well before and around the time of a
saccade obey an inverse-distance rule, while the neural architecture that mediates spatiotemporal
retinotopic shifts pRFs undergo – which give rise to this representation – includes pEFs.

In all, the observed transient shifts in gaze during fixation remain a curious point, since
they are directed away from the fixed saccade target. However, considering that humans were once
preys, it is possible that these miniature shifts away from the visual target of interest reflect an
evolutionarily conserved visual oculomotor avoidance or escape strategy the mammalian visual
system used to track a distal target (e.g., conspecific, predator) from a safe distance (e.g., corner
of the eye), later followed by a calculated movement (e.g., saccade) towards the retinotopic
location previously avoided (*47–49*).

AUTHOR CONTRUBITIONS

EL conceptualized the project. EL collected the data. EL preprocessed the behavioral data with the
assistance of XZ. EL analyzed the behavioral data. EL developed the computational model. EL
implemented the model. EL performed the model simulations. EL wrote the manuscript.

ACKNOWLEDGEMENTS

We would like to thank D. P. Melcher and S. Wirth for comments on the manuscript and B.
Wichterlová for editing the manuscript. We would also like to thank A. S. Nandy and M. Scudder
for their guidance on the experimental design and implementation.

COMPETING INTEREST

Authors has no competing financial interest.

DATA AVAILABILITY

The data that supports this study is available upon reasonable request.

CODE AVAILABILITY

The custom Matlab code that supports the figures reported in this study is available upon

reasonable request.

SUPPLEMENTAL TEXT

Fig 1. Behavioural protocol. (A) Markers in the illustration of visual space show all retinotopic

locations at which a probe was flashed before and after the deployment of attention, saccade

planning, and, finally, the execution of a saccade. Probes of interest, presented along the tangential

axes, are conspicuously highlighted in open circle markers. **(B)** Temporal sequence of a valid trial

across experiments with respect to target onset.

Fig 2. Behavioural data. (A-D) Probability density map of foveal gaze and saccade landing was

calculated using a bivariate gaussian kernel along with their bivariate contour ellipse area estimates

for each probability density map. Error estimates are standard error of means across observers. **(E-**

G) Shifts in visual sensitivity along tangential axes with respect to target onset; error estimates

were calculated using a 20-fold jackknife procedure. **(H)** Mean saccade trajectory deviation for

horizontal and oblique saccades across experiments. Positive values on the ordinate denote

trajectory deviation towards a clockwise location (cw), while negative values denote trajectory

deviation towards a counterclockwise location (ccw)

Fig 3. Frequency domain representation of visual sensitivity and cross-location phase

coherence estimates. (A-F) Representation of raw detrended visual sensitivity at less and more

eccentric locations in visual space. Shuffled (n=1000, dashed line) versus unshuffled (solid line)

data set were used to compute location-specific spectral signatures at a $p<0.05$. (**G-H**) For

parafoveal and peripheral clockwise and counterclockwise locations a cross-location phase

coherency analysis was performed.

Fig 4. Model architecture and neurobiological force field with simulated putative

population sensitivity. (A) For illustration purposes, a single eccentricity dependent population

receptive field (pRF) within a retinotopic field is highlighted in pink. A pRF is defined by its

center, size, and an eccentric dependent population elastic field (pEF). **(B)** MGα denotes gaze

transitional signals modelled as a virtual mass at infinity located in the direction of an upper

tangential location approximating the initial shift in gaze. MF denotes compressive signals

modelled as a varying mass located at a lower tangential location in the fovea. **(C)** Simulation

results in the foveal region when a pRF is under the influence of a resultant force exerted by (i)

MF and MGα (curved space, thick marker) (ii) MC (convergent signals modelled as a varying

mass at a stationary peripheral site; see Methods) and (iii) MT (saccade transitional signals

modelled as a virtual mass at infinity located in the direction of a straight saccade; see Methods).

(D) A pRF bounded by pEF is highlighted in pink for parafoveal simulations. **(E)** MGβ denotes

gaze translational signals modelled as a virtual mass at infinity located in the direction of a

downward shifts in gaze, MM denotes a varying mass at a mislocalized peripheral target, and

MS denotes the curved transitional signals modelled as a virtual mass at infinity located in the

direction of a curvilinear saccade. **(F)** Curved, convergent and translational simulation results.

Extended Data Fig. 1 Spatiotemporal retinotopic shifts for foveal simulations. (A)

hexagonal retinotopic field spanning the entire visual field. Retinotopic shifts are produced by

the perturbation of pRFs under an inverse force field in discrete time increments of 1ms by the

combination of external forces exerted by **(B)** corresponding masses ($M_{G\alpha}$ +M_F).

Extended Data Fig. 2 Spatiotemporal retinotopic shifts for parafoveal stimulations. (A)

Retinotopic shifts when the combination of external forces is exerted by **(B)** corresponding

masses ($M_{G\beta}$ +M_M + M_S).

METHODS

Human subjects

Six consenting human subjects (5 females and 1 male; 21-26 years old) with corrected-to-normal

visual acuity participated in the three experiments reported as it is standard within the field of

research. Three of the six subjects participated in the foveal and parafoveal experiments. The

remaining three subjects participated in the peripheral experiment. The Yale Ethics Review board

approved the protocol of these experiments and the storage of the data that was collected.

Visual stimuli and experimental protocol

Fig. 1 summarizes the protocol used across experiments. In a dimly lit room, stimuli were

presented on a gamma corrected display with a spatial resolution of 1400 x 1050 pixels and a mean

luminance of 38cd/m². Eye movements were recorded using an infrared video-based eye-tracker

sampled at 1kHz (I-Scan Inc., Woburn, MA). While tightly head fixed using a head rest and a

customized bite bar, subjects at the onset of each trial acquired fixation for 300ms. Fixation within

this temporal window was spatially aligned with the fixation point (0.5 dva in diameter) using a

custom-developed 9-point eye-calibration procedure in Picto. Next, depending on the subject, a

saccade target was presented at an eccentricity of 10° and at azimuth angles of 0° (for s_1), 45° (for

s_2) and 315° (for s_3) (Fig. 1A). Following target onset, fixation was maintained for another 300-

600ms within ~2dva fixation tolerance window, at the end of which the fixation spot was

extinguished, which served as the central motor cue for subjects to make an eye movement towards

the saccade target. For another 500ms after the central motor cue, the saccade target remained

visible. Finally, a low contrast probe (at a contrast level which the subject could detect with 50%

accuracy) were isotropically flashed, depending on the experiment, for 20ms either along the radial

or tangential axis at a random time during the 300-600ms window between the target onset and

the central motor cue (25% of trials) or at a random time between 0-340ms after the central motor

cue (50% of the trials). On the remaining 25% (which yielded a false alarm rate of 1.2%), no

probes were presented. All three conditions were randomly interleaved across trials. With respect

to target onset, the presentation of flashed probes fell within a time window of 0-1285ms after

target onset (Fig. 1B). Subject was asked to report, using a push button, whether they were able to

detect the flashed probe. While the responses along the radial axis were not analyzed, the

presentation of these probes was crucial in controlling for any response biases that may have

emerged along the tangential axes. To ensure that attentional and oculomotor resources was not

being driven by an asymmetric stimuli configuration as it is the case in previous studies, no

secondary salient spatial cue was presented. Along the tangential axes, a total of ~6010 trials were

collected across experiments, with a minimum of 650 trials per subject.

Statistical analyses

To investigate gaze dynamics, probability density maps and variance estimates were calculated

using median filtered edge preserving fixational data, which was rotated to an angle of 0° across

target locations. Next, I calculated a gaze probability density map for the foveal and periphery

regions using a bivariate Gaussian kernel (*50*). The estimated locus of gaze across subjects was

defined as the peak of each probability density map (i.e., the yellow contour lines). To compare

the distribution of gaze in the fovea during the 0 to 209ms window after target onset versus 210 to

820ms window after target onset, a Mann-Whitney U statistic was calculated at p value less than

0.05. Further, variance in gaze was calculated using a bivariate Contour Ellipse Area (BCEA)

approach which included 68% of fixational data around the mean (*51*). Accompanying error bars

were obtained which denote the standard error of the mean variance across subjects. To investigate

the extent saccade trajectories were curved across experiments at each saccade location. Horizontal

and vertical eye positions, with the time window starting point constituting the moment fixation

was maintained to the moment the eye landed in the periphery, was isolated. Next to obtain a

measurement of curvature, the curved area that emerged from the eye position during this time

window was calculated. To then obtain a ratio measure of curvature per unit amplitude, this area

was divided by the amplitude of the saccade (*52*). Accompanying calculated error bars denote the

standard error of the mean at each target location across experiments. To investigate normalized

changes in visual sensitivity across subjects, I first binned each subject's percent correct responses

for the foveal (Exp. 1), parafoveal (Exp. 2), and peripheral (Exp. 3) experiments, using a 15ms

sliding window that was moved in 45ms increments. Next, I obtained each subject's raw visual

sensitivity, using a 20-fold jackknife procedure by estimating sensitivity from 95% of the data.

This was repeated 20 times, each time leaving out 5% of the data. This produced a mean and an

error estimate of visual sensitivity at each tangential location for each experiment across subjects

(*53*). To control for any eccentric-dependent effect on raw visual sensitivity, these traces were

normalized by the average sensitivity over the initial 100ms in the data. To investigate frequency

domain and cross-location phase coherence dynamics, a fast Fourier transform (FFT) was applied

to the detrended raw visual sensitivity data. Next, we computed the mean of the spectral signatures

and corresponding standard error of the mean across subjects. To than assess whether the observed

periodicities were location specific, we randomly shuffled (n=1000) the probe location identities

across trials and calculated the sensitivity function and periodicity of the shuffled data using the

same procedures noted above. A two tailed paired-sample t-test at p-values of 0.05 was calculated

to assess the significance between shuffled and non-shuffled data. Finally, for the cross-location

phase coherence analysis, we applied an approach to the shuffled and non-shuffled detrended data

that is commonly used in assessing the underlying coherence in neural data (*54*). I then projected

the difference between phase angles of each subject between counterclockwise (ccw) and

clockwise (cw) locations and then calculated the resultant vector across subjects.

Novel computational framework - Retinotopic mechanics

The underlying architecture of this simple mechanical model is composed of a retinotopic field

(ϕ_r) and a retinotopic lever which is inspired by a neurobiological force field. The retinotopic field

is a two-dimensional overlapping hexagonal grid of eccentricity-dependent receptive fields - RF_i

- that tiled the foveal, parafoveal and peripheral regions in visual space (Extended Fig. 1-2). Each

RF_i was defined by three parameters: (a) p_i, the spatial extent of the RF center (x_i, y_i) at t_0 (when

there is no resultant force acting on the ϕ_r). p_i determines the eccentricity - e_i of RF_i, (b) s_i, the

radius of RF_i is proportional to e_i, and (c) the movement extent (M_{max}) of RF_i is its EF_i which is

also proportional to e_i. ϕ_r emerged from attentional and oculomotor signals actively modulated by

transient eccentric error signals that act as input and output forces of retinotopic lever in this field,

and transiently perturb a constellation of population receptive fields from equilibrium and thereby

modulate their putative population sensitivity (density readouts) as a function of space and time.

For simulations at the less eccentric ("foveal"), the resultant force $(\overrightarrow{F_{R1}})$ included an input and

output force $(\overrightarrow{F_I}$, and $\overrightarrow{F_O})$ and an internal force $(\overrightarrow{F_E})$ which RF_i at t_{0+n} experiences:

623
$$\overrightarrow{F_O} = \overrightarrow{U_{M_{G\alpha}}} \cdot M_{G\alpha} \cdot |G\alpha|$$

624
$$\overrightarrow{F_I} = \overrightarrow{U_{M_F}} \cdot M_F \cdot R_\beta^\varepsilon$$
625
626
$$\overrightarrow{F_{R1}} = \overrightarrow{F_I} \cdot k_I + \overrightarrow{F_O} + \overrightarrow{F_E}$$
627
628
where $M_{G\alpha}$ and M_F are the virtual mass at infinity in the upward direction which approximates an

initial shift in gaze and a foveal mass located at a lower tangential foveal retinotopic site which

approximates a compressed shift in gaze respectively. $\overrightarrow{U_{M_{G\alpha}}}$ and $\overrightarrow{U_{M_F}}$ are the unit vectors in the

direction of the virtual mass and the foveal mass respectively. $|G\alpha|$ is the magnitude in gaze

towards the upper tangential foveal retinotopic site, while R_β is the spatial difference between RF_i

and M_F raised to an epsilon scalar – ε. The value of ε, depending on the retinotopic mechanics

that best explains the phenomenon in question, can either be consistent with a proportional-

distance rule (i.e., >0) or an inverse-distance rule (i.e., <0). Further, k_I was obtained by calculating

the average of $R_{\beta 0}^{\varepsilon}$ across RF_i's and then taking its reciprocal $\frac{1}{\left(R_{\beta 0}^{\varepsilon}\right)}$, while $\overrightarrow{F_E}$ is proportional to

the movement of RF_i. For the more eccentric ("parafoveal") simulations, the resultant force $(\overrightarrow{F_{R2}})$

included an input, output, an additional orthogonally biased translational force $(\overrightarrow{F_I}, \overrightarrow{F_O}$ and $\overrightarrow{F_T})$

and an internal force $(\overrightarrow{F_E})$ which RF_i at t_{0+n} experiences:

$$\overrightarrow{F_I} = \overrightarrow{U_{M_\beta}} \cdot M_{G\beta} \cdot |G\beta|$$

$$\overrightarrow{F_O} = \overrightarrow{U_{M_M}} \cdot M_M \cdot R_\mu^\varepsilon$$

$$\overrightarrow{F_T} = \overrightarrow{U_{M_S}} \cdot M_S \cdot |S|$$

$$\overrightarrow{F_{R2}} = \overrightarrow{F_O} \cdot k_O + \overrightarrow{F_I} + \overrightarrow{F_T} + \overrightarrow{F_E}$$

where $M_{G\beta}$ and M_M are the virtual mass at infinity in the direction towards lower foveal tangential

retinotopic site which approximates a complete compression of gaze and the mislocalized mass

located at a peripheral retinotopic site that approximates a mislocalized saccade target respectively.

$\overrightarrow{U_{M_{G\beta}}}$ and $\overrightarrow{U_{M_M}}$ are the unit vectors in the direction of the virtual mass and the mislocalized

peripheral mass respectively. $|G\beta|$ is the magnitude of a fully compressed gaze which is roughly

proportional to the magnitude of a mislocalized target. R_μ is the spatial difference between RF_i

and M_M raised to ε, while $|S|$ approximates the magnitude of an impending saccade. The same

calculation used to obtain k_I was used to obtain k_O, however in this case, the average of $R_{\mu 0}^{\varepsilon}$

across RF_i's was used. Depending on the stimulation, a time varying resultant force $(\overrightarrow{F_{R1}}$ or $\overrightarrow{F_{R2}})$

perturbed each RF_i, which produced spatiotemporal retinotopic displacements (Extended Data Fig.

1-2). The population movements of the constellations of RF_i's manifested in time-varying

modulation of raw density, modeling each RF_i as a bivariate gaussian kernel function G *(55)*. A

probability density estimate (or putative population sensitivity readouts), $\hat{f}$, was obtained using

the following equation:

$$G\left(u\right) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2} u^{\mathrm{T}} u}$$

$$\hat{f}\left(\boldsymbol{x}, \mathbf{B}\right) = \frac{1}{k} \sum_{i=1}^{k} G_{\mathbf{B}}\left(|\boldsymbol{x} - \boldsymbol{x_i}|\right)$$

668 Finally, population sensitivity readouts were later normalized using the same approach used to

669 normalize the raw visual sensitivity traces.

670
REFERENCES

1. Ratliff F, Riggs LA. Involuntary motions of the eye during monocular fixation. *J Exp*
Psychol. 40:687–701. (1950).

2. Hafed, Z. M. Alteration of visual perception prior to microsaccades. *Neuron*. 77, 775–
786. (2013).

3. Shelchkova N. Poletti M. Modulations of foveal vision associated with microsaccade
preparation. *Proc. Natl. Acad. Sci. USA* 117: 11178-11183. (2020).

4. Chen, C. Y., Ignashchenkova, A., Thier, P., and Hafed, Z. M. Neuronal response gain
enhancement prior to microsaccades. *Curr. Biol.* 25, 2065–2074. (2015).

5. Tian, X., Yoshida, M., Hafed, ZM. A microsaccadic account of attentional capture and
inhibition of return in Posner cueing. *Front Syst Neurosci*. 10:23. (2016).

6. Zirnsak, M., Steinmetz, N. A., Noudoost, B., Xu, K. Z., Moore, T. Visual space is
compressed in prefrontal cortex before eye movements. *Nature*, 507, 504–507. (2014).

7. Tolias, A. S., Moore, T., Smirnakis, S. M., Tehovnik, E. J., Siapas, A. G., Schiller, P. H.
Eye movements modulate visual receptive fields of V4 neurons. *Neuron*, 29(3), 757–767.
(2001).

8. Connor, C. E., Gallant, J. L., Preddie, D. C., Van Essen, D. C. Responses in area V4
depend on the spatial relationship between stimulus and attention. *Journal of*
Neurophysiology, 75(3), 1306–1308. (1996).

9. Kaiser, M., Lappe M. Perisaccadic mislocalization orthogonal to saccade direction.
Neuron, 22;41(2), 293–300. (2004).

10. Zimmermann, E., Morrone, M. C., Burr, D. C. Visual mislocalization during saccades
sequences. *Exp Brain Res*, 233(2):577-585. (2014).

11. Duhamel, J. R., Colby, C. L., Goldberg, M. E. The updating of the representation of
visual space in parietal cortex by intended eye movements. *Science*, 255(5040), 90–92.
(1992).

12. Umeno, M.M., Goldberg, M.E. Spatial processing in the monkey frontal eye field. I.
Predictive visual responses. *J Neurophysiol,* 78:1373–138. (1997).

13. Wurtz R. H. Neuronal mechanisms of visual stability. *Vision research*, *48*(20), 2070–
2089. (2008).

14. Idrees, S., Baumann, M.P., Franke, F., Munch, T.A., and Hafed, Z.M. Perceptual
saccadic suppression starts in the retina. *Nat*. *Commun*. 11, 1977. (2020).

15. Hamker, FH., Zirnsak, M., Calow, D., Lappe, M. The peri-saccadic perception of objects
and space. *PLoS Comput Biol, 4: e31*. (2008).

16. Richard, A., Churan, J., Guitton, DE., Pack, CC. The geometry of perisaccadic visual
perception. *J Neurosci*, 29: 10160 –10170, (2009).

17. VanRullen, R. A. simple translation in cortical log-coordinates may account for the
pattern of saccadic localization errors. *Biol Cybern, 91*: 131–137. (2004).

18. Zirnsak, M., Lappe, M., Hamker, FH. The spatial distribution of receptive field changes
in a model of peri-saccadic perception: predictive remapping and shifts towards the
saccade target. *Vision Re*s, 50: 1328 –1337. (2010).
729

730   19. Hafed, Z.M., Chen, C.Y., Tian, X. Vision, perception, and attention through the lens of
731   microsaccades: Mechanisms and implications. *Front. Syst. Neurosci*, 9, 167. (2015).
732

733   20. Chen, C. Y., Hoffmann, K. P., Distler, C., Hafed, Z. M. The Foveal Visual
734   Representation of the Primate Superior Colliculus. *Current biology, 29*(13), 2109–
735   2119.e7. (2019).
736

737   21. Zirnsak, M., Moore, T. Saccades and shifting receptive fields: anticipating consequences
738   or selecting targets? *Trends in cognitive sciences*, 18,12, 621-628. (2014).
739

740   22. Arkesteijn, K., Belopolsky, A. V., Smeets, J., Donk, M. The Limits of Predictive
741   Remapping of Attention Across Eye Movements. *Frontiers in psychology*, 10, 1146.
742   (2019).
743

744   23. Hafed, Z. M., and Clark, J. J. Microsaccades as an overt measure of covert attention
745   shifts. *Vision Res*, 42(22):2533-45. (2002).
746

748   24. Hartmann, T.S., Zirnsak, M., Marquis, M., Hamker, F.H., Moore, T. Two types of
749   receptive field dynamics in area V4 at the time of eye movements? *Front. Syst. Neurosci*,
750   11, 1–7. (2017).
751

25. Andersen, R. A., Mountcastle, V. B. The influence of the angle of gaze upon the
excitability of the light-sensitive neurons of the posterior parietal cortex. *The Journal of*
Neuroscience, 3(3), 532–548. (1983).

26. Munoz, D. P., and Wurtz, R. H. Fixation cells in monkey superior colliculus. I.
Characteristics of cell discharge. *J. Neurophysiol*. 70, 559–575. (1993).

27. Hafed, Z. M., Goffart, L., Krauzlis, R. J. A neural mechanism for microsaccade
generation in the primate superior colliculus. *Science*, Vol. 323, No. 5916, pp. 940-943.
(2009).

28. Deubel, H., Bridgeman, B., Schneider, WX. Immediate post-saccadic information
mediates space constancy. *Vision Res* 38:3147–3159. (1998).

29. Assad, J. A., Maunsell, J. H. Neuronal correlates of inferred motion in primate posterior
parietal cortex. *Nature*, 373(6514), 518–521. (1995).

30. Zhou, Y., Liu, Y., Lu, H., Wu, S., Zhang, M. Neuronal representation of saccadic error in
macaque posterior parietal cortex (PPC). *eLife*, 5. (2016).

31. Connor, C. E., Gallant, J. L., Preddie, D. C., Van Essen, D. C. Responses in area V4
depend on the spatial relationship between stimulus and attention. *Journal of*
Neurophysiology, 75(3), 1306–1308. (1996).

32. Kaiser, M., Lappe M. Perisaccadic mislocalization orthogonal to saccade direction.
Neuron, 22;41(2), 293–300. (2004).

33. Zimmermann, E., Morrone, M. C., Burr, D. C. Visual mislocalization during saccades
sequences. *Exp Brain Res*, 233(2):577-585. (2014).

34. Castet, E., Jeanjean, S., Montagnini, A., Laugier, D., and Masson, G. S. Dynamics of
attentional deployment during saccadic programming. *Journal of Vision*, 6(3), 196–212.
(2006)

35. Montagnini A, Castet E. Spatiotemporal dynamics of visual attention during saccade
preparation: Independence and coupling between attention and movement planning. *J*
Vis. 7(14):8.1-16. (2007).

36. Neupane, S., Guitton, D., and Pack, C. C. Two distinct types of remapping in primate
cortical area V4. *Nat*. *Commun*.7:10402. (2016).

37. Rolfs, M., Jonikaitis, D., Deubel, H., and Cavanagh, P. Predictive remapping of attention
across eye movements. *Nat*. *Neurosci*. 14, 252–256. (2011).

38. Szinte, M., Jonikaitis, D., Rangelov, D., Deubel, H. Pre-saccadic remapping relies on
dynamics of spatial attention. *eLife*, 7:e37598. (2018).

39. Harmony, T. The functional significance of delta oscillations in cognitive processing.
Frontiers in Integrative Neuroscience, 7, 83. (2013).

40. Ioannides AA, Kostopoulos GK, Liu L, Fenwick PB. MEG identifies dorsal medial brain
activations during sleep. *Neuroimage* 44:455–468. (2009).

41. Benedetto, A., Morrone, M.C. Visual sensitivity and bias oscillate phase-locked to
saccadic eye movements. *J*. *Vis*. 19, 15. (2019).

42. Benedetto, A., and Morrone, M. C. Saccadic suppression is embedded within extended
oscillatory modulation of sensitivity. *The Journal of Neuroscience*, 37 (13), 3661–3670.
(2017).

43. Hogendoorn, H. Voluntary saccadic eye movements ride the attentional rhythm. *Journal*
of Cognitive Neuroscience, 28 (10), 1625–1635. (2016).

44. Fries, P. A mechanism for cognitive dynamics: neuronal communication through
neuronal coherence. *Trends in Cognitive Sciences*, 9(10), 474–480. (2005).

45. Honda, H. Eye movements to a visual stimulus flashed before, during, or after a
saccade. *Attention and Performance,* 13: 567–582. (1990)

46. Honda H. The time courses of visual mislocalization and of extraretinal eye position
signals at the time of vertical saccades. *Vision Res.*, 31(11):1915-21. (1991).

47. Porter, T. C. Contributions to the study of flicker II. *Proceedings of the Royal Society of*
London Series A, 70, 313–329. (1902).

48. Humphries, D., Driver, P. Protean defence by prey animals. *Oecologia*, 5 285–302.
(1970).

49. Furuichi, N. Dynamics between a predator and a prey switching two kinds of escape
motions. J *Theor Biol*, 217(2): 159–66. (2002).

50. Botev, Z. I., Grotowski, J. F., Kroese, D. P. Kernel density estimation via diffusion.
Annals of Statistics, 38:2916–2957. (2010).

51. Crossland, M.D., Sims, M., Galbraith, R.F., Rubin, G.S. Evaluation of a new quantitative
technique to assess the number and extent of preferred retinal loci in macular disease.
Vision Research, 44(13), 1537-1546. (2004).

52. Ludwig, C. J. H., Gilchrist, I. D. Measuring saccade curvature: A curve-fitting approach.
Behavior Research Methods, Instruments, & Computers, 34, 618-624. (2002).

53. Nandy, A. S., Nassi, J. J., Reynolds, J. H. Laminar Organization of Attentional
Modulation in Macaque Visual Area V4. *Neuron*, 4;93(1), 235-246. (2017).

54. Tass, P., Rosenblum, M.G., Weule, J., Kurths, J., Pikovsky, A., Volkmann, J., Schnitzler,
A., Freund, H. J. Detection of n:m Phase Locking from Noisy Data: Application to
Magnetoencephalography. *Physical review letters*, 81, 3291-3294. (1998).

55. Bowman, A.W., Azzalini, A. *Applied Smoothing Techniques for Data Analysis: the*
Kernel Approach with S-Plus Illustrations (Oxford University Press, Oxford, 1997).

Fig 1. Behavioural protocol. (A) Markers in the illustration of visual space show all retinotopic locations at which a probe was flashed before and after the deployment of attention, saccade planning, and, finally, the execution of a saccade. Probes of interest, presented along the tangential axes, are conspicuously highlighted in open circle markers. **(B)** Temporal sequence of a valid trial across experiments with respect to target onset.

Fig 2. Behavioural data. (A-D) Probability density map of foveal gaze and saccade landing was calculated using a bivariate gaussian kernel along with their bivariate contour ellipse area estimates for each probability density map. Error estimates are standard error of means across observers. **(E-G)** Shifts in visual sensitivity along tangential axes with respect to target onset; error estimates were calculated using a 20-fold jackknife procedure. **(H)** Mean saccade trajectory deviation for horizontal and oblique saccades across experiments. Positive values on the ordinate denote trajectory deviation towards a clockwise location (cw), while negative values denote trajectory deviation towards a counterclockwise location (ccw).

Fig 3. Frequency representation of visual sensitivity and cross-location phase coherence. (A-F) Representation of raw detrended visual sensitivity at less and more eccentric locations in visual space. Shuffled (n=1000, dashed line) versus unshuffled (solid line) data set were used to compute location-specific spectral signatures at a p<0.05. **(G-H)** For parafoveal and peripheral clockwise and counterclockwise locations a cross-location phase coherency analysis was performed.

Fig 4. Model architecture and neurobiological force field with simulated putative population sensitivity. (A) For illustration purposes, a single eccentricity dependent population receptive field (pRF) within a retinotopic field is highlighted in pink. A pRF is defined by its center, size, and an eccentric dependent population elastic field (pEF). **(B)** $M_{G\alpha}$ denotes gaze transitional signals modelled as a virtual mass at infinity located in the direction of an upper tangential location approximating the initial shift in gaze. M_F denotes compressive signals modelled as a varying mass located at a lower tangential location in the fovea. **(C)** Simulation results in the foveal region when a pRF is under the influence of a resultant force exerted by (i) M_F and $M_{G\alpha}$ (curved space, thick marker) (ii) M_C (convergent signals modelled as a varying mass at a stationary peripheral site; see Methods) and (iii) M_T (saccade transitional signals modelled as a virtual mass at infinity located in the direction of a straight saccade; see Methods). **(D)** A pRF bounded by pEF is highlighted in pink for parafoveal simulations. **(E)** $M_{G\beta}$ denotes gaze translational signals modelled as a virtual mass at infinity located in the direction of a downward shifts in gaze, M_M denotes a varying mass at a mislocalized peripheral target, and M_S denotes the curved transitional signals modelled as a virtual mass at infinity located in the direction of a curvilinear saccade. **(F)** Curved, convergent and translational simulation results.