Title: Rapid brain development and reduced neuromodulator titers correlate with host shifts

in *Rhagoletis pomonella*

Authors: Hinal Kharva[1,2], Jeffrey L. Feder[3], Daniel A Hahn[4], Shannon B. Olsson[1]

Affiliations:

[1]Naturalist-Inspired Chemical Ecology lab, National Centre for Biological Sciences, Tata Institute of
Fundamental Research, GKVK Campus, Bellary Road, Bangalore 560065, India.

[2]School of Life Sciences, The University of Trans-Disciplinary Health Sciences and Technology, 74/2,
Jarakabande Kaval, Post Attur via Yelahanka, Bangalore 560064, India.

[3] Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana 46556, USA.

[4]Department of Entomology and Nematology, University of Florida, Gainesville, Florida 32611, USA.

Correspondence to: hinal@nice.ncbs.res.in, shannon@nice.ncbs.res.in

Keywords: host choice, neurotransmitters, diapause, biogenic amines, sensory systems

Abstract:

Host shifts are considered a key generator of insect biodiversity. For insects, adaptation to
new host plants often requires changes in larval/pupal development and behavioural
preference towards new hosts. Neurochemicals play key roles in both development and
behaviour, and therefore provide a potential source for such synchronization. Here, we
correlated life history timing, brain development, and corresponding levels of 14
neurochemicals in *Rhagoletis pomonella* (Diptera: Tephritidae), a species undergoing
ecological speciation through an ongoing host shift from hawthorn to apple fruit. These races
exhibit differences in pupal diapause timing as well as adult behavioural preference with
respect to their hosts. This difference in behavioral preference is coupled with differences in
neurophysiological response to host volatiles. We found that apple race pupae exhibited
adult brain morphogenesis three weeks faster after an identical simulated winter than the
hawthorn race, which correlated with significantly lower titers of several neurochemicals. In
some cases, particularly biogenic amines, differences in titers were reflected in the mature
adult stage, when host preference is exhibited. In summary, life history timing,
neurochemical titre, and brain development can be coupled in this speciating system,
providing new hypotheses for the origins of new species through host shifts.

Introduction:

Adaptation to environmental and ecological factors has been shown to play an important role in population divergence and speciation in a large number of systems (1–3). For phytophagous insects shifting to new host plants, populations potentially need to adapt their growth and development to the new host, and at the same time modulate their behavioural preference to locate that host. Furthermore, populations on novel hosts must regulate their life-history timing to coincide with the new host phenology (4–11). How these multiple events are synchronized between ancestral and novel hosts with vastly different phenologies and characteristics is an area of intense study for understanding the genesis of insect biodiversity (7,12).

Rhagoletis pomonella (Diptera: Tephrtidae) provides a unique opportunity to identify associations between pupal development, life history timing, and adult host choice. This species of flies originally infested fruits of the native downy hawthorn (*Crataegus mollis*) until apples (*Malus pumila*) were introduced in Eastern USA around 180 years ago (12–14). This introduction of a new host facilitated a shift in host preference from their native host, downy hawthorn, to domesticated apples, eventually leading to two host races specific to each fruit (12,13). This host shift from hawthorn to apple fruit was facilitated by two factors. First, apple and hawthorn flies are univoltine and the adults are short lived, thus each host race must emerge synchronized with the fruiting time of their host plant. Due to the earlier fruiting time of apples, apple flies initiate their overwintering dormancy, termed diapause, earlier than hawthorn flies, and emerge as adults about one month earlier as well (13). Previous work shows that the earlier seasonal adult emergence of apple host-race flies is driven solely by the timing of the termination of pupal diapause (15). The difference in adult emergence timing that drives synchronization with host fruits results in a degree of mating isolation between the apple and hawthorn races (16). Second, adults of the two host races also exhibit distinct preferences for the volatiles of their respective host fruits, which serves as an important reproductive barrier because the flies mate directly on or near the ripe host fruit (17–19). Previous studies forcing hybridisation between the host races in the laboratory showed that F1 hybrids of these two populations have altered peripheral olfactory physiology suggesting developmental abnormalities regarding their response to the host volatiles.(20). Recent studies have shown that the change in host fruit olfactory preference between the two host races does not occur at the chemoreception stage in the antenna (21–23), but rather at the first synapse of the olfactory system in the brain, the antennal lobe (24). The coupled difference in pupal diapause timing, adult preference behaviour, and adult brain physiology in *R. pomonella* provides a unique opportunity to examine if there are corresponding differences in brain development between the races that could simultaneously impact both developmental rate and adult olfactory host choice.

Species from temperate regions like *R. pomonella* frequently use the timing of diapause to
avoid the stresses of winter and also to synchronize themselves with the phenology of their
hosts the next growing season (25). While diapause is often conceived of as a state of
developmental arrest, it is in fact a dynamic, physiologically regulated process with defined
phases of development including the diapause preparatory stage, diapause induction,
diapause maintenance, and the resumption of rapid development at the end of diapause
(26,27). In insects, the central nervous system (CNS) and associated endocrine glands
produce neurochemicals like neurotransmitters, neurohormones, and neuropeptides that
regulate diverse physiological events including the induction and termination of diapause
(28). Diapause can be regulated via changes in hormone/neurotransmitter titres, receptor
abundance, or regulation of specific neurochemical pathways across the stages of diapause,
such as has been found with the dopamine and serotonin pathways (29–31). Apart from life
history timing, many of these same neurochemicals also play important roles in insect
behaviour. Biogenic amines like dopamine, octopamine, and serotonin are known to have
profound impacts on adult insect behaviour across many taxa (32–35).
The dual role of neuromodulators in regulating both insect development and host-seeking
behavior suggests that changes in neuromodulator titres or production at specific life stages
could impact both life history timing and preference of a phytophagous insect for its host
plant through changes in brain development or differentiation. Further, the aforementioned
changes in neurophysiological response and preference for host volatiles between *R.*
pomonella races might be rooted in changes in brain development, neuromodulation, or
both. In this study, we use a variety of chemoanalytical, morphological, and
immunohistochemical techniques to examine pupal-adult brain development, life history
timing, and corresponding neurotransmitter levels in two closely related populations of *R.*
pomonella that differ both in diapause timing and adult preference for their respective hosts.
There are a number of potential neurochemical candidates and life history stages that could
play a role in this host shift, and also a general lack of knowledge regarding how
neuromodulators could impact this process. Therefore, our goal is to track brain
development from larval to adult stages in the apple and hawthorn host races of *R.*
pomonella, and identify which stages, if any, exhibit differences in neural development or
neuromodulation between the host races, and how these differences are reflected in the
mature adult fly, when host preference is exhibited.

(A) Life cycle of Rhagoletis pomonella

Figure 1: The life cycle of *Rhagoletis pomonella.* (A) Life stages of the fly from mating to oviposition from right to left. (B) Micrographic images of pupal, pharate-adult, and adult developmental stages from left to right. The upper stereomicrographs in the bottom panels show the head morphology while the lower confocal micrographs indicate the corresponding brain morphology of those stages using immunohistochemical nc82 staining across the different developmental stages (stages 1-8). The developmental stages are classified based on distinct head morphology, metabolic rate, or CNS development. Our imaging data showed that there are no morphological differences between stages 1 and 2 or between stages 7 and 8. The stages are as follows: **1**, Pre-winter diapause induction stage pupae still have high metabolic rates (Methods & Figure s1, s2 and Movie 1, apple race, stage 1); **2**, pre-winter diapause induction stage when pupae have entered metabolic depression (Figure s1, s2 Figure 1 and Movie 2, hawthorn race, stage 2); **3**, post-winter diapause maintenance phase (Figure s2 and Movie 3, apple race, stage 3); **4**, end of diapause (Figure s2. Movie 4, hawthorn race, stage 4); **5**, midway through pharate-adult development (Figure s2, Movie 5, apple race, stage 5); **6**, late pharate-adult development ((Figure s2, Movie 6, apple race, stage 6); **7**, sexually immature adult fly, less than 7 days old ((Figure s2, Movie 7, hawthorn race, stage 7); **8**, sexually mature fly, more than 12 days old;

Methods:

Insect collection and maintenance

Apple and hawthorn fruits naturally infested with larvae were collected from four different sites in Michigan, USA (Grant, Fennville, Cassopolis, Lansing) in August and September 2016, and flies were reared from larvae to adulthood following previously established

Rhagoletis husbandry methods (14). In May to August 2017, after leaving the pupae at room temperature for 15 days, they were shipped to India (with permit). This set of pupae were used to study brain development and quantification of neurotransmitters from adult flies. Eclosed adults were maintained on a diet of sugar and yeast on a 14L:10D light cycle at 25° C and 65% humidity. Post-eclosion, young flies 1-6 days old were classified as sexually immature whereas flies that were 12-14 days old were classified as sexually mature (36–38). To study the pre-winter and post-winter brain development as well as quantify neurotransmitters, a second set of pupae were collected in summer 2018. After collecting infested fruits from the above field sites, fruit were transferred to a tray with a wire mesh attached and kept in an insect-rearing room at $25\pm1°C$, 14L:10D light cycle. Every day newly emerged pupae were collected and transferred to petri dishes with damp vermiculite, and maintained within a chamber containing a saturated potassium chloride solution to maintain ~85% relative humidity. To differentiate diapausing and non-diapausing pupae during the diapause initiation stage, four different cohorts of pupae were set aside and subjected to metabolic rate measurements once they reached either 7 or 19 days after pupariation. Other cohorts of pupae at 10 days after pupariation were transferred to a dark refrigerator at $4°C$ with saturated KCL solution to stimulate over-wintering diapause for six months and study post winter development until they were hand-carried to India in November 2018.

Chemicals and Reagents

16% Paraformaldehyde EM grade (15710) was obtained from Electron Microscopy Sciences, Hatfield, PA, USA. Triton x and Bovine serum albumin were purchased from Sigma-Aldrich (Bangalore, India). All standards, ammonium acetate, Acetone, hydrochloric acid (HCl), boric acid, and reagents required for 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate (AQC) synthesis, were obtained from Sigma-Aldrich (Bangalore, India). Deuterated internal standards 14 were supplied by CDN isotope (Quebec, Canada). Ascorbic acid was obtained from Himedia (Bangalore, India), and Formic acid (FA) was obtained from Fisher Scientific (Bangalore, India). Reverse-phase solid phase extraction (RP-SPE) cartridges (Strata-X, 8B-S100-TAK) were obtained from Phenomenex, Inc. (Hyderabad, India). High-purity MS grade solvents (methanol, acetonitrile, and water) were obtained from Merck Millipore (Merck Millipore India Pvt. Ltd., Bangalore).

Pre-diapause metabolic rate measurement

Even though pupal diapause is ecologically obligate in *R. pomonella*, a small number of pupae avert diapause under laboratory conditions and directly develop into pharate adults in the prewinter period (11). To eliminate these non-diapausing individuals from our sampling before overwintering, we used a protocol adapted from Ragland *et* al 2009 & Powell *et* al 2021 to phenotype pupae as diapausing or non-diapausing by measuring g metabolic rates in the pre-winter period.

First we collected 7-day and 19-day-old pupae to measure their weight on an analytical

balance with 5 µg precision (Mettler XP6, Toledo, OH, USA). Pupae were transferred to a 5

ml syringe used as a respirometry chamber for checking metabolic rate as an indicator of

diapause or non-diapause status, held for 24h so adequate CO_2 could build up in the

chamber, and CO_2 was measured on the 8th and 20th day respectively. Syringes were

sealed with the plunger drawn back to produce a chamber of 1 ml internal volume. *R.*

pomonella pupae were small enough to fit into one arm of the luer valve, allowing the full

volume of the syringe to be injected. We also purged multiple empty syringes to serve as

controls/blanks. Every 24 h, the full volume of each syringe was injected into a flow-through

respirometry system consisting of a Li-Cor 7000 infrared CO_2 analyser (Lincoln, NE, USA)

with a resolution of 5 parts per million (ppm) CO_2 interfaced to a Sable Systems International

UI-2, recorded by Expedata data logging software (Las Vegas, NV, USA). The flow rate was

fixed at 150 ml/min using a Sierra Instruments mass flow controller (Monterey, CA, USA).

CO_2-free air, scrubbed with a dririte-ascarite-dririte column, served as the baseline for

measurements, and the system was routinely calibrated with CO_2-free air and a certified

standard mixture of 500 ppm CO_2 in nitrogen (Airgas, Jacksonville, FL, USA). For one

replicate a total of 30-40 pupae were used to calculate the metabolic rate and after that, the

brain samples were dissected on cold PBS(1x) and flash frozen with liquid nitrogen. For

further details, see Supplementary Materials, Extended Methods.

Brain morphology and immunohistochemistry

In early November 2018, diapausing pupae were hand-carried with ice packs from

Gainesville, FL, USA to Bangalore, India. After keeping pupae 6-months in the refrigerator

(October 2018 - March 2019), in March 2019 all four different cohorts of pupae of the apple

race were pulled out at their respective 6-month time points and left at room temperature

25°C, 14L:10D light cycle, and 65% humidity as described in Insect collection and

maintenance above. Every 5 days after removal from artificial overwintering, a few pupal

caps (n=5-35) were removed to observe development externally and brains were also

dissected from each individual to assess brain development (Figs. 1-2). Further, pupae were

individually photographed using an infinity HD camera (lumenera, model number N9033210)

attached to a stereomicroscope. After that, the brain was individually dissected using 50ul of

1X phosphate buffered saline solution, PBS (PH: 7.4) on ice with fine forceps, and stored at -

$80^{o}C$. Four different cohorts of pupae of the Hawthorn race were pulled out in April 2019 at

their respective 6 months of overwintering time points (November 2018 - April 2019) and

were photographed and then dissected using the above protocol.

To characterize brain morphology, brain samples were dissected as mentioned above and

then underwent immunohistochemical staining using a protocol adapted from those used in

Drosophila and Rhagoletis (24,39,40). Further details can be found in Supplementary

Materials, Extended Methods.

Quantification of neurotransmitters from the R. *pomonella* brain

After morphologically determining developmental stage, brain samples were removed from -

80^{o}C storage and five brains of the same stage were pooled for each sample replicate. The

sample processing was performed as in Ramesh *et* al, 2019 (41–43). After the brain

samples were pulled out, samples were immediately transferred to a vial containing 190 µl of

Acetone (with 0.1% Formic acid, FA) and 10 µl of 1% of ascorbic acid (1.76mg/ml), followed

by derivatization with 6-aminoquinolyl –N-hydroxysuccinimidyl carbamate (AQC) as in

Ramesh et al 2019. (41–43). Adult fly brains were dissected individually, pooled into a group

of five, and stored at -80^{o}C with190 µl of Acetone and 10 µl of 1% of ascorbic acid. After that

10 µl of internal standard (ISTD), a mixture of all 14 neurotransmitters, (0.5ug/ml i.e. 1ng on

column) was added to brain samples in screw cap vials. This mixture was sonicated for 1

min, and then homogenised using a plastic pestle. It was then immediately centrifuged

(13500 rpm 4^{o}C, 5 minutes), and the supernatant was transferred to a new tube.

Simultaneously, the vials for standard solutions for calibration curves were also prepared

(supplementary table s11). Serial dilutions of standard stocks were prepared with highest

concentration considered on column 100% to five points including 50%,25%,12.5% and

6,25% of the maximum quantity for each of the targeted compounds, for pre-winter and post-

winter brain samples. For sexually immature and sexually mature adults the standard stock

was diluted to prepare 200%, 160%, 80%, 40%, 20% of the targeted compounds. After that

10 µl internal standards with 190 µl acetone (0.1% FA) and 10 µl of ascorbic acid were

added to the standard solution tubes. All samples and standard tubes were dried in a speed-

vac for 1hr. Apart from these, one more vial containing 16 amino acids (1 µl of 10 µg/ml) was

added as an additional standard. These 16 amino acid standards helped to validate the

method and retention time used for mass spectrometry (MS) every time we ran a sample.

Once everything was dried in the speed-vac, 80 µl of borate buffer with 10 µl of ascorbic acid

(1.76 mg/ml) was added to all tubes and vortexed. Before analysis, 10 µl of 10 mg/ml AQC

(prepared in 100% Acetonitrile, ACN) was added and kept for 10 min at 55^{o}C. After that 3ul

of 100% formic acid FA was added and the tube was vortexed to stop the reaction. Tubes

were subsequently kept at room temperature until all SPE columns were rinsed and cleaned

with 100% methanol and 0.1% FA. After that, 500 µl of water was added to all the tubes,

then vortexed and the solution was loaded on SPE columns. Columns were washed twice

with 0.1% formic acid prepared with LC grade water. After that 1ml of ACN: MeOH 4:1 in

0.1% formic acid was added to the column and eluted in a new vial. All tubes were dried in a

speed vac for 3hrs. Dried samples were stored at -20^{o}C until they were run on LC-MS. Each

sample was thawed and reconstituted in 50 µl of 2% ACN prepared in 0.5% FA. The LC-MS instrument method and setup are described in detail in tables s9, s10, s11, s12 figure s6 and s7. The final quantification and analyses can be found in detail in Supplementary Materials, Extended Methods.

Statistical Analysis:

1) To compare the relative differences in timing of the proportion of apple vs. haw flies transitioning from pupal diapausing brain morphology to adult brain development post winter, we used a generalised linear model with a binomial, log-link function in R (V4.0.2) with ggplot, dplyr, and tidyr packed. In this logistic GLM host, days after winter was used to explain the proportion of individuals transitioning from stage 3 to stage 4 of brain development between apple and hawthorn flies, as well as to estimate 95% CIs around the logistic estimates. 2) Respiration and neuromodulator data were analysed using linear mixed effects models in R (V4.0.2) using the lmer function from the lme4 package. In the models, age and host race were used as a fixed effect whereas cohort / batch number was used as a random effect. Further, Tukey's HSD tests with correction for multiple comparisons were performed with pairwise multiple comparisons using the lmerTest package. Satterthwaite approximations on the lmerTest package were used to test the significance of the effects. Linear Discriminant Analysis (LDA) was performed on the same transformed data using the "MASS" package in R, with "leave-one-out cross-validation" to estimate the classification accuracy of the models. 3) The interaction between host race and days on amount of precursor neurochemicals at brain development stage 4 was done using, generalised linear model, Univariate analysis of variance in SPSS V.26. In the model host race and days were used as fixed factors and quantities of each neurochemical was used as the dependent variable. Further, Tukey's LSD tests were performed for pairwise comparisons with significance cut offs of 0.05.

Results:

***Rhagoletis pomonella* host races exhibit distinct brain development stages during pupation**

We observed external morphological and brain development in *Rhagoletis pomonella* pupae during diapause initiation, pre-winter diapause, post-winter diapause, diapause termination, pharate-adult, and eclosed adult developmental phases (Figure 1). The entire set of experiments were performed over a period of two seasons 2017 to 2019 with total of 2,091 individuals examined across both the host races. These phases were divided into 8 different stages according to changes in external head morphology, brain development, or metabolic rate (Figure 1 and s1). The pre-winter diapause initiation stage (stage 1, 8 days after puparium formation, n=130 total dissections, Methods & Figure s1, s2 and Movie 1, staining,

n=15, apple n=7,hawthorn n=8), and the pre-winter diapause maintenance stage (stage 2,
20 days after puparium formation, n=150, Figure s1, s2 Figure 1 and Movie 2, staining, n=6,
apple n=7,hawthorn n=8) were differentiated using respiration rates wherein pupae entering
diapause exhibited higher metabolic rates 8 days after pupariation that stabilized at low
levels of metabolic depression by 20 days after pupariation, when pupae were clearly in the
diapause maintenance phase [Figure s1,(15,44)]. During these two pre-winter stages, brain
morphology reflected that of the larval brain with a clearly identifiable ring gland and sub-
oesophageal ganglion (Figure s2, A and Movie 1, 2). After a six-month artificial overwintering
period at 4-5^{o}C, we observed substantial neural differentiation and remodelling had occurred
compared to pre-winter diapausing brains (stage 3, n=297, Figure s2, B and Movie 3,
staining, n=5, apple n=2, hawthorn n=3) in that the ring gland became disassociated and the
place between the hemispheres containing the dorsal vessel became thin (c.f. stages 1 & 2;
Figure s2, A, B & movie 2, 3).
Subsequent assessment of later stages also showed that adult brain (CNS) development
and differentiation occurred after overwintering, at a stage during pharate-adult development
where early antennal development and changes in head morphology were also observable
(stage 4, n=125, Figure s2, C Movie 4, staining, n=5, apple n=4, hawthorn n=1). At this stage
(4), brain region boundaries were apparent, but the regions themselves remained undefined
(Figure s2, C & movie 4). By stage 5, orange pigmentation had accumulated in the eyes and
a transparent antenna became well developed. At this stage, the central nervous system
was more defined with structures including the antennal lobe, mushroom body, and
suboesophageal ganglion (n=70, Figure s2, Movie 5, staining, n=3, apple n=1, hawthorn
n=2). The last three stages (including the sexually immature and mature eclosed adult
stages) were similar in both brain and external head morphology, just with greater
progression of bristle development and pigmentation of the pharate-adult cuticle (Stage 6,
n=60, Movie 6, staining, n=2, apple n=1, hawthorn n=1; stage 7, n=85, Movie 7, staining,
n=1, hawthorn n=1 and stage 8 n=85, Figure s2, D).
296
297   Apple race pupae exhibit more rapid onset of adult brain morphogenesis than the
298   hawthorn race
To compare post-winter brain development between the two races of *R. pomonella*, both
apple and hawthorn pupae were brought to room temperature (25^{o}C) after 6 months of
simulated overwintering (4-5^{o}C) synchronized to their respective diapause initiation timing
(Figure 2 A, B and Tables s1 and s2). By sampling a subset of pupae every 5 days from day
0 to day 70 after removal from artificial overwintering conditions, we observed that both host
races exhibited onset of adult neurogenesis during the post-winter phases, as shown in
Figure 1 and previously implied by Dowle et al. (2020). In the apple race, a substantial

number of pupae exhibited adult brain morphology (Stage 4-6 in Figure 1) starting from day 0 after removal from simulated winter and until day 25 when 42% of the pupae sampled exhibited adult brain morphology (57/135). Conversely, in the hawthorn race only 4/152 (3%) total pupae exhibited

adult brain morphology even until day 40 (Figure 2 A, B and Tables s1, s2). Logistic regression analysis (Figure 2 C) showed that apple race individuals began initiating adult brain development significantly more rapid than hawthorn race individuals (x^2day = 265.79, p<0.001; x^2host = 88.04, p<0.001; x^2day*host = 34.2, p<0.001). Diapausing apple race pupae began adult brain development approximately 24 days faster than diapausing hawthorn race pupae, with 18.34 ($\pm$0.56-0.42,95%CI) days for 50% pupae exhibiting adult brains in apple vs. 42.10 ($\pm$0.61-0.42,95%CI) days for hawthorn (Figure 2). Therefore, even though the artificial overwintering period was of the same duration for both the races, the transition from pupal to adult brain development began occurring significantly more rapid in the apple race vs. the hawthorn race. In other words, not only has the apple race shifted its overall life cycle to coincide with host phenology, but the rate of development of the pupal brain from the post winter to the initiation of adult brain morphogenesis has become more rapid.

Figure 2: Post-winter brain development over time between the two host races. Brains were dissected from pupae and assessed every 5 days after simulated overwintering. A) Apple race, n=138

327   larval brain morphology (Stages 3 in Figure 1) vs. n=142 adult brain morphology (Stage 4-6 in Figure
328   1); B) Hawthorn race, n=201 larval brain morphology vs. n=157 adult brain morphology). C) Logistic
329   regression curves with 95% confidence intervals for the proportion of adult brain morphology vs. larval
330   brain morphology observed in *R. pomonella* apple and hawthorn race pupae over time after artificial
331   overwintering.
332

Neuromodulator levels in developing Hawthorn race pupal brains are generally higher than in the apple race.

We next examined a total of 14 neurochemicals across 6 biochemical pathways in the
developing brains of both hawthorn and apple race of *R. pomonella.* These neurochemicals
included both precursor molecules and their products (Figures, 3, s3, s4 & s5). We used
linear mixed models to assess both within and between host race comparisons. All statistical
results have been corrected for multiple comparisons using Turkey's HSD. In general, the
titres of precursor chemicals increased first in the post-diapause stages as pharate-adult
brain development began (transition from stage 3 to 4), whereas product molecule levels did
not change until later stages, even as late as post-eclosion adult fly sexual maturation. Of
the 14 neurochemicals examined, 13 showed significant differences in titre between the host
races at one or more developmental stage. Of these, only three chemicals showed higher
levels in the apple race, (dopamine, stage 5, p= 0.0001, serotonin, stage 6, p= 0.0001, and
serine, stage 6, p= 0.009). For all other molecules, the hawthorn race exhibited higher levels
than the apple race. These chemicals further increased in titre as *R. pomonella* development
progressed, until the point of adult fly sexual maturity, when host preference first emerges.
At the stage of fly sexual maturity (Stage 8, Fig. 1), product neurochemicals from two major
pathways showed a difference between the host races involving histidine to histamine and
tyramine to octopamine, respectively (stage 7, apple race n=10, total of 50 brains; hawthorn
race n=10, total of 50 brains; histidine, p = 0.005; stage 8, apple race n=10, total of 50
brains; hawthorn race n=10, total of 50 brains; tyramine, p= 0.011; octopamine, p= 0.011;
histamine, p= 0.020). This indicates that the differences in neurochemical titre witnessed
during pupal development persist even into fly sexual maturity, particularly the octopamine
and histamine pathways.

Figure 3: Quantification of biogenic amines and their precursors from the onset of adult brain development (stage 4) all the way up to sexually mature fly (stage 8) as defined in Figure 1 (A-N). Line graph of neurochemical titres for both host races at different developmental stages with 4-15 samples per stage, containing a pool of five brains in each sample, symbols represent mean with SE: A) tyrosine; (B) DOPA; (C) dopamine; (D) tyramine; (E) octopamine; (F) histidine; (G) histamine; (H) glutamate; (I) GABA; (J) aspartate; (K) serine; (L) tryptophan; (M) tryptamine; and (N) serotonin; Asterisks above indicate differences between host races at the equivalent stage of brain development. P-values represented are < 0.05 *, < 0.01 **, and < 0.001 ***, linear mixed effect model, followed by Tukey's HSD correction for multiple comparisons.

Rapid onset of adult brain morphogenesis in the apple race corresponds to lower
levels of neuromodulators

Figure 4: Quantification of precursor neurochemicals during onset of adult brain morphogenesis. (A-
F) Line graph of neurochemical titres for both host races at different developmental stage 4 with 1-4
samples per time point, containing a pool of five brains in each sample, symbols represent mean with
SE: A) tyrosine; B) tyramine; C) DOPA; D) aspartate; E) glutamate; F) tryptophan; Asterisks above
indicate significant interactions between effects of host race*days for neurotransmitter titre. P values
represented are < 0.05 *, < 0.01 **, and < 0.001 ***, analysis using Two-way ANOVA, post hoc
Tukey's LSD test.

Given that in the apple race, the rate of adult brain development is significantly more rapid
than the hawthorn race (Figure 2), and also exhibits lower titres of several neuromodulators,
we next assessed whether the rate of development corresponded to lower levels of
neuromodulators, particularly at the first stage of adult brain development (Stage 4). Figures
3, s3, s4 and s5 show that six precursor molecules including tyrosine, tyramine, DOPA,
aspartate, glutamate, and tryptophan from four different biosynthetic pathways showed

significant differences between the host races at the first appearance of the adult brain
during pupal-pharate adult metamorphic development, stage 4 in Figure 1 (stage 4, apple
race n=14, total of 70 brains; hawthorn race n=11, total of 55 brains; tyrosine, p= 0001;
DOPA, p= 0.0004; tyramine, p= 0.0001; aspartate, p= 0.022; glutamate, p= 0.002;
tryptophan, p= 0.002;). For each chemical, the titres were higher in the hawthorn race
compared to the apple race. To compare how these precursor neurochemicals titres
changed over time between the developing hawthorn and apple race pupae, we compared
neurochemical titres of Stage 4 brains against the day they were sampled after winter.

A Univariate analysis of variance, UNIANOVA was conducted to examine the effects of host
race and days after winter on the titres of these six precursor neurochemicals. Except for the
neurochemical DOPA, there was a significant interaction between host race and day for
each neurochemical titre (table s7, tyrosine, $F_{(12,10)}$ =10.9, P = 0.0001; tyramine, $F_{(12,10)}$ = 4.95, p = 0.008; DOPA, $F_{(12,7)}$ = 0.82, p = 0.632; aspartate, $F_{(12,11)}$ = 10.1, p=
10) = 4.95, p = 0.008; DOPA, F (12, 7) = 0.82, p = 0.632; aspartate, F (12, 11) = 10.1, p=
0.0001; glutamate, $F_{(12,11)}$ = 6.47, p = 0.002; tryptophan, $F_{(12,10)}$ = 4.35, p = 0.013).
Therefore, neurochemical titres were significantly lower in rapidly developing apple race
pupae than at later time points, when hawthorn race pupae were beginning to develop adult
brain morphology. This was true for precursor chemicals even when accounting for pupae
that might have been impacted by the temperature effects of development during winter itself
(i.e. until day 20) (table s8, tyrosine, $F_{(8,9)}$ =38.14, P = 0.0001; tyramine, $F_{(7,10)}$ = 4.46, p
= 0.017; DOPA, $F_{(7,8)}$ = 0.84, p = 0.581; aspartate, $F_{(8,10)}$ = 14.25, p= 0.0001;
glutamate, $F_{(8,9)}$ = 25.36, p = 0.0001; tryptophan, $F_{(7,10)}$ = 7.19, p = 0.003). These
differences in titres between the races then carried through further developmental stages
particularly during the sexual maturation of the adult fly in some cases (note stages 7 and 8
in Figures 3, s3, s4 and s5). As a result, more rapidly developing brains in apple race pupae
exhibited significantly lower titres of neuromodulators than later developing hawthorn race
brains, especially biogenic amines, and these differences were also reflected in the adult fly
stage at which host preference is exhibited (36–38).

Discussion:
In this study, we have identified differences in rates of adult brain morphogenesis and levels
of several neurochemicals between the apple and hawthorn host races of *Rhagoletis*
pomonella across the transition from pupal diapause to post-diapause, pharate-adult
development, and adult fly host preference stages. These two closely related populations
differ in both life history timing and adult host preference behaviours, and are generally
considered a model for incipient ecological speciation. The divergence of these two
populations towards hosts with such different phenologies serves as an excellent system to

examine how multiple life history events are synchronized between ancestral and novel
hosts.

First, we found that while some neurogenesis occurred during the diapause maintenance
phase in both host races, substantial adult brain development and differentiation was
initiated only after winter and brain development progressed through several morphological
stages to adult emergence. When we dissected brains from apple and hawthorn race pupae
at regular intervals after they were removed from overwintering, almost no hawthorn pupae
showed brain development beyond stage 3 for at least a month after overwintering, and it
took up to 50 days for the majority of hawthorn pupae to terminate diapause and transition
from stage 3 to stage 4 where pharate adult neural development was apparent (Figure 2B).
In contrast, some apple race pupae were already progressing to stage 4 pharate adult
development as soon as they were removed from overwintering (Figure 2B). This finding
indicates not only has apple race shifted its entire life cycle to coincide with fruit phenology
but its brain developmental rate has become more rapid. This result is also in agreement
with recent transcriptomic work showing rapid up-regulation of growth and development-
related transcript genes in the apple race with shorter post-winter diapause duration as
compared to the longer post-winter diapause duration of hawthorn race (45).
To better understand how neurochemical signaling could be associated with the different
stages of development in *R. pomonella*, we examined six major biosynthetic pathways
known to impact brain development as well as behavior in insects (41,46). Our results
indicate that the titers of several neurochemicals were significantly reduced in the apple race
pupae across multiple developmental stages as compared to the hawthorn race, (Figures 3,
4, s3, s4 and s5). Further analysis showed that apple race adult brains contained lower titres
of several neurochemicals at the onset of adult brain development, particularly those in the
dopamine and octopamine pathways, than later developing hawthorn race adult brains, and
this difference is also apparent in adulthood, when host preference is exhibited. Given the
importance of these neurochemicals to both development and behaviour, we hypothesize
that this difference in neurochemical production between the host races could connect
diapause termination timing and subsequent host preference in the adult flies, a key
characteristic for host-associated divergence and ecological speciation in the *R. pomonella*
system.
In agreement with our hypothesis, a recent study comparing olfactory neurophysiology
between the apple and hawthorn races of *R. pomonella* identified a neuronal switch in the
chemosensory system in the adult brain associated with differential host choice behaviour
towards apple or hawthorn fruit (24). Such a switch in neurophysiology has similarly been
observed between the Z and E strains of the European Corn Borer, *Ostrinia nubilalis*, where

male preference for a particular isomer of the sex pheromone is controlled by cis-acting variation in a sex-linked transcription factor (bric à brac; bab) expressed in the developing male antenna (47). A recent study of the *Rhagoletis* brain transcriptome also shows variation in cis-regulatory elements associated with differentially expressed transcripts during diapause development and an important role of hub genes in transcriptional networks that differ during diapause development between the two host races (45). In our current study, we show that the onset of adult brain differentiation (Stage 4) also corresponds with morphogenesis and emergence of the adult antenna, and this stage is accompanied by several significant differences in neuromodulator levels between the host races, specifically members of the dopamine and octopamine pathways (Figures 3 and 4). Interestingly, in *Drosophila* cis-regulatory variation in two genes involved in these pathways, tyrosine hydroxylase and dopa decarboxylase (Figure 3A), have been shown to impact neurogenesis (46). This provides a new hypothesis that differences in titres of these neurochemicals expressed during the transition from pupal diapause to early pharate-adult brain development might impact, or be impacted by, the expression of developmental genes in *Rhagoletis* and ultimately lead to changes in host-fruit preference in reproductively mature adults. Future studies that measure the expression of enzymes involved in the production of dopamine and octopamine and selective pharmacological treatments that act as agonists and antagonists of these biogenic amines at early developmental stages in these two host races could indicate if these pathways are involved in the differentiation of these two host races.

To conclude, we have characterized the progression of neurogenesis from diapause onset to adult reproductive maturation in both the apple and hawthorn host races of *R. pomonella*, an important model for ecological speciation and diversification. We identified significantly lower neurochemical levels, particularly biogenic amines in the dopamine and octopamine pathways, in the apple race of *R. pomonella* that correspond to more rapid adult brain morphogenesis in this new host race. These differences in neurochemical titre between the races in the developing pupal stages are also apparent in the adult brains at stages when flies are exhibiting host preference. Genetic and behavioural analyses of hybrids between these two host races suggest that their olfactory divergence is based on differences at only a few loci (48). Furthermore, a recent study of a similar neurological change in pheromone perception in the European corn borer has implicated a cis-regularly element (47), several of which also influence biogenic amine pathways (46). Because biogenic amines have been implicated to impact both pupal diapause and adult behaviour, this study offers a new hypothesis to correlate life history timing and adult host preference through developmental differences in neuromodulation. This hypothesis must now be tested in further studies

assessing enzymatic expression and pharmacological manipulation of neuromodulator levels

in developing pupae. As previously suggested, connecting host preference and survival

through relatively simple changes could be a widespread mechanism for generating

biodiversity across phytophagous insects, contributing to the origin of the large number of

species observed (24).

Data accessibility: All data supporting this manuscript will be uploaded to Dryad upon

submission.

Authors' contribution: HK, SBO, and DAH conceived the study and designed experiments;

JLF provided the adult flies and reagents for the experiment. HK conducted experiments;

HK, DAH and SBO analyzed data; HK and SBO wrote the manuscript; all authors revised

and approved the manuscript.

Competing interests: The authors declare no competing interests.

Funding: This work was supported by NCBS-TIFR funding, Department of Atomic Energy,

Government of India, and a SERB Ramanujan Fellowship to SBO under project no. 12-R&D-

TFR-5.04-0800 and 12-R&D-TFR-5.04-0900; Infosys travel award to HK, as well as the US

National Science Foundation (DEB 1639005), the Florida Agricultural Experiment Station

(Hatch project FLA-ENY-005943), and the joint FAO/IAEA Coordinated Research Project on

Dormancy Management to Enable Mass-rearing to D.A.H.

Acknowledgements: We would like to thank Dr, Andrew Nguyen and Dr, Chao Chen for

helping with the analysis of metabolic rate. We would like to thank Dr, Cheyenne Tait for

sharing *the Rhagoletis* larvae image. We also thank Khushboo Patel for helping us with the

rearing of *Rhagoletis* and taking photographs of hawthorn pupal development. We thank the

NCBS CIFF facility for help with confocal imaging. We thank Dr. Divya Ramesh for helping

us with standardizing mass spectrometry methods and analysis.

References:

1.   Schluter D. Ecology and the origin of species. Trends Ecol Evol. 2001;16(7):372–80.

2.   Via S. Sympatric speciation in animals: the ugly duckling grows up. Trends Ecol Evol.
2001;16(7):381–90.

3.   Berlocher SH, Feder JL. Sympatric Speciation in Phytophagous Insects : Moving
Beyond Controversy ? Annu Rev Entomol. 2002;47:773–815.

4.   Craig TP, Itami JK, Abrahamson WG, Horner JD. Behavioral evidence for host-race
formation in Eurosta solidaginis. Evolution (N Y). 1993;47(6):1696–710.

5.   Feder JL, Hunt TA, Bush L. The effects of climate, host plant phenology and host
fidelity on the genetics of apple and hawthorn infesting races of Rhagoletis pomonella.
Entomol Exp Appl. 1993;69(2):117–35.

6.   Groman JD, Pellmyr O. Rapid evolution and specialization following host colonization

in a yucca moth. J Evol Biol. 2000;13(2):223–36.

7.   Itami JK, Craig TP, Horner JD. Factors Affecting Gene Flow between the Host Races
of Eurosta solidaginis. Genet Struct Local Adapt Nat Insect Popul. 1998;375–407.

8.   Pratt GF. Evolution of Euphilotes (Lepidoptera: Lycaenidae) by seasonal and host
shifts. Biol J Linn Soc. 1994;51(4):387–416.

9.   Smith DC. Heritable divergence of Rhagoletis pomonella host races by seasonal
asynchrony. Nature. 1988;336(6194):66–7.

10.   Wood TK, Keese MC. Host-Plant-Induced Assortative Mating in Enchenopa
Treehoppers. Evolution (N Y). 1990;44(3):619.

11.   Dambroski HR, Feder JL. Host plant and latitude-related diapause variation in
Rhagoletis pomonella: A test for multifaceted life history adaptation on different stages
of diapause development. J Evol Biol. 2007;20(6):2101–12.

12.   Bush GL. Sympatric Host Race Formation and Speciation in Frugivorous Flies of the
Genus Rhagoletis (Diptera, Tephritidae). Evolution (N Y). 1969;23(2):237.

13.   Feder JL, Chilcote CA, Bush GL. Genetic differentiation between sympatric host races
of the apple maggot fly Rhagoletis pomonella. Nature. 1988;336(6194):61–4.

14.   Walsh. B. The apple-worm and the apple maggot. Am J Hortic. 1867;338–43.

15.   Powell THQ, Nguyen A, Xia Q, Feder JL, Ragland GJ, Hahn DA. A rapidly evolved
shift in life-history timing during ecological speciation is driven by the transition
between developmental phases. J Evol Biol. 2020 Oct 1;33(10):1371–86.

16.   Feder JL, Filchak KE. It's about time: The evidence for host plant-mediated selection
in the apple maggot fly, Rhagoletis pomonella, and its implications for fitness trade-
offs in phytophagous insects. Entomol Exp Appl. 1999;91(1):211–25.

17.   Prokopy RJ, Bennett EW, Bush GL. Mating Behavior in Rhagoletis pomonellaa
(Diptera: Tephritidae) 1. Site of assembly. Can Entomol. 1971;103(10):1405–9.

18.   Linn CE, Dambroski HR, Feder JL, Berlocher SH, Nojima S, Roelofs WL. Postzygotic
isolating factor in sympatric speciation in Rhagoletis flies: Reduced response of
hybrids to parental host-fruit odors. Proc Natl Acad Sci U S A. 2004 Dec
21;101(51):17753–8.

19.   Linn C, Feder JL, Nojima S, Dambroski HR, Berlocher SH, Roelofs W. Fruit odor
discrimination and sympatric host race formation in Rhagoletis. Proc Natl Acad Sci U
S A. 2003 Sep 30;100(20):11490–3.

20.   Olsson SB, Linn CE, Michel A, Dambroski HR, Berlocher SH, Feder JL, et al.
Receptor expression and sympatric speciation: Unique olfactory receptor neuron
responses in F1 hybrid Rhagoletis populations. J Exp Biol. 2006;209(19):3729–41.

21.   Tait C, Batra S, Ramaswamy SS, Feder JL, Olsson SB. Sensory specificity and
speciation: A potential neuronal pathway for host fruit odour discrimination in
rhagoletis pomonella. Proc R Soc B Biol Sci. 2016;283(1845).

22.   Olsson SB, Linn CE, Roelofs WL. The chemosensory basis for behavioral divergence

involved in sympatric host shifts II: Olfactory receptor neuron sensitivity and temporal
firing pattern to individual key host volatiles. J Comp Physiol A Neuroethol Sensory,
Neural, Behav Physiol. 2006;192(3):289–300.

23.   Olsson SB, Linn CE, Roelofs WL. The chemosensory basis for behavioral divergence
involved in sympatric host shifts. I. Characterizing olfactory receptor neuron classes
responding to key host volatiles. J Comp Physiol A Neuroethol Sensory, Neural,
Behav Physiol. 2006;192(3):279–88.

24.   Tait C, Kharva H, Schubert M, Kritsch D, Sombke A, Rybak J, et al. A reversal in
sensory processing accompanies ongoing ecological divergence and speciation in
Rhagoletis pomonella. Proc R Soc B. 2021;288(1947).

25.   Denlinger DL, Hahn DA, Merlin C, Holzapfel CM, Bradshaw WE. Keeping time without
a spine: What can the insect clock teach us about seasonal adaptation? Philos Trans
R Soc B Biol Sci. 2017;372(1734).

26.   Kostal V. Eco-physiological phases of insect diapause. J Insect Physiol.
2006;52(2):113–27.

27.   Kostal V, Stetina T, Poupardin R, Korbelova J, Bruce AW. Conceptual framework of
the eco-physiological phases of insect diapause development justified by
transcriptomic profiling. Proc Natl Acad Sci U S A. 2017;114(32):8532–7.

28.   Denlinger DL. Regulation of Diapause. Annu Rev Entomol. 2002;47:93–122.

29.   Noguchi H, Hayakawa Y. Role of dopamine at the onset of pupal diapause in the
cabbage armyworm Mamestra brassicae. FEBS Lett. 1997;413(1):157–61.

30.   Wang Q, Hanatani I, Takeda M, Oishi K, Sakamoto K. D2-like dopamine receptors
mediate regulation of pupal diapause in Chinese oak silkmoth Antheraea pernyi.
Entomol Sci. 2015;18(2):193–8.

31.   Andreatta G, Kyriacou CP, Flatt T, Costa R. Aminergic Signaling Controls Ovarian
Dormancy in Drosophila. Sci Rep. 2018;8(1):1–14.

32.   Schulz DJ, Barron AB, Robinson GE. A role for octopamine in honey bee division of
labor. Brain Behav Evol. 2002;60(6):350–9.

33.   Schwaerzel M, Monastirioti M, Scholz H, Friggi-Grelin F, Birman S, Heisenberg M.
Dopamine and Octopamine Differentiate between Aversive and Appetitive Olfactory
Memories in Drosophila. J Neurosci. 2003;23(33):10495–502.

34.   Felix A, Kromann SH, Gramsbergen J, Hansson BS, Ignell R. Dopamine regulates the
mating switch in olfactory coding and preference in a female moth. 2008;1–9.

35.   Kromann A. Modulation of olfactory information in the antennal lobe of Spodoptera
littoralis. Modulation of olfactory information in the antennal lobe of Spodoptera
littoralis. 2012.

36.   Zhang A, Linn C, Wright S, Prokopy R, Reissig W, Roelofs W. Identification of a new
blend of apple volatiles attractive to the apple maggot, Rhagoletis pomonella. J Chem
Ecol. 1999;25(6):1221–32.

37.   Nojima S, Linn C, Morris B, Zhang A, Roelofs W. Identification of host fruit volatiles
from hawthorn (Crataegus spp.) attractive to hawthorn-origin Rhagoletis pomonella
flies. J Chem Ecol. 2003;29(2):321–36.

38.   Nojima S, Linn C, Roelofs W. Identification of host fruit volatiles from flowering
dogwood (Cornus florida) attractive to dogwood-origin Rhagoletis pomonella flies. J
Chem Ecol. 2003;29(10):2347–57.

39.   Ibba I, Angioy AM, Hansson BS, Dekker T. Macroglomeruli for fruit odors change
blend preference in Drosophila. Naturwissenschaften. 2010;97(12):1059–66.

40.   Singh AP, Das RN, Rao G, Aggarwal A, Diegelmann S, Evers JF, et al. Sensory
Neuron-Derived Eph Regulates Glomerular Arbors and Modulatory Function of a
Central Serotonergic Neuron. PLoS Genet. 2013;9(4).

41.   Ramesh D, Brockmann A. Mass Spectrometric Quantification of Arousal Associated
Neurochemical Changes in Single Honey Bee Brains and Brain Regions. ACS Chem
Neurosci. 2019;10(4):1950–9.

42.   Rangiah K, Palakodeti D. Comprehensive analysis of neurotransmitters from
regenerating planarian extract using an ultrahigh-performance liquid
chromatography/mass spectrometry/selected reaction monitoring method. Rapid
Commun Mass Spectrom. 2013;27(21):2439–52.

43.   Natarajan N, Ramakrishnan P, Lakshmanan V, Palakodeti D, Rangiah K. A
quantitative metabolomics peek into planarian regeneration. Analyst.
2015;140(10):3445–64.

44.   Ragland GJ, Fuller J, Feder JL, Hahn DA. Biphasic metabolic rate trajectory of pupal
diapause termination and post-diapause development in a tephritid fly. J Insect
Physiol. 2009;55(4):344–50.

45.   Dowle EJ, Powell THQ, Doellman MM, Meyers PJ, Calvert MB, Walden KKO, et al.
Genome-wide variation and transcriptional changes in diverse developmental
processes underlie the rapid evolution of seasonal adaptation. Proc Natl Acad Sci U S
A. 2020;117(38):23960–9.

46.   Monastirioti M. Biogenic amine systems in the fruit fly Drosophila melanogaster.
Microsc Res Tech. 1999;45(2):106–21.

47.   Unbehend M, Kozak GM, Koutroumpa F, Coates BS, Dekker T, Groot AT, et al. bric à
brac controls sex pheromone choice by male European corn borer moths. Nat
Commun. 2021;12(1):1–11.

48.   Dambroski HR, Linn C, Berlocher SH, Forbes AA, Roelofs W, Feder JL. The genetic
basis for fruit odor discrimination in Rhagoletis flies and its significance for sympatric
host shifts. Evolution (N Y). 2005;59(9):1953–64.