Dynamic Instability of Dendrite Tips Generates the Highly Branched Morphologies of Sensory Neurons

Sonal Shree[1]*, Sabyasachi Sutradhar[1]*, Olivier Trottier[1,2], Yuhai Tu[3], Xin Liang[4], and Jonathon Howard[1,2,5,#]

[1]Department of Molecular Biophysics & Biochemistry, Yale University, New Haven, CT 06511
[2]Department of Physics, Yale University, New Haven, CT 06511
[3]IBM T.J. Watson Research Center, Yorktown Heights, NY 10598
[4]Tsinghua-Peking Joint Center for Life Sciences, School of Life Sciences, Tsinghua University, 100084 Beijing, China
[5]Quantitative Biology Institute, Yale University, New Haven, CT 06511

*These authors contributed equally [#]Correspondence: joe.howard@yale.edu

Summary

The highly ramified arbors of neuronal dendrites provide the substrate for the high connectivity and computational power of the brain. Altered dendritic morphology is associated with neuronal diseases. Many molecules have been shown to play crucial roles in shaping and maintaining dendrite morphology. Yet, the underlying principles by which molecular interactions generate branched morphologies are not understood. To elucidate these principles, we visualized the growth of dendrites throughout larval development of *Drosophila* sensory neurons and discovered that the tips of dendrites undergo dynamic instability, transitioning rapidly and stochastically between growing, shrinking, and paused states. By incorporating these measured dynamics into a novel, agent-based computational model, we showed that the complex and highly variable dendritic morphologies of these cells are a consequence of the stochastic dynamics of their dendrite tips. These principles may generalize to branching of other neuronal cell-types, as well as to branching at the subcellular and tissue levels.

Key words

Branching morphogenesis, neuronal development, dendrite, arbor, dynamic instability, agent-based model

Introduction

Neurons are polarized cells (Cajal, 1906) whose axons and dendrites are often highly

branched. Branching provides the surface area necessary for dendrites to receive

inputs from thousands of other cells or from the environment (Lefebvre et al., 2013), and

for axons to output signals to multiple cells. In this way, branching facilitates the high

connectivity of the brain (Lefebvre et al., 2013). The morphology of the neurons,

together with their synaptic connections (Fornito et al., 2015; Kandel et al., 2013;

Seung, 2012; Zheng et al., 2018), defines the structure of the nervous system, the

connectome, which is viewed as a prerequisite for understanding brain function (Denk

et al., 2012). Knowing the rules of neuronal branching is expected to facilitate mapping

the connectome by providing anatomical constraints or laws, as well as giving insight

into genetic disorders that affect morphology (Forrest et al., 2018; Kapitein and

Hoogenraad, 2015; Koleske, 2013; Kulkarni and Firestein, 2012; Moughamian and

Holzbaur, 2017). Whereas much is known about the growth of axons, whose tips, the

growth cones, are guided by extracellular signals and guidepost cells (Lowery and

Vactor, 2009; Sánchez-Soriano et al., 2007; Stoeckli, 2018; Tessier-Lavigne and

Goodman, 1996), the mechanisms underlying the growth and branching of dendrites is

poorly understood.

While many molecules have been shown to play crucial roles in shaping and

maintaining dendrite morphology, the underlying principles by which molecular

interactions generate branched morphologies is not understood. To elucidate these

rules, we have focused on dendrite morphogenesis in class IV dendritic arborization

(da) neurons in *Drosophila*, a model system for dendritogenesis (Dong et al., 2015; Jan

and Jan, 2010; Singhania and Grueber, 2014). These nociceptive neurons form a highly

branched meshwork just under the cuticle that senses puncture of the larva by the

ovipositor barbs of parasitic wasps and initiates avoidance behaviors (Basak et al.,

2021; Robertson et al., 2013). Class IV cells are ideal for studying branching

morphogenesis because they grow rapidly over 5 days of larval development, their

branches are non-crossing due to self-avoidance mediated by the Down's syndrome

cell adhesion molecule (Hughes et al., 2007; Matthews et al., 2007; Soba et al., 2007)

and other molecules (Emoto et al., 2004; Parrish et al., 2009), and can be visualized

using cell-specific labeling (Grueber et al., 2003; Jan and Jan, 2010). Many molecules

that participate in dendrite morphogenesis have been identified: transcription factors

(Jinushi-Nakao et al., 2007); extracellular matrix and integrins (Han et al., 2012; Kim et

al., 2012); actin-associated proteins (Lee et al., 2003; Li and Gao, 2003; Scott et al.,

2003; Stürner et al., 2019); microtubule motors such as dynein and kinesin (Arthur et

al., 2015; Satoh et al., 2008; Zheng et al., 2008); microtubule regulators such as spastin

(Sherwood et al., 2004), katanin (Stewart et al., 2012) and γ-tubulin (Ori-McKenney et

al., 2012); and microRNAs such as *bantam* (Parrish et al., 2009). The major difficulty,

however, is that it is currently not possible to make a mechanistic connection between

developmental processes occurring at the molecular and subcellular levels, and the

morphology of the entire dendritic arbor.

While several theoretical and computational models can predict dendrite-like

branched morphologies, they are not grounded in molecular or development processes

data. Early models, designed to describe and classify neurons, reconstituted

morphologies based on the statistical properties of the observed arbors themselves

(Ascoli and Krichmar, 2000; Nanda et al., 2018). Optimization-based models that

minimize wiring (i.e., the total lengths of the branches) capture key features of neuronal

morphology (Baltruschat et al., 2020; Cuntz et al., 2010), but lack connection to the

molecular processes. Other models use hypothetical branching properties (Ganguly et

al., 2016; Palavalli et al., 2021), or abstract processes such as diffusion-limited

aggregation (Luczak, 2006), or Turing-like pattern formation (Sugimura et al., 2007). On

the other hand, branching models developed for tissues such as the lungs (Metzger et

al., 2008) and kidneys (Lefevre et al., 2017; Short et al., 2014), produce stereotyped

morphologies that differ from the highly variable morphologies of neurons (Kanari et al.,

2018). Stochastic models developed for other tissues, such as the mammary glands,

use properties that are specific to these systems, such as tip bifurcation (Hannezo et al.,

2017). Thus, current computational models fall short in providing a mechanistic

understanding of dendrite morphology.

To circumvent these limitations, we have formulated a computational model that

is based entirely on experimentally observed properties of dendrites measured over

their development. The data-based model takes as input tip-growth dynamics,

branching rates and self-avoidance measured using high-resolution, live-cell imaging in

the developing animal. The model successfully recapitulates class IV dendrite

morphogenesis and shows how the complex and variable morphology of dendritic

arbors emerge from the microscopic dynamics of dendrite tips and provides insights into

several mutant phenotypes.

Results

As *Drosophila* larvae grow (Figure 1A), the arbors of their class IV dendrites also grow (Figure 1B). By the end of larval development, the meshwork of branches covers the larval surface like chainmail and the individual dendritic arbors fill the eight abdominal segments with cell widths up to 500 μm. When the first instar larva hatches at 24 hr after egg-lay (egg-lay is defined as time zero), the dendritic arbors are smaller than the segments in which they are located. Over the next 24 hr, the arbors grow faster than the segments and touch the segment boundaries in a process called tiling (Grueber et al., 2002); at 72 hr the arbor densely fills the segment (Parrish et al., 2009) and then grows at the same rate as the segments (Figure 1, inset). We sought to understand how dendrites tile and infill.

Figure 1. Growth of larvae and class IV neurons over development. A Whole-mount, living larvae imaged by spinning-disk confocal microscopy at 24 hr to 120 hr (egg-lay defined as time zero). Anterior (A) is up and posterior (P) down. Left (L) and right (R) as viewed from the dorsal side (Methods). Class IV neurons are marked with the transmembrane protein CD4 tagged with GFP (;;ppkCD4-tdGFP). **B** Individual class IV cells from the A3 or A4 segments. An A3 segment is outlined in A (120 hr). **Inset** At 24 hr, the cell widths (filled circles and dashed lines) are smaller than the segment (filled lines). In the next 24 hr, they touch the growing segment boundaries (dark gray) and infill by 72 (light gray).

Uniform branch dilation cannot account for dendrite arbor growth

We first asked whether class IV arbors grow through dilation of their branches (Figure 2A, upper, purple dendrites)—that is, by the expansion of branches along their lengths (Yang and Chien, 2019). Dilation accounts for the growth of the overlying epidermal cells (Parrish et al., 2009) and of class I cells (Castro et al., 2020; Palavalli et al., 2021),

which expand concomitantly with the segments. To test the role of dilation in arbor

expansion, we reimaged the same neurons at discrete times over development, at 24 &

48 hr, and at 48 & 96 hr (Figure 2B). As the arbors grow, there is continuous addition

and removal of branches. Nevertheless, it was possible to identify shared structural

features of proximal, non-terminal branches (branches that lie between two branch

points as opposed to terminal branches that end in a tip) (Figure 2 C,D). The fractional

increases in lengths of these identified branches were considerably less than the

fractional increase in length of the AP and LR segments (Figure 2E), in agreement with

earlier measurements (Baltruschat et al., 2020). This finding implies that there must be

slow slippage between the class IV cells and the adjacent epidermal cells (Han et al.,

2012). Because branch dilation contributes only 3% to 11% of the overall growth of the

dendrites, other mechanisms must contribute to the bulk of arbor growth.

Figure 2. Branch dilation does not account for dendrite growth. A Two models for dendrite growth: dilation & infilling (upper) and branching & tip growth (lower). **B** Maximum-projection image of a GFP-labeled class IV neuron cell at 24 hr (magenta color) and same cell imaged at 48 hr (green color). 24 and 48hr images are combined with a leftward displacement of the latter. **C,D** Conserved non-terminal branches are marked with the same color and number. **E** The fractional length changes from 24 hrs to 48 hrs and from 48 hrs to 96 hrs were calculated for the AP and LR segments, and for the non-terminal (NT) branches (Methods). Each blue and green circle is a different larva; the red circles correspond to several branch measurements in each of 6 larvae.

Terminal dendrites grow from their tips and not from their bases

An alternative hypothesis to dilation is that the growth of the arbor is due to branching

and subsequent lengthening of the newly formed terminal branches (Figure 2A, lower).

To test this hypothesis, the behavior of terminal dendrites was examined. Following

birth by lateral branching from existing branches (Gao et al., 1999), terminal dendrites

can: lengthen; shorten spontaneously or following contact with another branch; and

pause (Figure 3A). Time-lapse imaging (Supplementary Movies S1-S5) suggests that

lengthening is due primarily to the addition of material near the tip. For example, branch

points and bends remain fixed during growth and shortening (Figure 3A, top-left panel;

Supplementary Movies S6-10), and the distance between these landmarks and the tips

do not change. These observations argue against growth at the proximal end or uniform

dilation along the length of terminal dendrites. Because new branches can form as close

as 2 μm to an existing tip, we estimate that growth occurs within ~2 μm of the distal

ends. Thus, tip growth, which occurs on timescales of minutes, may contribute to the

overall growth of the arbor.

Figure 3. **Dendrite tips transition between growing, shrinking and paused states A** Class IV dendrites growing (green star), birthing (yellow), colliding (blue), shrinking (red), and pausing (orange). Maximum projection spinning disk images were collected every 5 seconds. **B** The length of a dendrite as a function of time (see Methods). The black open circles represent the initial piecewise linear fit using $N/6$ segments, where N is the total number of frames. The gray dots show the fitting after the iterative merging process (see Methods). Green, orange, and red indicated periods of growth, pausing, and shrinkage. **C** The root-mean-squared error before (black) and after (gray) merging for 91 trajectories (24 hr). The average error is 50 nm. **D** The velocity distribution shows three distinct peaks representing the growing (G), paused (P), and shrinking (S) states. **E** Transition rates between the three states.

High-resolution tracking shows that tips transition between periods of constant growth velocity

To determine whether tip growth can account for arbor expansion, we tracked the

lengths of terminal dendrites over time with an accuracy of ~0.1 μm (Methods,

Supplementary Figure 1 A-F). To ensure that mounting and imaging larvae did not

interfere with growth, we restricted imaging to 20- to 30-minutes (Supplementary Figure

2A). Typical trajectories show that dendrite growth is highly dynamic, with large

fluctuations in velocity (Figure 3B, and Supplementary Figure 1 G, I, K). To analyze tip

trajectories, we first considered tip growth as a diffusion-with-drift process, a common

way to describe particles moving in a flow. However, we ruled out this description

because there were extended times of near-constant velocity: the green, red, and

orange lines in Figure 3B (and Supplementary Figure 1G, I, K) clearly indicate periods

of elongation, shortening, and stationarity. We therefore fit the trajectories to a

piecewise linear continuous function, for which fast algorithms exist (D'Errico, 2021).

This initial segmentation into regions of constant velocity provided a good fit to the

tracking data (Figure 3B and 3C, black circles), showing that tip trajectories can be

decomposed into sequential periods of linear growth or shrinkage.

Dendrite tips undergo dynamic instability

We then asked the more difficult question of whether tips undergo dynamic transitions

between growing (G), shrinking (S), and paused (P) states. In other words, can we

classify the regions of constant growth into just three states such that transitions only

occur between different states. Such a description is analogous to the dynamic

instability of microtubules (Mitchison and Kirschner, 1984), which transition

stochastically between two states: growing and shrinking.

To test whether a three-state dynamic model could account for tip growth, we

assigned each region to be in a growing, shrinking or paused state by fitting the

histogram of velocities with a three-peaked distribution, such as shown in Figure 3D, to

define velocity thresholds between growth and pause, and shrinkage and pause. We

then merged adjacent regions that belonged to the same state. Through an iterative

procedure (see Methods), we segmented the trajectories into growing, shrinking, and

paused states, with transitions only between different states. The resulting trajectory

(Figure 3B, gray lines) was an excellent fit to the data: the root-mean-squared error was

on average ~0.05 µm (Figure 3C), accounting for 85% to 99% of the variance

(Supplementary Figure 1 H, J, L). From these data, we calculated (i) the growing and

shrinking speeds (Figure 3D) and (ii) the rates of the transitions between the three

states (Figure 3E). At 24 hr, the growing and shrinking speeds were 1.61 μm/min and

1.52 μm/min, respectively, and the rates ranged from 0.31 to 0.95 per minute,

corresponding to average lifetimes of individual states between 0.6 to 1.5 minutes. The

net speed of dendrite elongation, ~0.034 μm/min (Table 1A), is much smaller than the

average speed in the growing state because the dendrites spend roughly equal times in

the growing and shrinking states.

Thus, the growth trajectories are in accord with a three-state kinetic scheme,

which provides a succinct yet comprehensive description of tip dynamics. This scheme

is a generalization of the Dogterom and Leibler model of microtubule dynamic instability

(Dogterom and Leibler, 1993), with inclusion of a third, paused state, and the growing

and shrinking states having distributions of speeds.

Dendrite dynamics and branching rates change over development time

Throughout development, the growing and shrinking speeds were roughly unchanged

(Table 1A). The main change over development was that the transition rates out of the

paused state decreased two- to four-fold and the transition rates into the paused state

increased by about 50%. As a result, the dendrites spend more time in the paused

state: they become less dynamic.

Branching, which always occurs on the sides of existing branches, also slowed

down over development. The branching rate per unit dendrite length decreased roughly

ten-fold from 24 to 96 hr (Figure 4A, Supplementary Table 1). This decrease is another

manifestation of dendrites becoming less dynamic over time. The geometry of

branching, however, remained constant over development: the mean angle of a new

(daughter) branch was close to 90° at all developmental stages (Figure 4B,

Supplementary Table 1), and the spatial distribution of branching remained roughly

uniform (Figure 4C and D). In summary, both growth and branching are highly

stochastic throughout development, though the transition and branching rates slow

down over time.

Figure 4. Dendrite branching over development. **A** The rate of appearance of new branches normalized by the total branch length is plotted against developmental time. Each symbol represents a neuron from a different larva (except at 24 hr where 3 neurons in each of 3 larvae were analyzed). The curve is an exponential fit with an offset (dotted line). Inset: The total branching rate per cell. **B** The distribution of branch angles between daughters at different developmental stages. The angle is zero when the new daughter grows parallel to the mother. Numbers of neurons: 6 (24 hr), 7 (36 hr), 4 (48 hr and 96 hr), 5 (72 hr) **C** Spatial distribution of branching events at 24 hr (9 neurons from 3 larvae). **D** Spatial distribution of branching events in 96 hr (6 neurons from 6 larvae). In both **C** and **D**, the soma positions are centered at the origin. Different colors represent different cells.

Dendrite tips retain memories: their dynamics are not Markovian

We tested whether the transitions between growth states are Markovian, meaning that they depend only on their current state: i.e., they do not depend on the history, and there is no memory of earlier states. Consistent with a Markov process, the lifetimes of the states were approximately exponential at long times (Supplementary Figure 3 D-F), and the probability of a transition in the sequence of occupied states (e.g., GPGSPGPSG…) did not depend on the previous state. For example, we found that the likelihood of G→P did not depend on the prior state: that is SG→P and PG→P were equally likely.

There were several violations of the Markov property, however. First, following contact of a tip with another branch, the growth dynamics change: the rates out of the growing state increase, so the dendrites spend less time growing (Table 1B). Therefore, in addition to contact-induced retraction, there is a long-lasting alteration of the dynamics: the average tip-growth rate changes from positive to negative (Table 1), and the post-contact dendrites shrink on average. This alteration implies that there is a long-lasting memory of the collision. Second, we found that the lifetimes of newly born dendrites were longer than expected for a Markov process. Following birth into the growing state, the transition rates, which were measured for older dendrites (>5 minutes after birth), predict that there will be an initial linear decrease in surviving dendrites due

to the chance that a growing dendrite stochastically switches into a shrinking state,

which then shortens and disappears. Instead, we found that the survival curve was

initially flat, consistent with an initial growing state of 0.3 minutes (Supplementary Figure

4A). Such a survival curve is another violation of the Markov process and implies that a

newborn dendrite retains a memory of birth. Third, growth and shrinkage events with

higher absolute speeds tended to have shorter lifetimes (Supplementary Figure 3A, B).

Branching also showed non-Markovian properties: for example, rebranching occurred

more often at points where an earlier dendrite had disappeared (Supplementary Figure

4B, C). Thus, while branching and growth are highly stochastic, they deviate from being

Markovian, indicating memory and "hidden variables" that influence the dynamics.

These hidden variables are likely to be long-lived biochemical states (e.g.,

phosphorylation) triggered by dendrite birth or tip contact with other dendrites, which

influence the dynamics.

An agent-based model incorporates measured tip properties

To test whether the dynamical properties of the dendrites measured above can account

for the observed morphology, we developed an agent-based computational model to

predict morphologies based on tip dynamics. The elementary "particle" in the model is

the dendrite tip, which serves as the agent (Figure 5A).

Figure 5: Agent-based computational model A Cartoon diagram depicted the different components of the model: tips are born by branching and transition between growing, shrinking, and paused states. Upon contact with another branch, the tips retract. **B** Diagram of the transitions. Parameters are listed in Table 1 and Supplementary Tables 1-2.

Tips were simulated using rules that closely followed the experimental

measurements (see Figure 5B, Methods). Terminal dendrites lengthen and shorten with

speeds sampled from the growth and shrinkage distributions (e.g., Figure 3D). They

transition between growing, shrinking, and paused states according to the measured

transition rates (Table 1), which were linearly interpolated between different

developmental stages. Tip birth occurred randomly in time and space along extant

branches with the measured branching rates (Figure 4A). The nascent daughter branch

was assumed to start in a growing state (Figure 5) with an initial length of $0.5\ \mu m$

(Supplementary Table 2) and included an initial lag of $0.3\ \min$ during which the transition

out of the growing state was forbidden, in accordance with our observations. Dendrite

death occurred when the last point disappeared during a shrinkage event. Contact,

defined as a tip getting closer than $0.15\ \mu m$ to another branch (roughly the branch

radius (Liao et al., 2021), switched a growing tip to a shrinking one with the post-contact

dynamics (Table 1B).

The initial larval morphology at 24 hr was established by (i) growing two to four

branches from a point (the origin) using the embryonic growth parameters at (18-20 hr)

and (ii) allowing growth until the total branch length and number reached their values at

24 hr. To model the segment boundary, we assumed that contact with neighboring

dendrites induced shrinkage (Parrish et al., 2009). Neighbor interaction was

implemented using a periodic boundary condition such that one side of a growing

neuron feels its opposite side as if growing on the topological equivalent of a torus. Over

time, we gradually increased the size of the boundary according to the measured

segment growth rates (Figure 1 inset).

Simulated dendritic trees recapitulate coarse-grained features of dendrite

morphology

We simulated dendritic trees using the parameters from Table 1 and Supplementary

Tables 1,2, all of which were measured or tightly constrained by experiments. The

simulations (Figure 6A) recapitulated key properties of real arbors (Figure 6B).

(i) **Tiling:** Simulated arbors tiled the segments, like real arbors. In the absence of

a boundary, the widths of simulated arbors initially grew at 10 μm/hr, which is faster

than the segment growth of 4 μm/hour; the simulated arbors, therefore, filled the

segments between 48 to 72 hr (Supplementary Figure 5Aii). After that, the simulated

arbors grew at 4 μm/hr, just sufficient to keep up with segment growth.

Figure 6. The agent-based model accords with overall neuronal growth A Example of a simulated
neuron using the parameters in Table 1 and Supplementary Tables 1-2 at different developmental stages.
Same scale bar as in B. **B** Example of the skeleton of a real neuron at 120 hr. **C** Total branch length over
developmental for simulated and real neurons. **D** Total number of branches. **E** Mean branch length (total
branch length/total number of branches).

(ii) **Total branch length and number**: The simulations predicted the observed

increases in total branch length (Figure 6C) and number (Figure 6D), as well as the

mean branch length (Figure 6E). The branch length distributions were roughly

exponential in the simulations and the data (Supplementary Figure 6 A-E). An

interesting feature of the branch number is the initial burst (24-36 hr) and subsequent

plateau (36-72 hr). The burst is predicted by the model and arises from two features of

tip growth: the high initial branching rate (Figure 4A), and the perseverance of the initial

growth of branches (i.e., a delay in transitioning out of the growing state). Without

perseverance, which is a memory of birth, the plateau is less pronounced, showing that

the initiation of branching is an essential determinant of arbor morphology.

There were some discrepancies between the data and the model. For example,

the branch number of the simulated arbors saturated at 120 hr, while that of the real

arbors continued to increase with an associated late decrease in mean branch length of

real arbors. These discrepancies may indicate that dynamical properties change after

96 hr. Also, the high density at AP boundaries (Figure 6B, right) arises from cell-cell

interactions, which we did not include. Nevertheless, we believe that the model is in

good agreement with the average properties of the arbors.

Simulated arbors capture the variability of dendrite morphology

In addition to predicting average branch properties, the model also predicted the

variation of these features. For example, the measured branch numbers are highly

variable, with coefficients of variation (SD/mean, CV) ranging from 0.11 to 0.22 over

development. This CV is even larger than that of a Poisson process, a prototypical

random process whose CV equals the inverse square root of the branch number

(expected range 0.02 to 0.06). This comparison to a Poisson process indicates that

branch number is highly variable from arbor to arbor, a manifestation of the stochasticity

of the morphology. Total branch length and average branch length were also highly

variable. Importantly, the simulations recapitulated this variability (Figure 6C-E). Thus,

the model predicted both the average properties and the stochasticity of the branch

number and length.

Simulated arbors recapitulate fine-scale properties of dendrite morphology

The branches of both the simulated and real arbors formed dense meshworks (Figure

7A,B). We estimated the extent to which the branches cover the arbor using the box-

counting method (Falconer, 1990) in which the number of boxes that contain a branch is

plotted against the size of the boxes (Figure 7C). We found that the logarithm of box

number was approximately proportional to the logarithm of box size, indicating that the

patterns have scale-free and fractal-like properties. The log-linear relation breaks down

at box sizes below 5 μm, the size of the "holes" in the pattern due to the average branch

size. We defined the fractal dimension as the log-log slope in the central region of the

plot encompassing the middle 50% of the points (Figure 7C dashed lines). The fractal

dimension increased from 1.4 to 1.8 from 24 hr to 120 hr for both the simulated and real

arbors (Figure 7D). Because a region containing a single line has a fractal dimension of

1 whereas a region completely filled region has a fractal dimension of 2, the dendritic

patterns are of intermediate dimension and at 120 hr nearly fill the plane (fractal

dimension 1.8). Though the fractal dimensions of real arbors were consistently lower by

about 0.1, we nevertheless conclude that the simulated arbors recapitulate the real

arbors in this metric.

Figure 7. The agent-based model recapitulates the fine-scale patterns of real arbors. A Model arbor at 72 hr. Light green branches indicate the outer 5% of the arbors excluded from the analysis to mitigate against spurious boundary effects (see Methods). Boxes with two different sizes are shown in the background. **B** Skeletonized real arbor at 72 hr. Light gray branches were excluded from the analysis. **C** The number of boxes that contain a branch (*y*-axis) is plotted against the box size (*x*-axis). The slopes over the central 50% of the data (vertical dashed lines) define the fractal dimensions. **D** Fractal dimensions of real and simulated arbors both increase over developmental. **E** Diagram defining the radial orientation. **F** Radial orientation of a simulated tree (green solid line) and a real arbor (black) at 72 hr. Simulated tree with pausing instead of retraction (green dotted curve).

Simulations predict the radial orientation of dendrite branches

We discovered that class-IV cells have an unexpected long-range order: branches are

not randomly oriented but instead tended to be parallel to the radial orientation (Figure

7E-F, Supplementary Figure 6 F-J). The simulations also displayed radial orientation.

Radial orientation is a consequence of contact-based retraction; if contact-based

retraction is replaced by contact-based pausing, the radial orientation was greatly

reduced (Figure 7F, dotted curve).

Discussion

We have discovered that the tips of *Drosophila* class IV dendrites transition

stochastically between three states - growing, shrinking, and paused. This allows

dendrites to explore extracellular space, analogous to the exploration of intracellular

space by microtubules undergoing dynamic instability (Kirschner and Mitchison, 1986).

Our modeling shows that these transitions, together with lateral branching, and contact-

mediated retraction, give rise to the complex and highly variable morphology of the

dendritic arbors, allowing them to fulfill their biological functions. The dense, almost

unbroken meshwork optimizes detection of the fine ovipositor barbs of parasite wasps

(Basak et al., 2021; Robertson et al., 2013). And the radial orientation of branches, a

form of long-range order that emerges from local interactions (Toner and Tu, 1995;

Vicsek et al., 1995), namely contact-induced retraction, reduces the path distance to the

cell body: this minimizes signaling delays and wiring costs (Baltruschat et al., 2020).

Thus, stochastic tip dynamics may partially solve the riddle of how "the morphological

features displayed by neurons appear to obey precise rules that are accompanied by

useful consequences" (Cajal, 1995).

Dendrite tips: an intermediate organizational principle of dendrite morphology

Our results strongly support the concept that the dendrite tip is a "branching engine …

that initiates, directs, and maintains branch outgrowth during development and

regrowth" (Lu and Werb, 2008). The dendrite tip, with a diameter of only 0.2 µm (Liao et

al., 2021) and with dynamics on the timescale of ~1 minute (the state lifetimes),

generates structures up to 500 µm in diameter (>1000-times larger sizes) over five days

(>1000-times longer times). Tips, therefore, are intermediate in length- and timescales

between molecules and morphology.

The concept of the dendrite tip as a branching machine has four important

implications. First, if the molecular basis of tip growth and branching can be elucidated,

our agent-based model will provide a full connection between genotype and phenotype,

with the caveats that morphogenesis is stochastic and some features such as three

dimensionality are not included. Second, altered tip dynamics due to mutations and

diseases may underly altered dendrite morphologies (see Introduction and the

Predictions of the model section). Third, tip rules may specify neuronal identity, often

defined by dendrite morphology (DeFelipe et al., 2013). And fourth, the stochastic

nature of the tip rules may facilitate the evolution of neuronal cell types. This is because

developmental stochasticity amplifies genetic variation by allowing a large class of

morphologies to be sampled for each genotype. While some morphological outliers may

function poorly, others might be beneficial, and genetic and/or epigenetic mechanisms

could selectively stabilize these beneficial morphologies.

Potential molecular mechanisms underlying dendrite tip dynamics and branching

Neurite elongation by tip growth also occurs in axons (Dent et al., 2011) and in other

dendrites, such as those in *C. elegans* PVD neurons (Liang et al., 2020). An important

difference between dendrite tips in class IV cells and the growth cones seen in these

other cells is that the tips of class IV dendrites are much smaller. The diameter of class

IV terminal dendrites is only ~200 nm (Liao et al., 2021), as small as a single filopodium,

the finest feature of growth cones observed under the light microscope. Thus, tip-growth

mechanisms in class IV dendrites likely differ from growth-cone-based growth. An

important open question is how the cytoskeleton and membranes reach the dendrite

tips: what are the relative contributions of diffusion, filament polymerization (Delandre et

al., 2016; Jinushi-Nakao et al., 2007; Nithianandam and Chien, 2018; Ori-McKenney et

al., 2012; Sears and Broihier, 2016; Yalgin et al., 2015), motor-driven transport (Arthur

et al., 2015; Weiner et al., 2016) and motor-driven sliding (Winding et al., 2016)?

Morphological predictions of the model

The agent-based model allowed us to explore which parameters are important for arbor

growth and mesh size (Supplementary Figure 5) and provides hypothesis for the

phenotypes of mutants.

A surprising finding was that branching drives overall arbor growth: increasing

the branching rate not only increased the number and density of branches, as expected,

but also increased arbor size. Indeed, setting the average velocity of dendrite extension
to zero still resulted in arbor growth, albeit slower, showing that branching without net tip
growth can drive arbor expansion. The modeling showed that increasing the fluctuations
in branch length (by reducing the tip-transition rates) also increased the growth rate.
This suggests an unexpected growth mechanism: length fluctuations are locked in by
the formation of new branches. The branching angle also affected arbor growth: if the
branch angle was decreased to 45° (i.e., growth towards the direction of the mother
branch), arbor growth increased, showing that outward growing branches are more
likely to survive. The persistence of branch growth after birth was also important.
Therefore, branching is essential not just for infilling the network but also for expansion.

Our model shows dendrite density is set by the interplay between branching and
self-avoidance, which acts as negative feedback. Inhibition of self-avoidance increases
the branch density as observed by several mutation studies (Emoto et al., 2004;
Hughes et al., 2007; Matthews et al., 2007; Soba et al., 2007). By allowing the dendrite
tips to ignore contact-based retraction with the neighboring branches, our model
produced high density of branches (see Supplementary Table 3).

Our observation that branching is an intensive property—the total rate of
formation of branches is almost independent of arbor size (Figure 4A, inset)—is
evidence that there are only a limited number of "branching factors" being produced in
the entire cell per unit time. The uniform distribution of new branches suggests that the
branching factors are dispersed throughout the cells, perhaps by molecular motors.
Several phenotypes of molecular motor mutants in class IV cells support this
hypothesis. The perturbation of molecular motors and their adapter proteins, including
dynein (Arthur et al., 2015; Satoh et al., 2008; Zheng et al., 2008) and kinesins (Kelliher
et al., 2018; Satoh et al., 2008), result in non-uniform branch densities, as expected if
the distribution of branching factors were disrupted (see Supplementary Table 3).

The transition rates between growth states are also important. For example, the
dendrite tips of Katanin (Kat-60L1) mutants spend less time in the paused states and
more time in the shrinking states compared to controls (Stewart et al., 2012). Our
simulations recapitulated the sparser morphologies reported in Stewart et al. (see
Supplementary Table 3).

General mechanisms of branching morphogenesis

Dendrite tips share features of branch tips in other systems. Tip cells drive branching in branched tissues such as the mammary glands (Lu and Werb, 2008). The growing ends of cytoskeletal filaments, with associated nucleation factors, drive branching of organelles such as the microtubule-based mitotic spindle (Decker et al., 2018; Petry et al., 2013) and the actin-based lamellipodium (Pollard and Borisy, 2003). In all three cases (dendrites, tissues and cytoskeleton), the tips operate at shorter length-scales and timescales than those of the structures they produce. Furthermore, they all respond to external signals: contact-based retraction of dendrite tips, cortex-induced catastrophe of microtubule ends (Komarova et al., 2002), and self-avoidance of mammary-gland branches (Lu and Werb, 2008). Finally, all three are stochastic and generate highly variable morphologies. Given these commonalities, it is likely that the principles that we have elucidated for dendrites generalize to other branched systems.

Acknowledgments

We thank all members of the Howard lab for many helpful discussions and suggestions. Special thanks to Dr. Ashley Arthur for proofreading the manuscript. We thank Hermann Cuntz, Damon Clark, and John Carlson for comments on an earlier version of the manuscript, and Chun Han and the Bloomington Drosophila Stock Center for fly stocks. O.T. was supported by the Fonds de Recherché du Quebec - Nature et Technologies; X.L. was supported by the National Natural Science Foundation of China (NSFC Grant 31671389, to X.L.) and the Max-Planck Partner Program; and J.H. was supported by NIH grants DP1 MH110065 and R01 NS118884.

Author Contributions

SSh, SSu and JH designed the experiments based on early results by XL. SSh conducted the experiments and analyzed branching. SSu, OT, YT and JH conceptualized the models, which were implemented by SSu, who performed image and tip tracking analysis and wrote the source code for the *in-silico* model, based on earlier work by OT. SSh, SSu and JH wrote the manuscript.

611 **Declaration of Interests**

612 The authors declare no competing interests.

Table 1

A: Pre-contact tip-dynamics parameters

Age (hr)	Tip velocity parameters			Tip transition rates						Corr.	Average tip velocity	
	μ_G, σ_G (µm/min)	μ_P, σ_P (µm/min)	μ_S, σ_S (µm/min)	k_{GP} (min⁻¹)	k_{GS} (min⁻¹)	k_{PG} (min⁻¹)	k_{PS} (min⁻¹)	k_{SG} (min⁻¹)	k_{SP} (min⁻¹)	r & p	v_d^{Tracks} (µm/min)	$v_d^{TranMat}$ (µm/min)
18-20 (E)	0.37, 0.34 $V_G = 1.53$	0, 0.36	0.43, 0.37 $V_S = 1.65$	0.696	0.509	0.423	0.296	0.669	0.71	0.017 0.09	0.097 ±0.0158	0.100 ±0.0151
24 (L1)	0.41, 0.36 $V_G = 1.61$	0, 0.34	0.35, 0.37 $V_S = 1.52$	0.784	0.64	0.335	0.314	0.598	0.946	0.008 0.65	0.034 ±0.0196	0.038 ±0.0191
48 (L2)	0.40, 0.39 $V_G = 1.61$	0, 0.25	0.0, 0.41 $V_S = 1.1$	0.933	0.435	0.155	0.235	0.282	1.251	0.045 0.24	0.020 ±0.1545	0.026 ±0.0177
96 (L3)	0.36, 0.52 $V_G = 1.64$	0, 0.25	0.19, 0.44 $V_S = 1.33$	0.923	0.799	0.116	0.117	0.575	1.276	0.002 0.9	0.008 ±0.0039	0.022 ±0.0177

B: Post-contact tip-dynamics parameters

Age (hr)	Tip velocity parameters			Tip transition rates						Corr	Average tip velocity	
	μ_G^*, σ_G^* (µm/min)	μ_P^*, σ_P^* (µm/min)	μ_S^*, σ_S^* (µm/min)	k_{GP}^* (min⁻¹)	k_{GS}^* (min⁻¹)	k_{PG}^* (min⁻¹)	k_{PS}^* (min⁻¹)	k_{SG}^* (min⁻¹)	k_{SP}^* (min⁻¹)	r & p	v_d^{Tracks} (µm/min)	$v_d^{TranMat}$ (µm/min)
18-20 (E)	0.53, 0.49 $V_G^* = 1.9$	0, 0.28	0.53, 0.54 $V_S^* = 1.9$	0.635	0.992	0.263	0.401	0.469	0.593	0.07 0.12	-0.42 ±0.07	-0.34 ±0.07
48 (L2)	0.40, 0.50 $V_G^* = 1.7$	0, 0.25	0, 0.38 $V_S^* = 1.1$	1.446	1.24	0.134	0.29	0.239	0.814	0.018 0.76	-0.23 ±0.05	-0.19 ±0.04

616 Errors are SE.
617

Methods

Fly Stocks and maintenance
The fly line *ppk-cd4-tdGFP* was used to image class IV dendritic arborization neurons and was a kind gift from Dr. Han Chun (Cornell University). Fly crosses were maintained in fly chambers at 25 °C, 60% humidity in a Darwin Chamber with 12 -hour light/dark cycles. An apple-agar plate was used to collect the fly embryos and a big drop of yeast paste was put in the center of the agar plate to induce egg-laying.

Apple-agar plates were made by mixing 4X apple juice concentrate (355 ml), water (300 ml), dextrose (155 gm), sucrose (80 g). This solution was stirred and heated to dissolve the sugars, and agar (Bacto agar, Becton Dickinson, 60 g) and 1.25N NaOH (70 ml) were added. The solution was covered loosely with foil and autoclaved in the liquid cycle for 30 min. The plating mixture contained 100 ml of this apple-agar concentrate, 197 ml of water, and 3 ml of Acid mix A—an equal mixture of propionic acid (100%, 83.6 ml, and 16.4 ml water) and phosphoric acid (100%, 8.3 ml and 91.7 ml water).

For neuron morphometrics, embryos were collected every 15 minutes and imaged when they reached the appropriate age **A**fter **E**gg **L**ay (AEL): 24hr, 48hr, 72hr, 96hr, and 120hr.

Sample preparation
For imaging, embryos of appropriate age (18 hr to 22 hr AEL) were collected from apple-agar plates and dechorionated by gently rolling them on a piece of double-sided tape stuck to a glass slide. The dechorionated embryos were then placed with their dorsal side down on a No. 1.5 coverslip, MatTex, with a small drop of halocarbon oil 700. A piece of wet Kim wipe was placed near the embryos to maintain humidity during imaging. No anesthetics were used for embryo imaging. For larvae imaging, larvae of ages 24 hr, 48 hr, 72 hr, 96 hr, and 120 hr were washed with 20% and 5% sucrose solution, anesthetized using FlyNap (Carolina Biologicals, Burlington, NC, USA), and transferred to apple-agar plates to recover for 1-5 minutes. After recovery, larvae were gently placed with their dorsal side up on a 1% agar bed adhered to a glass slide and imaged in a drop of 50% PBS, 50% Halocarbon oil 700 (Sigma Aldrich). Larvae were further immobilized by gently pressing them with a 22mm X 22mm coverslip lined with Vaseline or vacuum grease.

Imaging
Samples were imaged on a spinning disk microscope: a Yokogawa CSU-W1 disk (pinhole size 50 µm) built on a fully automated Nikon TI inverted microscope with perfect focus, 488nm laser illumination at 18-21 % laser power, either a 40X (1.25 NA, 0.1615-micron pixel size) or a 60X (1.20 NA, 0.106-micron pixel size) water immersion objective, an sCMOS camera (Zyla 4.2 plus), and Nikon Elements software. The temperature of the sample region was maintained using an objective space heater at 25°C (OKO labs stage heater). Samples were manually focused to identify abdominal third and fourth segments (A3 or A4 neurons) before image acquisition. Full-frame movies (2048×2048 pixels) containing 6 to 12 1-µm sections were collected every 4-6 s. Static images for morphometric studies were acquired using a 60X water immersion

664   objective for 24 hr and 40X objective for later stages. Images were stitched using in-
665   house code (https://github.com/oliviertrottier/neuron-stitch). Movies were curated for
666   subsequent offline analysis. Image analysis (segmentation, skeletonization, branching
667   analysis, and angle measurements) was done using ImageJ.
668

Controls for growth
Class IV neurons are susceptible to mechanical pressure, which, if too large, stops
growth and causes degeneration. Therefore, embryos were imaged without an overlying
coverslip. Larvae were immobilized with minimal pressure under a coverslip and were
placed on a 1% agar bed. As a control, we plotted the size of neurons throughout
imaging to confirm that they remain on the "standard" growth curve (Supplementary
Figure 2 A).
676

Segment length determination
The *Drosophila* larval abdomen is divided into eight segments (Supplementary Figure 2
B). Each segment on the dorsal side has two neurons, each occupying a hemi-segment.
Each hemi-segment is approximately rectangular with Anterior-Posterior (AP) and Left-
Right (LR) axes. The AP length was measured as the distance between the cell bodies
of the adjacent neurons on the anterior and posterior sides. The LR length was
measured as the distance between cell bodies in adjacent hemi-segments (across the
midline) corrected for the offset of the cell bodies, which are not in the centers of the
cells but displaced away from the midline. The AP and LR segment lengths were
measured for three larvae for all the respective stages.
687

688   ***Arbor skeletonization***
689   Images were maximally projected and individual dendrites manually segmented from
690   their neighbors. The segmented neurons were binarized using custom made algorithm
691   and skeletonized using MATLAB's 'bwmorph'. The identified branch points are then
692   subtracted from the skeleton to generate isolated individual branches.
693

694   ***Branch length measurements***
695   The pixel coordinates of all the branches were smoothed using a spline to calculate the
696   total length of all branches and their average length.
697

698   **Dendrite arbor width**
699   Arbors are approximately rectangular. If the mass of the dendrite skeleton is uniformly
700   distributed in a rectangle, then the widths, D_{AP} and D_{LR}, are:

701
$$D_{\mathrm{AP}} = \sqrt{12}\, R_{\mathrm{g}}^{\mathrm{AP}} \quad \text{and} \quad D_{\mathrm{LR}} = \sqrt{12}\, R_{\mathrm{g}}^{\mathrm{LR}}$$

where R_{g} is the radius of gyration defined by:

$R_{\mathrm{g}} = \sqrt{\frac{1}{N}\sum_{j=1}^{N}\left(r_j - \bar{r}\right)^2}$ where N is the total number of occupied pixels in the skeleton, r_j is
the projection onto the respective axis of the j^{th} occupied pixel and $\bar{r}$ is the mean
projected position of all occupied pixels. R_{g} is the standard deviation of the dendrite
pixels i.e., their spread from the center. We confirmed that the widths defined in this way
were good approximations to the 95% rectangles containing 95% of the skeletal mass.

708
Dilation analysis
Dilation analysis
To study the possible role of dilation in arbor growth, we imaged the same dorsal
neurons (A3, A4, and A5) every 24 hrs. Larvae were mounted and imaged as described
but without the use of anesthetics. Their movement was minimized by imaging at 4 °C
for 2-5 mins. They were then returned to the apple-agar plate in the Darwin Chamber.
The larvae were imaged using 20X and 40X objectives. For image analysis, the same
neurons at 24 & 48 hr, and 48 & 96hr were segmented and aligned using ImageJ to
identify conserved branches. These branches were proximal and non-terminal. The
fractional increase in branches and segment lengths were defined as

718
$$\text{Fractional length change (\%)} = \frac{\text{Final length} - \text{Initial length}}{\text{Initial length}}$$

719
Branching rate and branch angle
Branching rate and branch angle
To determine branching events, time-lapse movies of duration 20-30 minutes were
analyzed manually using ImageJ. A new protrusion of length >0.25 μm was scored as a
new branch. The total branching rate (min^{-1}) was calculated by dividing the total number
of branching events by the total time. The specific branching rate (μm^{-1}min^{-1}) was
calculated as the total branching rate divided by the total branch length. The spatial
distribution of all branching events was plotted using MATLAB with the soma at the
origin ($x = 0, y = 0$). The angle of new branches was measured using the angle tool of
ImageJ (zero angle defined as in the direction of the mother). The angle distribution
graph was plotted using Prism.

730
Fractal Dimension
Fractal Dimension
We used the box-counting method to calculate the fractal dimension. For each box
width, W, we measured the number of boxes, $N(W)$, needed to cover all the occupied
pixels of the skeleton (Figure 7A). $N(W)$ is approximately linear on a log-log plot (Figure
7C) indicative of a power law. The middle 50% of points (between the dashed lines in
Figure 7C) was fit to $N(W) = W^{-D_\text{f}}$ to obtain the fractal dimension D_f.

737
Tip-tracking algorithm
Tip-tracking algorithm
To quantify the dynamical properties of the tips, we developed an in-house algorithm to
track dendritic tips and determine dendrite length-time curves. Time-lapse movies were
stabilized, maximum-projected (ImageJ), and terminal dendrites selected for analysis
based on their separation from neighboring dendrites and the signal-to-background
ratio. Terminal dendrites were selected throughout the arbor. Examples are shown in
Figure 3 A. Extraneous objects were manually deleted.
To track the growing and shrinking tips, the algorithm determined the longitudinal
centerline of the terminal dendrite and the location of its end for each frame. The central
line was computed by fitting Gaussians to the cross-sectional intensity profiles at regular
intervals along the backbone of the dendrite using:

749
$$I(x) = I_0\, e^{\frac{-(x - x_\text{c})^2}{2\sigma^2}} + d$$

(Demchouk et al., 2011) where x is the position along a normal to the dendrite, I_0 is the
peak intensity value, x_c is the center of the Gaussian, σ is the standard deviation and d

is the measured camera offset (100) plus the background fluorescence. To compute the
location of the tip, $(x_\text{tip}, y_\text{tip})$, we fit a 2-dimensional Gaussian function convolved with
an error function using (Demchouk et al., 2011):

$$I(x,y) = I_0 \exp\left[\frac{-\left[(y - y_\text{tip})cos\theta - (x - x_\text{tip})sin\theta\right]^2}{2\sigma_v{}^2}\right] \cdot \text{erfc}\left[\frac{(y - y_\text{tip})cos\theta - (x - x_\text{tip})sin\theta}{\sigma_p}\right] + d$$

where I_0 is the peak intensity, θ is the angular direction of the tip, and σ_v and σ_p are the
standard deviations along the orthogonal and parallel directions. The length of the
dendrite in each frame was determined by fitting the center line and the tip with a cubic
spline. Length-time traces were smoothed with a median filter of size 3 to remove
glitches.
To estimate the precision of our tracking algorithm, we used two different
approaches. In the first, we tracked synthetic images of capped cylindrical tubes of
known length and radius with fluorophores placed randomly on their surfaces (10%
labeling density) and convolved with a point spread function (350 nm FWHM). To
ensure that our algorithm could perform robustly under a wide range of signal-to-
background ratios, we tested the tracking accuracy with decreasing signal to
background ratios. The typical precision was $\ll 1$ pixel (100 nm) even for low signal-to-
background ratios (Ruhnow et al., 2011). (ii) We tracked the position of *in-vivo* dendritic
tips that are in long-term paused states and found that the average standard deviation
of length was ~0.1 µm (< 1 pixel, 0.1615 µm) as shown in Supplementary Figure 1 D.
This accuracy is comparable to and, in some cases, better than available software, such
as, FIESTA (Ruhnow et al., 2011), JFilament (Smith et al., 2010), and Simple Neurite
Tracer (Longair et al., 2011). Using a parallelized method, several hundred tips can be
tracked simultaneously. A caveat of our method is that it can only track filaments that
are reasonably free of extraneous objects, excessive noise, and have no breaks,
discontinuities, or overlapping segments.
777
Calculating velocities and transition rates
Fitting piecewise linear functions to the data is an ill-posed problem (Hansen and
O'Leary, 1993) because a perfect fit can always be achieved with a large enough
number of segments (equal to the number of data points minus one). To circumvent this
problem, we defined a temporal resolution, *T* (in frames), that is necessary to
distinguish a transition event from the noise in the data. Then, the maximum number of
segments in each trajectory was calculated by dividing the total number of data points
(total number of frames) by the temporal resolution. We used simulated data to estimate
the temporal resolution that performs the best. We generated Markovian trajectories
with known and realistic velocity distribution (± 1.5 µm/min) and transition rates (0.5 min^{-1}
^{1}), similar to those shown the Figure 3. We then, added Gaussian white noise of
standard deviation 0.25 µm to the trajectories to mimic the experimental noise. To
analyze the trajectories, we used the following steps as shown in Supplementary Figure
7:
(i) We fit the trajectories with piecewise linear function considering N_data/T as the initial
number of segments (N_seg).
(ii) The velocity distribution (slope of the segments) was fitted to a lognormal-Gaussian-
lognormal distribution:

796
$$v_T = \frac{1 - C_G - C_S}{\sqrt{\{2\pi\sigma_P^2\}}} e^{\left\{-\frac{(v-\mu_P)^2}{2\sigma_P^2}\right\}} + (v > 0)\frac{C_G}{v\sqrt{\{2\pi\sigma_G^2\}}} e^{\left\{\frac{(\log(v)-\mu_G)^2}{2\sigma_G^2}\right\}}$$

797
$$+ (v < 0)\frac{C_S}{|v|\sqrt{\{2\pi\,\sigma_S^2\}}} e^{\left\{-\frac{(\log|v|-\mu_S)^2}{2\sigma_S^2}\right\}}$$

where the C's are the normalization constants, the μ's are the means, the σ's are the
standard deviations and the subscripts P, G, S stand for paused, growth, and shrinkage
states respectively. The first term is a Gaussian and denotes the paused state, whereas
2nd and 3rd terms are log-normal distributions representing the growth and shrinkage
states.
After fitting the above velocity distribution, the intersections between the paused and
growth distribution (I_G) and the paused and shrinkage distribution (I_S) were calculated,
and the segments labeled using these two velocity thresholds.
(iii) Consecutive segments with similar labels were then merged. This process
decreased the number of segments ($N^n{}_{seg}$). The trajectories were then refitted with the
new number of segments. Steps (ii) and (iii) were repeated until $N^n{}_{seg}$ stabilized.
(iv) Finally, the transition rates were calculated by counting the total number of
transitions from one state to another and then dividing that number by the total time
spent in that state. For example, if N_{GP} is the total number of transitions from G to P state
and T_G is the total time spent in the G state, then the transition rate from G to P is $K_{GP} =$
$\frac{N_{GP}}{T_G}$.
Supplementary Figure 7 shows the working protocol along with the validation of
our analysis method. To estimate the optimal frame resolution, we plotted the root-
mean-squared difference between the input and output transition rates as a function of
the frame resolution. Our analysis clearly shows that temporal resolution of 6 frames
generates the best results for rates ~0.5/min and noise ~0.25 μm. We used this frame
resolution for all our tip dynamics data analysis.
820
Average tip velocity & validation of segmentation
The segmentation yields a set of intervals, i, with associated distances, d_i, durations, t_i
and velocities, $v_i = d_i/t_i$. The distribution of velocities is fit to a lognormal-Gaussian-
lognormal model $p(v)$ with parameters $P_G, \mu_G, \sigma_G, P_P, 0, \sigma_P, P_S, \mu_S, \sigma_S$ such that
$\quad 1 = \int p(v)dv = P_G + P_P + P_S$ $\qquad\qquad$ (1)
and
$\quad \int L_{\mu_G,\sigma_G}(v)dv = \int L_{\mu_S,\sigma_S}(v)dv = \int G_{0,\sigma_P}(v)dv = 1$ $\qquad$ (2)
where $L_{\mu_G,\sigma_G}(v)$ and $L_{\mu_S,\sigma_S}(v)$ are the log-normal distributions corresponding to growth
and shrinkage and $G_{0,\sigma_P}(v)$ is the Gaussian distribution of the paused state. Transitions
can only occur between unlike states. This imposes an important structure on the data:
there are two threshold velocities v_+ and v_- such that if the segment velocity $v_i > v_+$
then it is assigned to be a growing segment. Likewise, if $v_i < v_-$ it is a shrinking
segment. The ones in the middle are paused.
The average velocity as:
$\quad v_d \equiv \frac{\sum_i d_i}{\sum_i t_i} = \frac{\sum_i v_i t_i}{\sum_i t_i} = \frac{n\bar{v}\bar{t} + n r_{v,t}\sigma_v\sigma_t}{n\bar{t}} = \bar{v} + r_{v,t}\sigma_v\frac{\sigma_t}{\bar{t}}$ $\qquad$ (3)

where $\bar{v}$ is the mean velocity, $\bar{t}$ is the mean time, and $r_{v,t}$ is the Pearson correlation
coefficient. We used the definition of the Pearson's correlation coefficient
$r_{x,y} = \sigma_{x,y}/\sigma_x\sigma_y$ and cross-correlation $\sigma_{x,y} = \left(\frac{1}{n}\right)\sum_i x_i y_i - \bar{x}\bar{y}$ to calculate equation 3.
The average velocity v_d has two parts: $\bar{v}$, calculated assuming t_i and v_i are
independent of each other, and a cross-correlation term, $r_{v,t}\sigma_v\frac{\sigma_t}{\bar{t}}$. $\bar{v}$ is given by:

$$\bar{v} = P_G \cdot \bar{V}_G + P_s \cdot \bar{V}_S \qquad (4)$$

where $\bar{V}_G$ and $\bar{V}_S$ are the first moments of the lognormal velocity distributions for growth
and paused states, and P_G and P_s can be calculated from the master equation
associated with the transition matrix:

$$\frac{dP_G(t)}{dt} = -(K_{GP} + K_{GS})P_G(t) + K_{PG}\,P(t) + K_{SG}\,P_s(t) \qquad (5)$$

$$\frac{dP_P(t)}{dt} = K_{GP}\,P_G(t) - (K_{PG} + K_{PS})\,P_P(t) + K_{SP}P_s(t) \qquad (6)$$

$$\frac{dP_s(t)}{dt} = K_{GS}\,P_G(t) + K_{PS}\,P_P(t) - (K_{SG} + K_{SP})P_s(t) \qquad (7)$$

848 where $K_{ij,j\neq i}\ i,j = \{G,P,S\}$ are the transition rates. The steady state solution, assuming
849 $\frac{dP_G(t)}{dt} = \frac{dP_P(t)}{dt} = \frac{dP_s(t)}{dt} = 0$ and using equation 1, is:

$$P_G = \frac{K_{PG} * K_{SG} + K_{PG} * K_{SP} + K_{PS} * K_{SG}}{D} \qquad (8)$$

$$P_P = \frac{K_{GP} * K_{SG} + K_{GP} * K_{SP} + K_{GS} * K_{SP}}{D} \qquad (9)$$

$$P_s = \frac{K_{GS} * K_{PG} + K_{GP} * K_{PS} + K_{GS} * K_{PS}}{D} \qquad (10)$$

852 where
853 $D = K_{GS} \cdot K_{PG} + K_{GP} \cdot K_{PS} + K_{GS} \cdot K_{PS} + K_{GP} \cdot K_{SG} + K_{GP} \cdot K_{SP} + K_{GS} \cdot K_{SP} + K_{PG} \cdot$
854 $K_{SG} + K_{PG} \cdot K_{SP} + K_{PS} \cdot K_{SG}.$

855
Finally, the average velocity as:

$$v_d = P_G \cdot \bar{V}_G + P_P.\bar{V}_P + P_s \cdot \bar{V}_S + r_{v,t}\sigma_v\frac{\sigma_t}{\bar{t}} \qquad (11)$$

The average velocity is a key parameter, which controls the growth of the simulated
arbor (Supplementary Figure 5 D & E). The average velocity calculated in this way from
the transition matrix agreed with that calculated directly from the raw traces at all
developmental stages (Table 1) considering the Pearson correlation coefficient $r_{v,t} = 0$
(Supplementary Figure 3 C) This validates of our segmentation scheme.
863
Computational model:
The agent-based two-dimensional computational model of dendritic growth incorporated
the fundamental processes that govern the morphogenesis of Class IV neurons: (i)

branching, (ii) tip dynamics, and (iii) contact-based retraction (non-overlapping). We
started our simulation with randomly oriented 2-4 branches emanating from the origin
(cell body). Each branch is a filament, and points (x, y) are added at 0.1 μm (Δl)
intervals as the branch grows. The simulation is divided into 0.1-minute time steps (Δt).
The details of individual processes are as follows:
Branching:
Assuming the branching is a random process, we visit all the branches randomly and
calculated the branching probability $P_b = 1 - e^{-L_b \omega_b \Delta t}$, where L_b is the length of the
branch and ω_b is the branching rate per unit time per unit length (Figure 4A). Then, a
uniformly distributed unit random number $R(0,1)$ is compared to P_b to spawn a nascent
branch from a random point on the mother branch. The branching angle is chosen
randomly from the measured branch angle distribution which is distributed normally with
a mean ~90° and standard deviation ~26° as shown in Figure 4B and Supplementary
Table 1. Each newly spawned branch is assumed to start in the growing phase with an
initial length of 0.5 μm.
Tip Dynamics:
Each branch with a free end (tip) follows a Markov process (after an initial lag, see
below), transitioning between growing (G), paused (P), and shrinking (S) states with
measured transition rates ($K_{ij, j \neq i}$ $i, j = \{G, P, S\}$;) and velocities ($V_{\{G,P,S\}}$) for free tips
(Figure 3H-J). The transition dynamics is implemented using a standard 'Monte-Carlo'
method. At each time step, the total probability of transition is calculated using $P_i = 1 -$
$e^{-K_{tot} \Delta t}$, where K_{tot} is the sum of the transition rates from one particular state: $K_{tot} =$
$\sum_{j=\{G,P,S\}}^{j \neq i} K_{ij}$. For example, the total transition rate from the growth state is $K_{tot} = K_{GP} +$
K_{GS}. Subsequently, P_i is compared with a uniform random number $R(0,1)$ to implement
the transition. If there is a transition, it happens maintaining the ratio K_{ij}/K_{tot}. After the
transition, the tip is assigned with corresponding state velocity, the magnitude of which
is randomly chosen from their respective velocity distributions (lognormal for growth and
shrinkage and normal for paused) (Figure 3). The growth process is implemented by
adding new points to the existing branch tip at each time step, Δt, as follows:

$$l_G = V_G \Delta t$$

$$x_{k+n} = x_{k+n-1} + \Delta l \, cos\varphi_{k+n-1}; \quad n = 1 \ldots \frac{l_G}{\Delta l}$$

$$y_{k+n} = y_{k+n-1} + \Delta l \, sin\varphi_{k+n-1}; \quad n = 1 \ldots \frac{l_G}{\Delta l}$$

$$\varphi_k^n = \varphi_{k-1}^n + \sqrt{\frac{2\Delta l}{l_p}} \Theta$$

Where, k is the index of the last point in the previous time step and l_p is the persistence
length of the branches (set to 150 μm, Supplementary Table 2). Additionally, the Δl
value of the last point is adjusted if l_G is not an integer multiple of Δl. Θ is a unit
Gaussian variable centered at zero.
While shrinking, points from the branches are removed until the shrinking length
$l_S = V_S \Delta t$ is reached.
Contact-based retraction:

It has been shown by several studies that Dscam molecules play an important role in
the self-avoidance of dendritic tips in Class IV neurons (Matthews et al., 2007; Soba et
al., 2007). To investigate the phenomenon, we measured the dynamics of the dendritic
tips after a collision/contact event has occurred. Interestingly, we observed that the
transition rates are altered after contact as shown in Table 1B and leads to an overall
shrinkage of the dendritic tips. To implement this observation in the model, we assumed
that the contact is achieved whenever a tip comes very close (<0.15 μm, ~average
radius of branches, Supplementary Table 2) to a nearby branch. Further, we used the
altered tip dynamical parameters after a tip makes a contact (post-contact). Because it
is difficult to measure how long the tips remain in their post-contact dynamics, we added
this as a free parameter in the model (Supplementary Table 2). It was chosen to be
longer than the expected lifetimes of post-contact dendrites (10-15 minutes).
Boundary Condition:
Individual Class IV neurons grow within the hemi-segments (Grueber et al., 2002). We
have experimentally measured the segment sizes in the AP and LR directions at
different developmental stages as shown in the inset to Figure 1. Linearly fits to the
growing region (24 to 96 hr) defined the growth rates of the boundary used in our
model. Because there is self-avoidance interaction between the neighboring neurons,
we used a periodic boundary condition.
Initial condition:
To avoid any ambiguity in the initial timepoint in the simulation, we divided the
simulation process into two halves. We started our simulation at 14 hr with 2-4 randomly
oriented branches of length ~15 μm and allowed them to grow with the 18 hr tip
dynamics data (Table 1 A) and the 24 hr branching rate. When the dendritic tree
reached the measured value of total branch number at 24 hr, we reassigned this time as
24 hr. In this way, we simulated larval morphogenesis.
Initial Lag
There is an unexpected paucity of dendrite deaths at short times. To analyze this, we
calculated the survival probability of the branches in the following way:
Suppose dendritic tips are born and die throughout the observation period T. Divide T
into small equal intervals dt. Let,
$f(0)$ = Number{die in time $[0, dt)$}/Number{alive at time 0}
$f(dt)$ = Number{die in time $[dt, 2dt)$}/Number{alive at time dt}
$f(2dt)$ = Number{die in time $[2dt, 3dt)$}/Number{alive at time $2dt$} …etc.
Then we can write the survival probability as:

942
$$D(t) = \prod_{i=0}^{t/dt} [1 - f(i)]$$

The survival probability of the experimentally observed dendritic tips was measured
manually and then calculated by using the above formula. This is shown by the solid
black line in Supplementary Figure 4 A. The survival curve does not decay exponentially
which led us to conclude that the tips have some initial period of sustained growth which
we termed as initial lag τ_{lag}. To estimate the amount of initial lag we simulated 1000
free tips with 48 hr tip dynamics data (because it is in the middle of the developmental
time) and implemented an initial lag (τ_{lag}) during which the tips did not switch into the

paused or shrinking states ($K_{GP} = K_{GS} = 0; t \leq \tau_{lag}$). We calculated the survival
probability by dividing number of alive branches by the total number of branches. The
survival probability increases with the initial lag τ_{lag} as shown by the dotted lines in
Supplementary Figure 4A. The dark blue is the best fit to the real data (τ_{lag} = 0.3 min,
Supplementary Table 2) and we used this value in our model.
955
Markovian tests of the tip trajectories
The tip dynamics is not a 'Markovian process'. The non-Markovian traits are shown by
the presence of initial lag during nascent branch formation (τ_{lag}) and the change of
dynamics after contact (Supplementary Table 2). However, the dynamics is a first-order
process as shown by the single exponential decay of phase durations as shown in
Supplementary Figure 3 D-F which points towards the fact that the dynamics don't have
any long-term memory. To confirm this, we calculated the state-state auto-correlation
function (data not shown). The average autocorrelation function quickly becomes
uncorrelated showing the absence of any long-term memory in the states.
965
966   **Key model parameters**
967
The agent-based model contains 67 parameters: 4 for the velocity distribution and 6 for
the transition rates (times 4 for 24, 48 and 96 hr and post-collision, Table 1 and
Supplementary Tables 1 -2), 3 for the exponential fit to branching rates (24, 36, 48, 72
and 96 hr, Supplementary Table 1), 8 from Supplementary Table 2, 2 for the initial
condition and 4 for the boundary condition. It is instructive to ask which parameters are
key for determining dendrite morphology. We therefore performed a sensitivity analysis.
The boundary was omitted to monitor the unconstrained arbor growth rate.
Branching increases arbor growth and density (Supplementary Figure 5 B).
When the branching frequency was increased 2-fold, with all other parameters equal,
the cell width increased ~20%, leading to a ~50% increase in cell area. The number of
branches increased 3-fold, the total branch length increased 2-fold and the average
branch length decreased 30%, with a small increase in the fractal dimension.
Decreasing branching frequency had the opposite effects.
The stochasticity of tip growth (its variability) increases arbor growth without
changing density (Supplementary Figure 5 B). When all the transition rates were
decreased 2-fold, which does not change in the average tip velocity but makes tips grow
more stochastically, the arbor width grew 40% faster, leading to a doubling of arbor
area. The branch number and total branch length also doubled, so the mean branch
length remained unchanged, as did the fractal dimension. Increasing the transition
rates, which decreases the stochasticity, had the opposite effect.
Increasing the average tip velocity increases arbor growth with little effect on
density (Supplementary Figure 5 C). When the average tip velocity was increased 2-fold
by increasing the tip growth velocity, the arbor width grew 50% faster, with a 2.2-fold
increase in area. There was a corresponding increase in branch points and total branch
length, and so no effect on mean branch length, though there was a small increase in
fractal dimension. Decreasing the average tip velocity 2-fold had the opposite effect.
When the tip velocity was set to zero (Supplementary Figure 5 C), the arbor still grew,
but only at half the rate of the control: despite the 4-fold decrease in area, there was

little effect on density, mean branch length, though the fractal dimension decreased.
When the tip velocity was made negative (-0.02 μm/min), the arbor failed to grow and
was more open, with an increase in mean branch length and a decrease in fractal
dimension.
The branching angle is also a key parameter (data not shown). Decreasing the
branching angle to 45° (i.e., more distal) increased the growth rate. Increasing the
branching angle to 135° has the opposite effect. We previously noted that several
parameters have little effect, including the persistence length, the post-contact time.
Furthermore, the interaction length, soma radius, and initial length are not expected to
have dramatic effects.
In summary, branching, a positive average tip velocity, and tip stochasticity all
drive arbor growth. Arbors can even grow when the average tip velocity is zero.
Branching increases the density of the arbor, whereas tip velocity (and its variability)
has little effect on density. The branching angle strongly affects arbor growth and the
prolonged growth after birth also contributes to the burst of branch formation at early
times (Figure 6D).
1012

Supplementary Figures

1016 **Supplementary Figure 1. Tracking dendrite tips**. **A** Example of a fluorescently labeled terminal
1017 dendrite. **B** The center of the dendrite was located by fitting the cross-sections (blue line in B) to a
1018 Gaussian. The precision is approximately 0.1 µm. **C** The position of the tip of the dendrite was calculated
1019 by fitting the end intensity profile (red box in A) to a 2D function corresponding to a Gaussian in the
1020 perpendicular direction and an error function the parallel direction (see Methods) **D** Montage of simulated
1021 images of cylindrical tubes (6 µm blue, 8 µm red) that are fluorescently labeled with 10% labeling density
1022 on the surface with signal-to-background ratio (SBR), defined as the mean signal divided by the standard
1023 deviation of background noise, varying from 33 (left) to 9 (right). The pixel size is 100 nm. The measured
1024 length distribution is shown in the bottom panel (200 independently generated images for each SBR). **E**
1025 The standard deviation of the measured length is plotted against the SBR. The accuracy is high even for
1026 the lowest SBRs. **F** The lengths of several live-imaged dendrites that were in their paused state as a
1027 function of time. **G** The standard deviation of the tracked lengths is ~1 pixel (108 nm). Examples of
1028 tracked dendritic length as a function of time: **G** 24 hr. **I** 48 hr. and **K** 96 hr. The green, orange, and red
1029 lines denote examples of growing, paused, and shrinking states. The tips tend to spend more time in the
1030 paused state over developmental time. **H**, **J**, and **L** show the statistics of the piecewise-linear fitting.

Supplementary Figure 2. A Control showing that imaging does not perturb growth. The growth of cells is assessed by measuring the rate of increase of their overall radius (calculated as √(area/π)) while imaging. Lines connect cells at the beginning and end of imaging. This shows that the imaging conditions do not retard growth. **B** Definitions of the anterior-posterior (**AP**) and left-right (**LR**) axes. A2, A3, A4, and A5 corresponds to the dorsal abdominal segments. The white dashed line is the dorsal midline and defines the AP axis. This larva is ~24 hr after egg lay.

1041

Supplementary Figure 3: Correlation between state velocities and lifetimes and state durations: A
The correlation between shrinking velocities (V_S) and shrinking lifetimes (T_S) shows a significant
correlation with Pearson's correlation coefficient r=0.064. **B** Similarly, a significant correlation is observed
between growth velocities (V_G) and growth lifetimes (T_G). **C** Pearson's correlation coefficient (r) between
state velocities (V_i) and lifetimes (T_i) for 18-20 hr data. The value of r is small (0.017) and there is no
significant correlation between V_i and T_i. The distribution durations for the shrinking (S), paused (P), and
growing (G) states at 24 hr (**A-F**), 48 hr (**G-I**), and 96 hr (**J-L**) plotted using semi-log axes. The
distributions are very close to exponentials (dotted lines) expected if switching among the states is first
order. The slope of the dotted lines is the inverse of the lifetimes spent in the states: for example, at 24 hr,
the sum of the transition rates from growing state is ($K_{GP} + K_{GS} = 0.696 + 0.509 = 1.205$ min⁻¹ (Table 1 A),
close to the slope of the lifetime distribution of 1.38 min⁻¹ (F).

1053

Supplementary Figure 4: Evidence for persistent growth after birth. A We manually measured the
time between branch initiation and branch death at different stages of the larva (24, 48,72, and 96 hr AEL,
3 movies for each stage). From these data, we calculated the survival probability by dividing the number
of alive branches by the total number of branches using the formula described in the Methods. The black
line is the average survival probability of the real dendrite tips. The survival probability does not start
decaying exponentially as one might expect if it were a Poisson process. Rather, it shows some initial lag.
This observation led us to believe that branch initiation is not a simple Poisson process. To estimate the
initial lag period, we simulated 1000 branches with initial length 0.5 μm and implemented a lag time (τ_{lag})
by preventing the tips to switch into the paused or shrinkage state ($K_{GP} = K_{GS} = 0; t \leq \tau_{lag}$). A branch is
deleted in the simulation when its length is <0.1 μm. The survival probability increases with the initial lag τ
as shown by the dotted lines. The dark blue is the best fit to the real data (τ_{lag} = 0.3 min). **B** An example
of branch retraction and regrowth from the same point. White arrows correspond to the region where a
branch exists and then disappear and branched again. **C** An example of a branch that collided with
another branch, retracted, and then followed the previous path during regrowth (white arrows). Red star
corresponds to shrinkage, yellow star corresponds to the birth of a branch, blue star corresponds to
collision event, and green star shows tip growth.
1070

1071

Supplementary Figure 5: **Sensitivity of morphology to branching and growth parameters**: **A** Control
with parameters from Table 1 and Supplementary Tables 1-2 without any boundary restriction. The black
and blue lines represent LR and AP sizes respectively. The solid lines represent the simulated segment
sizes over development. The simulation shows that initially (24-48 hr) the neurons grow faster than the
real segment and then grow with a constant rate equal to the segment growth rate (~0.06 μm/min) until 96
hr. The segment widths saturate after 96 hr even without a boundary. **B** Branching rate was doubled
(green) and halved (magenta) compared to the control, keeping all other parameters unchanged. All arbor
properties are normalized by the respective unconstrained controls. Fold change is plotted against time
for (ii) arbor size, (iii) branch number, (iv) branch length, (v) mean branch length, and (vi) fractal
dimension. **C** All transition rates were doubled (green) and halved (magenta): this leads to a decrease
and increase in the variability of growth. **D** The mean tip velocity (drift) was increased (green) and
decreases (magenta) 2-fold. **E** The average (drift) velocity was set to zero (green) and a negative value (-
0.02 /min, magenta). Shaded regions represent standard error of mean.

1085

1086

1087

1088

1089 **Supplementary Figure 6**: **Branch length and radial orientation distributions**: **A-E** Branch length
1090 distribution over different developmental stages for real and simulated arbors. **F-J** Radial orientation of
1091 branches over developmental time for real and simulated arbors. The branches are preferentially oriented
1092 in the radial direction. This preference is due to contact-based retraction. The dotted curves show
1093 diminished radial preference when branches are paused after contact in the simulation.

1094
1095

1096
1097
Supplementary Figure 7: Validation of trajectory analysis method. We simulated 200 Markov
trajectories with realistic input parameters shown in **A** and **B** and then added Gaussian white noise on the
individual points of the trajectories. We used the transition rate as 0.5 /min because of the observed fact
that the individual states last ~1 minute. **C** The flowchart of the trajectory analysis method. **D** We varied
the Frame resolution to find the optimal resolution. The root-mean-squared error between the input and
output transition rate matrix is plotted as a function of frame resolution. Frame resolution of 6 provides the
best result and we chose this value for all our analyses. **E &F** The output velocity distribution and
transition rate matrix using frame resolution 6. Our method of analysis produced a faithful reproduction of
the input parameters.
1107

Supplementary Tables

Supplementary Table 1: Branching rates

Age (hr)	Total branching rate[a] (min^{-1}) (Mean ± SD)	Linear branching rate[b] (μm^{-1}·min^{-1}) (Mean ± SD)	Branching angle[c] (°) (Mean ± SD)	Numbers (rates, angles)
18 (E)	4.26 ± 0.59	0.0109 ± 0.0015	85 ± 25	$n = 9, 5$
24 (L1)	7.59 ± 1.52	0.0095 ± 0.0017	88 ± 26	$n = 9, 7$
36 (L1)	5.77 ± 2.67	0.0031 ± 0.0009	85 ± 25	$n = 6, 7$
48 (L2)	4.86 ± 1.59	0.0019 ± 0.0007	86 ± 26	$n = 6, 4$
72 (L2)	8.31 ± 2.37	0.0011 ± 0.0004	90 ± 25	$n = 6, 5$
96 (L3)	11.12 ± 2.52	0.0011 ± 0.0006	91 ± 25	$n = 6, 4$

[a]Over the entire dendrite arbor
[b]Per total dendrite length
[c]Angle between dendrites is zero in the distal direction of the mother.
Standard deviation (SD).

Supplementary Table 2: Model parameters

Name	Description	Value
l_{initial}	Initial length of nascent branch	0.50 μm
$l_{\text{interaction}}$	Length scale of contact	0.15 μm
$\tau_{\text{post-contact}}$	Post-contact dynamics duration	15 min
τ_{lag}	Initial lag of nascent branch	0.3 min
l_{P}	Persistence length	150 μm
R_{soma}	Radius of soma	10 μm
μ_b^{θ}	Mean branching angle	$\pi/2$
σ_b^{θ}	Standard dev. of branching angle	$\pi/7$

Mutation	Reference	Morphology	Features of mutants	Simulated neuron	Parameters
Non-uniform branching					
Dynein intermediate light chain (Dlic)	(Arthur et al., 2015; Satoh et al., 2008; Zheng et al., 2008)	Fig 2C (Arthur et al., 2015)	Defective axonal and dendritic morphology, downsizing of the overall arbor, shift of branching activity to the proximal area within the arbor. Neurons fail to tile.		Non uniform branching, primary branch has basal branching rate.
Lis-1	(Liu et al., 2000; Satoh et al., 2008; Zheng et al., 2008)	Fig 1C (Zheng et al., 2008)	Reduced dendritic arborization in flies (similar to dlic) and mice. Neurons fails to tile.		
Change in tip dynamics					
katanin	(Stewart et al., 2012)	Fig 3B (Stewart et al., 2012)	Decreased dendritic branch number, length, and arbor coverage in class IV neurons		Terminal branches spend less time in the paused state and more time in the retracting phase.
Self-avoidance					
Trc	(Emoto et al., 2004)	Fig 1D (Emoto et al., 2004)	Excessive terminal branching and fail to avoid homologous dendritic branches, resulting in significant overlap of the dendritic fields.		Branches are allowed to cross 10 % of the time.
Dscam	(Hughes et al., 2007; Matthews et al., 2007; Soba et al., 2007)	Fig 1F (Soba et al., 2007)	Branch curling, crossing over, reduced self-avoidance, dendrite bundling. No effect on tiling.		

1121

1122 **Supplementary movies**

1123

1124

1125 **Supplementary movie 1. Time-lapse movie of a growing neuron**: Time lapse movie of an embryonic
1126 neuron at 17.5hr AEL was acquired using a spinning disk confocal microscope. The movie was full-frame
1127 (2048x 2048 pixels) and a complete stack of images (7um) was produced every 5 mins interval.

1128

1129

1130 **Supplementary movie 2. Tip growth and branching**

1131

1132

1133 **Supplementary movie 3. Self-avoidance and shrinkage**

1134

1135

1136 **Supplementary movie 4. Self-avoidance and growth**

1137

1138

1139 **Supplementary movie 5. Tip pause**

1140

1141

Supplementary movie 6-10. Tip growth, bending, and branching. Dendrite lengthening is likely due to
the addition of materials at the dendrite tip. The green stars show dendrite tip lengthening, the yellow star
is birth of new branch, and the red star is a shrinkage event. White arrows point to the bending of growing
tips. In example two, where the branch disappears, a sharp bend smoothens over time. In examples 3,4,
and 5, the white arrows correspond to structural features such as branches and bends that remain fixed
during growth and shortening.

References

Arthur, A.L., Yang, S.Z., Abellaneda, A.M., and Wildonger, J. (2015). Dendrite arborization requires the dynein cofactor NudE. J Cell Sci *128*, 2191–2201.

Ascoli, G.A., and Krichmar, J.L. (2000). L-neuron: A modeling tool for the efficient generation and parsimonious description of dendritic morphology. Neurocomputing *32–33*, 1003–1011.

Baltruschat, L., Tavosanis, G., and Cuntz, H. (2020). A developmental stretch-and-fill process that optimises dendritic wiring. Biorxiv 2020.07.07.191064.

Basak, R., Sutradhar, S., and Howard, J. (2021). Focal Laser Stimulation of Fly Nociceptors Activates Distinct Axonal and Dendritic Ca2+ Signals. Biophys J.

Cajal, S.R. (1995). Histology of the nervous system of man and vertebrates. History of Neuroscience (Oxford Univ Press, New York).

Cajal, S.R. y (1906). The Structure and Connexions of Neurons. Nobel Lecture.

Castro, A.F., Baltruschat, L., Stürner, T., Bahrami, A., Jedlicka, P., Tavosanis, G., and Cuntz, H. (2020). Achieving functional neuronal dendrite structure through sequential stochastic growth and retraction. Elife *9*, e60920.

Cuntz, H., Forstner, F., Borst, A., and Häusser, M. (2010). One Rule to Grow Them All: A General Theory of Neuronal Branching and Its Practical Application. Plos Comput Biol *6*, e1000877.

Decker, F., Oriola, D., Dalton, B., and Brugués, J. (2018). Autocatalytic microtubule nucleation determines the size and mass of Xenopus laevis egg extract spindles. ELife *7*, 6656.

DeFelipe, J., López-Cruz, P.L., Benavides-Piccione, R., Bielza, C., Larrañaga, P., Anderson, S., Burkhalter, A., Cauli, B., Fairén, A., Feldmeyer, D., et al. (2013). New insights into the classification and nomenclature of cortical GABAergic interneurons. Nature Reviews Neuroscience *14*, 202–216.

Delandre, C., Amikura, R., and Moore, A.W. (2016). Microtubule nucleation and organization in dendrites. Cell Cycle *15*, 1–8.

Demchouk, A.O., Gardner, M.K., and Odde, D.J. (2011). Microtubule Tip Tracking and Tip Structures at the Nanometer Scale Using Digital Fluorescence Microscopy. Cell Mol Bioeng *4*, 192–204.

Denk, W., Briggman, K.L., and Helmstaedter, M. (2012). Structural neurobiology: missing link to a mechanistic understanding of neural computation. Nat Rev Neurosci *13*, 351–358.

Dent, E.W., Gupton, S.L., and Gertler, F.B. (2011). The growth cone cytoskeleton in axon outgrowth and guidance. Csh Perspect Biol *3*, a001800–a001800.

D'Errico, J. (2021). SLM-Shape Language Modeling (Mathworks).

Dogterom, M., and Leibler, S. (1993). Physical aspects of the growth and regulation of microtubule structures. Physical Review Letters *70*, 1347–1350.

Dong, X., Shen, K., and Bülow, H.E. (2015). Intrinsic and Extrinsic Mechanisms of Dendritic Morphogenesis. Annu Rev Physiol *77*, 1–30.

Emoto, K., He, Y., Ye, B., Grueber, W.B., Adler, P.N., Jan, L.Y., and Jan, Y.-N. (2004). Control of Dendritic Branching and Tiling by the Tricornered-Kinase/Furry Signaling Pathway in Drosophila Sensory Neurons. Cell *119*, 245–256.

Falconer, K. (1990). Fractal Geometry: Mathematical Foundations and Applications (Chichester ; New York : Wiley).

Fornito, A., Zalesky, A., and Breakspear, M. (2015). The connectomics of brain disorders. Nat Rev Neurosci *16*, 159–172.

Forrest, M.P., Parnell, E., and Penzes, P. (2018). Dendritic structural plasticity and neuropsychiatric disease. Nat Rev Neurosci *19*, 215–234.

Ganguly, S., Trottier, O., Liang, X., Bowne-Anderson, H., and Howard, J. (2016). Morphology of Fly Larval Class IV Dendrites Accords with a Random Branching and Contact Based Branch Deletion Model.

Gao, F.-B., Brenman, J.E., Jan, L.Y., and Jan, Y.N. (1999). Genes regulating dendritic outgrowth, branching, and routing in Drosophila. Gene Dev *13*, 2549–2561.

Grueber, W.B., Jan, L.Y., and Jan, Y.N. (2002). Tiling of the Drosophila epidermis by multidendritic sensory neurons. Dev Camb Engl *129*, 2867–2878.

Grueber, W.B., Ye, B., Moore, A.W., Jan, L.Y., and Jan, Y.N. (2003). Dendrites of distinct classes of Drosophila sensory neurons show different capacities for homotypic repulsion. Current Biology *13*, 618–626.

Han, C., Wang, D., Soba, P., Zhu, S., Lin, X., Jan, L.Y., and Jan, Y.N. (2012). Integrins regulate repulsion-mediated dendritic patterning of drosophila sensory neurons by restricting dendrites in a 2D space. Neuron *73*, 64–78.

Hannezo, E., Scheele, C.L.G.J., Moad, M., Drogo, N., Heer, R., Sampogna, R.V.,
Rheenen, J. van, and Simons, B.D. (2017). A Unifying Theory of Branching
Morphogenesis. Cell *171*, 242-255.e27.

Hansen, P.C., and O'Leary, D.P. (1993). The Use of the L-Curve in the Regularization
of Discrete Ill-Posed Problems. Siam J Sci Comput *14*, 1487–1503.

Hughes, M.E., Bortnick, R., Tsubouchi, A., Bäumer, P., Kondo, M., Uemura, T., and
Schmucker, D. (2007). Homophilic Dscam Interactions Control Complex Dendrite
Morphogenesis. Neuron *54*, 417–427.

Jan, Y.-N., and Jan, L.Y. (2010). Branching out: mechanisms of dendritic arborization.
Nat Rev Neurosci *11*, 316–328.

Jinushi-Nakao, S., Arvind, R., Amikura, R., Kinameri, E., Liu, A.W., and Moore, A.W.
(2007). Knot/Collier and Cut Control Different Aspects of Dendrite Cytoskeleton and
Synergize to Define Final Arbor Shape. Neuron *56*, 963–978.

Kanari, L., Dłotko, P., Scolamiero, M., Levi, R., Shillcock, J., Hess, K., and Markram, H.
(2018). A Topological Representation of Branching Neuronal Morphologies.
Neuroinformatics *16*, 3–13.

Kandel, E.R., Markram, H., Matthews, P.M., Yuste, R., and Koch, C. (2013).
Neuroscience thinks big (and collaboratively). Nat Rev Neurosci *14*, 659–664.

Kapitein, L.C., and Hoogenraad, C.C. (2015). Building the Neuronal Microtubule
Cytoskeleton. Neuron *87*, 492–506.

Kelliher, M.T., Yue, Y., Ng, A., Kamiyama, D., Huang, B., Verhey, K.J., and Wildonger,
J. (2018). Autoinhibition of kinesin-1 is essential to the dendrite-specific localization of
Golgi outposts. J Cell Biol *217*, 2531–2547.

Kim, M.E., Shrestha, B.R., Blazeski, R., Mason, C.A., and Grueber, W.B. (2012).
Integrins Establish Dendrite-Substrate Relationships that Promote Dendritic Self-
Avoidance and Patterning in Drosophila Sensory Neurons. Neuron *73*, 79–91.

Kirschner, M., and Mitchison, T. (1986). Beyond self-assembly: from microtubules to
morphogenesis. *45*, 329–342.

Koleske, A.J. (2013). Molecular mechanisms of dendrite stability. Nat Rev Neurosci *14*,
536–550.

Komarova, Y.A., Vorobjev, I.A., and Borisy, G.G. (2002). Life cycle of MTs: persistent
growth in the cell interior, asymmetric transition frequencies and effects of the cell
boundary. Journal of Cell Science *115*, 3527–3539.

Kulkarni, V.A., and Firestein, B.L. (2012). The dendritic tree and brain disorders. Mol
Cell Neurosci *50*, 10–20.

Lee, A., Li, W., Xu, K., Bogert, B.A., Su, K., and Gao, F.-B. (2003). Control of dendritic
development by the Drosophila fragile X-related gene involves the small GTPase Rac1.
Development *130*, 5543–5552.

Lefebvre, J.L., Sanes, J.R., and Kay, J.N. (2013). Development of Dendritic Form and
Function. Annu Rev Cell Dev Bi *31*, 1–37.

Lefevre, J.G., Short, K.M., Lamberton, T.O., Michos, O., Graf, D., Smyth, I.M., and
Hamilton, N.A. (2017). Branching morphogenesis in the developing kidney is governed
by rules that pattern the ureteric tree. Development (Cambridge, England) *144*, 4377–
4385.

Li, W., and Gao, F.-B. (2003). Actin Filament-Stabilizing Protein Tropomyosin Regulates
the Size of Dendritic Fields. J Neurosci *23*, 6171–6175.

Liang, X., Kokes, M., Fetter, R.D., Sallee, M.D., Moore, A.W., Feldman, J.L., and Shen,
K. (2020). Growth cone-localized microtubule organizing center establishes microtubule
orientation in dendrites. Elife *9*, e56547.

Liao, M., Liang, X., and Howard, J. (2021). The narrowing of dendrite branches across
nodes follows a well-defined scaling law. Proc National Acad Sci *118*, e2022395118.

Liu, Z., Steward, R., and Luo, L. (2000). Drosophila Lis1 is required for neuroblast
proliferation, dendritic elaboration and axonal transport. Nat Cell Biol *2*, 776–783.

Longair, M.H., Baker, D.A., and Armstrong, J.D. (2011). Simple Neurite Tracer: open
source software for reconstruction, visualization and analysis of neuronal processes.
Bioinformatics *27*, 2453–2454.

Lowery, L.A., and Vactor, D.V. (2009). The trip of the tip: understanding the growth cone
machinery. Nat Rev Mol Cell Bio *10*, 332–343.

Lu, P., and Werb, Z. (2008). Patterning mechanisms of branched organs. Sci New York
N Y *322*, 1506–1509.

Luczak, A. (2006). Spatial embedding of neuronal trees modeled by diffusive growth. J
Neurosci Meth *157*, 132–141.

Matthews, B.J., Kim, M.E., Flanagan, J.J., Hattori, D., Clemens, J.C., Zipursky, S.L.,
and Grueber, W.B. (2007). Dendrite Self-Avoidance Is Controlled by Dscam. Cell *129*,
593–604.

Metzger, R.J., Klein, O.D., Martin, G.R., and Krasnow, M.A. (2008). The branching
programme of mouse lung development. Nature *453*, 745–750.

Mitchison, T., and Kirschner, M. (1984). Dynamic instability of microtubule growth.
Nature *312*, 237–242.

Moughamian, A.J., and Holzbaur, E.L.F. (2017). Dyneins. In Cytoplasmic Dynein
Dysfunction and Neurodegenerative Disease, S.M. King, ed. (Academic Press), pp.
286–315.

Nanda, S., Das, R., Bhattacharjee, S., Cox, D.N., and Ascoli, G.A. (2018).
Morphological determinants of dendritic arborization neurons in Drosophila larva. Brain
Struct Funct *223*, 1107–1120.

Nithianandam, V., and Chien, C.-T. (2018). Actin blobs prefigure dendrite branching
sites. J Cell Biol *217*, 3731–3746.

Ori-McKenney, K.M., Jan, L.Y., and Jan, Y.-N. (2012). Golgi Outposts Shape Dendrite
Morphology by Functioning as Sites of Acentrosomal Microtubule Nucleation in
Neurons. Neuron *76*, 921–930.

Palavalli, A., Tizón-Escamilla, N., Rupprecht, J.-F., and Lecuit, T. (2021). Deterministic
and Stochastic Rules of Branching Govern Dendrite Morphogenesis of Sensory
Neurons. Curr Biol *31*, 459-472.e4.

Parrish, J.Z., Xu, P., Kim, C.C., Jan, L.Y., and Jan, Y.N. (2009). The microRNA bantam
Functions in Epithelial Cells to Regulate Scaling Growth of Dendrite Arbors in
Drosophila Sensory Neurons. Neuron *63*, 788–802.

Petry, S., Groen, A.C., Ishihara, K., Mitchison, T.J., and Vale, R.D. (2013). Branching
microtubule nucleation in Xenopus egg extracts mediated by augmin and TPX2. *152*,
768–777.

Pollard, T.D., and Borisy, G.G. (2003). Cellular Motility Driven by Assembly and
Disassembly of Actin Filaments. Cell *112*, 453–465.

Robertson, J.L., Tsubouchi, A., and Tracey, W.D. (2013). Larval Defense against Attack
from Parasitoid Wasps Requires Nociceptive Neurons. Plos One *8*, e78704.

Ruhnow, F., Zwicker, D., and Diez, S. (2011). Tracking single particles and elongated
filaments with nanometer precision. Biophys J *100*, 2820–2828.

Sánchez-Soriano, N., Tear, G., Whitington, P., and Prokop, A. (2007). Drosophila as a
genetic and cellular model for studies on axonal growth. Neural Dev *2*, 9–9.

Satoh, D., Sato, D., Tsuyama, T., Saito, M., Ohkura, H., Rolls, M.M., Ishikawa, F., and Uemura, T. (2008). Spatial control of branching within dendritic arbors by dynein-dependent transport of Rab5-endosomes. Nat Cell Biol *10*, 1164–1171.

Scott, E.K., Reuter, J.E., and Luo, L. (2003). Small GTPase Cdc42 is required for multiple aspects of dendritic morphogenesis. J Neurosci Official J Soc Neurosci *23*, 3118–3123.

Sears, J.C., and Broihier, H.T. (2016). FoxO regulates microtubule dynamics and polarity to promote dendrite branching in Drosophila sensory neurons. Dev Biol *418*, 40–54.

Seung, S. (2012). Connectome: How the Brain's Wiring Makes Us Who We Are.

Sherwood, N.T., Sun, Q., Xue, M., Zhang, B., and Zinn, K. (2004). Drosophila Spastin Regulates Synaptic Microtubule Networks and Is Required for Normal Motor Function. Plos Biol *2*, e429.

Short, K.M., Combes, A.N., Lefevre, J., Ju, A.L., Georgas, K.M., Lamberton, T., Cairncross, O., Rumballe, B.A., McMahon, A.P., Hamilton, N.A., et al. (2014). Global Quantification of Tissue Dynamics in the Developing Mouse Kidney. Dev Cell *29*, 188–202.

Singhania, A., and Grueber, W.B. (2014). Development of the embryonic and larval peripheral nervous system of Drosophila. Wiley Interdiscip Rev Dev Biology *3*, 193–210.

Smith, M.B., Li, H., Shen, T., Huang, X., Yusuf, E., and Vavylonis, D. (2010). Segmentation and tracking of cytoskeletal filaments using open active contours. Cytoskelet Hoboken N J *67*, 693–705.

Soba, P., Zhu, S., Emoto, K., Younger, S., Yang, S.-J., Yu, H.-H., Lee, T., Jan, L.Y., and Jan, Y.-N. (2007). Drosophila Sensory Neurons Require Dscam for Dendritic Self-Avoidance and Proper Dendritic Field Organization. Neuron *54*, 403–416.

Stewart, A., Tsubouchi, A., Rolls, M.M., Tracey, W.D., and Sherwood, N.T. (2012). Katanin p60-like1 Promotes Microtubule Growth and Terminal Dendrite Stability in the Larval Class IV Sensory Neurons of Drosophila. J Neurosci *32*, 11631–11642.

Stoeckli, E.T. (2018). Understanding axon guidance: are we nearly there yet? Development *145*, dev151415.

Stürner, T., Tatarnikova, A., Mueller, J., Schaffran, B., Cuntz, H., Zhang, Y., Nemethova, M., Bogdan, S., Small, V., and Tavosanis, G. (2019). Transient localization of the Arp2/3 complex initiates neuronal dendrite branching in vivo. Development *146*, dev.171397.

Sugimura, K., Shimono, K., Uemura, T., and Mochizuki, A. (2007). Self-organizing
Mechanism for Development of Space-filling Neuronal Dendrites. Plos Comput Biol *3*,
e212.

Tessier-Lavigne, M., and Goodman, C.S. (1996). The Molecular Biology of Axon
Guidance. Science *274*, 1123–1133.

Toner, J., and Tu, Y. (1995). Long-Range Order in a Two-Dimensional Dynamical XY
Model: How Birds Fly Together. Physical Review Letters *75*, 4326–4329.

Vicsek, T., Czirók, A., Ben-Jacob, E., Cohen, I., and Shochet, O. (1995). Novel Type of
Phase Transition in a System of Self-Driven Particles. Phys Rev Lett *75*, 1226–1229.

Weiner, A.T., Lanz, M.C., Goetschius, D.J., Hancock, W.O., and Rolls, M.M. (2016).
Kinesin-2 and Apc function at dendrite branch points to resolve microtubule collisions.
Cytoskeleton (Hoboken, NJ) *73*, 35–44.

Winding, M., Kelliher, M.T., Lu, W., Wildonger, J., and Gelfand, V.I. (2016). Role of
kinesin-1-based microtubule sliding in Drosophila nervous system development. P Natl
Acad Sci Usa *113*, E4985-94.

Yalgin, C., Ebrahimi, S., Delandre, C., Yoong, L.F., Akimoto, S., Tran, H., Amikura, R.,
Spokony, R., Torben-Nielsen, B., White, K.P., et al. (2015). Centrosomin represses
dendrite branching by orienting microtubule nucleation. Nat Neurosci *18*, 1437–1445.

Yang, W.-K., and Chien, C.-T. (2019). Beyond being innervated: the epidermis actively
shapes sensory dendritic patterning. Open Biol *9*, 180257.

Zheng, Y., Wildonger, J., Ye, B., Zhang, Y., Kita, A., Younger, S.H., Zimmerman, S.,
Jan, L.Y., and Jan, Y.N. (2008). Dynein is required for polarized dendritic transport and
uniform microtubule orientation in axons. Nat Cell Biol *10*, 1172–1180.

Zheng, Z., Lauritzen, J.S., Perlman, E., Robinson, C.G., Nichols, M., Milkie, D., Torrens,
O., Price, J., Fisher, C.B., Sharifi, N., et al. (2018). A Complete Electron Microscopy
Volume of the Brain of Adult Drosophila melanogaster. Cell *174*, 730-743.e22.

1399