Title: The secreted neuronal signal Spock1 regulates the blood-brain barrier

Authors: Natasha M. O'Brown[1*], Nikit B. Patel[1], Ursula Hartmann[2], Allon M. Klein[1], Chenghua Gu[3], Sean G. Megason[1*]

Affiliations:
[1] Department of Systems Biology, Harvard Medical School, 200 Longwood Ave, Boston, MA 02115, U.S.A.
[2] University of Cologne, Joseph-Stelzmann-Str. 52 50931 Cologne, Germany
[3] Howard Hughes Medical Institute, Department of Neurobiology, Harvard Medical School, 220 Longwood Ave, Boston, MA 02115, USA

* denotes corresponding author

Correspondence:

Natasha M. O'Brown (Natasha_obrown@hms.harvard.edu)
Department of Systems Biology, Harvard Medical School
Boston, MA 02115, U.S.A.

Sean G. Megason (megason@hms.harvard.edu)
Department of Systems Biology, Harvard Medical School
Boston, MA 02115, U.S.A.
(Phone) 617-432-7441

ABSTRACT

The blood-brain barrier (BBB) is a unique set of properties of the brain vasculature which

severely restricts its permeability to proteins and small molecules. Classic chick-quail

chimera studies showed that these properties are not intrinsic to the brain vasculature but

rather are induced by surrounding neural tissue. Here we identify Spock1 as a candidate

neuronal signal for regulating BBB permeability in zebrafish and mice. Mosaic genetic

analysis shows that neuronally-expressed Spock1 is cell non-autonomously required for

a functional BBB. Leakage in *spock1* mutants is associated with altered extracellular

matrix (ECM), increased endothelial transcytosis, and altered pericyte-endothelial

interactions. Furthermore, a single dose of recombinant SPOCK1 partially restores BBB

function in *spock1* mutants by quenching gelatinase activity and restoring vascular

expression of BBB genes including *mcamb*. These analyses support a model in which

neuronally secreted Spock1 induces BBB properties by altering the ECM, thereby

regulating pericyte-endothelial interactions and downstream vascular gene expression.

One-Sentence Summary: Spock1 is a signal secreted by neurons that induces barrier properties

in the brain vasculature

INTRODUCTION

The blood-brain barrier (BBB) maintains a tightly controlled homeostatic environment in the brain

that is required for proper neural function. BBB breakdown has been implicated in multiple

neurodegenerative diseases including Alzheimer's, Parkinson's, and Huntington's Diseases

(Sweeney et al., 2018). Conversely, the BBB also serves as an obstacle for effective drug delivery

to the brain. Therefore there is great interest in generating a better understanding of how to

therapeutically regulate its permeability, both to restore BBB function in neurodegeneration and

to transiently open it for improved chemotherapeutic access for brain tumors. The BBB is a

specialized property of the brain vasculature, which is composed of a thin, continuous layer of

non-fenestrated endothelial cells with uniquely restrictive properties. Brain endothelial cells create

the barrier via two primary cellular mechanisms: 1) specialized tight junction complexes that block

the transit of small water-soluble molecules between cells and 2) reduced levels of vesicular

trafficking or transcytosis to restrict transit through endothelial cells (Reese and Karnovsky, 1967).

At a molecular level, functional barrier endothelial cells are differentiated from peripheral

endothelial cells by high expression of the tight junction molecule Claudin-5 (Campbell et al.,

2008) and the fatty acid transporter Mfsd2a (Andreone et al., 2017; Ben-Zvi et al., 2014), and an

absence of the fenestra and vesicle associated protein Plvap (Hallmann et al., 1995). Expression

of substrate-specific influx and efflux transporters, which dynamically regulate the intake of

necessary nutrients and the removal of metabolic waste products (Sanchez-Covarrubias et al.,

2014; Umans et al., 2017), further enhances the selectivity of the BBB.

The restrictive properties of the BBB are not intrinsic to brain endothelial cells, but are induced

and maintained by signals in the brain microenvironment during embryonic development (Stewart

and Wiley, 1981). When avascularized quail neural tissue was transplanted into embryonic

chicken gut cavities, the blood vessels ingressing from the non-barrier gut tissue obtained

functional BBB properties, specifically both tight junctions and reduced levels of vesicles, resulting

in Trypan blue tracer confinement within the blood vessels (Stewart and Wiley, 1981). Conversely,

when avascularized quail somite tissue was transplanted into embryonic chick brain, the blood

vessels ingressing from the neural tissue lost functional BBB properties, leaking tracer into the

quail graft due to increased endothelial transcytosis and loss of tight junction function. These data

show that signals from the neural microenvironment to the vasculature control barrier function.

Furthermore, these microenvironmental signals are also required to actively maintain barrier

properties throughout life (Guérit et al., 2021; Heithoff et al., 2021; Lyck et al., 2009; Urich et al.,

2012).

Mural cells called pericytes share the endothelial basement membrane and comigrate with

endothelial cells during neovascularization of the developing brain. This close interaction between

pericytes and endothelial cells is required for both the initiation and continued maintenance of

barrier properties (Armulik et al., 2010; Bell et al., 2010; Daneman et al., 2010). Astrocytes, glial

cells found exclusively in the central nervous system, completely ensheath the vasculature with

their endfeet in a polarized fashion (Abbott et al., 2006). Astrocytes arise late in embryonic

development, after neurogenesis is complete, with the vast majority of gliogenesis occurring

postnatally in rodents. Mammalian astrocytes are potent inducers of barrier properties in vitro and

in vivo, and are required for BBB maintenance (Guérit et al., 2021; Heithoff et al., 2021; Janzer

and Raff, 1987). While the necessity of these two cell types has been known for decades,

canonical Wnt signaling arising from both neuronal and astrocytic sources is the only

microenvironmental signal known to induce and maintain BBB function (Benz et al., 2019;

Daneman et al., 2009; Guérit et al., 2021; Liebner et al., 2008; Martin et al., 2022; Stenman et al.,

2008; Vanhollebeke et al., 2015; Wang et al., 2012, 2018). However in addition to its role in

regulating barrier properties, Wnt signaling also plays an important role in tip cell specification
and angiogenesis in the developing brain (Stenman et al., 2008; Vanhollebeke et al., 2015; Wang
et al., 2012). Furthermore, mouse endothelial cells are Wnt-responsive as early as E9.5 (Liebner
et al., 2008), preceding the acquisition of functional barrier properties (Daneman et al., 2009),
suggesting that other signaling may be acting in conjunction with or in addition to Wnt signaling
for the induction of BBB properties.

To further examine the molecular determinants of BBB development, we turned to the optically
transparent zebrafish system, which allows for imaging of the intact BBB in toto throughout
zebrafish development. Zebrafish brain endothelial cells express many of the same molecular
markers as mammalian brain endothelial cells, including Glut1, Cldn5, ZO-1, and Mfsd2a
(Guemez-Gamboa et al., 2015; Jeong et al., 2008; O'Brown et al., 2019; Umans et al., 2017;
Vanhollebeke et al., 2015; Xie et al., 2010). We previously characterized the molecular and
subcellular mechanisms of functional BBB development in zebrafish, and determined that the
zebrafish BBB becomes functionally mature by 5 days post fertilization (dpf) due to the
suppression of transcytosis rather than through the acquisition of tight junction function, which
was observed as early as 3 dpf (O'Brown et al., 2019). During these studies, we serendipitously
discovered a recessive viable mutant with profound forebrain and midbrain barrier leakage which
we identify here to be the neuronally produced and secreted proteoglycan Spock1. Using an array
of perturbations and imaging techniques, we uncover the range of Spock1 signaling to the
vasculature, and identify that mechanistically Spock1 signaling acts through control of the
pericyte-endothelial extracellular matrix (ECM) to control endothelial transcytosis.

RESULTS
Identification and mapping of a spontaneous leaky mutant to *spock1*
We found a spontaneous recessive mutant that leaks both an injected 1 kDa Alexa Fluor (AF)
405 NHS Ester dye and a transgenic 80 kDa serum protein DBP-EGFP (Xie et al., 2010) into the
forebrain and midbrain as early as 3 days post fertilization (dpf) with no improvement in BBB
function throughout larval development (Figure 1A-E) or adulthood (Figure 1G). This increased
BBB permeability in homozygous mutants occurs in the absence of hemorrhage or vascular
patterning defects (Figure S1), or any reduction in viability or fertility. Time lapse imaging of
injected 10 kDa Dextran revealed a steady accumulation of Dextran in the brain parenchyma of
leaky mutants over the course of an hour (Figure S2), closely resembling the dynamics of Dextran

leakage observed in *mfsd2aa* mutants with increased endothelial transcytosis (O'Brown et al.,

2019).

To identify the mutation responsible for this leakage phenotype, we performed linkage mapping

on 5 dpf mutant and wild type siblings using bulk segregant RNAseq (Miller et al., 2013) and

identified a single peak at chr14:2205271-3513919 (GRCz11; Figures 1H and S3). Of the 14

genes within the linkage region (Figure S3), 8 were expressed at 5 dpf. Two of these genes were

differentially expressed, *csf1ra* and *gstp2* (Table S1)*,* and one had several mutations that

segregated with the leakage phenotype, *spock1.* To test whether loss of any of these genes

conferred the increased BBB permeability, we assessed tracer leakage in mosaic crispants

(zebrafish larvae injected at the 1-cell stage with Cas9 protein and gene-specific sgRNAs) at 5

dpf and observed no BBB defects in 42 *gstp2* or 35 *csf1ra* crispants (data not shown). However,

when we assessed BBB function in *spock1* crispants, we observed a strong leaky phenotype in

26% of larvae (26/100 injected fish; Figure S3) and moderate leakage in 53% (53/100 injected

fish; Figure S3) as well as a few that displayed BBB leakage restricted to one midbrain

hemisphere (4/100 injected fish; Figure S3). Leakage in *spock1* crispants corresponded with a

loss of *spock1* expression (Figure S3). *Spock1* encodes a secreted protein of unclear function

named for its conserved protein domains: <u>SP</u>ARC (<u>O</u>steonectin), <u>C</u>wcv And <u>K</u>azal Like Domains

Proteoglycan <u>1</u> protein (also known as Testican-1). Spock1 has three predicted domains: 1) a

Kazal-type serine protease inhibitor, 2) an extracellular SPARC calcium-binding region, and 3) a

thyroglobulin type-1 repeat region, in addition to being decorated by both chondroitin sulfate (CS)

and heparan sulfate (HS) glycosaminoglycan (GAG) chains (Figure 1I) (Bonnet et al., 1996;

Edgell et al., 2004). Further sequencing of *spock1* confirmed a few point mutations in the SPARC

calcium-binding domain in the leaky fish (Figure 1I), including a T241A missense mutation and

two silent mutations at Q249 and G250, hereafter referred to as s*pock1^{hm41/hm41}* mutants.

To validate that loss of Spock1 function causes the leakage phenotype in the spontaneous

mutants, we generated a stable null allele designated *spock1^{hm43}* using CRISPR mutagenesis

with two sgRNAs to completely delete the 5' UTR and start codon of *spock1* (Figure 1I). As

expected based on the 550 bp genomic deletion, *spock1^{hm43/hm43}* fish do not express *spock1*

(Figure S4). Functional tracer leakage assays using 10 kDa Dextran revealed the same increased

BBB permeability specifically in the forebrain and the midbrain observed in the spontaneous

s*pock1^{hm41/hm41}* mutants (Figure 1J-L). This increased BBB permeability was accompanied with

increased expression of the pathological marker of leaky endothelial cells *plvapb:EGFP* (Umans

et al., 2017) compared to the negligible levels observed in wild type siblings with a functional BBB

(Figure 1M-O). Lastly, we crossed both lines to test for genetic complementation, and observed

the leaky phenotype in *spock1^hm41/hm43^* compound heterozygotes (Figure 1P-S), further indicating

that the initial spontaneous mutant acts via loss of Spock1 function.

Neuronal Spock1 signals to the vasculature within a short range

Given the necessity of Spock1 in determining BBB function, we next wanted to assess where

spock1 was expressed during BBB development. Using HCR fluorescent *in situ* hybridization, we

determined that *spock1* mRNA is expressed throughout the developing central nervous system

(CNS), including the entire brain, retina and spinal cord (Figures 2A and S4), and that this

expression is unaltered in the spontaneous *spock1^hm41/hm41^* mutants (Figure S4). Closer

examination of the fluorescent signal revealed co-localization with the neuronal marker *elavl3* and

not the vascular marker *kdrl* (Figure 2D-G), similar to its predominantly neural expression in the

developing mouse CNS (Charbonnier et al., 2000). Prior single-cell RNA sequencing (scRNA-

seq) data (Raj et al., 2020) and our own also showed expression of *spock1* primarily in neurons

and never in vascular endothelial cells (Figure S6). Taken together, these data indicate that we

have identified a neuronal signal, Spock1, that plays a role in establishing endothelial BBB

properties during development without altering vascular patterning.

To determine the range of the Spock1 signal, we performed cell transplantation experiments to

make genetically mosaic zebrafish embryos and assessed tracer leakage in relationship to the

closest transplanted cells at 5 dpf (Figure 2H). When we transplanted labeled wild type cells into

wild type host embryos, we observed a negligible level of tracer accumulation in the brain

parenchyma regardless of the proximity of the nearest donor cell (Figure 2I and 2L). In contrast,

when we transplanted *spock1^hm41/hm41^* mutant cells into mutant host embryos, we observed high

levels of tracer accumulation in the parenchyma regardless of the proximity of the nearest donor

cell (Figure 2J and 2L). Strikingly, when we transplanted wild type cells into *spock1* mutant hosts,

we observed a complete rescue of the mutant leakage if the wild type donor cell was within 10

µm of a blood vessel and no rescue if the donor cell was more than 20 µm away (Figure 2K-N),

indicating that Spock1 can functionally signal at a distance of 10 to 20 µm. Importantly, wild type

cells could rescue leakage in mutant hosts when they differentiate as neurons but not endothelial

cells (Figure 2M).

Mutant leakage arises from increased transcytosis and altered pericyte-endothelial interactions

To assess the subcellular mechanism of increased BBB permeability in $spock1^{hm41/hm41}$ mutants, we injected electron-dense NHS-gold nanoparticles (5 nm) into circulation, followed by transmission electron microscopy (TEM) imaging in 7 dpf $spock1$ mutant and wildtype siblings. These TEM leakage assays revealed no alterations in the cellular composition of the neurovascular unit (NVU) in $spock1$ mutants, with endothelial cells and pericytes sharing a basement membrane that is surrounded by neurons and glia (Figure 3A and 3B). A modest impairment in tight junction function was observed in $spock1$ mutants (52/59 functional tight junctions; Figure 3D) compared to wild type siblings (57/57; Figure 3C). However, $spock1$ mutants had a significant increase in both small (<100 nm diameter) non-clathrin coated vesicles (Figure 3D and 3G) and large (>200 nm diameter) vesicles (Figure 3F and 3H) suggesting that the leakage results primarily from an increase in vesicular trafficking across endothelial cells. In addition to the increase in total large vesicular abundance in $spock1$ mutants, we also observed several of these large vesicles fused to the abluminal membrane in multiple larvae with a few examples of the gold nanoparticles visibly spilling into the endothelial basement membrane (Figure 3F). Although pericyte coverage of the endothelium was unaltered (Figure S1), $spock1$ mutants displayed overall thinner pericyte-endothelial basement membranes with several pericytes having long stretches of apparently direct contact on endothelial cells (Figure 3I-K), corresponding with observations by live confocal microscopy (Figure S1). These data suggest that loss of Spock1 function alters the critical pericyte extracellular interactions with endothelial cells, resulting in loss of BBB properties as observed in pericyte-deficient mice and zebrafish (Armulik et al., 2011; Bell et al., 2010; Daneman et al., 2010; Wang et al., 2013).

One of the proposed roles of Spock1 is to regulate matrix metalloproteinase (MMP) activation, specifically gelatinases MMP-2 and MMP-9 (Du et al., 2020; Edgell et al., 2004; Nakada et al., 2001; Váncza et al., 2022; Ye et al., 2020). To assess whether $spock1^{hm41/hm41}$ mutants had altered gelatinase activity that corresponded to observed changes in vascular cell interactions, we performed $in vivo$ gelatin zymography by intracranial injections of highly quenched FITC-gelatin. Proteolytic cleavage of FITC-gelatin by either MMP-2 or 9 releases bright FITC peptides resulting in a non-reversible increase in fluorescence. This increase in fluorescence upon digestion is proportional to the proteolytic activity of the MMPs, allowing for the detection of gelatinase activity $in vivo$ (Garcia-Alloza et al., 2009; Underly et al., 2017). Using this assay, we were able to see that in wild type 5 dpf fish, the majority of the signal appears strongly in the cerebral spinal fluid

(CSF) and more subtly in many pericytes (Figure 3L). While *spock1[hm41/hm41]* mutants also

displayed strong gelatinase activity in the CSF, this was accompanied by a regional increase in

gelatinase activity specifically in the forebrain and midbrain (Figure 3M and 3N), the same regions

that were also prone to tracer accumulation (Figure 1). These data suggest that Spock1 helps

induce barrier function by keeping gelatinase activity in check.

Spock1 plays a conserved role in inducing barrier function in mammals

To determine if Spock1 plays a conserved role in determining vertebrate BBB properties, we

assessed BBB function in embryonic day 15.5 (E15.5) *Spock1[-/-]* mice (Röll et al., 2006) (Figure

4A), a stage when the cortex BBB is fully functional (Ben-Zvi et al., 2014). While wild type siblings

confined both 550 Da Sulfo-NHS-Biotin (Figure 4B) and 10 kDa Dextran (Figure 4D) within the

brain vasculature, *Spock1[-/-]* mice leaked both tracers into the cortex parenchyma (Figure 4C-G).

This increased BBB permeability was also associated with increased expression of the PLVAP

(Figure 4H-L), as seen in the zebrafish mutants (Figure 1N). These data indicate the Spock1 plays

a conserved role in inducing barrier properties during embryonic development. However, when

we repeated these leakage assays in adult mice, we observed full recovery in BBB function in

Spock1[-/-] mice (Figure S5). This contrasts with adult zebrafish *spock1[hm41/hm41]* mutants, which have

a defective BBB (Figure 1), indicating species-specific differences in the adult BBB.

Vascular cells are the only altered cell types in *spock1* mutant brains

To determine how Spock1, a neuronally produced and secreted proteoglycan, signals to and

regulates brain endothelial cell BBB properties, we turned to scRNA-seq of dissected 5 dpf

spock1[hm41/hm41] mutant and wild type brains, allowing us to globally assess cell type specific

molecular changes occurring in the mutant brains. With Leiden clustering of the scRNA-seq data

we defined cell clusters containing neurons, glial, and vascular cell types (Figure S6). We

performed differential gene expression (DGE) analysis for each cluster and did not observe any

changes in the neuronal or glial populations, but did observe significant changes in the vascular

cluster (Table S2). When we subclustered the vascular population, we were able to resolve

endothelial cells, pericytes and vascular smooth muscle cells (vSMCs; Figure 5A). Due to the

low vascular cell numbers present in the scRNA-seq data, we prioritized candidate genes for

subsequent validation by their fold-change in gene expression rather than statistical significance.

These analyses suggested molecular changes in the mutant endothelial cells indicative of a leaky

phenotype, with increased expression of *plvapb* and decreased expression of the tight junction

protein *cldn5b* (Figure 5B, Table S3). Interestingly, *spock1* mutants had decreased expression of

the melanoma cell adhesion molecule *mcamb* (also known as CD146) in both endothelial cells

and pericytes (Figure 5B and 5C). This decreased expression in the vasculature was validated

by HCR FISH, with minimal expression in mutant midbrain pericytes but normal levels in hindbrain

pericytes, which maintain BBB function in *spock1$^{hm41/hm41}$* mutants (Figure 5D-G). CD146 has

previously been shown to be required for BBB integrity, as *CD146$^{-/-}$* mice exhibit BBB breakdown

due to decreased pericyte coverage and downstream loss of endothelial expression of Cldn5

(Chen et al., 2017).

Exogenous SPOCK1 restores barrier function

To test the ability of Spock1 to rescue the mutant leakage phenotype, we injected recombinant

human SPOCK1 (rSPOCK1) protein directly into the brain of *spock1$^{hm41/hm41}$* mutant larvae at 5

dpf and assessed brain permeability at 6 dpf. While control animals injected with PBS maintained

high levels of brain permeability (Figure 6A), larvae that received at least 2.3 ng of rSPOCK1 per

mg fish body weight showed a 50% reduction in brain permeability following a single dose (Figure

6B and 6C). These results further confirm that SPOCK1 is able to act cell non-autonomously to

rescue the mutant BBB leakage, as these injections were targeted broadly to the neural tissue

rather than the endothelial cells, and demonstrate functional conservation of Spock1 from human

to zebrafish.

To determine whether this restoration of barrier function corresponds to Spock1's regulation of

the MMP activity, we repeated the intracranial injections of 11.5 ng rSPOCK1/mg of fish in

spock1$^{hm41/hm41}$ mutants at 4 dpf and assessed barrier function and gelatinase activity at 5 dpf.

While 5 dpf control injected *spock1$^{hm41/hm41}$* mutants continued to display high levels of BBB

permeability and gelatinase activity, rSPOCK1 reduced mutant leakage of 10 kDa Dextran about

50% (Figure 6E and 6F), just as it did when we injected rSPOCK1 at 5 dpf and assessed BBB

function at 6 dpf (Figure 6C). This restoration of BBB function was accompanied by a decrease

in gelatinase activity (Figure 6H and 6I). Finally, to determine whether Spock1 lies upstream of

vascular expression of the cell adhesion molecule *mcamb,* we assessed *mcamb* expression in

the mutant midbrain vasculature at 5 dpf with or without the addition of rSPOCK1. While

spock1$^{hm41/hm41}$ mutants injected with PBS displayed negligible levels of *mcamb* expression in the

midbrain (Figure 6J), mutants injected with rSPOCK1 had a 50% increase in *mcamb* expression

in the vasculature (Figure 6K and 6L), with a noticeable increase in *mcamb*+ pericytes.

Interestingly this 50% increase in *mcamb* expression directly correlates with the 50% reduction in

leakage and gelatinase activity following rSPOCK1 injections (Figure 6F and 6I).

Discussion

Together these data show that Spock1 is a secreted, neuronally expressed extracellular signal
that regulates BBB permeability in zebrafish and mouse without altering vascular patterning
during development (Figures 1 and 4). Mechanistically, this work suggests a model whereby
Spock1 regulates the brain extracellular environment via its inhibition of gelatinase activity,
thereby providing the appropriate chemical or mechanical signals for vascular pericytes and
endothelial cells to communicate properly with one another. This critical pericyte-endothelial
interaction thereby prompts a BBB transcriptional program, including increased expression of tight
junction (*cldn5b)* and cell adhesion (*mcamb*) molecules, decreased expression of leaky
endothelial genes (*plvapb*), and reduced transcytosis. Thus, loss of Spock1 function contributes
to increased BBB permeability by disruption of the complex extracellular scaffold that surrounds
and supports the vasculature during development. Together this work reveals how a novel signal
from the brain microenvironment can regulate the vasculature to give rise to the special properties
of the BBB and provides new targets for its therapeutic modulation.

Despite the ubiquitous expression of *spock1* throughout the CNS, *spock1* loss of function
zebrafish mutants display regional leakage in the forebrain and midbrain and a completely intact
hindbrain barrier. Spock1 is not alone in having region-specific regulation of barrier properties
despite pan-CNS expression. A recent study demonstrated that the pericyte-produced ECM
molecule vitronectin is required for functional retinal and cerebellar barriers despite its expression
by all CNS pericytes (Ayloo et al., 2022). Even mutants in canonical Wnt signaling have varying
levels or regionalized angiogenic or barrier defects despite wide-spread sources of Wnt signals
and receptors throughout the brain (Cho et al., 2017; Junge et al., 2009; Wang et al., 2012, 2018;
Xu et al., 2004; Zhou and Nathans, 2014; Zhou et al., 2014). Interestingly, the regions with the
highest susceptibility to single gene mutations in canonical Wnt signaling are the retina and the
cerebellum, while the cortex can maintain a functional BBB even with mutations in several genes
in the Wnt pathway (Wang et al., 2018). These data suggest that different regions of the brain are
either molecularly or cellularly (or both) distinct, perhaps in nuanced ways that are often masked
by current detection methods. For example, while the neurovascular unit of endothelial cells and
pericytes in close contact with astrocytic endfeet appears uniform throughout the brain by
immunostaining and EM, various scRNA-seq datasets have now shown that astrocytes are highly

regionally distinct at a molecular level (Herrero-Navarro et al., 2021; Ohlig et al., 2021; Zhu et al.,
2018) raising the possibility that they may respond to different signals. Pericytes in the brain,
unlike in the rest of the body, arise from two discrete origins, mesodermal and neural crest (Ando
et al., 2016; Trost et al., 2016). In zebrafish, the two origins have discrete domains, with the
hindbrain being exclusively derived from mesodermal origins and the midbrain and forebrain
being predominantly of neural crest origin (Ando et al., 2016) which interestingly matches the
leakage boundaries in *spock1* mutants.
338
Here we demonstrate that Spock1 regulates brain endothelial cell identity to initiate the barrier
program during development, both at molecular and subcellular resolution. However, this does
not preclude the fact that other brain barriers may be similarly affected by loss of this signal. In
addition to the BBB, the neural tissue is also protected by the blood-cerebrospinal fluid (CSF)
barrier, which is created by the choroid plexus epithelial cells lining the brain ventricles (Wolburg
and Paulus, 2010), and the arachnoid barrier, which is created by the epithelial cells beneath the
dura that completely enwrap the brain (Abbott, 2004). These various brain barriers mature on
different developmental timelines, with the blood-CSF barrier becoming impermeable by 4 dpf
(Henson et al., 2014), then the BBB by 5 dpf (O'Brown et al., 2019), and the arachnoid barrier
maturing after 9 dpf (Jeong et al., 2008) in zebrafish. While the focus of our study has been the
BBB, it is possible that these additional brain barriers also play a role in the observed leakage of
various tracers into the brain in *spock1* mutants.
351
The N-terminal region of Spock1 has previously been shown to regulate matrix metalloproteinase
(MMP) activity, specifically gelatinases MMP-2 and 9 (Du et al., 2020). We show here that leaky
spock1$^{hm41/hm41}$ mutants have increased gelatinase activity in the midbrain occurring in the same
place as the observed decrease in the endothelial-pericyte basement membrane (Figure 3 and
S1), suggesting that this excessive MMP activity may be responsible for the breakdown of the
basement membrane. Although all the cells in the neurovascular unit secrete MMP-2 and 9,
pericytes have recently been shown to produce MMP-9 in response to ischemia and TGF-β1
exposure, leading to increased barrier breakdown and neuronal damage (Takahashi et al., 2014;
Underly et al., 2017). For this reason, many have proposed gelatinases as a therapeutic target
for several neurological diseases that are accompanied with barrier breakdown including stroke,
multiple sclerosis, Alzheimer's diseases, and cerebral hemorrhage (Avolio et al., 2003; Kook et
al., 2013; Rosell et al., 2008; Underly et al., 2017). Given the ability of a single dose of rSPOCK1
to mitigate much of the leakage in the mutants and restore proper vascular expression of BBB

genes, we propose that Spock1 may serve as a new therapeutic target to reduce gelatinase
activity in these neurological diseases and help restore barrier function.
367

Acknowledgments

We thank members of the Megason laboratory for data discussion and comments on the
manuscript; Dr. Zach O'Brown for discussions and comments on the manuscript; Ignas Mazelis
for single cell demultiplexing software; the HMS Single Cell Core for single cell library preparation;
Dr. Bela Anand-Apte (Cleveland Clinic) for providing the transgenic *l-fabp:DBP-EGFP* fish line
(Xie et al., 2010) and Dr. Leonard Zon for providing the transgenic Tg(*kdrl:HRAS-mCherry*) line;
and the HMS Electron Microscopy Core Facility, with special thanks to Louise Trakimas for all of
her assistance in preparing the TEM samples. This work was supported by the Damon Runyon
Cancer Foundation (N.M.O.), NIH R01HD096755 (S.G.M), an Allen Distinguished Investigator
Award (C.G.), NIH R35NS116820 (C.G), and partially supported by Faculty Scholar grant from
the Howard Hughes Medical Institute (C.G.). C.G. is an investigator of the Howard Hughes
Medical Institute.

Author Contributions
N.M.O., S.G.M. and C.G. conceived the project and designed experiments. N.M.O. performed all
experiments and analyzed most data with the exception of the scRNA-seq which was performed
in collaboration with and analyzed by N.B.P. and A.M.K. U.H. provided the *Spock1*[-/-] mice. N.M.O.,
S.G.M. and C.G wrote the manuscript.

References and Notes:

Abbott, N.J. (2004). Evidence for bulk flow of brain interstitial fluid: significance for physiology and pathology. Neurochem Int *45*, 545–552. https://doi.org/10.1016/j.neuint.2003.11.006.

Abbott, N.J., Rönnbäck, L., and Hansson, E. (2006). Astrocyte–endothelial interactions at the blood–brain barrier. Nature Reviews. Neuroscience *7*, 41–53. https://doi.org/10.1038/nrn1824.

Ando, K., Fukuhara, S., Izumi, N., Nakajima, H., Fukui, H., Kelsh, R.N., and Mochizuki, N. (2016). Clarification of mural cell coverage of vascular endothelial cells by live imaging of zebrafish. Development (Cambridge, England) *143*, 1328–1339. https://doi.org/10.1242/dev.132654.

Andreone, B.J., Chow, B.W., Tata, A., Lacoste, B., Ben-Zvi, A., Bullock, K., Deik, A.A., Ginty, D.D., Clish, C.B., and Gu, C. (2017). Blood-Brain Barrier Permeability Is Regulated by Lipid Transport-Dependent Suppression of Caveolae-Mediated Transcytosis. Neuron *94*, 581-594.e5. https://doi.org/10.1016/j.neuron.2017.03.043.

Armulik, A., Genové, G., Mäe, M., Nisancioglu, M.H., Wallgard, E., Niaudet, C., He, L., Norlin, J., Lindblom, P., Strittmatter, K., et al. (2010). Pericytes regulate the blood-brain barrier. Nature *468*, 557–561. https://doi.org/10.1038/nature09522.

Armulik, A., Genové, G., and Betsholtz, C. (2011). Pericytes: Developmental, Physiological, and Pathological Perspectives, Problems, and Promises. Developmental Cell *21*, 193–215. https://doi.org/10.1016/j.devcel.2011.07.001.

Avolio, C., Ruggieri, M., Giuliani, F., Liuzzi, G.M., Leante, R., Riccio, P., Livrea, P., and Trojano, M. (2003). Serum MMP-2 and MMP-9 are elevated in different multiple sclerosis subtypes. J Neuroimmunol *136*, 46–53. https://doi.org/10.1016/s0165-5728(03)00006-7.

Ayloo, S., Lazo, C.G., Sun, S., Zhang, W., Cui, B., and Gu, C. (2022). Pericyte-to-endothelial cell signaling via vitronectin-integrin regulates blood-CNS barrier. Neuron https://doi.org/10.1016/j.neuron.2022.02.017.

Bell, R.D., Winkler, E.A., Sagare, A.P., Singh, I., LaRue, B., Deane, R., and Zlokovic, B.V. (2010). Pericytes control key neurovascular functions and neuronal phenotype in the adult brain and during brain aging. Neuron *68*, 409–427. https://doi.org/10.1016/j.neuron.2010.09.043.

Benz, F., Wichitnaowarat, V., Lehmann, M., Germano, R.F., Mihova, D., Macas, J., Adams, R.H., Taketo, M.M., Plate, K.-H., Guérit, S., et al. (2019). Low wnt/β-catenin signaling determines leaky vessels in the subfornical organ and affects water homeostasis in mice. Elife *8*, e43818. https://doi.org/10.7554/elife.43818.

Ben-Zvi, A., Lacoste, B., Kur, E., Andreone, B.J., Mayshar, Y., Yan, H., and Gu, C. (2014). Mfsd2a is critical for the formation and function of the blood-brain barrier. Nature *509*, 507–511. https://doi.org/10.1038/nature13324.

Bonnet, F., Périn, J.P., Charbonnier, F., Camuzat, A., Roussel, G., Nussbaum, J.L., and Alliel, P.M. (1996). Structure and cellular distribution of mouse brain testican. Association with the postsynaptic area of hippocampus pyramidal cells. The Journal of Biological Chemistry *271*, 4373–4380. https://doi.org/10.1074/jbc.271.8.4373.

Bresciani, E., Broadbridge, E., and Liu, P.P. (2018). An efficient dissociation protocol for
generation of single cell suspension from zebrafish embryos and larvae. Methodsx *5*, 1287–
1290. https://doi.org/10.1016/j.mex.2018.10.009.

Campbell, M., Kiang, A.-S., Kenna, P.F., Kerskens, C., Blau, C., O'Dwyer, L., Tivnan, A., Kelly,
J.A., Brankin, B., Farrar, G.-J., et al. (2008). RNAi-mediated reversible opening of the blood-
brain barrier. The Journal of Gene Medicine *10*, 930–947. https://doi.org/10.1002/jgm.1211.

Charbonnier, F., Chanoine, C., Cifuentes-Diaz, C., Gallien, C.L., Rieger, F., Alliel, P.M., and
Périn, J.P. (2000). Expression of the proteoglycan SPOCK during mouse embryo development.
Mechanisms of Development *90*, 317–321. https://doi.org/10.1016/s0925-4773(99)00255-5.

Chen, J., Luo, Y., Hui, H., Cai, T., Huang, H., Yang, F., Feng, J., Zhang, J., and Yan, X. (2017).
CD146 coordinates brain endothelial cell–pericyte communication for blood–brain barrier
development. Proc National Acad Sci *114*, E7622–E7631.
https://doi.org/10.1073/pnas.1710848114.

Chi, N.C., Shaw, R.M., Val, S.D., Kang, G., Jan, L.Y., Black, B.L., and Stainier, D.Y.R. (2008).
Foxn4 directly regulates tbx2b expression and atrioventricular canal formation. Genes &
Development *22*, 734–739. https://doi.org/10.1101/gad.1629408.

Cho, C., Smallwood, P.M., and Nathans, J. (2017). Reck and Gpr124 Are Essential Receptor
Cofactors for Wnt7a/Wnt7b-Specific Signaling in Mammalian CNS Angiogenesis and Blood-
Brain Barrier Regulation. Neuron *95*, 1056-1073.e5.
https://doi.org/10.1016/j.neuron.2017.07.031.

Choi, H.M.T., Schwarzkopf, M., Fornace, M.E., Acharya, A., Artavanis, G., Stegmaier, J.,
Cunha, A., and Pierce, N.A. (2018). Third-generation in situ hybridization chain reaction:
multiplexed, quantitative, sensitive, versatile, robust. Development (Cambridge, England) *145*.
https://doi.org/10.1242/dev.165753.

Daneman, R., Agalliu, D., Zhou, L., Kuhnert, F., Kuo, C.J., and Barres, B.A. (2009). Wnt/beta-
catenin signaling is required for CNS, but not non-CNS, angiogenesis. Proceedings of the
National Academy of Sciences of the United States of America *106*, 641–646.
https://doi.org/10.1073/pnas.0805165106.

Daneman, R., Zhou, L., Kebede, A.A., and Barres, B.A. (2010). Pericytes are required for blood-
brain barrier integrity during embryogenesis. Nature *468*, 562–566.
https://doi.org/10.1038/nature09513.

Du, Z., Lin, Z., Wang, Z., Liu, D., Tian, D., and Xia, L. (2020). SPOCK1 overexpression induced
by platelet-derived growth factor-BB promotes hepatic stellate cell activation and liver fibrosis
through the integrin $\alpha5\beta^2$1/PI3K/Akt signaling pathway. Laboratory Investigation 1–15.
https://doi.org/10.1038/s41374-020-0425-4.

Edgell, C.-J.S., BaSalamah, M.A., and Marr, H.S. (2004). Testican-1: a differentially expressed
proteoglycan with protease inhibiting activities. International Review of Cytology *236*, 101–122.
https://doi.org/10.1016/s0074-7696(04)36003-1.

Garcia-Alloza, M., Prada, C., Lattarulo, C., Fine, S., Borrelli, L.A., Betensky, R., Greenberg, S.M., Frosch, M.P., and Bacskai, B.J. (2009). Matrix metalloproteinase inhibition reduces oxidative stress associated with cerebral amyloid angiopathy in vivo in transgenic mice. J Neurochem *109*, 1636–1647. https://doi.org/10.1111/j.1471-4159.2009.06096.x.

Guemez-Gamboa, A., Nguyen, L.N., Yang, H., Zaki, M.S., Kara, M., Ben-Omran, T., Akizu, N., Rosti, R.O., Rosti, B., Scott, E., et al. (2015). Inactivating mutations in MFSD2A, required for omega-3 fatty acid transport in brain, cause a lethal microcephaly syndrome. Nature Genetics *47*, 809–813. https://doi.org/10.1038/ng.3311.

Guérit, S., Fidan, E., Macas, J., Czupalla, C.J., Figueiredo, R., Vijikumar, A., Yalcin, B.H., Thom, S., Winter, P., Gerhardt, H., et al. (2021). Astrocyte-derived Wnt growth factors are required for endothelial blood-brain barrier maintenance. Prog Neurobiol *199*, 101937. https://doi.org/10.1016/j.pneurobio.2020.101937.

Hallmann, R., Mayer, D.N., Berg, E.L., Broermann, R., and Butcher, E.C. (1995). Novel mouse endothelial cell surface marker is suppressed during differentiation of the blood brain barrier. Dev Dynam *202*, 325–332. https://doi.org/10.1002/aja.1002020402.

Heithoff, B.P., George, K.K., Phares, A.N., Zuidhoek, I.A., Munoz-Ballester, C., and Robel, S. (2021). Astrocytes are necessary for blood–brain barrier maintenance in the adult mouse brain. Glia *69*, 436–472. https://doi.org/10.1002/glia.23908.

Henson, H.E., Parupalli, C., Ju, B., and Taylor, M.R. (2014). Functional and genetic analysis of choroid plexus development in zebrafish. Frontiers in Neuroscience *8*, 364. https://doi.org/10.3389/fnins.2014.00364.

Herrero-Navarro, Á., Puche-Aroca, L., Moreno-Juan, V., Sempere-Ferràndez, A., Espinosa, A., Susín, R., Torres-Masjoan, L., Leyva-Díaz, E., Karow, M., Figueres-Oñate, M., et al. (2021). Astrocytes and neurons share region-specific transcriptional signatures that confer regional identity to neuronal reprogramming. Sci Adv *7*, eabe8978. https://doi.org/10.1126/sciadv.abe8978.

Janzer, R.C., and Raff, M.C. (1987). Astrocytes induce blood-brain barrier properties in endothelial cells. Nature *325*, 253–257. https://doi.org/10.1038/325253a0.

Jeong, J.-Y., Kwon, H.-B., Ahn, J.-C., Kang, D., Kwon, S.-H., Park, J.A., and Kim, K.-W. (2008). Functional and developmental analysis of the blood–brain barrier in zebrafish. Brain Research Bulletin *75*, 619–628. https://doi.org/10.1016/j.brainresbull.2007.10.043.

Junge, H.J., Yang, S., Burton, J.B., Paes, K., Shu, X., French, D.M., Costa, M., Rice, D.S., and Ye, W. (2009). TSPAN12 Regulates Retinal Vascular Development by Promoting Norrin- but Not Wnt-Induced FZD4/β-Catenin Signaling. Cell *139*, 299–311. https://doi.org/10.1016/j.cell.2009.07.048.

Klein, A.M., Mazutis, L., Akartuna, I., Tallapragada, N., Veres, A., Li, V., Peshkin, L., Weitz, D.A., and Kirschner, M.W. (2015). Droplet Barcoding for Single-Cell Transcriptomics Applied to Embryonic Stem Cells. Cell *161*, 1187–1201. https://doi.org/10.1016/j.cell.2015.04.044.

Kook, S.-Y., Hong, H.S., Moon, M., and Mook-Jung, I. (2013). Disruption of blood-brain barrier in Alzheimer disease pathogenesis. Tissue Barriers *1*, e23993. https://doi.org/10.4161/tisb.23993.

Li, B., and Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. BMC Bioinformatics *12*, 323. https://doi.org/10.1186/1471-2105-12-323.

Liebner, S., Corada, M., Bangsow, T., Babbage, J., Taddei, A., Czupalla, C.J., Reis, M., Felici, A., Wolburg, H., Fruttiger, M., et al. (2008). Wnt/β-catenin signaling controls development of the blood–brain barrier. J Cell Biology *183*, 409–417. https://doi.org/10.1083/jcb.200806024.

Lyck, R., Ruderisch, N., Moll, A.G., Steiner, O., Cohen, C.D., Engelhardt, B., Makrides, V., and Verrey, F. (2009). Culture-induced changes in blood-brain barrier transcriptome: implications for amino-acid transporters in vivo. Journal of Cerebral Blood Flow and Metabolism : Official Journal of the International Society of Cerebral Blood Flow and Metabolism *29*, 1491–1502. https://doi.org/10.1038/jcbfm.2009.72.

Martin, M., Vermeiren, S., Bostaille, N., Eubelen, M., Spitzer, D., Vermeersch, M., aci, C.P., Pozuelo, E., Toussay, X., Raman-Nair, J., et al. (2022). Engineered Wnt ligands enable blood-brain barrier repair in neurological disorders. Science *375*. https://doi.org/10.1126/science.abm4459.

McGinnis, C.S., Patterson, D.M., Winkler, J., Conrad, D.N., Hein, M.Y., Srivastava, V., Hu, J.L., Murrow, L.M., Weissman, J.S., Werb, Z., et al. (2019). MULTI-seq: sample multiplexing for single-cell RNA sequencing using lipid-tagged indices. Nat Methods *16*, 619–626. https://doi.org/10.1038/s41592-019-0433-8.

Miller, A.C., Obholzer, N.D., Shah, A.N., Megason, S.G., and Moens, C.B. (2013). RNA-seq-based mapping and candidate identification of mutations from forward genetic screens. Genome Research *23*, 679–686. https://doi.org/10.1101/gr.147322.112.

Nakada, M., Yamada, A., Takino, T., Miyamori, H., Takahashi, T., Yamashita, J., and Sato, H. (2001). Suppression of membrane-type 1 matrix metalloproteinase (MMP)-mediated MMP-2 activation and tumor invasion by testican 3 and its splicing variant gene product, N-Tes. Cancer Research *61*, 8896–8902. .

O'Brown, N.M., Megason, S.G., and Gu, C. (2019). Suppression of transcytosis regulates zebrafish blood-brain barrier function. ELife *8*, 41. https://doi.org/10.7554/elife.47326.

Ohlig, S., Clavreul, S., Thorwirth, M., Simon-Ebert, T., Bocchi, R., Ulbricht, S., Kannayian, N., Rossner, M., Sirko, S., Smialowski, P., et al. (2021). Molecular diversity of diencephalic astrocytes reveals adult astrogenesis regulated by Smad4. Embo J *40*, e107532. https://doi.org/10.15252/embj.2020107532.

Pomreinke, A.P., Soh, G.H., Rogers, K.W., Bergmann, J.K., Bläßle, A.J., and Müller, P. (2017). Dynamics of BMP signaling and distribution during zebrafish dorsal-ventral patterning. Elife *6*, e25861. https://doi.org/10.7554/elife.25861.

Raj, B., Farrell, J.A., Liu, J., Kholtei, J.E., Carte, A.N., Acedo, J.N., Du, L.Y., McKenna, A., Relić,
Đ., Leslie, J.M., et al. (2020). Emergence of Neuronal Diversity during Vertebrate Brain
Development. Neuron 1–24. https://doi.org/10.1016/j.neuron.2020.09.023.

Reese, T.S., and Karnovsky, M.J. (1967). Fine structural localization of a blood-brain barrier to
exogenous peroxidase. J Cell Biology *34*, 207–217. https://doi.org/10.1083/jcb.34.1.207.

Röll, S., Seul, J., Paulsson, M., and Hartmann, U. (2006). Testican-1 is dispensable for mouse
development. Matrix Biology *25*, 373–381. https://doi.org/10.1016/j.matbio.2006.05.004.

Rosell, A., Cuadrado, E., Ortega-Aznar, A., Hernández-Guillamon, M., Lo, E.H., and Montaner,
J. (2008). MMP-9–Positive Neutrophil Infiltration Is Associated to Blood–Brain Barrier
Breakdown and Basal Lamina Type IV Collagen Degradation During Hemorrhagic
Transformation After Human Ischemic Stroke. Stroke *39*, 1121–1126.
https://doi.org/10.1161/strokeaha.107.500868.

Sanchez-Covarrubias, L., Slosky, L.M., Thompson, B.J., Davis, T.P., and Ronaldson, P.T.
(2014). Transporters at CNS barrier sites: obstacles or opportunities for drug delivery? Current
Pharmaceutical Design *20*, 1422–1449. .

Stenman, J.M., Rajagopal, J., Carroll, T.J., Ishibashi, M., McMahon, J., and McMahon, A.P.
(2008). Canonical Wnt Signaling Regulates Organ-Specific Assembly and Differentiation of CNS
Vasculature. Science *322*, 1247–1250. https://doi.org/10.1126/science.1164594.

Stewart, P.A., and Wiley, M.J. (1981). Developing nervous tissue induces formation of blood-
brain barrier characteristics in invading endothelial cells: A study using quail-chick
transplantation chimeras. Dev Biol *84*, 183–192. https://doi.org/10.1016/0012-1606(81)90382-1.

Sweeney, M.D., Sagare, A.P., and Zlokovic, B.V. (2018). Blood–brain barrier breakdown in
Alzheimer disease and other neurodegenerative disorders. Nature Publishing Group *14*, 133–
150. https://doi.org/10.1038/nrneurol.2017.188.

Takahashi, Y., Maki, T., Liang, A.C., Itoh, K., Lok, J., Osumi, N., and Arai, K. (2014). p38 MAP
kinase mediates transforming-growth factor-β1-induced upregulation of matrix
metalloproteinase-9 but not -2 in human brain pericytes. Brain Res *1593*, 1–8.
https://doi.org/10.1016/j.brainres.2014.10.029.

Traag, V.A., Waltman, L., and Eck, N.J. van (2019). From Louvain to Leiden: guaranteeing well-
connected communities. Sci Rep-Uk *9*, 5233. https://doi.org/10.1038/s41598-019-41695-z.

Trost, A., Lange, S., Schroedl, F., Bruckner, D., Motloch, K.A., Bogner, B., Kaser-Eichberger,
A., Strohmaier, C., Runge, C., Aigner, L., et al. (2016). Brain and Retinal Pericytes: Origin,
Function and Role. Frontiers in Cellular Neuroscience *10*, 3171–13.
https://doi.org/10.3389/fncel.2016.00020.

Umans, R.A., Henson, H.E., Mu, F., Parupalli, C., Ju, B., Peters, J.L., Lanham, K.A., Plavicki,
J.S., and Taylor, M.R. (2017). CNS angiogenesis and barriergenesis occur simultaneously.
Developmental Biology *425*, 101–108. https://doi.org/10.1016/j.ydbio.2017.03.017.

Underly, R.G., Levy, M., Hartmann, D.A., Grant, R.I., Watson, A.N., and Shih, A.Y. (2017).
Pericytes as Inducers of Rapid, Matrix Metalloproteinase-9-Dependent Capillary Damage during
Ischemia. The Journal of Neuroscience *37*, 129–140. https://doi.org/10.1523/jneurosci.2891-
16.2016.

Urich, E., Lazic, S.E., Molnos, J., Wells, I., and Freskgård, P.-O. (2012). Transcriptional Profiling
of Human Brain Endothelial Cells Reveals Key Properties Crucial for Predictive In Vitro Blood-
Brain Barrier Models. PloS One *7*, e38149-16. https://doi.org/10.1371/journal.pone.0038149.

Váncza, L., Tátrai, P., Reszegi, A., Baghy, K., and Kovalszky, I. (2022). SPOCK1 with
unexpected function. The start of a new career. Am J Physiol-Cell Ph
https://doi.org/10.1152/ajpcell.00033.2022.

Vanhollebeke, B., Stone, O.A., Bostaille, N., Cho, C., Zhou, Y., Maquet, E., Gauquier, A.,
Cabochette, P., Fukuhara, S., Mochizuki, N., et al. (2015). Tip cell-specific requirement for an
atypical Gpr124- and Reck-dependent Wnt/β-catenin pathway during brain angiogenesis. ELife
4. https://doi.org/10.7554/elife.06489.

Wagner, D.E., Weinreb, C., Collins, Z.M., Briggs, J.A., Megason, S.G., and Klein, A.M. (2018).
Single-cell mapping of gene expression landscapes and lineage in the zebrafish embryo.
Science *108*, eaar4362-13. https://doi.org/10.1126/science.aar4362.

Wang, Y., Rattner, A., Zhou, Y., Williams, J., Smallwood, P.M., and Nathans, J. (2012).
Norrin/Frizzled4 signaling in retinal vascular development and blood brain barrier plasticity. Cell
151, 1332–1344. https://doi.org/10.1016/j.cell.2012.10.042.

Wang, Y., Pan, L., Moens, C.B., and Appel, B. (2013). Notch3 establishes brain vascular
integrity by regulating pericyte number. Development (Cambridge, England) *141*, 307–317.
https://doi.org/10.1242/dev.096107.

Wang, Y., Cho, C., Williams, J., Smallwood, P.M., Zhang, C., Junge, H.J., and Nathans, J.
(2018). Interplay of the Norrin and Wnt7a/Wnt7b signaling systems in blood–brain barrier and
blood–retina barrier development and maintenance. Proceedings of the National Academy of
Sciences of the United States of America *115*, E11827–E11836.
https://doi.org/10.1073/pnas.1813217115.

Weinreb, C., Wolock, S., and Klein, A.M. (2017). SPRING: a kinetic interface for visualizing high
dimensional single-cell expression data. Bioinformatics *34*, 1246–1248.
https://doi.org/10.1093/bioinformatics/btx792.

Westerfield, M. (1993). The zebrafish book: a guide for the laboratory use of zebrafish
(Brachydanio rerio).

Wolburg, H., and Paulus, W. (2010). Choroid plexus: biology and pathology. Acta
Neuropathologica *119*, 75–88. https://doi.org/10.1007/s00401-009-0627-8.

Xie, J., Farage, E., Sugimoto, M., and Anand-Apte, B. (2010). A novel transgenic zebrafish
model for blood-brain and blood-retinal barrier development. BMC Developmental Biology *10*,
76. https://doi.org/10.1186/1471-213x-10-76.

Xu, Q., Wang, Y., Dabdoub, A., Smallwood, P.M., Williams, J., Woods, C., Kelley, M.W., Jiang,
L., Tasman, W., Zhang, K., et al. (2004). Vascular Development in the Retina and Inner Ear
Control by Norrin and Frizzled-4, a High-Affinity Ligand-Receptor Pair. Cell *116*, 883–895.
https://doi.org/10.1016/s0092-8674(04)00216-8.

Ye, Z., Chen, J., Hu, X., Yang, S., Xuan, Z., Lu, X., and Zhao, Q. (2020). SPOCK1: a multi-
domain proteoglycan at the crossroads of extracellular matrix remodeling and cancer
development. Am J Cancer Res *10*, 3127–3137. .

Zhou, Y., and Nathans, J. (2014). Gpr124 Controls CNS Angiogenesis and Blood-Brain Barrier
Integrity by Promoting Ligand-Specific Canonical Wnt Signaling. Dev Cell *31*, 248–256.
https://doi.org/10.1016/j.devcel.2014.08.018.

Zhou, Y., Wang, Y., Tischfield, M., Williams, J., Smallwood, P.M., Rattner, A., Taketo, M.M., and
Nathans, J. (2014). Canonical WNT signaling components in vascular development and barrier
formation. J Clin Invest *124*, 3825–3846. https://doi.org/10.1172/jci76431.

Zhu, Q., Shah, S., Dries, R., Cai, L., and Yuan, G.-C. (2018). Identification of spatially
associated subpopulations by combining scRNA-seq and sequential fluorescence in situ
hybridization data. Nat Biotechnol 10.1038/nbt.4260. https://doi.org/10.1038/nbt.4260.

Figure 1. Spock1 induces BBB functional development.

(**A-E**) Fluorescent tracer leakage assays in larval zebrafish reveal BBB leakage in the forebrain and midbrain of both injected 1 kDa Alexa Fluor 405 NHS Ester (turquoise) and 80 kDa DBP-EGFP (green) of *hm41* homozygous larvae (C) compared to wild type controls (B) throughout larval development, as quantified in D and E. (**F-G**) Sagittal sections of adult brains show that the mutant leakage persists into adulthood (G). (**H**) Linkage mapping of the *hm41* mutant phenotype reveals tight linkage to *spock1* on chromosome 14. (**I**) *Spock1^hm41^* has several point mutations (T241A and several silent mutations) in the SPARC domain. *Spock1^hm43^* has a deletion of the 5' UTR and start codon. (**J-O**) Dextran leakage assays show *spock1^hm43/hm43^* mutants (K) have

642   increased BBB leakage (L). *spock1$^{hm43/hm43}$* mutants have increased expression of the leaky

643   vessel marker plvapb (N and O). (**P-S**) Compound *spock1$^{hm43/hm41}$* heterozygotes also display

644   increased NHS Ester (Q) and Dextran (S) leakage compared to *spock1$^{hm41/+}$* heterozygote

645   siblings, which confine both injected tracers at 5 dpf (P and R). Scale bars represent 50 μm (C,

646   N, S) and 200 μm (G). **** p<0.0001 by 2way ANOVA (D and E) and by t test (L and O).

Figure 2. Neuronal Spock1 regulates endothelial cells within a 10-20 µm range. (**A-C**) Whole-mount HCR *in situ* hybridization for *spock1* (A) reveals signal throughout the entire 5 dpf brain, which is outlined by the fixed DBP-EGFP serum transgene (B). (**D-G**) Slide HCR *in situ* hybridization for *spock1* (D), the vascular transcript *kdrl* (E), and the neuronal transcript *elavl3* (F) reveal the lack of *spock1* signal in the vasculature (outlined in white). (**H**) Schematic of transplantation experiments. Cells from donor embryos labeled with 10 kDa Dextran (red) at the single cell stage are transplanted into unlabeled host embryos at sphere stage. Leakage of the injected 1 kDa AF 405 NHS Ester (turquoise) into the midbrain parenchyma is then measured in relationship to the distance (d) from the blood vessel to the nearest donor cell in the 5 dpf chimeric

larvae. (**I-K**) Representative dorsal 20 µm thick maximum intensity projection confocal image of a

chimeric larva with transplanted wild type donor cells (red) into a wild type host (I, WT→WT),

spock1^{hm41/hm41} donor cells into a *spock1^{hm41/hm41}* mutant host (J, hm41→hm41), and wild type

donor cells into a *spock1^{hm41/hm41}* mutant host (K, WT→hm41). (**L**) Quantification of the NHS

leakage in WT→WT (black line) and hm41→hm41 (magenta line) reveals no change in tracer

leakage in relationship to the nearest donor cell, with wild type fish confining the tracer and mutant

fish leaking the tracer. However WT→hm41 (turquoise line) transplants reveal a full rescue of the

leakage in the mutant background when the transplanted cell is within 10 µm of a blood vessel

and no effect if the donor cell is further than 20 µm from the vessel. (**M-N**) Zoomed in images of

WT→hm41 transplants. The white arrows point to instances of local rescue of tracer (turquoise)

leakage when the wild type donor cell is close but not directly contacting the mutant vasculature

(magenta). The yellow arrowhead marks wild type donor cells that fall outside the range of

Spock1 signaling. Scale bars represent 50 µm (D and K) and 10 µm (G and N). * p=0.0495, ***

p=0.0001, **** p<0.0001 by 2way ANOVA compared to WT→WT transplants.

Figure 3. Spock1 mutants have increased endothelial vesicles. (**A-B**) The neurovascular unit
remains intact in *spock1^{hm41/hm41}* mutants (B) with a continuous single layer of endothelial cells
(pseudocolored magenta) enclosing the lumen (pseudocolored orange) and in close contact with
pericytes (pseudocolored green). (**C-H**) The majority of tight junctions (white arrowheads) are
functionally restrictive in the *spock1^{hm41}* mutant endothelial cells (88%). Mutant endothelial cells
displayed a significant increase in vesicular density, including both small flask shaped vesicles
(yellow stars, G) and larger vesicles greater than 200 nm in diameter (outlined by white dashed

line in F, H). (**I-K**) While pericyte coverage is unaltered in *spock1^{hm41}* mutants, the pericyte-
endothelial cell interactions are altered in the mutants, with several instances of direct pericyte-
endothelial cell contact (white arrows) and overall diminished average basement membrane
thickness between the two vascular cells (K). (**L-N**) In vivo gelatin zymography in wild type (L)
and *spock1$^{hm41/hm41}$* mutants (M) reveals significantly increased gelatinase activity in the mutant
midbrain compared to wild type siblings (N). Scale bars represent 1 µm (B), 200 nm (J), and 50
µm (M). ** p=0.0029 (H), **** p<0.0001 by nested t test (G and K) and by t test (N).

Figure 4. Spock1 plays a conserved role in inducing barrier function during embryonic development in mice. (A) Schematic of functional tracer leakage assays in embryonic day 15.5

(E15.5) mice. (**B-E**) Wild type mice confine both injected 550 Da NHS-Biotin (**B**) and 10 kDa

Dextran (**D**) within the vasculature, as previously reported. However, *Spock1$^{-/-}$* mice leak both

NHS-Biotin (**C**) and Dextran (**E**) into the brain parenchyma. (**F-G**) Quantification of the total area

of NHS-Biotin leakage (F) and Dextran leakage (G) normalized to vessel area where a ratio of 1

indicates no leakage reveals a significant increase in extravasation of both tracers in the *Spock1$^{-/-}$*

embryos (p=0.0213 (F) and p=0.0048 (G) by nested t test). (**H-K**) Zoomed in view of 10 kDa

Dextran (green) confined within the vasculature of wild type (H) vessels and leaked out into the

cortex of *Spock1$^{-/-}$* knockouts (I). This increased BBB permeability in *Spock1$^{-/-}$* mice is

accompanied by an increase in PLVAP (magenta) expression in the vasculature (K). (**L**)

Quantification of PLVAP expression within the vasculature reveals a significant increase in

PLVAP expression in the *Spock1$^{-/-}$* embryos (p<0.0001 by nested t test). N=4 embryos for each

genotype, marked by unique colors, with 5 sections analyzed per embryo. Scale bars represent

100 µm (E) and 50 µm (I).

Figure 5. *Spock1^hm41/hm41* mutant vascular cells have reduced expression of BBB

regulators. (**A**) UMAP of the subclustered vascular cells separated by cell type, with pericytes

(purple) separating from vascular smooth muscle cells (vSMCs, pink) and endothelial cells

(green). (**B-C**) Mean gene expression in wild type (WT, grey bars) and *spock1* mutant (hm41,

black bars) endothelial cells (B) and pericytes (C). Error bars represent SEM. Mutants appear to

have lower levels of *mcamb* in both pericytes and endothelial cells. (**D-G**) HCR FISH reveals

strong expression of *mcamb* (turquoise) in wild type vasculature (magenta) (D), both in the

midbrain and hindbrain. *Spock1* mutants have significantly reduced expression of *mcamb* in the

midbrain (E, F), but normal levels in the hindbrain (G), where no leakage is observed. Scale bar

represents 10 μm. ** p=0.0044, **** p<0.0001 by unpaired t test.

Figure 6. Spock1 induces BBB properties by modulating the brain microenvironment, altering vascular cell biology non-autonomously. (**A-C**) A single intracranial injection of human rSPOCK1 into the brain at 5 dpf reduces mutant leakage of DBP-EGFP at 6 dpf about 50% (B) compared to controls injected with PBS alone (A), quantified in C. (**D-F**) Similarly, injection of rSPOCK1 at 4 dpf reduces mutant leakage of 10 kDa Dextran at 5 dpf (E), quantified

720   in F. (**F-I**) The addition of rSPOCK1 reduces gelatinase activity in the mutant brain (H) compared

721   to control injected mutants (F), quantified in I. (**J-L**) Vascular expression of *mcamb* in the midbrain

722   is partially restored following rSPOCK1 intracranial injections (K), quantified in L. Scale bars

723   represent 50 µm (C, E, F) and 10 µm (K). **** p<0.0001 by 2way ANOVA compared to PBS

724   injected control *spock1*$^{hm41/hm41}$ mutants in C and by t test in I and L, *** p<0.001 by t test in F.

 Supplementary Materials:

Materials and Methods

Zebrafish Strains and Maintenance

Zebrafish were maintained at 28.5°C following standard protocols(Westerfield, 1993). All

zebrafish work was approved by the Harvard Medical Area Standing Committee on Animals under

protocol number IS00001263-3. Adult fish were maintained on a standard light-dark cycle from 8

am to 11 pm. Adult fish, age 3 months to 2 years, were crossed to produce embryos and larvae.

For imaging live larvae, 0.003% phenylthiourea (PTU) was used beginning at 1 dpf to inhibit

melanin production. These studies used the AB wild-type strains and the transgenic reporter

strains Tg(*l-fabp:DBP-EGFP*)[lri500] (Xie et al., 2010), (Tg(*kdrl:HRAS-mCherry*)[s896] (Chi et al., 2008),

abbreviated as Tg(*kdrl:mCherry*) in the text, TgBAC(*pdgfrb:EGFP*)[ncv22Tg] (Ando et al., 2016),

abbreviated as Tg(*pdgfrb:EGFP*), and Tg(*plvapb:EGFP*)[sj3Tg] (Umans et al., 2017), abbreviated

as Tg(*plvapb:EGFP*) in the text.

Spock1[hm41] Mutants

Spock1[hm41] mutants were maintained in the double transgenic Tg(*l-fabp:DBP-EGFP;*

kdrl:mCherry) background. Heterozygous fish were intercrossed for all leakage assays, with the

exception of the time lapse microscopy experiments, where a heterozygous fish was crossed to

a homozygous mutant, and the cell transplantation experiments, where homozygous mutants

were in-crossed. All larvae were imaged prior to genotyping to identify wild type and mutant fish.

The *spock1[hm41]* mutant line was genotyped using 5'-ACTGAGTGTTATTTTGTCATTGTGC-3' and

5'-TGATGCTGATCTGAGAAGTTTAGCC-3' primers followed by a HaeIII restriction digest, which

does not digest the wild-type product (327 bp).

Spock1[hm43] Mutants

Spock1[hm43] mutants were generated and maintained in the double transgenic Tg(*plvap:EGFP;*

kdrl:mCherry) background using CRISPR mutagenesis with two sgRNAs (5'-

GTTCGTCACTTTAAAAGGAG-3' and 5'-TCTCAGATGATGTTGAGTCT-3') to target the *spock1*

5' UTR and start codon. Heterozygous fish were intercrossed for all leakage assays. All larvae

were imaged prior to genotyping to identify wild type and mutant fish. The *spock1[hm43]* mutant line

was genotyped using 5'-TAAACTTCGATTGTGCTTTGGTTT G-3' and 5'-

AATGACTACAGACCTTTGAGGGAAC-3' primers and KAPA HiFi Hotstart polymerase

(Roche:KK2602) using a 58°C melting temperature. The mutant PCR product of 364 bp was
easily distinguishable by gel electrophoresis from the wild type product at 935 bp.

Fluorescent Zebrafish Tracer Injections and Live Imaging
Larvae were immobilized with tricaine and placed in an agarose injection mold with their hearts
facing upwards. 2.3 nl of Alexa Fluor 405 NHS Ester (Thermo Fisher: A30000) or Alexa Fluor 647
10 kDa Dextran (Thermo Fisher: D22914) fluorescently conjugated tracers (10 mg/ml) were
injected into the cardiac sac using Nanoject II (Drummond Scientific, Broomall, PA). Larvae were
then mounted with 1.5% low gelling agarose (Sigma: A9414) in embryo water on 0.17 mm
coverslips and imaged live within 2 hours post injection on a Leica SP8 laser scanning confocal
microscope using the same acquisition settings with 1 μm z-steps using a 25x water immersion
objective. All quantification was performed on blinded image sets. For static images, parenchymal
fluorescent tracer intensity was measured using ImageJ in the entire regional parenchyma outside
of the vasculature in 60 μm thick maximum intensity projections of the larval brains. These
projections began on average 15 μm below the mesencephalic vein to reduce the effects of
potential leakage diffusion from the surface vessels and had the vasculature masked and
removed for intensity quantification. These parenchymal tracer intensity values were then
background subtracted and normalized to the tracer intensity within the vasculature to account
for differential amounts of circulating tracer between fish. For time lapse imaging, Dextran
intensity was measured in six parenchymal regions of average intensity projections of the time
lapse videos and averaged as a single value per fish and similarly normalized to the average
blood vessel luminal fluorescence intensity at each time point. Pericyte and vascular double
transgenics were similarly imaged on the Leica SP8 for quantification of vascular defects.
Additional images were taken on a Zeiss LSM 980 with Airyscan 2 with 0.21 μm z-steps using a
20x water immersion objective for increased cellular resolution, allowing us to detect gaps
between the two cell types in vivo.

Linkage Mapping
Larvae from 2 separate crosses were screened for leakage of DBP-EGFP at 5 dpf and pooled
into 3 groups of 5 to 6 leaky or wild type heads. RNA was extracted from the pools using RNeasy
mini kit and ribo-depleted. RNA sequencing libraries were prepared using Wafergen Directional
RNA-Seq kits and sequenced on NextSeq High-Output sequencers producing 75 bp paired end
reads. Reads were mapped to the GRCz11 genome using tophat and bowtie2. Linkage mapping
was performed on the mapped reads using RNAmapper(Miller et al., 2013) with the following

specifications: zygosity=25, coverage=1, linkedRatio=0.96, neighbors=10. Differential gene
expression analysis was performed on these libraries using rsem(Li and Dewey, 2011).
796

CRISPR Mutants
Gstp2 crispant fish were generated by injection of Cas9 protein and 3 guide RNAs (5'-
CAGCTGCCTAAATTTGAAGA-3', 5'-GCGTTGGAAACTTACACATG-3', and 5'-GTGAG
AGTGTAGGGAGCCAC-3') into 1-cell fertilized double transgenic Tg(*l-fabp:DBP-EGFP;*
kdrl:mCherry) embryos. *Csf1ra* crispant fish were generated similarly with 4 guide RNAs (5'-
CTGCTCACCAACAGCCGAG-3', 5'-GTGTCTTCTGACCGACCCGG-3', 5'-CTCGTC
TTCATGCTTCACG-3', and 5'-AGTGACACCTTCTCCATGG-3'), as were *spock1* crispants (5'-
GTAGCCGACAGAAAGAGAGG-3', 5'-GAGTCGCAGGAGTTGAACAG-3', 5'-
GACAGTGAACCTTCATGCAG-3', and 5'-TGTCCGGGCAGGCAAGGGCA-3'). F0 crispants
were analyzed for leakage of the DBP-EGFP tracer outside of the kdrl:mCherry labeled
vasculature at 5 dpf.
808

HCR Fluorescent In Situ Hybridization (FISH)
Slide: HCR RNA in situ hybridization (Molecular Instruments) experiments on 14 µm cryosections
of fixed 5 dpf larvae were performed as previously described (Choi et al., 2018; O'Brown et al.,
2019). Briefly, sections were air dried and re-fixed in 4% paraformaldehyde (PFA) in PBS for 10
minutes at room temperature. Following fixation, slides were washed in PBS and then
permeabilized using 1 µg/ml Proteinase K (ThermoFisher) for 5 minutes, followed by PBS washes
and refixation. Tissues were further permeabilized by an ethanol dehydration series of 50%, 70%
and two rounds of 100% ethanol and air dried for 5 minutes. Dried slides were put into probe
hybridization solution for at least 10 minutes at 37°C. Subsequently, the probes for *spock1, kdrl,*
elavl3, and *mcamb* were added to the slides at a final concentration of 4 nM to hybridize overnight
at 37°C. The next day, slides were washed with a series of wash buffer to 5x SSCT (5x SSC with
0.1% Tween 20). Excess liquid was then removed and samples were immersed in amplification
buffer at room temperature for 30 minutes prior to hairpin amplification, which occurred overnight
at room temperature. Slides were then washed in 5x SSCT, washed in 5x SSC and mounted with
Fluoromount-G (Electron Microscopy Sciences).
824

Whole mount: Larvae were fixed in 4% PFA overnight and then stored in methanol at -20°C until
staining was performed. All staining was performed in PCR strip tubes. Larvae were rehydrated
in a series of methanol/PBST (0.1% Tween) washes and then permeabilized with 30 µg/ml

Proteinase K for 45 minutes at room temperature. Larvae were then refixed in 4% PFA for 20
minutes and treated with probe hybridization buffer at 37°C. Subsequently, the *spock1* probe was
added at a final concentration of 4 nM to hybridize overnight rocking at 37°C. The next day, larvae
were washed with a series of wash buffer to 5x SSCT (5x SSC with 0.1% Tween 20). Excess
liquid was then removed and samples were immersed in amplification buffer at room temperature
for 30 minutes prior to hairpin amplification, which occurred overnight rocking at room
temperature. Larvae were then washed and imaged in 5x SSCT.
835
Images for quantification were collected on a Leica SP8 laser scanning confocal microscope using
the same acquisition settings with 0.2 µm z-steps using a 25x water immersion objective or a 63x
oil immersion lens. All quantification was performed on blinded image sets of sections from a
single slide treated with all of the same reagents and imaged on the same day. To measure levels
of *mcamb* expression, we used ImageJ to first subtract tissue autofluorescence (captured in the
405 channel) from the *mcamb* expression in maximum intensity projections and then the
vasculature was manually traced and average intensity was measured. Additional images were
taken on a Zeiss LSM 980 with Airyscan 2 with 0.21 µm z-steps using either a 20x or a 40x water
immersion objective to further enhance the resolution of the *mcamb* signal.
845

Transplantation
Donor embryos were injected with 2.3 nl of 10 mg/ml Alexa Fluor 647 10 kDa Dextran (Thermo
Fisher: D22914) at the 1-cell stage to distinguish them from host cells. Following injection,
embryos were incubated at 28.5°C until transplantation. Host and donor embryos were
dechorinated with 1 mg/ml Pronase (Roche:11459643001) at oblong stage and transferred to
transplantation agarose dishes in 1/3 Ringer's buffer. Unfertilized or injured embryos were
discarded. To generate clonal sources secreting wild type or mutant Spock1, approximately 40–
80 cells were transplanted from sphere stage dextran labeled donor embryos into sphere stage
wild type and mutant hosts (similar to previous studies(Pomreinke et al., 2017)). Embryos
recovered overnight in 1/3 Ringer's buffer and were subsequently transferred to Danieau Buffer
with PTU. Tracer injections were performed at 5 dpf, as described above, with 1 kDa Alexa Flour
405 NHS Ester. Blinded image sets were analyzed using ImageJ. Individual larval brains were
segmented into 10 µm thick maximum intensity projections spanning the entire brain. Average
NHS tracer intensity was measured in 10 µm wide swaths connecting donor cells to the nearest
blood vessel and normalized to average blood vessel tracer intensity. The median tracer intensity
for a given distance (D) to the nearest donor cell was calculated for each individual fish.

Transmission Electron Microscopy (TEM)

Larvae (7 dpf) were anesthetized with tricaine and injected with 2.3 nl of 5 nm NHS-activated gold nanoparticles (Cytodiagnostics: CGN5K-5-1, ~1.1^{14} particles/ml in PBS) just as for the fluorescent tracer injections. After 5 minutes of circulation, the larvae were initially fixed by immersion in 4% paraformaldehyde (VWR:15713-S) /0.1M sodium-cacodylate (VWR:11653). Following this initial fixation, larvae were further fixed for 7 days in 2% glutaraldehyde (Electron Microscopy Sciences: 16320)/ 4% paraformaldehyde/ 0.1M sodium-cacodylate at room temperature. Following fixation, larvae were washed overnight in 0.1M sodium-cacodylate. Entire larval heads were post-fixed in 1% osmium tetroxide and 1.5% potassium ferrocyanide, dehydrated, and embedded in epoxy resin. Ultrathin sections of 80 nm were then cut from the block surface and collected on copper grids. Grids were imaged using a 1200EX electron microscope (JEOL) equipped with a 2k CCD digital camera (AMT) and quantified using ImageJ (NIH). Vesicular density values were calculated from the number of non-clathrin coated small vesicles less than 100 nm in diameter or large vesicles greater than 200 nm in diameter per μm^2 of endothelial area for each image collected. Average pericyte basement membrane (BM) thickness was quantified by measuring the total BM area divided by the length of the pericyte-endothelial contact. All images for analysis were collected at 12000x magnification. 10-15 vessels were quantified for each fish, with each color representing a different fish.

Intracranial rSPOCK1 Injections

Larvae were anesthetized with tricaine and placed in an agarose injection mold with their heads facing upwards. 2.3 nl of recombinant human SPOCK1 (rSPOCK1; R&D Systems: 2327-PI-050) at a range of 0.001-10 mg/ml was injected directly into the midbrain using Nanoject II (Drummond Scientific, Broomall, PA). Following injections, larvae recovered in normal embryo water for 1 day. The effects of rSPOCK1 were either assess in live fish using functional tracer leakage assays and in vivo gelatin zymography or in fixed samples followed by slide HCR in situ hybridization, as described above.

In vivo Gelatinase Zymography

Larvae were anesthetized and intracranially injected as for rSPOCK1 with 2.3 nl of FITC-Gelatin (Biovision: M1303) at 5 mg/ml. After one hour, larvae were also injected with 10 kDa Dextran in the cardiac sac to assess BBB function and then mounted with 1.5% low gelling agarose (Sigma: A9414) in embryo water on 0.17 mm coverslips and imaged live within 2 hours post injection on

a Leica SP8 laser scanning confocal microscope using the same acquisition settings with 1 μm
z-steps using a 25x water immersion objective. All quantification was performed on blinded image
sets. Parenchymal FITC-Gelatin probe intensity was measured using ImageJ in the entire
midbrain and normalized to the average intensity in the ventricles, which have high gelatinase
activity, to account for fish to fish injection variation. Regions of obvious damage at the injection
site were not quantified.
902

Mouse Maintenance

Spock1[-/-] knockout mice (Röll et al., 2006) were obtained from Ursula Hartmann's lab and were
backcrossed to and maintained on a C57Bl/6 background. Mice were genotyped with primers 5'-
GCCACTGGTCATTGTCTAGG-3', 5'- TGTGCCCAGTCATAGCCGAATAGC CTCTCC-3', and 5'-
GCTTGAGGTAGCCCTGTTGTCACC-3' using KAPA HiFi Hotstart polymerase (Roche:KK2602).
This PCR reaction produced a 185 bp wild type band and a 750 bp knockout band in heterozygous
animals. All animals were treated according to institutional and US National Institutes of Health
(NIH) guidelines approved by the Institutional Animal Care and Use Committee (IACUC) at
Harvard Medical School under protocol IS00000045-6.
912

Mouse Tracer Injections

Heterozygous *Spock1*[+/-] mice were intercrossed and used for tracer leakage assays performed at
embryonic day 15.5 (E15.5) as previously described(Ben-Zvi et al., 2014). In brief, 5 μl of tracer
cocktail (10 mg/ml EZ-Link NHS-Biotin (Thermo Fisher: 20217) and 5 mg/ml 10 kDa Dextran
Alexa Fluor 488 (Thermo Fisher: D22910)) was injected into the liver of each embryo and allowed
to circulate for 5 minutes. Embryonic heads were fixed by immersion in 4% paraformaldehyde
overnight at 4°C and then frozen in TissueTek OCT (Sakura). 20 μm thick sections were then
collected and immunostained with Streptavidin Alexa Fluor 405 (1:200; Thermo Fisher: S32351)
and rat anti-PLVAP (1:200; BD Biosciences:553849). All embryos were injected and fixed for
processing blind before genotyping. Adult mice were retroorbitally injected with 0.5 mg of EZ-Link
NHS-Biotin per g of mouse body weight. The NHS-Biotin circulated for 30 minutes prior to brain
dissections and fixation with 4% PFA. 30 μm thick cryosections were then collected and
immunostained with Streptavidin Alexa Fluor 568 (1:200; Thermo Fisher: S32351) and goat anti-
CD31 (1:100; R&D Systems: AF3628).
927

Single-cell RNA-sequencing

Larval zebrafish brains were dissected and split along the midbrain-hindbrain boundary in DMEM.
Brains were dissociated using a modified protocol(Bresciani et al., 2018). Briefly, chemical
dissociations were performed at 30.5°C using a mixture of 0.25% Trypsin-EDTA,
Collagenase/Dispase (8 mg/mL) and DNaseI (20 µg/mL) for 15-20 minutes with gentle pipetting
every few minutes and quenched with 10% fetal bovine serum in DMEM. Samples were hashed
using Multi-seq as previously described with slight modifications(McGinnis et al., 2019). For each
sample, 80 pmoles of lipid modified oligos (LMOs) were used to hash every 500k cells. The
hashing reaction was quenched using 1% BSA in PBS and barcoded samples were subsequently
pooled and washed with 1% BSA. The pooled cell mixture was resuspended in PBS + 0.1%BSA
+ 18% Optiprep at a final concentration of ~300k cells/mL prior to single-cell capture with inDrops.
The Single-cell Core (SCC) at Harvard Medical School captured single-cell transcriptomes and
prepared NGS libraries as previously described(Klein et al., 2015) with a target capture of 45k
cells per experiment. A summary of dissected tissues and corresponding sequencing information
are described in Table S4.

Gene expression and hashtag libraries were mixed (9:1 ratio) and sequenced on an Illumina
Nova-seq 6000 with the NovaSeq S2 kit. Reads were mapped onto the Zebrafish GRCz11
Release 101 genome assembly using previously described methods(Wagner et al., 2018).
Hashtags were identified using custom code available on:
https://github.com/AllonKleinLab/paper-data/tree/master/OBrown2021_ZebrafishBBB

Transcriptomes with greater than 350 UMIs were further filtered for viability by removing cells with
>20% mitochondrial reads. Cell demultiplexing was performed manually by applying thresholds
to delineate single cells from background and multiple populations. The resulting counts matrix
was normalized to the mean UMIs per cell in the dataset.

To visualize the data, we first mean-normalized the data and identified highly variable genes from
the wild type AB and RNF datasets (minimum of 3 transcripts per cell, minimum of 3 cells
expressing gene, minimum V-score percentile of 85%). We then Z-scored counts for each gene
and performed principal component analysis (PCA) with 50 components. Cells from the spock1
mutant libraries were projected into the same principal component subspace. A k-nearest
neighbor graph (k=10) was generated based on the Euclidean distance in gene expression
between cells within this subspace. The Leiden algorithm was used to cluster cells into
subgroups(Traag et al., 2019). The neighborhood graph was embedded and visualized using

UMAP and the data was explored interactively with SPRING(Weinreb et al., 2017) to aid in cell
cluster annotation.
965
Cells belonging to the liver, pharyngeal arches, skin, and muscle were removed prior to differential
gene expression (DGE) analysis. The filtered dataset was reanalyzed using the methods
described above. The Wilcoxon rank-sum test was used to generate a list of potential genes that
are differentially regulated (fold change > 2) between genotypes across each Leiden cluster
(Table S2). Genes were only considered for DGE analysis if they were expressed in at least 5%
of cells in a given cluster and had a minimum mean of 10 transcripts per cell. Cells belonging to
the AB and RNF background were treated as a single wild type genotype for the analysis. The
vascular cluster (Leiden 13) was subclustered and similarly analyzed for DGE (Table S3).
974
975 **Data availability**
976 The raw reads from the counts matrix and associated metadata from the single-cell experiments
977 have been deposited to GEO.
978
979 All code for the single-cell analysis is available as interactive Jupyter notebooks here:
980 https://github.com/AllonKleinLab/paper-data/tree/master/OBrown2021_ZebrafishBBB
981
982 Interactive exploration of the full single-cell data can be found here:
983 https://kleintools.hms.harvard.edu/tools/springViewer_1_6_dev.html?client_datasets/2021_OBr
984 own/2021_OBrown
985
986 Exploration of the vascular subcluster cells can be found here:
987 https://kleintools.hms.harvard.edu/tools/springViewer_1_6_dev.html?client_datasets/2021_OBr
988 own_Vasculature/2021_OBrown_Vasculature

     Supplemental Figures

Figure S1. ***Spock1$^{hm41/hm41}$ mutants have normal vascular patterning.** (**A-B**) Dorsal maximum

intensity projection of 5 dpf wild type (**A**) and mutant (**B**) brains with endothelial cells marked by

the kdrl:mCherry transgene (magenta) and pericytes marked by the pdgfrb:EGFP transgene

(green). (**C-D**) Zoomed in view of vessels reveals diminished basement membrane between

pericytes (green) and vessels (magenta) in mutants (**D**) compared to wild type siblings (**C**). This

is further highlighted in the higher magnification insets, with a visible gap present in the wild type

vessel (**C**) but not in the mutants (**D**). (**E-F**) Quantification of total vascular coverage of the brain

(kdrl:mCherry+ area/total brain area, **E**) and total pericyte coverage of the vasculature

(pdgfrb:EGFP+ area/kdrl:mCherry+ area, **F**), with each individual fish marked by a single point.

Scale bars represent 50 µm (**B**) and 10 µm (**C**). T-test comparison reveals no significant difference

between wild type and mutant fish for either vascular or pericyte coverage.

1002

Figure S2. Time lapse imaging reveals leakage dynamics in *spock1^hm41/hm41* mutants. (**A-**

B) Dorsal maximum intensity projection of a wild type (A) and *spock1^hm41/hm41* mutant (B) midbrain

reveals steady accumulation of 10 kDa Dextran (yellow) outside of the vasculature (magenta)

over the course of one hour in *spock1* mutants. (**C**) Quantification of Dextran accumulation in the

midbrain parenchyma outside of the vasculature over time in wild type fish (black line) and

spock1^hm41/hm41 mutants (magenta line). Scale bar represents 50 μm.

1009

1010

Figure S3. Leaky phenotype maps to *spock1* gene on chr14:2205271-3513919. (**A**)
Manhattan plot of linkage on chromosome 14 with the average mutant alleles per 20 neighbors
plotted in red. The region of highest linkage is highlighted in yellow. (**B**) Genome browser view of
the highest linked region to the leaky phenotype. Eight of the genes within this region were
expressed in the 5 dpf bulk RNAseq data: *fgf18a, spock1, bicc2, csf1ra, ndst1a, slc35a4, gstp1*
and *gstp2*. Two of these genes were differentially expressed in leaky mutants compared to wild
type (*csf1ra* and *gstp2*) and *spock1* (marked in red) had several SNPs that completely segregated
in the leaky fish. (**C**) Mosaic *spock1* crispants display increased leakage of the transgenic DBP-
EGFP tracer outside of the vasculature (outlined by white dashed line). (**D-E**) Dorsal view of whole
mount HCR in situ for *spock1* (white) reveals expression throughout the brain of uninjected
controls (D) but not in *spock1* crispants (E). (**F-G**) While controls confine the DBP-EGFP tracer
(green) within the blood vessels (magenta) in the brain (F), *spock1* crispants display increased
leakage into the brain parenchyma (G, outlined by white dashed line), corresponding with the loss
of *spock1* expression (E). Scale bars represent 50 µm.

1025
Figure S4. Expression of *spock1* is unaltered in *spock1^{hm41/hm41}* mutants and absent in
***spock1^{hm43/hm43}* mutants.** (**A-B**) *Spock1* expression (yellow) is found throughout the brain and
spinal cord (outlined by a white dashed line) in wild type (A) and *spock1^{hm41/hm41}* mutant fish (B) at
5 dpf. *Spock1* expression never colocalizes with vascular *Tg(kdrl:mCherry)* expression
(magenta). Tissue autofluorescence is depicted in turquoise. (**C-D**) Unlike *spock1^{hm41/hm41}* fish and
wild type fish (C), *spock1^{hm43/hm43}* mutants, which completely lack the 5' UTR and start codon of
spock1, do not have any detectable *spock1* expression in the brain (D). The remaining signal in
the yolk and blood vessels is due to tissue autofluorescence. Scale bar represents 50 µm.

1034

1035 **Figure S5. Adult *Spock1^{-/-}* mice recover BBB function.** (**A-B**) Tracer leakage assays in adult

1036 wild type (A) and *Spock1^{-/-}* mice (B) reveals that both genotypes confine the injected Sulfo-NHS-

1037 Biotin tracer (green) within the CD31+ vasculature (magenta), indicating a functional BBB. Scale

1038 bar represents 50 μm.

1039

Figure S6. scRNA-seq captures all neurovascular cells in both mutant and wild type larvae.
(**A-B**) UMAP of the whole brain dataset separated shows overlap between wild type (A) and
spock1^{hm41/hm41} mutant (B) cell type coverage, indicating that no cell type is absent in the mutant
background. (**C**) UMAP of the total data set separated by annotated cell type, with the vast
majority of sequenced cells being neuronal (blue). Our data does capture blood cells (red),
microglia (magenta), oligodendrocytes (aqua), radial glia (yellow), and vascular cells (green). (**D-
L**) Gene expression plots for *elavl3* (D) for neurons, *slc1a2b* (E) for radial glia, *fabp11a* (G) and
kdrl (H) for endothelial cells, *pdgfrb* (I) for pericytes and neurons, *sox10* (J) and *olig2* (K) for

1048   oligodendrocytes, and *lyve1b* (L) for lymphatic endothelial cells reveals that *spock1* (F) is primarily

1049   expressed by neurons and absent from the vascular cells.