Dietary super-doses of cholecalciferol in aged laying hens illustrates limitation of 24,25-dihydroxycholecalciferol conversion

Matthew F. Warren[1], Pete M. Pitman[2], Dellila D. Hodgson[1], Nicholas C. Thompson[3], and Kimberly A. Livingston[1]

[1]Prestage Department of Poultry Science, North Carolina State University, Raleigh, NC, USA

[2]Department of Biological Sciences, North Carolina State University, Raleigh, NC, USA

[3]Department of Animal Science, North Carolina State University, Raleigh, NC, USA

Sources of support: Funding for this research was supported by the US Poultry & Egg Association (KAL); the US Poultry & Egg Association had no involvement or restrictions regarding publication.

Corresponding Author Information: Matthew F. Warren, Department of Bacteriology, University of Wisconsin-Madison, 1550 Linden Drive, Madison, WI 53706-1205, (559) 816-0639, mwarren8@wisc.edu

Abbreviations used: $1,25\text{-}(OH)_2\text{-}D_3$, 1,25-dihydroxycholecalciferol; $24,25\text{-}(OH)_2\text{-}D_3$, 24,25-dihydroxycholecalciferol; $25\text{-}OH\text{-}D_3$, 25-hydroxycholecalciferol; CYP27C1, 1α-hydroxylase; CYP24A1, 24-hydroxylase; VDR, vitamin D receptor

Teaser Text: Feeding very high levels of dietary vitamin D_3 to aged laying hens leads to increased plasma and egg yolk vitamin D_3 metabolite levels.

ABSRACT

Background: Older humans who take high levels of vitamin D supplementation for a

prolonged time may be at risk of vitamin D toxicity. It is unclear how dietary super-doses

(10,000x greater than requirement) can affect vitamin D status in aged animals. Aged

laying hens could be a model to compare vitamin D supplementation effects with

women in peri- or postmenopausal stages of life.

Objective: We investigated dietary super-dose impacts of cholecalciferol (vitamin D_3)

on vitamin D status in aged laying hens in production.

Methods: Forty-eight 68 wk old Hy-Line Brown laying hens were individually housed in

cages with eight hens per dietary treatment for eleven weeks. Hens were randomly

assigned to one of six treatment groups of dietary vitamin D_3 supplementation and fed

ad libitum. Supplementation levels were 400, 800, 7,400, 14,000, 20,000, and 36,000 IU

D_3/kg of feed. At the end of the study, all hens were euthanized and tissue samples and

feces were collected. Plasma and egg yolk vitamin D metabolites, calcium and

phosphorus composition of multiple sites, and tissue gene expression levels were

measured.

Results: We observed that increasing dietary vitamin D_3 increased plasma vitamin D_3

and egg yolk vitamin D_3 ($p < 0.0001$ for both sites). We also observed an increase in

plasma 24,25-dihydroxycholecalciferol as dietary vitamin D_3 levels increased ($p <$

0.0001). The plasma 25-hydroxycholecalciferol:24,25-dihydroxycholecalciferol ratio

exhibited an asymptotic relationship starting at the 14,000 IU/kg D_3 treatment.

Conclusions: Dietary super-doses of vitamin D_3 led to greater plasma and egg yolk

vitamin D_3 levels to show that aged laying hens can deposit excess vitamin D_3 in egg

yolk. We suggest future research should explore how 24-hydroxylation mechanisms are

affected by vitamin D supplementation. Further understanding of 24-hydroxylation can

help ascertain ways to reduce risk of vitamin D toxicity.

Keywords: aged laying hen, chicken, dietary vitamin D_3, 25-hydroxycholecalciferol,

24,25-dihydroxycholecalciferol, egg, egg yolk vitamin D_3, plasma vitamin D_3, super-dose

Introduction

Older humans taking extremely high levels of vitamin D supplementation for a prolonged time may be at risk of vitamin D toxicity (1, 2). People tend to take vitamin D supplements to increase or maintain vitamin D levels (3). Also, older women take vitamin D supplements to manage the hormonal effects of menopause on bone resorption (4-6). While vitamin D toxicity is uncommon, vitamin D supplements and overfortified foods are the only known means of reaching intoxication levels (1, 7). Vitamin D supplementation can be administered through multiple means with the common routes being oral dose supplementation or dietary supplementation. There was a study that explored how human adults age 65+ given daily oral doses of 1,600 IU cholecalciferol (vitamin D_3) for 1 year did not exhibit signs of toxicity (8). This illustrates that taking oral doses of vitamin D_3 in a controlled manner are safe for maintaining vitamin D levels.

An important consideration is if extremely high supplementation of vitamin D_3 levels over an extended period of time would cause vitamin D toxicity. Chickens have been used extensively to explore questions involving vitamin D_3 supplementation (9-12). Effects of very high levels of dietary vitamin D_3 supplementation levels have been explored in laying hens over a 48-wk period and the hens did not exhibit histopathological effects related to vitamin D toxicity such as hypercalcemia (10). Considering older humans are likely to take vitamin D_3 supplements, further exploration of dietary vitamin D_3 supplementation effects in older hens is necessary. Characterizing how high levels of dietary vitamin D_3 affect vitamin D metabolism in older hens may help

to better understand how overfortified foods can potentially affect vitamin D metabolism

in older animals and humans.

Dietary vitamin D supplementation is important for laying hens in production

because their bone health is physiologically taxed from egg production (13, 14). Laying

hens in commercial farms are fed diets with supplemental vitamin D_3 beyond the

National Research Council (NRC) requirements (15, 16). This ensures the hens can lay

eggs and maintain adequate calcium (Ca) absorption for eggshell formation and

importantly, bone mineralization (12). There are a few studies which investigated how

very high levels of dietary vitamin D supplementation affected laying hen production and

the metabolic implications pertaining to vitamin D status (10, 11, 17). Altogether, the

aforementioned studies illustrate dietary vitamin D_3 supplementation results in vitamin

D-enriched eggs which may be a way to improve vitamin D intake for humans.

Understanding the impacts of very high dietary vitamin D supplementation in aged

laying hens have implications with older women and their vitamin D intake from food

fortification.

Our study examined dietary vitamin D_3 super-dose effects on plasma and egg

yolk vitamin D_3 metabolites and relative gene expression of vitamin D-related genes in

aged laying hens in production. We define "super-dose" as treatment doses at least

10,000x greater than requirement. We fed hens diets containing 400, 800, 7,400,

14,000, 20,000, and 36,000 IU D_3/kg of feed to ascertain vitamin D_3 supplementation

effects. Hens consuming diets with vitamin D_3 levels greater than 7,400 IU D_3/kg were

expected to have increased plasma 24,25-dihydroxycholecalciferol (24,25-$(OH)_2$-D_3).

High levels of plasma 24,25-$(OH)_2$-D_3 would suggest the hens reached vitamin D

saturation because 24,25-$(OH)_2$-D_3 is considered an inactive form of vitamin D_3 (18, 19).

Hens consuming super-doses of vitamin D_3 should also lay eggs with increased vitamin

D_3 content because they would deposit excess vitamin D_3 into the egg yolk (10).

Methods

Animal husbandry

The hens used in our study were from North Carolina State University's

maintained poultry flock. Forty-eight 68-wk old Hy-Line Brown laying hens were housed

at North Carolina State University, Raleigh, NC and fed experimental diets for eleven

weeks (**Figure 1**). Hens were individually housed in cages between two two-level (top

level and bottom level) battery cages with eight hens per treatment. The experimental

design was a randomized complete block design with six levels of dietary vitamin D_3

supplementation blocked by cage level. Vitamin D_3 supplementation levels were

formulated to be 250, 500, 1,500, 15,000, 30,000, and 60,000 IU D_3/kg of feed, but the

analyzed vitamin D_3 levels in the feed were 400, 800, 7,400, 14,000, 20,000, and

36,000 IU D_3/kg of feed (**Table 1 and Supplementary Table 1**). The source of vitamin

D_3 used in the study was the crystalline vitamin D_3 from Alfa Aesar (Ward Hill, MA). We

refer to the six different analyzed vitamin D_3 levels as the named treatment groups for

our study. In our study, the 400 and 800 IU/kg vitamin D_3 treatments were formulated to

meet the NRC (16) requirements for laying hens. The 14,000, 20,000, and 36,000 IU

D_3/kg treatments were dietary super-dose treatments for D_3 intake. Prior to the start of

the experiment, all hens were fed the same diet (400 IU D_3/kg of feed) for one month as

a washout period. Hens were fed *ad libitum* along with water. North Carolina State

University's Institutional Animal Care and Use Committee approved all methods for this

study, protocol ID number: 18-093-A.

Sample Collection

Egg collection started 24 hours after the hens were started on the experimental

diets. Eggs were collected every morning and stored at 7°C for egg quality analyses.

Starting on Mondays, the first egg laid for the week by each hen was selected for egg

yolk collection. Eggs were cracked open in a dim-lighted room to reduce

photodegradative impacts of light on vitamin D in yolk. The egg yolk was separated from

albumen and placed in a small plastic container wrapped in aluminum foil and stored at

4°C for a year until they were freeze-dried using a freeze-dryer (FreeZone 6 Liter

Benchtop Freeze Dry System, Labconco, Kansas City, MO). On weeks 0, 3, 6, and 10,

blood was collected from the brachial wing vein from all hens to measure ionized blood

Ca using an i-STAT™ blood analyzer (Abaxis, Union City, CA) using CG8+ cartridges

(Abaxis, Union City, CA). The remaining blood was spun down to collect plasma which

was stored at -80°C. All hens were euthanized by cervical dislocation and tissue

samples were collected from 43 hens (minimum of seven hens per treatment) due to

time constraints. The duodenum, ileum, liver and kidney were collected and stored in

RNAlater at -20°C until RNA extractions were performed and the humerus and tibia

bones were collected for measuring Ca and phosphorus (P) composition.

Calcium and phosphorus content of various sites

Eggshells from weeks 0, 3, 4, 6, and 9 were washed with warm water to remove

the shell membrane and dried for 48 h at room temperature. Dried eggshells were pre-

weighed and further dried at 68°C for 72 h using a dry oven (Blue M, Atlanta, GA) and

weighed again. Eggshells were crushed into fine powder and subjected to chemical

digestion to measure Ca composition of eggshells. Feces and ileal digesta were also

subjected to same steps as eggshells. Dried samples were weighed and then placed in

a muffle furnace at 500°C overnight to ash samples. Ashed samples were dehydrated in

2 mL of distilled water and 4 mL of 6 N hydrochloric acid. The resulting sample was

mixed and heated to warm to touch. Heated solution was poured into volumetric flask

and deionized water was added to 50 mL. The flask was inverted 12 times to mix and

the resulting solution was filtered using #40 filter paper into 15 mL centrifuge tubes for

analysis. Ca and P were measured by inductively coupled plasma optical emission

spectrometry.

The humerus and tibia were wrapped in petroleum ether-moistened cheesecloth

and placed in a desiccator for 72 h to extract fat and moisture from the bones. Fat-

extracted bones were pre-weighed and dried for 24 h at 100°C to evaporate petroleum

ether residues. Fat- and moisture-free bones were weighed and ashed using the same

methods as eggshell, feces, and ileal digesta for Ca and P composition and measured

by inductively coupled plasma atomic emission spectroscopy.

RNA extraction and qPCR

Total mRNA was extracted from duodenum, ileum, liver, and kidney using

Qiagen's RNeasy Mini Kit (Germantown, MD). The extracted RNA was diluted and

normalized to ~200 ng/µL for liver and 60 ng/µL for duodenum, ileum, and kidney. The

tissues' RNA was reverse transcribed to complementary DNA (cDNA) using Applied

Biosystems' High-Capacity cDNA Reverse Transcription Kit (ThermoFisher Scientific,

Waltham, MA) and their recommended steps to make a 20 µL working solution. Cycling

procedure for reverse transcription started with 25°C for 10 min, 37°C for 120 min, 85°C

for 5 mins, then held at 5°C indefinitely until storage or use.

Genes amplified for quantitative real time PCR (qPCR) were vitamin D receptor

(VDR), 1α-hydroxylase (CYP27C1), 24-hydroxylase (CYP24A1), and glyceraldehyde 3-

phosphate dehydrogenase (GAPDH) as the housekeeping gene (**Table 2**). qPCR was

conducted using PowerUP SYBR Master Mix (Life Technologies, Grand Island, NY)

following the manufacturer's protocol and using Applied Biosystems StepOnePlus Real-

Time PCR System (Carlsbad, CA). Cycling procedure started with 95°C for 10 min then

40 cycles of 95°C for 15 s for denaturing and 15 s at 60°C for annealing. All samples

were run in triplicates.

Vitamin D metabolites

Plasma from the week 10 timepoint from hens from the 400, 800, 14,000, and

36,000 IU D_3/kg groups were respectively pooled (n = 4/treatment) and sent to

Heartland Assays (Ames, IA) for measuring D_3, 25-OH-D_3, and 24,25-$(OH)_2$-D_3 using

LC-MS/MS. Freeze-dried egg yolk from the week 10 timepoint from the same hens were

pooled (15 g per sample) like plasma, but only samples from 400, 14,000, and 36,000

IU D_3/kg groups (n = 4/treatment) were analyzed for D_3 and 25-OH-D_3 from Heartland

Assays.

Statistical analysis

We conducted statistical analyses using general linear models using SAS 9.4$^®$

(Cary, NC) for all statistical tests and Tukey-Kramer test was used for multiple

comparisons for differences between dietary treatments. We utilized repeated measures

to account for temporal effects of weekly body weight, feed intake, egg production,

eggshell quality, and ionized blood Ca. Dietary vitamin D_3 level was the independent

variable for Ca and P composition, plasma and egg yolk vitamin D_3 metabolite

concentrations, and gene expression data. Plasma vitamin D_3 was below the detection

limit (< 0.5 ng/mL) for the 400 and 800 IU D_3/kg treatments so those samples were set

to 0.4 as an arbitrary value for statistics to account for model building. All vitamin D_3

metabolite data exhibited heteroscedasticity and were transformed using natural

logarithm function. Transformed data exhibited linear and homoscedastic relationships

and were used for statistics. We did not observe a cage level effect in any analysis so

the blocking variable was omitted from all statistical tests. All mRNA relative

expressions were normalized using $2^{-C\Delta\Delta T}$ with GAPDH as a housekeeper gene.

Statistical significance was established at $p < 0.05$.

Results

Hens' production performance was not influenced by dietary vitamin D_3

To determine if dietary super-doses of vitamin D_3 affected the hens' production

value, we measured the hens' weekly body weights and egg production. Dietary super-

doses of vitamin D_3 did not affect body weight of these laying hens ($p = 0.08$;

Supplementary Table 2), but there was an interaction between dietary vitamin D_3 level

over time on feed intake ($p < 0.0001$). However, there was constant feed wastage

throughout the study so this effect could be inflated. A dietary trend was observed for

egg production for the entire study duration ($p = 0.07$; **Supplementary Table 3**).

Eggshell strength and eggshell thickness were not affected by dietary vitamin D_3 levels

($p = 0.19$, $p = 0.72$, respectively; **Supplementary Table 3**). There was a trending

interaction between dietary vitamin D_3 level over time in which there was a decrease in

eggshell elasticity ($p = 0.07$).

Ionized blood calcium is affected by dietary vitamin D_3 levels

226   We also examined if ionized blood Ca in the hens was influenced by dietary

vitamin D_3 levels throughout the study. There was no temporal effect on ionized blood

Ca ($p = 0.65$). With week 0 considered as a covariate, there was a dietary effect on

ionized blood Ca ($p = 0.002$, **Supplementary Table 4**). It is not known why there is no

clear trend on ionized blood Ca relative to the dietary treatment, but one possibility

could be related to when the hen laid an egg prior to the blood collection which may

influence circulating Ca levels.

Fecal calcium is affected by dietary vitamin D_3 levels, but not ileal digesta or

eggshell calcium or phosphorus

236   We had the eggshell, ileal digesta, and fecal Ca and P measured to determine if

dietary vitamin D_3 fed to our hens would reduce the excretion or loss of Ca and P. There

was no dietary effect on eggshell Ca and P ($p = 0.64$ for both) and ileal digesta Ca or P

(p = 0.74 and 0.09, respectively). There was a dietary effect with fecal Ca with hens fed

the 14,000 IU/kg D_3 diet having 10.6±0.8% Ca by weight in their feces, whereas all

other treatments were 8.0-9.0%, with exception of 36,000 IU/kg D_3 fed hens which had

7.36±0.43% fecal Ca (p = 0.03, **Supplementary Table 5**). There was no dietary effect

on fecal P (p = 0.76).

Humerus is more calcium and phosphorus dense than tibia

We assessed if dietary vitamin D_3 would improve bone Ca and P in the hens.

However, no dietary effects of vitamin D_3 supplementation were observed on bone Ca

or P (p = 0.79 and 0.63, respectively). Humerus bones have more Ca and P than tibia

bones (p = 0.020 and 0.015, respectively; **Supplementary Figure 1**).

Dietary super-dosage levels of vitamin D increased plasma and egg yolk vitamin

D_3 metabolites

We had plasma and egg yolk vitamin D_3 metabolites measured by LC-MS/MS to

determine if dietary vitamin D_3 affected plasma and egg yolk concentrations. There was

a significant increase in plasma concentration of vitamin D_3, 25-OH-D_3, and 24,25-

$(OH)_2$-D_3 of hens fed dietary super-dose levels of vitamin D_3 (**D_3**: p = 0.0002; **25-OH-D_3**:

p < 0.0001; **24,25-OH-D_3**: p < 0.0001; **Figures 2A-2C**). Although plasma vitamin D_3

concentration was below limit of detection for the 400 and 800 IU D_3/kg treatments, the

plasma vitamin D_3 concentration had a strong positive correlation with 25-OH-D_3 and

24,25-$(OH)_2$-D_3 concentrations ($r = 0.95$, p < 0.0001; $r = 0.92$, p < 0.0001; respectively,

data not shown). Plasma vitamin D_3 and 25-OH-D_3 had similar concentration values

when dietary vitamin D_3 increased with both metabolites having ~85 ng/mL at the

36,000 IU D_3/kg treatment level. Although plasma $24,25\text{-}(OH)_2\text{-}D_3$ concentration was

lower than vitamin D_3 and 25-OH-D_3, $24,25\text{-}(OH)_2\text{-}D_3$ was affected by dietary vitamin D_3

and it exhibited a similar rate of increase relative to dietary vitamin D_3 levels like vitamin

D_3 and 25-OH-D_3. The percentage ratio of $24,25\text{-}(OH)_2\text{-}D_3$ to 25-OH-D_3 increased as

dietary vitamin D_3 levels increased and reached an asymptote at the super-dose levels

($p < 0.0001$; **Figure 2D**). The $24,25\text{-}(OH)_2\text{-}D_3$: 25-OH-$D_3$ ratio percentage ranged from

8.7% to 20.5% with all super-dose fed hens having a ratio of approximately 20%.

Egg yolk vitamin D_3 increased drastically as hens' dietary vitamin D_3 intake

increased ($p < 0.0001$; **Figures 3A and 3B**). Egg yolk 25-OH-D_3 was also significantly

increased in concentration as hens' dietary vitamin D_3 increased ($p < 0.0001$). Although

egg yolk 25-OH-D_3 had lower concentrations compared to egg yolk vitamin D_3, the rate

of increase for 25-OH-D_3 was similar to egg yolk vitamin D_3 as dietary vitamin D_3 levels

increased. Egg yolk vitamin D_3 was strongly and positively correlated with plasma

vitamin D_3 ($r = 0.99$, $p < 0.0001$, **Figure 4A**). Egg yolk 25-OH-D_3 also had a strong

positive correlation with plasma 25-OH-D_3 ($r = 0.96$, $p < 0.0001$, **Figure 4B**).

Dietary super-doses of vitamin D_3 intake affected vitamin D receptor expression

and kidney 24-hydroxylase expression

Considering VDR is the transcription factor responsible for exerting vitamin D's

physiological effects (20), we measured VDR expression in multiple tissues to

determine if dietary vitamin D_3 levels would affect VDR expression. Hens fed higher

levels of dietary vitamin D_3 had upregulated duodenal VDR expression ($p = 0.036$;

Figure 5A). There was no dietary effect on VDR expression from ileum, liver, and

kidney (p = 0.96, 0.17, 0.32; respectively; **Figures 5B-5D**). We also examined if vitamin

D_3 super-dosages would affect the gene expression of vitamin D hydroxylase enzymes

in the kidney. Unexpectedly, kidney 24-OHase expression was lower in hens fed diets

with super-dose levels of vitamin D_3 (p = 0.0006, **Supplementary Figure 2A**). No

differences were observed with kidney 1α-OHase expression (p = 0.81, **Supplementary**

Figure 2B).

Discussion

Our results suggest that dietary super-doses of vitamin D_3 greatly increased

plasma and egg yolk D_3 levels. Increased plasma vitamin D_3 indicates these hens

absorbed vitamin D_3 from their diets. Although the inactive vitamin D_3 metabolite,

plasma $24,25\text{-}(OH)_2\text{-}D_3$ had a lower measured value than vitamin D_3 and 25-OH-D_3,

$24,25\text{-}(OH)_2\text{-}D_3$'s rate of increase was similar to vitamin D_3 and 25-OH-D_3 as dietary

vitamin D_3 levels increased. Plasma $24,25\text{-}(OH)_2\text{-}D_3$ concentrations highlights that these

hens were likely trying to reduce their circulating vitamin D_3 levels **(Figure 6)**. In

addition, egg yolk vitamin D_3 drastically increased as dietary vitamin D_3 levels

increased. Egg yolk 25-OH-D_3 had a similar rate of increase like egg yolk vitamin D_3

even though egg yolk 25-OH-D_3 concentrations were lower. Altogether, there is a strong

association between dietary vitamin D_3 levels and plasma and egg yolk vitamin D_3

metabolite levels.

Although a laying hen's physiological status affects egg quality (21), high levels

of dietary vitamin D_3 was shown to reduce egg quality, but not affect egg production

(22). Mattila et al. (10) and Wen et al. (11) reported that laying hens fed diets with

greater levels of dietary vitamin D_3 throughout their production cycle increased yolk

vitamin D_3 content. The egg production and egg yolk vitamin D_3 data in our study were

similar to the two aforementioned studies, even though the hens in our study were

older. A novel finding we observed is that vitamin D_3 is deposited more readily into yolk

compared to 25-OH-D_3. One possible explanation is that excess circulating 25-OH-D_3

was transferred to the egg yolk as a way to lower circulating 25-OH-D_3 levels.

Supplementing laying hen diets with 25-OH-D_3 increased egg yolk 25-OH-D_3 and

reduced egg yolk vitamin D_3 (23). It seems likely that the vitamin D metabolite

composition in egg yolk is influenced by whatever dietary vitamin D_3 isoform is fed to

the laying hens.

The biological significance of 24,25-$(OH)_2$-D_3 is to reduce 25-OH-D_3's plasma

concentration (24). To our current knowledge, there is no research on 24,25-$(OH)_2$-D_3

plasma concentration in laying hens. The hens' plasma 24,25-$(OH)_2$-D_3 increased

relative to dietary vitamin D_3 levels. The rate of increase for plasma 24,25-$(OH)_2$-D_3

relative to plasma D_3 and plasma 25-OH-D_3 at super-dosage levels, levels off at 20%,

highlighting a possible asymptotic relationship. The asymptote suggests 24-

hydroxylation activity hit its maximal limit. It is not clear if VDR expression is associated

with plasma 24,25-$(OH)_2$-D_3 levels or 24-hydroxylation activity because there was little

difference in VDR expression across multiple tissues in this study.

Our study has several strengths that highlight its impact with advancing

nutritional knowledge. A significant strength of our study is plasma and egg yolk vitamin

D metabolite concentration ranges across treatment groups. This illustrates the

experimental design captured a broad range of dietary vitamin D_3 supplementation level

effects on plasma vitamin D metabolites that future research can focus on a specific

range to build off our findings. Our study provides novel observations of laying hen

plasma 25-OH-D_3 levels relative to dietary vitamin D_3 supplementation that can be

valuable for the poultry industry to consider with vitamin D status. The plasma 24,25-

$(OH)_2$-D_3 data is the most exciting finding of our study. Further understanding of the

asymptotic relationship of the super-dose levels with plasma 24,25-$(OH)_2$-D_3

concentrations can open new knowledge about 24,25-$(OH)_2$-D_3's value as a biomarker

for vitamin D metabolism.

There were a few limitations with this study that were realized when data was

collected. We should have investigated kidney histopathology of these hens because

soft-tissue calcification or renal kidney failure could result from the hens reaching

vitamin D toxicity (25, 26). However, Mattila et al. (10) did not observe any pathological

issues of kidneys from 67-wk old hens fed 15,000 IU D_3/kg of feed. The smaller sample

sizes and missing treatment groups from the qPCR results were because some tissue

samples would not yield RNA for cDNA synthesis, even after multiple extraction

attempts. It is important to note that the CYP27C1 and CYP24A1 genes may not

encode vitamin D-related hydroxylases (27) (Table 1 footnote). We also stored the egg

yolk in a refrigerator for about a year before the yolk was freeze-dried; however, our

findings are similar to Wen et al. (11) to hint towards minimal vitamin D_3 degradation

(**Table 3**). This could indicate how stable vitamin D_3 is when it is stored in cold, dark

conditions.

353   Our study exhibits that feeding super-doses of dietary vitamin D_3 to aged laying

354 hens increases their plasma and egg yolk vitamin D_3 metabolite levels. Importantly,

355 there is a possible metabolic limit of 24-hydroxylation to remove excess circulating

356 vitamin D_3. Investigating 24-hydroxylation mechanisms will be important to

357 understanding vitamin D supplementation impacts in geriatric animals for improving

358 bone health and vitamin D metabolism in older humans.

Acknowledgments

We thank Gavin Conant, Peter Ferket, Matt Koci, and Shannon Madden for their

comments and edits to the initial manuscript; Taylor Jones for her help with the RNA

extraction, cDNA synthesis, and qPCR experiments; David Dickey for statistical

assistance; Jeff Hall and Zach Spivey for their help with taking care of the hens and

collecting samples; Liza Lentz of the Environmental and Agricultural Testing Service

laboratory, Department of Crop and Soil Sciences, at North Carolina State University

who performed the mineral composition experiments and analyses; and John

Rathmacher of Heartland Assays for quantifying the plasma and egg yolk vitamin D_3.

The authors' responsibilities were as follows-MFW and KAL: designed research; MFW,

DDH, NCT, and KAL: conducted research; MFW and KAL: analyzed data; MFW and

KAL: wrote manuscript; MFW and KAL: prepared experimental diets for study; MFW

and PMP: prepared samples for ashing; MFW and DDH: prepared and shipped plasma

and egg yolk samples to Heartland Assays; MFW and KAL: had primary responsibility

for final content; and all authors have read and approved the final manuscript.

Conflict of Interest: All authors report no conflicts of interest.

Data availability

378 Data described in the current study will be made available from the corresponding

379 author upon request.

References

1.   Jones G. Pharmacokinetics of vitamin D toxicity. The American Journal of Clinical Nutrition 2008;88(2):582S-6S. doi: 10.1093/ajcn/88.2.582S.

2.   Marcinowska-Suchowierska E, Kupisz-Urbańska M, Łukaszkiewicz J, Płudowski P, Jones G. Vitamin D Toxicity–A Clinical Perspective. Frontiers in Endocrinology 2018;9. doi: 10.3389/fendo.2018.00550.

3.   Holick MF. Vitamin D deficiency. New England Journal of Medicine 2007;357(3):266-81.

4.   Tang BM, Eslick GD, Nowson C, Smith C, Bensoussan A. Use of calcium or calcium in combination with vitamin D supplementation to prevent fractures and bone loss in people aged 50 years and older: a meta-analysis. The Lancet 2007;370(9588):657-66.

5.   Bischoff-Ferrari HA, Willett WC, Wong JB, Stuck AE, Staehelin HB, Orav EJ, Thoma A, Kiel DP, Henschkowski J. Prevention of nonvertebral fractures with oral vitamin D and dose dependency: a meta-analysis of randomized controlled trials. Archives of internal medicine 2009;169(6):551-61.

6.   Bischoff-Ferrari HA, Willett WC, Orav EJ, Lips P, Meunier PJ, Lyons RA, Flicker L, Wark J, Jackson RD, Cauley JA. A pooled analysis of vitamin D dose requirements for fracture prevention. New England Journal of Medicine 2012;367(1):40-9.

7.   Klontz KC, Acheson DW. Dietary supplement–induced vitamin D intoxication. New England Journal of Medicine 2007;357(3):308-9.

8.   Binkley N, Gemar D, Engelke J, Gangnon R, Ramamurthy R, Krueger D, Drezner M. Evaluation of ergocalciferol or cholecalciferol dosing, 1,600 IU daily or 50,000 IU monthly in older adults. The Journal of Clinical Endocrinology & Metabolism 2011;96(4):981-8.

9.   Mattila P, Rokka T, Könkö K, Valaja J, Rossow L, Ryhänen E-L. Effect of cholecalciferol-enriched hen feed on egg quality. Journal of agricultural and food chemistry 2003;51(1):283-7.

10.   Mattila P, Valaja J, Rossow L, Venäläinen E, Tupasela T. Effect of vitamin D2- and D3-enriched diets on egg vitamin D content, production, and bird condition during an entire production period. Poultry Sci 2004;83(3):433-40.

11.   Wen J, Livingston K, Persia M. Effect of high concentrations of dietary vitamin D3 on pullet and laying hen performance, skeleton health, eggshell quality, and yolk vitamin D3 content when fed to W36 laying hens from day of hatch until 68 wk of age. Poultry Sci 2019;98(12):6713-20.

12.   Adhikari R, White D, House JD, Kim WK. Effects of additional dosage of vitamin D3, vitamin D2, and 25-hydroxyvitamin D3 on calcium and phosphorus utilization, egg quality and bone mineralization in laying hens. Poultry Sci 2020;99(1):364-73. doi: https://doi.org/10.3382/ps/pez502.

13.   Whitehead CC. Overview of bone biology in the egg-laying hen. Poultry Sci 2004;83(2):193-9. doi: 10.1093/ps/83.2.193.

14.   DeLuca HF. Overview of general physiologic features and functions of vitamin D. The American Journal of Clinical Nutrition 2004;80(6):1689S-96S. doi: 10.1093/ajcn/80.6.1689S.

15.   Whitehead C, Fleming R. Osteoporosis in cage layers. Poultry Sci 2000;79(7):1033-41.

16.   Council NR. Nutrient requirements of poultry: 1994: National Academies Press, 1994.

17.   Mattila P, Lehikoinen K, Kiiskinen T, Piironen V. Cholecalciferol and 25-hydroxycholecalciferol content of chicken egg yolk as affected by the cholecalciferol content of feed. Journal of Agricultural Food chemistry 1999;47(10):4089-92.

18.   van Leeuwen JPTM, van den Bemd G-JCM, van Driel M, Buurman CJ, Pols HAP. 24,25-Dihydroxyvitamin D3 and bone metabolism. Steroids 2001;66(3):375-80. doi: https://doi.org/10.1016/S0039-128X(00)00155-0.

19.   Tanaka Y, DeLuca H. Stimulation of 24, 25-Dihydroxyvitamin D3 Production by 1, 25-Dihydroxyvitamin D3. Science 1974;183(4130):1198-200.

20.   Haussler MR, Haussler CA, Bartik L, Whitfield GK, Hsieh J-C, Slater S, Jurutka PW. Vitamin D receptor: molecular signaling and actions of nutritional ligands in disease prevention. Nutrition reviews 2008;66(suppl_2):S98-S112.

21.   Travel A, Nys Y, Bain M. Effect of hen age, moult, laying environment and egg storage on egg quality. Edtion ed. Improving the safety and quality of eggs and egg products: Elsevier, 2011:300-29.

22.   Ameenuddin S, Sunde M, DeLuca H, Cook M. Excessive cholecalciferol in a layers diet: decline in some aspects of reproductive performance and increased bone mineralisation of progeny. Brit Poultry Sci 1986;27(4):671-7.

23.   Duffy SK, Rajauria G, Clarke LC, Kelly AK, Cashman KD, O'Doherty JV. The potential of cholecalciferol and 25-hydroxyvitamin D3 enriched diets in laying hens, to improve egg vitamin D content and antioxidant availability. Innovative Food Science & Emerging Technologies 2017;44:109-16.

24.   Bikle D. Vitamin D: Production, metabolism, and mechanisms of action. Edtion ed. Endotext [Internet]: MDText. com, Inc., 2017.

25.   Vieth R. Vitamin D toxicity, policy, and science. Journal of Bone and Mineral Research 2007;22(S2):V64-V8.

26.   Morrissey RL, Cohn RM, Empson Jr R, Greene HL, Taunton OD, Ziporin ZZ. Relative Toxicity and Metabolic Effects of Cholecalciferol and 25-Hydroxycholecalciferol in Chicks. The Journal of nutrition 1977;107(6):1027-34.

27.   Sinclair-Black M, Garcia RA, Ellestad LE. Physiological regulation of calcium and phosphorus utilization in laying hens. Frontiers in Physiology 2023;14. doi: 10.3389/fphys.2023.1112499.

Figure Legends

Figure 1. Study design. A) Forty-eight 68-wk Hy-Line Brown hens were used in the study and were randomly assorted into one of six treatment groups and fed the diet with the corresponding dietary vitamin D_3 supplementation level. **B)** Hens were individually housed in cages of a two-leveled battery cage in a randomized complete block design (n = 8/diet). **C)** The experimental timeline in which hens were fed the same basal diet (400 IU D_3/kg) for four weeks as a washout period (W-4). Hens were started on the experimental diets at week 0 (W0), eggs were collected daily, and weekly body weight was taken until the end of the study. On weeks 0, 3, 6, and 10, blood was collected from the brachial wing vein from all hens to measure ionized blood calcium using an i-STAT™ blood analyzer. Blood was spun down and plasma was collected to measure vitamin D_3 metabolite levels. The study ended on week 10 and all hens were sacrificed and sample tissues were collected.

Figure 2. Vitamin D_3 metabolite plasma concentrations of 78-wk Hy-Line Brown laying hens fed different levels of dietary vitamin D_3. A) Cholecalciferol (Vitamin D_3; 400 and 800 IU treatment concentrations were not determined) **B)** 25-hydroxycholecalciferol (25-OH-D_3) **C)** 24,25-dihydroxycholecalciferol (24,25-(OH)$_2$-D_3) **D)** Ratio of 24,25-(OH)$_2$-D_3/25-OH-D_3 presented as a percentage. Blue squares denote standard NRC range vitamin D_3 levels (400 and 800 IU D_3/kg) in diet (n = 4) and red diamonds denote super-dose levels of vitamin D_3 (14,000, 20,000, and 36,000 IU D_3/kg) in diet (n = 4). Samples were reported as means ± SEM. Samples with common letters were not statistically different from each other (General linear models, p < 0.0001).

Figure 3. Egg yolk vitamin D_3 metabolite concentrations from 78-wk Hy-Line Brown laying hens fed different levels of dietary vitamin D_3. A) Cholecalciferol (Vitamin D_3) **B)** 25-hydroxycholecalciferol (25-OH-D_3). Blue squares denote standard NRC range vitamin D_3 levels (400 IU D_3/kg) in diet (n = 4) and red diamonds denote super-dose levels of vitamin D_3 (14,000 and 36,000 IU D_3/kg) in diet (n = 4). Samples were reported as means ± SEM; however, error bar values were narrow and overlapped by the marker for each sample. Samples with common letters were not statistically different from each other (General linear models, p < 0.0001).

Figure 5. Relative gene expression of vitamin D receptor (VDR) in duodenum, ileum, liver, and kidney of 78-wk Hy-Line Brown laying hens fed different levels of dietary vitamin D_3. A) Duodenal VDR (n = 2-6) **B)** Ileal VDR (n = 2-5) **C)** Liver VDR (n = 2-4) **D)** Kidney VDR (n = 2-4). Tissues were analyzed using qPCR normalized against glyceraldehyde 3-phosphate dehydrogenase (GAPDH; housekeeping gene) expression. Blue bars with dots denote standard NRC range vitamin D_3 levels in diet and red bars denote super-dose levels of vitamin D_3 in diet. All samples ran in triplicates and reported

as means ± SEM. Bars with common letters were not statistically different from each other (General linear models, $p < 0.05$).

Figure 6. Dietary super-doses of vitamin D$_3$ fed to aged laying hens causes drastic increase in plasma and egg yolk vitamin D metabolites. Egg yolk vitamin D$_3$ levels were strongly correlated to the dietary levels of vitamin D$_3$ fed to the hens. Egg yolk 25-hydroxycholecalciferol (25-OH-D$_3$) levels were also dependent on dietary vitamin D$_3$; however, 25-OH-D$_3$ increased at a lower rate. For plasma vitamin D metabolites, vitamin D$_3$ and 25-OH-D$_3$ concentrations increased relative to dietary vitamin D$_3$ levels fed to the hens. Plasma 24,25-dihydroxycholecalciferol [24,25-(OH)$_2$] levels were also dependent on dietary vitamin D$_3$.

Supplementary Figure 1. Percent by weight of bone calcium and phosphorus from 78-wk Hy-Line Brown laying hens fed different levels of dietary vitamin D$_3$. A) Tibia and humerus calcium **B)** Tibia and humerus phosphorus. Blue squares with lines denote tibia and orange triangles with dashed lines denote humerus (n = 7). Humerus contain more calcium and phosphorus than tibia. Line graphs were expressed as means ± SEM (General linear models, $p < 0.05$).

Supplementary Figure 2. Relative gene expression of kidney vitamin D hydroxylases of 78-wk Hy-Line Brown laying hens fed different levels of dietary vitamin D$_3$. A) Kidney 24-hydroxylase (CYP24A1; n = 2-4) **B)** Kidney 1α-hydroxylase (CYP27C1; n = 2-5). Relative expression of vitamin D hydroxylase genes were compared to glyceraldehyde 3-phosphate dehydrogenase (GAPDH; housekeeping gene) expression. Blue bars denote standard NRC vitamin D$_3$ level range in diet and red bars denote super-dose levels of vitamin D$_3$ in diet. All samples were run in triplicates and reported as means ± SEM. Bars with common letters were not statistically different from each other (General linear models, $p < 0.05$).

Ingredient Name	%
Corn	63.40
Soybean meal, 46% crude protein	20.60
Calcium carbonate	9.20
Poultry fat	2.33
Dicalcium phosphate[1]	1.95
L-Lysine-hydrochloride, 78.8%	1.01
Sodium bicarbonate	0.57
Vitamin premix[2]	0.25
Mineral premix[3]	0.20

Table 1. Ingredient composition of the experimental basal diet

Choline chloride, 60% choline	0.20
DL-Methionine, 99%	0.14
Salt	0.10
Selenium premix[4]	0.05

---

[1]Dicalcium phosphate contains 19.79% calcium, 17.91% phosphorus, and 17.73% available phosphorus.

[2]Provided as milligrams per kilogram of diet: 125 mg ethoxyquin; 25 mg niacinamide; 10 mg calcium pantothenate; 6.7 mg DL-α-tocopherol; 3.6 mg riboflavin; 3 mg pyridoxine hydrochloride; 1.8 mg thiamine hydrochloride; 0.55 mg folic acid; 0.55 mg menadione sodium bisulfite; 0.516 mg retinol acetate; 0.15 mg biotin; 0.01 mg cyanocobalamin.

[3]Trace minerals provided per kg of premix: 60 g manganese sulfate; 60 g zinc sulfate; 40 g iron sulfate; 5 g copper sulfate; 1.25 g calcium iodate.

[4]Selenium premix provided selenium at 0.3 mg/kg of diet.

Table 2. Primer sequences for quantitative real-time PCR

Gene	Orientation	Primer Sequence (5' – 3')	Size (bp)	Accession #
VDR	Forward	TGCCTCCAGTCTGGCATCTC	297	NM_205098.1
	Reverse	GGTGATTTTGCAGTCCCCGT		
CYP27C1	Forward	ATGATTGGCGTCCCCTTCAG	177	XM_422077.4[1]
	Reverse	TCCACGCTTTCACTCACACA		
CYP24A1	Forward	AAACCCTGGAAAGCCTATCG	133	NM_204979.1[2]
	Reverse	CCAGTTTCACCACCTCCTTG		
GAPDH	Forward	TGTTGTTGACCTGACCTGCC	291	NM_204305.1
	Reverse	CTGGCTCACTCCTTGGATGC		

[1]The NCBI record for CYP27C1 was removed due to standard genome annotation processing. However, in our study, the CYP27C1 primers were made and used in 2019 when the record was available.

[2]The NCBI record for CYP24A1 was removed because of insufficient support for the transcript and protein. In our study, the CYP24A1 primers were made and used in 2019 when the record was available.

Table 3. Comparison of egg yolk vitamin D_3 from eggs laid by hens fed different dietary levels of vitamin D_3 in this study compared to Wen et al. 2019 study (19)

This study[1]	Wen et al. 2019 (19)[2]
IU/egg (Dietary treatment IU D_3/kg)	
5.13 (400)	12.6 (1,681)
199.7 (14,000)	214.3 (18,348)
605.9 (36,000)	435.5 (35,014)

[1]Values were calculated by converting the egg yolk vitamin D_3 concentration (ng/g) to IU and multiplying by 15g (amount used for LC-MS/MS).
[2]These values were reported in Figure 1 for (19).

Figure 1. Study design. A) Forty-eight 68-wk Hy-Line Brown hens were used in the study and were randomly assorted into one of six treatment groups and fed the diet with the corresponding dietary vitamin D_3 supplementation level. **B)** Hens were individually housed in cages of a two-leveled battery cage in a randomized complete block design (n = 8/diet). **C)** The experimental timeline in which hens were fed the same basal diet (400 IU D_3/kg) for four weeks as a washout period (W-4). Hens were started on the experimental diets at week 0 (W0), eggs were collected daily, and weekly body weight was taken until the end of the study. On week 0, 3, 6, and 10, blood was collected from the brachial (wing) vein from all hens to measure ionized blood calcium using an i-STAT™ blood analyzer. Blood was spun down and plasma was collected to measure vitamin D metabolite levels. The study ended on week 10 and all hens were sacrificed and sample tissues were collected.

Figure 2. Vitamin D_3 metabolite plasma concentrations of 78-wk Hy-Line Brown laying hens fed different levels of dietary vitamin D_3. A) Cholecalciferol (Vitamin D_3; 400 and 800 IU treatment concentrations were not determined) **B)** 25-hydroxycholecalciferol (25-OH-D_3) **C)** 24,25-dihydroxycholecalciferol (24,25-(OH)$_2$-D_3) **D)** Ratio of 24,25-(OH)$_2$-D_3/25-OH-D_3 presented as a percentage. Blue squares denote standard NRC range D_3 levels (400 and 800 IU D_3/kg) in diet (n = 4) and red diamonds denote super-dose levels of D_3 (14,000, 20,000, and 36,000 IU D_3/kg) in diet (n = 4). ND = not detected or below limit of detection. Samples reported as means ± SEM. Samples with common letters were not statistically different from each other (General linear models, p < 0.0001).

Figure 3. Egg yolk vitamin D$_3$ metabolite concentrations from 78-wk Hy-Line Brown laying hens fed different levels of dietary vitamin D$_3$. A) Cholecalciferol (Vitamin D$_3$) **B)** 25-hydroxycholecalciferol (25-OH-D$_3$). Blue squares denote standard NRC range D$_3$ levels (400 IU D$_3$/kg) in diet (n = 4) and red diamonds denote super-dose levels of D$_3$ (14,000 and 36,000 IU D$_3$/kg) in diet (n = 4). Samples were reported as means ± SEM; however, error bars values were narrow and overlapped by the marker for each sample. Samples with common letters were not statistically different from each other (General linear models, p < 0.0001).

Figure 4. Association between plasma and egg yolk vitamin D$_3$ metabolite concentrations from 78-wk Hy-Line Brown laying hens fed different levels of dietary vitamin D$_3$. A) Cholecalciferol (Vitamin D$_3$) **B)** 25-hydroxycholecalciferol (25-OH-D$_3$). Blue squares denote standard NRC range vitamin D$_3$ level (400 IU D$_3$/kg, n = 4) in diet; orange (14,000 IU D$_3$/kg, n = 4) and red diamonds (36,000 IU D$_3$/kg, n = 4) denote super-dose levels of vitamin D$_3$ in diet (Pearson correlation).

Figure 5. Relative gene expression of vitamin D receptor (VDR) in duodenum, ileum, liver, and kidney of 78-wk Hy-Line Brown laying hens fed different levels of dietary vitamin D$_3$. A) Duodenal VDR (n = 2-6) **B)** Ileal VDR (n = 2-5) **C)** Liver VDR (n = 2-4) **D)** Kidney VDR (n = 2-4). Relative expression of the VDR gene was compared to glyceraldehyde phosphate dehydrogenase (GAPDH; housekeeping gene) expression. Blue bars denote standard NRC range vitamin D$_3$ levels in diet and red bars denote super-dose levels of vitamin D$_3$ in diet. All samples ran in triplicates and reported as means ± SEM. ND = not detected or below limit of detection. Bars with common letters were not statistically different from each other (General linear models, $p < 0.05$).

Figure 6. Dietary super-doses of vitamin D$_3$ fed to aged laying hens increases plasma and egg yolk vitamin D metabolites. Egg yolk vitamin D$_3$ levels were strongly correlated to the dietary levels of vitamin D$_3$ fed to the hens. Egg yolk 25-hydroxycholecalciferol (25-OH-D$_3$) levels were also dependent on dietary vitamin D$_3$; however, 25-OH-D$_3$ increased at a lower rate. For plasma vitamin D metabolites, vitamin D$_3$ and 25-OH-D$_3$ concentrations increased relative to dietary vitamin D$_3$ levels fed to the hens. Plasma 24,25-dihydroxycholecalciferol [24,25-(OH)$_2$] levels were also dependent on dietary vitamin D$_3$, but rate of increase was low.