1 **Super-doses of dietary vitamin D_3 intake in aged laying hens**

2 **illustrates limitation of 24,25-dihydroxycholecalciferol conversion**

3 Matthew F. Warren[1], Pete M. Pitman[2], Dellila D. Hodgson[1], and Kimberly A. Livingston[1]

4 [1]Prestage Department of Poultry Science, North Carolina State University, Raleigh, NC,
5 USA
6
7 [2]Department of Biological Sciences, North Carolina State University, Raleigh, NC, USA
8

9 **Sources of support:** Funding for this research was supported by the US Poultry & Egg
10 Association (KAL).
11
12 **Conflict of Interest and Funding Disclosure:** All authors report no conflicts of interest.
13

14 Matthew F. Warren, Department of Bacteriology, University of Wisconsin-Madison, 1550
15 Linden Drive, Madison, WI 53706-1205, (559) 816-0639, mwarren8@wisc.edu
16

Word count: 4914 (Introduction – Discussion)

Running title: Dietary vitamin D_3 super-dose intake in aged laying hens

Abbreviations used: 1α-OHase, 1α-hydroxylase; 1,25-$(OH)_2$-D_3, 1,25-

dihydroxycholecalciferol; 24-OHase, 24-hydroxylase; 25-OHase, 25-hydroxylase; 24,25-

$(OH)_2$-D_3, 24,25-dihydroxycholecalciferol; 25-OH-D_3, 25-hydroxycholecalciferol; VDR,

vitamin D receptor

ABSRACT

Background: Humans take vitamin D supplements to reduce risk of vitamin D

deficiency and reduce the risk of osteoporosis. However, it is unclear how dietary super-

doses (10,000x greater than requirement) can affect vitamin D status in aged animals.

Aged laying hens could be a model to compare with women in peri- or postmenopausal

stages of life. The hens' bone health is physiologically taxed from egg production and

they are highly susceptible to osteoporosis.

Objective: We investigated dietary super-dose impacts of cholecalciferol (vitamin D_3)

on vitamin D status in aged laying hens in production.

Methods: Forty-eight 68 wk old Hy-Line Brown laying hens were individually housed in

cages with eight hens per dietary treatment for eleven weeks. Hens were randomly

assigned to one of six groups of dietary vitamin D_3 supplementation and fed *ad libitum*.

Supplementation levels were 400, 800, 7,400, 14,000, 20,000, and 36,000 IU D_3/kg of

feed. At termination of the study, all hens were euthanized and we collected tissue

samples and their feces. Tissue calcium, plasma and egg yolk vitamin D metabolites,

and gene expression levels were measured.

Results: We observed that increasing dietary vitamin D_3 increased plasma vitamin D_3,

25-hydroxycholecalciferol, and 24,25-dihydroxycholecalciferol concentrations ($p <$

0.0001 for all 3 metabolites). Also, egg yolk vitamin D_3 and 25-hydroxycholecalciferol

showed a similar effect like plasma vitamin D metabolites ($p < 0.0001$ for both

metabolites). We also observed super-dose fed hens had decreased kidney 24-

hydroxylase expression ($p = 0.0006$).

Conclusions: Dietary super-doses of vitamin D_3 led to greater plasma and egg yolk

vitamin D levels to show that aged laying hens can deposit excess vitamin D in egg

yolk. We suggest future research should explore how 24-hydroxylation mechanisms are

affected by vitamin D supplementation. Further understanding of 24-hydroxylation can

help ascertain ways to reduce risk of vitamin D toxicity.

Keywords: dietary vitamin D_3, aged laying hen, 25-hydroxycholecalciferol, 24,25-

dihydroxycholecalciferol, super-dose, 24-hydroxylase, 25-hydroxylase, 1α-hydroxylase,

egg yolk

Introduction

Vitamin D deficiency is directly linked to osteoporosis, with research

demonstrating that increasing vitamin D intake through diet or supplements is a means

to reduce risk and prevent vitamin D deficiency (1, 2). Older women in menopausal or

post-menopausal stages of life contend with increased demands of nutrient absorption

with onset of hormonal changes that are an effect of menopause (3-5). Reduction of

physiological turnover activity as an added effect of aging is another demand which

makes vitamin D deficiency more prevalent in older women (6). A recent study reported

how prevalent levels of vitamin D deficiency was in middle-aged and elderly people from

Lanzhou population in China (7). This population was regularly exposed to sunlight;

however, other factors such as smoking were correlated to contribute to vitamin D

deficiency. Culmination of vitamin D deficiency and aging can lead to increased

osteoporotic susceptibility and underlying potential health risks associated with bone

vulnerability (8).

Older people tend to take vitamin D supplements to increase or maintain vitamin

D levels (9). While vitamin D toxicity is uncommon, vitamin D supplements and

overfortified foods are the only known means of reaching intoxication levels (10, 11). A

study showed how adult men in winter months were given daily doses of D_3, ranging

from 0 - 250 μg for about 20 wk (12). The men had increased serum 25-

hydroxycholecalciferol (25-OH-D_3), even though their 25-OH-D_3 concentration was in a

safe range. There was also a study involving adult men and women given a single oral

dose that ranged from 1.5 - 10 μg 25-OH-D_3 /kg of body weight (13). Serum 25-OH-D_3

concentration of these adults quickly increased within 4-6 h. However, the half-life of 25-

OH-D$_3$ was long (~22 d) and 25-OH-D$_3$ can remain in circulation and potentially reach

toxic levels (13). Young laying hens fed diets with 15,000 IU D$_3$ /kg of feed for 48 wk did

not show signs of tissue pathologies from toxicity (14). There is value with assessing

how super-doses of dietary D$_3$ affects plasma vitamin D metabolite levels in older laying

hens. We hypothesized that older laying hens would have reduced vitamin D

metabolism that led to reduced circulating levels of hydroxylated vitamin D metabolites.

Understanding how high levels of dietary doses of D$_3$ would be valuable information for

ascertaining how overfortified foods can potentially affect vitamin D metabolism in older

animals.

Dietary vitamin D supplementation is important for laying hens in production

because their bone health is physiologically taxed from egg production (15, 16). Laying

hens in commercial farms are fed diets with supplemental vitamin D$_3$ beyond NRC

requirements (17). This ensures the hens are able to lay eggs and maintain adequate

Ca absorption for eggshell formation and importantly, bone mineralization. There are a

few studies which investigated how high dietary vitamin D supplementation affected

laying hen production and the metabolic implications pertaining to vitamin D status (14,

18, 19). There is need for studies that examined how super-doses of vitamin D$_3$ affect

gene expression and how vitamin D-related genes correlate to vitamin D metabolism in

laying hens. Understanding how gene expression and metabolic indicators, like blood

vitamin D metabolite levels, are affected by super-doses of dietary vitamin D$_3$ in aged

laying hens which has implications on the poultry industry, but also with human health.

We expect aged laying hens fed super-doses of dietary vitamin D$_3$ to have upregulated

expression of vitamin D-related genes, resulting in improved vitamin D metabolism.

Such findings have implications to justify increasing vitamin D intake from fortification in

food for older women to improve their vitamin D status.

Our study examined dietary vitamin D_3 super-dose effects on plasma and egg

yolk vitamin D_3 metabolites and relative gene expression of vitamin D-related genes in

aged laying hens in production. We fed hens diets containing 400, 800, 7,400, 14,000,

20,000, and 36,000 IU D_3/kg of feed to ascertain vitamin D_3 supplementation impacts.

Hens consuming diets with vitamin D_3 greater than 10,000 IU/kg were expected to have

increased plasma 24,25-dihydroxycholecalciferol (24,25-$(OH)_2$-D_3) because 24,25-

$(OH)_2$-D_3 is an inactive form to indicate they reached vitamin D saturation. Hens

consuming super-doses of D_3 should also lay eggs with increased vitamin D_3 content

because they would deposit excess D_3 into the egg yolk (14).

Methods

Birds and housing

The hens used in our study were from North Carolina State University's

maintained poultry flock. Forty-eight 68 wk old Hy-Line Brown laying hens were housed

at North Carolina State University, Raleigh, NC. Hens were individually housed in cages

between two two-level battery cages with eight birds per treatment. The experimental

design was a randomized complete block design with six levels of dietary vitamin D

supplementation blocked by cage level. Vitamin D_3 supplementation levels were

formulated to be 250, 500, 1,500, 15,000, 30,000, and 60,000 IU D_3/kg of feed, but the

analyzed vitamin D_3 in feed (and dietary treatments in this study are referred as) were

400, 800, 7,400, 14,000, 20,000, and 36,000 IU/kg of feed (**Tables 1 and 2**). Prior to the

start of the experiment, all hens were fed the same diet (400 IU D_3/kg of feed) for one

month as a washout period. Hens were fed *ad libitum* along with water. North Carolina

State University's Institutional Animal Care and Use Committee approved all methods

for this study, protocol ID number: 18-093-A.

Sample Collection

Eggs were collected every morning and stored at 7°C for egg quality analyses.

The first egg laid for the week by each hen was selected for egg yolk collection. Eggs

were cracked open in a dim-lighted room to reduce photodegradative impacts of light on

vitamin D in yolk. Yolk was separated from albumen and placed in a small plastic

container wrapped in aluminum foil and stored at 4°C for a year until they were freeze-

dried using a freeze-dryer (FreeZone 6 Liter Benchtop Freeze Dry System, Labconco,

Kansas City, MO). All 48 birds were bled to have their plasma vitamin D_3 metabolites

measured. All birds were sacrificed by cervical dislocation and tissue samples were

collected from 43 birds (minimum of seven birds per treatment) due to time constraints.

Liver and kidney were collected and stored in RNAlater at -20°C until RNA extractions

were performed.

Calcium and phosphorus content of various sites

Eggshells from week 0, 3, 4, 6, and 9 were washed with warm water to remove

shell membrane and dried for 48 h at room temperature. Dried eggshells were pre-

weighed and further dried at 68°C for 72 h using a dry oven (Blue M, Atlanta, GA) and

weighed again. Eggshells were crushed into fine powder and subjected to digestion to

measure Ca composition of eggshells. Feces and ileal digesta were also subjected to

same steps as eggshells. Dried samples were weighed and then placed in a muffle

furnace at 500°C overnight to ash samples. Ashed samples were dehydrated in 2 mL of

distilled water and 4 mL of 6 N hydrochloric acid. The resulting sample was mixed and

heated to warm to touch. Heated solution was poured into volumetric flask and

deionized water was added to 50 mL. The flask was inverted 12 times to mix and the

resulting solution was filtered using #40 filter paper into 15 mL centrifuge tubes for

analysis. Ca and P was measured by inductively coupled plasma optical emission

spectrometry.

Bones were wrapped in petroleum ether moistened cheesecloth and placed in a

desiccator for 72 h to extract fat from bones. Fat-extracted bones were pre-weighed and

dried for 24 h at 100°C to evaporate petroleum ether residues. Fat- and moisture-free

bones were weighed and ashed using same methods as eggshell, feces, and ileal

digesta for Ca and P composition and measured by inductively coupled plasma atomic

emission spectroscopy.

RNA extraction and qPCR

Total mRNA was extracted from duodenum, ileum, liver, and kidney using

Qiagen's RNeasy Mini Kit (Germantown, MD). The extracted RNA was diluted and

normalized to ~200 ng/µL for liver and 60 ng/µL for duodenum, ileum, and kidney. The

aforementioned tissues' RNA was reverse transcribed to complementary DNA (cDNA)

using Applied Biosystems' High-Capacity cDNA Reverse Transcription Kit

(ThermoFisher Scientific, Waltham, MA) and their recommended steps to make a 20 µL

working solution. Cycling procedure for reverse transcription started with 25°C for 10

min, 37°C for 120 min, 85°C for 5 mins, then held at 5°C indefinitely until storage or use.

Genes amplified for quantitative real time PCR (qPCR) were vitamin D receptor

(VDR), 1α-hydroxylase (1α-OHase), 25-hydroxylase (25-OHase), 24-hydroxylase (24-

OHase), and glyceraldehyde 3-phosphate dehydrogenase (GAPDH) as the

housekeeping gene (**Table 3**). qPCR was conducted using PowerUP SYBR Master Mix

(Life Technologies, Grand Island, NY) following manufacturer's protocol and using

Applied Biosystems StepOnePlus Real-Time PCR System (Carlsbad, CA). Cycling

procedure started with 95°C for 10 min then 40 cycles of 95°C for 15 s for denaturing

and 15 s at 60°C for annealing. All samples were run in triplicates.

Vitamin D metabolites

Plasma from specific-fed hens from 400, 800, 14,000, and 36,000 IU D_3 groups

were respectively pooled (n = 4/treatment) and sent to Heartland Assays (Ames, IA) for

measuring D_3, 25-OH-D_3, and 24,25-dihydroxycholecalciferol [24,25-$(OH)_2$-D_3, inactive

form of D_3] using LC/MS/MS. Freeze-dried egg yolk from the same hens were pooled

(15 g per sample) like plasma, but only samples from 400, 14,000, and 36,000 IU D_3

groups (n = 4/treatment) were analyzed for D_3 and 25-OH-D_3 from Heartland Assays.

Statistical analysis

We conducted statistical analyses using general linear models using SAS 9.4®

for all statistical tests and Tukey-Kramer test was used for multiple comparisons for

differences between dietary treatments. We utilized repeated measures to account for

temporal effects of weekly body weight, feed intake, egg production, eggshell quality,

and ionized blood Ca. Dietary vitamin D_3 level was the independent variable for Ca and

P composition, plasma and egg yolk vitamin D metabolite concentrations, and gene

expression data. We did not observe a cage effect in any analysis so the blocking

variable was omitted from all statistical tests. All mRNA relative expressions were

normalized using $2^{-C\Delta\Delta T}$ with GAPDH as a housekeeper gene. Statistical significance

was established at $p < 0.05$.

Results

Hens' production performance was not influenced by dietary vitamin D_3

To determine if dietary super-doses of vitamin D_3 affected the hens' production

value, we measured the hens' weekly body weights and egg production. Dietary super-

doses of vitamin D_3 did not affect body weight of these laying hens ($p = 0.08$; **Table S1**).

There was an interaction between dietary vitamin D_3 level over time ($F = 5.45$, $p <$

0.0001); however, there was constant feed wastage throughout the study so this effect

could be inflated. A dietary effect was observed for egg production for the entire study

duration ($F = 2.52$, $p = 0.047$; **Table S2**) with hens from 14,000 IU D_3 treatment group

having highest production rate of 86.7%. Eggshell strength and eggshell thickness were

not affected by dietary vitamin D_3 levels ($p = 0.19$, $p = 0.72$, respectively; **Table S2**).

There was a trending interaction between dietary vitamin D_3 level over time in which

there was a decrease in eggshell elasticity ($p = 0.07$).

Ionized blood calcium is affected by dietary vitamin D_3 levels

We also examined if ionized blood Ca in the hens was influenced by dietary

vitamin D_3 levels throughout the study. There was no temporal effect on ionized blood

Ca (p = 0.65). With week 0 considered as a covariate, there was a strong dietary effect

on ionized blood Ca (p = 0.002, **Table S3**). Even so, it is unclear on how dietary levels

of D_3 influenced ionized blood Ca concentration because their levels are both high and

low across standard-fed D_3 hens and super-dose-fed D_3 hens.

Fecal calcium is affected by dietary vitamin D_3 levels, but not ileal digesta or

eggshell calcium or phosphorus

We had the eggshell, ileal digesta, and fecal Ca and P measured to determine if

dietary vitamin D_3 fed to our hens would reduce the excretion or loss of Ca and P. There

was no dietary effect on eggshell Ca and P (p = 0.64 for both) and ileal digesta Ca or P

(p = 0.74 and 0.09, respectively). There was a dietary effect with fecal Ca with hens fed

14,000 IU D_3 diet had 10.6±0.8% Ca by weight in their feces, whereas all other

treatments were 8-9%, with exception of 36,000 IU D_3 fed hens which had 7.36±0.43%

fecal Ca (p = 0.03, **Table S4**). There was no dietary effect on fecal P (p = 0.76).

Feeding hens diets containing super-dose levels of vitamin D_3 did not affect P in all

measured samples and hens fed the highest level of D_3 had the lowest Ca excretion in

feces.

Humerus is more calcium and phosphorus dense than tibia

We assessed if dietary vitamin D_3 would improve bone Ca and P in the hens.

However, no dietary effects of vitamin D supplementation were observed on bone Ca or

P (p = 0.79, p = 0.63, respectively). Humerus bones have more Ca and P than tibia

bones (p = 0.020, p = 0.015, respectively; **Figure S1**). We showed that feeding aged

laying hens super-dose levels of vitamin D_3 does not affect Ca and P of humerus and

tibia.

Dietary super-dosage levels of vitamin D increased plasma and egg yolk vitamin

D_3 metabolites

We had plasma and egg yolk vitamin D_3 metabolites measured by LC-MS/MS to

determine if dietary vitamin D_3 affected plasma and egg yolk concentrations. Plasma

vitamin D_3 was below readable range for 400 and 800 IU treatments and those samples

were considered zero and included in statistics to account for model building. All vitamin

D_3 metabolite data exhibited heteroskedasticity and were transformed using natural

logarithm function. Transformed data exhibited linear and homoskedastic relationships

and were used for statistics.

There was a significant increase in plasma concentration of D_3, 25-OH-D_3, and

24,25-$(OH)_2$-D_3 for hens fed dietary super-dose levels of vitamin D_3 (**D_3**: $F = 27.2$, $R^2 =$

0.86, p = 0.0002; **25-OH-D_3**: $F = 495.1$, $R^2 = 0.99$, p < 0.0001; **24,25-OH-D_3**: $F = 239.4$,

$R^2 = 0.98$, p < 0.0001). Although plasma D_3 concentration could not be read for 400 and

800 IU treatments, D_3 concentration had a strong positive correlation with 25-OH-D_3 and

24,25-$(OH)_2$-D_3 concentrations ($r = 0.95$, p < 0.0001; $r = 0.92$, p < 0.0001; respectively).

Plasma D_3 and 25-OH-D_3 had similar concentration values when dietary vitamin D_3

increased with both metabolites having ~85 ng/mL at 36,000 IU treatment (**Figures 1A**

and 1B). Although plasma 24,25-$(OH)_2$-D_3 concentration was statistically affected by

dietary treatment, its rate of increase relative to dietary treatment, compared to the D_3

and 25-OH-D_3, was much lower with 17.4 ng/mL at 36,000 IU (**Figure 1C**). The

percentage ratio of 24,25-$(OH)_2$-D_3 to 25-OH-D_3 increased as dietary D_3 levels

increased and reached an asymptote at the super-dose levels ($F = 18.8$, $R^2 = 0.83$, $p <$

0.0001; **Figure 1D**). The percentage ranged from 8.7% to 20.5% with all super-dose fed

hens having a similar ratio of about 20%.

Egg yolk D_3 increased drastically as hens' dietary D_3 intake increased ($F = 537.8$,

$p < 0.0001$; **Figures 2A and 2B**). Egg yolk 25-OH-D_3 was also significantly increased in

concentration as hens' dietary D_3 increased ($F = 44.8$, $p < 0.0001$), but the rate of

increase was much lower compared to egg yolk D_3 (D_3 ranged from 8.55 – 1009.9 ng/g;

25-OH-D_3 ranged from 3.48 – 51.6 ng/g). Yolk D_3 was greatly positively correlated with

plasma D_3 and yolk 25-OH-D_3 ($r = 0.90$, $p = 0.002$; $r = 0.96$, $p < 0.0001$; respectively).

Altogether, the plasma and egg yolk vitamin D_3 data highlights a strong dose-

dependent response towards dietary vitamin D_3 fed to aged laying hens. All measured

metabolites displayed increasing concentrations as dietary vitamin D_3 levels increased.

Dietary super-doses of D_3 intake upregulated duodenal vitamin D receptor

expression

Considering VDR is the transcription factor responsible for exerting vitamin D's

physiological effects (20), we measured VDR expression in multiple tissues to

determine if dietary D_3 levels would affect VDR expression. Hens fed higher levels of

dietary D_3 had upregulated duodenal VDR expression ($F = 3.54$, $p = 0.036$; **Figure 3A**).

There was no dietary effect on VDR expression from ileum, liver, and kidney ($p = 0.96$,

0.17, 0.32; respectively; **Figures 3B-3D**). These data suggest that duodenal VDR

expression is responsive to dietary vitamin D_3.

Dietary super-doses of D_3 decreased kidney 24-OHase expression

We examined if vitamin D_3 super-dosages would affect the gene expression of

vitamin D hydroxylase enzymes in the liver and kidney. Unexpectedly, kidney 24-OHase

expression was lower in hens fed diets with super-dose levels of D_3 ($F = 8.42$, $R^2 =$

0.73, $p = 0.0006$) with 400 and 800 IU hens having about 1.32 expression and super-

dose hens having 0.80 - 1.00 expression (**Figure 4A**). There was no dietary effect on

liver 25-OHase expression ($p = 0.95$; **Figure 4B**). Also, no differences were seen with

kidney 1α-OHase expression ($p = 0.81$; **Figure 4C**). Our findings highlighted that liver

25-OHase and the kidney 1α-OHase expression were not affected by dietary super-

doses of vitamin D_3; however, the super-dosages led to downregulation in kidney 24-

OHase expression.

Discussion

Our results suggest that dietary super-doses of D_3 greatly increased plasma and

egg yolk D_3 levels. Increased plasma vitamin D_3 indicates these birds absorbed the

vitamin D_3 from their diets. However, the inactive vitamin D_3 metabolite, $24,25\text{-}(OH)_2\text{-}D_3$

increased at a much lower rate than D_3 and $25\text{-}OH\text{-}D_3$. Increasing plasma $24,25\text{-}(OH)_2\text{-}$

D_3 concentrations highlights that these hens were likely trying to reduce their circulating

vitamin D_3 levels. In addition, egg yolk D_3 drastically increased while yolk $25\text{-}OH\text{-}D_3$ had

a smaller rate of increase. Therefore, we speculate that maximal 24-hydroxylation

activity was achieved and these hens deposited excess D_3 in egg yolks to avoid vitamin

D toxicity.

Vitamin D, regardless of its biogenic or dietary origins, is transported and

circulated in blood which is why vitamin D is also considered a hormone (**Figure 5)**

(16). Circulating vitamin D_3 is transported to the liver to have a hydroxyl group added to

C-25 by 25-OHase which results in 25-OH-D_3. 25-OH-D_3 moves to kidney to have

another hydroxyl group added at C-1 by 1α-OHase to become 1,25-

dihydroxycholecalciferol [1,25-$(OH)_2$-D_3], the most active form of vitamin D_3 that exerts

vitamin D's biological effects via binding to VDR. However, there are regulatory

pathways for 25-OH-D_3 that can occur in the kidney with 24-OHase adding a hydroxyl

group to C-24 to result in 24,25-$(OH)_2$-D_3 (16). 24,25-$(OH)_2$-D_3 is considered an inactive

form of vitamin D that is excreted as a means for the body to remove excess vitamin D

from circulation (21).

Vitamin D hydroxylases are part of a cytochrome P450 family that regulate

vitamin D metabolism (21). The liver is a major site of 25-hydroxylation via 25-OHase in

liver mitochondria and microsomes (22). There are extrahepatic sources of 25-OHase

like kidney (23) and intestine (24); however, it is not clear how extrahepatic sources

contribute physiologically to vitamin D metabolism (22). To our knowledge, there is only

one paper that described the avian cytochrome P450 family in chicken liver (25). Aged

laying hens fed diets containing super-doses of D_3 did not result in any differential liver

25-OHase expression. This could signify liver 25-OHase may not be under metabolic

control in avian species. Studies with rats and humans showed that increased plasma

25-OH-D_3 concentration is related to dietary vitamin D_3 intake (24, 26). Half-life of

plasma 25-OH-D_3 is at least 18 d (27). Therefore, it is possible that our observations of

increased 25-OH-D_3 concentration may not be necessarily caused by increased 25-

OHase expression.

VDR is important for exerting biological effects of vitamin D with increasing

intestinal Ca absorption being a prominent effect (28). Knockout rats that could not

express VDR had decreased intestinal Ca absorption and almost no Ca-related

transport protein expression (28, 29). In our study, hens fed diets with higher super-

dose levels of D_3 had upregulated VDR expression in duodenum (**Figure 3A**). This is

further supported by an increase in responsiveness of VDR to D_3 earlier in the intestine

for increasing Ca absorptive capacity (30, 31). Super-dose levels of vitamin D_3 intake

did not affect liver VDR expression of the hens in this study likely because VDR has

little to no expression in the liver (30, 32, 33). However, the distal part of intestine and

colon were shown to have increased VDR expression in rats fed diets with $1,25\text{-(OH)}_2$-

D_3 (34).

Our hens were fed the experimental diets for eleven weeks and no discernable

changes in egg production or body weight were observed (**Table S1**). Although egg

quality was shown to not be affected by dietary super-doses of D_3 in our study, a laying

hen's physiological status affects egg quality (35). Another study reported how laying

hens fed extremely high levels of D_3 also had no changes in egg production, except egg

qualities like eggshell quality were reduced (36). Our findings agree with Mattila et al.

(14) and a recent study by our colleagues (19) that laying hens fed diets with greater

levels of dietary D_3 throughout their production cycle increases yolk D_3 content. An

interesting finding in our study was that 25-OH-D_3 in yolk increased with increased

dietary D_3 even though it was much lower than D_3. Mattila et al. (18) also observed the

same phenomenon with D_3 and $25\text{-OH-}D_3$ in egg yolk being affected by dietary D_3 fed to

hens. Although $25\text{-OH-}D_3$ is a more potent source for vitamin D activity, more research

is needed to explore the transfer rate of $25\text{-OH-}D_3$ from hens fed crystalline $25\text{-OH-}D_3$ in

comparison to hens fed D_3. Our findings indicated that D_3 is deposited more readily into

yolk versus $25\text{-OH-}D_3$. One assumption is that hens fed $25\text{-OH-}D_3$ have regulatory

feedback mechanisms acting to reduce plasma $25\text{-OH-}D_3$ levels. Therefore, it is

possible that excess $25\text{-OH-}D_3$ was transferred to the egg yolk instead of D_3 as

observed with hens from our study.

Plasma $25\text{-OH-}D_3$ concentration is important as a vitamin D status indicator (37,

38). Although plasma $25\text{-OH-}D_3$ concentration ranges have not been fully characterized

in laying hens, vitamin D status can still be ascertained by observing $25\text{-OH-}D_3$

concentration relative to D_3 and $24,25\text{-(OH)}_2\text{-}D_3$. In our study, the 400 and 800 IU D_3

treatments were formulated to meet NRC requirements for laying hens. Whereas the

14,000, 20,000, and 36,000 IU D_3 treatments were dietary super-dose treatments for D_3

intake. Higher dietary vitamin D_3 levels led to increasing plasma concentrations of

vitamin D_3 and $25\text{-OH-}D_3$ (**Figures 1A and 1B**). The fascinating element of this data

was the large range for plasma $25\text{-OH-}D_3$. The range was 7.15 ng/mL $25\text{-OH-}D_3$ for 400

IU hens to 85.2 ng/mL $25\text{-OH-}D_3$ for 36,000 IU hens. Yolk $25\text{-OH-}D_3$ levels were similar

to plasma with the concentration range, 3.48 ng/g $25\text{-OH-}D_3$ for 400 IU hens to 51.6

ng/g $25\text{-OH-}D_3$ for 36,000 IU hens. Supplementing laying hen diets with $25\text{-OH-}D_3$

increased egg yolk $25\text{-OH-}D_3$, although it resulted in lower egg yolk vitamin D_3 (39).

Taking into consideration with our findings and what Duffy et al. (39) reported, the

vitamin D metabolite composition in egg yolk is influenced by whatever dietary vitamin

D_3 isoform is fed to the laying hens.

Prior research characterized 24,25-$(OH)_2$-D_3's function as an inactive vitamin D

metabolite that is excreted from an animal's body (40, 41). The biological significance of

24,25-$(OH)_2$-D_3 is to reduce 25-OH-D_3's plasma concentration (42). Hydroxylation of C-

24 is significant because it can also occur with 1,25-$(OH)_2$-D_3 to lead to 1,24,25-

trihydroxylcholecalciferol which is also excreted (41). To our current knowledge, there is

no research on 24,25-$(OH)_2$-D_3 plasma concentration in laying hens. Comparable to

plasma 25-OH-D_3, the hens' plasma 24,25-$(OH)_2$-D_3 increased as their dietary D_3 intake

increased. However, unlike plasma D_3 and 25-OH-D_3, there was a large difference in

the rate of increase with 24,25-$(OH)_2$-D_3 having a tiny increase overall. Plasma 24,25-

$(OH)_2$-D_3 in hens fed 14,000, 20,000, and 36,000 IU D_3 /kg of feed was 6.8, 11.1, and

17.4 ng/mL, respectively. The rate of increase for 24,25-$(OH)_2$-D_3 relative to D_3 and 25-

OH-D_3 at super-dosage levels off at 20%, highlighting a possible asymptotic

relationship. The asymptote began at the 14,000 IU D_3 level and can be potentially

explained by 24-hydroxylation activity achieving its maximal limit. The combination of

the lack of an exponential increase for plasma 24,25-$(OH)_2$-D_3, unlike D_3 and 25-OH-D_3,

and the aforementioned asymptote is a monumental finding. It should be noted that

there is a paradox with how hens fed super-dose levels of dietary vitamin D_3 in this

study had decreased kidney expression of 24-OHase. A study with rats reported how

kidney 24-OHase expression was reduced by 1,25-$(OH)_2$-D_3 administration (43). The

decreased 24-OHase expression in hens fed super-dose levels of D_3 is possibly linked

to increased levels of 1,25-$(OH)_2$-D_3. Another possible explanation is the long half-life of

plasma 24,25-$(OH)_2$-D_3 (over 15 d (44)) caused 24,25-$(OH)_2$-D_3 to build up

concentration in super-dose fed hens.

Our study has a number of strengths that highlight its impact with advancing

nutritional knowledge. A significant strength of our study is the distinct separation of

plasma and egg yolk vitamin D metabolite data between treatment groups. This

emphasized the experimental design captured a broad range of dietary vitamin D_3

supplementation level effects on plasma vitamin D metabolites that future research can

focus on a specific range to build off our findings. Another strength of our study was

the plasma 25-OH-D_3 measurements to help assess the vitamin D status of the hens.

Our study provides novel observations of laying hen plasma 25-OH-D_3 levels relative to

dietary vitamin D_3 supplementation that can be valuable for the poultry industry to

consider with vitamin D status. The plasma 24,25-$(OH)_2$-D_3 data is the most exciting

novel finding of our study. Further understanding of the asymptotic relationship of the

super-dose levels with plasma 24,25-$(OH)_2$-D_3 concentrations can open up new

knowledge about 24,25-$(OH)_2$-D_3's value as a biomarker for vitamin D metabolism.

There were a few limitations with this study that were realized when data was

collected. We should have investigated kidney histopathology of these birds because

soft-tissue calcification or renal kidney failure could result from the hens reaching

vitamin D toxicity (45, 46). However, Mattila et al. (14) examined histological samples of

younger hens fed 15,000 IU D_3/kg of feed and they did not observe any pathological

issues with their hens' kidneys, leading them to conclude that vitamin D toxicity had not

occurred. Wen et al. (19) noted how 17-wk old pullets fed diets containing 68,348 IU D_3

had lower body weight and suggested that the pullets were likely affected by Ca toxicity.

No weight loss was observed with the hens in our study and there was no interactive

effect with ionized calcium between dietary vitamin D_3 level overtime. So, it is plausible

the hens may not have reached vitamin D toxicity. The smaller sample sizes and

missing treatment groups from the qPCR results were because some tissue samples

would not yield RNA for cDNA synthesis, even after multiple extraction attempts. All the

tissue samples, except for plasma, had to be temporarily stored in a 7°C cold room until

freezer space was made available which could have caused a reduction in RNA quality.

Although statistical power may be lacking as a result of reduced sample size, the

biological effect with 24-OHase looks promising and is still important for future research

for vitamin D metabolism. We also stored the egg yolk in a refrigerator for about a year

before the yolk was freeze dried. When compared to Wen et al. (19) results which used

hens close to the age of the hens used in our study, our findings hinted towards minimal

D_3 degradation (**Table 4**). The amount of egg yolk vitamin D_3 for similar dietary

treatments between the two studies were comparable. This could indicate how stable D_3

is when it is stored in cold, dark conditions. If this study was repeated, then using aged

roosters, in addition to aged hens, would provide a perspective of what dietary vitamin

D_3 super-doses could result in vitamin D metabolism because the roosters cannot lay

eggs and remove excess vitamin D through egg yolk deposition like hens. If roosters

had soft-tissue calcification or kidney failure, then that would illustrate vitamin D super-

doses led to toxicity and the hens avoided the toxicity by laying eggs containing the

excessive vitamin D.

Accounting for excessive yolk D_3 levels, we speculate that hens fed super-doses

of D_3 converted their 25-OH-D_3 to 24,25-$(OH)_2$-D_3 to avoid vitamin D toxicity, but 24-

hydroxylation reached its maximal activity rate. Hens would shunt excess D_3 and 25-

OH-D_3 to egg yolk as a consequence of increasing plasma vitamin D concentration.

This rationale is further validated because these hens can only transfer so much D_3 to

yolk. This suggests why super-dose fed hens had higher levels of plasma D_3 even

though they transferred a lot of D_3 to egg yolk. Our novel finding with plasma 24-OH-D_3

concentration has important implications for geriatric humans who take supplements.

Except for lactating women who can transfer excess vitamin D to milk, there is a

possibility that older people taking high levels of vitamin D supplementation than

intended could be at risk of vitamin D toxicity. To our knowledge, there are no studies

that explored 24,25-$(OH)_2$-D mechanisms when vitamin D is administered in large

doses. Future research should elucidate vitamin D mechanisms with great levels of

intake and how 24-hydroxylation is influenced.

Our study exhibits that feeding super-doses of dietary D_3 to aged laying hens

increases their plasma and egg yolk D_3 metabolite concentrations (**Figure 6**).

Importantly, there is a possible metabolic limit of 24-hydroxylation to remove excess

circulating D_3. Investigating 24-hydroxylation mechanisms will be important to

understanding vitamin D supplementation impacts in geriatric animals for improving

bone health and vitamin D metabolism in older humans.

Acknowledgments

We thank Gavin Conant, Meghan Connelly, Peter Ferket, Laura Hernandez, Matt Koci,

Shannon Madden, Celeste Sheftel, and Teresa Walker for their comments and edits to

the initial manuscript; Taylor Jones and Nick Thompson for their help with the RNA

extraction, cDNA synthesis, and qPCR experiments; David Dickey for statistical

assistance; Jeff Hall and Zach Spivey for their help with taking care of the hens and

collecting samples; and Liza Lentz of the Environmental and Agricultural Testing

Service laboratory, Department of Crop and Soil Sciences, at North Carolina State

University who performed the mineral composition experiments and analyses. The

authors' responsibilities were as follows-MFW and KAL: designed research; MFW,

DDH, and KAL: conducted research; MFW: analyzed data; MFW and KAL: wrote

manuscript; MFW and KAL: prepared experimental diets for study; MFW and PMP:

prepared samples for ashing; MFW and DDH: prepared and shipped plasma and egg

yolk samples to Heartland Assays; MFW and KAL: had primary responsibility for final

content; and all authors have read and approved the final manuscript.

References

1.   Dawson-Hughes B, Harris SS, Krall EA, Dallal GE. Effect of calcium and vitamin D supplementation on bone density in men and women 65 years of age or older. New England Journal of Medicine 1997;337(10):670-6.

2.   Ooms ME, Roos JC, Bezemer PD, Van Der Vijgh W, Bouter LM, Lips P. Prevention of bone loss by vitamin D supplementation in elderly women: a randomized double-blind trial. The Journal of Clinical Endocrinology & Metabolism 1995;80(4):1052-8.

3.   Tang BM, Eslick GD, Nowson C, Smith C, Bensoussan A. Use of calcium or calcium in combination with vitamin D supplementation to prevent fractures and bone loss in people aged 50 years and older: a meta-analysis. The Lancet 2007;370(9588):657-66.

4.   Bischoff-Ferrari HA, Willett WC, Wong JB, Stuck AE, Staehelin HB, Orav EJ, Thoma A, Kiel DP, Henschkowski J. Prevention of nonvertebral fractures with oral vitamin D and dose dependency: a meta-analysis of randomized controlled trials. Archives of internal medicine 2009;169(6):551-61.

5.   Bischoff-Ferrari HA, Willett WC, Orav EJ, Lips P, Meunier PJ, Lyons RA, Flicker L, Wark J, Jackson RD, Cauley JA. A pooled analysis of vitamin D dose requirements for fracture prevention. New England Journal of Medicine 2012;367(1):40-9.

6.   Holick MF. Vitamin D status: measurement, interpretation, and clinical application. Annals of epidemiology 2009;19(2):73-8.

7.   Zhen D, Liu L, Guan C, Zhao N, Tang X. High prevalence of vitamin D deficiency among middle-aged and elderly individuals in northwestern China: its relationship to osteoporosis and lifestyle factors. Bone 2015;71:1-6.

8.   Heaney RP. Nutrition and risk for osteoporosis. Edtion ed. Osteoporosis: Elsevier, 2001:669-700.

9.   Holick MF. Vitamin D deficiency. New England Journal of Medicine 2007;357(3):266-81.

10.   Klontz KC, Acheson DW. Dietary supplement–induced vitamin D intoxication. New England Journal of Medicine 2007;357(3):308-9.

11.   Jones G. Pharmacokinetics of vitamin D toxicity. The American Journal of Clinical Nutrition 2008;88(2):582S-6S. doi: 10.1093/ajcn/88.2.582S.

12.   Heaney RP, Davies KM, Chen TC, Holick MF, Barger-Lux MJ. Human serum 25-hydroxycholecalciferol response to extended oral dosing with cholecalciferol. The American journal of clinical nutrition 2003;77(1):204-10.

13.   Haddad JG, Jr., Rojanasathit S. Acute Administration of 25-Hydroxycholecalciferol in Man. The Journal of Clinical Endocrinology & Metabolism 1976;42(2):284-90. doi: 10.1210/jcem-42-2-284.

14.   Mattila P, Valaja J, Rossow L, Venäläinen E, Tupasela T. Effect of vitamin D2- and D3-enriched diets on egg vitamin D content, production, and bird condition during an entire production period. Poultry Sci 2004;83(3):433-40.

15.   Whitehead CC. Overview of bone biology in the egg-laying hen. Poultry Sci 2004;83(2):193-9. doi: 10.1093/ps/83.2.193.

16.   DeLuca HF. Overview of general physiologic features and functions of vitamin D. The American Journal of Clinical Nutrition 2004;80(6):1689S-96S. doi: 10.1093/ajcn/80.6.1689S.

17.   Whitehead C, Fleming R. Osteoporosis in cage layers. Poultry Sci 2000;79(7):1033-41.

18.   Mattila P, Lehikoinen K, Kiiskinen T, Piironen V. Cholecalciferol and 25-hydroxycholecalciferol content of chicken egg yolk as affected by the cholecalciferol content of feed. Journal of Agricultural Food chemistry 1999;47(10):4089-92.

19.   Wen J, Livingston K, Persia M. Effect of high concentrations of dietary vitamin D3 on pullet and laying hen performance, skeleton health, eggshell quality, and yolk vitamin D3 content when fed to W36 laying hens from day of hatch until 68 wk of age. Poultry Sci 2019;98(12):6713-20.

20.   Haussler MR, Haussler CA, Bartik L, Whitfield GK, Hsieh J-C, Slater S, Jurutka PW. Vitamin D receptor: molecular signaling and actions of nutritional ligands in disease prevention. Nutrition reviews 2008;66(suppl_2):S98-S112.

21.   Horst R, Reinhardt T, Reddy GS. Vitamin D metabolism. Vitamin D 2005;1:15-36.

22.   Zhu J, DeLuca HF. Vitamin D 25-hydroxylase–Four decades of searching, are we there yet? Archives of biochemistry and biophysics 2012;523(1):30-6.

23.   Gascon-Barré M, Demers C, Ghrab O, Theodoropoulos C, Lapointe R, Jones G, Valiquette L, Ménard D. Expression of CYP27A, a gene encoding a vitamin D-25 hydroxylase in human liver and kidney. Clinical endocrinology 2001;54(1):107-15.

24.   Theodoropoulos C, Demers C, Mirshahi A, Gascon-Barré M. 1, 25-Dihydroxyvitamin D3 downregulates the rat intestinal vitamin D3-25-hydroxylase CYP27A. American Journal of Physiology-Endocrinology and Metabolism 2001;281(2):E315-E25.

25. Watanabe KP, Kawai YK, Ikenaka Y, Kawata M, Ikushiro S-I, Sakaki T, Ishizuka M. Avian cytochrome P450 (CYP) 1-3 family genes: isoforms, evolutionary relationships, and mRNA expression in chicken liver. PLoS One 2013;8(9).

26. Thomas MK, Lloyd-Jones DM, Thadhani RI, Shaw AC, Deraska DJ, Kitch BT, Vamvakas EC, Dick IM, Prince RL, Finkelstein JS. Hypovitaminosis D in medical inpatients. New England Journal of Medicine 1998;338(12):777-83.

27. Jones K, Assar S, Harnpanich D, Bouillon R, Lambrechts D, Prentice A, Schoenmakers I. 25 (OH) D2 half-life is shorter than 25 (OH) D3 half-life and is influenced by DBP concentration and genotype. The Journal of Clinical Endocrinology & Metabolism 2014;99(9):3373-81.

28. Van Cromphaut SJ, Dewerchin M, Hoenderop JG, Stockmans I, Van Herck E, Kato S, Bindels RJ, Collen D, Carmeliet P, Bouillon R. Duodenal calcium absorption in vitamin D receptor–knockout mice: functional and molecular aspects. Proceedings of the National Academy of Sciences 2001;98(23):13324-9.

29. Song Y, Kato S, Fleet JC. Vitamin D receptor (VDR) knockout mice reveal VDR-independent regulation of intestinal calcium absorption and ECaC2 and calbindin D9k mRNA. The Journal of nutrition 2003;133(2):374-80.

30. Wang Y, Zhu J, DeLuca HF. Where is the vitamin D receptor? Archives of biochemistry and biophysics 2012;523(1):123-33.

31. Boos A, Riner K, Hässig M, Liesegang A. Immunohistochemical demonstration of vitamin D receptor distribution in goat intestines. Cells Tissues Organs 2007;186(2):121-8.

32. Sandgren ME, Brönnegärd M, DeLuca HF. Tissue distribution of the 1, 25-dihydroxyvitamin D3 receptor in the male rat. Biochemical and biophysical research communications 1991;181(2):611-6.

33. Pike JW, Goozé LL, Haussler MR. Biochemical evidence for 1, 25-dihydroxyvitamin D receptor macromolecules in parathyroid, pancreatic, pituitary, and placental tissues. Life sciences 1980;26(5):407-14.

34. Chow EC, Maeng HJ, Liu S, Khan AA, Groothuis GM, Pang KS. 1α, 25-Dihydroxyvitamin D3 triggered vitamin D receptor and farnesoid X receptor-like effects in rat intestine and liver in vivo. Biopharmaceutics & drug disposition 2009;30(8):457-75.

35. Travel A, Nys Y, Bain M. Effect of hen age, moult, laying environment and egg storage on egg quality. Edtion ed. Improving the safety and quality of eggs and egg products: Elsevier, 2011:300-29.

36. Ameenuddin S, Sunde M, DeLuca H, Cook M. Excessive cholecalciferol in a layers diet: decline in some aspects of reproductive performance and increased bone mineralisation of progeny. Brit Poultry Sci 1986;27(4):671-7.

37. Dawson-Hughes B, Heaney RP, Holick MF, Lips P, Meunier PJ, Vieth R. Estimates of optimal vitamin D status. Springer, 2005.

38. Intakes IoMSCotSEoDR. Dietary reference intakes. Edtion ed. Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride: National Academies Press (US), 1997.

39. Duffy SK, Rajauria G, Clarke LC, Kelly AK, Cashman KD, O'Doherty JV. The potential of cholecalciferol and 25-hydroxyvitamin D3 enriched diets in laying hens, to improve egg vitamin D content and antioxidant availability. Innovative Food Science & Emerging Technologies 2017;44:109-16.

40. Holick M, Schnoes H, DeLuca H, Gray R, Boyle I, Suda T. Isolation and identification of 24, 25-dihydroxycholecalciferol, a metabolite of vitamin D3 made in the kidney. Biochemistry 1972;11(23):4251-5.

41. Tanaka Y, Castillo L, DeLuca H. The 24-hydroxylation of 1, 25-dihydroxyvitamin D3. Journal of Biological Chemistry 1977;252(4):1421-4.

42. Bikle D. Vitamin D: Production, metabolism, and mechanisms of action. Edtion ed. Endotext [Internet]: MDText. com, Inc., 2017.

43. Shinki T, Jin CH, Nishimura A, Nagai Y, Ohyama Y, Noshiro M, Okuda K, Suda T. Parathyroid hormone inhibits 25-hydroxyvitamin D3-24-hydroxylase mRNA expression stimulated by 1 alpha, 25-dihydroxyvitamin D3 in rat kidney but not in intestine. Journal of Biological Chemistry 1992;267(19):13757-62.

44. Russell R, Kanis J, Smith R, Adams N, Bartlett M, Cundy T, Cochran M, Heynen G, Warner G. Physiological and pharmacological aspects of 24, 25-dihydroxycholecalciferol in man. Edtion ed. Homeostasis of Phosphate and Other Minerals: Springer, 1978:487-503.

45. Vieth R. Vitamin D toxicity, policy, and science. Journal of Bone and Mineral Research 2007;22(S2):V64-V8.

46. Morrissey RL, Cohn RM, Empson Jr R, Greene HL, Taunton OD, Ziporin ZZ. Relative Toxicity and Metabolic Effects of Cholecalciferol and 25-Hydroxycholecalciferol in Chicks. The Journal of nutrition 1977;107(6):1027-34.

Figure Legends

Figure 1. Vitamin D₃ metabolite plasma concentrations of 78 wk Hy-Line Brown laying hens fed different levels of dietary vitamin D₃. A) Cholecalciferol (Vitamin D₃; 400 and 800 IU treatment concentrations were not determined) **B)** 25-hydroxycholecalciferol (25-OH-D₃) **C)** 24,25-dihydroxycholecalciferol (24,25-(OH)₂-D₃) **D)** Ratio of 24,25-(OH)₂-D₃/25-OH-D₃ presented as a percentage. White squares denote standard NRC range D₃ levels (400 and 800 IU D₃/kg) in diet (n = 4) and gray diamonds denote super-dose levels of D₃ (14,000, 20,000, and 36,000 IU D₃/kg) in diet (n = 4). Samples were reported as means ± SEM. Samples with common letters were not statistically different from each other (General linear models, $p < 0.0001$).

Figure 2. Egg yolk vitamin D₃ metabolite concentrations from 78 wk Hy-Line Brown laying hens fed different levels of dietary vitamin D₃. A) Cholecalciferol (Vitamin D₃) **B)** 25-hydroxycholecalciferol (25-OH-D₃). White squares denote standard NRC range D₃ levels (400 and 800 IU D₃/kg) in diet (n = 4) and gray diamonds denote super-dose levels of D₃ (14,000, 20,000, and 36,000 IU D₃/kg) in diet (n = 4). Samples were reported as means ± SEM; however, error bars values were narrow and overlapped by the marker for each sample. Samples with common letters were not statistically different from each other (General linear models, $p < 0.0001$).

Figure 3. Relative gene expression of vitamin D receptor (VDR) in duodenum, ileum, liver, and kidney of 78 wk Hy-Line Brown laying hens fed different levels of dietary vitamin D₃. A) Duodenal VDR (n = 2-6) **B)** Ileal VDR (n = 2-5) **C)** Liver VDR (n = 2-4) **D)** Kidney VDR (n = 2-4). Tissues were analyzed using qPCR normalized against glyceraldehyde phosphate dehydrogenase (GAPDH; housekeeping gene) expression. White bars with dots denote standard NRC range D₃ levels in diet and gray bars with stripes denote super-dose levels of D₃ in diet. All samples ran in triplicates and reported as means ± SEM. Bars with common letters were not statistically different from each other (General linear models, $p < 0.05$).

Figure 4. Relative gene expression of vitamin D hydroxylases in liver and kidney of 78 wk Hy-Line Brown laying hens fed different levels of dietary vitamin D₃. A) Kidney 24-hydroxylase (24-OHase; n = 2-4) **B)** Liver 25-hydroxylase (25-OHase; n = 4-7) **C)** Kidney 1α-hydroxylase (1α-OHase; n = 2-5). Tissues were analyzed using qPCR normalized against glyceraldehyde phosphate dehydrogenase (GAPDH; housekeeping gene) expression White bars with dots denote standard NRC range D₃ levels in diet and gray bars with stripes denote super-dose levels of D₃ in diet. All samples ran in triplicates and reported as means ± SEM. Bars with common letters were not statistically different from each other (General linear models, $p < 0.05$).

Figure 5. Vitamin D_3 metabolism, enzymes, and two possible endpoints of vitamin D metabolism. Vitamin D_3 is converted to 25-hydroxycholecalciferol (25-OH-D_3) in the liver by 25-hydroxylase (25-OHase). As a vitamin D status biomarker, 25-OH-D_3 can be further hydroxylated at the C-1 or C-24 positions to become 1,25-dihydroxycholecalciferol [1,25-$(OH)_2$-D_3] or 24,25-dihydroxycholecalciferol [24,25-$(OH)_2$-D_3], respectively by 1α-hydroxylase (1α-OHase) or 24-hydroxylase (24-OHase) which are expressed in the kidney. 1,25-$(OH)_2$-D_3 is the biologically active form of D_3 that binds to vitamin D receptor to exert biological effects and 24,25-$(OH)_2$-D_3 is an inactive vitamin D_3 metabolite that is excreted from the body.

Figure 6. Dietary super-doses of vitamin D_3 fed to aged laying hens causes drastic increase in plasma and egg yolk vitamin D metabolites. Egg yolk vitamin D_3 levels were strongly correlated to the dietary levels of vitamin D_3 fed to the hens. Egg yolk 25-hydroxycholecalciferol (25-OH-D_3) levels were also dependent on dietary vitamin D_3; however, 25-OH-D_3 increased at a lower rate. For plasma vitamin D metabolites, vitamin D_3 and 25-OH-D_3 concentrations increased relative to dietary vitamin D_3 levels fed to the hens. Plasma 24,25-dihydroxycholecalciferol [24,25-$(OH)_2$] levels were also dependent on dietary vitamin D_3, but rate of increase was low. Dietary super-doses of vitamin D_3 caused downregulation of kidney 24-hydroxylase (24-OHase) expression.

Table 1. Ingredient composition of the experimental basal diet

Ingredient Name	%
Corn	63.40
Soybean meal, 46% CP	20.60
Calcium carbonate	9.20
Poultry fat	2.33
Dicalcium phosphate[1]	1.95
L-Lysine-HCl, 78.8%	1.01
Sodium bicarbonate	0.57
Vitamin premix[2]	0.25
Mineral premix[3]	0.20
Choline chloride, 60% choline	0.20
DL-Methionine, 99%	0.14
Salt	0.10
Selenium premix	0.05

[1]Dicalcium phosphate contains 19.79% calcium, 17.91% phosphorus, and 17.73% available phosphorus.

[2]Provided as milligrams per kilogram of diet: Thiamin HCL, 1.8; Riboflavin, 3.6; Ca

pantothenate, 10; Niacin, 25; Pyridoxine HCL, 3; Folacin, 0.55; Biotin, 0.15; B12, 0.01; Vitamin A 1500 IU/g, 6; Vitamin D3 400 IU/g, 1; Vitamin E 10 IU/g, 40; Vitamin K, 0.55; Ethoxyquin, 125.

[3]Trace minerals provided per kg of premix: 60 g manganese (Mn SO4); 60 g zinc (ZnSO4); 40 g iron (FeSO4); 5 g copper (CuSO4); 1.25 g iodine [Ca(IO3)2].

Table 2. Nutrient content of the experimental diet

Nutrient	%
Dry matter	90.46
Moisture	9.54
Crude protein	16.00
Fat	4.98
Fiber	1.81
Calcium	4.00
Total phosphorus	0.67
Ash	12.44
Available phosphorus	0.46
Methionine+Cysteine	0.64
Lysine	1.57
Methionine	0.40
Cysteine	0.26
Lysine	1.57
Tryptophan	0.19
Threonine	0.60
Isoleucine	0.72
Histidine	0.43
Valine	0.83
Leucine	1.43
Arginine	1.00
Phenylalanine	0.83
Metabolizable Energy, kcal/kg	2,900
Dietary electrolyte balance, mEq/100 g	202

Table 3. Primer sequences for quantitative real-time PCR (qPCR)

Gene	Orientation	Primer Sequence (5' – 3')	Size (bp)	Accession #
VDR	Forward	TGCCTCCAGTCTGGCATCTC	297	NM_205098.1
	Reverse	GGTGATTTTGCAGTCCCCGT		
1α-OHase	Forward	ATGATTGGCGTCCCCTTCAG	177	XM_422077.4
	Reverse	TCCACGCTTTCACTCACACA		
25-OHase	Forward	GCTGTCACTGGGATTCTTTGC	160	NM_001277354.1
	Reverse	CCAACCGAAAGGCACAAGTC		
24-OHase	Forward	AAACCCTGGAAAGCCTATCG	133	NM_204979.1
	Reverse	CCAGTTTCACCACCTCCTTG		
GAPDH	Forward	TGTTGTTGACCTGACCTGCC	291	NM_204305.1
	Reverse	CTGGCTCACTCCTTGGATGC		

Table 4. Comparison of egg yolk vitamin D_3 from eggs laid by hens fed different dietary levels of vitamin D_3 in this study compared to Wen et al. 2019 study (19). The closest experimental treatments by dietary vitamin D_3 levels were compared.

This study[1]	Wen et al. 2019 (19)[2]
IU/egg (Dietary treatment IU/kg)	
5.13 (400)	12.6 (1,681)
199.7 (14,000)	214.3 (18,348)
605.9 (36,000)	435.5 (35,014)

[1]Values were calculated by converting the egg yolk vitamin D_3 concentration (ng/g) to IU and multiplying by 15g (amount used for LC-MS/MS).
[2]These values were reported in Figure 1 for (19).

A

B

C

D

A

B

A

B

C

D

A

B

C