Angiomotin isoform 2 promotes binding of PALS1 to KIF13B at the base of primary cilia and

suppresses ciliary elongation

Running title: Ap80 promotes PALS1 binding to KIF13B

Stine Kjær Morthorst[1,5], Camilla Nielsen[1,5], Pietro Farinelli[1,5], Zeinab Anvarian[1], Christina Birgitte

R. Rasmussen[1], Andrea Serra-Marques[2,6], Ilya Grigoriev[2], Maarten Altelaar[3], Nicoline

Fürstenberg[1], Alexander Ludwig[3], Anna Akhmanova[2], Søren Tvorup Christensen[1] and Lotte Bang

Pedersen[1, *]

[1]Department of Biology, University of Copenhagen, Universitetsparken 13, DK-2100 Copenhagen

Ø, Denmark.

[2]Cell Biology, Department of Biology, Faculty of Science, Utrecht University, Utrecht,

Netherlands.

[3]Biomolecular Mass Spectrometry and Proteomics, Bijvoet Center for Biomolecular Research and

Utrecht Institute for Pharmaceutical Sciences, Utrecht University, Padualaan 8, Utrecht 3584 CH,

Netherlands.

[4]School of Biological Sciences and Institute of Structural Biology, Nanyang Technological

University, Singapore.

[5]Co-first authors.

[6]Current address: Immunology Discovery, Genentech, Inc., South San Francisco, CA, USA.

*Corresponding author

Lotte B. Pedersen
Department of Biology, University of Copenhagen
Universitetsparken 13
DK-2100 Copenhagen OE
Denmark
Tel: +45 3532 1690
E-mail: lbpedersen@bio.ku.dk

Abstract

The kinesin-3 motor KIF13B functions in endocytosis, vesicle transport, and regulation of ciliary length and signaling. Direct binding of the membrane-associated guanylate kinase (MAGUK) DLG1 to KIF13B's MAGUK-binding stalk (MBS) domain relieves motor autoinhibition and promotes microtubule plus end-directed cargo transport. Here we characterize Angiomotin isoform 2 (Ap80) as a novel KIF13B interactor that promotes binding of another MAGUK, the polarity protein and Crumbs complex component PALS1, to KIF13B. Live cell imaging analysis indicated that Ap80 is concentrated at the base of primary cilia and recruits PALS1 to this site, but is not itself a cargo of KIF13B. Consistent with a ciliary function for Ap80, its depletion led to elongated primary cilia while its overexpression caused ciliary shortening. Our results suggest that Ap80 may specifically activate KIF13B cargo binding at the base of primary cilia to regulate ciliary length.

Introduction

Kinesin motors mediate microtubule-based transport of diverse intracellular cargoes, including organelles and vesicles (Hirokawa and Tanaka, 2015). In humans 8 kinesin genes code for kinesin-3 family members, which are highly processive microtubule plus-end directed motors involved in vesicle transport and endocytosis (Siddiqui and Straube, 2017). Kinesin-3 member KIF13B (also known as GAKIN; guanylate kinase-associated kinesin) was initially identified as a direct binding partner of the scaffold protein DLG1 in lymphocytes (Hanada et al., 2000). Subsequent studies implicated KIF13B in multiple cellular and physiological processes e.g. regulation of neuronal polarity and axon formation (Horiguchi et al., 2006, Yoshimura et al., 2010); Schwann cell and oligodendrocyte myelination (Bolis et al., 2009, Noseda et al., 2016); cholesterol uptake and CAV1-dependent endocytosis of LRP1 in hepatocytes (Kanai et al., 2014, Mills et al., 2019); Golgi to plasma membrane trafficking of VEGFR2 in endothelial cells (Yamada et al., 2014) and of Rab6 exocytotic vesicles in HeLa cells (Serra-Marques et al., 2020); germ cell migration and Planar Cell Polarity signaling in *Xenopus laevis* (Tarbashevich et al., 2011, Ossipova et al., 2015); and regulation of ciliary CAV1 distribution and Sonic hedgehog (Shh) signaling in hTERT-immortalized retinal pigment epithelial (hereafter RPE1) cells (Schou et al., 2017).

KIF13B function requires tight temporal and spatial control of its motor activity, which is mediated by binding partners that recognize specific domains in KIF13B to regulate its conformation and dimerization (Soppina et al., 2014, Siddiqui and Straube, 2017, Morthorst et al., 2018, Ren et al., 2018). The best characterized KIF13B regulator and cargo is DLG1, a Scribble polarity complex component of the membrane-associated guanylate kinase (MAGUK) homologue family. DLG1 binds directly to the MAGUK-binding stalk (MBS) domain of KIF13B, which relieves motor autoinhibition and promotes KIF13B-mediated transport of DLG1 along microtubules (Hanada et al., 2000, Asaba et al., 2003, Yamada et al., 2007, Zhu et al., 2016). Similarly, other MAGUKs including DLG4 and PALS1, a core component of the Crumbs complex required for development and maintenance of epithelial cell polarity in diverse tissues, bind directly to the KIF13B MBS domain (Zhu et al., 2016). Furthermore, we identified an interaction between KIF13B and the ciliary transition zone (TZ) protein NPHP4 (Schou et al., 2017), which also associates with PALS1 and PATJ (Sang et al., 2011, Delous et al., 2009). However, it is unknown whether interaction between KIF13B and PALS1 is relevant in a ciliary context.

Primary cilia are microtubule-based sensory organelles present on the surface of many vertebrate cell types, where they coordinate e.g. Shh (Bangs and Anderson, 2017) and receptor

tyrosine kinase signaling (Christensen et al., 2017). The ciliary axoneme extends from the basal
body and is surrounded by a membrane, which is continuous with but compositionally distinct from
the plasma membrane. This compartmentalization is crucial for ciliary function, and is regulated by
the TZ at the ciliary base (Garcia-Gonzalo and Reiter, 2017), as well as by intraflagellar transport
(IFT) that via kinesin-2 and dynein 2 motors, respectively, moves trains of IFT particles with ciliary
cargoes up and down the cilium (Taschner and Lorentzen, 2016). Nematodes additionally express a
kinesin-3 motor, KLP-6, that regulates ciliary membrane composition and function specifically in
male sensory neurons by promoting release of extracellular vesicles from the ciliary tip (Akella and
Barr, 2021). Recently, we demonstrated that KIF13B displays bursts of bidirectional intraciliary
movement within primary cilia of RPE1 cells and is occasionally released from the ciliary tip (Juhl
et al., 2021). How KIF13B intraciliary movement is regulated and the identity of ciliary cargoes of
KIF13B are not known.

Here we characterize Angiomotin isoform 2 (Ap80) as a KIF13B interactor and show
that Ap80 promotes the binding of KIF13B to PALS1. We show that Ap80 is concentrated at and
recruits PALS1 to the ciliary base of RPE1 cells, and that its depletion and overexpression cause
ciliary elongation and shortening, respectively. Thus Ap80 may activate KIF13B cargo binding at
the ciliary base to regulate ciliary length.

Results and discussion

KIF13B interacts with the N-terminus of Ap80. To identify KIF13B interactors we performed
streptavidin pull-down assays with BirA- and GFP-tagged (BirAGFP) KIF13B constructs combined
with mass spectrometry (Table 1). We used the deletion mutants BirAGFP-KIF13B Tail 2 (residues
607-1826) and BirAGFP-KIF13B Tail 3 (residues 752-1826), which contain and lack the MBS
domain, respectively (Serra-Marques et al., 2020), reasoning that these may reveal MBS-specific
interactors. We identified several significant hits that had more peptides in the Tail 2 mutant sample
compared to Tail 3 (Table 1). These included DLG1, and its known interactors MPP7, LIN7C and
CASK (Bohl et al., 2007, Lee et al., 2002), and utrophin, a large cytoskeletal adaptor that binds to
KIF13B in a complex mediating endocytosis of LRP1 (Kanai et al., 2014). Known interactors of
utrophin, syntrophin (Kramarcy et al., 1994) and dystrobrevins (Peters et al., 1997), were also
present in the pull down. The most prominent potential novel binding partners of KIF13B, showing
significantly stronger association with Tail 2 compared to Tail 3, were kinase D-interacting
substrate of 220 kDa (KIDINS220), a conserved transmembrane molecule implicated in signaling

(Neubrand et al., 2012), and an adaptor protein belonging to the motin family, angiomotin (AMOT)

(Moleirinho et al., 2014) (Table 1).

We focused on AMOT due to its reported interaction with PALS1 (Wells et al., 2006,

Tan et al., 2020), which associates with KIF13B (Zhu et al., 2016) and the KIF13B interactor

NPHP4 (Delous et al., 2009, Schou et al., 2017). AMOT is found in two splice variants; angiomotin

isoform 1 (Ap130) contains a 409 amino acid N-terminal extension absent in isoform 2, Ap80

(Ernkvist et al., 2006). We previously reported that KIF13B binds Ap80 (Schou et al., 2017), but

the interaction was not characterized in detail. Here, we confirmed by FLAG immunoprecipitation

(IP) in HEK293T cells that FLAG-Ap80 interacts with HA-tagged full-length or motorless KIF13B

(KIF13B-Δmotor-HA; residues 393-1826), but FLAG-Ap130 does not (Figure 1A). Supportively,

FLAG-Ap80 co-IPed with GFP-KIF13B (Figure 1B). IP with different FLAG-Ap80 truncations

identified the first 34 residues in Ap80 as essential for KIF13B binding, since Ap80-NB (residues

1-245) comprising the N-terminus and BAR domain region, bound to KIF13B-Δmotor-HA, but

Ap80-B (residues 35-245) did not (Figure 1C, D). Ap80 residues 1-245 are also present in Ap130,

which failed to bind KIF13B (Figure 1A, C), suggesting that the KIF13B binding site in AMOT is

blocked by the extended N-terminus of Ap130.

Ap80 interacts with KIF13B stalk and tail region and promotes KIF13B-PALS1 association.

To map the region of KIF13B that binds Ap80-NB, we generated plasmids encoding GFP-KIF13B

truncations covering most of the C-terminal stalk and tail region (Figure 2A; (Schou et al., 2017)).

Co-expression with FLAG-Ap80-NB followed by FLAG IP identified residues 561-1500 (GFP-Tail

11) as the minimal Ap80-NB binding site in KIF13B (Figure 2B; Figure S1A, B); reciprocal GFP

IP confirmed this result (Figure S1C). Full length FLAG-Ap80 interacted with an even shorter

KIF13B fusion, GFP-Tail 8, which contains the MBS domain but lacks residues 1327-1500 present

in GFP-Tail 11 (Figure 2A,C; Figure S1D), suggesting that the MBS domain is necessary for

interaction with Ap80-NB and that residues 1327-1500 are also important for binding. However, the

latter region is dispensable for interaction with full length Ap80. The results are consistent with our

mass spectrometry data as the sequences of Tail 8 and Tail 11 are comprised within the BirAGFP-

KIF13B Tail 2 fusion that interacts with AMOT (Table 1; (Serra-Marques et al., 2020)).

Additionally, immunofluorescence microscopy (IFM) of RPE1 cells co-expressing FLAG-Ap80-

NB or FLAG-Ap80 with different GFP-KIF13B truncations showed that those binding to FLAG-

Ap80-NB (e.g. Tail 7 and 11) or FLAG-Ap80 (Tail 8) in IP (Fig. 2B, C), also co-localize with these

fusions, whereas non-binding truncations (e.g. Tail 9 and 12) do not (Fig. S1E, F). Co-expression of

FLAG-Ap80 with Ap80-binding GFP-KIF13B truncations increased the total cellular number of

FLAG-Ap80 positive vesicles (Fig. S1G), possibly due to their fission.

148   The KIF13B MBS domain binds directly to the guanylate kinase (GUK) domains of

DLG1/DLG4 and PALS1 (Yamada et al., 2007, Zhu et al., 2016), while residues 1327-1500 bind

MARK2 (Yoshimura et al., 2010). We therefore asked if Ap80 binds KIF13B indirectly via these

proteins. However, GFP-DLG1 and GFP-MARK2 did not co-IP with FLAG-Ap80, whereas HA-

PALS1 did (Figure 2D; Figure S2A). PALS1 binds to the Ap80 C-terminus, presumably via its

PDZ domain (Figure 2E; Figure S2A), while KIF13B binds to the Ap80 N-terminus (Ap80-NB;

Figure 1C, D). Thus, Ap80 interacts independently with both KIF13B and PALS1. Interestingly,

HA-PALS1 did not co-IP with motorless KIF13B (GFP-KIF13B-Δmotor) alone, but did upon

simultaneous co-expression with FLAG-Ap80 (Figure 2F). Similar results were obtained with full-

length GFP-KIF13B (Figure S2B). The inability of PALS1 to bind full-length or motorless KIF13B

could be due to intramolecular interaction in KIF13B blocking access to the MBS domain.

Supportively, IP using GFP- or HA-KIF13B truncations revealed that KIF13B residues 1289-1499

interact with the MBS domain (Figure S2C, D), consistent with previous work (Yamada et al.,

2007). Our results suggest that Ap80 promotes association of PALS1 with KIF13B by inducing a

conformational change in KIF13B that allows PALS1 to access the MBS region.

163   Our protein interaction data is summarized in Figure 2G. First, we suggest that the

interaction (a) between the NB region of Ap80 and MBS domain in KIF13B is necessary, but not

sufficient for these proteins to interact. Two additional supportive interactions can occur: one (b)

between the Ap80 NB region and KIF13B residues 1327 to 1500, which seem important for a

strong interaction between the proteins, and another (c) between the MBS domain and the C-

terminus of Ap80, likely mediated by PALS1. Binding of Ap80 to KIF13B presumably induces a

conformational change in KIF13B that permits binding of the MBS domain to PALS1.

Co-localisation of Ap80, PALS1 and KIF13B at the base of primary cilia. Next, we asked if

Ap80, KIF13B and PALS1 co-localize in cells. First, we examined the level of endogenous Ap80

and Ap130 in mouse fibroblasts (NIH 3T3 cells) and different mouse or human epithelial cell lines

(IMCD3, HEK293T, RPE1 cells), by immunoblotting with AMOT antibody. We chose RPE1 cells

for further analysis as they specifically expressed Ap80, but not Ap130. Further, in these cells Ap80

and KIF13B appeared to be upregulated by high cell confluency and serum deprivation (Figure

S3A), conditions that promote primary cilia formation (Pugacheva et al., 2007). Previously, we showed that GFP-KIF13B localizes to and moves within primary cilia of RPE1 cells (Schou et al., 2017, Juhl et al., 2021). We examined the subcellular localization of Ap80-GFP by live imaging in serum-deprived, ciliated RPE1 cells stably expressing the ciliary membrane marker SMO-tRFP (Lu et al., 2015). We found that GFP-tagged Ap80 is concentrated at the cilia base in most cells (89% ± 8%, n=67 cells) and exhibits little, if any, movement (Figure 3A, Figure S3G, Movie 1), even when co-expressed with mCherry-KIF13B that co-localized with Ap80-GFP at the centrosome (Movie 2, Movie 3). However, we cannot rule out that the GFP tag affects Ap80 motility. In fixed cells, FLAG-Ap80 and FLAG-Ap80-NB were also detected at the ciliary base (Figure S3B, E), but recruitment of Ap80 to this site was independent of KIF13B, as centrosomal levels of FLAG-Ap80 in $KIF13B^{-/-}$ cells was analogous to that of wild type (WT) cells (Figure S3C, D). GFP-tagged (Figure 3B) and endogenous PALS1 (Figure S3F) were mostly dispersed localized in RPE1 cells, but upon co-expression with FLAG-Ap80 (Figure 3C) or GFP-Ap80 (Figure S3G) PALS1 accumulated at the ciliary base and at cytoplasmic Ap80-positive puncta. Furthermore, in agreement with our biochemical data GFP-Ap80 and endogenous PALS1 both co-localized with KIF13B-Δmotor-HA in intracellular puncta (Figure S3H). Together, these results indicate that Ap80 localizes to the base of cilia and where it recruits PALS1, and Ap80 is likely not a cargo of KIF13B, but Ap80, KIF13B and PALS1 co-localize within cells. In MDCK-II cells expressing PALS1 fused to APEX2-EGFP (PALS1-A2E) (Tan et al., 2020), PALS1-A2E localized around the base (Figure S3I) and proximal region of the primary cilium (Figure S3J), suggesting that such localization is conserved in cell types other than RPE1.

Ap80 regulates ciliary length in RPE1 cells. Next, we set out to knockout the *AMOT* gene (coding for both Ap80 and Ap130) in RPE1 cells using CRISPR/Cas9 methodology (Ran et al., 2013). We generated a clonal $AMOT^{-/-}$ line containing a homozygous 1 bp deletion in the first exon resulting in a premature stop codon at position 184-186 (Figure S4A) and loss of detectable Ap80 protein (Figure 3D). Interestingly, IFM with antibodies for ciliary markers revealed that $AMOT^{-/-}$ RPE1 cells have significantly longer cilia than WT cells (Figure 3E, F). The $AMOT^{-/-}$ cells also seemed to have fewer cilia than controls, but this difference was statistically insignificant (Figure S4B). For technical reasons, rescue experiments in the $AMOT^{-/-}$ cells were unsuccessful, but partial depletion of Ap80 using endoribonuclease-prepared small interfering RNA (esiRNA) recapitulated the long cilia phenotype of $AMOT^{-/-}$ RPE1 cells (Figure S4C-E). Conversely, cells expressing GFP-Ap80

had significantly shorter cilia, but similar ciliation frequency, compared to GFP expressing cells (Figure 3G; Figure S4F). This was not simply due to protein over expression because the average relative expression level of GFP-Ap80 was about 5 times lower (18.2% ±2.0%, n=3) than that of GFP (100%, n=3), whereas average transfection efficiencies were 25.1% ±7.2% (n=3) for GFP-Ap80 and 58.8% ±6.0% (n=3) for GFP, respectively. Collectively, these results indicate that Ap80 negatively regulates ciliary length, but the underlying mechanism and functional consequences thereof remain to be clarified.

Conclusion

We characterized Ap80 as a KIF13B interactor that promotes binding of KIF13B to PALS1. Ap80 does not appear to be a KIF13B cargo, but is localized to the ciliary base where it recruits PALS1 and controls ciliary length. Ap80 was previously shown to bind RICH1, a GTPase-activating protein for the Rho GTPases Cdc42 and Rac1 (Richnau and Aspenstrom, 2001), and mediate its targeting to a protein complex at epithelial tight junctions containing PALS1, PATJ and PAR3 (Wells et al., 2006). This promoted re-localization of PALS1, PATJ and PAR3 from tight junctions to recycling endosomes, causing loss of tight junction integrity and induction of epithelial-mesenchymal transition (Wells et al., 2006, Heller et al., 2010). Similarly, in endothelial cells Ap80 promoted breakdown of cellular junctions to stimulate endothelial to mesenchymal transition (Bratt et al., 2005, Levchenko et al., 2004, Troyanovsky et al., 2001). Our work suggests that Ap80 may have an additional role in controlling the trafficking of PALS1 or even the entire Crumbs complex (CRB3, PALS1, PATJ, LIN7C) to the ciliary base. Supportively, components of the Crumbs complex have been detected in cilia, and CRB3 is required for ciliogenesis (Fan et al., 2004, Bazellieres et al., 2018).

Although more work is needed to determine how Ap80 regulates ciliary length, and the functional consequences thereof, the mechanisms could include local regulation of actin dynamics, e.g. through ciliary recruitment of RICH1 (Wells et al., 2006). The RICH1 effector Cdc42 localizes at the ciliary base and regulates ciliary length and signaling by modulating actin dynamics, and by recruiting PAR3/PAR6/aPKC (Drummond et al., 2018). Ap80 also associates physically with the PAR3/PAR6/aPKC complex (Wells et al., 2006), which regulates ciliogenesis (Sfakianos et al., 2007, Pruliere et al., 2011) and KIF13B activity (Yoshimura et al., 2010). In future studies it will be interesting to explore these interactions further from a ciliary perspective.

Materials and Methods

Antibodies. For immunoblotting, the following primary antibodies were used (catalog numbers and dilutions in parenthesis): rabbit anti-FLAG (Sigma, cat# F7425; 1:1,000), rabbit anti-GFP (Santa Cruz, cat# sc-8334; 1:1,000), rabbit anti-HA (SantaCruz, cat# sc-805; 1:500), mouse anti-DCTN1 (BD Biosciences, cat# 610474; 1:1,000), mouse anti-PALS1 (Santa Cruz, cat#sc-365411; 1:200), rabbit anti-GAPDH (Cell Signaling Technology, cat# 2118S; 1:1,000), rabbit anti-AMOT (kindly provided by Dr. Joseph Kissil, Scripps Research Institute, Jupiter, FL, USA; 1:1,000), rabbit anti-IFT88 (ProteinTech, cat# 13967-1-AP; 1:2,000). Secondary antibodies used for immunoblotting were Horseradish Peroxidase (HRP)-conjugated swine anti-rabbit (DAKO, cat# P0399; 1:4,000) or goat anti-mouse (DAKO, cat# P0447; 1:4,000).

For IFM analysis, the following primary antibodies were used: mouse anti-FLAG (Sigma cat# F1804; 1:1,000), mouse anti-PALS-1 (Santa Cruz, cat#sc-365411; 1:50), goat anti-MYC (Abcam, cat# ab9132; 1:500), chicken anti-GFP (Abcam, cat# ab13970; 1:2,000), rabbit anti-ARL13B (ProteinTech cat# 17711-1-AP; 1:500), rabbit anti-AMOT (kindly provided by Joseph Kissil, Scripps Research Institute, Jupiter, FL, USA; 1:200), mouse anti-DCTN1 (BD Biosciences, cat# 610473; 1:500), mouse anti-acetylated α-tubulin (Sigma, cat# T7451; 1:2,000), rabbit anti-CEP164 (Sigma, cat# HPA037606; 1:500). Secondary antibodies used for IFM were all from Invitrogen and diluted 1:600: Alexa Fluor 488-conjugated donkey anti-mouse (A-21202), donkey anti-goat (A-11055), donkey anti-rabbit (A-21206) and goat anti-chicken (A11039); Alexa Fluor 568-conjugated donkey anti-mouse (A-10037), and donkey anti-rabbit (A-10042); Alexa Fluor 647-conjugated donkey anti-mouse (A-31571), donkey anti-rabbit (A-31573).

PCR, cloning procedures and plasmids. All PCR and cloning was done following standard procedures; primer sequences are listed in Table S1 and plasmids used are listed in Table S2. Plasmids encoding truncated versions of GFP-KIF13B (Tail 7, 8, 9, 11 and 12) were generated by PCR with relevant primers using pEGFP-C1-KIF13B as template (Asaba et al., 2003), followed by cloning into pEGFP-C1 (Clontech). Plasmids encoding truncated KIF13B-HA fusions (Tail 1 and 16) were generated by PCR with relevant primers using pcDNA3-KIF13B-HA plasmid (Lamason et al., 2010) as template. Plasmids encoding truncated versions of FLAG-Ap80 (-NB, -B, -C and -BC) were generated by PCR with relevant primers using pCMV-Ap80 as template, followed by cloning into pFLAG-CMV2 (Sigma). Plasmid encoding GFP-Ap80 or Ap80-GFP was generated by PCR with relevant primers using pCMV-Ap80 as template, followed by cloning into pEGFP-C1 or

pEGFP-N1, respectively (Clontech). For CRISPR-Cas9 mediated knockout of AMOT, 25 nt oligos
targeting exon 1 in AMOT (5'- AATACCGTGGTCCCTCCACTTGG-3') were cloned into
pSpCas9(BB)-2A-Puro (PX459) using the procedure in (Ran et al., 2013). *Escherichia coli* DH10B
was used for transformation and plasmid amplification. Plasmid purification was performed using
NucleoBond Xtra Midi EF Kit from Macherey-Nagel. Inserts were sequenced at Eurofins MWG
Operon, Ebersberg, Germany.

Cell culture and transfections. Unless otherwise stated, cells were grown at 37°C, 5% CO_2 and
95% humidity in Dulbecco's modified Eagle's medium (DMEM, Gibco) with 10% heat inactivated
fetal bovine serum (FBS, Gibco) and 10 ml 1^{-1} penicillin-streptomycin (Gibco) and cell cultures
were passaged every 3-4 days. Cell lines used were: HEK293T cells (laboratory stock, originally
derived from American Type culture collection (ATCC), clone CRL-3216); hTERT-RPE1 cells
(laboratory stock; originally derived from the immortalized hTERT-RPE1 cell line, ATCC, clone
CRL-4000); hTERT-RPE1 cells lacking KIF13B (*KIF13B$^{-/-}$*; (Schou et al., 2017)); hTERT-RPE1
cells stably expressing SMO-tRFP (Lu et al., 2015); swiss NIH3T3 mouse fibroblasts (laboratory
stock, originally derived from ATCC clone CRL-1658); Mouse Inner Medullary Collecting Duct
(IMCD3) cells (laboratory stock, originally derived from ATCC, clone CRL-2123); and PALS1-
A2E MDCK-II cells (Tan et al., 2020). IMCD3 cells were grown in 45% DMEM and 45% F-12
(Ham; Sigma) with 10% FBS and 10 ml 1^{-1} penicillin-streptomycin whereas the PALS1-A2E
MDCK-II cells were grown and subjected to APEX2-mediated biotinylation as described
previously (Tan et al., 2020). Generation of *AMOT$^{-/-}$* RPE1 clone was done using the CRISPR-Cas9
method described in (Ran et al., 2013) with plasmid pSpCas9(BB)-2A-Puro (PX459) encoding
AMOT-specific guide RNA targeting the first exon of AMOT (5'-
AATACCGTGGTCCCTCCACTTGG -3'). Screening of 91 clones by immunoblot analysis led to
identification of a single clone with no detectable Ap80 expression. Sequencing of PCR-amplified
genomic DNA from this clone revealed a 1 bp deletion in exon 1 resulting in a premature stop
codon. All cell lines were routinely tested for contamination.
For plasmid transfections cells were grown to 40-50% confluency and transfections
were performed using FuGene6 (Promega, E2692) according to the manufacturer's protocol. For IP
experiments, HEK293T cells were grown in 10 cm diameter petri dishes and transfected with 4 µg
DNA. Six hours after transfection the medium was changed to fresh growth medium and the cells
were incubated for 16 hrs before harvest. For IFM experiments, RPE1 cells were grown on glass

coverslips in 20 mm diameter dishes and transfected with 1 µg DNA. Six hours after transfection

the medium was changed to either growth or serum-deprived medium for 16-24 hrs. For

immunoblot experiments, RPE1 cells were grown in 60 mm diameter dishes and transfected with 2

µg DNA. Six hours after transfection the medium was changed to either growth or serum-deprived

medium for 16-24 hrs.

310   All transfections with esiRNA (Sigma MISSION® esiRNA human AMOT cat#

EHU129771) and control siRNA (5'-UAAUGUAUUGGAAUGCAUA(dTdT)-3' from Eurofins

MWG Operon) were carried out as double-transfections using DharmaFECT Duo (Dharmacon).

RPE1 cells were grown to 80% confluency and transfected according to the manufacturer's manual.

Six hours after transfection the medium was changed to fresh growth medium. The following day

the cells were transfected again. 36 hrs later, the cells were split and an appropriate number of cells

were seeded for IFM or immunoblot analysis. Prior to fixation or cell lysis the cells were serum-

deprived for 16-24 hours.

Streptavidin pulldown assays and mass spectrometry. HEK293 cells were transfected with

BirAGFP-KIF13B Tail 2 or Tail 3 constructs (designated C2 and C3, respectively, in (Serra-

Marques et al., 2020)) using Polyethylenimine (PEI; Mw 2500; Polysciences) at a 3:1 PEI:DNA

ratio (w/w). Cells were harvested 24 hours after transfection, by scraping the cells in ice-cold PBS

and lysing cell pellets in the lysis buffer (20 mM Tris-HCl, pH 7.5, 100 mM NaCl, 1.0% Triton X-

100, and protease inhibitors; Roche). Supernatants and pellet fractions were separated by

centrifugation at maximum speed for 20 minutes. Supernatants were mixed with an equal amount of

Dyna M-280 Streptavidin beads (Life Technologies). Samples were incubated for 2 hours while

rotating at 4°C, collected with a magnet and pellets were washed 5-7 times with the wash buffer (20

mM Tris-HCl, pH 7.5, 100 mM NaCl, 0.1 % Triton X-100). Samples were eluted in SDS sample

buffer and 30 µl of each sample was run on a 12% Bis-Tris 1D SDS-PAGE gel (Biorad) for 1 cm

and stained with colloidal Coomassie dye G-250 (Gel Code Blue Stain Reagent, Thermo Scientific).

Each lane was cut into 1 band, which was treated with 6.5 mM dithiothreitol (DTT) for 1 hour at 60

°C for reduction and 54 mM iodoacetamide for 30 min for alkylation. The proteins were digested

overnight with trypsin (Promega) at 37°C. The peptides were extracted with acetonitrile (ACN) and

dried in a vacuum concentrator. The data were acquired using an Orbitrap Q Exactive mass

spectrometer. Peptides were first trapped (Dr Maisch Reprosil C18, 3 µm, 2 cm x 100 µm) before

being separated on an analytical column (Zorbax SB-C18, 1.8 µm, 40 cm x 50 µm), using a gradient

of 60 min at a column flow of 150 nl min⁻¹. Trapping was performed at 8 μl/min for 10 min in

solvent A (0.1 M acetic acid in water) and the gradient was as follows 7-30% solvent B (0.1 M

acetic acid in acetonitrile) in 31 min, 30-100% in 3 min, 100% solvent B for 5 min, and 7% solvent

B for 13 min. Full scan MS spectra from m/z 350-1500 were acquired at a resolution of 35,000 at

m/z 400 after accumulation to a target value of 3e6. Up to ten most intense precursor ions were

selected for fragmentation. HCD fragmentation was performed at normalized collision energy of

25% after the accumulation to a target value of 5e4. MS/MS was acquired at a resolution of 17,500.

In all cases nano-electrospray was performed at 1.7 kV using an in-house made gold-coated fused

silica capillary (o.d. 360 μm; i.d. 20 μm; tip i.d. 10 μm). Raw files were processed using Proteome

Discoverer 1.3 (Thermo Scientific, Bremen, Germany). The database search was performed against

the Swissprot human database, taxonomy (version November 2012) using Mascot (version 2.3,

Matrix Science, UK) as search engine. Carbamidomethylation of cysteines was set as a fixed

modification and oxidation of methionine was set as a variable modification. Trypsin was specified

as enzyme and up to two miss cleavages were allowed. Data filtering was performed using

percolator, resulting in 1% false discovery rate (FDR). Additional filter was Mascot ion score >20.

Raw files corresponding to one sample were merged into one result file.

Immunoprecipitation. 24 hours before IP HEK293T cells were transfected with relevant plasmids.

Cells were harvested using ice cold EBC buffer (140 mM NaCl, 50 mM Tris-HCl, 0.5% NP-40, 5

mM EDTA and protease inhibitor cocktail (Roche)) and briefly sonicated prior to 20 min

centrifugation at 20,000 *g*, 4°C. For FLAG, GFP and HA IP, cleared cell lysates were incubated

rotating for 1 hr at 4°C with 20 μl bead slurry of Anti-FLAG (M2) conjugated beads (Sigma, cat#

A2220) or 25 μl bead slurry of GFP-Trap®_A beads (Chromotek, cat# gta-20) or 20 μl bead slurry

of Pierce™ Anti-HA Agarose Conjugate (Thermo Fisher Scientific, cat#26181), respectively. The

immunocomplexes were washed 4 times with EBC buffer and eluted with SDS-PAGE sample

buffer. The eluted proteins were then subjected to SDS-PAGE and immunoblotting. All IP

experiments were repeated independently at least three times, and figures show representative

results from these independent repeats.

Protein concentration determination, SDS-PAGE and immunoblotting. The BioRad *DC*

Protein Assay was used to determine the protein concentration of cell lysates. SDS-PAGE and

immunoblotting was performed as described in (Schou et al., 2017).

Immunofluorescence microscopy. For IFM analysis, RPE1 cells were grown on glass coverslips, and subjected to 16-24 hrs serum-deprivation to induce ciliogenesis. Cells were washed in ice cold PBS, fixed with 4% paraformaldehyde (PFA) solution for 15 min at room temperature (RT) and cell membranes were permeabilized by incubation in 1x PBS with 0.2% (v/v) Triton X-100 and 1% (w/v) bovine serum albumin (BSA) for 12 min at RT. A blocking step of 30 min incubation at RT in 1x PBS with 2% BSA was done to avoid unspecific binding of antibodies, followed by incubation with primary antibodies overnight at 4°C. After three washing steps with BSA, cells were incubated in appropriate Alexa Fluor-conjugated secondary antibodies, diluted in BSA, for 45 min at RT and nuclei were labeled with DAPI. Coverslips were mounted using 1x PBS, 90% glycerol and 2% n-Propyl Gallate on glass slides and edges were sealed with nail polish. Fluorescence images were captured on a fully motorized Olympus BX63 upright microscope with an Olympus DP72 color, 12.8-megapixel, 4.140 x 3.096-resolution camera and with a fully motorized and automated Olympus IX83 Inverted microscope with a Hamamatsu ORCA-Flash 4.0 camera (C11440-22CU). The software used was Olympus CellSens dimension. Images were processed for publication using Adobe Photoshop version CS6. For quantifications of the centrosomal FLAG-Ap80 levels images were analyzed using ImageJ software. An outline was drawn around each centrosome and using the measurement- and integrated density (IntDens) functions, the Mean Green Fluorescence Intensity was measured in this area along with a background reading. The corrected mean fluorescence intensity in the centrosome was calculated by subtracting the corresponding background value. Procedures for confocal microscopy analysis of PALS1-A2E MDCK-II cells are described in (Tan et al., 2020).

Live cell imaging

Transfection and live cell imaging of RPE1 cells stably expressing SMO-tRFP were done as described previously (Juhl et al., 2021). Live cell imaging of Ap80-GFP and mCherry-KIF13B was performed using Total Internal Reflection Fluorescence (TIRF) microscopy. Inverted research microscope Nikon Eclipse Ti-E (Nikon), equipped with the perfect focus system (Nikon), Nikon Apo TIRF 100x N.A. 1.49 oil objective (Nikon) and iLas3 system (Dual Laser illuminator for azimuthal spinning TIRF (or Oblique) illumination and Simultaneous Targeted Laser Action including PhotoAblation; Gataca Systems) was used. The system was also equipped with ASI motorized stage MS-2000-XY (ASI), Photometrics Evolve 512 Delta EMCCD back illuminated

camera (Teledyne Photometrics) and controlled by the MetaMorph 7.8 software (Molecular

Devices). Stradus 488 nm (150 mW, Vortran) and OBIS 561 nm (100 mW, Coherent) lasers were

used as the light sources. We used ZT405/488/561/640rpc ZET405/488/561/635m filter set

(TRF89901, Chroma) together with Optosplit III beamsplitter (Cairn Research Ltd, UK) equipped

with double emission filter cube configured with ET525/50m, ET630/75m and T585LPXR

(Chroma). 16-bit images were projected onto the EMCCD chip with intermediate lens 2.5X (Nikon

C mount adapter 2.5X) at a magnification of 0.065 μm/pixel. To keep cells at 37°C we used stage

top incubator (model INUBG2E-ZILCS, Tokai Hit).

Statistical analysis. Statistical analysis was performed using GraphPad Prism 6 software.

Acknowledgements

We thank Søren Johansen, Emilie Baldram, Ida Hemmingsen, Signe Aagard and Sara Enevoldsen

for technical assistance, and Joseph Kissil, Athar Chisthi, Joel Pomerantz, Sadanori Watanabe,

Hiroaki Miki, Ronald Roepman, Francesc Garcia-Gonzalo, Lars Ellgaard, Dannel McCollum,

Christopher Westlake, and Jeffrey Miner for reagents. Supported by grants from Novo Nordisk

Foundation (NNF14OC0011535, NNF15OC0016886), Hartmann Fonden (A31662), Danish Cancer

Society (R146-A9590-16-S2), the University of Copenhagen (UCPH) Excellence Programme for

Interdisciplinary Research (LBP, STC), Department of Biology, UCPH (SKM), and the Netherlands

X-omics Initiative partially funded by NWO (project 184.034.019 to AA).

Gene name*	Protein ID*	KIF13B Tail 2		KIF13B Tail 3	
		Unique Peptide	PSM	Unique Peptide	PSM
KIF13B	Q9NQT8	147	294	102	177
MOV10	Q9HCE1	27	27	24	24
MARK2	Q7KZI7	26	31	17	19
HSPA5	P11021	22	24	16	18
KIDINS220	**Q9ULH0**	**22**	**22**	**8**	**8**
HADHA	P40939	20	21	11	11
UTRN	**P46939**	**19**	**19**	**2**	**2**
LRPPRC	P42704	17	17	10	10
MARK3	P27448	16	18	17	18
CUL3	Q13618	16	17	12	13
DLG1	**Q12959**	**15**	**16**	**0**	**0**
AMOT	**Q4VCS5**	**14**	**15**	**0**	**0**
MARK1	Q9P0L2	12	12	6	6
HADHB	P55084	11	12	10	10
ATAD3C	Q5T2N8	11	11	10	11
SNTB2	**Q13425**	**11**	**11**	**4**	**4**
NCL	P19338	9	9	9	9
MPP7	**Q5T2T1**	**9**	**9**	**0**	**0**
KLHL12	Q53G59	8	9	7	7
OSBPL8	Q9BZF1	8	9	5	5
PPP2R1A	P30153	5	5	4	4
ILF2	Q12905	5	7	3	3
CCT6A	P40227	5	5	3	3
TCP1	P17987	5	5	3	3
HNRNPH1	**P31943**	**5**	**5**	**2**	**2**
RPS11	**P62280**	**5**	**5**	**1**	**1**
PPP2R2A	**P63151**	**5**	**6**	**1**	**1**
CCT4	P50991	4	4	4	4
HNRNPA1	P09651	4	4	4	4
PKN3	Q6P5Z2	4	4	4	4
CSNK2A1	P68400	4	4	3	3
DDX6	P26196	4	4	3	3
LRRC59	Q96AG4	4	4	2	2
LIN7C	**Q9NUP9**	**4**	**4**	**0**	**0**
PDP1	**Q9P0J1**	**4**	**4**	**0**	**0**
CASK	**O14936**	**4**	**4**	**0**	**0**
KLHL7	Q8IXQ5	3	3	2	2
FAM98A	Q8NCA5	3	3	2	2
CCT7	Q99832	3	4	1	1
AIFM1	O95831	3	3	1	1
IRS4	O14654	3	3	1	1
DDX3X	O00571	3	3	1	1
DTNB	**O60941**	**3**	**4**	**0**	**0**
FBXW11	Q9UKB1	3	3	0	0
XPR1	Q9UBH6	3	3	0	0
DTNA	**Q9Y4J8**	**3**	**3**	**0**	**0**
CEP104	O60308	3	3	0	0
GLG1	Q92896	3	3	0	0

Table 1. Binding partners of the indicated BirAGFP-KIF13B constructs in HEK293T cells

identified by mass spectrometry analysis.

*The highlighted genes/proteins preferentially bind the KIF13B Tail 2 over Tail 3 and are discussed

in the text. Only proteins identified by at least three unique peptides in the BirAGFP-KIF13B Tail 2

sample are included. PSM, peptide spectrum matches.

Figure legends

Figure 1. Ap80 interacts with KIF13B via its N-terminus. (**A, D**) FLAG IP of cells co-

expressing FLAG-tagged Ap80 or Ap130 fusions and KIF13B-HA or KIF13B-Δmotor-HA. (**B**)

GFP IP of cells co-expressing FLAG-Ap80 and GFP-KIF13B. Input (in) and IP pellet (IP) fractions

were subjected to Immunoblotting with indicated antibodies. (**C**) Diagram of full length Ap80 and

Ap130 and FLAG-Ap80 truncations used. CC/BAR, coiled-coil/BAR domain; ABD, angiostatin

binding domain; PDZ-BD, PDZ-binding domain.

Figure 2. Ap80 binds KIF13B tail region and promotes KIF13B-PALS1 association. (**A**)

Diagram of full length and truncated KIF13B fusions; those binding to Ap80 are in bold. FHA,

Forkhead-associated domain; DUF, domain of unknown function; red asterisks, MARK2

phosphorylation sites; PRR, proline-rich region; CG, CAP-Gly domain. Redrawn from (Yoshimura

et al., 2010, Soppina et al., 2014, Yamada et al., 2014, Zhu et al., 2016, Schou et al., 2017). (**B**)

FLAG IP of cells co-expressing FLAG-Ap80-NB and GFP (Vector) or different GFP-KIF13B

truncations. Input (in) and IP pellet (IP) fractions were analyzed by immunoblotting using indicated

antibodies. (**C-F**) FLAG (**C-E**) or GFP (**F**) IP of cells expressing indicated fusion proteins, analyzed

with indicated antibodies. GFP-KIF13BΔm: GFP-KIF13B-Δmotor. (**G**) Model for Ap80-KIF13B

interaction.

Figure 3. Ap80 localizes to the ciliary base and regulates ciliary length in RPE1 cells. (**A**)

Frames of time-lapse movie of a cell co-expressing SMO-tRFP and Ap80-GFP. Asterisk: ciliary

base. Time in seconds. (**B, C**) Snapshots of live cells co-expressing SMO-tRFP and PALS1-GFP in

the absence (**B**) or presence (**C**) of FLAG-Ap80. Asterisk: ciliary base. (**D**) Immunoblot of WT and

AMOT[-/-] cells using indicated antibodies. (**E**) IFM of serum-deprived WT (parental) and *AMOT*[-/-]

cells using acetylated α-tubulin (AcTub) and ARL13B as cilia markers (arrows). DNA was stained

with DAPI. (**F**) Cilia length quantification in WT and *AMOT*[-/-] cells; >69 cilia measured in total per

condition (n=3). **(G)** Cilia length quantification in GFP or GFP-Ap80 expressing cells; 18-35 cilia

measured per condition (n=3). Data in (F) and (G) show mean□±□SD. ****, P≤0.0001 (unpaired,

two-tailed Student's *t* test).

References

AKELLA, J. S. & BARR, M. M. 2021. The tubulin code specializes neuronal cilia for extracellular
vesicle release. *Dev Neurobiol,* 81**,** 231-252.

ASABA, N., HANADA, T., TAKEUCHI, A. & CHISHTI, A. H. 2003. Direct Interaction with a
Kinesin-related Motor Mediates Transport of Mammalian Discs Large Tumor Suppressor
Homologue in Epithelial Cells. *Journal of Biological Chemistry,* 278**,** 8395-8400.

BANGS, F. & ANDERSON, K. V. 2017. Primary Cilia and Mammalian Hedgehog Signaling. *Cold*
Spring Harb Perspect Biol, 9.

BAZELLIERES, E., AKSENOVA, V., BARTHELEMY-REQUIN, M., MASSEY-HARROCHE,
D. & LE BIVIC, A. 2018. Role of the Crumbs proteins in ciliogenesis, cell migration and actin
organization. *Semin Cell Dev Biol,* 81**,** 13-20.

BOHL, J., BRIMER, N., LYONS, C. & VANDE POL, S. B. 2007. The stardust family protein
MPP7 forms a tripartite complex with LIN7 and DLG1 that regulates the stability and localization
of DLG1 to cell junctions. *J Biol Chem,* 282**,** 9392-400.

BOLIS, A., COVIELLO, S., VISIGALLI, I., TAVEGGIA, C., BACHI, A., CHISHTI, A. H.,
HANADA, T., QUATTRINI, A., PREVITALI, S. C., BIFFI, A. & BOLINO, A. 2009. Dlg1, Sec8,
and Mtmr2 regulate membrane homeostasis in Schwann cell myelination. *J Neurosci,* 29**,** 8858-70.

BRATT, A., BIROT, O., SINHA, I., VEITONMAKI, N., AASE, K., ERNKVIST, M. &
HOLMGREN, L. 2005. Angiomotin regulates endothelial cell-cell junctions and cell motility. *J*
Biol Chem, 280**,** 34859-69.

CHRISTENSEN, S. T., MORTHORST, S. K., MOGENSEN, J. B. & PEDERSEN, L. B. 2017.
Primary Cilia and Coordination of Receptor Tyrosine Kinase (RTK) and Transforming Growth
Factor beta (TGF-beta) Signaling. *Cold Spring Harb Perspect Biol,* 9.

DELOUS, M., HELLMAN, N. E., GAUDE, H. M., SILBERMANN, F., LE BIVIC, A.,
SALOMON, R., ANTIGNAC, C. & SAUNIER, S. 2009. Nephrocystin-1 and nephrocystin-4 are
required for epithelial morphogenesis and associate with PALS1/PATJ and Par6. *Hum Mol Genet,*
18**,** 4711-23.

DRUMMOND, M. L., LI, M., TARAPORE, E., NGUYEN, T. T. L., BAROUNI, B. J., CRUZ, S.,
TAN, K. C., ORO, A. E. & ATWOOD, S. X. 2018. Actin polymerization controls cilia-mediated
signaling. *J Cell Biol,* 217**,** 3255-3266.

ERNKVIST, M., AASE, K., UKOMADU, C., WOHLSCHLEGEL, J., BLACKMAN, R.,
VEITONMAKI, N., BRATT, A., DUTTA, A. & HOLMGREN, L. 2006. p130-angiomotin
associates to actin and controls endothelial cell shape. *Febs j,* 273**,** 2000-11.

FAN, S., HURD, T. W., LIU, C. J., STRAIGHT, S. W., WEIMBS, T., HURD, E. A., DOMINO, S.
E. & MARGOLIS, B. 2004. Polarity proteins control ciliogenesis via kinesin motor interactions.
Curr Biol, 14**,** 1451-61.

GARCIA-GONZALO, F. R. & REITER, J. F. 2017. Open Sesame: How Transition Fibers and the
Transition Zone Control Ciliary Composition. *Cold Spring Harb Perspect Biol,* 9.

HANADA, T., LIN, L., TIBALDI, E. V., REINHERZ, E. L. & CHISHTI, A. H. 2000. GAKIN, a
novel kinesin-like protein associates with the human homologue of the Drosophila discs large tumor
suppressor in T lymphocytes. *J Biol Chem,* 275**,** 28774-84.

HELLER, B., ADU-GYAMFI, E., SMITH-KINNAMAN, W., BABBEY, C., VORA, M., XUE, Y.,
BITTMAN, R., STAHELIN, R. V. & WELLS, C. D. 2010. Amot recognizes a juxtanuclear
endocytic recycling compartment via a novel lipid binding domain. *J Biol Chem,* 285**,** 12308-20.

HIROKAWA, N. & TANAKA, Y. 2015. Kinesin superfamily proteins (KIFs): Various functions
and their relevance for important phenomena in life and diseases. *Exp Cell Res,* 334**,** 16-25.

HORIGUCHI, K., HANADA, T., FUKUI, Y. & CHISHTI, A. H. 2006. Transport of PIP_3 by
GAKIN, a kinesin-3 family protein, regulates neuronal cell polarity. *J Cell Biol,* 174**,** 425-436.

JUHL, A. D., ANVARIAN, Z., BERGES, J., WÜSTNER, D. & PEDERSEN, L. B. 2021. Transient
accumulation, bidirectional movement and tip release of KIF13B in primary cilia. *bioRxiv***,**
2021.07.09.451768.

KANAI, Y., WANG, D. & HIROKAWA, N. 2014. KIF13B enhances the endocytosis of LRP1 by
recruiting LRP1 to caveolae. *J Cell Biol,* 204**,** 395-408.

KRAMARCY, N. R., VIDAL, A., FROEHNER, S. C. & SEALOCK, R. 1994. Association of
utrophin and multiple dystrophin short forms with the mammalian M(r) 58,000 dystrophin-
associated protein (syntrophin). *J Biol Chem,* 269**,** 2870-6.

LAMASON, R. L., KUPFER, A. & POMERANTZ, J. L. 2010. The dynamic distribution of
CARD11 at the immunological synapse is regulated by the inhibitory kinesin GAKIN. *Mol Cell,* 40**,**
798-809.

LEE, S., FAN, S., MAKAROVA, O., STRAIGHT, S. & MARGOLIS, B. 2002. A novel and
conserved protein-protein interaction domain of mammalian Lin-2/CASK binds and recruits SAP97
to the lateral surface of epithelia. *Mol Cell Biol,* 22**,** 1778-91.

LEVCHENKO, T., BRATT, A., ARBISER, J. L. & HOLMGREN, L. 2004. Angiomotin expression
promotes hemangioendothelioma invasion. *Oncogene,* 23**,** 1469-73.

LU, Q., INSINNA, C., OTT, C., STAUFFER, J., PINTADO, P. A., RAHAJENG, J., BAXA, U.,
WALIA, V., CUENCA, A., HWANG, Y. S., DAAR, I. O., LOPES, S., LIPPINCOTT-
SCHWARTZ, J., JACKSON, P. K., CAPLAN, S. & WESTLAKE, C. J. 2015. Early steps in
primary cilium assembly require EHD1/EHD3-dependent ciliary vesicle formation. *Nat Cell Biol,*
17**,** 228-240.

MILLS, J., HANADA, T., HASE, Y., LISCUM, L. & CHISHTI, A. H. 2019. LDL receptor related
protein 1 requires the I3 domain of discs-large homolog 1/DLG1 for interaction with the kinesin
motor protein KIF13B. *Biochim Biophys Acta Mol Cell Res,* 1866**,** 118552.

MOLEIRINHO, S., GUERRANT, W. & KISSIL, J. L. 2014. The Angiomotins--from discovery to
function. *FEBS Lett,* 588**,** 2693-703.

MORTHORST, S. K., CHRISTENSEN, S. T. & PEDERSEN, L. B. 2018. Regulation of ciliary
membrane protein trafficking and signalling by kinesin motor proteins. *FEBS J,* 285**,** 4535-4564.

NEUBRAND, V. E., CESCA, F., BENFENATI, F. & SCHIAVO, G. 2012. Kidins220/ARMS as a
functional mediator of multiple receptor signalling pathways. *J Cell Sci,* 125**,** 1845-54.

NOSEDA, R., GUERRERO-VALERO, M., ALBERIZZI, V., PREVITALI, S. C., SHERMAN, D.
L., PALMISANO, M., HUGANIR, R. L., NAVE, K. A., CUENDA, A., FELTRI, M. L., BROPHY,
P. J. & BOLINO, A. 2016. Kif13b Regulates PNS and CNS Myelination through the Dlg1 Scaffold.
PLoS Biol, 14**,** e1002440.

OSSIPOVA, O., CHU, C. W., FILLATRE, J., BROTT, B. K., ITOH, K. & SOKOL, S. Y. 2015.
The involvement of PCP proteins in radial cell intercalations during Xenopus embryonic
development. *Dev Biol,* 408**,** 316-27.

PETERS, M. F., ADAMS, M. E. & FROEHNER, S. C. 1997. Differential association of syntrophin
pairs with the dystrophin complex. *J Cell Biol,* 138**,** 81-93.

PRULIERE, G., COSSON, J., CHEVALIER, S., SARDET, C. & CHENEVERT, J. 2011. Atypical
protein kinase C controls sea urchin ciliogenesis. *Mol Biol Cell,* 22**,** 2042-53.

PUGACHEVA, E. N., JABLONSKI, S. A., HARTMAN, T. R., HENSKE, E. P. & GOLEMIS, E.
A. 2007. HEF1-dependent Aurora A activation induces disassembly of the primary cilium. *Cell,*
129**,** 1351-63.

RAN, F. A., HSU, P. D., WRIGHT, J., AGARWALA, V., SCOTT, D. A. & ZHANG, F. 2013.
Genome engineering using the CRISPR-Cas9 system. *Nat Protoc,* 8**,** 2281-2308.

REN, J., WANG, S., CHEN, H., WANG, W., HUO, L. & FENG, W. 2018. Coiled-coil 1-mediated
fastening of the neck and motor domains for kinesin-3 autoinhibition. *Proc Natl Acad Sci U S A,*
115**,** E11933-E11942.

RICHNAU, N. & ASPENSTROM, P. 2001. Rich, a rho GTPase-activating protein domain-
containing protein involved in signaling by Cdc42 and Rac1. *J Biol Chem,* 276**,** 35060-70.

SANG, L., MILLER, J. J., CORBIT, K. C., GILES, R. H., BRAUER, M. J., OTTO, E. A., BAYE,
L. M., WEN, X., SCALES, S. J., KWONG, M., HUNTZICKER, E. G., SFAKIANOS, M. K.,
SANDOVAL, W., BAZAN, J. F., KULKARNI, P., GARCIA-GONZALO, F. R., SEOL, A. D.,
O'TOOLE, J. F., HELD, S., REUTTER, H. M., LANE, W. S., RAFIQ, M. A., NOOR, A., ANSAR,
M., DEVI, A. R., SHEFFIELD, V. C., SLUSARSKI, D. C., VINCENT, J. B., DOHERTY, D. A.,
HILDEBRANDT, F., REITER, J. F. & JACKSON, P. K. 2011. Mapping the NPHP-JBTS-MKS
protein network reveals ciliopathy disease genes and pathways. *Cell,* 145**,** 513-28.

SCHOU, K. B., MOGENSEN, J. B., MORTHORST, S. K., NIELSEN, B. S., ALELIUNAITE, A.,
SERRA-MARQUES, A., FURSTENBERG, N., SAUNIER, S., BIZET, A. A., VELAND, I. R.,
AKHMANOVA, A., CHRISTENSEN, S. T. & PEDERSEN, L. B. 2017. KIF13B establishes a

CAV1-enriched microdomain at the ciliary transition zone to promote Sonic hedgehog signalling. *Nat Commun,* 8**,** 14177.

SERRA-MARQUES, A., MARTIN, M., KATRUKHA, E. A., GRIGORIEV, I., PEETERS, C. A., LIU, Q., HOOIKAAS, P. J., YAO, Y., SOLIANOVA, V., SMAL, I., PEDERSEN, L. B., MEIJERING, E., KAPITEIN, L. C. & AKHMANOVA, A. 2020. Concerted action of kinesins KIF5B and KIF13B promotes efficient secretory vesicle transport to microtubule plus ends. *Elife,* 9.

SFAKIANOS, J., TOGAWA, A., MADAY, S., HULL, M., PYPAERT, M., CANTLEY, L., TOOMRE, D. & MELLMAN, I. 2007. Par3 functions in the biogenesis of the primary cilium in polarized epithelial cells. *J Cell Biol,* 179**,** 1133-40.

SIDDIQUI, N. & STRAUBE, A. 2017. Intracellular Cargo Transport by Kinesin-3 Motors. *Biochemistry (Mosc),* 82**,** 803-815.

SOPPINA, V., NORRIS, S. R., DIZAJI, A. S., KORTUS, M., VEATCH, S., PECKHAM, M. & VERHEY, K. J. 2014. Dimerization of mammalian kinesin-3 motors results in superprocessive motion. *Proceedings of the National Academy of Sciences of the United States of America,* 111**,** 5562-5567.

TAN, B., YATIM, S., PENG, S., GUNARATNE, J., HUNZIKER, W. & LUDWIG, A. 2020. The Mammalian Crumbs Complex Defines a Distinct Polarity Domain Apical of Epithelial Tight Junctions. *Curr Biol,* 30**,** 2791-2804 e6.

TARBASHEVICH, K., DZEMENTSEI, A. & PIELER, T. 2011. A novel function for KIF13B in germ cell migration. *Dev Biol,* 349**,** 169-78.

TASCHNER, M. & LORENTZEN, E. 2016. The Intraflagellar Transport Machinery. *Cold Spring Harb Perspect Biol,* 8.

TROYANOVSKY, B., LEVCHENKO, T., MANSSON, G., MATVIJENKO, O. & HOLMGREN, L. 2001. Angiomotin: an angiostatin binding protein that regulates endothelial cell migration and tube formation. *J Cell Biol,* 152**,** 1247-54.

WELLS, C. D., FAWCETT, J. P., TRAWEGER, A., YAMANAKA, Y., GOUDREAULT, M., ELDER, K., KULKARNI, S., GISH, G., VIRAG, C., LIM, C., COLWILL, K., STAROSTINE, A., METALNIKOV, P. & PAWSON, T. 2006. A Rich1/Amot complex regulates the Cdc42 GTPase and apical-polarity proteins in epithelial cells. *Cell,* 125**,** 535-48.

YAMADA, K. H., HANADA, T. & CHISHTI, A. H. 2007. The Effector Domain of Human Dlg Tumor Suppressor Acts as a Switch That Relieves Autoinhibition of Kinesin-3 Motor GAKIN/KIF13B. *Biochemistry,* 46**,** 10039-10045.

YAMADA, K. H., NAKAJIMA, Y., GEYER, M., WARY, K. K., USHIO-FUKAI, M., KOMAROVA, Y. & MALIK, A. B. 2014. KIF13B regulates angiogenesis through Golgi to plasma membrane trafficking of VEGFR2. *J Cell Sci,* 127**,** 4518-30.

YOSHIMURA, Y., TERABAYASHI, T. & MIKI, H. 2010. Par1b/MARK2 phosphorylates kinesin-like motor protein GAKIN/KIF13B to regulate axon formation. *Mol Cell Biol,* 30**,** 2206-19.

606   ZHU, J., SHANG, Y., XIA, Y., ZHANG, R. & ZHANG, M. 2016. An Atypical MAGUK GK
607   Target Recognition Mode Revealed by the Interaction between DLG and KIF13B. *Structure,* 24**,**
608   1876-1885.

Figure 1

Figure 2

Figure 3