A Genomic Perspective on the Evolutionary Diversification of Turtles

Simone M. Gable[1*], Michael I. Byars[1*], Robert Literman[2], Marc Tollis[1]

¹School of Informatics, Computing, and Cyber Systems, Northern Arizona University, PO Box

5693, Flagstaff, AZ 8601, USA

²Department of Biological Sciences, University of Rhode Island, 120 Flagg Road, Kingstown, RI,

0288, USA

*These authors contributed equally.

Corresponding author: Marc Tollis, School of Informatics, Computing, and Cyber Systems,

Northern Arizona University, PO Box 5693, Flagstaff, AZ, 86011; marc.tollis@nau.edu

Running header: Phylogenomic Heterogeneity in Turtles

ABSTRACT

To examine phylogenetic heterogeneity in turtle evolution, we collected thousands of high-

confidence single-copy orthologs from 19 genome assemblies representative of extant turtle

diversity and estimated a phylogeny with multispecies coalescent and concatenated partitioned

methods. We also collected next-generation sequences from 26 turtle species and assembled

millions of biallelic markers to reconstruct phylogenies based on annotated regions from the

western painted turtle (*Chrysemys picta bellii*) genome (coding regions, introns, untranslated

regions, intergenic, and others). We then measured gene tree-species tree discordance, as well as

gene and site heterogeneity at each node in the inferred trees, and tested for temporal patterns in

phylogenomic conflict across turtle evolution. We found strong and consistent support for all

bifurcations in the inferred turtle species phylogenies. However, a number of genes, sites, and

genomic features supported alternate relationships between turtle taxa. Our results suggest that

gene tree-species tree discordance in these datasets is likely driven by population-level processes

such as incomplete lineage sorting. We found very little effect of substitutional saturation on

species tree topologies, and no clear phylogenetic patterns in codon usage bias and compositional

heterogeneity. There was no correlation between gene and site concordance, node age, and DNA

substitution rate across most annotated genomic regions. Our study demonstrates that

heterogeneity is to be expected even in well resolved clades such as turtles, and that future

phylogenomic studies should aim to sample as much of the genome as possible in order to obtain

accurate phylogenies for assessing conservation priorities in turtles.

Keywords: turtles, genomes, phylogeny, discordance

Introduction

Next-generation sequencing has greatly increased the numbers of sampled loci for phylogenomic

analyses (Lemmon et al. 2012; Faircloth et al. 2013; Edwards et al. 2016), shedding new light on

the branching order of diversification across the history of life. However, phylogenetic studies

based on genome-scale data often contain a great deal of heterogeneity in the inferred patterns

(Hime et al. 2021; Lopes et al. 2021; Morales-Briones et al. 2021; Singhal et al. 2021),

particularly in the form of discordance among sites and gene trees (Kumar et al. 2012). Gene

tree-species tree discordance occurs when gene trees are in conflict with the underlying species

relationships, while site discordance occurs when a subset of sites support different bifurcations

in a species tree, and both factors can lead to biased phylogenetic estimates. Understanding

patterns and potential sources of phylogenomic heterogeneity will not only settle phylogenetic

arguments and reduce bias in tree building, but also provide better models with which to

understand trait evolution and help define conservation priorities through comparative methods

(Garland et al. 2005; Colston et al. 2021).

Biological factors driving phylogenomic heterogeneity include: differing evolutionary

rates across the genome, incomplete lineage sorting, and hidden paralogy (Pamilo and Nei 1988;

Galtier and Daubin 2008). Artifacts that drive heterogeneity include sequencing errors and

substitution model violations such as genetic saturation, compositional heterogeneity, and codon

usage bias (Foster 2004; Cooper 2014; Cox et al. 2014). Accounting for gene and site

discordance in phylogenomic estimation often requires partitioning schemes which try to account

for differing evolutionary rates and model parameters (Kubatko and Degnan 2007), methods that

are consistent with the multispecies coalescent accounting for population-level processes such as

incomplete lineage sorting (Edwards et al. 2016), as well as strict filtering of sequence

alignments that aim to reduce potential errors. The application of hundreds or even thousands of

loci is often an attempt to overcome biases stemming from or driving phylogenomic conflict, and

these studies often boast strong branch support as measured by bootstrap replicates or posterior

probabilities, likely as a result of reduced sampling variance in large datasets (Minh et al. 2020a).

However, competing phylogenomic studies of the same clades that use different datasets and tree

building methods can produce conflicting results, despite high levels of statistical support (see

modern birds: Jarvis et al. 2014; Prum et al. 2015).

An emerging advantage of the genomic era is the theoretical ability to access the whole

genome in order to sample loci representing a diversity of evolutionary rates and coalescent

histories (Wolf et al. 2002; Rokas et al. 2003). The most widely used sampling methods include

the use of large numbers of coding sequences (Singhal et al. 2021), anchored hybrid enrichment

loci (AHE; Lemmon et al. 2012), ultra-conserved elements (UCEs; Faircloth et al. 2013), and

transcriptomes (Irisarri et al. 2017), but these are each still reduced representations of a relatively

small proportion of the genome (Lynch 2007). Meanwhile, the development of bioinformatics

tools and sequencing technologies have enabled chromosome-scale genome assemblies for non-

model organisms (see http://dnazoo.org, Dudchenko et al. 2017), potentially providing insights

to the genomic regions or types of loci that drive heterogeneous phylogenetic results. Thus, by

accessing complete genomes and studying heterogeneity closely, we can more fully comprehend

conflicting results across gene trees, obtain more accurate perspectives of species relationships,

and gain a fuller understanding of molecular evolution across the genome.

Here, we examined the importance of genome-wide heterogeneity in the phylogenomic

estimation of turtles (Order Testudines). Turtles are a near-globally distributed and

morphologically distinct clade of shelled reptiles, with a rich fossil history dating from the

Triassic Period (Gaffney 1980; Joyce 2007; Joyce et al. 2021). Despite their persistence in the

fossil record and across modern ecosystems, almost half of turtle species are listed in the

International Union for Conservation of Nature (IUCN) as endangered, critically endangered, or

vulnerable to extinction (IUCN, 2021). The approximately 350 extant species of turtles are

classified into two main groups, Pleurodira ("side-necked") and Cryptodira ("hidden-necked"),

which are further divided into 8 superfamilies and 14 families (Uetz et al. 2021). Early molecular

phylogenetic results disagreed in the placement of some turtle taxa; in particular, the position of

the monotypic big-headed turtle (*Platysternon*) and its relationships with or within Testudinoidea

(Shaffer et al. 1997; Krenz et al. 2005). Other questions about deeper turtle relationships such as

the position of soft-shell turtles (Trionychoidea), the Americhelydiae, and the position of turtles

in the amniote phylogeny persisted into the next-generation sequencing era (Chiari et al. 2012;

Crawford et al. 2012, 2015; Brown and Thomson 2017). Since then, phylogenomic studies of

turtles have produced fully resolved species trees with 100% statistical support at every node and

near complete agreement across studies as measured by bootstrap replicates or posterior

probabilities (Crawford et al. 2015; Shaffer et al. 2017).

Phylogenomic heterogeneity has been examined in vertebrate groups such as mammals

(Tarver et al. 2016; Liu et al. 2017) squamates (Burbrink et al. 2020; Singhal et al. 2021), birds

(Jarvis et al. 2014; Prum et al. 2015), and amphibians (Hime et al. 2021), but there has not yet

been an investigation into how gene tree-species tree discordance across the genome affects the

inference of relationships between the major turtle lineages. While phylogenetic conflict often

belies more traditional means of measuring certainty via bootstrap support or posterior

probabilities (Minh et al. 2020a), the study of turtle systematics has not yet truly taken advantage

of the genomic era; this is despite ample genomic resources being available for a wide range of

turtle species (Shaffer et al. 2013; Wang et al. 2013; Tollis et al. 2017; Quesada et al. 2019).

Since evolutionary distinctiveness is linked to conservation status in turtles (Colston et al. 2020),

and many turtle species have adaptations of physiological and developmental importance such as

longevity, resistance to hypoxia, cold tolerance, and the anatomical changes responsible for the

turtle shell, a study using genome-scale data to measure phylogenomic heterogeneity is needed in

order to obtain the most accurate models of turtle evolution. Our goals in this study were

threefold: (1) to revisit discoveries about higher turtle systematics using substantial proportions

of the genome; (2) to measure heterogeneity at genes and sites in important events in turtle

evolution; and (3) determine what drives heterogeneity in turtle phylogenomics in terms of

mechanisms and locus type.

119   To accomplish these goals, we generated two distinct yet overlapping genomic datasets

for turtles. The first comprises 5,310 high-confidence single-copy orthologs bioinformatically

extracted from the genome assemblies of 19 turtles plus three outgroups, aligned and

phylogenetically analyzed using multilocus coalescent-consistent and concatenated partition-

based methods. The second dataset (26 species, including all turtle species from the first method)

consisted of 1,655,675 biallelic parsimony-informative sites that we extracted using mapped

Illumina sequence reads. We assembled the reads into contigs that were further mapped to a

reference turtle genome (the western painted turtle, *Chrysemys picta bellii*). Based on genome

annotations, we partitioned sites by locus type including coding regions (CDS), introns, 5'-UTR,

3'-UTR, intergenic, pseudogenes, lncRNA, and smRNA, and reconstructed separate phylogenies

using each locus type. With the first dataset we assessed patterns and sources of heterogeneity in

turtle phylogenomics with a large number of DNA sequence alignments, while with the second

dataset we compared topologies reconstructed with parsimony-informative sites extracted from

different annotated regions across the turtle genome. Taken together, this scope of analyses is

unprecedented in turtle phylogenomics. We confirm that heterogeneity in phylogenomic datasets

is to be expected, even in well-resolved clades such as turtles, and suggest that a combination of

processes is driving the incongruence between previous studies of turtle relationships.

Methods and Materials

Phylogenomic analysis of high-confidence single-copy orthologs

We downloaded the complete genome assemblies of 19 turtle species plus three outgroup taxa

from NCBI. The species, assembly accession numbers, assembly lengths, and assembly

contiguities measured by scaffold N50 are shown in Table 1. Using BUSCO v3 (Waterhouse et

al. 2018), we extracted the nucleotide sequences of 5,310 OrthoDB v9 tetrapod orthologs

(Zdobnov et al. 2017) from each genome assembly. BUSCO orthologs are high-confidence genes

that persist in eukaryotic genomes as single-copy, which precludes downstream problems in

phylogenetics stemming from gene duplication such as hidden paralogy (Waterhouse et al.

2018). We aligned each set of orthologs with MAFFT v7.475 (Katoh and Standley 2013), and

removed erroneous columns in the alignments with the heuristic method (*automated 1*) in

TrimAL v1.4.1 (Capella-Gutierrez et al. 2009). Summary statistics including number of taxa,

alignment length, missing data percentage, number and proportion of variable sites, number and

proportion of parsimony-informative sites, and GC content for each alignment were estimated

with AMAS v3.04 (Borowiec 2016). Trimmed alignments were filtered at a cut-off length

≥1,500 bp and a minimum of 50% taxa representation. We estimated the average pairwise

Kimura 2-parameter corrected distance for each alignment with MEGAX (Kumar et al. 2018).

Table 1. Genomic data for turtles analyzed in this study.

Suborder	Superfamily/Clade	Family	Species Name	Assembly Accession	Assembly Length (Gb)	Scaffold N50 (Mb)	Trimmed Read Depth*
Cryptodira	Chelonioidea	Cheloniidae	*Chelonia mydas*	GCF_000344595.1	2.2	3.9	12.8X
		Dermochelyidae	*Dermochelys coriacea*	GCA_006547105.1	2.1	0.12	31.3X
	Chelydridae	Chelydridae	*Chelydra serpentina*	GCA_007922165.1	2.5	21	16.1X
	Kinosternoidea	Dermatemydidae	*Dermatemys mawii*	GCA_007922305.1	1.9	34.4	39.7X
		Kinosternidae	*Sternotherus carinatus*	NA	NA	NA	45.9X
	Testudinoidea	Emydidae	*Chrysemys picta*	GCA_000241765.3	2.4	16	10.3X
			Malaclemys terrapin	GCA_001728815.2	2.4	0.44	9.3X
			Terrapene mexicana	GCF_002925995.2	2.6	24.2	12.4X
			Emys orbicularis	NA	NA	NA	1.3X
			Trachemys scripta	NA	NA	NA	40.2X
			Actinemys marmorata	NA	NA	NA	41.4X
		Geoemydidae	*Cuora amboinensis*	GCA_004028625.2	2.1	0.25	23.1X
			Cuora mccordi	GCA_003846335.1	2.4	32.6	70.6X
			Mauremys reevesii	NA	NA	NA	34.1X
			Batagur trivittata	NA	NA	NA	1.1X
		Platysternidae	*Platysternon megacephalum*	GCA_003942145.1	2.3	7.2	16.7X
		Testudinidae	*Chelonoidis abingdonii*	GCA_003597395.1	2.3	1.3	17.3X
			Gopherus agassizii	GCA_002896415.1	2.2	0.228	40.5X
			Aldabrachelys gigantea	NA	NA	NA	21.9X
	Trionychoidea	Carettochelyidae	*Carettochelys insculpta*	GCA_007922185.1	2.3	45.9	49.7X
		Trionychidae	*Apalone spinifera*	GCA_000385615.1	1.9	2.3	10.3X
			Pelodiscus sinensis	GCF_000230535.1	2.2	3.4	21.1X

Pleurodira	Chelidoidea	Chelidae	*Emydura subglobosa*	GCA_007922225.1	2	44.8	40.6X
			Mesoclemmys tuberculata	GCA_007922155.1	2	46.4	49.0X
	Pelomedusoidea	Pelomedusidae	*Pelusios castaneus*	GCA_007922175.1	2	14.1	47.5X
		Podocnemididae	*Podocnemis expansa*	GCA_007922195.1	2.4	37.1	10.5X

Gb=gigabases; Mb=megabases; *Short Read Archive Accession numbers are in Supplementary Data; NA=no

genome assembly was analyzed in the current study

Gene trees were reconstructed for each locus using RAxML v1.8 (Stamatakis 2014). For

each ortholog, we generated 10 maximum likelihood trees with the GTRGAMMA substitution

model, and performed 100 rapid bootstrap replicates on the best tree. We then collected the best

tree for each ortholog to use as evidence to construct a species tree with ASTRAL-III (Zhang et

al. 2018), assessing branch support with local posterior probabilities (Sayyari and Mirarab 2016).

We used Snakemake (Mölder et al. 2021) to automate a reproducible bioinformatics pipeline for

BUSCO extraction, filtering, alignment, gene tree and species tree estimation, available at

https://github.com/mibyars/busco_phylo_pipeline. We also concatenated the single copy

orthologs into a supermatrix with AMAS and performed a partitioned maximum likelihood

phylogenetic reconstruction with IQ-TREE v2.1.2 (Minh et al. 2020b). We used ModelFinder

(Kalyaanamoorthy et al. 2017) set to TESTMERGE to combine partitions where necessary. We

assessed branch support on the concatenated partitioned maximum likelihood tree using 1,000

ultrafast bootstrap replicates and the Shimodaira-Hasegawa-like (SH-like) approximate

likelihood ratio test (aLRT, Shimodaira 2002).

Phylogenomic analysis of biallelic markers

In addition to single-copy orthologs from turtle and outgroup genome assemblies, we constructed

sets of biallelic single nucleotide polymorphisms (SNPs) using paired-end sequence data from

multiple turtle species downloaded from the NCBI Short Read Archive (SRA, Table 1,

Supplementary Data). We assessed each SRA dataset with FastQC v0.115 (S. Andrews -

http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and trimmed the raw reads with

BBDuk v.37.41 (B. Bushnell - sourceforge.net/projects/bbmap/, last accessed December 2020).

We used SISRS to generate de novo orthologs from the processed reads (Schwartz et al. 2015;

Literman and Schwartz 2021). After subsampling all sequences to a target depth of ~10X per

taxon (except for two low-coverage species, Table 1, Supplementary Data), we assembled a

composite genome using Ray v2.2.3-devel (Boisvert et al. 2010) with default parameters and a k-

value of 31. We then used SISRS to map the trimmed reads from each species to the composite

genome, retaining only uniquely mapped reads to avoid false positives from duplicated or

repetitive regions. Bases in the composite genome were replaced according to mapped data from

each species if sites were covered by at least three reads and if there was no within-taxon

variation; sites not meeting these criteria were masked as "N".

For downstream ortholog annotation, we mapped each scaffold from the composite

sequences against the reference genome for the western painted turtle (*Chrysemys picta bellii,*

GCA_000241765.3) with Bowtie2 v2.3.4 (Langmead and Salzberg 2012), removing contigs that

either failed to map or mapped to multiple positions, and removing sites resulting from

overlapping coverage of independent scaffolds. We then annotated the mapped composite

genome sites parsed from the painted turtle genome as belonging to the following genomic

regions (locus types) with BEDTools v2.2.6 (Quinlan and Hall 2010): (1) coding sequences

(CDS); (2) 3'-UTR; (3) 5'-UTR; (4) introns; (5) other genic regions not annotated as CDS, UTR,

or intronic; (6) long-noncoding RNAs (lncRNAs); (7) pseudogenes; (8) smRNAs (including

miRNAs, ncRNAs, rRNAs, scRNAs, smRNAs, snoRNAs, snRNAs, tRNAs, and vaultRNAs).;

and (9) intergenic regions. We defined "intergenic" as all regions not included within the other

locus types.

We used SISRS to identify parsimony-informative sites along the mapped composite

contigs and to remove invariant sites and singletons, as well as any sites containing "Ns" (i.e.

polymorphic within species) and overlapping indels or gaps. For the nine annotated genomic

regions as well as a set of the complete mapped data (=10 locus types), we concatenated biallelic

parsimony-informative sites and inferred locus type phylogenies via SVDquartets (Chifman and

Kubatko 2014) in PAUP* v4.0a (build 168) (Swofford 2003) using 1,000 bootstrap replicates

and estimated majority consensus trees.

Heterogeneity analyses

To measure similarity across genealogies, we calculated Robinson-Foulds (RF) distances

(Robinson and Foulds 1981) with PhyKIT (Steenwyk et al. 2021). First, we measured the RF

distance between each inferred single-copy ortholog gene tree and the inferred species tree. In

order to take into account missing species across some of the gene trees, we normalized the RF

distances by taking the raw RF distance and dividing it by $2(n\text{-}3)$ where n is the number of tips in

the phylogeny. We also calculated multiple pairwise RF distances between the trees inferred

from SISRS data mapped to each genome annotation (CDS, introns, intergenic, etc.). Finally, we

calculated the RF distances of each SISRS phylogeny from the topology inferred with all

mapped sites across the turtle genome.

We computed gene and site concordance factors for each branch in the turtle phylogeny

using the trimmed and length-filtered single-copy ortholog alignments, their gene trees, and the

ASTRAL tree with IQ-TREE (Minh et al. 2020a). A branch's gene concordance factor

represents the percentage of decisive gene trees that also contain that branch, and a branch's site

concordance factor is the percentage of decisive alignment sites supporting that branch. We also

computed gene concordance factor and site concordance factor for each branch in the

concatenated single-copy ortholog tree by computing individual locus trees based on the

partitions in IQ-TREE. Locus trees were inferred with default settings, and the species tree was

inferred using an edge-linked proportional partition model with an SH-like approximate

likelihood ratio test and ultrafast bootstrap analysis of 10,000 replicates each (-alrt 10000 -B

10000). The greedy algorithm of PartitionFinder (Lanfear et al. 2012) was applied to find the

best-fit partitioning scheme which was then used in the subsequent tree reconstruction step (-m

TESTMERGE). We also calculated site concordance factors for every branch in the

SVDquartets trees inferred from biallelic parsimony-informative sites within locus types.

Discordance between gene trees and species trees can be caused by introgression,

incomplete lineage sorting, erroneously inferred gene trees stemming from model violations

and/or other artifacts, or combinations of all of these factors, and we aimed to determine how

these issues affect phylogenomic inference in turtles. One expectation of incomplete lineage

sorting is a roughly equal proportion of topologies supporting alternative relationships, while

introgression would cause some minor gene trees to be more frequent than others, reflecting the

direction of gene flow (Huson et al. 2005; Green et al. 2010). We collected the two most

common minor topologies and each node (gDF1 and gDF2 resulting from the concordance factor

analysis in IQ-TREE) and used a chi-squared test to determine if they were similar in terms of

their proportions of all gene trees, where a rejection of the null hypothesis was interpreted as the

presence of introgression in addition to incomplete lineage sorting and potential model

violations. Nodes where we failed to reject the null hypothesis were interpreted as divergence

patterns in turtle evolution driven primarily by either incomplete lineage sorting or model

violations.

Analysis of potential model violations via substitutional saturation, compositional heterogeneity,

and codon usage bias

Phylogenetic signal can be obscured by the accumulation of multiple substitutions at the same

site over time, leading to model violations and increased gene tree-species tree discordance,

particularly at deep evolutionary timescales (Jeffroy et al. 2006; Philippe et al. 2011). Therefore,

to further distinguish between the effects of incomplete lineage sorting and model violations at

nodes in the turtle species tree, we estimated levels of substitutional saturation in the single-copy

ortholog dataset. First, we calculated the slope of a linear regression between the computed raw

pairwise distances and corrected pairwise distances under a TN93 substitution model for 685

single-copy ortholog alignments with complete taxon sampling using the ape 5.5 R package

(Paradis and Schliep 2019). For genes with high levels of saturation, the slope of this regression

will be closer to 0 while less saturated genes will have a slope approaching 1 (Philippe et al.

1994). We then plotted a histogram of all slopes and characterized each alignment as having

slopes above ("unsaturated") and below ("saturated") the mean. To determine the effect of

substitutional saturation on phylogenomic inference in turtles, we compared ASTRAL species

trees inferred from the RAxML trees inferred from unsaturated and saturated genes.

To analyze substitutional saturation at codon positions and assess their effects on

phylogenomic inference in turtles, we realigned the 685 single-copy orthologs with complete

taxon sampling using MACSE (Ranwez et al. 2011), and concatenated alignments that were

partitioned based on (1) first and second codon positions and (2) third codon positions with

AMAS. We then computed the corrected and uncorrected pairwise distances and estimated the

regression slopes for the two codon-based partitions in R as above. We also further separated

these codon partitions according to whether they belonged to the unsaturated or saturated gene

sets, inferred maximum likelihood phylogenies on the four resulting datasets in IQ-TREE with

model testing and 1,000 ultrafast bootstrap replicates, and calculated site concordance factors for

each inferred tree from codon partitions. Finally, to account for substitutional saturation we

computed a maximum likelihood tree from the concatenated and partitioned MACSE amino acid

alignments of the 685 single-copy orthologs using model testing in IQ-TREE. Maximum

likelihood tree topologies, branch support, and concordance factors from codon-partitioned

saturated and unsaturated alignments as well as the amino acids were compared to the ASTRAL

and concatenated supermatrix trees based on single-copy orthologs ≥1,500bp.

A chi-squared test for lineage-specific compositional heterogeneity is computed before

each IQ-TREE run, where the frequency of each nucleotide in a given taxon is compared to the

overall frequency of each nucleotide in the entire dataset. We compared the results of this test

across IQ-TREE analyses of the saturated and unsaturated concatenated partitions based on

codon position. To analyze codon usage bias in our single copy ortholog dataset, we estimated

how often a codon is observed relative to its predicted occurrence in the absence of codon usage

bias, by computing the Relative Synonymous Codon Usage (RSCU) of the concatenated 685

complete-taxon MACSE alignments using CodonW v1.4.4 (Peden 1999) followed by factor

analysis in FactoMineR (Lê et al. 2008) in R.

Divergence time estimation

We estimated divergence times with single-copy ortholog alignments using Bayesian modeling

implemented in BEAST v.2.6.3 (Bouckaert et al. 2019). We subsampled 685 alignments with

complete taxon sampling, from which we further randomly sampled three replicate sets of 10

genes. Based on model testing with ModelFinder, we applied the HKY+Gamma+Invariant site

model to all partitions, and utilized a relaxed lognormal molecular clock and Calibrated Yule tree

prior. To calibrate the time trees, we used 11 calibration priors from the literature (Joyce et al.

2013) which overlapped with our taxon sampling (Supplementary Data). To account for

uncertainty in fossil dating, we used hard minimum constraints and set soft maximum constraints

by placing the maximum ages in the 97.5% quantile of a lognormal prior distribution. Markov

chain Monte Carlo (MCMC) analyses were sampled every 50,000 generations. For each

replicate, we assessed convergence of parameter estimates across the MCMC by monitoring

effective sample sizes with Tracer v.1.7.2 (Rambaut et al. 2018), and we estimated a maximum

clade credibility tree with TreeAnnotator v2.6.3.

We also estimated divergence times in turtle evolution using penalized likelihood with

r8s v1.8.1 (Sanderson 1997, 2002, 2003), using the same fossil calibrations as above for

minimum and maximum node ages; the main difference in this analysis was we fixed the time to

most recent common ancestor (TMRCA) for amniotes at 312 million years ago (Ma) (Donoghue

and Benton 2007) and the TMRCA for Testudines at 220 Ma (Thomson et al. 2021). First, we

concatenated the 685 complete partitions and inferred a maximum likelihood tree with model

testing for each partition as above in IQ-TREE. We then used the inferred phylogeny with

branches in terms of substitutions per site and the 1,836,182 concatenated bases for penalized

likelihood estimation, with cross validation to optimize the smoothing parameter which

quantifies the deviation from the molecular clock.

We used linear regression to determine the relationship between the estimated divergence

time for each node and the concordance factors computed with the inferred trees from each locus

type in the SISRS analysis, the gene trees and the species tree based on single-copy orthologs,

and the locus trees and partitioned maximum likelihood tree based on single-copy orthologs. We

also tested for a correlation between the concordance factors and the estimated rates of molecular

evolution at each branch. We repeated these regressions using estimated divergence times from

Thomson et al. (2020).

Results

A very large number of informative sites for turtle phylogenomics

We analyzed publicly available genome assemblies to extract single copy orthologs from 19

turtles representing 13 out of 14 extant families and 7 of 8 superfamilies, plus 3 outgroup taxa

(Table 1). On average, 4,609 (83%, range 1,911-5,099) conserved tetrapod orthologs were

complete in the genome assemblies, 112 (2.1%, range 7-496) were duplicated, 170 (3.2%, range

55-485) were fragmented, and 419 (7.9%, range 143-2,907) were missing (Fig. 1). Fifty-five

percent of conserved tetrapod orthologs were missing from the genome assembly of the

trionychid *Apalone spinifera*; therefore, we omitted this species from downstream analyses of

single-copy orthologs due to concerns about assembly quality (Waterhouse et al. 2018). Each

single-copy ortholog was present in an average of 19 out of 21 species, averaged 1,902 bp in

length, and contained an average of 53% variable sites and 26% parsimony-informative sites.

Alignment length was correlated with the number of parsimony-informative sites (R^2=0.69,

P=2.2e-16, Fig. 2a). GC content was consistent across the single-copy orthologs and averaged

48% (standard deviation 6.2%, Fig. 2b), as was average pairwise Kimura 2-parameter distance

(mean 0.154, median 01.39, standard deviation 0.072, Fig. 2c). The subset of 2,513 alignments

≥1,500 bp in length had very similar characteristics to the complete set of loci; the average large

locus length was 1,903 bp and the relationship between alignment length and the number of

parsimony-informative sites held (R^2=0.73, P=2.2e-16).

Figure 1. Summary of single-copy orthologs. Percent of tetrapod orthologs from orthoDBv9 present in each

genome assembly as complete single copy, complete duplicated, fragmented, or missing.

Figure 2. Description of single copy ortholog alignments. (a) Scatterplot showing the relationship between

alignment length and the number of parsimony-informative sites for trimmed alignments ≥1,500 bp. (b) Histogram

showing the distribution of GC content in all single copy orthologs after trimming. Solid vertical line represents the

mean value (0.476). (c) Histogram showing the distribution of average pairwise Kimura 2-parameter (K2P)

distances for single copy orthologs after trimming. Solid vertical line represents the mean value (0.154).

We downloaded 38 SRA datasets from 26 turtle species representing 7 of 8 superfamilies

and 14 families (Table 1; Supplementary Data), including the 19 species with complete genome

assemblies from the first dataset. Based on FastQC results, we selected 32 of the SRA datasets

for further analysis ranging from ~1–70X post-trimming coverage. The composite genome

consisted of 5,555,666 contigs with an N50 of 2.4 Mb. We successfully mapped 302,355,876

bases to 2,110,354 contigs in the composite genome which covered 80% of the painted turtle

genome (Table 2), and called 1,655,675 parsimony-informative biallelic SNPs when allowing

missing data for up to two species at a given site. The number of parsimony-informative biallelic

SNPs was much larger in mapped datasets containing only cryptodires (2,787,072 bases) and

testudinoideans (4,696,637 bases) (Supplementary Data), as expected based on Literman et al.

(2021). We collected the highest number of biallelic SNPs from intergenic regions (746,724

bases), followed by introns (533,336 bases) and CDS (136,823 bases). The relatively low

number of parsimony-informative biallelic SNPs in pseudogenic (427) and smRNA regions

(137) is likely due to a small number of annotations for these features in the western painted

turtle genome.

Table 2. Summary of short read data from 26 turtle species mapped to the western painted turtle (*Chrysemys*

***picta*) genome annotations.**

Locus Type	Annotated Bases in the Genome	Bases Mapped by SISRS	Number of biallelic parsimony-informative sites*	Percent of Annotated Bases Mapped
All Mapped	2,365,766,571	302,355,876	1,655,675	12.78%
Intergenic	1,382,145,112	161,445,195	746,724	11.68%
Intronic	812,010,672	115,297,781	533,336	14.20%
lncRNA	64,450,450	9,451,244	52,207	14.66%
Genic Other	65,764,053	9,052,113	36,604	13.76%
CDS	33,486,790	5,718,074	136,823	17.08%
3'-UTR	8,034,514	1,510,783	15,150	18.80%
5'-UTR	5,096,255	708,214	6,519	13.90%
Pseudogenic	670,023	59,058	427	8.81%
smRNA	133,894	20,076	137	14.99%

*When allowing up to two missing species per site, see Supplementary Data for numbers of sites with varying levels

of missing data.

Bioinformatic analysis of turtle genomes yielded well-resolved species relationships

The inferred relationships among turtle lineages based on single-copy orthologs were consistent

with previous studies using phylogenomic markers (Crawford et al. 2015; Shaffer et al. 2017),

with full statistical support for all resulting branches based on posterior probabilities in the

ASTRAL tree, and bootstrap replicates and SH-like approximate likelihood in the supermatrix

tree (Fig. 3, Supplementary Fig. 1, Supplementary Fig. 2). Hereafter we refer to these results

collectively as the turtle species tree. The turtle species tree supports the reciprocal monophyly

of Pleurodira and Cryptodira (Fig. 4). Within cryptodires, there is a split between the

Trionychoidea (soft-shelled turtles) and the Durocryptodira (Crawford et al. 2015), and a further

subdivision of Durocryptodira into the Testudinoidea (pond turtles, big-headed turtles,

geoemyds, and tortoises) and Americhelydia (sea turtles, snapping turtles, mud turtles, and

Dermatemys; Joyce et al. 2013). Within the Testudinoidea, the turtle species tree supports

Testuguria (geoemyds and tortoises; Joyce et al. 2004), and a *Platysternon* + Emydidae clade

("Platyemidae"; Barley et al. 2010; Crawford et al. 2015).

Figure 3. Species trees for turtles based on high-confidence single-copy orthologs. We used 2,513 single-copy

orthologs ≥1,500 bp in length to infer a multilocus coalescent-consistent tree based on gene genealogies (ASTRAL,

left) and a maximum likelihood tree (ML, right) from a concatenated and partitioned supermatrix in IQ-TREE. All

branches received 100% support from local posterior probabilities (ASTRAL), 1,000 bootstrap replicates (IQ-

TREE), and approximate likelihood ratio test (IQ-TREE). Node labels indicate gene and site concordance factors

before and after the slash, respectively. Branch lengths are in terms of coalescent units (ASTRAL) and substitutions

per site (ML). Outgroups (*Gallus, Alligator, Homo*) are not shown.

Figure 4. The phylogenetic relationships of higher turtle taxa. Cladogram depicting the turtle species tree in agreement with results from ASTRAL and concatenated partitioned maximum likelihood analyses as well as most SVDquartets analyses using the mapped data (although these analyses lacked outgroups). Superfamilies or clades are indicated by horizontal shaded text. Additional key clades are labelled with letters in boxes. (a) Durocryptodira, (b) Americhelydia, (c) Testuguria, (d) Emydidae. Additional outgroup *Homo sapiens* not shown.

SVDquartets trees from the genome-wide, all-mapped data as well as from CDS, introns, and intergenic regions were consistent with the species tree based on single-copy orthologs (Fig. 4, Supplementary Fig. 3-11; Supplementary Data). When there was disagreement between trees from mapped regions and the species tree, the difference was often in the placement of *Platysternon* (Fig. 5). For instance, when allowing zero missing gapless parsimony-informative

sites from the all-turtle data, 5' UTR placed *Platysternon* as the outgroup of Geoemydidae +

Testudinidae (435 sites). When allowing up to two missing gapless parsimony-informative sites

from the all-turtle data, pseudogenic regions (426 sites) also placed *Platysternon* as the outgroup

of Geoemydidae + Testudinidae, while smRNA (137 sites) agreed with the species tree and all-

mapped topologies. Meanwhile, at multiple levels of allowed missing data the Testudinoidea-

only mapped sites weakly supported *Platysternon* as the sister taxon to Geoemydidae +

Testudinidae + Emydidae (Supplementary Data).

Figure 5. Alternative topologies in turtle phylogenetics based on markers from different regions of the

genome. (a) *Platysternon* as the sister taxon to Emydidae to the exclusion of Testuguria (Geoemydidae +

Testudinidae), consistent with the turtle species tree (current study), Crawford et al. (2015), and Shaffer et al.

(2017). (b) *Platysternon* as the sister taxon to Testuguria with the exclusion of Emydidae, consistent with 17% of the

single-copy ortholog genes trees, mapped 5'-UTR parsimony-informative sites data with zero missing data, mapped

pseudogene parsimony-informative sites when allowing 2 missing taxa per site, and Krenz et al. (2005). (c)

Platysternon as the sister taxon to Geoemydidae + Testudinidae + Emydidae, consistent with 20% of single-copy

ortholog gene trees and Shaffer et al. (1997).

Heterogeneity is prevalent at well resolved nodes in the turtle phylogeny

While results from our single-copy orthologs and biallelic sites analyses of turtles are consistent

and largely in agreement with other studies based on 100% branch support (Crawford et al. 2015;

Shaffer et al. 2017), some genomic regions, either from single-copy orthologs or from genome

feature annotations, yielded genealogies conflicting with the turtle species tree. For instance,

based on normalized RF distances most gene trees from single-copy orthologs were similar to the

species tree, although there was a long tail to this distribution indicating gene trees with greater

dissimilarity (Fig. 6a). SVDquartets trees using mapped data from CDS, introns, and intergenic

regions were consistent with both the single-copy ortholog species tree and a SVDquartets tree

derived from the all-mapped data. In contrast, mapped pseudogenic DNA, 5' UTR, and smRNA

deviated from the all-mapped data in terms of their RF distances (Supplementary Fig. 12).

439   We also found a high degree of heterogeneity based on gene and site concordance factors

throughout the turtle phylogeny (Fig. 3, Fig. 6, Supplementary Fig. 1, Supplementary Fig. 2). For

instance, while local posterior probabilities were uniformly 100% across branches of the

ASTRAL tree, gene concordance factors ranged from 44 to 97 and site concordance factors

ranged from 30 to 71 (Fig. 6b-d). Gene and site concordance factors based on the ASTRAL tree

were correlated with each other (Figure 6b; R^2=0.55, P=0.0002) and with branch lengths (Fig.

5c-d; R^2=0.88, 3.88e-09 and R^2=0.49, P=0.0007, respectively). We also estimated gene and site

concordance factors based on the locus trees with respect to the maximum likelihood tree

inferred from the concatenated supermatrix, and found a similar range of gene concordance

factors (34 to 92) and site concordance factors (30 to 71) despite 100% bootstrap support, as well

as a strong correlation between the two concordance measures (R^2=0.60, P=0.0001). Gene and

site concordance factors based on the concatenated partitioned maximum likelihood tree were

also correlated with branch lengths (R^2=0.90, 1.02e-09 and R^2=0.58, P=0.0001, respectively).

Figure 6. Phylogenomic heterogeneity in turtles. (a) Histogram of RF distances between gene trees and species

tree. (b-c) Concordance factors calculated using single-copy ortholog alignments, their gene trees, and the ASTRAL

species tree. Results based on locus trees and concatenated supermatrix maximum likelihood tree are in the

supplement. (b) Scatterplot depicting the relationship between gene and site concordance factors at each node. (c)

Scatterplot depicting the relationship between gene concordance factor and branch length. (d) Scatterplot depicting

the relationship between site concordance factor and branch length.

The ancestral branch leading to turtles was assigned a relatively large gene concordance

factor and site concordance factor across analyses based on the ASTRAL, supermatrix, and

SVDquartets trees, and there was high concordance for reciprocally monophyletic Pleurodira and

Cryptodira across genes and sites (Fig. 3, Fig. 4, Supplementary Figs. 3-10). The placement of

Trionychoidea as sister taxon to all other cryptodires was assigned a relatively large gene

concordance factor but small site concordance factor, suggesting that there are a large number of

sites supporting alternative bifurcations. One area with particularly high discordance (i.e.

relatively small gene concordance factor and site concordance factor across all datasets) was the

placement of the big-headed turtle (*Platysternon*) with respect to Emydidae. Our chi-squared test

rejected introgression at this node (Table 3), favoring incomplete lineage sorting or potential

model violations as sources of discordance in the position of *Platysternon*. There was also a high

degree of discordance associated with the americhelydian taxa (Chelonioidea + Chelydridae +

Kinosternoidea), particularly with the ancestral branch leading to Americhelydia as well as with

the sister taxon relationship between *Dermatemys* (Kinosternoidea) and *Chelydra* (Chelydridae).

Based on single-copy orthologs, minor gene tree frequencies were significantly different at these

nodes (Table 4), suggesting that introgression in addition to incomplete lineage sorting and/or

model violations may be driving phylogenomic discordance in these groups.

Table 3. Gene concordance factors for clades represented by nodes in the ASTRAL species tree phylogeny

where the minor gene tree frequencies were similar according to a chi-squared test.

Clade description	Number of possible supporting gene trees	Gene concordance factor	Percent of gene trees supporting first alternative topology	Percent of gene trees supporting second alternative topology
Durocryptodira	2,387	88.73	0.54	0.21
Chelonioidea	1,755	89.46	1.48	0.8
Testudinoidea	2,435	80.66	0.57	0.7
Testuguria	2,255	84.7	0.62	0.8
Testudinidae	1,989	85.22	4.17	5.38
Cuora	1,932	90.94	2.69	2.07
Emydidae + Platysternidae	2,270	46.92	17.05	19.52
Malachlemys + *Chrysemys*	1,730	86.99	5.03	4.39
Pelomedusoidea	2,195	95.81	1.32	1.14
Chelidoidea	2,138	96.49	0.8	1.08

480   **Table 4. Gene concordance factors for clades represented by nodes in the ASTRAL species tree phylogeny**

481   **where the minor gene tree frequencies were statistically different according to a chi-squared test.**

Clade description	Number of possible supporting gene trees	Gene concordance factor	Percent of gene trees supporting first alternative topology	Percent of gene trees supporting second alternative topology
Archosauria	2,144	44.73	10.21	43.38
Testudines	2,254	94.5	0.35	1.24

Cryptodira	2,390	84.56	1.63	5.98
Trionychoidea	1,920	87.6	2.24	3.85
Americhelydia	2,335	43.94	17.69	13.83
Kinosternoidea + Chelydridae	1,970	55.33	14.01	9.95
Emydidae	1,924	83.26	3.12	1.87
Pleurodira	2,396	94.45	0.29	1.34

Patterns of substitutional saturation, compositional heterogeneity, and codon usage bias in turtle

phylogenomics

We measured the slope of the line in a regression between uncorrected and corrected genetic

distances for 685 single-copy ortholog alignments with complete taxon sampling. The slopes

ranged from $y=0.47x$ (most saturated) to $y=0.91x$ (least saturated), but were skewed towards low

levels of saturation, with a mean of $y=0.76x$, a first quartile of $y=0.72x$, and a third quartile of

$y=0.81x$ (Supplementary Fig. 13). When analyzed by codon position, we found that first and

second codon positions were unsaturated ($y=0.86x$) while the slope of the line between corrected

and uncorrected distances at third codon positions were consistent with saturation ($y=0.57x$)

(Supplementary Fig. 14). Saturation did not have an effect on the topology of the species tree, as

ASTRAL trees derived from the most and least saturated sets of genes were identical

(Supplementary Data). In addition, all codon partitions regardless of their saturation levels

produced maximum likelihood trees with intra-Testudines relationships that were identical in

topology to the species tree, and the maximum likelihood tree estimated based on amino acid

partitions was also identical in topology to the species tree (Supplementary Data). There was

some disagreement in the placement of turtles in the amniote phylogeny that appeared to be

driven by substitutional saturation; for instance, the maximum likelihood trees based on first and

second codon positions from saturated genes and all sets of third codon positions supported

turtles as the sister taxon to crocodilians (Supplementary Data; see Discussion).

We found that compositional heterogeneity was most prominent in the saturated third

position and first and second position datasets, as well as in the saturated first and second

position dataset (Supplementary Data). The unsaturated alignment of first and second codon

positions exhibited the least amount of compositional heterogeneity, with five species differing

significantly from the rest of the alignment; however, these species were distributed randomly

across the turtle species tree. Correspondence analyses of the RSCU revealed that some codons,

particularly CCG (Proline), ACG (Threonine), and GCG (Alanine) were favored in different

species (Supplementary Fig. 15). However, there was little evidence for a phylogenetic signal in

this pattern within turtles since codon usage bias was dispersed randomly among species, and the

only outlier along the second dimension was an outgroup taxon (*Gallus*).

Heterogeneity at different timescales in turtle evolution

Results from the replicate BEAST analyses consistently underestimated turtle divergence times

with respect to results from the penalized likelihood analysis as well as those of Thomson et al.

(2020) (Supplementary Data), although are closer to estimates from Chiari et al. (2012) and

Pereira et al. (2017). Across locus types from the SISRS analyses, we found that site

concordance factors were not strongly correlated with estimated node ages (Fig. 7,

Supplementary Fig. 16). For most locus types the relationship between site concordance factor

and divergence time at a given node trended slightly negatively, suggesting a pattern of higher

concordance at younger nodes, although these comparisons were non-significant (Fig. 7b,

$P > 0.1$). The one exception to this trend was for pseudogenes, for which the relationship between

site concordance factors and estimated divergence times trended positive albeit without

significance or with only moderate significance (P=0.072 using Thomson et al. 2020's

divergence time estimates). The relationship between site concordance factors and estimated

DNA substitution rates trended slightly positive for all locus types from the SISRS analysis, but

was non-significant in all comparisons (Supplementary Fig. 17).

We also found a nonsignificant negative trend in the relationship between site

concordance factor and divergence time at a given node based on both the ASTRAL and

supermatrix trees based on single-copy orthologs (Fig. 8a-c; Supplementary Fig. 18. However,

we found a positive (albeit non-significant) trend in the relationship between the estimated

divergence time at a given node and its gene concordance factor when using both trees derived

from single-copy orthologs (Fig. 8e-g; Supplementary Fig. 18). There was no clear relationship

between the site concordance factor of a given node and the estimated substitution rate at the

preceding branch (Fig. 8d), and a moderately significant positive trend (P≤0.05) for the

relationship between the gene concordance factor of a given node and the estimated substitution

rate at the preceding branch (Fig. 8h; Supplementary Fig. 18).

Figure 7. Site concordance and estimated node ages in turtle evolution. (a) Chronogram showing estimated

divergence times in millions of years based on penalized likelihood and fossil calibrations of labelled nodes. 1.

Testudines, 2. Cryptodira, 3. Trionychoidea, 4. Durocryptodira, 5. Geoemydidae, 6. Platyemidae, 7. Emydidae, 8.

Americhelydia, 9. Chelonioidea, 10. Pleurodira, 11. Pelomedusoidea. (b) Site concordance factors (sCF) for each

node in phylogenies inferred from nine locus types and the estimated node age in millions of years ago (Ma) based

on penalized likelihood.

Figure 8. Concordance at genes and sites, estimated node ages, and DNA substitution rates in turtle evolution.

Results shown are from comparisons of single-copy ortholog alignments and their gene trees to the species

tree inferred using ASTRAL. There was a negative trend between site concordance factors (sCF) and estimated

node age (in millions of years or Ma) based on penalized likelihood (PL) from the current study (a), estimated node

ages in Ma from Thomson et al. (2020) (b), estimated node ages in Ma from the BEAST analysis in the current

study (c), and no discernable trend between sCF and estimated DNA substitution rates in terms of substitutions per

site per Ma (d). There was a positive trend between gene concordance factors (gCF) and estimated node ages (e, f, g)

and a moderately significant positive relationship between gene concordance factors and estimated DNA

substitution rates (h). Results from comparisons of the locus trees and the maximum likelihood tree inferred from

the concatenated supermatrix are shown in Supplementary Materials.

Discussion

Heterogeneity has been recognized as an important feature of phylogenomic datasets, and the

exploration of the patterns and sources of gene tree and site discordance has become an essential

step when reconstructing the branching order of life with molecular data (Kumar et al. 2012;

Minh et al. 2020a). To our knowledge, while the role of gene tree-species tree discordance in

estimating the position of turtles within the amniote phylogeny has been the focus of a few

studies (Chiari et al. 2012), there has not been an assessment of phylogenomic heterogeneity and

its effects on the assessment of intra-turtle relationships until now. Here, we have constructed a

robust phylogenomic dataset for turtles based on whole genomes in order to sample the major

turtle lineages. We identified splits in the turtle phylogeny and regions of the turtle genome

associated with gene tree and site heterogeneity, and investigated how potential sources of this

discordance such as introgression, incomplete lineage sorting, and technical issues leading to

model violations may have contributed to conflicting results in phylogenetic estimation of

turtles. The results of our study demonstrate that heterogeneity can be rampant, even in

phylogenomic datasets which have yielded fully resolved species trees.

Protein-coding genes such as the single-copy orthologs used in the current study are

common markers used for phylogenetics, due to their straightforward amplification via PCR or

RNA-Seq, as well as relatively straightforward methods for their orthology assignment and

alignment. However, the use of coding sequences in phylogenetics is complicated by difficulties

with accurate evolutionary modeling due to multiple hits at ancient divergences, a diversity of

substitution rates across the genome, and selective constraint potentially limiting the number of

variant sites. These problems often scale with the size of phylogenomic datasets and can lead to

biased results (Philippe et al. 2011). Therefore, in addition to the single-copy orthologs we also

generated a complementary dataset for turtle phylogenomics consisting of millions of

parsimony-informative biallelic sites using SISRS. Employment of these sites has multiple

advantages: (1) they are easily extracted from multiple sequence alignments; (2) they do not

require complicated evolutionary modeling (i.e. they either support or do not support a splitting

event); and (3) the sheer number of potential sites allows strict quality filtering without loss of

signal (Literman et al. 2021). SISRS resulted in a considerable number of sites collected from

both coding, noncoding, and regulatory parts of the genome, including intergenic regions which

yielded almost 750,000 parsimony-informative sites for turtles. These locus types sample a wide

range of substitution rates, whereas many other phylogenomic methods focus on a much smaller

range of locus types (such as UCEs, AHE, or coding genes). Our results show that many

different regions of the turtle genome yield consistent support for the turtle species tree, leaving a

solid phylogenetic hypothesis for higher turtle systematics. However, inferences from some

regions such as pseudogenes and smRNA were based on a relatively small number of parsimony-

informative sites, highlighting the need for more comprehensive annotations of noncoding

portions of turtle genomes.

The relationships between turtles and other reptiles has been a longstanding debate in

evolutionary biology, due to conflicting results between studies that use fossils and molecules

(Benton et al. 2005; Chiari et al. 2012; Brown and Thomson 2017). Paleontological studies have

placed turtles either at the root of the amniote phylogeny, or as sister taxa with various other

reptilian groups, some of which are long extinct (Gaffney 1980). In the genomic era, analyses of

nucleotides and protein sequences have been mostly in agreement, placing turtles as the sister

taxon to archosaurs (birds and crocodilians) (Shaffer et al. 2013; Wang et al. 2013). Based on

2,513 long single-copy orthologs, our ASTRAL tree results supported a turtles + archosaurs

clade, while the maximum likelihood tree based on the concatenated and partitioned supermatrix

supported a turtles + crocodilians clade that excludes birds. Several lines of evidence suggest that

the latter result is due to poorly fit evolutionary models. First, the maximum likelihood

phylogeny inferred using partitioned amino acids, which accounts for saturation since proteins

are less degenerate than nucleotides, yielded the turtles + archosaurs relationship. Second, we

found that first and second codon positions from non-saturated genes supported turtles +

archosaurs, while first and second codon positions from saturated genes as well as all third codon

positions agreed with turtles + crocodilians (Supplementary Data). Brown and Thomson (2017)

found that duplicated genes in the dataset published by Chiari et al. (2012) dataset were more

likely to reject a turtles + archosaurs clade. Meanwhile, our reliance on single-copy orthologs

from OrthoDB greatly reduces the role of paralogy in the conflict at this node. We suggest that

the correct placement of turtles within the amniote phylogeny using genomic data has been

hampered by substitutional saturation at such ancient divergences, and that when accounting for

these artifacts the data support a turtles + archosaurs clade (Archelosauria; see Crawford et al.

2015).

When gene tree-species tree discordance is driven largely by technical errors such as

saturation, it should be most apparent at older nodes in phylogeny, which are more affected by

homoplasy, poor alignments, or model misspecifications. However, with the exception of the

turtle-archosaur split, across both single-copy ortholog and SISRS datasets we found no

significant correlation between concordance factors and estimated node ages. A slightly negative

relationship between site concordance factors calculated from the single-copy ortholog dataset

and node age suggests that these deeper splits in turtle evolution might be affected by multiple

substitutions; however, this pattern is obscured a slightly positive relationship between gene

concordance factors and node age using the same data. In addition, inferred turtle relationships

were identical when analyzing different sets of genes separated by levels of substitutional

saturation and partitioned by codon position. Thus, while it may be possible that multiple

substitutions or other model violations are driving discordance at ancient nodes in the turtle

phylogeny, their effects are either extremely weak or nonexistent within and among most turtle

lineages, and the observed patterns may have more to do with errors in rate estimation using

molecular data and fossil calibrations.

Our results suggest that population-level processes such as introgression and/or

incomplete lineage sorting have played a key role at multiple stages of turtle diversification,

leading to confusion in the literature about the placement of at least some taxa. In particular, the

phylogenetic position of *Platysternon* has been highly contentious (Joyce et al. 2004, 2021), with

early molecular studies in disagreement (Shaffer et al. 1997; Krenz et al. 2005; Parham et al.

2006; Barley et al. 2010). Our turtle species tree results mirror Crawford et al. (2015) and

Shaffer et al. (2017), fully supporting *Platysternon* as the sister taxon to Emydidae. However, we

found that only 47% of single-copy ortholog gene trees are in agreement with a *Platysternon* +

Emydidae clade, and 37% of the remaining gene trees support only two alternative relationships:

of these, 17% of gene trees support *Platysternon* as the sister taxon to Testuguria (Geoemydidae

+ Testudinidae) following Krenz et al. (2005), and 20% support *Platysternon* as the sister taxon

to Testudinoidea (Emydidae + Geoemydidae + Testudinidae) following Shaffer et al. (1997).

The proportion of gene trees supporting these alternative topologies was similar according to our

chi-squared test (Table 3), and when we analyzed gene trees according to their levels of

substitutional saturation, we saw no effect on the placement of *Platysternon*. As the most

frequent alternate gene trees mirror published alternative hypotheses concerning the relationship

of *Platysternon* to other testudinoideans, we suggest that incomplete lineage sorting is a

previously unrecognized driver of conflicting phylogenetic results in the higher classification of

these groups.

Another subject of conflict across previous phylogenetic studies of turtles is the

relationship between Chelonioidea, Kinosternoidea, and the Chelydridae, all of which are now

considered to be in the monophyletic Americhelydia (Barley et al. 2010; Joyce et al. 2013;

Crawford et al. 2015). We found consistent support for Americhelydia across datasets and

methods, as well as a sister taxon relationship between Chelydridae and Kinosternoidea, yet with

considerable gene tree-species tree discordance with single-copy orthologs (Fig. 3) and site

discordance with SISRS data (Supplementary Fig. 3-11). For instance, only 44% of decisive

gene trees from single-copy orthologs contained an Americhelydia clade, and only 55% of

decisive gene trees supported Chelydridae + Kinosternoidea. As with *Platysternon*, comparison

of trees when accounting for saturation or codon position did not affect the placement of

americhelydians. The frequencies of minor gene trees disagreeing with these splits were

statistically different (Table 4), suggesting both introgression and incomplete lineage sorting may

have played a role in the diversification of these taxa. To our knowledge, the role of

hybridization during the speciation events at the deeper splits of the turtle phylogeny has not

been investigated, although contemporary hybridization between sea turtle genera with up to ~65

Ma of divergence is well documented (Karl et al. 1995; Arantes et al. 2020). As these

occurrences are associated with low reproductive success (Arantes et al. 2020), it will be

important to determine if episodic introgression is a feature of the long-term evolution of turtles.

Some recent molecular analyses of turtle relationships are limited in terms of the size of

the molecular dataset, yet have more complete taxonomic sampling than our study, including up

to 85% of extant species (Pereira et al. 2017; Colston et al. 2020; Thomson et al. 2021). A denser

sampling of lineages maximizes the molecular signal for testing hypotheses about past

diversification events as well as the relationship between evolutionary distinctiveness and

extinction risk. In contrast, our study is limited in terms of taxonomic sampling due to our

reliance on complete genomic sequences and assemblies for turtles, which are lagging compared

to other amniote groups. As our results demonstrate widespread discordance in turtle

phylogenomic datasets, additional lineages should be targeted for genomic sequencing in order

to improve our ability to accurately estimate the evolutionary distinctiveness of certain species.

These studies will be crucial in setting future conservation priorities for turtles.

Acknowledgements

This research was funded by the National Science Foundation (#1812291 to RL) and by startup

funds provided to MT by the School of Informatics, Computing, and Cyber Systems at Northern

Arizona University. We would like to acknowledge the Monsoon computing cluster at Northern

Arizona University (https://nau.edu/high-performance-computing/) for providing the

computational resources necessary to carry out this study.

Data Availability

The bioinformatic workflows used to extract and analyze single-copy orthologs from genome

assemblies are available at https://github.com/mibyars/busco_phylo_pipeline. Data generated for

this study including alignments and trees are available at the Harvard Dataverse

(https://doi.org/10.7910/DVN/WYQRPC).

Literature Cited

Arantes L.S., Ferreira L.C.L., Driller M., Repinaldo Filho F.P.M., Mazzoni C.J. and Santos F.R.

2020. Genomic evidence of recent hybridization between sea turtles at Abrolhos

699   Archipelago and its association to low reproductive output. Sci. Rep. 10:12847.

700 Barley A.J., Spinks P.Q., Thomson R.C., Shaffer H.B. 2010. Fourteen nuclear genes provide

701   phylogenetic resolution for difficult nodes in the turtle tree of life. Mol. Phylogenet. Evol.

702   55:1189–1194.

703 Benton M. 2005. Vertebrate Paleontology. Oxford, UK: Blackwell Science Ltd.

704 Boisvert S., Laviolette F., Corbeil J. 2010. Ray: Simultaneous Assembly of Reads from a Mix of

705   High-Throughput Sequencing Technologies. J. Comput. Biol. 17:1519–1533.

706 Borowiec M.L. 2016. AMAS: a fast tool for alignment manipulation and computing of summary

707   statistics. PeerJ. 4:e1660.

708 Bouckaert R., Vaughan T.G., Barido-Sottani J., Duchêne S., Fourment M., Gavryushkina A.,

709   Heled J., Jones G., Kühnert D., Maio N.D., Matschiner M., Mendes F.K., Müller N.F.,

710   Ogilvie H.A., Plessis L. du, Popinga A., Rambaut A., Rasmussen D., Siveroni I., Suchard

711   M.A., Wu C.-H., Xie D., Zhang C., Stadler T., Drummond A.J. 2019. BEAST 2.5: An

712   advanced software platform for Bayesian evolutionary analysis. PLOS Comput. Biol.

713   15:e1006650.

714 Brown J.M., Thomson R.C. 2017. Bayes Factors unmask highly variable information content,

715   bias and extreme influence in phylogenomic analyses. Syst. Biol. 66:517–530.

716 Burbrink F.T., Grazziotin F.G., Pyron R.A., Cundall D., Donnellan S., Irish F., Keogh J.S.,

717   Kraus F., Murphy R.W., Noonan B., Raxworthy C.J., Ruane S., Lemmon A.R., Lemmon

718   E.M., Zaher H. 2020. Interrogating Genomic-Scale Data for Squamata (Lizards, Snakes,

719   and Amphisbaenians) Shows no Support for Key Traditional Morphological

720   Relationships. Syst. Biol. 69:502-520.

721 Chiari Y., Cahais V., Galtier N., Delsuc F. 2012. Phylogenomic analyses support the position of

turtles as the sister group of birds and crocodiles (Archosauria). BMC Biol. 65 (2012).

Chifman J., Kubatko L. 2014. Quartet Inference from SNP Data Under the Coalescent Model.

Bioinformatics. 30:3317–3324.

Cooper E.D. 2014. Overly simplistic substitution models obscure green plant phylogeny. Trends

Plant Sci. 19:576–582.

Colston T.J., Kulkarni P., Jetz W., Pyron R.A. 2020. Phylogenetic and spatial distribution of

evolutionary diversification, isolation, and threat in turtles and crocodilians (non-avian

archosauromorphs). BMC Evol. Biol. 20:81.

Cox C.J., Li B., Foster P.G., Embley T.M., Civáň P. 2014. Conflicting Phylogenies for Early

Land Plants are Caused by Composition Biases among Synonymous Substitutions. Syst.

Biol. 63:272–279.

Crawford N.G., Faircloth B.C., McCormack J.E., Brumfield R.T., Winker K., Glenn T.C. 2012.

More than 1000 ultraconserved elements provide evidence that turtles are the sister group

of archosaurs. Biol. Lett. 8:783–786.

Crawford N.G., Parham J.F., Sellas A.B., Faircloth B.C., Glenn T.C., Papenfuss T.J., Henderson

J.B., Hansen M.H., Simison W.B. 2015. A phylogenomic analysis of turtles. Mol.

Phylogenet. Evol. 83:250–257.

Donoghue P.C.J., Benton M.J. 2007. Rocks and clocks: calibrating the Tree of Life using fossils

and molecules. Trends Ecol. Evol. 22:424–431.

Dudchenko O., Batra S.S., Omer A.D., Nyquist S.K., Hoeger M., Durand N.C., Shamim M.S.,

Machol I., Lander E.S., Aiden A.P., Aiden E.L. 2017. De novo assembly of the *Aedes*

aegypti genome using Hi-C yields chromosome-length scaffolds. Science. 356:92–95.

Edwards S.V., Xi Z., Janke A., Faircloth B.C., McCormack J.E., Glenn T.C., Zhong B., Wu S.,

Lemmon E.M., Lemmon A.R., Leaché A.D., Liu L., Davis C.C. 2016. Implementing and

testing the multispecies coalescent model: A valuable paradigm for phylogenomics. Mol.

Phylogenet. Evol. 94:447–462.

Faircloth B.C., Sorenson L., Santini F., Alfaro M.E. 2013. A Phylogenomic Perspective on the

Radiation of Ray-Finned Fishes Based upon Targeted Sequencing of Ultraconserved

Elements (UCEs). PLOS ONE. 8:e65923.

Foster P.G. 2004. Modeling Compositional Heterogeneity. Syst. Biol. 53:485–495.

Galtier N., Daubin V. 2008. Dealing with incongruence in phylogenomic analyses. Philos. Trans.

R. Soc. B Biol. Sci. 363:4023–4029.

Garland T., Bennett A.F., Rezende E.L. 2005. Phylogenetic approaches in comparative

physiology. J. Exp. Biol. 208:3015-3035.

Gaffney E. 1980. Phylogenetic relationships of the major groups of amniotes. In: Panchen A.

editor. The terrestrial environment and the origin of land vertebrates. London: Academic

Press. P. 593-610.

Green R.E., Krause J., Briggs A.W., Maricic T., Stenzel U., Kircher M., Patterson N., Li H., Zhai

W., Fritz M.H.-Y., Hansen N.F., Durand E.Y., Malaspinas A.-S., Jensen J.D., Marques-

Bonet T., Alkan C., Prüfer K., Meyer M., Burbano H.A., Good J.M., Schultz R., Aximu-

Petri A., Butthof A., Höber B., Höffner B., Siegemund M., Weihmann A., Nusbaum C.,

Lander E.S., Russ C., Novod N., Affourtit J., Egholm M., Verna C., Rudan P., Brajkovic

D., Kucan Ž., Gušic I., Doronichev V.B., Golovanova L.V., Lalueza-Fox C., Rasilla M.

de la, Fortea J., Rosas A., Schmitz R.W., Johnson P.L.F., Eichler E.E., Falush D., Birney

E., Mullikin J.C., Slatkin M., Nielsen R., Kelso J., Lachmann M., Reich D., Pääbo S.

2010. A Draft Sequence of the Neandertal Genome. Science. 328:710–722.

Hime P.M., Lemmon A.R., Lemmon E.C.M., Prendini E., Brown J.M., Thomson R.C., Kratovil

J.D., Noonan B.P., Pyron R.A., Peloso P.L.V., Kortyna M.L., Keogh J.S., Donnellan

S.C., Mueller R.L., Raxworthy C.J., Kunte K., Ron S.R., Das S., Gaitonde N., Green

D.M., Labisko J., Che J., Weisrock D.W. 2021. Phylogenomics Reveals Ancient Gene

Tree Discordance in the Amphibian Tree of Life. Syst. Biol. 70:48-66.

Huson D.H., Klöpper T., Lockhart P.J., Steel M.A. 2005. Reconstruction of reticulate networks

from gene trees. Springer, Berlin, Heidelberg. P. 233-249.

Irisarri I., Baurain D., Brinkmann H., Delsuc F., Sire J.-Y., Kupfer A., Petersen J., Jarek M.,

Meyer A., Vences M., Philippe H. 2017. Phylotranscriptomic consolidation of the jawed

vertebrate timetree. Nat. Ecol. Evol. 1:1370–1378.

IUCN 2021. The IUCN Red List of Threatened Species. Version 2021-2.

https://www.iucnredlist.org. Downloaded on November 1, 2021.

Jarvis E.D., Mirarab S., Aberer A.J., Li B., Houde P., Li C., Ho S.Y.W., Faircloth B.C., Nabholz

B., Howard J.T., Suh A., Weber C.C., da Fonseca R.R., Li J., Zhang F., Li H., Zhou L.,

Narula N., Liu L., Ganapathy G., Boussau B., Bayzid M.S., Zavidovych V., Subramanian

S., Gabaldón T., Capella-Gutiérrez S., Huerta-Cepas J., Rekepalli B., Munch K.,

Schierup M., Lindow B., Warren W.C., Ray D., Green R.E., Bruford M.W., Zhan X.,

Dixon A., Li S., Li N., Huang Y., Derryberry E.P., Bertelsen M.F., Sheldon F.H.,

Brumfield R.T., Mello C.V., Lovell P.V., Wirthlin M., Schneider M.P.C., Prosdocimi F.,

Samaniego J.A., Velazquez A.M.V., Alfaro-Núñez A., Campos P.F., Petersen B.,

Sicheritz-Ponten T., Pas A., Bailey T., Scofield P., Bunce M., Lambert D.M., Zhou Q.,

Perelman P., Driskell A.C., Shapiro B., Xiong Z., Zeng Y., Liu S., Li Z., Liu B., Wu K.,

Xiao J., Yinqi X., Zheng Q., Zhang Y., Yang H., Wang J., Smeds L., Rheindt F.E., Braun

M., Fjeldså J., Orlando L., Barker F.K., Jønsson K.A., Johnson W., Koepfli K.-P.,

O'Brien S., Haussler D., Ryder O.A., Rahbek C., Willerslev E., Graves G.R., Glenn T.C.,

McCormack J., Burt D., Ellegren H., Alström P., Edwards S.V., Stamatakis A., Mindell

D.P., Cracraft J., Braun E.L., Warnow T., Jun W., Gilbert M.T.P., Zhang G. 2014.

Whole-genome analyses resolve early branches in the tree of life of modern birds.

Science. 346:1320–1331.

Jeffroy O., Brinkmann H., Delsuc F., Philippe H. 2006. Phylogenomics: the beginning of

incongruence? 22:225-231.

Joyce W.G. 2007. Phylogenetic Relationships of Mesozoic Turtles. Bull. Peabody Mus. Nat.

Hist. 48:3–102.

Joyce W.G., Anquetin J., Cadena E.-A., Claude J., Danilov I.G., Evers S.W., Ferreira G.S.,

Gentry A.D., Georgalis G.L., Lyson T.R., Pérez-García A., Rabi M., Sterli J., Vitek N.S.,

Parham J.F. 2021. A nomenclature for fossil and living turtles using phylogenetically

defined clade names. Swiss J. Palaeontol. 140:5.

Joyce W.G., Parham J.F., Gauthier J.A. 2004. Developing a protocol for the conversion of rank-

based taxon names to phylogenetically defined clade names, as exemplified by turtles. J.

Paleontol. 78:989–1013.

Joyce W.G., Parham J.F., Lyson T.R., Warnock R.C.M., Donoghue P.C.J. 2013. A divergence

dating analysis of turtles using fossil calibrations: an example of best practices. J.

Paleontol. 87:612–634.

Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., von Haeseler A., Jermiin L.S. 2017.

ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods.

14:587–589.

Katoh K., Standley D.M. 2013. MAFFT Multiple Sequence Alignment Software Version 7:

815   Improvements in Performance and Usability. Mol. Biol. Evol. 30:772–780.

816   Karl S.A., Bowen B.W., Avise J.C. 1995. Hybridization among the ancient mariners:

817   Characterization of marine turtle hybrids with molecular genetic assays. J. Hered.

818   86:262–268.

819   Krenz J.G., Naylor G.J.P., Shaffer H.B., Janzen F.J. 2005. Molecular phylogenetics and

820   evolution of turtles. Mol. Phylogenet. Evol. 37:178–191.

821   Kubatko L.S., Degnan J.H. 2007. Inconsistency of Phylogenetic Estimates from Concatenated

822   Data under Coalescence. Syst. Biol. 56:17–24.

823   Kumar S., Filipski A.J., Battistuzzi F.U., Kosakovsky Pond S.L., Tamura K. 2012. Statistics and

824   Truth in Phylogenomics. Mol. Biol. Evol. 29:457–472.

825   Kumar S., Stecher G., Li M., Knyaz C., Tamura K. 2018. MEGA X: Molecular Evolutionary

826   Genetics Analysis across Computing Platforms. Mol. Biol. Evol. 35:1547–1549.

827   Lanfear R., Calcott B., Ho S.Y.W., Guindon S. 2012. Partitionfinder: combined selection of

828   partitioning schemes and substitution models for phylogenetic analyses. Mol. Biol. Evol.

829   29:1695–1701.

830   Langmead B., Salzberg S.L. 2012. Fast gapped-read alignment with Bowtie 2. Nat. Methods.

831   9:357–359.

832   Lê S., Josse J., Husson F. 2008. FactoMineR: An R Package for Multivariate Analysis. J. Stat.

833   Softw. 25: 1–18.

834   Lemmon A.R., Emme S.A., Lemmon E.M. 2012. Anchored Hybrid Enrichment for Massively

835   High-Throughput Phylogenomics. Syst. Biol. 61:727–744.

Literman R., Schwartz R. 2021. Genome-Scale Profiling Reveals Noncoding Loci Carry Higher

Proportions of Concordant Data. Mol. Biol. Evol.

Liu L., Zhang J., Rheindt F.E., Lei F., Qu Y., Wang Y., Zhang Y., Sullivan C., Nie W., Wang J.,

Yang F., Chen J., Edwards S.V., Meng J., Wu S. 2017. Genomic evidence reveals a

radiation of placental mammals uninterrupted by the KPg boundary. Proc. Nat. Acad. Sci.

E7282-E7290.

Lopes F., Oliveira L.R., Kessler A., Beux Y., Crespo E., Cárdenas-Alayza S., Majluf P.,

Sepúlveda M., Brownell R.L. Jr., Franco-Trecu V., Páez-Rosas D., Chaves J., Loch C.,

Robertson B.C., Acevedo-Whitehouse K., Elorriaga-Verplancken F.R., Kirkman S.P.,

Peart C.R., Wolf J.B.W., Bonatto S.L. 2021. Phylogenomic Discordance in the Eared

Seals is best explained by Incomplete Lineage Sorting following Explosive Radiation in

the Southern Hemisphere. Syst. Biol. 70:786–802.

Lynch M. 2007. The Origins of Genome Architecture. Sinauer Associates Sunderland, MA.

Minh B.Q., Hahn M.W., Lanfear R. 2020a. New Methods to Calculate Concordance Factors for

Phylogenomic Datasets. Mol. Biol. Evol. 37:2727–2733.

Minh B.Q., Schmidt H.A., Chernomor O., Schrempf D., Woodhams M.D., von Haeseler A.,

Lanfear R. 2020b. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. Mol. Biol. Evol. 37:1530–1534.

Mölder F., Jablonski K.P., Letcher B., Hall M.B., Tomkins-Tinch C.H., Sochat V., Forster J.,

Lee S., Twardziok S.O., Kanitz A., Wilm A., Holtgrewe M., Rahmann S., Nahnsen S.,

Köster J. 2021. Sustainable data analysis with Snakemake.

https://f1000research.com/articles/10-33.

Morales-Briones D.F., Kadereit G., Tefarikis D.T., Moore M.J., Smith S.A., Brockington S.F.,

Timoneda A., Yim W.C., Cushman J.C., Yang Y. 2021. Disentangling Sources of Gene

Tree Discordance in Phylogenomic Data Sets: Testing Ancient Hybridizations in

Amaranthaceae s.l. Syst. Biol. 70:219-235.

Pamilo P., Nei M. 1988. Relationships between gene trees and species trees. Mol. Biol. Evol.

5:568–583.

Parham J.F., Feldman C.R., Boore J.L. 2006. The complete mitochondrial genome of the

enigmatic bigheaded turtle (Platysternon): description of unusual genomic features and

the reconciliation of phylogenetic hypotheses based on mitochondrial and nuclear DNA.

BMC Evol. Biol. 6:11.

Paradis E., Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary

analyses in R. Bioinformatics. 35:526-528.

Peden J. 1999. Analysis of Codon Usage. PhD diss., University of Nottingham.

Pereira A.G., Sterli J., Moreira F.R.R., Schrago C.G. 2017. Multilocus phylogeny and statistical

biogeography clarify the evolutionary history or major lineages of turtles. Syst. Biol.

113:59-66.

Philippe H., Brinkmann H., Lavrov D.V., Littlewood D.T.J., Manuel M., Wörheide G., Baurain

D. 2011. Resolving difficult phylogenetic questions: why more sequences are not enough.

PLOS Biology. 9:e1000602.

Philippe H., Sörhannus U., Baroin A., Perasso R., Gasse F., Adoutte A. 1994. Comparison of

molecular and paleontological data in diatoms suggests a major gap in the fossil record. J.

Evol. Biol. 7:247-265.

Prum R.O., Berv J.S., Dornburg A., Field D.J., Townsend J.P., Lemmon E.M., Lemmon A.R.

2015. A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA

sequencing. Nature. 526:569–573.

Quesada V., Freitas-Rodríguez S., Miller J., Pérez-Silva J.G., Jiang Z.-F., Tapia W., Santiago-

Fernández O., Campos-Iglesias D., Kuderna L.F.K., Quinzin M., Álvarez M.G., Carrero

D., Beheregaray L.B., Gibbs J.P., Chiari Y., Glaberman S., Ciofi C., Araujo-Voces M.,

Mayoral P., Arango J.R., Tamargo-Gómez I., Roiz-Valle D., Pascual-Torner M., Evans

B.R., Edwards D.L., Garrick R.C., Russello M.A., Poulakakis N., Gaughran S.J., Rueda

D.O., Bretones G., Marquès-Bonet T., White K.P., Caccone A., López-Otín C. 2019.

Giant tortoise genomes provide insights into longevity and age-related disease. Nat. Ecol.

Evol. 3:87–95.

Quinlan A.R., Hall I.M. 2010. BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics. 26:841–842.

Rambaut A., Drummond A.J., Xie D., Baele G., Suchard M.A. 2018. Posterior summarization in

Bayesian phylogenetics using Tracer 1.7. Syst. Biol. 67:901–904.

Ranwez V., Harispe S., Delsuc F., Douzery E.J.P. 2011. MACSE: Multiple Alignment of Coding

SEquences accounting for frameshifts and stop codons. PLOS ONE. 6:e22594.

Robinson D.F., Foulds L.R. 1981. Comparison of phylogenetic trees. Math. Biosci. 53:131–147.

Rokas A., Williams B.L., King N., Carroll S.B. 2003. Genome-scale approaches to resolving

incongruence in molecular phylogenies. Nature. 425:798–804.

Sanderson M.J. 1997. A nonparametric approach to estimating divergence times in the absence

of rate constancy. Mol. Biol. Evol.

Sanderson M.J. 2002. Estimating absolute rates of molecular evolution and divergence times: a

penalized likelihood approach. Mol. Biol. Evol. 19:101–109.

Sanderson M.J. 2003. r8s: inferring absolute rates of molecular evolution and divergence times

in the absence of a molecular clock. Bioinformatics. 19:301–302.

Sayyari E., Mirarab S. 2016. Fast Coalescent-Based Computation of Local Branch Support from

907   Quartet Frequencies. Mol. Biol. Evol. 33:1654–1668.

908   Schwartz R.S., Harkins K.M., Stone A.C., Cartwright R.A. 2015. A composite genome approach

909   to identify phylogenetically informative data from next-generation sequencing. BMC

910   Bioinformatics. 16:193.

911   Shaffer H.B., McCartney-Melstad E., Near T.J., Mount G.G., Spinks P.Q. 2017. Phylogenomic

912   analyses of 539 highly informative loci dates a fully resolved time tree for the major

913   clades of living turtles (Testudines). Mol. Phylogenet. Evol. 115:7–15.

914   Shaffer H.B., Meylan P., McKnight M.L. 1997. Tests of turtle phylogeny: molecular,

915   morphological, and paleontological approaches. Syst. Biol. 46:235–268.

916   Shaffer H.B., Minx P., Warren D.E., Shedlock A.M., Thomson R.C., Valenzuela N., Abramyan

917   J., Amemiya C.T., Badenhorst D., Biggar K.K., Borchert G.M., Botka C.W., Bowden

918   R.M., Braun E.L., Bronikowski A.M., Bruneau B.G., Buck L.T., Capel B., Castoe T.A.,

919   Czerwinski M., Delehaunty K.D., Edwards S.V., Fronick C.C., Fujita M.K., Fulton L.,

920   Graves T.A., Green R.E., Haerty W., Hariharan R., Hernandez O., Hillier L.W.,

921   Holloway A.K., Janes D., Janzen F.J., Kandoth C., Kong L., de Koning A.P.J., Li Y.,

922   Literman R., McGaugh S.E., Mork L., O'Laughlin M., Paitz R.T., Pollock D.D., Ponting

923   C.P., Radhakrishnan S., Raney B.J., Richman J.M., St John J., Schwartz T., Sethuraman

924   A., Spinks P.Q., Storey K.B., Thane N., Vinař T., Zimmerman L.M., Warren W.C.,

925   Mardis E.R., Wilson R.K. 2013. The western painted turtle genome, a model for the

926   evolution of extreme physiological adaptations in a slowly evolving lineage. Genome

927   Biol. 14:R28.

Shimodaira H. 2002. An Approximately Unbiased Test of Phylogenetic Tree Selection. Syst.

Biol. 51:492–508.

Singhal S., Colston T.J., Grundler M.R., Smith S.A., Costa G.C., Colli G.R., Moritz C., Pyron

R.A., Rabosky D.L. 2021. Congruence and Conflict in the Higher-Level Phylogenetics of

Squamate Reptiles: An Expanded Phylogenomic Perspective. Syst. Biol. 70:542–557.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. Bioinformatics. 30:1312–1313.

Steenwyk J.L., Buida T.J. III, Labella A.L., Li Y., Shen X.-X., Rokas A. 2021. PhyKIT: a

broadly applicable UNIX shell toolkit for processing and analyzing phylogenomic data.

Bioinformatics. btab096.

Swofford D.L. 2003. PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods).

Sinauer associates Sunderland, MA.

Tarver J.E., dos Reis M., Mirarab S., Moran R.J., Parker S., O'Reilly J.E., King B.L., O'Connell

M.J., Asher R.J., Warnow T., Peterson K.J., Donoghue P.C.J., Pisani D. 2016. The

interrelationships of placental mammals and the limits of phylogenetic inference.

Genome Biol. Evol. 8:330-344.

Thomson R.C., Spinks P.Q., Shaffer H.B. 2021. A global phylogeny of turtles reveals a burst of

climate-associated diversification on continental margins. Proc. Natl. Acad. Sci. 118.

Tollis M., Denardo D.F., Cornelius J.A., Dolby G.A., Edwards T., Henen B.T., Karl A.E.,

Murphy R.W., Kusumi K. 2017. The Agassiz's desert tortoise genome provides a

resource for the conservation of a threatened species. PloS One. 12:e0177708.

Townsend J.P., López-Giráldez F, Friedman R. 2008. The phylogenetic informativeness of

nucleotide and amino acid sequences for reconstructing the vertebrate tree. J. Mol. Evol.

67:437-447.

Uetz P., Freed P., Aguilar R., Hošek J. 2021. The Reptile Database. (last accessed August 2021)

Wang Z., Pascual-Anaya J., Zadissa A., Li W., Niimura Y., Wang J., Huang Z., Li C., White S.,

Xiong Z., Fang D., Wang B., Ming Y., Chen Y., Zheng Y., Kuraku S., Pignatelli M.,

Herrero J., Beal K., Nozawa M., Li Q., Wang J., Zhang H., Yu L., Shigenobu S., Wang

J., Liu J., Flicek P., Searle S., Wang J., Kuratani S., Yin Y., Aken B., Zhang G., Irie N.

2013. The draft genomes of soft-shell turtle and green sea turtle yield insights into the

development and evolution of the turtle-specific body plan. Nat. Genet. 45:701–706.

Waterhouse R.M., Seppey M., Simão F.A., Manni M., Ioannidis P., Klioutchnikov G.,

Kriventseva E.V., Zdobnov E.M. 2018. BUSCO Applications from Quality Assessments

to Gene Prediction and Phylogenomics. Mol. Biol. Evol. 35:543–548.

Wolf Y.I., Rogozin I.B., Grishin N.V., Koonin E.V. 2002. Genome trees and the tree of life.

Trends Genet. TIG. 18:472–479.

Zdobnov E.M., Tegenfeldt F., Kuznetsov D., Waterhouse R.M., Simão F.A., Ioannidis P.,

Seppey M., Loetscher A., Kriventseva E.V. 2017. OrthoDB v9.1: cataloging evolutionary

and functional annotations for animal, fungal, plant, archaeal, bacterial and viral

orthologs. Nucleic Acids Res. 45:D744–D749.

Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree

reconstruction from partially resolved gene trees. BMC Bioinformatics. 19:153.