A Genomic Perspective on the Evolutionary Diversification of Turtles

Simone M. Gable[1*], Michael I. Byars[1*], Robert Literman[2], Marc Tollis[1]

[1]School of Informatics, Computing, and Cyber Systems, Northern Arizona University, PO Box

5693, Flagstaff, AZ 86011

[2]Department of Biological Sciences, University of Rhode Island, 120 Flagg Road, Kingstown, RI,

02881

*These authors contributed equally.

Corresponding author: Marc Tollis, School of Informatics, Computing, and Cyber Systems,

Northern Arizona University, PO Box 5693, Flagstaff, AZ, 86011; marc.tollis@nau.edu

Running header: Phylogenomic Heterogeneity in Turtles

ABSTRACT

To examine phylogenetic heterogeneity in turtle evolution, we collected thousands of high-

confidence single-copy orthologs from 19 genome assemblies representative of extant turtle

diversity and estimated a phylogeny with multispecies coalescent and concatenated partitioned

methods. We also collected next-generation sequences from 26 turtle species and assembled

millions of biallelic markers to reconstruct phylogenies from annotated regions (coding regions,

introns, untranslated regions, intergenic, and others) of the western painted turtle (*Chrysemys*

picta bellii) genome. We then measured gene tree-species tree discordance, as well as gene and

site heterogeneity at each node in the inferred trees, and tested for incomplete lineage sorting and

temporal patterns in phylogenomic heterogeneity across turtle evolution. We found 100%

support for all bifurcations in the inferred turtle species phylogenies. However, a number of

genes, sites, and genomic features supported alternate relationships between turtle taxa, and

some nodes in the turtle phylogeny were well-explained by incomplete lineage sorting. There

was no clear pattern between site concordance, node age, and DNA substitution rate across most

annotated genomic regions, suggesting a relatively uniform proportion of informative sites drive

phylogenetic inference across the evolution of turtles. We found more gene concordance at older

nodes in the turtle phylogeny, and suggest that, in addition to incomplete lineage sorting, an

overall lack of gene informativeness stemming from a slow rate of evolution can confound

inferred patterns in turtle phylogenomics, particularly at more recent divergences. Our study

demonstrates that heterogeneity is to be expected even in well resolved clades such as turtles,

and that future phylogenomic studies should aim to sample as much of the genome as possible.

Keywords: turtles, genomes, phylogeny, discordance

Introduction

Next-generation sequencing has revolutionized the understanding of evolutionary relationships

by greatly increasing the numbers of sampled loci (Lemmon et al. 2012; Faircloth et al. 2013;

Edwards et al. 2016), shedding new light on the branching order of diversification across the

history of life. However, phylogenetic studies based on genome-scale data often contain a great

deal of heterogeneity in the inferred patterns (Hime et al. 2021; Lopes et al. 2021; Morales-

Briones et al. 2021; Singhal et al. 2021), particularly in the form of discordance among sites and

gene trees (Kumar et al. 2012). Gene tree-species tree discordance occurs when gene trees are in

conflict with the underlying species relationships, while site discordance occurs when a subset of

sites support different bifurcations in a species tree, and both factors can lead to biased

phylogenetic estimates. Biological factors driving this heterogeneity include: differing

evolutionary rates across the genome, incomplete lineage sorting, gene duplication or loss, and

hidden paralogy (Pamilo and Nei 1988; Galtier and Daubin 2008). Artifactual factors driving

heterogeneity include sequencing errors, as well as substitution model violations such as

saturation or multiple hits (Foster 2004; Cooper 2014; Cox et al. 2014). Accounting for gene and

site discordance often requires the application of multiple tools in phylogenetic analysis,

including partitioning schemes which try to account for differing evolutionary rates and model

parameters (Kubatko and Degnan 2007), and multispecies coalescent methods accounting for

population-level processes such as incomplete lineage sorting (Edwards et al. 2016). The

application of hundreds or even thousands of loci is often an attempt to overcome errors

stemming from or driving phylogenomic conflicts, with the goal of reconstructing better-

resolved phylogenies by reducing the overall sampling variance (Minh et al. 2020a).

63   An emerging advantage of the genomic era is the theoretical ability to access the whole

64 genome in order to sample loci representing a diversity of evolutionary rates and coalescent

65 histories, potentially overcoming problems from discordance in phylogenomics (Wolf et al.

66 2002; Rokas et al. 2003). The most widely used sampling methods include the use of large

67 numbers of coding sequences (Singhal et al. 2021), anchored hybrid enrichment loci (Lemmon et

68 al. 2012), ultra-conserved elements (Faircloth et al. 2013), and transcriptomes (Irisarri et al.

69 2017), which encompass hundreds or thousands of markers yet are still reduced representations

70 of a small proportion of the genome (Lynch 2007). The development of bioinformatics tools and

71 sequencing technologies have enabled chromosome-scale genome assemblies for non-model

72 organisms (see http://dnazoo.org, Dudchenko et al. 2017), facilitating the understanding of not

73 only which diversification events are associated with discordance, but also the genomic regions

74 or types of loci driving heterogeneous phylogenetic results.

75   Only recently have investigators looked closely at patterns of heterogeneity in order to

76 understand how molecular evolution across the genome affects inferred species relationships,

77 mainly in clades where there has been considerable phylogenetic and taxonomic debate such as

78 squamate reptiles (Burbrink et al. 2020; Singhal et al. 2021), birds (Jarvis et al. 2014; Prum et al.

79 2015), amphibians (Hime et al. 2021), and land plants (Morales-Briones et al. 2021). Meanwhile,

80 discordance in well-resolved phylogenies with high statistical support among inferred branches

81 and agreement across studies has been largely overlooked, leaving a gap in knowledge about the

82 expected range of signals and amount of disagreement between loci when there are differing

83 levels of phylogenetic uncertainty. This would be critical for informing the designs of future

84 phylogenomic studies.

Here, we examined the importance of heterogeneity in the phylogenomic estimation of

turtles (Order Testudines). Turtles are shelled reptiles with a rich fossil history dating from the

Triassic Period (Joyce 2007; Joyce et al. 2021). The approximately 350 extant species of turtles

are classified into two main groups, Pleurodira and Cryptodira, which are further divided into 8

superfamilies and 14 families (Uetz et al. 2021). Early molecular phylogenetic results disagreed

in the placement of some turtle taxa; in particular, the position of the monotypic big-headed

turtle (*Platysternon*) and its relationships with or within Testudinoidea (Shaffer et al. 1997;

Krenz et al. 2005). Other questions about deeper turtle relationships such as the position of soft-

shell turtles (Trionychoidea), the Americhelydiae, and the position of turtles in the amniote

phylogeny persisted into the next-generation sequencing era (Crawford et al. 2012, 2015). Since

then, phylogenomic studies of turtles have produced fully resolved species trees with 100%

statistical support at every node and near complete agreement across studies (Crawford et al.

2015; Shaffer et al. 2017).

Despite the harmony among phylogenomic studies of turtles using next-generation

sequencing data, the amount of underlying heterogeneity in these datasets has not been

examined. Also, while complete genome assemblies are available for a wide range of turtle

species (Shaffer et al. 2013; Wang et al. 2013; Tollis et al. 2017; Quesada et al. 2019),

phylogenomic studies have not taken advantage of this genome-wide information. Our goals

were threefold: (1) to revisit discoveries about higher turtle systematics using more substantial

proportions of the genome than previous studies; (2) to measure heterogeneity at genes and sites

in important events in turtle evolution; and (3) determine what drives heterogeneity in turtle

phylogenomics in terms of mechanisms and locus type. We show that heterogeneity in

phylogenomic datasets is to be expected, even in well-resolved clades such as turtles, and

suggest that a combination of incomplete lineage sorting and lack of gene informativeness due to

an overall slow rate of evolution may be driving incongruence between studies of turtle

relationships.

Methods and Materials

Phylogenomic analysis of high-confidence single-copy orthologs

We downloaded the complete genome assemblies of 19 turtle species plus three outgroup taxa

from NCBI. The species, assembly accession numbers, assembly lengths, and assembly

contiguities measured by scaffold N50 are shown in Table 1. Using Benchmarking Universal

Single-Copy Orthologs (BUSCO) v3 (Waterhouse et al. 2018), we extracted the nucleotide

sequences of 5,310 OrthoDB v9 tetrapod orthologs (Zdobnov et al. 2017) from each genome

assembly. BUSCO orthologs are high-confidence genes that persist in eukaryotic genomes as

single-copy, which precludes downstream problems in phylogenetics stemming from gene

duplication and hidden paralogy (Waterhouse et al. 2018). We aligned each set of orthologs with

MAFFT v7.475 (Katoh and Standley 2013), and removed erroneous columns in the alignments

with the heuristic method (*automated 1*) in TrimAL v1.4.1 (Capella-Gutierrez et al. 2009).

Summary statistics including number of taxa, alignment length, missing data percentage, number

and proportion of variable sites, number and proportion of parsimony-informative sites, and GC

content for each alignment were estimated with AMAS v3.04 (Borowiec 2016). Trimmed

alignments were filtered at a cut-off length ≥1,500 bp and a minimum of 50% taxa

representation. We estimated the average pairwise Kimura 2-parameter corrected distance for

each alignment with MEGAX (Kumar et al. 2018).

Table 1: Turtle genome assemblies analyzed in this study.

Suborder	Superfamily	Family	Species Name	Accession Number	Assembly length (Gb)	Scaffold N50 (Mb)
Cryptodira	Chelonioidea	Cheloniidae	*Chelonia mydas*	GCF_000344595.1	2.2	3.9
		Dermochelyidae	*Dermochelys coriacea*	GCA_006547105.1	2.1	0.12
	Chelydroidea	Chelydridae	*Chelydra serpentina*	GCA_007922165.1	2.5	16
	Kinosternoidea	Dermatemydidae	*Dermatemys mawii*	GCA_007922305.1	1.9	34.4
	Testudinoidea	Emydidae	*Chrysemys picta*	GCA_000241765.3	2.4	0.44
			Malaclemys terrapin	GCA_001728815.2	2.4	0.44
			Terrapene mexicana	GCF_002925995.2	2.6	24.2
		Geoemydidae	*Cuora amboinensis*	GCA_004028625.2	2.1	0.47
			Cuora mccordi	GCA_003846335.1	2.4	32.6
		Platysternidae	*Platysternon megacephalum*	GCA_003942145.1	2.3	7.2
		Testudinidae	*Chelonoidis abingdonii*	GCA_003597395.1	2.3	1.3
			Gopherus agassizii	GCA_002896415.1	2.2	0.228
	Trionychoidea	Carettochelyidae	*Carettochelys insculpta*	GCA_007922185.1	2.3	45.9
		Trionychidae	*Apalone spinifera*	GCA_000385615.1	1.9	2.3
			Pelodiscus sinensis	GCF_000230535.1	2.2	3.4
Pleurodira	Chelidoidea	Chelidae	*Emydura subglobosa*	GCA_007922225.1	2	44.8
			Mesoclemmys tuberculata	GCA_007922155.1	2	46.4
	Pelomedusoidea	Pelomedusidae	*Pelusios castaneus*	GCA_007922175.1	2	14.1
		Podocnemididae	*Podocnemis expansa*	GCA_007922195.1	2.4	37.1

Gb=gigabases; Mb=megabases

Gene trees were reconstructed for each locus using RAxML v1.8 (Stamatakis 2014). For

each ortholog, we generated 10 maximum likelihood trees with the GTRGAMMA substitution

model, and performed 100 rapid bootstrap replicates on the best tree. We then collected the best

tree for each ortholog to use as evidence to construct a species tree using ASTRAL-III (Zhang et

al. 2018), assessing branch support with local posterior probabilities (Sayyari and Mirarab 2016).

We used Snakemake (Mölder et al. 2021) to automate a reproducible bioinformatic pipeline for

BUSCO extraction, filtering, alignment, gene tree and species tree estimation, available at

https://github.com/mibyars/busco_phylo_pipeline. We also concatenated the single copy

orthologs into a supermatrix with AMAS and performed a partitioned maximum likelihood

phylogenetic reconstruction with IQ-TREE v2.1.2 (Minh et al. 2020b). We used ModelFinder

(Kalyaanamoorthy et al. 2017) set to TESTMERGE to combine partitions where necessary. We

assessed branch support on the concatenated partitioned maximum likelihood tree using 1,000

ultrafast bootstrap replicates and the Shimodaira-Hasegawa-like (SH-like) approximate

likelihood ratio test (aLRT, Shimodaira 2002).

Phylogenomic analysis of biallelic markers

In addition to single-copy orthologs from turtle and outgroup genome assemblies, we constructed

sets of biallelic single nucleotide polymorphisms (SNPs) across annotated features of the turtle

genome based on paired-end sequence data downloaded from the NCBI Short Read Archive

(SRA, Supplementary Data). We assessed each short read dataset with FastQC v0.115 (S.

Andrews - http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and trimmed the raw

reads with BBDuk v.37.41 (B. Bushnell - sourceforge.net/projects/bbmap/, last accessed

December 2020). We used SISRS to generate de novo orthologs from the processed reads

(Schwartz et al. 2015; Literman and Schwartz 2021). After subsampling all sequences to a target

depth of ~10X per taxon (except for two low-coverage species, Supplementary Data), we

assembled a composite genome using Ray v2.2.3-devel (Boisvert et al. 2010) with default

parameters and a k-value of 31. We then used SISRS to map the trimmed reads from each

species to the composite genome, retaining only uniquely mapped reads to avoid false positives

from duplicated or repetitive regions. Bases in the composite genome were replaced according to

mapped data from each species if sites were covered by at least three reads and if there was no

within-taxon variation; sites not meeting these criteria were masked as "N".

For downstream ortholog annotation, we mapped the composite sequences from each

species to the reference genome for the western painted turtle (*Chrysemys picta bellii,*

GCA_000241765.3) with Bowtie2 v2.3.4 (Langmead and Salzberg 2012), removing contigs that

either failed to map or mapped to multiple positions, and removing sites resulting from

overlapping coverage of independent scaffolds. We then annotated the mapped composite

genome sites parsed from the painted turtle genome as belonging to the following genomic

regions (locus types) with BEDTools v2.2.6 (Quinlan and Hall 2010): (1) coding sequences

(CDS); (2) 3'-UTR; (3) 5'-UTR; (4) introns; (5) other genic regions not annotated as CDS, UTR,

or intronic; (6) long-noncoding RNAs (lncRNAs); (7) pseudogenes; (8) smRNAs (including

miRNAs, ncRNAs, rRNAs, scRNAs, smRNAs, snoRNAs, snRNAs, tRNAs, and vaultRNAs).;

and (9) intergenic regions. We defined "intergenic" as all regions not included within the other

locus types.

We then used SISRS to identify parsimony-informative sites along the mapped composite

contigs and to remove invariant sites and singletons, as well as any sites containing "Ns" (i.e.

polymorphic within species) and overlapping indels or gaps. For the nine annotated genomic

regions as well as a set of the complete mapped data (=10 locus types), we concatenated biallelic

parsimony-informative sites and inferred locus type phylogenies via SVDQuartets (Chifman and

Kubatko 2014) in PAUP* v4.0a (build 168) (Swofford 2003) using the default 100 bootstrap

replicates to estimate a majority consensus tree, and maximum likelihood inference in IQ-TREE.

A schematic of all bioinformatic analyses used to extract homologous markers is found in Figure

1.

Figure 1. Bioinformatics pipelines. Schematic representing the workflows used in this study to extract homologous

markers from genome assemblies and whole genome shotgun sequences for phylogenomic inference of turtles.

BUSCO = Benchmarking Single Copy Orthologs; ML=maximum likelihood; RF = Robinson-Fould's.

Heterogeneity analyses

To measure similarity across gene genealogies, we calculated Robinson-Foulds (RF) distances

(Robinson and Foulds 1981) with PhyKIT (Steenwyk et al. 2021). First, we measured the RF

distance between each inferred single-copy ortholog gene tree and the inferred species tree. In

order to take into account missing species across some of the gene trees, we normalized the RF

distances by taking the raw RF distance and dividing it by $2(n-3)$ where n is the number of tips in

the phylogeny. We also calculated multiple pairwise RF distances between the locus type

phylogenies inferred from data mapped to each genome annotation feature with SISRS (CDS,

introns, intergenic, etc.). Finally, we calculated the RF distances of each locus type phylogeny

from the topology inferred with all mapped sites across the turtle genome.

We computed gene and site concordance factors for each branch in the turtle phylogeny

using the trimmed and filtered single-copy ortholog alignments, their gene trees, and the

coalescent-based species tree in IQ-TREE (Minh et al. 2020a). A branch's gene concordance

factor represents the percentage of decisive gene trees that also contain that branch, and a

branch's site concordance factor is the percentage of decisive alignment sites supporting that

branch. We also computed gene concordance factor and site concordance factor for each branch

in the concatenated single-copy ortholog tree of alignments >1,500 bp by computing individual

locus trees based on the partitions in IQ-TREE. Locus trees were inferred with default settings,

and the species tree was inferred using an edge-linked proportional partition model with an SH-

like approximate likelihood ratio test and ultrafast bootstrap analysis of 10,000 replicates each (-

alrt 10000 -B 10000). The greedy algorithm of PartitionFinder (Lanfear et al. 2012) was applied

to find the best-fit partitioning scheme which was then used in the subsequent tree reconstruction

step (-m TESTMERGE). We also calculated site concordance factor for every branch in the

maximum likelihood trees inferred from biallelic parsimony informative sites within locus types.

Concordance factors for genes and sites allow for complementary approaches to measure

the consistency of loci that agree with particular branches in a phylogeny, as well as

opportunities to test for incomplete lineage sorting, which we applied to each branch in the

single-copy ortholog species tree. One expectation of incomplete lineage sorting is a roughly

equal proportion of topologies supporting alternative relationships (Green et al. 2010). We used a

chi-square test to determine if similar proportions of gene trees supported conflicting

relationships at each branch, where a failure to reject the null hypothesis supported incomplete

lineage sorting at a given branch.

Divergence Time Estimation

We estimated divergence times with single-copy ortholog alignments using Bayesian modeling

implemented in BEAST v.2.6.3 (Bouckaert et al. 2019). We subsampled 685 alignments with

complete taxon sampling, from which we further randomly sampled three replicate sets of 10

genes. Based on model testing with ModelFinder, we applied the HKY+Gamma+Invariant site

model to all partitions, and utilized a relaxed lognormal molecular clock and Calibrated Yule tree

prior. To calibrate the time trees, we used 11 calibration priors from the literature (Joyce et al.

2013) which overlapped with our taxon sampling (Supplementary Data). To account for

uncertainty in fossil dating, we used hard minimum constraints and set soft maximum constraints

by placing the maximum ages in the 97.5% quantile of a lognormal prior distribution. Markov

chain Monte Carlo (MCMC) analyses were sampled every 50,000 generations. For each

replicate, we assessed convergence of parameter estimates across the MCMC by monitoring

effective sample sizes with Tracer v.1.7.2 (Rambaut et al. 2018), and we estimated a maximum

clade credibility tree with TreeAnnotator v2.6.3.

We also estimated divergence times in turtle evolution using penalized likelihood with

r8s v1.8.1 (Sanderson 1997, 2002, 2003), using the same fossil calibrations as above for

minimum and maximum node ages; the main difference in this analysis was we fixed the time to

most recent common ancestor (TMRCA) for amniotes at 312 million years ago (Ma) (Donoghue

and Benton 2007) and the TMRCA for Testudines at 220 Ma (Thomson et al. 2021). First, we

concatenated the 685 complete partitions and inferred a maximum likelihood tree with model

testing for each partition as above in IQ-TREE. We then used the inferred phylogeny with

branches in terms of substitutions per site and the 1,836,182 concatenated bases for penalized

likelihood estimation, with cross validation to optimize the smoothing parameter which

quantifies the deviation from the molecular clock.

We used linear regression to determine the relationship between the estimated divergence

time for each node and the gene and site concordance factors in the inferred trees from each

locus type in the SISRS analysis, the gene trees and the species tree based on single-copy

orthologs, and the locus trees and partitioned maximum likelihood tree based on single-copy

orthologs. We also tested for a correlation between the concordance factors and the estimated

rates of molecular evolution at each branch. We repeated these regressions using estimated

divergence times from Thomson et al. (2020).

Results

A very large number of informative sites for turtle phylogenomics

We analyzed publicly available genome assemblies to extract single copy orthologs from 19

turtles representing 13 out of 14 extant families and 7 of 8 superfamilies, plus 3 outgroup taxa

(Table 1). On average, 4,609 (83%, range 1,911-5,099) conserved tetrapod orthologs were

complete in the genome assemblies, 112 (2.1%, range 7-496) were duplicated, 170 (3.2%, range

55-485) were fragmented, and 419 (7.9%, range 143-2,907) were missing. Fifty-five percent of

conserved tetrapod orthologs were missing from the genome assembly of the trionychid *Apalone*

spinifera; therefore, we omitted this species from downstream analyses of single-copy orthologs

due to concerns about assembly quality, of which BUSCO is known to be a reliable indicator

(Waterhouse et al. 2018). Each single-copy ortholog was present in an average of 19 out of 21

species, averaged 1,902 bp in length, and contained an average of 53% variable sites and 26%

parsimony-informative sites. Alignment length was correlated with the number of parsimony

informative sites (R^2=0.69, P=2.2e-16, Fig. 3a). GC content was consistent across the single-

copy orthologs and averaged 48% (standard deviation 6.2%, Fig. 3b), as was average pairwise

Kimura 2-parameter distance (mean 0.154, median 01.39, standard deviation 0.072, Fig. 3c). The

subset of 2,513 alignments ≥1,500 bp in length had very similar characteristics to the complete

set of loci; the average large locus length was 1,903 bp and the relationship between alignment

length and the number of parsimony-informative sites held (R^2=0.73, P=2.2e-16).

Figure 2. Summary of single-copy orthologs. Percent of tetrapod orthologs from orthoDBv9 present in each

genome assembly as complete single copy, complete duplicated, fragmented, or missing.

Figure 3. Description of single copy ortholog alignments. (a) Scatterplot showing the relationship between alignment length and the number of parsimony informative sites for trimmed alignments ≥1,500 bp. (b) Histogram showing the distribution of GC content in all single copy orthologs after trimming. Solid vertical line represents the mean value (0.476). (c) Histogram showing the distribution of average pairwise Kimura 2-parameter (K2P) distances for single copy orthologs after trimming. Solid vertical line represents the mean value (0.154).

We downloaded 38 SRA datasets from 26 turtle species representing 7 of 8 superfamilies and 14 families (Supplementary Data). Based on FastQC results, we selected 32 of the datasets for further analysis ranging from ~1–70X post-trimming coverage. The composite genome consisted of 5,555,666 contigs with an N50 of 2.4 Mb. We successfully mapped 302,355,876 bases to 2,110,354 contigs in the composite genome which covered 80% of the painted turtle genome (Table 2), and called 1,655,675 parsimony-informative biallelic SNPs when allowing missing data for up to two species at a given site. The number of parsimony-informative biallelic SNPs was much larger in mapped datasets containing only cryptodires (2,787,072 bases) and testudinoideans (4,696,637 bases) (Supplementary Data), as expected based on Literman et al. (2021). We collected the highest number of biallelic SNPs from intergenic regions (746,724 bases), followed by introns (533,336 bases) and CDS (136,823 bases). The relatively low

299   number of parsimony-informative biallelic SNPs in pseudogenic (427) and smRNA regions

300   (137) is likely due to a small number of annotations for these features in the western painted

301   turtle genome.

303   **Table 2. Summary of data mapped to the western painted turtle (*Chrysemys picta*) genome annotations.**

Locus Type	Annotated Bases in the Genome	Based Mapped by SISRS	Percent of Annotated Bases Mapped
All Mapped	2,365,766,571	302,355,876	12.78%
Intergenic	1,382,145,112	161,445,195	11.68%
Intronic	812,010,672	115,297,781	14.20%
lncRNA	64,450,450	9,451,244	14.66%
Genic Other	65,764,053	9,052,113	13.76%
CDS	33,486,790	5,718,074	17.08%
3'-UTR	8,034,514	1,510,783	18.80%
5'-UTR	5,096,255	708,214	13.90%
Pseudogenic	670,023	59,058	8.81%
smRNA	133,894	20,076	14.99%

Bioinformatic analysis of turtle genomes yielded well-resolved species relationships

The inferred relationships among turtle lineages based on single-copy orthologs were consistent

with previous studies using phylogenomic markers (Crawford et al. 2015; Shaffer et al. 2017),

with full statistical support for all resulting branches based on posterior probabilities in the

multilocus coalescent tree, and bootstrap replicates and SH-like approximate likelihood in the

supermatrix tree (Fig. 4, Supplementary Fig. 1, Supplementary Fig. 2). Hereafter we refer to

these results collectively as the turtle species tree. The turtle species tree supports the reciprocal

monophyly of Pleurodira and Cryptodira. Within cryptodires, there is a split between the

Trionychoidea (soft-shelled turtles) and the Durocryptodira (Crawford et al. 2015), and a further

subdivision of Durocryptodira into the Testudinoidea (pond turtles, big-headed turtles,

geoemyds, and tortoises) and Americhelydia (sea turtles, snapping turtles, mud turtles; Joyce et

al. 2013). Within the Testudinoidea, the turtle species tree supports Testuguria (geoemyds and

tortoises; Joyce et al. 2004), and a *Platysternon* + Emydidae clade ("Platyemidae"; Crawford et

al. 2015).

Figure 4. Multilocus coalescent species tree of turtles based on high-confidence single-copy orthologs.
Topology of the species tree inferred from 2,513 single-copy orthologs ≥1,500 bp in length. All branches received
100% local posterior support. Numbers on branches indicate gene and site concordance factors before and after the
slash, respectively. Superfamilies are indicated by vertical text. Key clades are labelled with letters in boxes. (a)
Durocryptodira, (b) Americhelydia, (c) Testuguria, (d) Emydidae. Additional outgroup *Homo sapiens* not shown.

Maximum likelihood and quartets trees from the mapped data, particularly for all-
mapped, CDS, introns, and intergenic regions, were consistent with the species tree based on
single-copy orthologs (Supplementary Fig. 3-12) although this was highly dependent on the
amount of allowed missing data and gapless sites, and was variable between the taxonomic
specificity of the data used to map sites (Supplementary Data). When there was disagreement

between trees from mapped regions and the species tree, the difference was often in the

placement of *Platysternon* (Fig. 5). For instance, when allowing zero missing gapless parsimony-

informative sites from the all-turtle data, 5' UTR placed *Platysternon* as the outgroup of

Geoemydidae + Testudinidae. When allowing up to two missing gapless parsimony-informative

sites from the all-turtle data, pseudogenic regions (426 sites) also placed *Platysternon* as the

outgroup of Geoemydidae + Testudinidae, while smRNA (137 sites) agreed with the species tree

and all-mapped topologies. Meanwhile, at multiple levels of allowed missing data the

Testudinoidea-only mapped sites weakly supported *Platysternon* as the sister taxon to

Geoemydidae + Testudinidae + Emydidae (Supplementary Data).

Figure 5. Alternative topologies in turtle phylogenetics based on markers from different regions of the

genome. (a) *Platysternon* as the sister taxon to Emydidae to the exclusion of Testuguria (Geoemydidae +

Testudinidae), consistent with the turtle species tree (current study), Crawford et al. (2015), and Shaffer et al.

(2017). (b) *Platysternon* as the sister taxon to Testuguria with the exclusion of Emydidae, consistent with 17% of the

single-copy ortholog genes trees, mapped 5'-UTR parsimony-informative sites data with zero missing data, mapped

pseudogene parsimony-informative sites when allowing 2 missing taxa per site, and Krenz et al. (2005). (c)

Platysternon as the sister taxon to Geoemydidae + Testudinidae + Emydidae, consistent with 20% of single-copy

ortholog gene trees and Shaffer et al. (1997).

Heterogeneity is prevalent at well resolved nodes in the turtle phylogeny

While results from our multilocus and biallelic sites analyses of turtles are consistent and largely

in agreement with other studies based on 100% branch support, some genomic regions, either

from single-copy orthologs or from genome feature annotations, yielded genealogies conflicting

with the turtle species tree. While most gene trees from single-copy orthologs matched the

species tree based on normalized RF distances, there was a long tail to this distribution indicating

gene trees with greater dissimilarity (Fig. 6a). Quartet trees using mapped data from CDS,

introns, and intergenic regions were consistent with both the single-copy ortholog species tree

and a quartet tree derived from the all-mapped data, except for pseudogenic DNA, 5' UTR, and

smRNA which deviated from the all-mapped data in terms of RF distances (Supplementary Fig.

13).

We also found a high degree of heterogeneity based on gene and site concordance factors

throughout the turtle phylogeny (Fig. 1, Supplementary Fig. 1, Supplementary Fig. 2). For

instance, while local posterior probabilities were uniformly 100% across branches of the

multilocus coalescent species tree, gene concordance factors ranged from 44 to 97 and site

concordance factors ranged from 30 to 71 (Fig. 6b-d). Gene and site concordance factors based

on the multilocus species tree were correlated with each other (Figure 6b; R^2=0.55, P=0.0002)

and with branch lengths (Fig. 6c-d; R^2=0.88, 3.88e-09 and R^2=0.49, P=0.0007, respectively). We

also estimated gene and site concordance factors based on the locus trees with respect to the

maximum likelihood tree inferred from the concatenated supermatrix, and found a similar range

of gene concordance factors (34 to 92) and site concordance factors (30 to 71) despite 100%

bootstrap support, as well as a strong correlation between the two concordance measures

(R^2=0.60, P=0.0001). Gene and site concordance factors based on the concatenated partitioned

maximum likelihood tree were also correlated with branch lengths (R^2=0.90, 1.02e-09 and

R^2=0.58, P=0.0001, respectively).

Figure 6. Phylogenomic heterogeneity in turtles. (a) Histogram of RF distances between gene trees and species

tree. (b-c) Concordance factors calculated using single-copy ortholog alignments, their gene trees, and the

multilocus coalescent species tree. Results based on locus trees and concatenated supermatrix maximum likelihood

tree are in Supplementary Materials. (b) Scatterplot depicting the relationship between gene and site concordance

factors at each node. (c) Scatterplot depicting the relationship between gene concordance factor and branch length.

(d) Scatterplot depicting the relationship between site concordance factor and branch length.

The root of the turtle species tree was assigned a relatively large gene concordance factor

(94.5) and site concordance factor (64.2), suggesting strong support for reciprocally

monophyletic Pleurodira and Cryptodira across genes and sites. We rejected incomplete lineage

sorting at the root of the turtle phylogeny and the roots of Cryptodira (P=8.05E-04) and

Pleurodira (P=2.32E-04). The placement of Trionychoidea as sister taxon to all other cryptodires

was assigned relatively large gene concordance factor (84.56) but small site concordance factor

(39.26), suggesting that there are a large number of sites supporting alternative bifurcations. One

area with particularly high discordance (i.e. relatively small gene concordance factor=46.9 and

site concordance factor=43.8) was the placement of the big-headed turtle (*Platysternon*) with

respect to Emydidae, for which we could not reject incomplete lineage sorting (see Table 3, Fig.

5 and Discussion).

Table 3. Gene concordance factors for nodes in the species tree phylogeny where incomplete lineage sorting

was not rejected.

Clade description	Node	Number of possible supporting gene trees	Gene concordance factor	Percent of gene trees supporting first alternative topology	Percent of gene trees supporting second alternative topology
Durocryptodira	28	2387	88.73	0.54	0.21
Chelonoidea	30	1755	89.46	1.48	0.8
Testudinoidea	32	2435	80.66	0.57	0.7
Testuguria	33	2255	84.7	0.62	0.8
Testudinidae	34	1989	85.22	4.17	5.38
Cuora	35	1932	90.94	2.69	2.07
Emydidae + Platysternidae	36	2270	46.92	17.05	19.52
Malachlemys + Chrysemys	38	1730	86.99	5.03	4.39
Pelomedusoidea	40	2195	95.81	1.32	1.14
Chelidoidea	41	2138	96.49	0.8	1.08

Heterogeneity at different timescales in turtle evolution

Results from the replicate BEAST analyses consistently underestimated turtle divergence times

with respect to results from the penalized likelihood analysis as well as those of Thomson et al.

(2020), even though the TMRCA for all taxa (i.e. the time of the mammal-reptile split) was

consistent (Supplementary Data). Across locus types from the SISRS analysis, we found that site

concordance factors were not strongly correlated with estimated node ages, regardless of whether

or not we used our estimated divergence times or those from Thomson et al. (2021) (Fig. 7,

Supplementary Fig. 14). For most locus types the relationship between site concordance factor

and divergence time at a given node trended slightly negatively, suggesting a pattern of higher

concordance at younger nodes, although these comparisons were non-significant (Fig. 7b,

P>0.1). The one exception to this trend was for pseudogenes, for which the relationship between

site concordance factors and estimated divergence times trended positive albeit without

significance or with only moderate significance (P=0.072 using Thomson et al. 2020's

divergence time estimates). The relationship between site concordance factors and estimated

DNA substitution rates trended slightly positive for all locus types from the SISRS analysis, but

was non-significant in all comparisons (Supplementary Fig. 15).

We also found a negative trend in the relationship between site concordance factor and

divergence time at a given node based on both the multilocus species tree and the supermatrix

tree based on single-copy orthologs, similar to the patterns observed using the locus types from

SISRS (Fig. 8a-c; Supplementary Fig. 16). However, we found a positive (albeit non-significant)

trend in the relationship between the estimated divergence time at a given node and its gene

concordance factor when using both trees derived from single-copy orthologs (Fig. 8e-g;

Supplementary Fig. 16). There was no clear relationship between the site concordance factor of a

given node and the estimated substitution rate at the preceding branch (Fig. 8d), and a

moderately significant positive trend (P≤0.05) for the relationship between the gene concordance

factor of a given node and the estimated substitution rate at the preceding branch (Fig. 8h;

Supplementary Fig. 16).

Figure 7. Site concordance and estimated node ages in turtle evolution. (a) Chronogram showing estimated

divergence times in millions of years based on penalized likelihood and fossil calibrations of labelled nodes. 1.

Testudines, 2. Cryptodira, 3. Trionychoidea, 4. Durocryptodira, 5. Geoemydidae, 6. Platyemidae, 7. Emydidae, 8.

Americhelydia, 9. Chelonioidea, 10. Pleurodira, 11. Pelomedusoidea. (b) Site concordance factors (sCF) for each

node in phylogenies inferred from nine locus types and the estimated node age in millions of years ago (Ma) based

on penalized likelihood.

Figure 8. Concordance at genes and sites, estimated node ages, and DNA substitution rates in turtle evolution.

Results shown are from comparisons of single-copy ortholog alignments and their gene trees to the multilocus

coalescent species tree inferred using ASTRAL. There was a negative trend between site concordance factors (sCF)

and estimated node age (in millions of years or Ma) based on penalized likelihood (PL) from the current study (a),

estimated node ages in Ma from Thomson et al. (2020) (b), estimated node ages in Ma from the BEAST analysis in

the current study (c), and no discernable trend between sCF and estimated DNA substitution rates in terms of

substitutions per site per Ma (d). There was a positive trend between gene concordance factors (gCF) and estimated

node ages (e, f, g) and a moderately significant positive relationship between gene concordance factors and

estimated DNA substitution rates (h). Results from comparisons of single-copy ortholog alignments, locus trees, and

the maximum likelihood tree inferred from the concatenated supermatrix are shown in Supplementary Materials.

Discussion

While heterogeneity has been recognized as an important feature of phylogenomic datasets, and

the exploration of the patterns and sources of gene tree and site discordance has become an

essential step when reconstructing the branching order of life with molecular data (Kumar et al.

451 2012; Minh et al. 2020a), to our knowledge there has not been an assessment of phylogenomic

452 heterogeneity in turtles until now. Here, we have constructed a robust dataset based on whole

453 genomes with the goal of disentangling conflicting hypotheses about turtle relationships. Single-

454 copy orthologs yielded more than double the variation and triple the number of parsimony

455 informative sites for turtles compared to other high-throughput methods (Crawford et al. 2015;

456 Shaffer et al. 2017). In addition, the mapped data included millions of parsimony informative

457 sites that were further parsed by locus type. We identified splits in the turtle phylogeny and

458 regions of the turtle genome associated with gene tree and site heterogeneity, and investigated

459 potential sources of this discordance. We found that heterogeneity in turtle phylogenomics can

460 be explained by a combination of factors, such as incomplete lineage sorting and lack of gene

461 informativeness, and has contributed to conflicting results in phylogenetic studies of turtles. The

462 results of our study demonstrate that heterogeneity can be rampant, even in phylogenomic

463 datasets which have yielded fully resolved species trees.

464   Gene tree-species tree discordance in turtles may be partly due to an overall slow

465 substitution rate compared to other amniotes (Tollis et al. 2018; Gemmell et al. 2020), resulting

466 in a lack of gene informativeness at more recent divergences. We found no significant

467 relationship between site concordance and node age across most locus types, a uniform

468 distribution of site concordances across the range of estimated DNA substitution rates across

469 most annotated genomic regions, and that the average pairwise Kimura 2-parameter distances

470 across single-copy orthologs were also fairly uniform. This suggests relatively consistent

471 substitution rates across the turtle genome, as well as across turtle evolution. Where we did

472 estimate elevated DNA substitution rates along branches in the turtle phylogeny, we found that

473 gene concordance factors were significantly larger. This suggests that shifts in the rate of

evolution may lead to a comparatively rapid accumulation of mutations during certain time

periods, resulting in more gene trees in agreement with the species phylogeny. Meanwhile, the

gene concordances at older nodes were larger (albeit non-significantly), suggesting that

heterogeneity at more recent nodes may be more driven by stochastic error due to limited

information at slowly evolving loci. Our results are consistent with the longstanding prediction

that slowly evolving loci, which includes much of the turtle genome, provide disproportionally

greater phylogenetic information for older splits (Townsend et al. 2008).

Interestingly, site concordance in pseudogenes was positively associated (albeit non-

significantly) with node age, suggesting that even slightly elevated substitution rates in some

noncoding regions of the turtle genome can lead to more informative sites supporting the species

tree topology at older nodes. This is somewhat different from what was reported in Literman et

al. (2021), who showed that noncoding regions such as introns and intergenic regions supported

more recent splits in the mammalian phylogeny, and may be again related to the fact that

estimated substitution rates in mammals are consistently higher than for other amniotes such as

turtle and crocodilians. However, our inferences from pseudogenes are based on a relatively

small number of parsimony informative sites (427). More comprehensive annotations for

noncoding regions of turtle genomes are needed to investigate this further.

Our tests for incomplete lineage sorting reveal that population-level processes likely also

have played a key role at multiple stages of turtle diversification. In particular, the phylogenetic

position of *Platysternon* has been highly contentious (Joyce et al. 2004, 2021), with early

molecular studies in disagreement (Shaffer et al. 1997; Krenz et al. 2005; Parham et al. 2006;

Barley et al. 2010). Our turtle species tree results mirror Crawford et al. (2015) and Shaffer et al.

(2017), fully supporting *Platysternon* as the sister taxon to Emydidae. However, only 47% of

single-copy ortholog gene trees are in agreement with a *Platysternon* + Emydidae clade, and

37% of gene trees support only two alternative relationships: of these, 17% of gene trees support

Platysternon as the sister taxon to Testuguria (Geoemydidae + Testudinidae) following Krenz et

al. (2005), and 20% support *Platysternon* as the sister taxon to Testudinoidea (Emydidae +

Geoemydidae + Testudinidae) following Shaffer et al. (1997). The proportion of gene trees

supporting these alternative topologies was similar according to our chi-square test (Table 3),

and supports incomplete lineage sorting rather than introgression or statistical noise (Green et al.

2010). As these loci support published alternative hypotheses concerning the relationship of

Platysternon to other Testudinoidea, we suggest that incomplete lineage sorting is a previously

unrecognized driver of conflicting phylogenetic results in the higher classification of turtles.

Acknowledgements

This research was funded by the National Science Foundation (#1812291 to RL) and by startup

funds provided to MT by the School of Informatics, Computing, and Cyber Systems at Northern

Arizona University. We would like to acknowledge the Monsoon computing cluster at Northern

Arizona University (https://nau.edu/high-performance-computing/) for providing the

computational resources necessary to carry out this study.

Data Availability

The bioinformatic workflows used to extract and analyze single-copy orthologs from genome

assemblies are available at https://github.com/mibyars/busco_phylo_pipeline. Data generated for

this study including alignments and trees are available at the Harvard Dataverse

(https://doi.org/10.7910/DVN/WYQRPC).

Literature Cited

Barley A.J., Spinks P.Q., Thomson R.C., Shaffer H.B. 2010. Fourteen nuclear genes provide

phylogenetic resolution for difficult nodes in the turtle tree of life. Mol. Phylogenet. Evol.

55:1189–1194.

Boisvert S., Laviolette F., Corbeil J. 2010. Ray: Simultaneous Assembly of Reads from a Mix of

High-Throughput Sequencing Technologies. J. Comput. Biol. 17:1519–1533.

Borowiec M.L. 2016. AMAS: a fast tool for alignment manipulation and computing of summary

statistics. PeerJ. 4:e1660.

Bouckaert R., Vaughan T.G., Barido-Sottani J., Duchêne S., Fourment M., Gavryushkina A.,

Heled J., Jones G., Kühnert D., Maio N.D., Matschiner M., Mendes F.K., Müller N.F.,

Ogilvie H.A., Plessis L. du, Popinga A., Rambaut A., Rasmussen D., Siveroni I., Suchard

M.A., Wu C.-H., Xie D., Zhang C., Stadler T., Drummond A.J. 2019. BEAST 2.5: An

advanced software platform for Bayesian evolutionary analysis. PLOS Comput. Biol.

15:e1006650.

Burbrink F.T., Grazziotin F.G., Pyron R.A., Cundall D., Donnellan S., Irish F., Keogh J.S.,

Kraus F., Murphy R.W., Noonan B., Raxworthy C.J., Ruane S., Lemmon A.R., Lemmon

E.M., Zaher H. 2020. Interrogating Genomic-Scale Data for Squamata (Lizards, Snakes,

and Amphisbaenians) Shows no Support for Key Traditional Morphological

Relationships. Syst. Biol. 69:502-520.

Chifman J., Kubatko L. 2014. Quartet Inference from SNP Data Under the Coalescent Model.

Bioinformatics. 30:3317–3324.

Cooper E.D. 2014. Overly simplistic substitution models obscure green plant phylogeny. Trends

543   Plant Sci. 19:576–582.

544 Cox C.J., Li B., Foster P.G., Embley T.M., Civáň P. 2014. Conflicting Phylogenies for Early

545   Land Plants are Caused by Composition Biases among Synonymous Substitutions. Syst.

546   Biol. 63:272–279.

547 Crawford N.G., Faircloth B.C., McCormack J.E., Brumfield R.T., Winker K., Glenn T.C. 2012.

548   More than 1000 ultraconserved elements provide evidence that turtles are the sister group

549   of archosaurs. Biol. Lett. 8:783–786.

550 Crawford N.G., Parham J.F., Sellas A.B., Faircloth B.C., Glenn T.C., Papenfuss T.J., Henderson

551   J.B., Hansen M.H., Simison W.B. 2015. A phylogenomic analysis of turtles. Mol.

552   Phylogenet. Evol. 83:250–257.

553 Donoghue P.C.J., Benton M.J. 2007. Rocks and clocks: calibrating the Tree of Life using fossils

554   and molecules. Trends Ecol. Evol. 22:424–431.

555 Dudchenko O., Batra S.S., Omer A.D., Nyquist S.K., Hoeger M., Durand N.C., Shamim M.S.,

556   Machol I., Lander E.S., Aiden A.P., Aiden E.L. 2017. De novo assembly of the *Aedes*

557   *aegypti* genome using Hi-C yields chromosome-length scaffolds. Science. 356:92–95.

558 Edwards S.V., Xi Z., Janke A., Faircloth B.C., McCormack J.E., Glenn T.C., Zhong B., Wu S.,

559   Lemmon E.M., Lemmon A.R., Leaché A.D., Liu L., Davis C.C. 2016. Implementing and

560   testing the multispecies coalescent model: A valuable paradigm for phylogenomics. Mol.

561   Phylogenet. Evol. 94:447–462.

562 Faircloth B.C., Sorenson L., Santini F., Alfaro M.E. 2013. A Phylogenomic Perspective on the

563   Radiation of Ray-Finned Fishes Based upon Targeted Sequencing of Ultraconserved

564   Elements (UCEs). PLOS ONE. 8:e65923.

565 Foster P.G. 2004. Modeling Compositional Heterogeneity. Syst. Biol. 53:485–495.

Galtier N., Daubin V. 2008. Dealing with incongruence in phylogenomic analyses. Philos. Trans.

R. Soc. B Biol. Sci. 363:4023–4029.

Gemmell N.J., Rutherford K., Prost S., Tollis M., Winter D., Macey J.R., Adelson D.L., Suh A.,

Bertozzi T., Grau J.H., Organ C., Gardner P.P., Muffato M., Patricio M., Billis K., Martin

F.J., Flicek P., Petersen B., Kang L., Michalak P., Buckley T.R., Wilson M., Cheng Y.,

Miller H., Schott R.K., Jordan M.D., Newcomb R.D., Arroyo J.I., Valenzuela N., Hore

T.A., Renart J., Peona V., Peart C.R., Warmuth V.M., Zeng L., Kortschak R.D., Raison

J.M., Zapata V.V., Wu Z., Santesmasses D., Mariotti M., Guigó R., Rupp S.M., Twort

V.G., Dussex N., Taylor H., Abe H., Bond D.M., Paterson J.M., Mulcahy D.G., Gonzalez

V.L., Barbieri C.G., DeMeo D.P., Pabinger S., Van Stijn T., Clarke S., Ryder O.,

Edwards S.V., Salzberg S.L., Anderson L., Nelson N., Stone C., Ngatiwai Trust Board.

2020. The tuatara genome reveals ancient features of amniote evolution. Nature.

584:403–409.

Green R.E., Krause J., Briggs A.W., Maricic T., Stenzel U., Kircher M., Patterson N., Li H., Zhai

W., Fritz M.H.-Y., Hansen N.F., Durand E.Y., Malaspinas A.-S., Jensen J.D., Marques-

Bonet T., Alkan C., Prüfer K., Meyer M., Burbano H.A., Good J.M., Schultz R., Aximu-

Petri A., Butthof A., Höber B., Höffner B., Siegemund M., Weihmann A., Nusbaum C.,

Lander E.S., Russ C., Novod N., Affourtit J., Egholm M., Verna C., Rudan P., Brajkovic

D., Kucan Ž., Gušic I., Doronichev V.B., Golovanova L.V., Lalueza-Fox C., Rasilla M.

de la, Fortea J., Rosas A., Schmitz R.W., Johnson P.L.F., Eichler E.E., Falush D., Birney

E., Mullikin J.C., Slatkin M., Nielsen R., Kelso J., Lachmann M., Reich D., Pääbo S.

2010. A Draft Sequence of the Neandertal Genome. Science. 328:710–722.

Hime P.M., Lemmon A.R., Lemmon E.C.M., Prendini E., Brown J.M., Thomson R.C., Kratovil

J.D., Noonan B.P., Pyron R.A., Peloso P.L.V., Kortyna M.L., Keogh J.S., Donnellan

S.C., Mueller R.L., Raxworthy C.J., Kunte K., Ron S.R., Das S., Gaitonde N., Green

D.M., Labisko J., Che J., Weisrock D.W. 2021. Phylogenomics Reveals Ancient Gene

Tree Discordance in the Amphibian Tree of Life. Syst. Biol. 70:48-66.

Irisarri I., Baurain D., Brinkmann H., Delsuc F., Sire J.-Y., Kupfer A., Petersen J., Jarek M.,

Meyer A., Vences M., Philippe H. 2017. Phylotranscriptomic consolidation of the jawed

vertebrate timetree. Nat. Ecol. Evol. 1:1370–1378.

Jarvis E.D., Mirarab S., Aberer A.J., Li B., Houde P., Li C., Ho S.Y.W., Faircloth B.C., Nabholz

B., Howard J.T., Suh A., Weber C.C., da Fonseca R.R., Li J., Zhang F., Li H., Zhou L.,

Narula N., Liu L., Ganapathy G., Boussau B., Bayzid M.S., Zavidovych V., Subramanian

S., Gabaldón T., Capella-Gutiérrez S., Huerta-Cepas J., Rekepalli B., Munch K.,

Schierup M., Lindow B., Warren W.C., Ray D., Green R.E., Bruford M.W., Zhan X.,

Dixon A., Li S., Li N., Huang Y., Derryberry E.P., Bertelsen M.F., Sheldon F.H.,

Brumfield R.T., Mello C.V., Lovell P.V., Wirthlin M., Schneider M.P.C., Prosdocimi F.,

Samaniego J.A., Velazquez A.M.V., Alfaro-Núñez A., Campos P.F., Petersen B.,

Sicheritz-Ponten T., Pas A., Bailey T., Scofield P., Bunce M., Lambert D.M., Zhou Q.,

Perelman P., Driskell A.C., Shapiro B., Xiong Z., Zeng Y., Liu S., Li Z., Liu B., Wu K.,

Xiao J., Yinqi X., Zheng Q., Zhang Y., Yang H., Wang J., Smeds L., Rheindt F.E., Braun

M., Fjeldså J., Orlando L., Barker F.K., Jønsson K.A., Johnson W., Koepfli K.-P.,

O'Brien S., Haussler D., Ryder O.A., Rahbek C., Willerslev E., Graves G.R., Glenn T.C.,

McCormack J., Burt D., Ellegren H., Alström P., Edwards S.V., Stamatakis A., Mindell

D.P., Cracraft J., Braun E.L., Warnow T., Jun W., Gilbert M.T.P., Zhang G. 2014.

Whole-genome analyses resolve early branches in the tree of life of modern birds.

Science. 346:1320–1331.

Joyce W.G. 2007. Phylogenetic Relationships of Mesozoic Turtles. Bull. Peabody Mus. Nat.

Hist. 48:3–102.

Joyce W.G., Anquetin J., Cadena E.-A., Claude J., Danilov I.G., Evers S.W., Ferreira G.S.,

Gentry A.D., Georgalis G.L., Lyson T.R., Pérez-García A., Rabi M., Sterli J., Vitek N.S.,

Parham J.F. 2021. A nomenclature for fossil and living turtles using phylogenetically

defined clade names. Swiss J. Palaeontol. 140:5.

Joyce W.G., Parham J.F., Gauthier J.A. 2004. Developing a protocol for the conversion of rank-

based taxon names to phylogenetically defined clade names, as exemplified by turtles. J.

Paleontol. 78:989–1013.

Joyce W.G., Parham J.F., Lyson T.R., Warnock R.C.M., Donoghue P.C.J. 2013. A divergence

dating analysis of turtles using fossil calibrations: an example of best practices. J.

Paleontol. 87:612–634.

Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., von Haeseler A., Jermiin L.S. 2017.

ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods.

14:587–589.

Katoh K., Standley D.M. 2013. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. Mol. Biol. Evol. 30:772–780.

Krenz J.G., Naylor G.J.P., Shaffer H.B., Janzen F.J. 2005. Molecular phylogenetics and

evolution of turtles. Mol. Phylogenet. Evol. 37:178–191.

Kubatko L.S., Degnan J.H. 2007. Inconsistency of Phylogenetic Estimates from Concatenated

Data under Coalescence. Syst. Biol. 56:17–24.

Kumar S., Filipski A.J., Battistuzzi F.U., Kosakovsky Pond S.L., Tamura K. 2012. Statistics and

Truth in Phylogenomics. Mol. Biol. Evol. 29:457–472.

Kumar S., Stecher G., Li M., Knyaz C., Tamura K. 2018. MEGA X: Molecular Evolutionary

Genetics Analysis across Computing Platforms. Mol. Biol. Evol. 35:1547–1549.

Lanfear R., Calcott B., Ho S.Y.W., Guindon S. 2012. Partitionfinder: combined selection of

partitioning schemes and substitution models for phylogenetic analyses. Mol. Biol. Evol.

29:1695–1701.

Langmead B., Salzberg S.L. 2012. Fast gapped-read alignment with Bowtie 2. Nat. Methods.

9:357–359.

Lemmon A.R., Emme S.A., Lemmon E.M. 2012. Anchored Hybrid Enrichment for Massively

High-Throughput Phylogenomics. Syst. Biol. 61:727–744.

Literman R., Schwartz R. 2021. Genome-Scale Profiling Reveals Noncoding Loci Carry Higher

Proportions of Concordant Data. Mol. Biol. Evol.

Lopes F., Oliveira L.R., Kessler A., Beux Y., Crespo E., Cárdenas-Alayza S., Majluf P.,

Sepúlveda M., Brownell R.L. Jr., Franco-Trecu V., Páez-Rosas D., Chaves J., Loch C.,

Robertson B.C., Acevedo-Whitehouse K., Elorriaga-Verplancken F.R., Kirkman S.P.,

Peart C.R., Wolf J.B.W., Bonatto S.L. 2021. Phylogenomic Discordance in the Eared

Seals is best explained by Incomplete Lineage Sorting following Explosive Radiation in

the Southern Hemisphere. Syst. Biol. 70:786–802.

Lynch M. 2007. The Origins of Genome Architecture. Sinauer Associates Sunderland, MA.

Minh B.Q., Hahn M.W., Lanfear R. 2020a. New Methods to Calculate Concordance Factors for

Phylogenomic Datasets. Mol. Biol. Evol. 37:2727–2733.

Minh B.Q., Schmidt H.A., Chernomor O., Schrempf D., Woodhams M.D., von Haeseler A.,

Lanfear R. 2020b. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. Mol. Biol. Evol. 37:1530–1534.

Mölder F., Jablonski K.P., Letcher B., Hall M.B., Tomkins-Tinch C.H., Sochat V., Forster J.,

Lee S., Twardziok S.O., Kanitz A., Wilm A., Holtgrewe M., Rahmann S., Nahnsen S.,

Köster J. 2021. Sustainable data analysis with Snakemake.

https://f1000research.com/articles/10-33.

Morales-Briones D.F., Kadereit G., Tefarikis D.T., Moore M.J., Smith S.A., Brockington S.F.,

Timoneda A., Yim W.C., Cushman J.C., Yang Y. 2021. Disentangling Sources of Gene

Tree Discordance in Phylogenomic Data Sets: Testing Ancient Hybridizations in

Amaranthaceae s.l. Syst. Biol. 70:219-235.

Pamilo P., Nei M. 1988. Relationships between gene trees and species trees. Mol. Biol. Evol.

5:568–583.

Parham J.F., Feldman C.R., Boore J.L. 2006. The complete mitochondrial genome of the

enigmatic bigheaded turtle (Platysternon): description of unusual genomic features and

the reconciliation of phylogenetic hypotheses based on mitochondrial and nuclear DNA.

BMC Evol. Biol. 6:11.

Prum R.O., Berv J.S., Dornburg A., Field D.J., Townsend J.P., Lemmon E.M., Lemmon A.R.

2015. A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA

sequencing. Nature. 526:569–573.

Quesada V., Freitas-Rodríguez S., Miller J., Pérez-Silva J.G., Jiang Z.-F., Tapia W., Santiago-

Fernández O., Campos-Iglesias D., Kuderna L.F.K., Quinzin M., Álvarez M.G., Carrero

D., Beheregaray L.B., Gibbs J.P., Chiari Y., Glaberman S., Ciofi C., Araujo-Voces M.,

Mayoral P., Arango J.R., Tamargo-Gómez I., Roiz-Valle D., Pascual-Torner M., Evans

B.R., Edwards D.L., Garrick R.C., Russello M.A., Poulakakis N., Gaughran S.J., Rueda

D.O., Bretones G., Marquès-Bonet T., White K.P., Caccone A., López-Otín C. 2019.

Giant tortoise genomes provide insights into longevity and age-related disease. Nat. Ecol.

Evol. 3:87–95.

Quinlan A.R., Hall I.M. 2010. BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics. 26:841–842.

Rambaut A., Drummond A.J., Xie D., Baele G., Suchard M.A. 2018. Posterior summarization in

Bayesian phylogenetics using Tracer 1.7. Syst. Biol. 67:901–904.

Robinson D.F., Foulds L.R. 1981. Comparison of phylogenetic trees. Math. Biosci. 53:131–147.

Rokas A., Williams B.L., King N., Carroll S.B. 2003. Genome-scale approaches to resolving

incongruence in molecular phylogenies. Nature. 425:798–804.

Sanderson M.J. 1997. A nonparametric approach to estimating divergence times in the absence

of rate constancy. Mol. Biol. Evol.

Sanderson M.J. 2002. Estimating absolute rates of molecular evolution and divergence times: a

penalized likelihood approach. Mol. Biol. Evol. 19:101–109.

Sanderson M.J. 2003. r8s: inferring absolute rates of molecular evolution and divergence times

in the absence of a molecular clock. Bioinformatics. 19:301–302.

Sayyari E., Mirarab S. 2016. Fast Coalescent-Based Computation of Local Branch Support from

Quartet Frequencies. Mol. Biol. Evol. 33:1654–1668.

Schwartz R.S., Harkins K.M., Stone A.C., Cartwright R.A. 2015. A composite genome approach

to identify phylogenetically informative data from next-generation sequencing. BMC

Bioinformatics. 16:193.

Shaffer H.B., McCartney-Melstad E., Near T.J., Mount G.G., Spinks P.Q. 2017. Phylogenomic

analyses of 539 highly informative loci dates a fully resolved time tree for the major

704   clades of living turtles (Testudines). Mol. Phylogenet. Evol. 115:7–15.

705   Shaffer H.B., Meylan P., McKnight M.L. 1997. Tests of turtle phylogeny: molecular,

706   morphological, and paleontological approaches. Syst. Biol. 46:235–268.

707   Shaffer H.B., Minx P., Warren D.E., Shedlock A.M., Thomson R.C., Valenzuela N., Abramyan

708   J., Amemiya C.T., Badenhorst D., Biggar K.K., Borchert G.M., Botka C.W., Bowden

709   R.M., Braun E.L., Bronikowski A.M., Bruneau B.G., Buck L.T., Capel B., Castoe T.A.,

710   Czerwinski M., Delehaunty K.D., Edwards S.V., Fronick C.C., Fujita M.K., Fulton L.,

711   Graves T.A., Green R.E., Haerty W., Hariharan R., Hernandez O., Hillier L.W.,

712   Holloway A.K., Janes D., Janzen F.J., Kandoth C., Kong L., de Koning A.P.J., Li Y.,

713   Literman R., McGaugh S.E., Mork L., O'Laughlin M., Paitz R.T., Pollock D.D., Ponting

714   C.P., Radhakrishnan S., Raney B.J., Richman J.M., St John J., Schwartz T., Sethuraman

715   A., Spinks P.Q., Storey K.B., Thane N., Vinař T., Zimmerman L.M., Warren W.C.,

716   Mardis E.R., Wilson R.K. 2013. The western painted turtle genome, a model for the

717   evolution of extreme physiological adaptations in a slowly evolving lineage. Genome

718   Biol. 14:R28.

719   Shimodaira H. 2002. An Approximately Unbiased Test of Phylogenetic Tree Selection. Syst.

720   Biol. 51:492–508.

721   Singhal S., Colston T.J., Grundler M.R., Smith S.A., Costa G.C., Colli G.R., Moritz C., Pyron

722   R.A., Rabosky D.L. 2021. Congruence and Conflict in the Higher-Level Phylogenetics of

723   Squamate Reptiles: An Expanded Phylogenomic Perspective. Syst. Biol. 70:542–557.

724   Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

725   large phylogenies. Bioinformatics. 30:1312–1313.

Steenwyk J.L., Buida T.J. III, Labella A.L., Li Y., Shen X.-X., Rokas A. 2021. PhyKIT: a

broadly applicable UNIX shell toolkit for processing and analyzing phylogenomic data.

Bioinformatics. btab096.

Swofford D.L. 2003. PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods).

Sinauer associates Sunderland, MA.

Thomson R.C., Spinks P.Q., Shaffer H.B. 2021. A global phylogeny of turtles reveals a burst of

climate-associated diversification on continental margins. Proc. Natl. Acad. Sci. 118.

Tollis M., Denardo D.F., Cornelius J.A., Dolby G.A., Edwards T., Henen B.T., Karl A.E.,

Murphy R.W., Kusumi K. 2017. The Agassiz's desert tortoise genome provides a

resource for the conservation of a threatened species. PloS One. 12:e0177708.

Tollis M., Hutchins E.D., Stapley J., Rupp S.M., Eckalbar W.L., Maayan I., Lasku E., Infante

C.R., Dennis S.R., Robertson J.A., May C.M., Crusoe M.R., Bermingham E., Denardo

D.F., Hsieh S.T., Kulathinal R.J., McMillan W.O., Menke D.B., Pratt S.C., Rawls J.A.,

Sanjur O., Wilson Rawls J., Wilson Sayres M.A., Fisher R.E., Kusumi K. 2018.

Comparative Genomics Reveals Accelerated Evolution in Conserved Pathways during

the Diversification of Anole Lizards. Genome Biol. Evol. 10:489–506.

Townsend J.P., López-Giráldez F, Friedman R. 2008. The phylogenetic informativeness of

nucleotide and amino acid sequences for reconstructing the vertebrate tree. J. Mol. Evol.

67:437-447.

Uetz P., Freed P., Aguilar R., Hošek J. 2021. The Reptile Database. (last accessed August 2021)

Wang Z., Pascual-Anaya J., Zadissa A., Li W., Niimura Y., Wang J., Huang Z., Li C., White S.,

Xiong Z., Fang D., Wang B., Ming Y., Chen Y., Zheng Y., Kuraku S., Pignatelli M.,

Herrero J., Beal K., Nozawa M., Li Q., Wang J., Zhang H., Yu L., Shigenobu S., Wang

J., Liu J., Flicek P., Searle S., Wang J., Kuratani S., Yin Y., Aken B., Zhang G., Irie N.

2013. The draft genomes of soft-shell turtle and green sea turtle yield insights into the

development and evolution of the turtle-specific body plan. Nat. Genet. 45:701–706.

Waterhouse R.M., Seppey M., Simão F.A., Manni M., Ioannidis P., Klioutchnikov G.,

Kriventseva E.V., Zdobnov E.M. 2018. BUSCO Applications from Quality Assessments

to Gene Prediction and Phylogenomics. Mol. Biol. Evol. 35:543–548.

Wolf Y.I., Rogozin I.B., Grishin N.V., Koonin E.V. 2002. Genome trees and the tree of life.

Trends Genet. TIG. 18:472–479.

Zdobnov E.M., Tegenfeldt F., Kuznetsov D., Waterhouse R.M., Simão F.A., Ioannidis P.,

Seppey M., Loetscher A., Kriventseva E.V. 2017. OrthoDB v9.1: cataloging evolutionary

and functional annotations for animal, fungal, plant, archaeal, bacterial and viral

orthologs. Nucleic Acids Res. 45:D744–D749.

Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree

reconstruction from partially resolved gene trees. BMC Bioinformatics. 19:153.