Propionate ameliorates diabetes-induced neurological dysfunction through regulating the
PI3K/Akt/eNOS signaling pathway

Qin Wu[2†], Jiajun Dong[2†], Yahong Cheng[1]*, Gaofeng Jiang[3]*

[1]Key Laboratory of Combinatorial Biosynthesis and Drug Discovery, Ministry of Education, Wuhan University

School of Pharmaceutical Sciences, Wuhan University, Wuhan, Hubei, P. R. China

[2]Zhejiang University-University of Edinburgh Institute, Zhejiang University, Haining, Zhejiang, P. R. China

[3]Center for Translational Medicine, Tianyou Hospital, Wuhan University of Science and Technology, Wuhan,

Hubei, P. R. China

☐ Corresponding authors

E-mail address: yahongcheng1106@163.com (Yahong Cheng); jianggf@wust.edu.cn (Gaofeng Jiang);

Phone: +86-18627009067

[†] These authors contributed to the work equally and should be regarded as co-first authors.

Abstract

A large body of research has established diabetes-related cognitive deterioration, sometimes
known as "diabetic encephalopathy." Current evidence supports that oxidative stress, neuronal
apoptosis, and cerebral microcirculation weakness are associated with cognition deficits induced
by diabetes. The present study explores the effect of propionate on neurological deficits, cerebral
blood flow, and oxidative stress in diabetic mice. Propionate can markedly improve neurological
function, which is correlated with its capabilities of stimulating nitric oxide (NO) production,
increasing cerebral microcirculation, suppressing oxidative stress and reducing neuron loss in the
hippocampus. In addition, the results of Western Blotting indicated that the brain-protective
function of propionate in streptozocin (STZ)-induced insulin-dependent diabetes mellitus (IDDM)
mice is related to phosphoinositide 3-kinase (PI3K)/serine-threonine protein kinase
(Akt)/endothelial nitric oxide synthase(eNOS) signaling pathway. In a diabetic mouse model,
propionate reduces cerebral microcirculation, hippocampus apoptosis, and neurological
impairment. Thus, propionate, now employed as a food preservative, may also help slow
diabetes-induced cognitive loss.

Highlights

● Propionate ameliorates diabetic-induced depression-like behavior, spatial learning and
memory deficits.

● Propionate increases cerebral microcirculation by enhancing NO production in diabetic mice.

● Propionate suppresses oxidative stress and reduces neuron loss in the hippocampus.

● The brain-protective function of propionate is related to PI3K/Akt/eNOS signaling pathway.

Keywords: Propionate; IDDM; Cerebral microcirculation; Neurological dysfunction;
PI3K/Akt/eNOS

Abbreviations: **IDDM,** insulin-dependent diabetes mellitus**; DM**, diabetes mellitus; **STZ**,
streptozocin; **Met**, metformin; **Pro**, propionate; **NO,** nitric oxide; **PYY,** peptide YY; **GLP-1,**
glucagon like peptide-1; **ROS**, reactive oxygen species; **SOD**, superoxide dismutase; **CAT**,
catalase; **MDA**, malondialdehyde; **ROI**, region-of-interest; **LCBF**, local cerebral blood flow;
ELISA, enzyme-linked immunosorbent assay; **PI3K**, phosphoinositide 3-kinase; **Akt**,
serine-threonine protein kinase; **eNOS**, endothelial nitric oxide synthase; **PDK1**, 3-inositol
phosphate-dependent protein kinase-1; **TST**, tail suspension test; **FST**, force swimming test;

MWM, Morris water maze; **FBG**, fasting blood glucose; **IPGTT**, intraperitoneal glucose tolerance test; **SDS-PAGE**, sodium dodecyl sulfate-polyacrylamide gel electrophoresis; **PVDF**, polyvinylidene fluoride; **Ctrl**, control; **ANOVA**, analysis of variance; **SD**, standard deviation; **AUC**, area under the curve.

1. Introduction

Cognitive disorder is very common among diabetic patients. Clinical research shows that both type 1 and type 2 diabetes patients are always accompanied by varying degrees of neurological dysfunction [1, 2], the incidence of which is related to the course of the disease, the selection of hypoglycemic drugs and other factors [3]. Slight cognitive impairment may occur in mild cases, and dementia may develop in severe cases [4]. Therefore, to find effective drugs for the prevention and treatment of diabetic learning nerve function impairment is a research hotspot in this field. Dietary supplementation with nutrients that protect brain tissue from damage may be a promising treatment approach for neurological complications caused by diabetes.

Propionic acid is a kind of short-chain fatty acid (SCFA), which is usually added to cheese, bread, animal feed and artificial spices as a preservative [5], and the concentration of propionic acid added as a food additive ranges from 1.0-2.5g/kg. It is reported that propionate plays an important biological role by acting as an antioxidant, anti-inflammatory, antiviral, anti-cancer, etc [6-8]. Some studies have shown that propionate can inhibit lipid synthesis in rat hepatocytes [9]. When the concentration of propionic acid was 0.1mmol/L, the synthesis of 50% cholesterol and triglyceride was inhibited [10]. In addition to reducing cholesterol and fat production, propionate may also be involved in weight control by suppressing appetite [11]. Propionate enhanced the release of PYY and GLP-1 from human colonic cells considerably in a human-based study. Postprandial plasma PYY and GLP-1 were considerably elevated after ingesting 10 grams of inulin-propionate ester acutely. The enteric release of PYY and GLP-1 creates senses of satiation and satiety in the appetite centers of the brain. Besides, it lessened energy uptake [11].

In addition to its role in peripheral nerves, more and more evidence indicates that propionic acid may play an important role in the central nervous system [11]. Propionic acid was found to enter the brain from the bloodstream in a concentration-dependent manner [12]. Propionic acid is catalyzed to propionyl coenzyme A (CoA) by thiokinase. Then propionyl CoA reacts with carbon dioxide to generate methylmalonyl CoA under the catalysis of propionyl CoA carboxylase. Finally, methylmalonyl CoA is catalyzed to succinyl CoA by the mutase, thus participating in the tricarboxylic acid cycle and ATP synthesis, which provides energy for the brain [13].

However, the therapeutic effect and underlying mechanisms of propionate on treating diabetes-related neurological complications have not been studied. Although the pathologic mechanisms of diabetes-induced nervous system impairment are not fully understood, an increasing number of studies have illustrated that decreased angiogenesis induced by oxidative stress is associated with neurological dysfunction [14-16]. Oxidative stress is the essential pathologic factor in the early phase [17]. Oxygen radical damages endothelial cells and induce their apoptosis, which greatly restricts the angiogenesis in brain tissue and limits the blood supply to the brain [18-20]. It also impacts the secretion of neurotrophic factors by endothelial cells, which exhibit a repairing effect to the function of impaired neurons [21]. Therefore, it will be a novel and effective therapeutic strategy for treating diabetic neurological complications to prevent oxidative stress injury and thus protect endothelial cells and promote angiogenesis.

Excitatory amino acid toxicity, immune-inflammatory reaction, free radicals and nitric oxide
damage are the main factors leading to neuron apoptosis [22]. Recently, the PI3K/Akt signaling
pathway has attracted much attention due to its regulatory role in neuronal apoptosis [23, 24].
Chemoattractant receptors are known to activate PI3K, a signaling enzyme. When PI3K is
activated, it produces the secondary messenger plasma membrane intrinsic protein 3, which
interacts with the signaling proteins AKT and phosphoinositide dependent kinase-1 (PDK1). AKT
is activated as a result of the combination causing PDK1 to phosphorylate Ser308 in AKT. It is
well known that AKT is a significant PI3K effector. [25]. Phosphorylation of Akt and nitric oxide
synthetase is an essential pathway of NO synthesis in endothelial cells [26-28]. However, whether the
intervention of propionate is capable of influencing PI3K/Akt/eNOS signaling pathway is unclear.
The effects of propionate in animal models with diabetic-induced neuronal dysfunction need
to be explored. Therefore, the present study focuses on evaluating the effects of propionate on
blood glucose level and neuronal function in streptozocin (STZ)-induced diabetic mice. The
effects of propionate on cerebral blood flow, oxidative stress, neuronal apoptosis, and
PI3K/Akt/eNOS signaling pathway were evaluated at the cellular or protein level through basic
pathological tests and biochemical tests enzyme-linked immunosorbent assays, and western blot.
This study provides not only a theoretical and experimental foundation on propionate effects of
alleviating diabetes-related neurological complications, but also a novel economical, effective and
side-effect-free candidate compound for the treatment of diabetic patients with a neurological
disorder.

2. Materials and methods
2.1. Reagents and Chemicals
Sinopharm Chemical Reagents Co., Ltd supplied sodium propionate (> 99 percent purity) (Wuhan,
China). Sigma-Aldrich provided the streptozotocin (STZ) and metformin (St. Louis, Mo, USA).
STZ was dissolved in citrate buffer, while metformin and sodium propionate were dissolved in
distilled water.
2.2. Animals
Sixty male Kunming adult mice weighted 18 to 22 grams were used. Every five mice shared one
cage and all mice had free access to food and water, under a 12-hour light/12-hour dark condition.
Mice were cared strictly compliant with the guidance of the National Institutes of Health Guide
for the Care and Use for Laboratory Animals and all animal experiments were performed with the
approval of the Institutional Animal Care and Used Committee of Wuhan University (AUP no.
S2020110121). Mice were randomly divided into 6 groups, 10 mice for each group. Mice in the
normal group (N=10) were not intervened, and 50 successfully modelled mice were randomly
divided into five groups of ten mice each: model group, STZ + propionate (37.5 mg/kg), STZ +
propionate (75 mg/kg), STZ + propionate (150 mg/kg) and STZ + metformin (100 mg/kg) (positive
drug control). The drug intervention groups mentioned above were given high, middle, low dosages
of propionate or metformin, respectively. Animals were administered treatments in a randomized,
double-blind design, and treatments were randomized across cages to avoid cage level effects.
2.3. Diabetes experimentally induced and treatment schedule
STZ (35mg/kg, i.p.) produced in citrate buffer (pH 4.4, 0.1M) was used to develop diabetes,
whereas control mice received an identical volume of citrate buffer. After seven days of STZ
injection, blood glucose levels were evaluated using blood glucose test strips. Mice with fasting
plasma glucose levels greater than 11.1mmol/L were chosen for the subsequent study. Once

diabetes was proven, mice were given either sodium propionate (37.5, 75, or 150 mg/kg), metformin (100 mg/kg), or vehicle (0.1 ml/10 g distilled water) orally once daily (9:00 a.m.) for 36 days. Fig. 1 displays the experiment's timeline.

Propionate levels in foods such as bread range between 1.0 and 2.5 g per kilogram. The dose of propionate used in the present study was determined based on the dose applied as a food preservative. The dosing range for propionate was determined based on our preliminary tests, which demonstrated that propionate was ineffective at doses less than 37.5 mg/kg. However, animals would have difficulty taking propionate at doses more than 75 mg/kg. Thus, we established three dosage groups: high, moderate, and low in order to assess the anti-epileptic effect of the three dosage groups.

2.4. Blood glucose and glucose tolerance test

Before the experiment, mice were fasted for 16 hours. Blood was obtained by drawing blood from a vein which is 2 mm from the distal end of the tail and baseline glucose levels were assessed using a glucometer [29]. Mice were intraperitoneally injected with 100 mg/ml glucose solution in sterile water at 2 g/kg body weight. The blood was taken to measure glucose levels at 0, 30, 60, 120 minutes after the sugar was given [30].

2.5. Body weight recording

All the experimental mice were weighed at 14:00 to 15:00 once per 3 days throughout the experiment.

2.6. Nissl staining

To evaluate the change of neuronal numbers, Nissl staining was performed. Mice were euthanized (N=6 per group) and perfused with 0.9% NaCl solution and 4% paraformaldehyde solution for brain fixation. The entire brain was then isolated and further fixated in paraformaldehyde for more than 24 hours. The brain was embedded in paraffin and sliced into 5 μm thick coronal sections. Paraffin sections were dewaxed and rehydrated before being stained with toluidine blue. Three sections were randomly selected from each mouse, and the stained sections were observed and photographed under a light microscope (Nikon, Tokyo, Japan) in the CA3 region of the hippocampus area. The number of neurons in this region was counted at high magnification (200×) [31].

2.7. Cerebral blood flow measurement

A moorFLPI-2 Full-Field Laser Perfusion Imager (Moor Instruments, Axminster, UK) was utilized to measure local cerebral blood flow. Anesthesia was induced with 4% chloral hydrate, following which mice (N=5 per group) were fixed on a stereotaxic frame plate with a constant temperature of 37□. An incision of skin was made to expose the skull surface, where ultrasound gel was applied. local cerebral blood flow (LCBF) was conducted for 5 min, 25 Hz for sampling frequency and 1 s per frame. The data images were analyzed with moorFLPI2 Full-Field Laser Perfusion Imager Review v5.0 software. For each animal, brain region-of-interest (ROI) was drawn, and ROI flux data was obtained from 19 images per scan. The mean value was used for subsequent analysis. All the analysis was blinded to experimental conditions [32].

2.8. Determination of oxidative stress-related indicators

2.8.1. Measurement of catalase (*CAT*), superoxide dismutase *(SOD) and* malondialdehyde (*MDA*)

Following completion of the behavioral studies, mice (n = 4) were killed. The tissue was subsequently removed from the brain's hippocampus region. These tissues (10%, w/v) were immediately homogenized in a freezing potassium phosphate buffer (pH 7.4) and centrifuged at

3000 rpm for 10 minutes at 4 °C. The supernatants obtained were collected and subjected to
biochemical analysis.
Protein content was quantified with a commercially available BCA protein assay kit (Solarbio
Science and Technology Co., Beijing, China). Malondialdehyde (MDA) level, as a marker of lipid
peroxidation, was assessed as contents of thiobarbituric acid-reactive substances (TBARS) by
spectrophotometry; the level of superoxide dismutase (SOD) was determined by xanthine oxidase
method as the index of the antioxidant system (Nanjing Jiancheng Bioengineering Institute).
Based on the operation instruction, catalase (CAT) determination was performed by using a
commercial CAT assay kit (Nanjing Jiancheng Bioengineering Institute) [33].
2.8.2. Detection of reactive oxygen species (ROS) level
Mouse ROS enzyme-linked immunosorbent assay (ELISA) kit was used to determine ROS
concentration. The hippocampal tissues of mice (N=5 per group) were extracted, homogenized,
centrifuged, and the supernatant was collected. 40µl supernatant was added to 96-well plate, and
reagents were applied according to the manufacturer's instruction. The absorbance of the mixture
at 450 nm was determined by a microplate analyzer. The standard curve was obtained by plotting
the concentration of the standard solution with the corresponding absorbance. The sample
concentration was determined based on the absorbance of the sample to be tested and expressed as
U/mgprot.
2.9. Assessment of NO level
NO levels were determined using the supernatant of hippocampal homogenates. NO level was
assessed for the purpose of evaluating endothelial cell function in the brain vasculature of diabetic
mice. In the body or in an aqueous solution, NO is extremely easy to be oxidized into nitrite ion
(NO_2^-). Low acidic pH value promotes the reaction of NO_2^- with diazosalt amine sulfonate to form
diazo compound, which further couples with naphthalene vinyl diamine to form a product is
characterized by an absorption peak of 550 nm. The level of NO can be determined by measuring
the absorption value. The NO concentration in brain tissue was determined using a NO assay kit
and following the instruction of the manufacture's (Caozhiyuan Biotechnology Co., Ltd.).
2.10. Western blotting
Western blotting was performed following a standard protocol. 20 µg of protein sample from the
hippocampus per lane was separated through sodium dodecyl sulfate-polyacrylamide gel
electrophoresis (SDS-PAGE). The proteins in the gel were subsequently transferred onto
polyvinylidene fluoride (PVDF) membranes and then immunological visualized. Primary
antibodies of anti-PI3K (1:1000), anti-p-PI3K (1:5000), anti-Akt (1:1000), and anti-p-Akt (1:5000)
were used. β-actin (1:5000) was used as an internal reference. The membrane was incubated
overnight at 4 □, and the chemiluminescence signal of interest was visualized using an imager
(BIO-RAD Gel Doc XR, America) [34].
2.11. Functional tests
Previous studies indicated that STZ induction resulted in the appearance of behavioral
abnormalities such as depression and motor deficits in TIDM mice. In this study, animals were
rotated through each test on different days. The test sequence remained constant to guarantee that
all animals were subjected to similar test circumstances. All tests were conducted in a blinded
fashion, with observers unaware of the treatments administered to mice.
2.11.1. Open field test
A black open-field apparatus (50 × 50 × 40 cm, with 25 equal squares) made of lacrylic box was

used to assess the exploratory willingness and spontaneous locomotor activity. The center of the
arenas was defined as the central nine squares. Mice (N=10 per group) were placed individually
into the center area and allowed 5-min free behave. Animal behavior was recorded using a video
tracking system and was analyzed automatically using OFT-100 software. Total distance traveled,
the time spent in the center and quiescent, and rearing times were calculated. The apparatus was
cleaned with 70% alcohol and dried between trials [35].

2.11.2. Tail suspension test

A tail suspension test (TST) was performed to evaluate the depression index of animals by
observing the time of immobility and agitation [36]. The instrument for this test is a grey acrylic box
(54 cm height) with an open front and a metallic hook at the top. We use laboratory sticky tape to
hang mice (N=10 per group) by the tails from the hook. Their movements were captured for 6
minutes via a camera complete with tracking software. The total immobility period and climbing
times were calculated by using the event count mode of RWD software. Longer immobility time
was an indicator of depression [37].

2.11.3. Forced swimming test

The FST was performed for the purpose of determining behavior related to depression [38]. Each
mouse was placed alone in a vertical plexiglass cylinder (140 mm in diameter, 200 mm high) that
contained water at a depth of 150 mm and a temperature of 23-25°C, for a period of 8 min.
Recorded for the same period, a blinded experimenter scored each mouse's immobility time for
the last 6 min. Immobility was recognized when mice were floating and moving a minimal amount
to keep its head and body from becoming submerged. The researcher remained in the room for the
duration of the test, and the water was changed between the introduction of each new mouse to
prevent odors (N=10 per group) [39].

2.11.4 Elevated plus maze test

The elevated plus-maze test is to investigate the anxiety state of animals by using the
contradictory behaviors of exploration and the fear of the high position [40]. The elevated plus-maze
has a pair of open arms (28.5×7cm) and two closed arms (28.5×7×14 cm) they are separated by a
central platform (7cm×7cm). The apparatus is 50 cm high. Each mouse was placed on the central
platform. The number of entries is recorded, and the time spent in the open arm is recorded over a
5 min time session (N=10 per group).

2.11.5. Morris water maze (MWM) task

MWM was performed to test the spatial learning and memory of the animals [41]. The basic device
of the water maze is a circular pool with a diameter of 1.2 meters and a height of 0.6 meters. The
water inside is colored milky white with non-fat milk powder. The water temperature in the pool
was maintained at 25±1□.

The pool was evenly divided into four quadrants (called south, east, north, west). An invisible
platform with a diameter of 8 cm was hidden 1 cm below the water surface in the target quadrant.
The whole test contains 2 phases, navigation test and probe test. During the navigation phase,
mice (N=10 per group) were released into the pool from randomly assigned release points, one
trial lasted 60 s at most, and would be guided to the platform beyond the time limit and allowed to
stay for 15 s. Four trials were run for the MWM navigation phase every day and these were
conducted on days 1 to 4. On day 5, the platform was withdrawn from the pool and probing tests
were conducted.   During the probe phase, the platform was removed from the pool, mice were
released for 1 min to swim freely. Tracks of mice in the pool were recorded with the Smart3.0

software during the training and probe phase. There were also 60-second intervals used to record
and analyze time spent in the target quadrant (the initial platform quadrant), number of platform
crossings, and travel velocity. [42].

2.12. Statistical analysis

All statistics were analyzed using SPSS statistics version 25, and graphs were generated by Prism
6.0 (GraphPad Software, Inc.). Data for the glucose tolerance test (GTT), body weight and escape
latencies in the MWM test were analyzed with a two-way analysis of variance (ANOVA). The
other outcomes' measurement data were compared using one-way analysis of variance (ANOVA).
The t-test was used to examine the significance of variable comparisons between groups.
Arithmetic mean ± standard deviation (SD) was used to indicate statistical differences, where p
less 0.05 was considered statistically significant.

3. Results

3.1. Reduction of fasting blood glucose and improvement of glucose intolerance by propionate
treatment

After 16 hours of fasting, diabetic mice's fasting blood glucose (FBG) climbed considerably.
($p<0.001$). However, propionate (75 mg/kg) and metformin (100 mg/kg) treatments significantly
decreased FBG ($p<0.001$) and there was no significant difference among the low dose, middle
dose and STZ groups (Fig. 2A). In the intraperitoneal glucose tolerance test (IPGTT), the area
under the curve (AUC) of blood glucose was calculated and found to be considerably higher in the
STZ group than in the Ctrl group ($p<0.001$) (Fig. 2B-C; Table 1). Comparing with STZ
administrated group, propionate administration reduced the AUC ($p<0.05$ in 37.5 and 150 mg/kg
groups; $p<0.001$ in 75 mg/kg group), and treatment of 75 mg/kg propionate has significant
advantages over 37.5 mg/kg and 150 mg/kg of propionate gavage groups. The data clearly showed
administration of propionate at a dose of 75 mg/kg effectively decreased the FBG and improved
glucose intolerance in IDDM mice. In the remaining experiments, propionate at a dose of 75
mg/kg was selected for intervention.

3.2. Propionate ameliorated STZ-induced body weight loss

The control group exhibited a gradual increase in body weight from 19.9±2.50g to 43.8±3.25g.
(Fig. 3). Meanwhile, the streptozotocin-induced mice exhibited a significant decrease in body
weight from 20.3±3.27g to 33.1±4.28g ($p<0.001$) in comparison to the former group. Conversely,
when compared to the model group, the animals who received propionate ($p<0.001$) or metformin
($p<0.01$) exhibited a significant enhancement in their body weight, from 20.3±2.81g to 40.2±2.94g
and from 19.2±1.28g to 37.4±1.62g, respectively. The results showed that streptozotocin
significantly reduced body weight in the animals compared to the control group, while propionate
therapy increased body weight, reversing the weight loss caused by streptozotocin.

3.3. Improvement of local cerebral blood flow (LCBF) by propionate intervention

We determined the effects of propionate and metformin on the LCBF of mice since it can provide
objective indicators for cerebral circulation or obstruction. LCBF was measured by Doppler
flowmetry (Fig. 4). The STZ-administered group exhibited a compromised LCBF level with
approximately 35.11% decrement ($p<0.001$). The pre-treatment of propionate significantly
increased LCBF comparing with the model group ($p<0.01$), while metformin treatment failed to
reverse LCBF decrement caused by STZ. These observations indicated that propionate could
rescue decreased LCBF levels induced by diabetes.

3.4. Oxidative stress partially ameliorated by pretreated propionate

3.4.1. Propionate enhanced CAT and SOD levels
The effect of propionate on antioxidase was further studied by measuring CAT and SOD activities.
As figure 5A showed, CAT activity was significantly reduced after STZ intervention when
compared to the non-treated group ($p<0.01$). While CAT activity was significantly increased after
propionate treatments compared with the STZ group ($p<0.01$). Furthermore, CAT in the
propionate treated group was found to be higher than in metformin-treated group ($p<0.05$). In
addition, SOD activity was obviously decreased in STZ-exposed mice in comparison to the
control group ($p<0.01$). The activity of SOD was restored in mice following propionate or
metformin treatment ($p<0.05$), and no substantial difference between the metformin and
propionate groups was found (Fig. 5B).
3.4.2. Inhibition of MDA and reactive oxygen species (ROS) accumulation by propionate
To ascertained the potential anti-oxidative effect of propionate, we also measured MDA (a marker
of lipid peroxidation) and ROS content in mice hippocampal tissue. The MDA contents in mice
hippocampus are shown in Fig. 5C. MDA formation was detected to be significantly elevated in
the STZ group compared to the control group ($p<0.001$). In contrast, significant decreases in MDA
levels were found in propionate or metformin treatment groups comparing to the STZ group
($p<0.01$). No statistical difference was presented between propionate and metformin groups.
The effects of increased production of ROS involve brain damage. ROS can produce membrane
damage, change inner proteins, induce lipids denaturation, and damage DNA. ROS concentration
was detected in mice brain tissues. Mice administrated by STZ showed significantly higher ROS
levels than the control group ($p<0.01$) (Fig. 5D). ROS concentration was remarkably reversed by
propionate intaking compared with the STZ group ($p<0.01$). While metformin treatments showed
little benefit effects. Propionate application had a distinct advantage over metformin
administration in reducing ROS concentration ($p<0.05$). In summary, propionate showed positive
effects in scavenging excessive reactive oxygen levels in mice brains.
3.5. Propionate enhanced ATP levels
The model group's ATP level was substantially lower than the control group's ($p < 0.05$). It was
significantly elevated following propionate pretreatment in comparison to the STZ-induced group.
However, metformin had no discernible effect on the level of ATP (Fig. 6A).
3.6. Enhancement of nitric oxide (NO) release by propionate therapy
We examed brain NO level to evaluate NO production in brain vasculature. NO concentration was
significantly lower in the STZ group ($p<0.001$): less than two-thirds that of the control group (Fig.
6B). Pre-treatment of propionate or metformin to the STZ-exposed mice raised NO levels by 59.2%
and 31.5%, respectively compared with the STZ group ($p<0.001$ and $p<0.01$, respectively). The
NO level in the propionate group was also elevated in comparison with that in the metformin
group ($p<0.05$). The increased NO concentration indicated an effect of propionate to promote
angiogenesis in the brain, with a significantly higher efficacy than metformin.
Propionate can significantly increase the activity of antioxidant enzymes and reduce the level of
malonaldehyde as an antioxidant. Although NO is also a product from oxidative stress, it has
protective effect on brain in this study. The experimental results showed that the main role of NO
is to dilate blood vessels, protect vascular endothelium and increase blood supply to the brain. It
can be seen that the protective effect of NO is greater than its oxidative damage. The experimental
results obtained and presented are reasonable.
3.6. Administration of propionate rescues STZ-induced neuronal loss in the hippocampus

Thirty-six days after STZ administration, we utilized Nissl staining to evaluate the structures of
the hippocampus in each group through comparing the number of neurons in optical fields of the
CA3 regions across different groups. STZ administration induced obvious lesions in neurons in
the CA3 and caused significant neuronal degeneration ($p<0.001$). On the contrary, metformin and
propionate rescued the degradation as demonstrated by the results that numbers of neurons in STZ
+ propionate (Pro) and STZ + metformin (Met) groups are significantly higher than that in the
STZ group ($p<0.001$). Noteworthily, propionate exhibited a superior effect than metformin
($p<0.01$). The data indicated that pre-treatment of propionate could persistently alleviate neuronal
necrosis resulted from STZ-induction.

3.7. Improvement of exploratory and locomotor activity following propionate administration

It is well documented that exploratory willingness and spontaneous locomotor activity can be
assessed by the open field tests via measuring distance moved, number of rearing, and time
duration in the center and immobility time. Here, we firstly found that propionate administration
was able to reverse reduced distance traveled, number of rearing, and time spent at the center of
the open field ($p<0.05$) caused by STZ treatments ($p<0.05$) (Fig. 8A, B and C). No statistical
significance was presented on these three indexes between propionate and metformin groups. The
immobility time was shown in Fig. 8D. Mice in the STZ group spent more immobility time than
those in the control group ($p<0.001$). Nevertheless, propionate reversed this result ($p<0.001$).
Notably, propionate-treated mice presented remarkably less immobility time than
metformin-administrated mice ($p<0.05$). Therefore, the results from open field tests suggested that
propionate could enhance the exploratory willingness of STZ-induced diabetic mice.

3.8. Antidepressant effects of propionate treatment

Next, these results were further confined in an elevated plus-maze, which is wildly utilized to
investigate the anxiety state of animals. As illustrated in Fig. 8, mice treated with STZ were less
inclined to enter or remain in open arms ($p<0.01$ and $p<0.05$, respectively). However,
propionate-treated or metformin-treated mice entered open arms more frequently than those in the
STZ group (Fig. 9A, $p<0.05$). In addition, the retention time in open arms was significantly longer
in the propionate-treated group than STZ model group ($p<0.05$). Interestingly, mice in the
propionate group tarried for a longer period in open arms than mice in the metformin group
($p<0.05$). By contrast, metformin-treated mice showed no significant difference in time in open
arms with those in the STZ group (Fig. 9B). Thus, the results in open arm tests indicated that
propionate treatment attenuated anxiety in the STZ-induced IDDM model.

Moreover, the effects of propionate on ameliorating mice's depression-like behaviors were
estimated in tail suspension tests and forced swimming tests. As shown in Fig. 10A, STZ-induced
diabetic mice presented significantly longer immobility time when suspended by tails ($p<0.05$).
Both propionate and metformin diminished the immobility time ($p<0.05$ and $p<0.01$, respectively).
STZ exposure reduced the intention of tail climbs (Fig. 10B, $p<0.001$), while propionate or
metformin treatments significantly reversed these tendencies ($p<0.001$ and $p<0.05$, respectively).
Interestingly, propionate showed a better effect in increasing tail climbing behaviors than
metformin ($p<0.05$). Furthermore, mice in the STZ-treated model group showed a remarkable
increase in immobility time than mice in the control group in forced swimming tests (Fig. 10C,
$p<0.05$), which was significantly reduced with the presence of propionate or metformin
administrations ($p<0.05$), although the immobility time in STZ + Pro group was still longer than
STZ + Met group ($p<0.05$). Therefore, the results implied that the administration of propionate

could ameliorate the depression-related status of mice induced by STZ induction.

3.9. Improvement of learning and memory by propionate

Spatial memory functions of mice were commonly evaluated by water maze tests. As shown in
figure 11A and Table 2, we found a significantly longer escape latency of the STZ group than the
control group on day 2 and day 3-4 ($p<0.05$ and $p<0.001$, respectively), although not on day 1.
Compared with the STZ model group, propionate treatment significantly reduced the escape
latency of mice on day 2-4 ($p<0.01$ on day 2 and 3; $p<0.001$ on day 4). Metformin also decreased
the latency on day 2 ($p<0.001$) and day 4 ($p<0.05$) but the propionate group showed a significantly
shorter latency time than the metformin group in the last two days ($p<0.05$ for day 3 and $p<0.01$
for day 4). In summary, propionate and metformin could positively alter spatial learning of
diabetic mice, while propionate was found to have better efficiency.

The number of crossing into the former location of the platform was also recorded (Fig. 12A). The
STZ treated group showed a significant decrease in crossing number in comparison with the
control group ($p<0.05$). Both metformin treatment and propionate treatment significantly restored
the crossing number ($p<0.05$). Another indicator, time spent in the target quadrant in probe task,
was significantly lower in STZ treated group in comparison to the control group ($p<0.01$) (Fig.
12B). While both the propionate-treated mice and metformin-treated mice spent a significantly
longer time in the target quadrant than the STZ group ($p<0.001$ and $p<0.01$, respectively). Besides,
the time that propionate-treated mice spent was longer than the metformin-treated group ($p<0.05$).
As for the swimming velocity, there had been a falling-off in STZ treated group ($p<0.05$) (Fig.
12C). In contrast, the propionate-treated mice were faster ($p<0.05$), while the metformin-treated
mice showed no significant acceleration compared to the propionate pretreated mice.

3.10. Potent activation of the PI3K/Akt/eNOS pathway by propionate

Propionate activates the PI3K/Akt/eNOS pathway in the hippocampus of diabetic mice. Western
blotting was implemented in this study for the purpose of identifying the relative expression of
phosphorylated PI3K, Akt, and eNOS. From the panel, the protein levels of PI3K, Akt and eNOS
remained unaltered across all 4 groups. While, the STZ significantly reduced the
phosphorylation levels of PI3K, Akt and eNOS. Both metformin and propionate presented rescue
effects on phosphorylation level of PI3K, Akt and eNOS comparing with the STZ group ($p<0.001$
for PI3K and Akt; $p<0.05$ for eNOS). Propionate had exserted stronger effects on restoring
phosphorylation levels of PI3K and Akt from the comparison between the propionate group and
metformin group ($p<0.05$).

Discussion

Diabetes mellitus is the most common chronic metabolic syndrome is characterized by absolute or
relative insulin deficiency. The major symptom is chronic hyperglycemia caused by disturbance of
saccharides, lipid, and protein metabolism [43]. According to an epidemiological survey in 2015,
approximately 8% of the global population aged between 20 to 79 suffers from diabetes, and the
proportion is anticipated to rise to 9.9%. Abundant studies consistently suggest that diabetes
mellitus (DM) affects the nervous system. Due to the prolonged life expectancy of DM patients,
cognitive impairment and dementia have become new striking complications of DM [44].
DM-induced cognitive deficits are also known as diabetic encephalopathy [45]. Presently, many
studies on diabetic encephalopathy worldwide focus on peripheral neuropathy, while the
pathophysiological mechanisms of neuropathy caused by diabetic encephalopathy are less
explored. However, with the significant increase in the life expectancy of DM patients, the

prevalence of diabetic cognitive dysfunction is bound to increase. Thus, illuminating the
pathogenesis of diabetic cognitive dysfunction and seeking effective protective or therapeutic
strategies are urgently required.

This study contributes to this research field by discovering the therapeutic effect of propionate on
IDDM, which has not been discussed previously. Firstly, we noticed that propionate intervention
decreased blood glucose levels in STZ-induced type I diabetic mice, reversed their weight loss,
increased their local cerebral blood flow, and indirectly improved their neural function. Secondly,
propionate showed good antioxidant activity and could alleviate the loss of neurons in the
hippocampus in IDDM mice. Moreover, we discovered that the neural protective function of
propionate might be associated with the modulation of NO biosynthesis and improved blood
supply to brain tissues. Furthermore, we found that the brain-protective function of propionate in
STZ-induced IDDM mice was related or at least partly related to PI3K/Akt/eNOS signaling
pathway.

STZ is a broad-spectrum antibiotic. It has properties of antibacterial, antitumor, and causing
diabetes, and it has high-selective toxicity of pancreatic β-cells in experimental animals. Its
activity in inducing diabetes may be through free radical damage to islet β-cells, which impairs the
insulin-producing function of islet β-cells and leads to diabetes. Intraperitoneal injection of STZ
has recently been applied in many diabetes-related studies to specifically destroy pancreatic
β-cells, resulting in secretion disorder and persistent hyperglycemia without toxicity to brain cells
[46]. In this study, after a single intraperitoneal injection of STZ at 55 mg/kg to mice, their body
weight was significantly decreased, and the fasting blood glucose level could reach up to or more
than 20 mmol/L. Mice treated with STZ were generally in poor conditions, and some of them died
before the terminal of the study.

Even in pediatric patients, metformin is the most commonly given treatment for hyperglycemia. [47,
48], It belongs to the biguanide family of drugs and works to lower blood sugar levels by reducing
hepatic gluconeogenesis and enhancing insulin sensitivity. While metformin has been used to treat
diabetes for more than 60 years, new evidence suggests that it could be used to treat a variety of
other diseases, including cancer, cardiovascular disorders, neurodegenerative diseases [49] and more
recently, intellectual disability-related neurodevelopmental diseases [50, 51]. Metformin's mechanism
of action is not totally understood. Metformin's activity is clearly focused on energy metabolism,
and accumulating data shows that its therapeutic potential may be derived from its ability to target
mitochondrial abnormalities. [52]. In fact, chronic metformin administration enhances mitochondrial
activity and antioxidant protection. Consistent with metformin's propensity to cross the
blood-brain barrier, these therapeutic effects have been observed in both the peripheral and central
nervous systems. [53]. Thus, metformin was used as a positive control drug in this study.

Insufficient insulin secretion leads to hyperglycemia. Persistent high blood glucose may cause
thickening of capillary basement membrane and narrowing of vessel lumen [54]. In combination
with increased blood viscosity and slowed blood flow caused by metabolic disturbance of
saccharides and lipids in diabetic patients, the structure changes of capillaries would decrease
cerebral blood flow [55]. Some research has proved that the decrease of cerebral blood flow could
cause dysfunction of the brain in information processing, manipulating, and integrating [56]. It
would also impact cognitive response and processing ability, and eventually lead to impairment in

learning and memory. In this study, cerebral blood flow was measured, besides the body weight
and blood glucose level of mice. According to the results, in STZ-induced diabetic mice, their
blood glucose level was markedly increased, and body weight was significantly decrease related to
the control mice 36 days after STZ induction. As expected, results from cerebral blood flow
studies also demonstrated obvious lower blood flow in high-glucose mice (36 days after STZ
administration) than control mice. In contrast, after sodium propionate treatment, the blood
glucose level of STZ-induced IDDM mice was significantly reduced. The sodium propionate
exhibited the best effect on controlling blood glucose with a dose of 75 mg/kg. The weight gain of
mice in the sodium propionate group was improved compared with the simple model group.
Moreover, the cerebral blood flow of the diabetic mice was significantly increased after 36 days
compared with the model mice, as showed by results from cerebral blood flow measurement.
Studies have found that long-term hyperglycemia is closely related to diabetic neurological
complications, and diabetic models have exhibited varying degrees of behavioral and cognitive
dysfunction [57]. In this study, the behavior of experimental mice was evaluated by animal ethology
methods, in order to study the improvement brought by propionate in neural function. The
open-field test was used to evaluate the activity and exploration desire of mice, the tail suspension
test and forced swimming test was used to evaluate the depression-like behaviors, and the water
maze test was used to detect spatial learning and memory ability of mice. Experimental results
showed that long-term hyperglycemia-induced significant cognitive and behavioral impairments,
which are concretely showed in depression, a decline of learning and memory, and the results
were consistent with that of previous studies about cognitive and behavioral impairments in
diabetic models [58].
It has been reported that high glucose leads to excessive production of reactive oxygen species
(ROS) in vascular endothelial cells [59]. Excessive ROS formation affects the hypoxia signaling
pathway and further aggravates vascular endothelial disorders [60]. It causes damage to lipids,
proteins and mitochondrial DNA on the mitochondrial intima of neurons, resulting in intracellular
energy failures and diabetic brain injury. At the same time, oxidative stress will result in metabolic
disorders. The results of this study showed that the hippocampus CA3 area of the experimental
diabetes model mice was significantly damaged, and the neuron loss was severe. The SOD and
CAT activities in the brain tissue were significantly decreased, while the MDA content increased.
The preventive intervention of propionate can significantly ameliorate the morphological changes
of brain tissue and reduce the loss of neurons in the hippocampus, and antioxidant effects were
also significantly enhanced. These findings indicate that propionate has a significant anti-oxidative
function and can relieve hyperglycemia-induced brain damage in mice while promoting the
regeneration of neurons.
Studies have shown that vascular endothelial cells can secrete a variety of vasoactive substances,
including vasodilator factor NO. As a previous study suggested, NO synthesis is inhibited due to
impaired vascular endothelial function in diabetic patients. Hyperglycemic environment leads to a
reduction in endothelial cell function and reduction in the bioavailability of NO. In addition, high
blood sugar culture conditions will accelerate the senescence of vascular endothelial cells, leading
to impaired proliferation [61]. This study found that the NO level of model mice was significantly
reduced, and propionate treatment significantly increased the NO level of mice. It is suggested that
the therapeutic effect of propionate may be related to the regulation of NO synthesis and secretion.
Despite the fact that NO is a byproduct of oxidative stress, it was found to have a protective

impact on the brain in this study. The findings of the experiments revealed that the primary function of NO is to widen blood vessels, protect vascular endothelium, and boost blood supply to the brain. It may be observed that the preventive impact of NO outweighs its oxidative harm.

Other researchers have found that inhibition of the PI3K/Akt/eNOS signaling pathway can engender vascular endothelial cell dysfunction, and a decrease in NO synthesis is observed when eNOS activity decreases since the eNOS is mainly activated by the PI3K/Akt signaling pathway. In this study, propionate treatment was shown to significantly upregulate the expression levels of PI3K/Akt/eNOS signaling pathway-related proteins, suggesting that propionate's brain protection function in STZ-induced type I diabetic mice is associated with PI3K/Akt/eNOS signaling pathway activation.

Our findings demonstrated that propionate (75 mg/kg) has a beneficial effect on the injured brains of mice. As a result, approximately 0.5 g propionate should be administered to a 60 kg individual via dose conversion from mice to human. Propionate levels in bread range between 1.0 and 2.5 g per kilogram. We believe that the recommended dosage of propionate is easily achieved from food supply. Supplements in the diet, such as propionate, appear to be promise for safeguarding the brain against harm in IDDM. However, given the difference between humans and mice, further rigorous studies examining the safety and efficacy of such high doses of propionate in different animal models of epilepsy are required prior to initiating clinical trials.

Conclusion

Taken together, strong evidence has been provided that propionate acts to ameliorate cerebral microcirculation to improve the cognitive abilities in diabetic mice, which may be attributed to its anti-diabetic, antioxidant and modulated by PI3K/Akt/eNOS pathway. The results of this study emphasize the importance of propionate on NO release, cerebral circulation, oxidative stress, and PI3K/Akt/eNOS pathway against high glucose insult.

Funding source

This study did not get any funding from funding agencies

Conflict of interest

The authors acknowledge no conflict of interest pertaining to the publication of the manuscript.

Acknowledgements

Thanks to Shiqi Fan and Jinjin Li for their help in preparing samples and in measuring biochemical indices.

Author contributions

Yahong Cheng: Conceptualization, Writing-original draft and Writing-review & editing.

Yuan Jiang and Xinying Bai: Data curation and Methodology.

Qin Wu and Jiajun Dong: Conceptualization, Software, Data curation and analysis, Methodology, Writing-original draft, drawing graphic abstract, and revising the manuscript.

Gaofeng Jiang: Supervision, Validation and Methodology.

References

1.	Moheet A, Mangia S, Seaquist ER. Impact of diabetes on cognitive function and brain structure. Ann N Y Acad Sci. 2015;1353: 60-71.

2.	Biessels GJ, Staekenborg S, Brunner E, Brayne C, Scheltens P. Risk of dementia in diabetes mellitus: a systematic review. The Lancet Neurology. 2006;5(1): 64-74.

3.	Barbara G, Mégarbane B, Argaud L, et al. Functional outcome of patients with prolonged hypoglycemic encephalopathy. Ann Intensive Care. 2017;7(1): 54.

4.	Albai O, Frandes M, Timar R, Roman D, Timar B. Risk factors for developing dementia in type 2 diabetes mellitus patients with mild cognitive impairment. Neuropsychiatr Dis Treat. 2019;15: 167-175.

5.	Yun SS, Kim J, Lee SJ, et al. Naturally occurring benzoic, sorbic, and propionic acid in vegetables. Food Addit Contam Part B Surveill. 2019;12(3): 167-174.

6.	Filippone A, Lanza M, Campolo M, et al. The Anti-Inflammatory and Antioxidant Effects of Sodium Propionate. Int J Mol Sci. 2020;21(8).

7.	Luu M, Weigand K, Wedi F, et al. Regulation of the effector function of CD8(+) T cells by gut microbiota-derived metabolite butyrate. Sci Rep. 2018;8(1): 14430.

8.	Nurdin SU, Le Leu RK, Young GP, Stangoulis JC, Christophersen CT, Abbott CA. Analysis of the Anti-Cancer Effects of Cincau Extract (Premna oblongifolia Merr) and Other Types of Non-Digestible Fibre Using Faecal Fermentation Supernatants and Caco-2 Cells as a Model of the Human Colon. Nutrients. 2017;9(4).

9.	Hu Y, Chen D, Yu B, et al. Effects of dietary fibres on gut microbial metabolites and liver lipid metabolism in growing pigs. J Anim Physiol Anim Nutr (Berl). 2020;104(5): 1484-1493.

10.	Park S, Kang J, Choi S, et al. Cholesterol-lowering effect of Lactobacillus rhamnosus BFE5264 and its influence on the gut microbiome and propionate level in a murine model. PLoS One. 2018;13(8): e0203150.

11.	Chambers ES, Viardot A, Psichas A, et al. Effects of targeted delivery of propionate to the human colon on appetite regulation, body weight maintenance and adiposity in overweight adults. Gut. 2015;64(11): 1744-1754.

12.	Ostendorf F, Metzdorf J, Gold R, Haghikia A, Tönges L. Propionic Acid and Fasudil as Treatment Against Rotenone Toxicity in an In Vitro Model of Parkinson's Disease. Molecules. 2020;25(11).

13.	Bui D, Ravasz D, Chinopoulos C. The Effect of 2-Ketobutyrate on Mitochondrial Substrate-Level Phosphorylation. Neurochem Res. 2019;44(10): 2301-2306.

14.	Liyanagamage D, Martinus RD. Role of Mitochondrial Stress Protein HSP60 in Diabetes-Induced Neuroinflammation. Mediators Inflamm. 2020;2020: 8073516.

15.	Kim KA, Shin YJ, Kim JH, et al. Dysfunction of endothelial progenitor cells under diabetic conditions and its underlying mechanisms. Arch Pharm Res. 2012;35(2): 223-234.

16.	Dhalla NS, Shah AK, Tappia PS. Role of Oxidative Stress in Metabolic and Subcellular Abnormalities in Diabetic Cardiomyopathy. Int J Mol Sci. 2020;21(7).

17.	Nan W, Zhonghang X, Keyan C, Tongtong L, Wanshu G, Zhongxin X. Epigallocatechin-3-Gallate Reduces Neuronal Apoptosis in Rats after Middle Cerebral Artery Occlusion Injury via PI3K/AKT/eNOS Signaling Pathway. Biomed Res Int. 2018;2018: 6473580.

18.	Yang J. The role of reactive oxygen species in angiogenesis and preventing tissue injury after brain ischemia. Microvasc Res. 2019;123: 62-67.

19.   Yang Y, Torbey MT. Angiogenesis and Blood-Brain Barrier Permeability in Vascular
Remodeling after Stroke. Curr Neuropharmacol. 2020;18(12): 1250-1265.
20.   Jambusaria A, Hong Z, Zhang L, et al. Endothelial heterogeneity across distinct vascular beds
during homeostasis and inflammation. Elife. 2020;9.
21.   Wen C, Huang C, Yang M, et al. The Secretion from Bone Marrow Mesenchymal Stem Cells
Pretreated with Berberine Rescues Neurons with Oxidative Damage Through Activation of the
Keap1-Nrf2-HO-1 Signaling Pathway. Neurotox Res. 2020;38(1): 59-73.
22.   Calcerrada P, Peluffo G, Radi R. Nitric oxide-derived oxidants with a focus on peroxynitrite:
molecular targets, cellular responses and therapeutic implications. Current pharmaceutical design.
2011;17(35): 3905-3932.
23.   Tang JF, Wen Q, Sun J, Zhang WM, Zhu HL. Advances in the researches on the biological
activities and inhibitors of phosphatidylinositol 3-kinase. Anti-cancer agents in medicinal
chemistry. 2014;14(5): 673-687.
24.   Yang F, Zhang Y, Tang Z, Shan Y, Wu X, Liu H. Hemin treatment protects neonatal rats from
sevoflurane-induced neurotoxicity via the phosphoinositide 3-kinase/Akt pathway. Life sciences.
2020;242: 117151.
25.   Dangelmaier C, Manne BK, Liverani E, Jin J, Bray P, Kunapuli SP. PDK1 selectively
phosphorylates Thr(308) on Akt and contributes to human platelet functional responses. Thromb
Haemost. 2014;111(3): 508-517.
26.   Xu L, Wang S, Li B, Sun A, Zou Y, Ge J. A protective role of ciglitazone in ox-LDL-induced
rat microvascular endothelial cells via modulating PPARγ-dependent AMPK/eNOS pathway. J
Cell Mol Med. 2015;19(1): 92-102.
27.   Ishii M, Nakahara T, Ikeuchi S, Nishimura M. β-Amyrin induces angiogenesis in vascular
endothelial cells through the Akt/endothelial nitric oxide synthase signaling pathway. Biochem
Biophys Res Commun. 2015;467(4): 676-682.
28.   Lee JH, Parveen A, Do MH, Lim Y, Shim SH, Kim SY. Lespedeza cuneata protects the
endothelial dysfunction via eNOS phosphorylation of PI3K/Akt signaling pathway in HUVECs.
Phytomedicine. 2018;48: 1-9.
29.   Liu X, Bai X, Li M, Li H, Zhang Y, Yang B. Detecting Establishment of Shared Blood
Supply in Parabiotic Mice by Caudal Vein Glucose Injection. J Vis Exp. 2020(156).
30.   Dinger K, Mohr J, Vohlen C, et al. Intraperitoneal Glucose Tolerance Test, Measurement of
Lung Function, and Fixation of the Lung to Study the Impact of Obesity and Impaired Metabolism
on Pulmonary Outcomes. J Vis Exp. 2018(133).
31.   Wang R, Weng G, Yu S, Dai S, Zhang W, Zhu F. Diffusion-weighted imaging detects early
brain injury after hypothermic circulatory arrest in pigs. Interactive cardiovascular and thoracic
surgery. 2018;26(4): 687-692.
32.   An Y, Kang Y, Lee J, Ahn C, Kwon K, Choi C. Blood flow characteristics of diabetic patients
with complications detected by optical measurement. Biomed Eng Online. 2018;17(1): 25.
33.   Ahangarpour A, Oroojan AA, Khorsandi L, Kouchak M, Badavi M. Solid Lipid
Nanoparticles of Myricitrin Have Antioxidant and Antidiabetic Effects on
Streptozotocin-Nicotinamide-Induced Diabetic Model and Myotube Cell of Male Mouse.
Oxidative medicine and cellular longevity. 2018;2018: 7496936.
34.   Zeng Y, Wang X, Yin B, et al. Role of the stromal cell derived factor-1/CXC chemokine
receptor 4 axis in the invasion and metastasis of lung cancer and mechanism. Journal of thoracic

disease. 2017;9(12): 4947-4959.

35. Sayapina NV, Sergievich AA, Kuznetsov VL, et al. Influence of multi-walled carbon nanotubes on the cognitive abilities of Wistar rats. Experimental and therapeutic medicine. 2016;12(3): 1311-1318.

36. Cryan JF, Mombereau C. In search of a depressed mouse: utility of models for studying depression-related behavior in genetically modified mice. Mol Psychiatry. 2004;9(4): 326-357.

37. Ide S, Takahashi T, Takamatsu Y, et al. Distinct Roles of Opioid and Dopamine Systems in Lateral Hypothalamic Intracranial Self-Stimulation. The international journal of neuropsychopharmacology. 2017;20(5): 403-409.

38. Yankelevitch-Yahav R, Franko M, Huly A, Doron R. The forced swim test as a model of depressive-like behavior. J Vis Exp. 2015(97).

39. Stieglitz MS, Fenske S, Hammelmann V, et al. Disturbed Processing of Contextual Information in HCN3 Channel Deficient Mice. Frontiers in molecular neuroscience. 2017;10: 436.

40. Walf AA, Frye CA. The use of the elevated plus maze as an assay of anxiety-related behavior in rodents. Nat Protoc. 2007;2(2): 322-328.

41. Vorhees CV, Williams MT. Morris water maze: procedures for assessing spatial and related forms of learning and memory. Nat Protoc. 2006;1(2): 848-858.

42. Arika WM, Kibiti CM, Njagi JM, Ngugi MP. Modulation of Cognition: The Role of Gnidia glauca on Spatial Learning and Memory Retention in High-Fat Diet-Induced Obese Rats. Neural plasticity. 2019;2019: 2867058.

43. Poznyak A, Grechko AV, Poggio P, Myasoedova VA, Alfieri V, Orekhov AN. The Diabetes Mellitus-Atherosclerosis Connection: The Role of Lipid and Glucose Metabolism and Chronic Inflammation. Int J Mol Sci. 2020;21(5).

44. Umegaki H. Therapeutic Potential of Antidiabetic Medications in the Treatment of Cognitive Dysfunction and Dementia. Drugs & aging. 2016;33(6): 399-409.

45. Wang S, Zhou SL, Min FY, et al. mTOR-mediated hyperphosphorylation of tau in the hippocampus is involved in cognitive deficits in streptozotocin-induced diabetic mice. Metabolic brain disease. 2014;29(3): 729-736.

46. Kim J, Shin SH, Kang JK, Kim JW. HX-1171 attenuates pancreatic β-cell apoptosis and hyperglycemia-mediated oxidative stress via Nrf2 activation in streptozotocin-induced diabetic model. Oncotarget. 2018;9(36): 24260-24271.

47. Brufani C, Fintini D, Nobili V, Patera PI, Cappa M, Brufani M. Use of metformin in pediatric age. Pediatr Diabetes. 2011;12(6): 580-588.

48. Thomas I, Gregg B. Metformin; a review of its history and future: from lilac to longevity. Pediatr Diabetes. 2017;18(1): 10-16.

49. Lv Z, Guo Y. Metformin and Its Benefits for Various Diseases. Front Endocrinol (Lausanne). 2020;11: 191.

50. Gantois I, Khoutorsky A, Popic J, et al. Metformin ameliorates core deficits in a mouse model of fragile X syndrome. Nat Med. 2017;23(6): 674-677.

51. Greger HK, Hanem LGE, Østgård HF, Vanky E. Cognitive function in metformin exposed children, born to mothers with PCOS - follow-up of an RCT. BMC Pediatr. 2020;20(1): 60.

52. Soukas AA, Hao H, Wu L. Metformin as Anti-Aging Therapy: Is It for Everyone? Trends Endocrinol Metab. 2019;30(10): 745-755.

53. Huang YC, Hsu CC, Lin WC, et al. Effects of metformin on the cerebral metabolic changes

in type 2 diabetic patients. ScientificWorldJournal. 2014;2014: 694326.

54. Wimmer RA, Leopoldi A, Aichinger M, et al. Human blood vessel organoids as a model of
diabetic vasculopathy. Nature. 2019;565(7740): 505-510.

55. Huang JY, Li LT, Wang H, et al. In vivo two-photon fluorescence microscopy reveals
disturbed cerebral capillary blood flow and increased susceptibility to ischemic insults in diabetic
mice. CNS Neurosci Ther. 2014;20(9): 816-822.

56. Ogoh S. Relationship between cognitive function and regulation of cerebral blood flow. The
journal of physiological sciences : JPS. 2017;67(3): 345-351.

57. Zhang S, Xue R, Hu R. The neuroprotective effect and action mechanism of polyphenols in
diabetes mellitus-related cognitive dysfunction. European journal of nutrition. 2020;59(4):
1295-1311.

58. Bagatini PB, Xavier LL, Bertoldi K, et al. An evaluation of aversive memory and
hippocampal oxidative status in streptozotocin-induced diabetic rats treated with resveratrol.
Neuroscience letters. 2017;636: 184-189.

59. Dymkowska D, Drabarek B, Podszywałow-Bartnicka P, Szczepanowska J, Zabłocki K.
Hyperglycaemia modifies energy metabolism and reactive oxygen species formation in
endothelial cells in vitro. Archives of biochemistry and biophysics. 2014;542: 7-13.

60. Karki P, Birukov KG. Rho and Reactive Oxygen Species at Crossroads of Endothelial
Permeability and Inflammation. Antioxidants & redox signaling. 2019;31(13): 1009-1022.

61. Kang H, Ma X, Liu J, Fan Y, Deng X. High glucose-induced endothelial progenitor cell
dysfunction. Diabetes & vascular disease research. 2017;14(5): 381-394.

Tables

Table 1

Blood glucose in the IPGTT test.

Data was expressed as mean ± SD (N = 10). ###$p < 0.001$ vs. Ctrl group, *$p < 0.05$, **$p < 0.01$, and

***$p < 0.001$ vs. STZ group. The two-way ANOVA for IPGTT indicated a significant

group-by-min interaction.

Table 2

The exact escape latency in morris water maze values in seconds.

Data was expressed as mean ± SD (N = 10). #$p < 0.05$, ##$p < 0.01$, and ###$p < 0.001$ vs. Ctrl group,

*$p < 0.05$, **$p < 0.01$, and ***$p < 0.001$ vs. STZ group, $^{\$}p < 0.05$ and $^{\$\$}p < 0.01$ vs. STZ + Met

group. The two-way ANOVA for escape latency indicated a significant group-by-day interaction.

Fig. 1 The study design is depicted graphically, encompassing the delivery of drugs, neurobehavioral assessment, histological assessment, biochemical analysis, and protein expression processes.

Fig. 2. Effects of propionate on FBG and IPGTT. (A) FBG levels; (B) The AUC of blood glucose in the IPGTT test; (C) IPGTT test 36 days after STZ induction. The data are expressed as the mean $\pm$ SD (N = 10). ###$p < 0.001$ vs Ctrl group; *$p < 0.05$, **$p < 0.01$ and ***$p < 0.001$ vs STZ group. Data for the IPGTT were analyzed with a two-way analysis of variance (ANOVA).

Fig. 3. The body weight data of the untreated control group, the streptozotocin exposure group, and the groups treated with propionate and metformin. All data were expressed as mean $\pm$ SD (N = 10). ###$p < 0.001$ vs Ctrl group; **$p < 0.01$ and ***$p < 0.001$ vs STZ group. Data for the body weight were analyzed with a two-way analysis of variance (ANOVA).

Fig. 4. The effect of propionate on LCBF in mice cerebral cortex. (A) Representative images of LCBF in different groups. The magnitude of LCBF is represented by different colors, with blue to red indicating low to high. (B) Quantitative analysis of LCBF in all groups. Data were expressed as mean $\pm$ SD (n = 6). ###$p < 0.001$ vs. Ctrl group, **$p < 0.01$ vs. STZ group, n.s., no significant difference.

Fig. 5. Levels of oxidative stress. (A and B) STZ decreased CAT and SOD. Pretreatment with propionate attenuated these decrements. (C and D) STZ increased MDA and ROS levels in mice brains. Pretreatment with propionate attenuated these increments. Data were expressed as mean $\pm$ SD (n = 6). ###$p < 0.001$ and ##$p < 0.01$ vs. Ctrl group, *$p < 0.05$ and **$p < 0.01$ vs. STZ group, $^{\$}p < 0.05$ vs. STZ + Met group, n.s., no significant difference.

Fig. 6. Effects of propionate on ATP level and NO release in brain tissues. Data were expressed as mean $\pm$ SD (N = 6). #$p < 0.05$ and ###$p < 0.001$ vs. Ctrl group, *$p < 0.05$, **$p < 0.01$ and ***$p < 0.001$ vs. STZ group, $^{\$}p < 0.05$ vs. STZ + Met group. n.s., no significant difference.

Fig. 7. Cellular structures in the hippocampus are visible after Nissl staining. (A-D) Representation images of Ctrl, STZ, STZ + Pro, and STZ + Met, 36 days after IDDM induction. (E) Quantitative of neurons in the CA3. The red arrows indicate normal neurons, while the black arrows indicate necrosis neurons. Scale bar = 100 μm. Data were expressed as mean $\pm$ SD (N = 3). ###$p < 0.001$ vs. Ctrl group, ***$p < 0.001$ vs. STZ group, $^{\$\$}p < 0.01$ vs. STZ + Met group.

Fig. 8. Animal behavior analysis in the open field test. In the OFT, the travel distance (A), number of standing (B), time spent at the center (C) and immobility time (D) were recorded and analyzed. Data were expressed as mean $\pm$ SD (N = 10). #$p < 0.05$ and ###$p < 0.001$ vs. Ctrl group, *$p < 0.05$ and ***$p < 0.001$ vs. STZ group, $^{\$}p < 0.05$ vs. STZ + Met group, n.s., no significant difference.

Fig. 9. Results of elevated plus maze test. Open arms entry times (A) and retention time in open arms (B) were recorded and analyzed. Data were expressed as mean $\pm$ SD (N = 10). #$p < 0.05$ and ##$p < 0.01$ vs. Ctrl group, *$p < 0.05$ vs. STZ group, $^{\$}p < 0.05$ vs. STZ + Met group, n.s., no significant difference.

Fig. 10. The influence of propionate on STZ-induced depression-like behavior evaluated by tail suspension test (TST) and force swimming test (FST). Quiescent time in TST (A), number of climbing times (B), and immobility time in FST (C) were measured and analyzed. Data were expressed as mean $\pm$ SD (N = 10). #$p < 0.05$ and ###$p < 0.001$ vs. Ctrl group, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ vs. STZ group, $^{\$}p < 0.05$ vs. STZ + Met group, n.s., no significant difference.

Fig. 11. Escape latencies are shown as the mean value of each trail session for four sequential days

(A). Swimming tracks are recorded via a video tracking camera system (B). Data for the escape

latencies were analyzed with a two-way analysis of variance (ANOVA). Data was expressed as

mean ± SD (N = 10). #$p < 0.05$ vs. Ctrl group, *$p < 0.05$ and **$p < 0.01$ vs. STZ group, n.s., no

significant difference.

Fig. 12. Influence of propionate on STZ-induced cognition and memory dysfunction measured by

MWM test. Crossing into the former location of the platform (A), time spent in the target quadrant

in probe task (B) and Swimming speed (C) were recorded and analyzed. Data were expressed as

mean ± SD (N = 10). #$p < 0.05$ and ##$p < 0.01$ vs. Ctrl group, *$p < 0.05$, **$p < 0.01$, and ***$p <$

0.001 vs. STZ group, $^\$p < 0.05$ vs. STZ + Met group, n.s., no significant difference.

Fig. 13. Phosphorylation of PI3K/AKT/eNOS signaling pathway proteins, as measured by western

blotting. Representative western blot images (A) and the ratio of p-PI3K/PI3K (B), p-AKT/AKT

(C), and p-eNOS/eNOS. GAPDH served as an internal control. Data are presented as the mean ±

standard deviation (n=3 independent experiments). #$p < 0.05$ vs. Ctrl group, *$p < 0.05$ and ***$p <$

0.001 vs. STZ group, $^\$p < 0.05$ vs. STZ + Met group, n.s., no significant difference.

Table 1

Groups	Time (min)			
	0	30	60	120
Ctrl	5.9±0.74	8.5±0.93	7.6±1.09	5.4±0.63
STZ	21.7±3.35###	25.6±4.22###	22.7±4.45###	20.7±4.40###
STZ+Pro-37.5	17.8±5.14	19.7±4.59**	18.8±3.88	16.4+2.54*
STZ+Pro-75	11.5±3.37***	14.5±3.43***	12.5±3.21***	10.4±3.53***
STZ+Pro-150	18.7±4.09	20.2±4.28*	18.9±4.93	16.8±4.05
STZ+Met	12.4±2.57***	15.1±3.44***	11.7±2.42***	9.5±2.68***

791   Table 2

Groups	Escape latency (sec)			
	Day1	Day2	Day3	Day4
Ctrl	54.9±5.31	46.5±7.08	37.7±2.64	27.1±2.91
STZ	58.7±2.05	57.3±3.38$^{\#}$	51.0±5.48$^{\#\#}$	44.7±4.17$^{\#\#\#}$
STZ + Pro-75	52.8±7.87	45.7±4.43**	35.9±4.98**, $^{\$}$	25.7±2.59***, $^{\$\$}$
STZ + Met	53.6±7.59	48.8±3.11**	46.4±7.43	36.0±5.51*

793   Graphical Abstract

795   Fig. 1

797   Fig. 2

799   Fig. 3

801 Fig. 4

A

B

 Fig. 5

 Fig. 6

    Fig. 8

811   Fig. 9

    Fig. 10

815   Fig. 11

A

Escape latencies during 4 days of training

B

817   Fig. 12