Identification of binding interaction between *Curcumin derivative* and *PERK13 Proline rich receptor like protein kinase* protein using *In silico* docking techniques – to help plants tackle salt water

Prakash Vaithyanathan*

* Science Teacher, SACROSANCT SERVICES, Chennai 600041, India.

Corresponding Email: pvaithyanathan@gmail.com

Abstract

Introduction

Arabidopsis thaliana, mouse ear cress or thale cress are small flowering plants included in the cruciferae family. They comprise of various characteristics such as diploid genetics, small genome size, rapid growth cycle, and relatively low repetitive DNA content, making it a perfect model for plant genome projects.

Objective

The aim of this present *Insilico* research study is to carry out molecular drug docking studies between PERK13 Proline rich receptor like protein kinase and De O Acetylated Curcumin Di Galactose or di–galactosylated curcumin, a derivative of curcumin. PERK13 protein is considered to play a significant role in helping plants tackle salinity levels in water.

Methods

In this study, protein modelling tools and servers are used to model the 3D structure and the same is validated using Protein structure validation tools. Automated drug docking servers were used to dock the modelled protein with the chemical compound to analyse the electrostatic (H bond) interaction between PERK13 and De O Acetylated Curcumin Di Galactose, a better water soluble compound than curcumin. The docked structure was visualized using an advanced molecular visualization tool.

Results and Discussion

The overall results obtained from this study on De O Acetylated Curcumin Di Galactose and PERK13 protein shows that De O Acetylated Curcumin Di Galactose, directly binds with the active site and other potentially binding regions of PERK13. Hence, it is concluded that De O Acetylated Curcumin Di Galactose could potentially play a vital role in future research related to the problem of helping the plants tackle increased saline levels in water.

Keywords: *A. thaliana***,** De O Acetylated Curcumin Di Galactose, Drug Docking, *Insilico*, PERK13, Protein Modelling.

Introduction

Arabidopsis is an efficient model to carry out research. Since many years, *Arabidopsis* has been used to understand about gravitropism (Kiss *et al.,* 2000), genetics (Zheng 2006), and genomics (Brooks *et al.* 2011). Mutants can be obtained for reverse genetics projects from *Arabidopsis* Biological Resource Center (https://abrc.osu.edu/) or the Nottingham *Arabidopsis* Stock Centre (http://www.arabidopsis.info), which includes sequence indexed collections of above 30,000 homozygous T DNA insertional lines (O'Malley and Ecker 2010) from many thousands of insertion events (Alonso *et al.,* 2003)[6]. Full length complementary DNAs (Yamada *et al.,* 2003) can also be obtained for most genes from the *Arabidopsis* Biological Resource Center. Additionally, collections of *Arabidopsis* expression vectors (Curtis and Grossniklaus, 2003; Earley *et al.* 2006) and yeast two hybrid vectors (Trigg *et al.,* 2017) are available which can be made use of to conduct overexpression, localization, and interaction studies.

The Arabidopsis proline rich extensin like receptor kinase (PERK) family includes 15 predicted receptor kinases. A comprehensive analysis of expression was performed for identifying the overlapping and unique expression patterns within this family related to their phylogeny. PERK13 protein is believed to play a significant role to help plants tackle salinity levels in water.

Methodology

Protein target analysis: In this study, the gene coded protein sequence of **PERK13, (UniProt ID:** Q9CAL8) was retrieved from UniProt database (https://www.uniprot.org/uniprot/Q9CAL8) for performing sequence analysis and protein modelling. The sequence was converted into FASTA format. **Sequence analysis:** The retrieved protein was analysed using a UniProt Proteomics tool named MobiDB (https://mobidb.bio.unipd.it/) for identifying the functional domains present in PERK13. **Molecular modelling**: The selected amino acid sequence of the target protein (PERK13) was converted into 3D structure using an automated protein modelling server, SWISS MODEL (https://swissmodel.expasy.org/). The modelled structure was validated using ProCheck server (https://www.ebi.ac.uk/thornton srv/software/PROCHECK/) to analyse the Ramachandran Plot. **3D structure visualization**: The modelled protein was visualized in 3D

form using an advanced molecular visualization tool called Discovery Studio software in order to clearly view the intramolecular structural details of PERK13. **Molecular drug docking**: In this study, the drug compound De O Acetylated Curcumin Di Galactose was selected and was docked with the modelled PERK13 protein using an automated molecular modelling and drug docking server called PatchDock (https://bioinfo3d.cs.tau.ac.il/PatchDock/) for analysing the drug binding score and the interaction between De O Acetylated Curcumin Di Galactose and PERK13.

Results

```
>sp|Q9CAL8|PEK13_ARATH Proline rich receptor like protein kinase PERK13
OS=Arabidopsis thaliana OX=3702 GN=PERK13 PE=1 SV=1
MSDSPTSSPPAPSADSAPPPDTSSDGSAAPPPTDSAPPPSPPADSSPPPALPSLPPAVFS
PPPTVSSPPPPPLDSSPPPPPDLTPPPSSPPPPDAPPPIPIVFPPPIDSPPPESTNSPPP
PEVFEPPPPPADEDESPPAPPPPEQLPPPASSPQGGPKKPKKHHPGPATSPPAPSAPATS
PPAPPNAPPRNSSHALPPKSTAAGGPLTSPSRGVPSSGNSVPPPANSGGGYQGKTMAGFA
IAGFAVIALMAVVFLVRRKKKRNIDAYSDSQYLPPSNFSIKSDGFLYGQNPTKGYSGPGG
YNSQQQSNSGNSFGSQRGGGGYTRSGSAPDSAVMGSGQTHFTYEELTDITEGFSKHNILG
EGGFGCVYKGKLNDGKLVAVKQLKVGSGQGDREFKAEVEIISRVHHRHLVSLVGYCIADS
ERLLIYEYVPNQTLEHHLHGKGRPVLEWARRVRIAIGSAKGLAYLHEDCHPKIIHRDIKS
ANILLDDEFEAQVADFGLAKLNDSTQTHVSTRVMGTFGYLAPEYAQSGKLTDRSDVFSFG
VVLLELITGRKPVDQYQPLGEESLVEWARPLLHKAIETGDFSELVDRRLEKHYVENEVFR
MIETAAACVRHSGPKRPRMVQVVRALDSEGDMGDISNGNKVGQSSAYDSGQYNNDTMKFR
KMAFGFDDSSDSGMYSGDYSVQDSRKGSNGASSEFTRNETENRNFNNRRY
```

Fig. 1 FASTA sequence of PERK13

The above picture shows the amino acid sequence of PERK13 protein

Fig. 2 Sequence analysis of PERK13 using MobiDB Database

In the above picture, the coloured region indicates the various functional domains present in PERK13.

Fig. 3 Sequence analysis of PERK13 using MobiDB Database

In the above picture, the blue coloured region represents predominant functional domains of PERK13 (356 626aa).

Fig. 4 Protein modelling – SWISS MODEL server

The above picture is the Home Page of SWISS MODEL server.

Fig. 5 3D structure visualization of PERK13

The above image shows the 3D structure of PERK13 viewed in "Solid Ribbon Model" using Discovery Studio Software.

Fig. 6 3D structure visualization of PERK13

The above image shows the 3D structure of PERK13 viewed in "Cartoon Model" using Discovery Studio Software.

Fig 7 3D structure visualization of PERK13

Fig. 8 a) 3D structure validation ProCheck server

Plot statistics

Residues in most favoured regions [A,B,L]	222	87.1%
Residues in additional allowed regions [a,b,l,p]	29	11.4%
Residues in generously allowed regions [~a,~b,~l,~p]	3	1.2%
Residues in disallowed regions	1	0.4%
	----	------
Number of non-glycine and non-proline residues	255	100.0%
Number of end-residues (excl. Gly and Pro)	2	
Number of glycine residues (shown as triangles)	24	
Number of proline residues	9	

Total number of residues	290	

Based on an analysis of 118 structures of resolution of at least 2.0 Angstroms and R-factor no greater than 20%, a good quality model would be expected to have over 90% in the most favoured regions.

Fig. 8 b) Plot statistics ProCheck server

The above images show the assessment of Ramachandran Plot for the modelled PERK13 protein.

Receptor	Ligand		Complex Type	Cluste
C1.pdb	Q9CAL8_Protein.pdb		drug	4.0

Solution No	Score	Area	ACE
1	6318	756.90	-116.36
2	5784	797.30	-304.26
3	5742	796.60	-318.80
4	5698	844.00	-153.74
5	5570	1092.40	-431.21
6	5552	825.30	-234.33
7	5546	761.00	-179.79
8	5444	902.10	-217.50
9	5370	670.70	-124.91
10	5320	756.60	90.53
11	5298	653.30	-60.10
12	4946	667.00	-117.20
13	4872	627.10	11.82
14	4860	782.20	-270.83
15	4858	622.70	-184.08
16	4828	726.00	-252.64
17	4788	778.80	-235.84
18	4776	597.20	-143.28
19	4746	589.40	-248.52
20	4740	530.10	-165.91

Fig 9 Molecular drug docking – PatchDock results

The above picture is the result page of PatchDock server. It shows the Atomic Contact Energy (ACE) value of De O Acetylated Curcumin Di Galactose with PERK13 protein complex which is 431.21.

Fig. 10 Molecular drug docking (De O Acetylated Curcumin Di Galactose with PERK13)

The above picture represents the 3D form of the De O Acetylated Curcumin Di Galactose   PERK13 complex. The yellow coloured spacefill model represents De O

Acetylated Curcumin Di Galactose which binds with PERK13 and is viewed using Discovery Studio Software.

Fig. 11 Drug Protein binding sites prediction

(De O Acetylated Curcumin Di Galactose with PERK13)

The above picture represents the 3D form of the De O Acetylated Curcumin Di Galactose   PERK13 complex. Green coloured structure in Stick model represents De O Acetylated Curcumin Di Galactose which interacts with PERK13 at the amino acid binding regions viewed using Discovery Studio Software.

Fig. 12 Drug Protein binding sites prediction

(De O Acetylated Curcumin Di Galactose with PERK13)

The above picture represents the 3D form of the De O Acetylated Curcumin Di Galactose   PERK13 complex. Green coloured structure in Stick model represents De O

Acetylated Curcumin Di Galactose which interacts with PERK13 (Surface model) at the amino acid binding regions viewed using Discovery Studio Software.

```
<HBondMonitor>
:UNK0:O61 - A:VAL385:O
A:GLY388:N - A:GLY386:O
A:GLN389:NE2 - :UNK0:O10
A:LYS500:NZ - :UNK0:N4
A:LYS500:NZ - :UNK0:O10
A:THR505:OG1 - A:ALA499:O
A:HIS508:ND1 - :UNK0:N4
A:ARG512:N - A:VAL509:O
A:ARG512:NE - :UNK0:O16
A:ARG512:NH2 - :UNK0:O16
A:ARG512:NH2 - :UNK0:O58
A:ARG512:NH2 - A:GLY363:O
A:VAL513:N - :UNK0:O16
A:VAL513:N - :UNK0:O58
A:VAL513:N - A:VAL509:O
A:THR516:N - :UNK0:O26
A:THR516:N - :UNK0:O29
A:THR516:OG1 - :UNK0:O26
A:THR516:OG1 - :UNK0:O29
A:SER563:OG - :UNK0:O38
A:ARG569:NE - :UNK0:O50
A:ARG569:NE - :UNK0:O56
A:ARG569:NH1 - :UNK0:O50
A:ARG569:NH1 - :UNK0:O56
A:ARG569:NH2 - :UNK0:O56
```

Fig 12 H bond interaction of De O Acetylated Curcumin Di Galactose with PERK13

The above picture shows the Electrostatic and H bond interaction between De O Acetylated Curcumin Di Galactose and PERK13.

Table 1: Summary of Molecular Docking results

Target	**Compound :** *De O Acetylated Curcumin Di Galactose*
PERK13 **(UniProt ID:** Q9CAL8 **)**	431.21

The above table shows the ACE (Atomic Contact Energy) value when De O Acetylated Curcumin Di Galactose binds with PERK13.

Discussion

PERK13 protein was retrieved in FASTA format using UniProt database. The total length of the protein was 710 aa and Mass (Da):75,372 (Fig. 1). In first step involved of

protein analysis, we use UniProt Prosite scan database which clearly shows that the amino acid range 353 – 619 aa falls within the functional domains (protein kinases) of PERK13.

Protein kinases (Hunter, 1991; Hanks and Quinn, 1991; Hanks, 1991; Hanks et al., 1988) are enzymes that makeup a large family of proteins which share sequence homology. Their conserved catalytic core is common to serine/threonine as well as to tyrosine protein kinases. There are several conserved regions present in the catalytic domain of protein kinases. Two of these regions were selected to construct signature patterns. The first region, which can be spotted in the N terminal end of the catalytic domain, is a stretch of residues, which is rich in glycine, and which is present near a lysine residue. It has also been indicated to be concerned with ATP binding (Fig. 2).

In the second step, mobiDB Database (https://mobidb.bio.unipd.it/Q9CAL8) was used for domain analysis. Transmembrane (TM) α helices comprise of a few residues which are strongly hydrophilic when compared to the general composition of proteins (Marticorena et al., 1993; Jones et al., 1994, Arkin and Brunger, 1998; Tourasse and Li, 2000). This can be partly explained by the high loss of energy when polar side chains are buried in a hydrophobic environment (Engelman and Steitz, 1984). Polar residues are highly conserved, particularly in multi spanning TM proteins, which suggests molecular interactions that favor these residues either functionally or structurally (Gray and Matthews, 1984; Gromiha, 1999). Functional roles of certain polar residues are noticed in protein structures, such as binding of prosthetic groups (H and E) in the photosynthetic reaction center (Deisenhofer et al., 1985), proton transport (D) and retinal binding (K) in bacteriorhodopsin (Henderson et al., 1990; Luecke et al., 1999), binding of hemes (H) in cytochrome c oxidase (Iwata et al.,1995) and the cytochrome bc_1 complex, and Ca^{2+} binding (N, D, E, and T) in the Ca^{2+} ATPase (Toyoshima et al., 2000). The results obtained from MobiDB Database for PERK13 protein shows that another functional domain of PERK13 lies within 676 700 polar residues (Fig. 3).

In this research study, SWISS MODEL was used to convert the amino acid sequence of PERK13 into 3D structure [Fig (4)]. SWISS MODEL (Bienert et al., 2017; Guex et al., 2009; Studer et al., 2020; Bertoni et al., 2017; Laskowski et al., 1993) was used to elaborately analyse the molecular and structural details of PERK13 for the purpose of docking. SWISS MODEL is a server for automated comparative modelling of three dimensional (3D) protein structures. Waterhouse et al. computed models by the SWISS MODEL server homology modelling pipeline which relies on ProMod3, an in house comparative modelling engine

based on Open Structure. (Laskowski et al., 1996; Duhovny et al., 2002) [Fig. 5]. The modelled 3D protein was completely evaluated using ProCheck server for assessment of Ramachandran Plot. The results for PERK13 protein obtained from ProCheck server show a Ramachandran plot score of 87.1% which indicates that there is no error present in the modelled structure.

After modelling the 3D structure of PERK13, the protein was viewed using Discovery Studio Software [Fig (6, 7, and 8)]. Discovery Studio is a software which is currently used by the majority of Bioinformaticians and Cheminformaticians. It is used for simulating small molecule and macromolecule systems. It is developed and distributed by Dassault Systemes BIOVIA.

Molecular Drug Docking

The selected test compound, *De O Acetylated Curcumin Di Galactose*, was docked with PERK13 and the molecular H bond interaction between them was analysed. Proteins with small ligands are docked using Patchdock server (Duhovnyet al., 2005; Duhovny et al., 2003; Mashiach et al., 2010). Hence, docking, for our study, was performed using this server. In molecular docking studies, the compound, *De O Acetylated Curcumin Di Galactose* was docked with PERK13. H bond interactions between the drug and the compound were observed along with binding affinity scores. Table 1 shows the drug docking result with ACE value.

Fig.7 shows the PatchDock result scores of PERK13 with *De O Acetylated Curcumin Di Galactose* and the 3D structure of H bond interactions with the respective drug binding amino acid pockets (VAL:385, GLY:388, GLY:386, GLN:389, LYS:500, LYS:500, THR:505, ALA:499, HIS:508, ARG:512, VAL:509, ARG:512, ARG:512, ARG:512, ARG:512, GLY:363) (Fig 9,10,11,12).

The PatchDock results of *De O Acetylated Curcumin Di Galactose* with PERK13 show an ACE (Atomic Contact Energy) value of 432.19. According to theory, higher negative value indicates more binding affinities between the target and the compound. Here, it has been proved that *De O Acetylated Curcumin Di Galactose* is an ideal candidate to interact with PERK13.

Interestingly, the results obtained from docking shows that *De O Acetylated Curcumin Di Galactose* directly binds with the functional domain regions of PERK13 protein. The

functional domains were analysed using ScanProsite and MobiDB. The following amino acids shows the positions where the drug *De O Acetylated Curcumin Di Galactose* interacts with the PERK13 protein: "VAL: 385,GLY:388,GLY:386,GLN:389,LYS:500,LYS:500,THR:505,ALA:499,HIS:508,ARG:512,VAL:509,ARG:512,ARG:512,ARG:512,ARG:512,GLY:363. Database references mention that the Nucleotide binding regions of amino acids are from **359 367** positions (https://prosite.expasy.org/rule/PRU00159) and the Functional Domains residues of amino acids are from (https://mobidb.bio.unipd.it/Q9CAL8) **356 626.**

Conclusion

In this current *Insilico* drug docking research study, the selected compound, *De O Acetylated Curcumin Di Galactose* efficiently binds with the target protein, PERK13 (Proline rich receptor like protein kinase). The current docking study clearly proves that the Domains and the polar residue amino acid positions (Drug Binding Pockets) of PERK13 exhibit H bond interaction with *De O Acetylated Curcumin Di Galactose*. Hence, the selected compound efficiently down regulates PERK13 and can be used as an efficient antagonist agent in future studies to help plants tolerate saline levels.

Acknowledgement

Thanks to the late Sri V Balasubramanian for his help over the years and his son Mr. B V Narayanan (currently residing in U.K.) for funding this project. I would like to extend my thanks to Dr. Balaji Munivelan, Ph.D., Senior Bioinformatician at ABS Geno informatics, Chennai (bioinfobalaji@gmail.com, Mobile: +91 9444345906) to help me learn the basics of docking and for his support in the verification of *Insilico* Docking studies. I thank my student Dr. Harishchander Anandaram, Amrita Vishwa Vidyapeetham for all the basic lessons in bioinformatics.

References

1. Kiss, J. Z., Weise, S. E., Kiss, H. G. (2000). How can plants tell which way is up? *Am Biol Teach.*, *62*(1):59 63.
2. Zheng, Z. L. (2006). Use of the gl1 Mutant & the CA rop2 Transgenic Plants of Arabidopsis thaliana in the Biology Laboratory Course. *Am Biol Teach., 68*(9), e148 e153.

3. Brooks, E., Dolan, E., Tax, F. (2011). Partnership for Research & amp; Education in Plants (PREP): Involving High School Students in Authentic Research in Collaboration with Scientists. *Am Biol Teach., 73*(3): 10.1525/abt.2011.73.3.4.

4. O'Malley, R.C., Ecker, J. R. (2010). Linking genotype to phenotype using the Arabidopsis unimutant collection, *Plant J., 61*(6), 928 40.

5. *A*lonso, J. M., Stepanova, A. N., Leisse, T. J., et al., (2003). Genome wide insertional mutagenesis of *Arabidopsis thaliana*, *Science, 301*(5633), 653 7.

6. Yamada, K., Lim. J., Dale, J. M., et al., (2003). Empirical analysis of transcriptional activity in the Arabidopsis genome, *Science, 302*(5646), 842 6.

7. Curtis, M. D., Grossniklaus, U. (2003). A gateway cloning vector set for high throughput functional analysis of genes in plants. *Plant Physiol., 133*(2), 462 9.

8. Earley, K. W., Haag, J. R., Pontes, O., et al., (2006). Gateway compatible vectors for plant functional genomics and proteomics. *Plant J., 45*(4), 616 29.

9. Trigg, S. A., Garza, R. M., MacWilliams, A., et al., (2017). CrY2H seq: a massively multiplexed assay for deep coverage interactome mapping. *Nat Methods., 14*(8), 819 825.

10. Hanks, S. K., Hunter, T. (1995). Protein kinases 6. The eukaryotic protein kinase superfamily: kinase (catalytic) domain structure and classification. *FASEB J. 9,* 576 596.

11. Hunter, T. (1991). Protein kinase classification. *Methods Enzymol., 200*, 3 37.

12. Hanks, S. K., Quinn, A. M. (1991). Protein kinase catalytic domain sequence database: identification of conserved features of primary structure and classification of family members. *Methods Enzymol. 200*, 38 62.

13. Hanks, S. K. (1991). Eukaryotic protein kinases, Current Opinion in Structural Biology, *Curr. Opin. Struct. Biol.*, *1*(3), 369 383.

14. Hanks, S.K., Quinn, A. M., Hunter, T. (1988). The protein kinase family: conserved features and deduced phylogeny of the catalytic domains. *Science*, *241,* 42 52.

15. Landolt Marticorena, C., Williams, K. A., Deber, C. M. & Reithmeier, R. A. (1993). Non random distribution of amino acids in the transmembrane segments of human type I single span membrane proteins. *J. Mol. Biol.*, *229*(3), 602–608.

16. Jones, D. T., Taylor, W. R. & Thornton, J. M. (1994). A mutation data matrix for transmembrane proteins, *FEBS Letters*, *339*(3), 269–275.

17. Arkin, I. T. & Brunger, A. T. (1998). Statistical analysis of predicted transmembrane alpha helices. *Biochim. Biophys. Acta.*, *1429*(1), 113–128.

18. Tourasse, N. J. & Li, W. H. (2000). Selective constraints, amino acid composition, and the rate of protein evolution. *Mol. Biol. Evol*. 17, 656–664.

19. Engelman, D. M. & Steitz, T. A. (1984) in The Protein Folding Problem, ed. Wetlaufer, D. B. (Westview, Boulder, CO), pp. 87–113.

20. Gray, T. M. & Matthews, B. W. (1984). Intrahelical hydrogen bonding of serine, threonine and cysteine residues within alpha helices and its relevance to membrane bound proteins, *J. Mol. Biol.* 175(1), 75–81.

21. Gromiha, M. M. (1999). A simple method for predicting transmembrane α helices with better accuracy, *Protein Eng. 12*(7), 557–561.

22. Deisenhofer, J., Epp, O., Miki, K., Huber, R. & Michel, H. (1985). Structure of the protein subunits in the photosynthetic reaction centre of *Rhodopseudomonas viridis* at 3Å resolution. *Nature (London)*, 318, 618–624.

23. Henderson, R., Baldwin, J. M., Ceska, T. A., Zemlin, F., Beckmann, E. & Downing, K. H. (1990). Model for the structure of bacteriorhodopsin based on high resolution electron cryo microscopy. *J. Mol. Biol. 213*(4), 899–929.

24. Luecke, H., Schobert, B., Richter, H. T., Cartailler, J. P. & Lanyi, J. K. (1999). Structure of bacteriorhodopsin at 1.55 □ resolution. *J. Mol. Biol., 291*, 899–911.

25. Iwata, S., Ostermeier, C., Ludwig, B. & Michel, H. (1995). Structure at 2.8 A resolution of cytochrome c oxidase from *Paracoccus denitrificans*. Nature (London) *376*(6542), 660–669.

26. Toyoshima, C., Nakasako, M., Nomura, H. & Ogawa, H. (2000). Crystal structure of the calcium pump of *sarcoplasmic reticulum* at 2.6 A resolution. *Nature, 405*(6787), 647–655.

27. Waterhouse, A., Bertoni, M., Bienert, S., et al., (2018). SWISS MODEL: homology modelling of protein structures and complexes. *Nucleic Acids Res., 46*(W1), W296 W303.

28. Bienert, S., Waterhouse, A., de Beer, T. A. P., et al., (2017). The SWISS MODEL Repository new features and functionality. *Nucleic Acids Res.*, *45*(D1), D313 D319.

29. Guex, N., Peitsch, M. C., Schwede, T. (2009). Automated comparative protein structure modeling with SWISS MODEL and Swiss PdbViewer: A historical perspective. *Electrophoresis*, *30*(1), S162 S173.

30. Studer, G., Rempfer, C., Waterhouse, A.M., et al., (2020). QMEANDisCo distance constraints applied on model quality estimation. *Bioinformatics, 36*, 1765 1771.

31. Bertoni, M., Kiefer, F., Biasini, M., et al., (2017). Modeling protein quaternary structure of homo and hetero oligomers beyond binary interactions by homology. *Scientific Reports,* 7, 10480.

32. Laskowski, R. A., MacArthur, M. W., Moss, D. S., et al., (1993). PROCHECK a program to check the stereochemical quality of protein structures. *J. App. Cryst.*, 26, 283 291.

33. Laskowski, R. A., Rullmannn, J. A., MacArthur, M. W., et al., (1996). AQUA and PROCHECK NMR: programs for checking the quality of protein structures solved by NMR. *J Biomol NMR*, *8*, 477 486.

34. Duhovny, D., Nussinov, R., Wolfson, H. J. Efficient Unbound Docking of Rigid Molecules. In Gusfield et al., Ed. Proceedings of the 2'nd Workshop on Algorithms in Bioinformatics(WABI) Rome, Italy, Lecture Notes in Computer Science 2452, pp. 185 200, Springer Verlag, 2002.

35. Schneidman Duhovny, D., Inbar, Y., Nussinov, R., et al., (2005) PatchDock and SymmDock: servers for rigid and symmetric docking. *Nucl. Acids. Res.*, *33*, W363 367.

36. Mashiach, E., Schneidman Duhovny, D., Peri, A., et al., (2010). An integrated suite of fast docking algorithms. *Proteins*, *78* (15), 3197 204.

37. https://www.nature.com/articles/s41598 020 70921 2.

38. https://patents.google.com/patent/WO2017217508A1/en