Anti-TRAP/SSP2 monoclonal antibodies can inhibit sporozoite infection and enhance protection of anti-CSP monoclonal antibodies

Brandon K. Wilder*[1]†#, Vladimir Vigdorovich*[1], Sara Carbonetti[1], Nana Minkah[1], Nina Hertoghs[1], Andrew Raappana[1], Hayley Cardamone[1], Brian G. Oliver[1], Olesya Trakhimets[1], Sudhir Kumar[1], Nicholas Dambrauskas[1], Silvia A. Arredondo[1], Nelly Camargo[1], Stefan H.I. Kappe*[1,2,3] and D. Noah Sather*[1,2,3]

[1]Center for Global Infectious Disease Research, Seattle Children's Research Institute, Seattle, WA, USA

[2]Department of Pediatrics, University of Washington, Seattle, WA, USA

[3]Department of Global Health, University of Washington, Seattle, WA, USA

†Present address: Vaccine and Gene Therapy Institute, Oregon Health & Science University, Beaverton, Oregon, 97006, USA

*Equal contribution

Correspondence: D. Noah Sather: noah.sather@seattlechildrens.org and

Stefan H.I. Kappe: stefan.kappe@seattlechildrens.org

<u>Summary</u>

Vaccine-induced sterilizing protection from infection with the *Plasmodium* parasite, the

pathogen that causes malaria, will be an essential tool in the fight against malaria as it

would prevent both malaria-related disease and transmission. Stopping the relatively

small number of parasites injected by the mosquito before they can migrate from the

skin to the liver is an attractive goal. Antibody-eliciting vaccines have been used to

pursue this objective by targeting the major parasite surface protein present during this

stage, the circumsporozoite protein (CSP). While CSP-based vaccines have recently

had encouraging success in disease reduction, this was only achieved with extremely

high antibody titers and appeared less effective for a complete block of infection. While

such disease reduction is important, these results also indicate that further

improvements to vaccines based solely on CSP will likely yield diminishing benefits

towards the goal of durable, infection-blocking immunity. Here, we show that

monoclonal antibodies (mAbs) recognizing the sporozoite protein TRAP/SSP2 across

the major protein domains exhibit a range of inhibitory capacity and that these mAbs

can augment CSP-based protection despite delivering no sterile protection on their own.

Therefore, pursuing a multivalent subunit vaccine immunization is a promising strategy

for improving infection-blocking malaria vaccines.

Introduction

The last four years have marked a disheartening milestone as the first time in a

generation without a reduction in the global burden of malaria (WHO) [1]. The

interventions that have provided much of the previous progress, such as insecticide-

treated bednets and large scale treatment programs, are highly susceptible to

interruptions due to political or economic instability. This was starkly illustrated by the

resurgence of malaria in Venezuela after near-elimination and also as predicted

following the COVID-19 pandemic [1]. Therefore, it is likely that long-lasting, infection-

blocking interventions (e.g. vaccines, long-lasting monoclonal antibodies or

chemoprophylactics) will be required to drive malaria to elimination.

Developing such an intervention is hampered by the complex life cycle of the parasite,

which begins when an infected mosquito injects tens to hundreds of the "sporozoite"

forms of the parasite into the dermis [2]. From here, sporozoites actively traverse through

multiple cell types in search of an endothelial cell through which they will gain access to

the blood [3]. They are carried to the liver within minutes, where they traverse multiple cell

types in the liver parenchyma and eventually establish infection in a hepatocyte [4].

Following ~7–10 days of asexual replication and transformation (~2 days in rodent

malaria models), each successful liver stage releases 30,000–50,000 "merozoites" that

cyclically infect, replicate within and lyse red blood cells [5,6]. It is only during this blood

stage of infection when symptomatic disease occurs and is also where a subset of

sexually differentiated parasite forms can be picked up by a new mosquito host to

continue the transmission cycle. Each step in the infection cycle presents an opportunity

for intervention, although vaccines targeting the "pre-erythrocytic" stages in the skin and liver have yielded the most promising results [7] .

The most advanced pre-erythrocytic vaccine is RTS,S—an antibody-eliciting subunit vaccine targeting the major sporozoite surface protein circumsporozoite protein (CSP) [8]. Vaccines based on attenuated live sporozoites that arrest in the liver and function by a combination of T cells and antibodies have also demonstrated robust protection[9]. Unfortunately, despite significant efficacy from both approaches in controlled human malaria infection (CHMI) studies in malaria-naïve volunteers, both vaccines have markedly reduced efficacy in field trials and have not met the goals of 75–80% protection for vaccines [10] or injectable anti-malarials [11]. Recent encouraging Phase II results with the R21 CSP particle in Matrix-M adjuvant do meet the WHO goal of 75% protection against clinical disease for one year [12]. However, this protection appears less robust in preventing infection and depends on high antibody titers requiring yearly boosters that are vulnerable to the same interruptions as current interventions. These realities highlight the significant room for improvement in both T cell and antibody-eliciting vaccines, with the latter more amenable to iterative improvement due to available *in vitro* and *in vivo* preclinical assays [13–16].

Of the hundreds of proteins expressed at the sporozoite stage, at least 47 are surface-exposed [17–19] and therefore potentially accessible to antibodies. However, few of these proteins have been rigorously investigated for their use in antibody-eliciting vaccines [20]. In addition to CSP, the thrombospondin-related anonymous protein (TRAP, also known

as sporozoite surface protein 2 or SSP2) has been pursued as a vaccine candidate due

to its high abundance, essentiality for sporozoite infection and correlation with protection

in naturally-immune persons [21,22]. The TRAP ectodomain consists of 3 main domains: a

von Willebrand factor A-like domain (vWA), the thrombospondin repeat (TSR) domain

and a repeat domain [21]. The most advanced TRAP vaccine candidate is an

adenovirus/MVA-vectored vaccine eliciting strong T cell responses that has had low or

mixed efficacy results in CHMI trials [23,24] and field trials [25] but has been improved in

mice following targeting of the T cell response to the liver [26]. A combination protein

TRAP/RTS,S immunization failed to show significant protection [27], while a fusion-protein

approach using TRAP and CSP resulted in complete protection [28]. Antibody function in

experiments involving immunization with TRAP-derived peptides yielded mixed results

ranging from significant sporozoite inhibition in vitro [29] to no protection in vivo [30] . In

summary, the results using TRAP alone or in combination with CSP are difficult to

interpret due to the diversity of vaccine platforms used, the possibility of immune

interference in studies combining platforms, and the unclear dominance of roles for

antibodies and T cells in protection [31]. Whether a more targeted TRAP antibody

response could contribute to protection either alone or in combination with CSP remains

poorly defined.

Here, we used both active immunization and passive transfer of monoclonal antibodies

raised against either *Plasmodium yoelii* (rodent malaria) and *Plasmodium falciparum*

(human malaria) TRAP to more directly explore the potential efficacy of anti-TRAP

antibodies. We found that anti-TRAP antibodies modestly prevent liver infection in a

manner dependent on the TRAP domain recognized. Importantly, however, we

demonstrate proof of concept that anti-TRAP antibodies with minimal protective

capacity of their own can augment anti-CSP antibodies, providing additive protection

that raises their protective efficacy above the desired threshold of 75–80% protection.

Together, these findings argue for further investigation of rationally designed antibody-

eliciting malaria vaccines that targets multiple antigens and might include CSP as well

as non-CSP targets such as TRAP.

Results

PyTRAP polyclonal antibodies can prevent parasite infection of hepatocytes in vitro and

in vivo.

To elicit potentially functional anti-TRAP antibodies, we generated full-length

ectodomains and fragments of both rodent (*P. yoelii*) and human (*P. falciparum*) malaria

TRAP proteins (**Fig. 1A; Suppl. Table 1**) and verified their purity (**Fig. 1B**). Serum from

mice immunized with the rodent malaria *P. yoelii* TRAP ectodomain (PyTRAP)

recognized *Py* sporozoites by immunofluorescence in a pattern consistent with

micronemal localization, indicating the antigenic fidelity of the recombinant protein (**Fig.**

2A). We further tested this serum in an inhibition of sporozoite traversal and invasion

(ISTI) assay. Compared to control serum, anti-PyTRAP serum was able to modestly but

significantly (p=0.028) reduce sporozoite invasion of Hepa1-6 hepatoma cells *in vitro* at

a level similar to serum from mice immunized with the recombinant PyCSP ectodomain,

although the latter failed to reach significance (p=0.106) (**Fig. 2B**). In contrast,

sporozoite traversal of Hepa1-6 cells was not affected by anti-PyTRAP serum

(p=0.125), whereas anti-PyCSP serum did significantly reduce traversal (p=0.0057)

(**Fig. 2C**). The known inhibitory anti-PyCSP mAb 2F6 reduced both inhibition and

traversal in this assay as expected (**Fig. 2B and C**).

PyTRAP-immunized mice were then challenged with *Py* sporozoites via mosquito bite to

determine if these antibodies could function *in vivo* to reduce liver infection. We utilized

a PyGFPluc parasite, which expresses luciferase, enabling the measurement of liver

stage parasite burden by bioluminescence imaging. In contrast to mice immunized with

a non-specific control protein (Env), which showed no reduction in parasite liver stage

burden following challenge compared with naive mice, mice immunized with the

PyTRAP ectodomain demonstrated a significant 62% reduction of parasite liver stage

burden. Mice immunized with PyCSP ectodomain had a 91% reduction relative to naive

controls (**Fig. 2D**). Together, these data indicate that anti-PyTRAP antibodies can

function *in vitro* and *in vivo* to reduce parasite infection of hepatocytes.

PyTRAP monoclonal antibodies display a diverse array of functions in vitro and can

provide additive protection to anti-CSP antibodies in vivo.

Serum polyclonal antibodies, as studied above, are a mixture of many antibody

specificities, making it difficult to characterize the relative contribution to functional

activity of responses directed at different domains. To enable such a characterization of

the repertoire of PyTRAP-elicited antibodies, we produced a panel of 15 mAbs. When

tested in ISTI at 10, 50 and 100 µg/mL, 12 of these mAbs significantly inhibited invasion

or traversal at one or more concentrations, with mAbs TY03 and TY11 showing the

most consistent and potent inhibition (**Fig. 3A**, **Suppl. Fig. 1**).

Overall, the mAbs demonstrated a wide range of binding affinities to recombinant

PyTRAP (**Fig. 3B**, **Suppl. Table 2**) and recognized epitopes in the vWA, TSR and

repeat regions (**Suppl. Table 3**), thus covering the entire protein ectodomain. Among

the 15 mAbs recovered, 10 mAbs bound to the vWA domain. Six of these (TY02, TY05,

TY06, TY10, TY11, TY20) shared variable-segment assignments for both heavy and

light chains, had closely related complementarity-determining-region (CDR) sequences

and had 88.4–96.7% and 93.9–96.9% sequence identity in the variable-region

sequences of their heavy and light chain, respectively (**Suppl. Table 4 and Suppl. Fig.**

2). As expected, these antibodies were functionally similar in that they bound

specifically to the vWA domain (**Suppl. Table 3**) and inhibited sporozoite infection *in*

vitro (**Fig. 3A, Suppl. Fig. 1**). Two mAbs specifically recognized the TSR domain, and

the remaining three mAbs bound epitopes in the repeat region (**Suppl. Table 3**). These

non-vWA antibodies had only modest or no sporozoite inhibition of infection *in vitro* (**Fig.**

3A). This panel of mAbs also showed a wide range of binding kinetics, with all strongly

inhibitory mAbs having a k_{on} of $>10^5$ M^{-1}s^{-1} and a k_{dis} of $<10^{-2}$ s^{-1} (**Figure 3B**, note the

red box, **Suppl. Table 4**). Together, these data demonstrate that, similar to polyclonal

antibodies, anti-PyTRAP mAbs can mediate anti-parasitic function *in vitro*, and that

inhibitory function likely depends on fast and stable binding to the vWA domain,

although within the vWA domain there might exist epitopes that show higher

vulnerability to blocking of infection.

We next wanted to determine whether an anti-PyTRAP mAb could provide sterilizing

protection *in vivo* on its own or in combination with an anti-CSP mAb. For this, we chose

three vWA domain-binding anti-PyTRAP mAbs: TY03 and TY11, which were the top-

performing mAbs in ISTI, and TY12, which failed to demonstrate efficacy in ISTI. The

anti-PyTRAP mAbs were given at 300 µg/mouse (~15 mg/kg) alone or with a partially

protective dose of 100 µg/mouse (~5 mg/kg) of anti-PyCSP mAb 2F6 prior to mosquito

bite challenge [32]. As shown in **Fig. 3C** and **Table 1**, mice administered anti-PyCSP mAb

2F6 showed significant sterile protection, with 9/18 (50%) remaining parasite-free,

compared to 2/15 (13.3%) for mice receiving non-specific murine IgG (p=0.032). Neither

TY11 nor TY12 showed any protection (2/13 or 15.4% non-infected) despite TY11

demonstrating the most robust inhibition *in vitro*. Administration of the mAb TY03

resulted in 7/16 mice (43.7%) remaining parasite-free which did not reach significance

(p=0.072). When combined with the anti-CSP mAb, only the addition of TY03 afforded

significant sterile protection (87.5% or 14/16 mice) over the control group (p<0.001)

which, importantly, was a significant improvement over protection observed with anti-

PyCSP mAb alone (p=0.025). Together these data indicate that while *in vitro* testing of

mAbs can be useful for identifying non-functional mAbs (e.g. TY12), they should be

validated *in vivo* for function. Importantly, these data provide proof of concept that non-

CSP antibodies can provide at least additive protection to anti-CSP antibodies, even

when those non-CSP antibodies only provide partial levels of protection on their own.

Antibodies targeting the human malaria parasite P. falciparum TRAP can function

against sporozoite invasion of hepatocytes.

We next wanted to determine if antibodies directed against TRAP/SSP2 from the

human malaria parasite, *P. falciparum*, could also function to prevent sporozoite

infection. Serum from mice immunized with the ectodomain of *P. falciparum* TRAP

(PfTRAP) was able to recognize *Pf* sporozoites in IFA (**Fig. 4A**) and demonstrated

consistent inhibition of *Pf* sporozoite invasion *in vitro* at a level similar to serum from

mice immunized with the ectodomain of *P. falciparum* CSP (PfCSP) (**Fig. 4B**). Inhibition

of sporozoite traversal *in vitro* was more modest as compared to anti-PfCSP polyclonal

serum (**Fig. 4C**). The known inhibitory anti-PfCSP mAb 2A10 demonstrated robust

inhibition of both invasion and traversal (**Fig. 4A and B**).

Using a similar approach to the anti-PyTRAP work described above, we isolated 7 anti-

PfTRAP mAbs from immunized mice. Of these mAbs, 5 recognized the vWA domain

and 2 recognized the TSR domains (**Suppl. Table 3**). In contrast to the high proportion

of functional anti-PyTRAP mAbs (12 of 15), only 2 of 7 anti-PfTRAP mAbs, both

recognizing the vWA domain, showed any sporozoite-inhibitory function *in vitro*: AKBR-

4 and AKBR-10. Further, only AKBR-4 demonstrated significant inhibition of both

invasion and traversal (**Fig. 5A**), despite having unremarkable binding properties with

the PfTRAP ectodomain (**Fig. 5B**). Surprisingly, mAb AKBR-7, which had the best

binding properties of the set (K_d ~0.15±0.04 nM, **Suppl. Table 2**), demonstrated the

worst inhibitory properties (**Fig. 5B**). Similar to the case with the anti-PyTRAP mAb

panel described above, our data suggest that the PfTRAP vWA domain contains

epitopes exposing vulnerability to inhibition, however lack of mAbs that strongly bind

other portions of PfTRAP make it difficult to discount the roles that these domains may

play in inhibition in vivo.

A vWA-directed anti-PfTRAP mAb increases the protection afforded by a protective

CSP mAb.

Because *Pf* sporozoites do not infect murine livers, the only means to test the activity of

anti-*Pf* antibodies against sporozoite infection *in vivo* is by either challenging passively

or actively immunized wild-type mice with transgenic rodent parasites expressing the *Pf*

proteins of interest [33–35] or by passive immunization of immune-deficient humanized liver

mice (FRG huHep), which can be challenged with *Pf* sporozoites [14]. We chose to utilize

the latter as it is an established model of antibody-mediated protection against *Pf*

infection [14,36–41] and allows testing of any future combination of anti-*Pf* antibodies

without the need for generating combinatorial transgenic parasites. In this model,

humanized-liver mice receive a passive transfer of antibodies and are then infected with

Pf via mosquito bite. Six days later, mice are injected with human red blood cells, which

can then be infected by emerging merozoites and infection quantified by qRT-PCR on

days 7 and 9.

Using this method, we tested the ability of the anti-PfTRAP mAb AKBR-4 to provide

sterile protection against *Pf* mosquito-bite infection alone or in combination with a

partially-protective anti-PfCSP mAb CIS43 [37]. We chose a dose of 50 µg/mouse (~2.5

mg/kg) for each mAb, as this provides partial protection with an anti-PfCSP mAb [37] and

gives a serum concentration of ~10 µg/mL at the time of infection, which is achievable

by both active vaccination and passive transfer of long-lasting mAbs [42,43]. We previously

conducted passive administration, mosquito bite challenge in 2 replicate experiments [37]

that showed that 50 µg/mouse dose of anti-PfCSP mAb CIS43 was protective (5/7 and

5/8 protected in each experiment), compared to control mice (0/7 and 0/7 protected). To

avoid unnecessary duplication of FRG huHep experiments, we included those cohorts

in our overall analysis of mAbs in this study. We conducted a third replicate experiment

with an additional 8 CIS43 mAb mice and 5 control mAb mice in the current study,

where 5/8 CIS43 treated mice were protected and 0/5 control mice were unprotected,

bringing our total experimental yield to a total of 15/23 protected (65%), which was

significant compared to 0/19 of control mice protected (0%, $p<0.0001$; **Table 2**, **Fig.**

5C). This protection was not improved in a single experiment of 5 FRG huHep mice in

which the dose was increased 3-fold to 150 µg/mouse (3/5, 60% protected; $p=0.002$

over control).

On its own, passive administration of 50 µg/mouse of AKBR-4 failed to provide any

sterile protection (0/11, 0%). Yet, when 50 µg/mouse of AKBR-4 was combined with 50

µg/mouse of the anti-PfCSP mAb (100 µg mAb/mouse total), 14/16 (88%) mice were

sterilely protected. The improvement afforded by the AKBR-4/anti-PfCSP mAb

combination over the efficacy of the anti-PfCSP mAb alone (up to 65% protection with

up to 150 µg/mouse) trended toward, but did not reach statistical significance at this

group size ($p=0.131$), despite still being statistically protective compared to the control

antibody treatment ($p<0.0001$). Nevertheless, together these results strongly support

proof of concept that polyclonal and monoclonal antibodies directed against PfTRAP

can reduce Pf sporozoite invasion and traversal of hepatocytes *in vitro* and are able to

raise the protective capacity of anti-PfCSP antibodies above the threshold of 75%

protection.

Discussion

Here, we have shown that combining antibodies recognizing TRAP and CSP resulted in an additive protective effect when assessed by a mosquito-bite challenge. To date, the most extensive and clear proof-of-concept that non-CSP antigens can be combined with CSP to provide superior sterile protection was from a recently published rodent malaria study [44]. The protective contributions of antibodies in that study were not specifically investigated, as antibodies were deemed irrelevant for protection compared to the role of T cells. This was determined in the context of a vectored vaccine immunization, which is biased towards the elicitation of T cells, and using an intravenous sporozoite challenge, which has been shown to bypass the role of antibodies in the skin and interstitial tissues [45,46]. Therefore, although promising, this study was not designed to assess the substantial role of antibodies in protection from pre-erythrocytic infection as has been previously reported [45–47]. However, these findings, along with our data, suggest that enhanced protection over CSP-alone immunization is possible by way of multivalent subunit vaccination targeting distinct sporozoite surface-exposed proteins. Thus, further efforts should be directed at identifying additional immunogen combinations in preclinical studies.

In light of the complex life cycle of *Plasmodium*, dissection of antibody function requires the use of polyclonal or monoclonal antibodies *in vitro* or in passive transfer *in vivo* followed by a mosquito-bite or intradermal sporozoite challenge. The challenge mode may be particularly important to take advantage of the protective effects elicited by antibodies while the parasite is traversing the skin [46]. In a further complication, studies

with CSP have shown that within a functional polyclonal antibody response, ineffective

antibodies may make up the majority and only a subset are potent antibodies

distinguished by nuanced binding properties [36,37,48–53]. Understanding these complex

activities can identify superior mAb products and vaccine immunogens, yet such studies

have not been previously performed for TRAP or other non-CSP pre-erythrocytic

antibody targets. Here, we show that the polyclonal antibody response to full-length

PyTRAP can substantially reduce parasite infection of hepatocytes *in vitro* and *in vivo*.

We confirmed these observations using monoclonal antibodies and conclude that this

effect is likely driven by vWA and TSR-specific antibodies. These findings clearly

support TRAP as a functional antibody target and is in line with previous studies using

antibodies against TRAP protein fragments [29], yet are in contrast to others that failed to

see significant inhibition [30]. Our data with PfTRAP were more limited but the only mAb

that was functional *in vitro* recognized the vWA domain. The difference in the

proportions of functional antibodies we observed between PyTRAP and PfTRAP may

be due to subtle differences in protein localization [54,55] and, potentially, binding partners

[56] between the two species.

Achievable and minimal functional antibody levels are important parameters in vaccine

and monoclonal antibody product development. In our *in vivo Py* inhibition studies, the

combination of mAbs brought the total antibody dose to four times that of the anti-

PyCSP antibody alone (100 µg/mouse of anti-PyCSP alone to 400 µg/mouse total with

the combination). It is reasonable to expect that simply increasing the dose of the anti-

CSP mAb would have also yielded higher levels of protection in this mouse model. Yet

experience with RTS,S—which elicits extremely high levels of anti-CSP antibodies—as

well as published data using highly potent anti-CSP mAbs [37,49,57] suggest that

increasing anti-CSP titers alone provides diminishing returns. Furthermore, sustaining

such high levels of antibodies over years may be unachievable by vaccines or mAb

prophylaxis. The first CHMI trial using passive transfer of the same CIS43 anti-PfCSP

mAb used in this study indeed showed that mAbs can provide sterilizing protection

against *P. falciparum* mosquito bite infection at serum concentrations between ~50–500

µg/mL. However, prophylactic mAb products will likely need increased potency in order

to provide long-lasting protection at lower concentrations against more diverse and

likely more fit parasites in the field. Rather than increasing the dose or frequency of

administration of anti-CSP mAbs to sustain high levels of serum concentrations, long-

term maintenance of moderate or low titers against two or more antigens with additive

protection properties may be more easily achieved. Our experiments using *Pf* mosquito-

bite challenge in FRGhuHep mice support such a strategy, as we were able to achieve

high levels of sterile protection (88%) using a combination of anti-PfCSP and anti-

PfTRAP mAbs. Importantly, this level of protection was achieved using a low total dose

of mAb (100 µg/mouse or ~5 mg/kg) and brought the combination above the 75–80%

protection threshold desired for vaccines [10] or injectable anti-malarials [11]. Furthermore,

our total combination dose of 100 µg/mouse (50 µg/mouse each of anti-PfCSP and anti-

PfTRAP mAb) gave a total circulating mAb concentration of ~20 µg/mL—a level that

can be achieved for ~36 weeks with a single 20 mg/kg injection of long-lasting mAbs

[42,58,59] or ~4 years via active vaccination [60] . On the other hand, our data showing that

tripling the dose of the anti-PfCSP mAb alone (150 µg/mouse, 7.5 mg/kg) failed to

improve protection, therefore, suggests that reaching desired thresholds of protection at

lower antibody concentrations may be more efficiently achieved by adding additional

antibody targets rather than increasing the anti-CSP antibody concentration.

In summary, we provide proof of concept that antibodies targeting TRAP can contribute

to sterile protection when used in combination with anti-CSP antibodies. These findings

support vaccine and mAb strategies where multiple antigens are targeted to enhance

the protection afforded by CSP. Specifically, we posit that rationally designed non-CSP

proteins or mAbs may be powerful tools for eliciting high levels of long-lasting protection

at lower antibody titers, as our data suggest they can enhance CSP-based protection

even when they do not elicit substantial sterile protection on their own. Such a

multivalent approach has previously been hampered by the difficulty in generating and

combining multiple protein-in-adjuvant formulations, but may be more easily achieved

by the use of mRNA-based vaccines which have proven adept as a multi-antigen

vaccine platform in preclinical studies [61,62]. Alternatively, pre-selected, multivalent mAbs

could be used as a cocktail, although such an approach is currently limited by cost [63].

Our results indicate that efforts to develop a long-lasting, infection-blocking malaria

intervention would greatly benefit from identifying non-CSP antibody targets that

enhance CSP-elicited protection.

<u>Materials and Methods</u>

Recombinant protein production

Recombinant proteins were produced in transiently transfected suspension culture of

FreeStyle 293 cells (Thermo), as previously described [64]. Briefly, codon-optimized

constructs encoding the ectodomain or deletions of *Plasmodium falciparum* (PfTRAP)

and *Plasmodium yoelii* TRAP (PyTRAP) were generated as fusions to the 8xHis and

AviTag [65] sequences (Table 1). Following transfection using the high-density PEI

method [66] and the subsequent 5-day incubation, cells were removed by centrifugation

and the culture supernatants were supplemented with NaCl (+350 mM) and sodium

azide (0.02%). Treated culture supernatants were passed by gravity through NiNTA

agarose, washed with Wash Buffer (10 mM Tris-HCl, pH 8, 300 mM NaCl, 10 mM

imidazole), and eluted with Elution Buffer (10 mM Tris-HCl, pH 7.4, 300 mM NaCl, 200

mM imidazole). Further purification was performed by size-exclusion chromatography

using a calibrated Superdex 200 (10/600) column. The HIV Env gp120 control protein

was produced, as previously described [67].

16
Monoclonal antibody cloning and production

Monoclonal antibodies were cloned and produced, as previously described [64]. Briefly,

ectodomain PfTRAP and PyTRAP constructs were used as immunogens, and their

biotinylated versions were used to isolate antigen-specific B cells by flow cytometry.

Following culture, wells containing B cells producing antigen-binding IgG were identified

by ELISA, immunoglobulin-encoding transcripts were amplified by RT-PCR and used for

the generation of heavy- and light-chain constructs for recombinant mAb expression.

The sequences were annotated using IgBLAST [68].

To express recombinant mAbs, the plasmid DNA was used to transfect suspension

cultures of FreeStyle 293 cells (Thermo), as described above. After five days in culture,

cells were removed by centrifugation and the cultures were supplemented with NaCl

(+350 mM) and sodium azide (0.02%). Treated culture supernatants were passed by

gravity through Protein G resin equilibrated in Wash Buffer (10 mM HEPES, pH 7, 300

mM NaCl, 2 mM EDTA), washed with Wash Buffer, and eluted with 100 mM glycine, pH

2.7. Resulting eluates were buffer-exchanged by repeated centrifugal ultrafiltration with

HBS-E (10 mM HEPES, pH 7, 150 mM NaCl, 2 mM EDTA).

Binding kinetics measurements

Binding properties of mAbs were characterized using biolayer interferometry (BLI)

measurements on an Octet QK^e instrument (Sartorius), as previously described [64].

Briefly, antibodies in culture supernatants were immobilized on anti-Mouse IgG Fc

Capture biosensors and allowed to associate with antigen serially diluted (in the range

of 1–1000 nM) in 10x Kinetics Buffer (10xKB: PBS + 0.1% BSA, 0.02% Tween-20 and

0.05% sodium azide) followed by dissociation in 10x KB. Resulting sensorgram data

was evaluated using ForteBio Data Analysis software (version 7.0.1.5) to generate a fit

to the 1:1 binding model and provide estimates for the k_{on} and k_{dis} rate constants.

Coarse epitope mapping by ELISA

Domain specificity of the mAbs was characterized by enzyme-linked immunosorbent

assay (ELISA) using TRAP ectodomain and fragments from PfTRAP and PyTRAP, as

previously described [64].

Sporozoite production

For rodent parasite (*P. yoelii*), female Swiss Webster (SW) mice for parasite cycles

were purchased from Envigo laboratories and injected intraperitoneally (i.p.) with blood

stage PyGFPluc [69]. Three days later, gametocyte exflagellation was confirmed and the

infected mice were used to feed female *Anopheles stephensi* mosquitoes. Fourteen to

16 days after the feed, salivary gland sporozoites were isolated from the mosquitoes

and used in mouse infections.

For human malaria (*P. falciparum*) experiments, infected *A. stephensi* mosquitoes were

produced, as previously described [70].

Animal studies ethics statement

All procedures involving animals were performed in adherence to protocols of the

Institutional Animal Care and Use Committee at the Seattle Children's Research

Institute. The study was carried out in compliance with the ARRIVE guidelines

(https://arriveguidelines.org/).

Mouse active immunization and challenge

To generate polyclonal serum and a source of mouse monoclonal antibodies, six to

eight week-old BALBc/J mice were purchased from Jackson laboratories and injected

intramuscularly three times at days 0, 14 and 38 using Adjuplex mixed with 20–25 µg of

target protein. Mice immunized with recombinant *Py* proteins were then challenged by

the bite of 15 PyGFPluc-infected mosquitos, as published [32]. Forty-two hours later,

parasite liver burden was assessed by bioluminescent imaging, as previously published

[32]. Mice were then immediately sacrificed and splenocytes collected and cryopreserved

for B cell isolation and mAb production.

Mice immunized with *Pf* proteins were immunized as above with the exception that mice

were additionally boosted with IV protein three days prior to sacrifice and collection and

cryopreservation of splenocytes.

For both, serum was collected from immunized mice by collecting whole blood in BD

microtainer serum tubes, allowing blood to clot at room temperature for at least 30

minutes and then centrifuged according to manufacturer's instructions to separate

serum for storage and use in *in vitro* assays.

Sporozoite immunofluorescence microscopy

Fresh-dissected *Py* or *Pf* sporozoites were stained, as previously published [71]. Briefly,

fresh-dissected *Py* or *Pf* sporozoites were fixed with 4% PFA and air-dried onto glass

slides overnight. These were then permeabilized with 0.1% Triton-X and stained with

polyclonal (serum at 1:200 dilution) or monoclonal (10 µg/mL) antibodies. Sporozoites

were identified by co-staining with either anti-MTIP or anti-CSP mAbs as well as DAPI

for nuclear localization. Images were acquired using an Olympus IX-70 DeltaVision

deconvolution microscope at 100X magnification.

In vitro inhibition of sporozoite traversal and invasion (ISTI)

In vitro ISTI was performed, as previously published for both *Py* [45] and *Pf* [14]. Briefly,

fresh-dissected sporozoites were added to hepatoma cells (Hepa1-6 for *Py* and HC04

for *Pf*) plated the day prior in 96-well plates in the presence of antibodies and FITC-

dextran in technical duplicates or triplicates. After 90 minutes, cells were fixed,

permeabilized and stained with fluorescently-labeled anti-CSP mAbs and analyzed by

flow cytometry. Invaded cells were identified by the presence of CSP and traversed

cells by the uptake of FITC-dextran. Within each experimental replicate, antibody-

treated wells were normalized to the invasion and traversal of wells treated with pre-

immune serum or non-specific mouse IgG which was set to 100%.

Py passive transfer studies

Six to eight week-old BALBc/J mice were intravenously injected with indicated doses of

monoclonal antibody 24 hours prior to challenge by bite of 5 PyGFPluc-infected

mosquitos. Mice were followed up for infection by Giemsa-stained thin blood smear

every day from days 3–7 and every other day from days 8–14 for identification of blood

stage parasites. Mice in which we failed to identify parasites in 40,000 red blood cells

over the entire period were considered negative and sterilely protected. Control mice

were administered non-specific polyclonal mouse IgG at a dose equivalent to the

highest dose in experimental groups.

Pf passive transfer in FRG humanized liver mice

Mice repopulated with human hepatocytes (FRGhuHep) were purchased from Yecuris,

Inc. and infected with *Pf* via mosquito bite, as previously published [14,36]. Briefly,

indicated doses of mAb were intravenously injected into mice 24 hours prior to

challenge by the bite of 5 *Pf*-infected mosquitos. On day 6 post-infection, mice were

intravenously injected with 400 µL of human red blood cells and on days 7 and 9

peripheral blood was collected and assessed for parasites by qRT-PCR for *Pf*18s rRNA.

Any mouse with a Ct value above the no template control at either day was considered

positive for parasitemia.

Statistics

Statistical analyses and plotting were carried out in Prism (GraphPad) or in R (version

4.0.2) using packages Exact (version 2.1), ggpubr (version 0.4.0), ggstatsplot (version

0.7.2). Statistical tests and outcomes are noted in the figure legend for each figure. For

all tests, a p-value of <0.05 was considered significant and values not specifically

labeled were above this threshold.

Data availability

DNA sequences encoding the mAbs described here have been deposited in GenBank

(accession numbers OK484322–OK484365).

1 **<u>Acknowledgements</u>**

2 We would like to thank the vivarium staff at Seattle Children's Research Institute for

3 their support of animal studies. We would also like to thank Drs. Nevile Kisalu and

4 Robert Seder of the NIH VRC for their provision of monoclonal antibody CIS43. This

5 study was funded by NIH R01 AI117234 to DNS and SHK.

Author contributions

B.K.W. and V.V. contributed equally to this work.

Conceptualization and experimental design: B.K.W., V.V., S.H.I.K. and D.N.S.

Investigation: B.K.W., V.V., S.C., N.M., N.H., A.R., H.C, B.G.O., O.T., S.K., N.D., S.A.A. and N.C.

Data analysis and visualization: B.K.W., V.V., N.H., N.M.

Writing — Original draft: B.K.W.

Writing — Review and editing: B.K.W., V.V., S.H.I.K. and D.N.S.

Resources: S.H.I.K. and D.N.S.

Supervision, Project Administration and Funding Acquisition: B.K.W., S.H.I.K., D.N.S.

1 **<u>Competing Interests statement</u>**

2 The authors declare no competing interests.

References

1. Geneva: World Health Organization. World malaria report 2020: 20 years of global progress and challenges. (2020).
2. Medica, D. L. & Sinnis, P. Quantitative dynamics of Plasmodium yoelii sporozoite transmission by infected anopheline mosquitoes. *Infect. Immun.* **73**, 4363–4369 (2005).
3. Ejigiri, I. & Sinnis, P. Plasmodium sporozoite-host interactions from the dermis to the hepatocyte. *Curr. Opin. Microbiol.* **12**, 401–407 (2009).
4. Vaughan, A. M. & Kappe, S. H. I. Malaria Parasite Liver Infection and Exoerythrocytic Biology. *Cold Spring Harb. Perspect. Med.* **7**, (2017).
5. Cowman, A. F., Healer, J., Marapana, D. & Marsh, K. Malaria: Biology and Disease. *Cell* **167**, 610–624 (2016).
6. Lindner, S. E., Miller, J. L. & Kappe, S. H. I. Malaria parasite pre-erythrocytic infection: preparation meets opportunity. *Cell. Microbiol.* **14**, 316–324 (2012).
7. Duffy, P. E. & Patrick Gorres, J. Malaria vaccines since 2000: progress, priorities, products. *NPJ Vaccines* **5**, 48 (2020).
8. Tinto, H. *et al.* Long-term incidence of severe malaria following RTS,S/AS01 vaccination in children and infants in Africa: an open-label 3-year extension study of a phase 3 randomised controlled trial. *Lancet Infect. Dis.* **19**, 821–832 (2019).
9. Epstein, J. E. *et al.* Protection against Plasmodium falciparum malaria by PfSPZ Vaccine. *JCI Insight* **2**, e89154 (2017).
10. Malaria Vaccine Funders Group. *Malaria Vaccine Technology Roadmap.* https://www.who.int/publications/m/item/malaria-vaccine-technology-roadmap (2013).
11. Macintyre, F. *et al.* Injectable anti-malarials revisited: discovery and development of new agents to protect against malaria. *Malar. J.* **17**, 402 (2018).
12. Datoo, M. S. *et al.* Efficacy of a low-dose candidate malaria vaccine, R21 in adjuvant Matrix-M, with seasonal administration to children in Burkina Faso: a randomised controlled trial. *Lancet* **397**, 1809–1818 (2021).
13. Roth, A. *et al.* A comprehensive model for assessment of liver stage therapies targeting Plasmodium vivax and Plasmodium falciparum. *Nat. Commun.* **9**, 1837 (2018).
14. Sack, B. K. *et al.* Humoral protection against mosquito bite-transmitted Plasmodium falciparum infection in humanized mice. *NPJ Vaccines* **2**, 27 (2017).
15. Steel, R. W. J. *et al.* An Opsonic Phagocytosis Assay for Plasmodium falciparum Sporozoites. *Clin. Vaccine Immunol.* **24**, (2017).
16. Boyle, M. J. *et al.* Human antibodies fix complement to inhibit Plasmodium falciparum invasion of erythrocytes and are associated with protection against malaria. *Immunity* **42**, 580–590 (2015).
17. Swearingen, K. E. *et al.* Proteogenomic analysis of the total and surface-exposed proteomes of Plasmodium vivax salivary gland sporozoites. *PLoS Negl. Trop. Dis.* **11**, e0005791 (2017).
18. Swearingen, K. E. *et al.* Interrogating the Plasmodium Sporozoite Surface: Identification of Surface-Exposed Proteins and Demonstration of Glycosylation on CSP and TRAP by Mass Spectrometry-Based Proteomics. *PLoS Pathog.* **12**, (2016).
19. Lindner, S. E. *et al.* Total and putative surface proteomics of malaria parasite salivary gland sporozoites. *Mol. Cell. Proteomics* **12**, 1127–1143 (2013).
20. Sack, B., Kappe, S. H. I. & Sather, D. N. Towards functional antibody-based vaccines to prevent pre-erythrocytic malaria infection. *Expert Rev. Vaccines* **16**, 403–414 (2017).
21. Dolo, A. *et al.* Thrombospondin related adhesive protein (TRAP), a potential malaria vaccine candidate. *Parassitologia* **41**, 425–428 (1999).
22. John, C. C. *et al.* Correlation of high levels of antibodies to multiple pre-erythrocytic

Plasmodium falciparum antigens and protection from infection. *Am. J. Trop. Med. Hyg.* **73**, 222–228 (2005).

23. Hodgson, S. H. *et al.* Evaluation of the efficacy of ChAd63-MVA vectored vaccines expressing circumsporozoite protein and ME-TRAP against controlled human malaria infection in malaria-naive individuals. *J. Infect. Dis.* **211**, 1076–1086 (2015).

24. Ewer, K. J. *et al.* Protective CD8+ T-cell immunity to human malaria induced by chimpanzee adenovirus-MVA immunisation. *Nat. Commun.* **4**, 2836 (2013).

25. Ogwang, C. *et al.* Prime-boost vaccination with chimpanzee adenovirus and modified vaccinia Ankara encoding TRAP provides partial protection against Plasmodium falciparum infection in Kenyan adults. *Sci. Transl. Med.* **7**, 286re5 (2015).

26. Gola, A. *et al.* Prime and target immunization protects against liver-stage malaria in mice. *Sci. Transl. Med.* **10**, (2018).

27. Kester, K. E. *et al.* Sequential Phase 1 and Phase 2 randomized, controlled trials of the safety, immunogenicity and efficacy of combined pre-erythrocytic vaccine antigens RTS,S and TRAP formulated with AS02 Adjuvant System in healthy, malaria naïve adults. *Vaccine* **32**, 6683–6691 (2014).

28. Lu, C. *et al.* Design and assessment of TRAP-CSP fusion antigens as effective malaria vaccines. *PLoS One* **15**, e0216260 (2020).

29. Charoenvit, Y. *et al.* Development of two monoclonal antibodies against Plasmodium falciparum sporozoite surface protein 2 and mapping of B-cell epitopes. *Infect. Immun.* **65**, 3430–3437 (1997).

30. Gantt, S. *et al.* Antibodies against thrombospondin-related anonymous protein do not inhibit Plasmodium sporozoite infectivity in vivo. *Infect. Immun.* **68**, 3667–3673 (2000).

31. Rampling, T. *et al.* Safety and High Level Efficacy of the Combination Malaria Vaccine Regimen of RTS,S/AS01B With Chimpanzee Adenovirus 63 and Modified Vaccinia Ankara Vectored Vaccines Expressing ME-TRAP. *J. Infect. Dis.* **214**, 772–781 (2016).

32. Sack, B. K. *et al.* Model for in vivo assessment of humoral protection against malaria sporozoite challenge by passive transfer of monoclonal antibodies and immune serum. *Infect. Immun.* **82**, 808–817 (2014).

33. Atcheson, E. *et al.* Tailoring a Plasmodium vivax Vaccine To Enhance Efficacy through a Combination of a CSP Virus-Like Particle and TRAP Viral Vectors. *Infect. Immun.* **86**, (2018).

34. Longley, R. J. *et al.* Assessment of the Plasmodium falciparum Preerythrocytic Antigen UIS3 as a Potential Candidate for a Malaria Vaccine. *Infect. Immun.* **85**, (2017).

35. Longley, R. J. *et al.* Comparative assessment of vaccine vectors encoding ten malaria antigens identifies two protective liver-stage candidates. *Sci. Rep.* **5**, 11820 (2015).

36. Tan, J. *et al.* A public antibody lineage that potently inhibits malaria infection through dual binding to the circumsporozoite protein. *Nat. Med.* **24**, 401–407 (2018).

37. Kisalu, N. K. *et al.* A human monoclonal antibody prevents malaria infection by targeting a new site of vulnerability on the parasite. *Nat. Med.* **24**, 408–416 (2018).

38. Triller, G. *et al.* Natural Parasite Exposure Induces Protective Human Anti-Malarial Antibodies. *Immunity* **47**, 1197–1209.e10 (2017).

39. Kublin, J. G. *et al.* Complete attenuation of genetically engineered Plasmodium falciparum sporozoites in human subjects. *Sci. Transl. Med.* **9**, (2017).

40. Ishizuka, A. S. *et al.* Protection against malaria at 1 year and immune correlates following PfSPZ vaccination. *Nat. Med.* **22**, 614–623 (2016).

41. Foquet, L. *et al.* Vaccine-induced monoclonal antibodies targeting circumsporozoite protein prevent Plasmodium falciparum infection. *J. Clin. Invest.* **124**, 140–144 (2014).

42. Gaudinski, M. R. *et al.* Safety and pharmacokinetics of the Fc-modified HIV-1 human monoclonal antibody VRC01LS: A Phase 1 open-label clinical trial in healthy adults. *PLoS Med.* **15**, e1002493 (2018).

43. Ledgerwood, J. E. *et al.* Safety, pharmacokinetics and neutralization of the broadly neutralizing HIV-1 human monoclonal antibody VRC01 in healthy adults. *Clin. Exp. Immunol.* **182**, 289–301 (2015).

44. Daniel, S. *et al.* Plasmodium pre-erythrocytic vaccine antigens enhance sterile protection in mice induced by circumsporozoite protein. *Infect. Immun.* IAI0016521 (2021).

45. Keitany, G. J. *et al.* Immunization of mice with live-attenuated late liver stage-arresting Plasmodium yoelii parasites generates protective antibody responses to preerythrocytic stages of malaria. *Infect. Immun.* **82**, 5143–5153 (2014).

46. Flores-Garcia, Y. *et al.* Antibody-Mediated Protection against Plasmodium Sporozoites Begins at the Dermal Inoculation Site. *MBio* **9**, (2018).

47. Dups, J. N., Pepper, M. & Cockburn, I. A. Antibody and B cell responses to Plasmodium sporozoites. *Front. Microbiol.* **5**, 625 (2014).

48. Alanine, D. G. W. *et al.* Human Antibodies that Slow Erythrocyte Invasion Potentiate Malaria-Neutralizing Antibodies. *Cell* **178**, 216–228.e21 (2019).

49. Wang, L. T. *et al.* A Potent Anti-Malarial Human Monoclonal Antibody Targets Circumsporozoite Protein Minor Repeats and Neutralizes Sporozoites in the Liver. *Immunity* (2020) doi:10.1016/j.immuni.2020.08.014.

50. Livingstone, M. C. *et al.* In vitro and in vivo inhibition of malaria parasite infection by monoclonal antibodies against Plasmodium falciparum circumsporozoite protein (CSP). *Sci. Rep.* **11**, 5318 (2021).

51. Douglas, A. D. *et al.* A defined mechanistic correlate of protection against Plasmodium falciparum malaria in non-human primates. *Nat. Commun.* **10**, 1953 (2019).

52. Scally, S. W. *et al.* Rare PfCSP C-terminal antibodies induced by live sporozoite vaccination are ineffective against malaria infection. *J. Exp. Med.* **215**, 63–75 (2018).

53. Vijayan, K. *et al.* Antibody interference by a non-neutralizing antibody abrogates humoral protection against Plasmodium yoelii liver stage. *Cell Rep.* **36**, 109489 (2021).

54. Rogers, W. O., Rogers, M. D., Hedstrom, R. C. & Hoffman, S. L. Characterization of the gene encoding sporozoite surface protein 2, a protective Plasmodium yoelii sporozoite antigen. *Mol. Biochem. Parasitol.* **53**, 45–51 (1992).

55. Rogers, W. O. *et al.* Characterization of Plasmodium falciparum sporozoite surface protein 2. *Proc. Natl. Acad. Sci. U. S. A.* **89**, 9176–9180 (1992).

56. Steel, R. W. J. *et al.* Platelet derived growth factor receptor β (PDGFRβ) is a host receptor for the human malaria parasite adhesin TRAP. *Sci. Rep.* **11**, 11328 (2021).

57. Kisalu, N. K. *et al.* Enhancing durability of CIS43 monoclonal antibody by Fc mutation or AAV delivery for malaria prevention. *JCI Insight* **6**, (2021).

58. Gaudinski, M. R. *et al.* Safety and pharmacokinetics of broadly neutralising human monoclonal antibody VRC07-523LS in healthy adults: a phase 1 dose-escalation clinical trial. *Lancet HIV* **6**, e667–e679 (2019).

59. Gaudinski, M. R. *et al.* A Monoclonal Antibody for Malaria Prevention. *N. Engl. J. Med.* **385**, 803–814 (2021).

60. Minassian, A. M. *et al.* Reduced blood-stage malaria growth and immune correlates in humans following RH5 vaccination. *Med (N Y)* **2**, 701–719.e19 (2021).

61. Chahal, J. S. *et al.* Dendrimer-RNA nanoparticles generate protective immunity against lethal Ebola, H1N1 influenza, and Toxoplasma gondii challenges with a single dose. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E4133–42 (2016).

62. Vogel, A. B. *et al.* Self-Amplifying RNA Vaccines Give Equivalent Protection against Influenza to mRNA Vaccines but at Much Lower Doses. *Mol. Ther.* **26**, 446–455 (2018).

63. Pelfrene, E., Mura, M., Cavaleiro Sanches, A. & Cavaleri, M. Monoclonal antibodies as anti-infective products: a promising future? *Clin. Microbiol. Infect.* **25**, 60–64 (2019).

64. Carbonetti, S. *et al.* A method for the isolation and characterization of functional murine monoclonal antibodies by single B cell cloning. *J. Immunol. Methods* **448**, 66–73 (2017).

65. Fairhead, M. & Howarth, M. Site-specific biotinylation of purified proteins using BirA. *Methods Mol. Biol.* **1266**, 171–184 (2015).
66. Backliwal, G., Hildinger, M., Hasija, V. & Wurm, F. M. High-density transfection with HEK-293 cells allows doubling of transient titers and removes need for a priori DNA complex formation with PEI. *Biotechnol. Bioeng.* **99**, 721–727 (2008).
67. Sather, D. N. *et al.* Factors associated with the development of cross-reactive neutralizing antibodies during human immunodeficiency virus type 1 infection. *J. Virol.* **83**, 757–769 (2009).
68. Ye, J., Ma, N., Madden, T. L. & Ostell, J. M. IgBLAST: an immunoglobulin variable domain sequence analysis tool. *Nucleic Acids Res.* **41**, W34–40 (2013).
69. Miller, J. L. *et al.* Quantitative bioluminescent imaging of pre-erythrocytic malaria parasite infection using luciferase-expressing Plasmodium yoelii. *PLoS One* **8**, e60820 (2013).
70. Vaughan, A. M. *et al.* A transgenic Plasmodium falciparum NF54 strain that expresses GFP-luciferase throughout the parasite life cycle. *Mol. Biochem. Parasitol.* **186**, 143–147 (2012).
71. Sack, B. K. *et al.* Mechanisms of stage-transcending protection following immunization of mice with late liver stage-arresting genetically attenuated malaria parasites. *PLoS Pathog.* **11**, e1004855 (2015).

Tables

Table 1. Combination of anti-PyCSP and anti-PyTRAP can improve sterile protection from mosquito bite challenge.

	Sterile Protection				Comparison p value	
	Exp 1	Exp 2	Exp 3	Combined	vs. mIgG	vs. αCSP + mIgG p value
mIgG	1/5 (20%)	0/5 (0%)	1/5 (20%)	2/15 (13.3%)	–	**0.056**
αCSP + mIgG	4/8 (50%)	3/5 (60%)	2/5 (40%)	9/18 (50%)	**0.056**	–
TY11 + mIgG	1/3 (33%)	0/5 (0%)	1/5 (20%)	2/13 (15.4%)	0.79	0.048
TY12 + mIgG	2/4 (50%)	0/5 (0%)	0/4 (0%)	2/13 (15.4%)	0.79	0.048
TY03 + mIgG	5/6 (83.3%)	0/5 (0%)	2/5 (40%)	7/16 (43.7%)	0.072	0.734
αCSP + TY11	2/4 (50%)	1/5 (20%)	1/5 (20%)	4/14 (28.6%)	0.355	0.278
αCSP + TY12	3/4 (75%)	0/5 (0%)	2/5 (40%)	5/14 (35.7%)	0.211	0.586
αCSP + TY03	6/6 (100%)	4/5 (90%)	4/5 (80%)	14/16 (87.5%)	**0.000034**	**0.025**

Mice were injected with 100 µg/mouse of anti-CSP mAb (2F6), 300 µg/mouse of an anti-PyTRAP mAb or a combination of both 24h prior to challenge by 5 Py-infected mosquitos. Where only one mAb was injected, mice were also given non-specific mIgG to total 400 µg/mouse. Mice were tracked for 14 days for parasitemia by thin blood smear and those which remained parasite-free at day 14 were considered sterilely protected. Comparisons between groups were carried out via Barnard's exact test with the resulting p-value, as indicated.

Table 2. Combination of anti-PfCSP and anti-PfTRAP can improve sterile protection
from mosquito bite challenge.

	Sterile Protection				Comparison p value	
	Exp 1	Exp 2	Exp 3	Combined	vs. mIgG	vs. αCSP
150µg mIgG	0/7 (0%) [#]	0/7 (0%) [#]	0/5 (0%)	0/19 (0%)	–	**<0.0001**
50µg AKBR-4	0/6 (0%)	–	0/5 (0%)	0/11 (0%)	1	**0.0002**
50µg αCSP	5/7 (71%) [#]	5/8 (63%) [#]	5/8 (63%)	15/23 (65%)	**<0.0001**	–
150µg αCSP	–	3/5 (60%)	–	3/5 (60%)	**0.002**	0.88
50µg AKBR-4 + 50µg αCSP	–	6/7 (86%)	8/9 (89%)	14/16 (88%)	**<0.0001**	0.131

Mice were injected with indicated doses of either non-specific mIgG, anti-CSP mAb
CIS43, anti-PfTRAP mAb AKBR-4 or a combination of anti-CSP and AKBR-4 24h prior
to challenge with 5 Pf-infected mosquito bites. Mice were injected with human red blood
cells at day 5 and 6, and then blood sampled at days 7 and 9 to detect blood stage
parasitemia by qPCR. Number and percentages of mice protected across 3
independent experiments are shown. The p values for comparisons to either control
group (mIgG) or anti-CSP alone using Barnard's exact test are shown. [#] indicates
results previously reported in Kisalu *et al.* [49].

1   **Figures**

3   *Figure 1. TRAP domain organization and constructs used.*

4   Ectodomain and deletion constructs for PyTRAP and PfTRAP generated using the
5   domain boundaries (**A**) were recombinantly expressed and purified alongside the
6   control CSP ectodomain proteins (**B**).

Figure 2. Polyclonal antibodies to PyTRAP inhibit parasite invasion, traversal and in vivo

infection.

Mice were immunized three times with PyTRAP and PyCSP ectodomains. **A**) Immune
serum was used to verify binding to *Py* sporozoite via immunofluorescence. Shown are
fixed, permeabilized sporozoites labeled with a 1:800 dilution of polyclonal mouse
serum followed by anti-mouse IgG (red channel) and DAPI nuclear stain (blue). Immune
serum was then assessed for function in vitro for inhibition of invasion (**B**) and traversal
(**C**). In **B** and **C**, each data point is the average "% of pre-immune" invasion or traversal
for pooled serum from a cohort of 5 mice across independent assays (number shown
above each bar). Each bar indicates the mean "% of pre-immune" parasite invasion or
traversal with percent change from 100% (shown as dashed line) shown above.
Asterisks indicate a significant difference from 100% as determined by a two-tailed one-
sample *t*-test. **D**) Immunized mice were challenged by the bite of 15 *Py*GFPluc-infected
mosquitos and assessed for parasite liver burden by bioluminescent imaging. Each data
point represents an individual mouse with each color corresponding to an independent
immunization-challenge experiment (total number of animals shown above each bar).
Each data point is normalized to the mean luminescent signal from "naive" mice within

1   each challenge experiment while "control" mice are an additional group immunized with

2   HIV Env gp120 protein. Asterisks indicate significance as determined by ANOVA with

3   Kruskall-Wallis post-test. For **B-D**, * is $p \leq 0.05$; ** is $p \leq 0.01$; and *** is $p \leq 0.0001$.

Figure 3. Effects of PyTRAP monoclonal antibodies on parasite activity.

Each mAb was assessed for in vitro function of inhibition of invasion and traversal (**A**).
In each case, mean values of % inhibition (i.e., 100% – invasion or traversal value) from
the 100-µg/mL mAb concentrations (bar plots with these and additional conditions
shown in Suppl. Fig. 1) are represented on a color axis. Binding kinetics for each mAb
was measured by BLI and shown as kinetic maps (**B**) with gray dashed diagonal
contour lines labeled with the corresponding K_d values and symbols representing the
characterized epitopes for invasion (*left*) and traversal (*right*) inhibition. Higher-affinity
(i.e., those possessing lower K_d values) mAbs are closer to the upper-right corner of this

plot. Symbol color coding represents "strong" inhibition for mean values ≤ 50%, "mild"

inhibition for values ≤ 70% and "none" for mean values >70% observed at the 100-

µg/mL concentration. Red box highlights the region of the kinetic plots containing the

values for mAbs that showed strong inhibition in invasion and traversal assays.

Summarized sterile protection ratios following passive-transfer-challenge experiments

(**C**; number of animals in each group is shown below the corresponding bar, individual

values shown in Table 1).

Figure 4. Polyclonal antibodies to PfTRAP inhibit parasite invasion, traversal in vitro.

Mice were immunized three times with full-length PfTRAP and PfCSP. **A**) Immune
serum was used to verify binding to *Pf* sporozoite via immunofluorescence. Shown are
fixed, permeabilized sporozoites labeled with a 1:800 dilution of polyclonal mouse
serum followed by anti-mouse IgG (green channel), anti-PfCSP monoclonal antibody
2A10 (red channel, right image) and DAPI nuclear stain (blue, right image). Immune
serum was then assessed for function in vitro for inhibition of invasion (**B**) and traversal
(**C**). In **B** and **C**, each data point is the average % of Pre-immune invasion or traversal
from technical triplicates in independent experiments. Each bar indicates the mean % of
Pre-immune parasite invasion or traversal ±SEM with exact mean above. Asterisks
indicate a significant difference from 100% as determined by one-way t test where * is
p≤0.05; ** is p≤0.01; and *** is p≤0.0001.

Figure 5. Monoclonal antibodies to PfTRAP inhibit parasite invasion, traversal in vitro.

Each mAb was assessed for in vitro function of inhibition of invasion and traversal (**A**).
In each case, mean values of % inhibition (i.e., 100% – invasion or traversal value) from
the 100-µg/mL mAb concentrations (bar plots with these and additional conditions
shown in Suppl. Fig. 3) are represented on a color axis. Binding kinetics for each mAb
was measured by BLI and shown as kinetic maps (**B**) with gray dashed diagonal
contour lines labeled with the corresponding K_d values and symbols representing the
characterized epitopes for invasion (*left*) and traversal (*right*) inhibition. Higher-affinity
(i.e., those possessing lower K_d values) mAbs are closer to the upper-right corner of this
plot. Symbol color coding represents "strong" inhibition for mean values ≤ 50%, "mild"
inhibition for values ≤ 70% and "none" for mean values >70% observed at the 100-
µg/mL concentration. Summarized sterile protection breakdowns following passive-
transfer-challenge experiments (**C**; number of animals in each group is shown below the
corresponding bar, individual values shown in Table 2).