Anti-TRAP/SSP2 monoclonal antibodies can inhibit sporozoite infection and

enhance protection of anti-CSP monoclonal antibodies

Brandon K. Wilder*[1][†], Vladimir Vigdorovich*[1], Sara Carbonetti[1], Nana Minkah[1], Nina

Hertoghs[1], Andrew Raappana[1], Hayley Cardamone[1], Brian G. Oliver[1], Olesya

Trakhimets[1], Sudhir Kumar[1], Nicholas Dambrauskas[1], Silvia A. Arredondo[1], Nelly

Camargo[1], Stefan H.I. Kappe*[1,2,3] and D. Noah Sather*[1,2,3]

[1]Center for Global Infectious Disease Research, Seattle Children's Research Institute,

Seattle, WA, USA

[2]Department of Pediatrics, University of Washington, Seattle, WA, USA

[3]Department of Global Health, University of Washington, Seattle, WA, USA

[†]Present address: Vaccine and Gene Therapy Institute, Oregon Health & Science

University, Beaverton, Oregon, 97006, USA

*Equal contribution

Correspondence: D. Noah Sather: noah.sather@seattlechildrens.org and

17   Stefan H.I. Kappe: stefan.kappe@seattlechildrens.org

Running title: Anti-TRAP mAbs enhance protection of anti-CSP mAbs

<u>Abstract</u>

Vaccine-induced sterilizing protection from infection by *Plasmodium* parasites, the

pathogens that cause malaria, will be essential in the fight against malaria as it would

prevent both malaria-related disease and transmission. Stopping the relatively small

number of parasites injected by the mosquito before they can migrate from the skin to

the liver is an attractive means to this goal. Antibody-eliciting vaccines have been used

to pursue this objective by targeting the major parasite surface protein present during

this stage, the circumsporozoite protein (CSP). While CSP-based vaccines have

recently had encouraging success in disease reduction, this was only achieved with

extremely high antibody titers and appeared less effective for a complete block of

infection (i.e. sterile protection). While such disease reduction is important, these and

other results indicate that strategies focusing on CSP alone may not achieve the high

levels of sterile protection needed for malaria eradication. Here, we show that

monoclonal antibodies (mAbs) recognizing another sporozoite protein, TRAP/SSP2,

exhibit a range of inhibitory activity and that these mAbs can augment CSP-based

protection despite conferring no sterile protection on their own. Therefore, pursuing a

multivalent subunit vaccine immunization is a promising strategy for improving infection-

blocking malaria vaccines.

<u>Introduction</u>

The last few years have marked a disheartening milestone as the first period in a

generation without a reduction in the global burden of malaria [1]. The interventions that

have provided much of the previous progress, such as insecticide-treated bednets and

large-scale treatment programs, are highly susceptible to interruptions due to political or

economic instability. This was starkly illustrated by the resurgence of malaria in

Venezuela in recent years after near-elimination; and in 2020, more globally, due to

interruptions in eradication efforts during the COVID-19 pandemic [1]. Therefore, it is

likely that durable, infection-blocking interventions (e.g. vaccines, long-lasting mAbs or

chemoprophylactics) will be required to drive malaria to elimination.

Developing such an intervention is hampered by the complex life cycle of the parasite,

which begins when an infected mosquito injects tens to hundreds of the "sporozoite"

forms of the parasite into the dermis [2]. From here, sporozoites actively traverse through

multiple cell types in search of an endothelial cell through which they will gain access to

the blood [3]. Upon entering the bloodstream, a sporozoite is carried to the liver within

minutes, where it traverses multiple cell types in the liver parenchyma and eventually

establishes infection in a hepatocyte [4]. Following ~7–10 days of development and

genome replication (~2 days in rodent malaria models), each successful liver stage

releases 30,000–50,000 "merozoites" that cyclically infect, replicate within and lyse red

blood cells [5,6]. It is only during this blood stage of infection when symptomatic disease

occurs, and is also where a subset of sexually differentiated parasite forms can be

picked up by a new mosquito host to continue the transmission cycle. Each step in the

infection cycle presents opportunities for intervention, although vaccines targeting the

"pre-erythrocytic" stages in the skin and liver have yielded the most promising results [7] .

The most advanced pre-erythrocytic vaccine is RTS,S [8]—an antibody-eliciting subunit

vaccine targeting the major sporozoite surface protein circumsporozoite protein (CSP),

which has been recently recommended by the WHO [1,9]. Vaccines based on attenuated

live sporozoites that arrest in the liver and function by a combination of T cells and

antibodies have also demonstrated robust protection [10]. Unfortunately, despite

significant efficacy from both approaches in controlled human malaria infection (CHMI)

studies in malaria-naive volunteers, both vaccines have markedly reduced efficacy in

field trials and have not met the goals of 75% protection against clinical disease for one

year as expressed by the WHO [11]. Recent encouraging Phase II results with the R21

CSP particle in Matrix-M adjuvant do meet this goal [12]. However, protection with R21

appears dependent on high antibody titers, which would require yearly boosters that are

vulnerable to interruptions, and protection is less robust against *infection*. If a vaccine or

other intervention (e.g. a mAb or an injectable chemoprophylactic) is to be used as a

tool to achieve malaria eradication, it will likely need at least 80% efficacy against

infection to have a significant and sustained impact on transmission [13–15]. These

realities highlight the significant room for improvement in both T cell and antibody-

eliciting vaccines, with the latter more amenable to iterative improvement due to

available *in vitro* and *in vivo* preclinical assays [16–19].

Of the hundreds of proteins expressed at the sporozoite stage, at least 47 are surface-

exposed [20–22] and therefore potentially accessible to antibodies. However, few of these

proteins have been rigorously investigated for their use in antibody-eliciting vaccines [23].

In addition to CSP, the thrombospondin-related anonymous protein (TRAP, also known

as sporozoite surface protein 2 or SSP2) has been pursued as a vaccine candidate.

Similar to CSP, TRAP is essential for sporozoite infectivity [24,25], antibodies against it

correlate with protection [26,27] and the protein is abundant [21] during the skin stage when

parasites are particularly susceptible to antibody mediated inhibition. The TRAP

ectodomain consists of 3 main domains: a von Willebrand factor A-like domain (vWA),

the thrombospondin repeat (TSR) domain and a repeat domain [28]. The most advanced

TRAP vaccine candidate is an adenovirus/MVA-vectored vaccine eliciting strong T cell

responses that has had low or mixed efficacy results in CHMI trials [29,30] and field trials [31]

but has been improved in mice following targeting of the T cell response to the liver [32].

Antibody function in experiments involving immunization with TRAP-derived peptides

have yielded mixed results ranging from significant sporozoite inhibition in vitro [33] to no

protection in vivo [34] . A combination protein TRAP/RTS,S immunization failed to show

significant protection in a clinical trial [35], while a fusion-protein approach using TRAP

and CSP resulted in complete protection in a small mouse study [36]. These results using

TRAP alone or in combination with CSP are difficult to interpret due to the diversity of

vaccine platforms used, the possibility of immune interference in studies combining

platforms, and the unclear dominance of roles for antibodies and T cells in protection [37].

Whether a more targeted TRAP antibody response could contribute to protection either

alone or in combination with CSP remains poorly defined.

Here, we used both active immunization and passive transfer of mAbs raised against

either *Plasmodium yoelii* (rodent malaria) or *Plasmodium falciparum* (human malaria)

TRAP to more directly explore the potential efficacy of anti-TRAP antibodies. We found

that anti-TRAP antibodies modestly prevent liver infection in a manner dependent on

the TRAP domain recognized. Importantly, we also demonstrate a proof-of-concept that

anti-TRAP antibodies with minimal protective capacity of their own can augment anti-

CSP antibodies, providing additive protection that raises their protective efficacy above

80% sterile protection. Together, these findings argue for further investigation of

rationally designed multi-antigen, antibody-eliciting malaria vaccines or mAb

prophylactics that target multiple antigens and might include CSP as well as non-CSP

targets such as TRAP.

Results

PyTRAP polyclonal antibodies can prevent parasite infection of hepatocytes in vitro and

in vivo.

To elicit potentially functional anti-TRAP antibodies, we generated full-length

ectodomains and fragments of both rodent (*P. yoelii*) and human (*P. falciparum*) malaria

TRAP proteins (**Fig. 1A; Suppl. Table 1**) and verified their purity (**Fig. 1B**). Serum from

mice immunized with the rodent malaria *P. yoelii* TRAP ectodomain (PyTRAP)

recognized *Py* sporozoites by immunofluorescence in a pattern consistent with

micronemal localization, indicating the antigenic fidelity of the recombinant protein (**Fig.**

2A). We further tested this serum in an inhibition of sporozoite cell traversal and

invasion (ISTI) assay. Compared to control serum, anti-PyTRAP serum was able to

modestly but significantly (p=0.028) reduce sporozoite invasion of Hepa1-6 hepatoma

cells *in vitro* at a level similar to serum from mice immunized with the recombinant

PyCSP ectodomain, although the latter failed to reach significance (p=0.106) (**Fig. 2B**).

In contrast, sporozoite traversal of Hepa1-6 cells was not affected by anti-PyTRAP

serum (p=0.125), whereas anti-PyCSP serum did significantly reduce traversal

(p=0.0057) (**Fig. 2C**). The known inhibitory anti-PyCSP mAb 2F6 [38,39] reduced both

inhibition and traversal in this assay, as expected (**Fig. 2B and C**).

PyTRAP-immunized mice were then challenged with *Py* sporozoites via mosquito bite to

determine if these antibodies could function *in vivo* to reduce liver infection. We utilized

a PyGFPluc parasite, which expresses luciferase, enabling the measurement of liver

stage parasite burden by bioluminescence imaging. Mice immunized with a non-specific

control protein (Env) showed no reduction in parasite liver stage burden following

challenge compared with naive mice. In contrast, mice immunized with the PyTRAP

ectodomain showed a significant 62% reduction of parasite liver stage burden. Mice

immunized with PyCSP ectodomain showed a 91% reduction relative to naive controls

(**Fig. 2D**). Together, these data indicate that anti-PyTRAP antibodies can function *in*

vitro and *in vivo* to reduce parasite infection of hepatocytes.

PyTRAP mAbs display a diverse array of functions in vitro and can provide additive

protection to anti-CSP antibodies in vivo.

Serum polyclonal antibodies, as studied above, are a mixture of many antibody

specificities, making it difficult to characterize the relative contribution to functional

activity of responses directed at different domains. To enable such a characterization of

the repertoire of PyTRAP-elicited antibodies, we produced a panel of 15 mAbs. When

tested in ISTI at 10, 50 and 100 µg/mL, 12 of these mAbs significantly inhibited invasion

or cell traversal at one or more concentrations, with mAbs TY03 and TY11 showing the

most consistent and potent inhibition (**Fig. 3A**, **Suppl. Fig. 1**).

Overall, the mAbs demonstrated a wide range of binding affinities to recombinant

PyTRAP (**Fig. 3B**, **Suppl. Table 2**) and recognized epitopes in the vWA, TSR and

repeat regions (**Suppl. Table 3**), thus covering the entire protein ectodomain. Among

the 15 mAbs recovered, 10 mAbs bound to the vWA domain. Six of these (TY02, TY05,

TY06, TY10, TY11, TY20) shared variable-segment assignments for both heavy and

light chains, had closely related complementarity-determining-region (CDR) sequences

and had 88.4–96.7% and 93.9–96.9% sequence identity in the variable-region

sequences of their heavy and light chain, respectively (**Suppl. Table 4 and Suppl. Fig.**

2A,B). As expected, these antibodies were functionally similar in that they bound

specifically to the vWA domain (**Suppl. Table 3**), clustered in the same epitope bin

(**Suppl. Fig. 3A,B**) and inhibited sporozoite infection *in vitro* (**Fig. 3A, Suppl. Fig. 1**).

Two mAbs specifically recognized the TSR domain, and the remaining three mAbs

bound epitopes in the repeat region (**Suppl. Table 3**). These non-vWA antibodies

showed only modest or no sporozoite inhibition of infection *in vitro* (**Fig. 3A**). Binding

interference experiments allowed for the assignment of several distinct epitope bins

(**Suppl. Fig. 3A,B**) in addition to the one largely formed by the aforementioned group of

mAbs sharing high sequence identity. This likely indicates that the mAbs in our panel

bind several distant epitopes on PyTRAP. In addition, this panel of mAbs showed a

wide range of binding kinetics, with all strongly inhibitory mAbs having a k_{on} of $>10^5$ M$^-$

^{1}s^{-1} and a k_{dis} of $<10^{-2}$ s^{-1} (**Figure 3B**, note the red box, **Suppl. Table 4**). Together,

these data demonstrate that, similar to polyclonal antibodies, anti-PyTRAP mAbs can

mediate anti-parasitic function *in vitro*, and that inhibitory function likely depends on fast

and stable binding to the vWA domain. However, within the vWA domain some epitopes

show higher correlation between binding and blocking of infection compared to others.

We next wanted to determine whether an anti-PyTRAP mAb could provide sterilizing

protection *in vivo* on its own or in combination with an anti-CSP mAb. For this, we chose

three vWA domain-binding anti-PyTRAP mAbs from distinct epitope bins: TY03 and

TY11, which were the top-performing mAbs in ISTI, and TY12, which failed to

demonstrate efficacy in ISTI. Neither the anti-PyTRAP mAbs nor the anti-CSP mAb

showed significant binding to the mismatched Ag in vitro (**Suppl. Fig. 4A, B**), indicating

target specificity. The anti-PyTRAP mAbs were given at 300 µg/mouse (~15 mg/kg)

alone or with a partially protective dose of 100 µg/mouse (~5 mg/kg) of anti-PyCSP

mAb 2F6 prior to mosquito bite challenge [38]. As shown in **Fig. 3C** and **Table 1**, mice

administered anti-PyCSP mAb 2F6 showed significant sterile protection, with 9/18

(50%) remaining blood stage parasitemia-free, compared to 2/15 (13.3%) for mice

receiving non-specific murine IgG (p=0.032; this value was not corrected for multiple

comparisons due to small sample size and a small number of predefined comparisons

being made). Neither TY11 nor TY12 showed any protection (2/13 or 15.4% non-

infected) despite TY11 demonstrating the most robust inhibition *in vitro*. Administration

of the mAb TY03 resulted in 7/16 mice (43.7%) remaining parasitemia-free, not

reaching statistical significance. When combined with the anti-CSP mAb, only the

addition of TY03 afforded significant sterile protection (87.5% or 14/16 mice) over the

control group (p<0.001), which, importantly, was a significant improvement over

protection observed with anti-PyCSP mAb alone (p=0.025; again not corrected for

multiple comparisons as above). Together these data indicate that while *in vitro* testing

of mAbs can be useful for identifying non-functional mAbs (e.g. TY12), they should be

validated *in vivo* for function. Importantly, these data provide proof of concept that non-

CSP antibodies can provide additive protection to anti-CSP antibodies.

Antibodies targeting the human malaria parasite P. falciparum TRAP can function

against sporozoite invasion of hepatocytes.

We next wanted to determine if antibodies directed against TRAP/SSP2 from the

human malaria parasite, *P. falciparum*, could also function to prevent sporozoite

infection. Serum from mice immunized with the ectodomain of *P. falciparum* TRAP

(PfTRAP) was able to recognize *Pf* sporozoites in IFA (**Fig. 4A**) and demonstrated

consistent inhibition of *Pf* sporozoite invasion *in vitro* at a level similar to serum from

mice immunized with the ectodomain of *P. falciparum* CSP (PfCSP) (**Fig. 4B**). Inhibition

of sporozoite traversal *in vitro* was more modest as compared to anti-PfCSP polyclonal

serum (**Fig. 4C**). The known inhibitory anti-PfCSP mAb 2A10 [40] demonstrated robust

inhibition of both invasion and traversal (**Fig. 4B and C**).

Using a similar approach to the anti-PyTRAP work described above, we isolated 7 anti-

PfTRAP mAbs from immunized mice. Of these, 5 mAbs recognized the vWA domain

with AKBR-3, AKBR-4 and AKBR-6 likely recognizing adjacent epitopes (**Suppl. Fig.**

3C,D), and 2 mAbs recognized the TSR domains (**Suppl. Table 3**). In contrast to the

high proportion of functional anti-PyTRAP mAbs (12 of 15), only 2 of 7 anti-PfTRAP

mAbs, both recognizing the vWA domain, showed any sporozoite-inhibitory function *in*

vitro: AKBR-4 and AKBR-10. Further, only AKBR-4 demonstrated significant inhibition of

both invasion and traversal (**Fig. 5A, Suppl. Fig. 5**), despite having unremarkable

binding properties with the PfTRAP ectodomain (**Fig. 5B**). Surprisingly, mAb AKBR-7,

which had the best binding properties of the set (K_d ~0.15±0.04 nM, **Suppl. Table 2**),

demonstrated the worst inhibitory properties (**Fig. 5B**). Similar to the case with the anti-

PyTRAP mAb panel described above, our data suggest that the PfTRAP vWA domain

contains epitopes exposing vulnerability to inhibition, however lack of mAbs that

strongly bind other portions of PfTRAP make it difficult to discount the roles that these

domains may play in inhibition *in vivo*.

A vWA-directed anti-PfTRAP mAb increases the protection afforded by a protective

CSP mAb.

Because *Pf* sporozoites do not infect murine livers, the only means to test the activity of

anti-*Pf* antibodies against sporozoite infection *in vivo* is by either challenging passively

or actively immunized wild-type mice with transgenic rodent parasites expressing the *Pf*

proteins of interest [41–43] or by passive immunization of immune-deficient humanized liver

mice (FRGhuHep) that can be challenged with wild type *Pf* sporozoites [17]. We chose to

utilize the latter as it is an established model of antibody-mediated protection against *Pf*

infection [17,44–49] and allows testing of any future combination of anti-*Pf* antibodies

without the need for generating combinatorial transgenic parasites. In this model,

humanized-liver mice receive a passive transfer of antibodies and are then infected with

Pf sporozoites via mosquito bite. Six days later, mice are injected with human red blood

cells, which can then be infected by merozoites emerging from the liver, and blood

stage infection can be quantified by qRT-PCR on days 7 and 9. In this model, detection

of parasites by qRT-PCR on either days 7 or 9 has proven to be a stringent and

sensitive means of detecting the presence of blood stage parasites [17,50,51] . Therefore,

we define sterile protection in this model as the absence of parasites in the blood above

the limit of detection at either day 7 or 9.

Using this method, we tested the ability of the anti-PfTRAP mAb AKBR-4 to provide

sterile protection against *Pf* mosquito-bite infection alone or in combination with a

partially-protective anti-PfCSP mAb CIS43 [45]. Neither the anti-PfTRAP mAb nor the anti-

CSP mAb showed significant binding to the mismatched Ag in vitro (**Suppl. Fig. 4C, D**),

indicating target specificity. We chose a dose of 50 µg/mouse (~2.5 mg/kg) for each

mAb as this provides partial protection with an anti-PfCSP mAb [45] and gives a serum

concentration of ~10 µg/mL at the time of infection, which is achievable by both active

vaccination and passive transfer of long-lasting mAbs [52,53]. We previously conducted

passive administration, mosquito bite challenge in 2 independent experiments [45] , which

showed that 50 µg/mouse dose of anti-PfCSP mAb CIS43 was protective (5/7 and 5/8

protected in each experiment), compared to control mice (0/7 and 0/7 protected). To

avoid unnecessary repetition of FRGhuHep experiments, we included those cohorts in

our overall analysis of mAbs in this study and conducted a third independent experiment

with the control and 50 µg/mouse dose of anti-PfCSP mAb CIS43 groups in each. In

these experiments, 50 µg/mouse mAb CIS43 yielded a total of 15/23 protected (65%),

which was significant compared to 0/19 of control mice protected (0%, p<0.0001; **Table**

2, **Fig. 5C**). To determine if the protection afforded by CIS43 would scale linearly with

dose and possibly reach 100%, we included a group of 5 FRGhuHep mice in a single

experiment, in which the dose was increased 3-fold to 150 µg/mouse. This resulted in 3

of 5 mice protected (60%; p=0.002 over control) but was not significantly different from

the groups that received 50 µg/mouse.

On its own, passive administration of 50 µg/mouse of AKBR-4 failed to provide any

sterile protection over two of these experiments (0/11, 0%). Yet, when 50 µg/mouse of

AKBR-4 was combined with 50 µg/mouse of the anti-PfCSP mAb (100 µg mAb/mouse

total), 14/16 (88%; p<0.0001 over control) mice were sterilely protected over two

independent experiments. The improvement afforded by the AKBR-4/anti-PfCSP mAb

combination over the efficacy of the anti-PfCSP mAb alone trended toward, but did not

reach statistical significance at this group size (p=0.131). Together, these results

provide the proof of concept that antibodies directed against PfTRAP can reduce *Pf*

sporozoite cell traversal and invasion of hepatocytes *in vitro*, and potentially enhance

the protection of anti-CSP mAbs when used in combination with the latter.

<u>Discussion</u>

Studies examining CSP-elicited antibody responses have shown that within a polyclonal

antibody population only a subset are highly potent antibody clones, and their

distinguishing binding properties can be quite nuanced [39,44,45,54–58]. Understanding the

characteristics associated with protection is crucial for the development of superior mAb

products and vaccine immunogens, yet such studies have not been previously

performed for TRAP or other non-CSP pre-erythrocytic antibody targets. Here, we show

that the polyclonal antibody response to PyTRAP ectodomain can substantially reduce

parasite infection of hepatocytes *in vitro* and use mAbs to conclude that this effect is

likely driven by vWA and TSR-specific antibodies. These findings are in line with some

previous work using antibodies against TRAP protein fragments [33], yet they contrast

other observations that failed to see significant inhibition [34]. Our data with PfTRAP were

more limited but the only mAb that was functional *in vitro* also recognized the vWA

domain. Taken together, our data with polyclonal and monoclonal antibodies clearly

demonstrate that TRAP is a viable antibody target, and that its vWA domain contains

sites of vulnerability.

Critical for any vaccine or mAb product that can be used for malaria eradication will be

achieving high levels of sterile protection at sustainable antibody levels. Experience with

RTS,S—which elicits extremely high peak levels of anti-CSP antibodies—as well as

published data describing the activity of potent anti-CSP mAbs in animal models

suggest that increasing anti-CSP antibody titers can increase protection [45,55,59]. The first

CHMI trial using passive transfer of the anti-PfCSP mAb CIS43 (also used in this study)

showed that mAbs can provide sterilizing protection against *P. falciparum* mosquito-bite

infection at serum concentrations between ~50–500 µg/mL [60]. However, maintenance

of such high antibody titers for over a year may not be sustainable for active or passive

immunization strategies. As an alternative to frequent vaccine boosting or mAb

injections to sustain high titers, it may be possible to achieve high levels of protection at

lower antibody titers using multivalent vaccination or multiple mAbs recognizing distinct

protein targets. However, there have been no studies to date directly addressing this

question, which is best examined using passive transfer of antibodies followed by

mosquito bite challenge, as done here.

Our data in the *Py* model show that an anti-PyTRAP mAb, offering no significant

protection by itself, can improve protection against mosquito-bite infection of a partially-

protective anti-CSP mAb regimen. Our experiments using *Pf* mosquito-bite challenge in

FRGhuHep mice, which received a combination of anti-PfCSP and anti-PfTRAP mAbs,

did not show a statistically significant improvement over anti-PfCSP mAb treatment

alone. However, the fact that this combination was the only regimen to deliver strong

protection in repeated experiments, as well as the strong statistical trend observed, offer

support for such an approach against *P. falciparum*. Importantly, the 88% sterile

protection was achieved using a low total dose of mAb (100 µg/mouse or ~5 mg/kg).

This total dose of 100 µg/mouse (50 µg/mouse each of anti-PfCSP and anti-PfTRAP

mAb) is expected to give a total circulating mAb concentration of ~20 µg/mL [45] —a level

that can be achieved for ~36 weeks with a single 20 mg/kg injection of long-lasting

mAbs [52,60,61] or ~4 years via active vaccination [62] . Although it remains to be seen how

accurately these animal models translate to the clinic, these data suggest that reaching

the 80% sterile protection threshold needed for vaccines [11] or injectable anti-malarials [15]

that can be used as eradication tools may be achieved by targeting multiple proteins

rather than by increasing the concentration of antibodies recognizing CSP alone.

Intriguingly, our data showed that some anti-TRAP mAbs, when combined with anti-

CSP mAbs, resulted in enhanced protection despite providing no statistically significant

sterile protection on their own. These observations may be explained by the fact that the

sterile protection readout requires the prevention of all parasites from successfully

infecting the liver, effectively introducing a threshold effect. Therefore, it is possible that

a weakly inhibitory mAb would have a more pronounced effect in combination with a

partially protective regimen (e.g., that of a suboptimal dose of an anti-CSP mAb) than

would be predicted by single-mAb experiments when using sterile protection as a

readout. Additional studies clarifying the additive vs. synergistic nature of this or any

multivalent approach will be needed to determine the utility of combining CSP with other

immunogens, but will require large group sizes and experiments designed specifically to

test such hypotheses.

In summary, we present a proof-of-concept that antibodies targeting TRAP can

contribute to sterile protection when used in combination with anti-CSP antibodies.

These findings support vaccine and mAb strategies involving multiple *Plasmodium* pre-

erythrocytic-stage antigens, and argue that efforts to develop a long-lasting, infection-

blocking malaria intervention would greatly benefit from identifying non-CSP antibody

targets that can enhance CSP-elicited protection. Although such a multivalent approach

can be achieved with mAbs, it is currently limited by cost [63]. Active vaccination with

multiple antigens has been hampered by challenges of generating and combining

multiple protein-in-adjuvant formulations, although this may be more easily achieved by

the use of mRNA-based vaccines, which have proven adept as a multi-antigen vaccine

platform in preclinical studies [64,65] . Our data, which demonstrate that enhanced

protection over CSP-only strategies is possible by way of multivalent subunit

vaccination or delivery of mAbs, provide the impetus to pursue such strategies in

preclinical studies that better define additive protection and identify additional targets.

Materials and Methods

Recombinant protein production

Recombinant proteins were produced in transiently transfected suspension culture of

FreeStyle 293 cells (Thermo Fisher Scientific, Waltham, MA, USA), as previously

described [66]. Briefly, codon-optimized constructs encoding the ectodomains or deletion

constructs of *Plasmodium falciparum* TRAP (PfTRAP), *Plasmodium yoelii* CSP (PyCSP)

and *Plasmodium yoelii* TRAP (PyTRAP) were generated as fusions to the 8xHis and

AviTag [67] sequences (**Suppl. Table 1**). Following transfection using the high-density

PEI method [68] and the subsequent 5-day incubation, cells were removed by

centrifugation and the culture supernatants were supplemented with NaCl (+350 mM)

and sodium azide (0.02%). Treated culture supernatants were passed by gravity

through NiNTA agarose, washed with Wash Buffer (10 mM Tris-HCl, pH 8, 300 mM

NaCl, 10 mM imidazole), and eluted with Elution Buffer (10 mM Tris-HCl, pH 7.4, 300

mM NaCl, 200 mM imidazole). Further purification was performed by size-exclusion

chromatography using a calibrated Superdex 200 (10/600) column (Cytiva,

Marlborough, MA, USA). The HIV Env gp120 control protein was produced, as

previously described [69]. When required, site-specific biotinylation with BirA and buffer-

exchange by gel filtration was performed, as previously described [66].

19
Antibody cloning and production

Antibodies were cloned and produced, as previously described [66]. Briefly, ectodomain

PfTRAP and PyTRAP constructs were used as immunogens, and their biotinylated

versions were used to isolate antigen-specific B cells by flow cytometry (see sample

gating strategy in **Suppl. Fig. 6**). Following culture previously described medium [66]

modified by the addition of 1.5 µM CpG (ODN-1826) (Integrated DNA Technologies,

Coralville, IA, USA), wells containing B cells producing antigen-binding IgG were

identified by ELISA, immunoglobulin-encoding transcripts were amplified by RT-PCR

and used for the generation of heavy- and light-chain constructs for recombinant mAb

expression. The sequences were annotated using IgBLAST [70].

To express recombinant mAbs, the plasmid DNA was used to transfect suspension

cultures of FreeStyle 293 cells (Thermo), as described above. After five days in culture,

cells were removed by centrifugation and the cultures were supplemented with NaCl

(+350 mM) and sodium azide (0.02%). Treated culture supernatants were passed by

gravity through Protein G resin equilibrated in Wash Buffer (10 mM HEPES, pH 7, 300

mM NaCl, 2 mM EDTA), washed with Wash Buffer, and eluted with 100 mM glycine, pH

2.7. Resulting eluates were buffer-exchanged by repeated centrifugal ultrafiltration with

HBS-E (10 mM HEPES, pH 7, 150 mM NaCl, 2 mM EDTA).

Binding properties of mAbs

Binding kinetics measurements were characterized using biolayer interferometry (BLI)

measurements on an Octet QK^e instrument (Sartorius, Göttingen, Germany), as

previously described [66]. Briefly, antibodies in culture supernatants were immobilized on

anti-Mouse IgG Fc Capture biosensors and allowed to associate with antigen serially

diluted (in the range of 1–1000 nM) in 10x Kinetics Buffer (10xKB: PBS + 0.1% Bovine

Serum Albumin, 0.02% Tween-20 and 0.05% sodium azide) followed by dissociation in

10x KB. Resulting sensorgram data was evaluated using ForteBio Data Analysis

software (version 7.0.1.5) to generate a fit to the 1:1 binding model and provide

estimates for the k_{on} and k_{dis} rate constants (see sample sensorgrams and fitted curves

in **Suppl. Fig. 7**).

The relative specificity of Ag recognition by the mAbs was assayed using biotinylated

Ags (30 µg/mL, with the exception of PfCSP, which was used at 10 µg/mL) immobilized

on streptavidin biosensors, and incubated with mAbs (50 µg/mL, with the exception of

anti-PfCSP, which was used at 10 µg/mL) diluted in 10xKB.

Epitope bins within anti-TRAP mAb panels were assigned based on the interference

patterns similar to previous work [71,72]. First, His-tagged PyTRAP or PfTRAP (30 µg/mL)

was immobilized on NiNTA biosensors in HBS-NPM buffer (20 mM HEPES, pH 7, 150

mM NaCl, 1 mM $MgCl_2$, 0.1 mg/mL Bovine Serum Albumin, 0.05% NaN_3, 0.02% Tween-

20). Interference for each pair of mAbs was assessed by binding the first mAb (mAb1)

(50 µg/mL, except for TY14 and TY15, which were used at 100 µg/mL) to saturation

before allowing the binding from the second mAb (mAb2) (50 µg/mL) to take place.

Magnitude of the signal for each mAb2 binding event was corrected by subtracting the

signal for the corresponding mAb1 binding step. Additionally, the signal for each mAb2

binding was collected in absence of pre-bound mAb1 (i.e., "blank" HBS-NPM buffer was

used in place of mAb1 solution) and used to normalize the corrected mAb2 signal.

Finally, the normalized mAb2 values were collected for each mAb1 and the resulting

interference pattern sets were used to calculate the Pearson correlation coefficients

using R (version 4.0.2) and plotted using the R packages pheatmap (1.0.12). Network

graphs were plotted using R package igraph (version 1.2.10) with edges connecting

pairs of nodes with a Pearson correlation coefficient >0.7; and clusters of

interconnected nodes are referred to as epitope bins.

Coarse epitope mapping by ELISA

Domain specificity of the mAbs was characterized by enzyme-linked immunosorbent

assay (ELISA) using TRAP ectodomain and fragments from PfTRAP and PyTRAP, as

previously described [66].

Sporozoite production

For rodent parasite (*P. yoelii*), female Swiss Webster mice for parasite maintenance

were purchased from Envigo (Livermore, CA, USA) and injected intraperitoneally (i.p.)

with blood stage PyGFPluc [73]. Three days later, gametocyte exflagellation was

confirmed and the infected mice were used to feed female *Anopheles stephensi*

mosquitoes. Fourteen to 16 days after the feed, salivary gland sporozoites were isolated

from the mosquitoes and used in mouse infections.

For human malaria (*P. falciparum*) experiments, infected *A. stephensi* mosquitoes were

produced, as previously described [74].

Animal studies ethics statement

All procedures involving animals were performed in adherence to protocols of the

Institutional Animal Care and Use Committee (IACUC) at the Seattle Children's

Research Institute.

Mouse active immunization and challenge

To generate polyclonal serum and a source of mouse mAbs, six to eight week-old

BALBc/J mice were purchased from Jackson Laboratories (Bar Harbor, ME, USA) and

injected intramuscularly three times at days 0, 14 and 38 using Adjuplex mixed with 20–

25 µg of target protein. Mice immunized with recombinant *Py* proteins were then

challenged by the bite of 15 PyGFPluc-infected mosquitos, as described [38]. Briefly, the

proportion of mosquitos infected with *Py* was determined by presence of midgut oocysts

on days 7–12. This proportion was used to prepare a cage with 15 infected mosquitoes

per animal (i.e. if 50% of mosquitoes had oocysts, 30 mosquitos/animal were used).

These mosquitos were then exposed to anesthetized mice for 10 minutes with lifting of

mice every minute to encourage active probing as opposed to blood feeding. Forty-two

hours later, parasite liver burden was assessed by bioluminescent imaging, as

previously described [38]. Mice were then immediately sacrificed and splenocytes

collected and cryopreserved for B cell isolation and mAb production.

Mice immunized with *Pf* proteins were immunized as above with the exception that mice

were additionally boosted with IV protein three days prior to sacrifice and collection and

cryopreservation of splenocytes.

For both, serum was collected from immunized mice by collecting whole blood in BD

microtainer serum tubes (Becton-Dickinson, Franklin Lakes, NJ, USA), allowing blood to

clot at room temperature for at least 30 minutes and then centrifuged according to

manufacturer's instructions to separate serum for storage and use in *in vitro* assays.

Sporozoite immunofluorescence microscopy

Py or *Pf* sporozoites were stained as previously described [75]. Briefly, freshly dissected

Py or *Pf* sporozoites were fixed with 4% PFA and air-dried onto glass slides overnight.

These were then permeabilized with 0.1% Triton X-100 and stained with polyclonal

(serum at 1:100-1:800 dilution) or monoclonal (10 μg/mL) antibodies. Sporozoites were

identified by co-staining with anti-CSP mAbs as well as DAPI for nuclear localization.

Images were acquired using an Olympus IX-70 DeltaVision deconvolution microscope

at 100X magnification.

In vitro inhibition of sporozoite traversal and invasion (ISTI)

In vitro ISTI was performed, as previously described for both *Py* and *Pf* [76]. Briefly,

freshly-dissected sporozoites were added to hepatoma cells (Hepa1-6 for *Py* and HC-04

for *Pf*) plated a day prior in 96-well plates in the presence of antibodies and FITC-

dextran in technical duplicates or triplicates. After 90 minutes, cells were fixed,

permeabilized and stained with Alexa Fluor 647-labeled anti-CSP mAbs and analyzed

by flow cytometry. Invaded cells were identified by the presence of CSP and traversed

cells by the uptake of FITC-dextran with gating set to uninfected, stained wells. Within

each experimental replicate, antibody-treated wells were normalized to the invasion and

traversal of wells treated with pre-immune serum or non-specific mouse IgG, which was

set to 100%.

Anti-Py mAb passive transfer and challenge

Six to eight week-old BALBc/J mice were intravenously injected with indicated doses of

mAbs 24 hours prior to challenge by bite of 5 PyGFPluc-infected mosquitos following

the same methods as described above for "Mouse active immunization and challenge".

Mice were followed up for infection by Giemsa-stained thin blood smear every other day

from days 3–14 for identification of blood stage parasites. Mice in which we failed to

identify parasites in 40,000 red blood cells over the entire period were considered

negative and sterilely protected. Control mice were administered non-specific polyclonal

mouse IgG at a dose equivalent to the highest dose in experimental groups.

anti-Pf mAb passive transfer in FRG humanized liver mice

Mice repopulated with human hepatocytes (FRGhuHep) were purchased from Yecuris,

Inc. (Tualatin, OR, USA) and infected with *Pf* via mosquito bite, as previously described

[17,45]. Briefly, indicated doses of mAb were intravenously injected into mice 24 hours

prior to challenge by the bite of 5 *Pf*-infected mosquitos using the same criteria and

methods as above. On day 6 post-infection, mice were intravenously injected with 400

µL of human red blood cells. On days 7 and 9 post-infection, 100 µL of peripheral blood

was collected, immediately added to Nuclisens Buffer (bioMerieux, Inc., Durham, NC,

USA) for lysis and storage. Parasite presence was quantified by qRT-PCR for *Pf* 18s

rRNA, as previously described [77]. Any mouse with a Ct value above the "no template"

control at either day was considered positive for parasitemia.

Statistics

Statistical analyses and plotting were carried out in Prism (version 9.2.0) (GraphPad

Software, San Diego, CA, USA) or in R (version 4.0.2) using packages Exact (version

2.1), ggpubr (version 0.4.0), ggstatsplot (version 0.7.2). Statistical tests and outcomes

are noted in the figure legend for each figure. For all tests, a p-value of <0.05 was

considered significant and values not specifically labeled were above this threshold.

Data availability

DNA sequences encoding the mAbs described here have been deposited in GenBank

(accession numbers OK484322–OK484365).

Acknowledgements

We would like to thank the vivarium staff at Seattle Children's Research Institute for their support of animal studies, and Weldon DeBusk for his assistance with the flow cytometry experiments. We would like to thank Dr. Sean C. Murphy of the University of Washington for performing the *Pf* qPCR analysis. Additionally, we would like to thank Dr. Paul T. Edlefsen of the Fred Hutch Cancer Research Center for the helpful discussions of statistical analysis and Drs. Nevile Kisalu and Robert Seder of the NIH VRC for their provision of mAb CIS43. This study was funded by NIH R01 AI117234 to S.H.K. and D.N.S.

<u>Author contributions</u>

B.K.W. and V.V. contributed equally to this work.

Conceptualization and experimental design: B.K.W., V.V., S.H.I.K. and D.N.S.

Investigation: B.K.W., V.V., S.C., N.M., N.H., A.R., H.C, B.G.O., O.T., S.K., N.D., S.A.A.

and N.C.

Data analysis and visualization: B.K.W., V.V., N.H., N.M.

Writing — Original draft: B.K.W.

Writing — Review and editing: B.K.W., V.V., S.H.I.K. and D.N.S.

Resources: S.H.I.K. and D.N.S.

Supervision, Project Administration and Funding Acquisition: B.K.W., S.H.I.K., D.N.S.

1 **<u>Competing Interests statement</u>**

2 The authors declare no competing interests.

References

1. World Health Organization. *World malaria report 2021*. (World Health Organization, 2021).
2. Medica, D. L. & Sinnis, P. Quantitative dynamics of Plasmodium yoelii sporozoite transmission by infected anopheline mosquitoes. *Infect. Immun.* **73**, 4363–4369 (2005).
3. Ejigiri, I. & Sinnis, P. Plasmodium sporozoite-host interactions from the dermis to the hepatocyte. *Curr. Opin. Microbiol.* **12**, 401–407 (2009).
4. Vaughan, A. M. & Kappe, S. H. I. Malaria Parasite Liver Infection and Exoerythrocytic Biology. *Cold Spring Harb. Perspect. Med.* **7**, (2017).
5. Cowman, A. F., Healer, J., Marapana, D. & Marsh, K. Malaria: Biology and Disease. *Cell* **167**, 610–624 (2016).
6. Lindner, S. E., Miller, J. L. & Kappe, S. H. I. Malaria parasite pre-erythrocytic infection: preparation meets opportunity. *Cell. Microbiol.* **14**, 316–324 (2012).
7. Duffy, P. E. & Patrick Gorres, J. Malaria vaccines since 2000: progress, priorities, products. *NPJ Vaccines* **5**, 48 (2020).
8. Tinto, H. *et al.* Long-term incidence of severe malaria following RTS,S/AS01 vaccination in children and infants in Africa: an open-label 3-year extension study of a phase 3 randomised controlled trial. *Lancet Infect. Dis.* **19**, 821–832 (2019).
9. Vogel, G. WHO gives first malaria vaccine the green light. *Science* **374**, 245–246 (2021).
10. Epstein, J. E. *et al.* Protection against Plasmodium falciparum malaria by PfSPZ Vaccine. *JCI Insight* **2**, e89154 (2017).
11. Malaria Vaccine Funders Group. *Malaria Vaccine Technology Roadmap*. https://www.who.int/publications/m/item/malaria-vaccine-technology-roadmap (2013).
12. Datoo, M. S. *et al.* Efficacy of a low-dose candidate malaria vaccine, R21 in adjuvant Matrix-M, with seasonal administration to children in Burkina Faso: a randomised controlled trial. *Lancet* **397**, 1809–1818 (2021).
13. White, M. T., Verity, R., Churcher, T. S. & Ghani, A. C. Vaccine approaches to malaria control and elimination: Insights from mathematical models. *Vaccine* **33**, 7544–7550 (2015).
14. Penny, M. A., Camponovo, F., Chitnis, N., Smith, T. A. & Tanner, M. Future use-cases of vaccines in malaria control and elimination. *Parasite Epidemiol Control* **10**, e00145 (2020).
15. Macintyre, F. *et al.* Injectable anti-malarials revisited: discovery and development of new agents to protect against malaria. *Malar. J.* **17**, 402 (2018).
16. Roth, A. *et al.* A comprehensive model for assessment of liver stage therapies targeting Plasmodium vivax and Plasmodium falciparum. *Nat. Commun.* **9**, 1837 (2018).
17. Sack, B. K. *et al.* Humoral protection against mosquito bite-transmitted Plasmodium falciparum infection in humanized mice. *NPJ Vaccines* **2**, 27 (2017).
18. Steel, R. W. J. *et al.* An Opsonic Phagocytosis Assay for Plasmodium falciparum Sporozoites. *Clin. Vaccine Immunol.* **24**, (2017).
19. Boyle, M. J. *et al.* Human antibodies fix complement to inhibit Plasmodium falciparum invasion of erythrocytes and are associated with protection against malaria. *Immunity* **42**, 580–590 (2015).
20. Swearingen, K. E. *et al.* Proteogenomic analysis of the total and surface-exposed proteomes of Plasmodium vivax salivary gland sporozoites. *PLoS Negl. Trop. Dis.* **11**, e0005791 (2017).
21. Swearingen, K. E. *et al.* Interrogating the Plasmodium Sporozoite Surface: Identification of Surface-Exposed Proteins and Demonstration of Glycosylation on CSP and TRAP by Mass Spectrometry-Based Proteomics. *PLoS Pathog.* **12**, (2016).
22. Lindner, S. E. *et al.* Total and putative surface proteomics of malaria parasite salivary gland sporozoites. *Mol. Cell. Proteomics* **12**, 1127–1143 (2013).
23. Sack, B., Kappe, S. H. I. & Sather, D. N. Towards functional antibody-based vaccines to

prevent pre-erythrocytic malaria infection. *Expert Rev. Vaccines* **16**, 403–414 (2017).

24. Sultan, A. A. *et al.* TRAP is necessary for gliding motility and infectivity of plasmodium sporozoites. *Cell* **90**, 511–522 (1997).

25. Klug, D. *et al.* Evolutionarily distant I domains can functionally replace the essential ligand-binding domain of Plasmodium TRAP. *Elife* **9**, (2020).

26. John, C. C. *et al.* Correlation of high levels of antibodies to multiple pre-erythrocytic Plasmodium falciparum antigens and protection from infection. *Am. J. Trop. Med. Hyg.* **73**, 222–228 (2005).

27. Scarselli, E. *et al.* Analysis of the human antibody response to thrombospondin-related anonymous protein of Plasmodium falciparum. *Infect. Immun.* **61**, 3490–3495 (1993).

28. Dolo, A. *et al.* Thrombospondin related adhesive protein (TRAP), a potential malaria vaccine candidate. *Parassitologia* **41**, 425–428 (1999).

29. Hodgson, S. H. *et al.* Evaluation of the efficacy of ChAd63-MVA vectored vaccines expressing circumsporozoite protein and ME-TRAP against controlled human malaria infection in malaria-naive individuals. *J. Infect. Dis.* **211**, 1076–1086 (2015).

30. Ewer, K. J. *et al.* Protective CD8+ T-cell immunity to human malaria induced by chimpanzee adenovirus-MVA immunisation. *Nat. Commun.* **4**, 2836 (2013).

31. Ogwang, C. *et al.* Prime-boost vaccination with chimpanzee adenovirus and modified vaccinia Ankara encoding TRAP provides partial protection against Plasmodium falciparum infection in Kenyan adults. *Sci. Transl. Med.* **7**, 286re5 (2015).

32. Gola, A. *et al.* Prime and target immunization protects against liver-stage malaria in mice. *Sci. Transl. Med.* **10**, (2018).

33. Charoenvit, Y. *et al.* Development of two monoclonal antibodies against Plasmodium falciparum sporozoite surface protein 2 and mapping of B-cell epitopes. *Infect. Immun.* **65**, 3430–3437 (1997).

34. Gantt, S. *et al.* Antibodies against thrombospondin-related anonymous protein do not inhibit Plasmodium sporozoite infectivity in vivo. *Infect. Immun.* **68**, 3667–3673 (2000).

35. Kester, K. E. *et al.* Sequential Phase 1 and Phase 2 randomized, controlled trials of the safety, immunogenicity and efficacy of combined pre-erythrocytic vaccine antigens RTS,S and TRAP formulated with AS02 Adjuvant System in healthy, malaria naïve adults. *Vaccine* **32**, 6683–6691 (2014).

36. Lu, C. *et al.* Design and assessment of TRAP-CSP fusion antigens as effective malaria vaccines. *PLoS One* **15**, e0216260 (2020).

37. Rampling, T. *et al.* Safety and High Level Efficacy of the Combination Malaria Vaccine Regimen of RTS,S/AS01B With Chimpanzee Adenovirus 63 and Modified Vaccinia Ankara Vectored Vaccines Expressing ME-TRAP. *J. Infect. Dis.* **214**, 772–781 (2016).

38. Sack, B. K. *et al.* Model for in vivo assessment of humoral protection against malaria sporozoite challenge by passive transfer of monoclonal antibodies and immune serum. *Infect. Immun.* **82**, 808–817 (2014).

39. Vijayan, K. *et al.* Antibody interference by a non-neutralizing antibody abrogates humoral protection against Plasmodium yoelii liver stage. *Cell Rep.* **36**, 109489 (2021).

40. Deal, C. *et al.* Vectored antibody gene delivery protects against Plasmodium falciparum sporozoite challenge in mice. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 12528–12532 (2014).

41. Atcheson, E. *et al.* Tailoring a Plasmodium vivax Vaccine To Enhance Efficacy through a Combination of a CSP Virus-Like Particle and TRAP Viral Vectors. *Infect. Immun.* **86**, (2018).

42. Longley, R. J. *et al.* Assessment of the Plasmodium falciparum Preerythrocytic Antigen UIS3 as a Potential Candidate for a Malaria Vaccine. *Infect. Immun.* **85**, (2017).

43. Longley, R. J. *et al.* Comparative assessment of vaccine vectors encoding ten malaria antigens identifies two protective liver-stage candidates. *Sci. Rep.* **5**, 11820 (2015).

44. Tan, J. *et al.* A public antibody lineage that potently inhibits malaria infection through dual

binding to the circumsporozoite protein. *Nat. Med.* **24**, 401–407 (2018).

45. Kisalu, N. K. *et al.* A human monoclonal antibody prevents malaria infection by targeting a new site of vulnerability on the parasite. *Nat. Med.* **24**, 408–416 (2018).

46. Triller, G. *et al.* Natural Parasite Exposure Induces Protective Human Anti-Malarial Antibodies. *Immunity* **47**, 1197–1209.e10 (2017).

47. Kublin, J. G. *et al.* Complete attenuation of genetically engineered Plasmodium falciparum sporozoites in human subjects. *Sci. Transl. Med.* **9**, (2017).

48. Ishizuka, A. S. *et al.* Protection against malaria at 1 year and immune correlates following PfSPZ vaccination. *Nat. Med.* **22**, 614–623 (2016).

49. Foquet, L. *et al.* Vaccine-induced monoclonal antibodies targeting circumsporozoite protein prevent Plasmodium falciparum infection. *J. Clin. Invest.* **124**, 140–144 (2014).

50. Goswami, D. *et al.* A replication-competent late liver stage-attenuated human malaria parasite. *JCI Insight* **5**, (2020).

51. Foquet, L. *et al.* Plasmodium falciparum Liver Stage Infection and Transition to Stable Blood Stage Infection in Liver-Humanized and Blood-Humanized FRGN KO Mice Enables Testing of Blood Stage Inhibitory Antibodies (Reticulocyte-Binding Protein Homolog 5) In Vivo. *Front. Immunol.* **9**, 524 (2018).

52. Gaudinski, M. R. *et al.* Safety and pharmacokinetics of the Fc-modified HIV-1 human monoclonal antibody VRC01LS: A Phase 1 open-label clinical trial in healthy adults. *PLoS Med.* **15**, e1002493 (2018).

53. Ledgerwood, J. E. *et al.* Safety, pharmacokinetics and neutralization of the broadly neutralizing HIV-1 human monoclonal antibody VRC01 in healthy adults. *Clin. Exp. Immunol.* **182**, 289–301 (2015).

54. Alanine, D. G. W. *et al.* Human Antibodies that Slow Erythrocyte Invasion Potentiate Malaria-Neutralizing Antibodies. *Cell* **178**, 216–228.e21 (2019).

55. Wang, L. T. *et al.* A Potent Anti-Malarial Human Monoclonal Antibody Targets Circumsporozoite Protein Minor Repeats and Neutralizes Sporozoites in the Liver. *Immunity* (2020) doi:10.1016/j.immuni.2020.08.014.

56. Livingstone, M. C. *et al.* In vitro and in vivo inhibition of malaria parasite infection by monoclonal antibodies against Plasmodium falciparum circumsporozoite protein (CSP). *Sci. Rep.* **11**, 5318 (2021).

57. Douglas, A. D. *et al.* A defined mechanistic correlate of protection against Plasmodium falciparum malaria in non-human primates. *Nat. Commun.* **10**, 1953 (2019).

58. Scally, S. W. *et al.* Rare PfCSP C-terminal antibodies induced by live sporozoite vaccination are ineffective against malaria infection. *J. Exp. Med.* **215**, 63–75 (2018).

59. Kisalu, N. K. *et al.* Enhancing durability of CIS43 monoclonal antibody by Fc mutation or AAV delivery for malaria prevention. *JCI Insight* **6**, (2021).

60. Gaudinski, M. R. *et al.* A Monoclonal Antibody for Malaria Prevention. *N. Engl. J. Med.* **385**, 803–814 (2021).

61. Gaudinski, M. R. *et al.* Safety and pharmacokinetics of broadly neutralising human monoclonal antibody VRC07-523LS in healthy adults: a phase 1 dose-escalation clinical trial. *Lancet HIV* **6**, e667–e679 (2019).

62. Minassian, A. M. *et al.* Reduced blood-stage malaria growth and immune correlates in humans following RH5 vaccination. *Med (N Y)* **2**, 701–719.e19 (2021).

63. Pelfrene, E., Mura, M., Cavaleiro Sanches, A. & Cavaleri, M. Monoclonal antibodies as anti-infective products: a promising future? *Clin. Microbiol. Infect.* **25**, 60–64 (2019).

64. Chahal, J. S. *et al.* Dendrimer-RNA nanoparticles generate protective immunity against lethal Ebola, H1N1 influenza, and Toxoplasma gondii challenges with a single dose. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E4133–42 (2016).

65. Vogel, A. B. *et al.* Self-Amplifying RNA Vaccines Give Equivalent Protection against Influenza to mRNA Vaccines but at Much Lower Doses. *Mol. Ther.* **26**, 446–455 (2018).

66. Carbonetti, S. *et al.* A method for the isolation and characterization of functional murine monoclonal antibodies by single B cell cloning. *J. Immunol. Methods* **448**, 66–73 (2017).

67. Fairhead, M. & Howarth, M. Site-specific biotinylation of purified proteins using BirA. *Methods Mol. Biol.* **1266**, 171–184 (2015).

68. Backliwal, G., Hildinger, M., Hasija, V. & Wurm, F. M. High-density transfection with HEK-293 cells allows doubling of transient titers and removes need for a priori DNA complex formation with PEI. *Biotechnol. Bioeng.* **99**, 721–727 (2008).

69. Sather, D. N. *et al.* Factors associated with the development of cross-reactive neutralizing antibodies during human immunodeficiency virus type 1 infection. *J. Virol.* **83**, 757–769 (2009).

70. Ye, J., Ma, N., Madden, T. L. & Ostell, J. M. IgBLAST: an immunoglobulin variable domain sequence analysis tool. *Nucleic Acids Res.* **41**, W34–40 (2013).

71. Abdiche, Y. N. *et al.* High-throughput epitope binning assays on label-free array-based biosensors can yield exquisite epitope discrimination that facilitates the selection of monoclonal antibodies with functional activity. *PLoS One* **9**, e92451 (2014).

72. Mast, F. D. *et al.* Highly synergistic combinations of nanobodies that target SARS-CoV-2 and are resistant to escape. *Elife* **10**, (2021).

73. Miller, J. L. *et al.* Quantitative bioluminescent imaging of pre-erythrocytic malaria parasite infection using luciferase-expressing Plasmodium yoelii. *PLoS One* **8**, e60820 (2013).

74. Vaughan, A. M. *et al.* A transgenic Plasmodium falciparum NF54 strain that expresses GFP-luciferase throughout the parasite life cycle. *Mol. Biochem. Parasitol.* **186**, 143–147 (2012).

75. Sack, B. K. *et al.* Mechanisms of stage-transcending protection following immunization of mice with late liver stage-arresting genetically attenuated malaria parasites. *PLoS Pathog.* **11**, e1004855 (2015).

76. Douglass, A. N., Metzger, P. G., Kappe, S. H. I. & Kaushansky, A. Flow Cytometry-Based Assessment of Antibody Function Against Malaria Pre-erythrocytic Infection. *Methods Mol. Biol.* **1325**, 49–58 (2015).

77. Murphy, S. C. *et al.* Real-time quantitative reverse transcription PCR for monitoring of blood-stage Plasmodium falciparum infections in malaria human challenge trials. *Am. J. Trop. Med. Hyg.* **86**, 383–394 (2012).

1 **Tables**

2 *Table 1. Combination of anti-PyCSP and anti-PyTRAP can improve sterile protection*
3 *from mosquito bite challenge.*
4

	Sterile Protection[a]				Comparison p-value[b]	
	Exp 1	Exp 2	Exp 3	Combined	vs. mIgG	vs. αCSP + mIgG
400 mIgG[c]	1/5 (20%)	0/5 (0%)	1/5 (20%)	2/15 (13.3%)	–	**0.033**
100 µg αCSP + 300 µg mIgG[c]	4/8 (50%)	3/5 (60%)	2/5 (40%)	9/18 (50%)	**0.033**	–
300 µg TY11 + 100 µg mIgG[c]	1/3 (33%)	0/5 (0%)	1/5 (20%)	2/13 (15.4%)	0.956	**0.049**
300 µg TY12 + 100 µg mIgG[c]	2/4 (50%)	0/5 (0%)	0/4 (0%)	2/13 (15.4%)	0.956	**0.049**
300 µg TY03 + 100 µg mIgG[c]	5/6 (83.3%)	0/5 (0%)	2/5 (40%)	7/16 (43.7%)	0.072	0.734
100 µg αCSP + 300 µg TY11	2/4 (50%)	1/5 (20%)	1/5 (20%)	4/14 (28.6%)	0.355	0.279
100 µg αCSP + 300 µg TY12	3/4 (75%)	0/5 (0%)	2/5 (40%)	5/14 (35.7%)	0.211	0.586
100 µg αCSP + 300 µg TY03	6/6 (100%)	4/5 (90%)	4/5 (80%)	14/16 (87.5%)	**0.000034**	**0.025**

5
6 [a] Sterile protection: mice that remain parasite-free (via microscopic blood-smear
7 monitoring) throughout the experimental time course.
8 [b] Barnard's exact test *p*-values shown were not adjusted for multiple comparisons due
9 to small group sizes and a small number of predefined comparisons being made.
10 [c] mIgG: normal mouse IgG control
11
Mice were injected with 100 µg/mouse of anti-CSP mAb (2F6), 300 µg/mouse of an
anti-PyTRAP mAb or a combination of both 24h prior to challenge by 5 *Py*-infected
mosquitos. Where only one mAb was injected, mice were also given non-specific mIgG
to total 400 µg/mouse. Mice were tracked for 14 days for parasitemia by thin blood
smear and those remaining parasite-free at day 14 were considered sterilely protected.
Number and percentages of mice protected across 3 independent experiments are
shown.
19

1 *Table 2. Combination of anti-PfCSP and anti-PfTRAP can improve sterile protection*
2 *from mosquito bite challenge.*
3

	Sterile Protection[a]				Comparison p-value[b]	
	Exp 1	**Exp 2**	**Exp 3**	**Combined**	**vs. mIgG**	**vs. 50 µg αCSP**
150 µg mIgG[c]	0/7 (0%) [#]	0/7 (0%) [#]	0/5 (0%)	0/19 (0%)	–	**<0.0001**
50 µg AKBR-4	0/6 (0%)	–	0/5 (0%)	0/11 (0%)	1	**0.0002**
50 µg αCSP	5/7 (71%) [#]	5/8 (63%) [#]	5/8 (63%)	15/23 (65%)	**<0.0001**	–
150 µg αCSP	–	3/5 (60%)	–	3/5 (60%)	**0.002**	0.88
50 µg AKBR-4 + 50 µg αCSP	–	6/7 (86%)	8/9 (89%)	14/16 (88%)	**<0.0001**	0.131

[a] Sterile protection: mice that remain parasite-free (via microscopic blood-smear
monitoring) throughout the experimental time course.
[b] Barnard's exact test *p*-values shown were not adjusted for multiple comparisons due
to small group sizes and a small number of predefined comparisons being made.
[c] mIgG: normal mouse IgG control
[#] indicates results previously reported in Kisalu *et al.* [55].
10
Mice were injected with indicated doses of either non-specific mIgG, anti-CSP mAb
CIS43, anti-PfTRAP mAb AKBR-4 or a combination of anti-CSP and AKBR-4 24h prior
to challenge with 5 *Pf*-infected mosquito bites. Mice were injected with human red blood
cells at day 5 and 6, and then blood sampled at days 7 and 9 to detect blood stage
parasitemia by qPCR. Number and percentages of mice protected across 3
independent experiments are shown.
17
18
19
20

1 **Figures**

3 *Figure 1. TRAP domain organization and constructs used.*

4 Ectodomain and deletion constructs for PyTRAP and PfTRAP generated using the
5 domain boundaries (**A**) were recombinantly expressed and purified alongside the
6 control CSP ectodomain proteins (**B**).
7

Figure 2. Polyclonal antibodies to PyTRAP inhibit parasite invasion, traversal and in vivo

infection.

Mice were immunized three times with PyTRAP or PyCSP ectodomains. (**A**) Immune
sera were used to verify binding to *Py* sporozoites via immunofluorescence. Shown is a
representative example of fixed, permeabilized sporozoites labeled with a 1:100 dilution
of polyclonal mouse serum from PyTRAP immunization. The anti-mouse IgG (anti-
TRAP serum) is in the red channel shown alone on the left and in combination with anti-
CSP mAb 2F6 (green channel) and a DAPI nuclear stain (blue channel); 10-μm scale
bars are shown. Immune sera were then assessed for function in vitro for inhibition of
invasion (**B**) and traversal (**C**). In **B** and **C**, pooled serum from cohorts of n=5 mice
(number of cohorts indicated above each bar) was tested in three independent assays.
Each data point represents the average "% of pre-immune" invasion or traversal of
these independent assays for each cohort pool. Each bar indicates the group mean,
with error bars representing standard error of the mean. Values representing percent
changes from 100% (indicated by dotted lines) are shown above. Asterisks indicate a
significant difference from 100% as determined by a two-tailed one-sample *t*-test. (**D**)
Immunized mice were challenged by the bite of 15 PyGFPluc-infected mosquitos and

assessed for parasite liver burden by bioluminescent imaging. Each data point represents an individual mouse with each color corresponding to an independent immunization-challenge experiment (total number of animals shown above each bar). Each data point was normalized to the mean flux from "naive" mice within each challenge experiment, while "control" mice were an additional group immunized with HIV Env gp120 protein. Each bar indicates the group mean, with error bars representing standard error of the mean. Values representing percent changes from 100% (indicated by a dotted line) are shown above. Asterisks indicate significance as determined by ANOVA with Kruskall-Wallis post-test. For **B-D**, * is $p \leq 0.05$; ** is $p \leq 0.01$; and **** is $p \leq 0.0001$.

Figure 3. Effects of PyTRAP monoclonal antibodies on parasite activity.

(**A**) Each mAb was assessed for in vitro function of inhibition of invasion and traversal.
In each case, mean values of % inhibition (i.e., 100% – invasion or traversal value) from
the 100-µg/mL mAb concentrations (bar plots with these and additional conditions
shown in **Suppl. Fig. 1**) are represented on a color axis. (**B**) Binding kinetics for each
mAb was measured by BLI and shown as kinetic maps with gray dashed diagonal
contour lines labeled with the corresponding K_d values and symbols representing the
characterized epitopes for invasion (*left*) and traversal (*right*) inhibition. Higher-affinity

(i.e., those possessing lower K_d values) mAbs are closer to the upper-right corner of this

plot. Symbol color coding represents "strong" inhibition for mean values ≤ 50%, "mild"

inhibition for values ≤ 70% and "weak" for mean values >70% observed at the 100-

µg/mL concentration. Red box highlights the region of the kinetic plots containing the

values for mAbs that showed strong inhibition in invasion and traversal assays. (**C**)

Summarized sterile protection ratios following passive-transfer-challenge experiments

(number of animals in each group is shown below the corresponding bar, individual

values shown in Table 1). For **C**, * is p≤0.05 and **** is p≤0.0001; values reported were

not adjusted for multiple comparisons due to small group size and limited comparisons.

Figure 4. Polyclonal antibodies to PfTRAP inhibit parasite invasion and traversal in vitro.

Mice were immunized three times with PfTRAP or PfCSP ectodomains. (**A**) Immune sera were used to verify binding to *Pf* sporozoites via immunofluorescence. Shown are fixed, permeabilized sporozoites labeled with a 1:800 dilution of polyclonal anti-PfTRAP mouse serum (followed by anti-mouse IgG secondary; green channel), fluorescently labeled anti-PfCSP monoclonal antibody 2A10 (red channel, right image) and DAPI nuclear stain (blue channel, right image); 10-µm scale bars are shown. Immune serum was then assessed for function in vitro for inhibition of invasion (**B**) and traversal (**C**). In **B** and **C**, each data point is the average "% of pre-immune" invasion or traversal from technical triplicates in independent experiments; two separate immunization experiment sets are represented as "PfTRAP cohort 1" and "PfTRAP cohort 2". Each bar indicates the group mean, with error bars representing standard error of the mean and percent change from 100% (shown as dashed line) shown above. Asterisks indicate a

significant difference from 100% as determined by two-tailed one-sample *t*-test where * is p≤0.05; ** is p≤0.01; and **** is p≤0.0001.

Figure 5. Monoclonal antibodies to PfTRAP inhibit parasite invasion, traversal in vitro.

(**A**) Each mAb was assessed for in vitro function of inhibition of invasion and traversal.
In each case, mean values of % inhibition (i.e., 100% – invasion or traversal value) from
the 100-µg/mL mAb concentrations (bar plots with these and additional conditions
shown in Suppl. Fig. 5) are represented on a color axis. (**B**) Binding kinetics for each
mAb was measured by BLI and shown as kinetic maps with gray dashed diagonal
contour lines labeled with the corresponding K_d values and symbols representing the
characterized epitopes for invasion (*left*) and traversal (*right*) inhibition. Higher-affinity
(i.e., those possessing lower K_d values) mAbs are closer to the upper-right corner of this
plot. Symbol color coding represents "mild" inhibition for values ≤ 70% and "weak" for
mean values >70% observed at the 100-µg/mL concentration. (**C**) Summarized sterile
protection breakdowns following passive-transfer-challenge experiments (number of
animals in each group is shown below the corresponding bar, individual values shown in
Table 2). For **C**, ** is p≤0.01, *** is p≤0.001, and **** is p≤0.0001; values reported were
not adjusted for multiple comparisons due to small group size and limited comparisons.
17