Hair phenotype diversity across Indriidae lemurs

Running Title: Hair evolution in Indriids

*Elizabeth Tapanes[1,I], Rachel L. Jacobs[1], Ian Harryman[1,II], Edward E. Louis Jr.[2], Mitchell T.

Irwin[3], Jason M. Kamilar[4,5], Brenda J. Bradley[1]

[1] Center for the Advanced Study of Human Paleobiology, Department of Anthropology, The

George Washington University

[2] Conservation Genetics Department, Omaha's Henry Doorly Zoo and Aquarium

[3] Department of Anthropology, Northern Illinois University

[4] Department of Anthropology, University of Massachusetts Amherst

[5] Graduate Program in Organismic and Evolutionary Biology, University of Massachusetts

Amherst

*= Corresponding author:

Elizabeth.tapanes@gmail.com

[I] Present address of Elizabeth Tapanes: University of California San Diego, Department of Biological Sciences
[II] Present address of Ian Harryman: Stanford University, Department of Earth System Science

Abstract

Objectives

Hair (i.e., pelage/fur) is a salient feature of primate (including human) diversity and evolution—serving functions tied to thermoregulation, protection, camouflage, and signaling—but wild primate pelage evolution remains relatively understudied. Specifically, assessing multiple hypotheses across distinct phylogenetic scales is essential but is rarely conducted. We examine whole body hair color and density variation across Indriidae (*Avahi*, *Indri*, *Propithecus*)—a lineage that, like humans, exhibits vertical posture (i.e., their whole bodies are vertical to the sun).

Materials and Methods

Our analyses consider multiple phylogenetic scales (family-level, genus-level) and hypotheses (e.g., Gloger's rule, the body cooling hypotheses). We obtain hair color and density from museum and/or wild animals, opsin genotypes from wild animals, and climate data from WorldClim. To analyze our data, we use phylogenetic generalized linear mixed models (PGLMM) using Markov chain Monte Carlo algorithms.

Results

Our results show that across the Indriidae family, darker hair is typical in wetter regions. However, within *Propithecus*, dark black hair is common in colder forest regions. Results also show pelage redness increases in populations exhibiting enhanced color vision. Lastly, we find follicle density on the crown and limbs increases in dry and open environments.

Discussion

This study highlights how different selective pressures across distinct phylogenetic scales

have likely acted on primate hair evolution. Specifically, our data across *Propithecus* may

implicate thermoregulation and is the first empirical evidence of Bogert's rule in mammals. Our

study also provides rare empirical evidence supporting an early hypothesis on hominin hair

evolution.

Keywords: lemurs, coloration, opsins, integument, adaptation

Introduction

Hair (i.e., pelage, fur) is a salient feature of primate diversity and evolution and is the product of natural and sexual selection. Hair has been found to play a role in conspecific signaling (Allen & Higham, 2013), thermoregulation (Fratto & Davis, 2011), camouflage (Barrett et al., 2019; Hoekstra et al., 2005) and to possibly provide protection from pathogens/parasites (Paus & Foitzik, 2004). Across all mammals, primates—including humans—arguably exhibit the most diversity in hair color (e.g., bright reds) and growth patterns (e.g., mustaches) between and within species (Caro, 2005; Rowe, 1996). For example, within Colobinae, some *Colobus* spp. sport a long black and white dorsal cape, and *Pygathrix* spp. are often tri-colored and short-haired (Rowe, 1996). Similarly, human hair color and texture vary throughout the human body, between individuals, between populations, and can be correlated with ancestry and ethnicity (Lasisi et al., 2016; Seibert & Steggerda, 1999; Steggerda & Seibert, 1941). Nevertheless, our knowledge about hair evolution is based mainly on non-primate mammals. Primate hair remains relatively understudied (with some notable exceptions: (e.g., (Allen & Higham, 2013; Bell et al., 2021; Kamilar et al., 2013; Rakotonirina et al., 2017; Santana et al., 2012)), with most research taking a macro-scale approach and sampling broadly across the primate order (e.g., all catarrhines).

These previous studies indicate that variation in primate hair color likely evolved in response to multiple distinct but not mutually exclusive pressures associated with ecology, visual systems, activity cycles, and sociality (Allen & Higham, 2013; Bell et al., 2021; Kamilar et al., 2013; Rakotonirina et al., 2017; Santana et al., 2012). For example, across primates, melanic (i.e., black) colors often co-occur with humid regions—which may be due to camouflage, pleiotropy, pathogen resistance, or selection on a related trait (i.e., Gloger's rule) (Bell et al.,

2021; Kamilar & Bradley, 2011; Santana et al., 2012; 2013). Gloger's rule generally predicts

birds and mammals will be more pigmented in tropical regions where it is warm and humid, such

as tropical rainforests (Gloger, 1833). However, without accounting for variation within families,

genera, and species, other subtle trends or relationships may not be detected. For example, recent

work in birds suggests a distinct rule (Bogert's rule) may explain some melanic variation in

endotherms depending on the phylogenetic and geographic scale sampled (Bogert, 1949; Delhey

et al., 2019). According to Bogert's rule, melanic colors should co-occur with cold regions as

dark colors may absorb heat better and aid in thermoregulation (Bogert, 1949; Delhey et al.,

2019). However, there is no current empirical evidence for this rule in primates (or any

mammals). Yet across human populations, variation in eumelanin exceeds that seen in red,

blonde, or lighter hair (Lasisi et al., 2016), which suggests that important biological variation

may be identified if the scale of analyses was smaller than previously considered (i.e., within a

few populations vs. across all populations) (Norton et al., 2016).

Across non-human primates, red hues (and facial patterning) may act as a means to aid

conspecific communication (Cuthill et al., 2017; Santana et al., 2012; 2013; Winters et al., 2020).

For example, diurnal activity and the ability to perceive a wider range of colors compared to

other mammals may have contributed to the enhanced color diversity in specific primate clades

(Bell et al., 2021). However, the evidence for coevolution of hair coloration, specifically red hair

and color vision in primates, remains mixed (Fernandez & Morris, 2007; Kamilar et al., 2013).

Our understanding of primate hair color may benefit from broader sampling across wild

populations and/or analyses conducted on smaller phylogenetic scales.

Similarly, our understanding of pelage density evolution remains limited to extensive

macro-scale studies. Across primates, the primary trend is that hair density decreases as body

size increases (Sandel, 2013; Schwartz & Rosenblum, 1981). One early hypothesis argued that a

reduction in body hair density across hominins coupled to increases in activity level, sweat

glands, sweat production, and a change to a dry savannah environment may have aided

thermoregulation (i.e., "body-cooling hypothesis") (Jablonski & Chaplin, 2000; Post et al., 1975;

Wheeler, 1992). Despite various physiological models testing evidence of the body-cooling

hypothesis (Chaplin et al., 1994; Schwartz & Rosenblum, 1981; Wheeler, 1992), there is only

rare empirical evidence for this phenomenon within primates (Best et al., 2019). There also

remains a poor understanding of how hair density in primates may be impacted by other factors,

such as climate. Because hair does not preserve in the fossil record, studies on wild primates can

provide critical insights into the evolutionary mechanisms underlying one of the most interesting

traits among humans.

This study examined whole-body pelage color and hair/follicle density across Indriidae—

a diverse clade of lemurs. Since the Indriidae clade varies in critical aspects known to drive

primate (including human) hair variation, they make a good proxy for testing multiple hair

hypotheses that have not yet been jointly assessed in one study. Our overarching goals were to 1)

examine if Indriidae pelage variation varies across distinct phylogenetic scales and 2) determine

if multiple, non-exclusive selective pressures can impact the same trait. Specifically, we aimed to

jointly assess the impacts of climate, body size, and color vision on hair evolution at distinct

phylogenetic scales. Collectively, this family (comprising three genera: nocturnal *Avahi* and

diurnal *Indri* and *Propithecus*) inhabits almost every bio-climate of Madagascar (e.g., humid

rainforest, temperate forests, open spiny forest, mangroves) (Mittermeier et al., 2010). The hair

of the 13 species in our study varies intra- and interspecifically in both color (e.g., melanic,

tricolored orange) and growth morphologies (e.g., long silky hairs, short hairs) (King et al., 2014;

Mittermeier et al., 2010; Petter & Peyrieras, 1972; Rakotonirina et al., 2014; Tattersall, 1986).

Indriidae also vary in body size (≈1kg (*Avahi*), ≈3-8kg (*Propithecus*), ≈10kg (*Indri*))—and

phylogenetic relatedness likely impacts body size evolution (Lehman, 2007). Thus, we

specifically (1a\2a) aimed to uncover how climate and body size influences: pelage brightness

(i.e., black color), hue (i.e., red color), and hair follicle density across Indriidae and within

Propithecus. We expected that climate impacts pelage color and density evolution distinctly

across different phylogenetic scales. We predicted that body size only impacts hair/follicle

density, based on evidence from other primates (Sandel, 2013). Diurnal Indriidae (*Indri* and

Propithecus) also vary in their color vision. In many species, some females have trichromatic

color vision (ability to distinguish reddish and greenish hues), while males and other females are

red-green color blind (polymorphic color vision; Jacobs et al., 2017). Although this feature

occurs in some other day-active lemurs, Indriidae has been found to exhibit greater genetic

variation in color vision across species (Jacobs et al., 2017). Accordingly, we (2b) aimed to

understand how this variation in color vision impacts pelage hue in diurnal genera. We predicted

that populations with enhanced capacity for trichromatic color vision would have individuals

with redder pelage (Sumner & Mollon, 2003). Our results provide critical insight into how

multiple variables impact primate hair evolution across scales. Importantly, this family consists

solely of vertical clingers and leapers, which means they exhibit upright locomotory posture

similar to humans (via bipedal hopping (Schmidt, 2011)). Thus, these results (specifically those

on hair density) may provide insight into human hair evolution.

Materials and Methods

Hair morphology and body size data collection

We used hair color and density measurements from museum research skins ($N_{Total} = 63$;

$N_{Museum} = 59$) and living animals in one wild population ($N_{Wild} = 4$, *P. diadema* from

Tsinjoarivo) (SI 1). Museum samples spanned three museum collections (SI 2). In total, we

sampled six *Avahi* populations (Figure 1A), 23 *Propithecus* populations (Figure 1B), and four

Indri populations (Figure 1C). Our sampling included nocturnal and diurnal genera (Figure 1D)

across all significant bioclimatic regions in Madagascar (Figure 1E). Many individuals did not

have available age or sex-class data. Hence, we did not test for differences between age classes

or sex. Although some preliminary evidence suggests there may be subtle pelage dichromatism,

sifakas are unlikely highly sexually dichromatic in terms of coloration. Instead, sifakas exhibit

more significant visible variation within sexes (i.e., bimorphism in males) (Lewis & Van Schaik,

2007; Mittermeier et al., 2010; Spriggs, 2017; Tapanes et al., 2018). Measurements of hair from

wild sifakas consisted of following the same protocol as museum individuals. We also extracted

body length (mm), measured from head to tip of tail, from all individuals as a proxy for body

size using Photoshop CC 6. We standardized all measurements using the known length of a ruler

in the photos. Wild sifakas were captured for ongoing parallel research using a standard capture

protocol established by the Prosimian Biomedical Survey Project (PBSP) Team (used on 750+

lemurs, 16+ sites, and 35+ species since 2000) (Dutton et al., 2003; Irwin et al., 2010). Groups

were habituated before capture. The capture team was led by a trained professional, and captures

were performed at close range using Pneu-dart 9mm disposable non-barbed darts containing

tiletamine/zolazepam (Telazol) for immobilization.

<u>Hair coloration collection</u>

We used color-calibrated photographs of all animals (N = 63 individuals; SI 1) and a

Canon EOS Rebel SL1/EOS 100D mounted with a Canon EF-S18-55mm f/3.5-5.6 IS STM lens.

We calibrated each image against a color standard with 24 color squares of known reflectance

values. The camera color mode was set to 'faithful.' The camera was calibrated with a grey

standard to set a manual white balance. All individuals were photographed against dull

backgrounds using the adjacent method (Bergman & Beehner, 2008; Kendal et al., 2013). We

followed established protocols for calibrating photos in previous studies for both museum and

wild specimens (Stevens et al., 2007; 2009).

We used the Red-Green-Blue (RGB) color model and extracted all RGB values using

Photoshop CC 6 across 14 body regions. Due to the high degree of variation across the pelage of

the *Propithecus* tail, ventral surface, and dorsal surface, we divided measurements of these

regions into three quadrants (lower, mid, and upper) (SI 3). If one of these regions was damaged,

we took measurements from the center as the "average." Primate hair occupies a relatively small

color space, primarily varying along two axes. The first relates to variation in luminance and

exhibits little chromatic variation and is captured by similar red, green, and blue values. Thus,

we used RGB averages here as a proxy for 'brightness' in a hair patch to indicate if the hair is

white, a shade of grey, or black. The second axis of variation is hue and indicates variation along

a red/brown axis and can be captured by a ratio of red to green. The latter is because the green

and blue channels are most similar, while the red channel exhibits higher values on average. As

the Red-Green ratio increases, hair appears increasingly red (Kamilar et al., 2013). We ran a

correlation matrix across color values and removed any that were highly correlated with each

other (r > .90). This reduced our hair color variables to RGB averages and RG ratios for the (1)

crown, (2) cheek, (3) dorsal torso, (4) ventral torso, (5) forelimb, (6) hindlimb, and (7) tail. In

total, we collected 882 brightness and 882 hue measurements (63 individuals x 14 sampling

regions). Previous work on museum skins indicated no effect of storage time on pelage

173 coloration from 142 museum skins collected over 100 years (1898 and 1998) (Kamilar et al.,

174 2013).

176 <u>Hair and follicle density collection</u>

177  We used a protocol that borrows from dermatological methods and instrumentation, which

178 automates hair density measurements (Avram & Rogers, 2009). We used CompareViewHair®

179 software (STR Technologies) tools combined with a specialized hand-held digital light microscope

180 (ProScope HR2©; Bodelin) with a 100x lens. The software included a built-in procedure for

181 calibrating the lens and software before use. The hair of an animal was parted, and the lens was

182 placed on the skin. A light within the microscope illuminated the skin and provided needed light

183 for subsequently photographing the area. We collected microscopy images (detailing transects of

184 hair follicles and shafts) and compiled semi-automated measurements in the laboratory using this

185 system. We examined two hair morphology properties: hair density (shafts/cm^2) and follicle

186 density (infundibula/cm^2).

187  We collected three microscopy photos (replicates) from five body regions (SI 4) of each

188 animal (n = 47 individuals) (SI 1) using the hand-held microscope, adhering to a previously

189 established protocol (Bradley et al., 2014). The method provided a visual transect/plot of follicles

190 (i.e., infundibula) and the number of hair shafts emerging from each. We used the 'density tool'

191 in CompareViewHair® software to estimate the number of infundibula per cm^2 (i.e., follicle

192 density). We marked each follicle strand manually and used that number to estimate the final

193 value in any given region (given the specific magnification). However, this protocol assumed

194 that the entire photograph was visible, which was not always the case. Therefore, in our final

195 measurements for 'follicle density,' we used the value the software provided of density per cm^2

196 and divided it by the visible photo's proportion. Similarly, for our final measurement for 'hair

density,' we used the value the software calculated for density per cm^2 and multiplied it by the

average number of hairs emerging from each follicle. In total, we collected 235 hair and 235

follicle density measurements (47 individuals x 5 body regions). Previous work on museum skins

shows no effect of storage time on pelage density for skins collected over 100 years (Bradley et

al., 2014).

Climatic and phylogenetic data collection

We collected climate and environmental data from 33 sampling locations across

Madagascar (Figure 1). Specifically, we extracted data from WorldClim (version 1) that

consisted of averages from 1960 to 1990. This likely created some noise in the data given

museum data spanned the early 1900s to late 1990s. However, Indriidae lifespans (≈20+ years)

and generation times are long (≈3-10 years) (Lawler et al., 2009; Godfrey et al., 2001; Guevara

et al. 2018). Thus, the data are likely representative of accurate averages across bioclimates and

phenotypes. Localities were obtained from museum records, and we used Google Earth to obtain

geographic coordinates of the nearest forested area in the species range. Localities were listed as

forests or towns with directional coordinates (e.g., 40 km W of Tamatave). We used R packages

'raster' and 'sp' to extract all localities' climate data in the WorldClim database (Hijmans &

Etten, 2012; Pebesma & Bavand, n.d.). We ran a correlation matrix for all 19 bioclimatic

variables and removed any highly correlated variables (r > .80). In the end, we used five

bioclimatic variables in our comparative analysis as predictor variables (SI 5).

DNA extractions for opsin genotypes

We categorized diurnal Indriidae color vision status (e.g., polymorphic color vision,

dichromatic – red-green colorblind) based on X-linked opsin genotypes (Jacobs et al., 2017,

2019). Our analysis included diurnal Indriidae represented by a subset of species/populations for

which pelage data were also available ($N_{species}$ = 7; $N_{populations}$ = 10; SI 6). Opsin genotypes for

populations of diademed sifaka (*Propithecus diadema*), Milne-Edwards's sifaka (*Propithecus*

edwardsi), golden crowned sifaka (*Propithecus tattersalli*), Verreaux's sifaka (*Propithecus*

verreauxi), and *Indri indri* were previously published (Jacobs et al., 2017). Data for the silky

sifaka (*Propithecus candidus*) and Perrier's sifaka (*Propithecus perrieri*) were based on opsin

genotypes obtained from five individuals from Anjanaharibe-Sud ($N_{X\ chromosomes}$ = 8) and

Analamera ($N_{X\ chromosomes}$ = 8) populations, respectively, following (Jacobs et al., 2019). Briefly,

we extracted genomic DNA from blood/tissue samples using phenol-chloroform-isoamyl

extraction protocols (Sambrook et al., 1989). We used quantitative PCR (Mic qPCR Cycler, Bio

Molecular Systems; Rotor-Gene Q, Qiagen) to amplify exons 3 and 5 of the X-linked opsin gene.

Amplification was followed by high-resolution melt analysis (HRMA) (*following* (Jacobs et al.,

2016)), and we determined genotypes for each exon based on the observed melt curve shape and

temperature compared to positive controls ($N_{Replicates}$ = 2 per exon). To further confirm genotype

calls, we Sanger sequenced qPCR amplicons from at least two individuals representing each

observed melt curve. All sequences matched scored genotypes based on melt curves.

We estimated the peak spectral sensitivity of observed opsin alleles in each population

(*following* (Jacobs et al., 2017)) and determined the difference between the two alleles with

maximum and minimum peak spectral sensitivities. These values, along with a total number of

observed alleles, were used in downstream analyses as proxies for enhanced trichromatic color

vision. Trichromatic color vision is most optimal with more widely separated photopigments

(e.g., (Osorio et al., 2004)). All else being equal, a greater number of opsin alleles results in an

increased proportion of heterozygous (trichromatic) females in a population.

Analyses

<u>Climate and body size effects on pelage brightness and hue</u>

We conducted four principal component analyses (PCAs): two across Indriidae and two

within *Propithecus*. One was for all the brightness variables across body regions and a second for

all the hue variables across body regions for Indriidae, to test specific hypotheses related to each

and to reduce the dimensionality of the data. We then ran two additional albeit similar PCAs

within only the *Propithecus* genus—given that the *Propithecus* genus is more widely

geographically distributed than either *Avahi* or *Indri*. We removed any variables with strongly

correlated vectors in the PCAs and then re-ran them to obtain the final PC values. We only used

the resulting PC values with eigenvalues over one as our dependent variables.

To test the effects of climate on pelage color while accounting for the evolutionary

relatedness of taxa, we used phylogenetic generalized linear mixed models (PGLMM) using

Markov chain Monte Carlo algorithms (MCMC) via the 'MCMCglmm' package in R (Hadfield,

2010). Our Indriidae phylogeny was based on a previously published lemur phylogeny that

contains a complete picture of Indriidae evolution (Herrera & Dávalos, 2016). We assigned the

phylogeny to the 'pedigree' argument in MCMCglmm, and the resulting matrix was used to

weigh the error structure of the model. Similar methodologies have been used to test the effects

of climate variables on bumblebee body size (Ramírez-Delgado et al., 2016) and frog vocal

signals (McLean et al., 2013). Each model included one color PC value as the dependent variable

and the bioclimatic variables (SI 5) and body size as independent variables, with species and

phylogeny considered random factors. We log-transformed PC values, bioclimatic variables, and

body size to achieve normality. All models ran for 5,000,000 iterations after a burn-in of 500,000

iterations and a thinning interval of 500 using flat priors. We assured effective sampling sizes

were adequate for all our models (i.e., ESS virtually equal to MCMC sample size) (Villemereuil

& Nakagawa, 2014).

We assigned statistical significance when pMCMC ≤ 0.05 (which is equivalent to a p-

value (Villemereuil & Nakagawa, 2014)) and when the confidence interval for a given variable

did not include zero (Ramírez-Delgado et al., 2016). To estimate what fraction of the pelage

variation was attributable to phylogenetic relatedness, we calculated H^2 from Lynch (Lynch,

1991) (equivalent to Pagels's λ) which corresponds to the degree of variation in a given trait

attributable to phylogenetic relatedness (Hansen & Orzack, 2005; Kamilar & Cooper, 2013). We

ran the models across the whole Indriidae family and within the *Propithecus* genus. Using

G*Power analysis (Faul et al., 2009), we estimated the PGLMMs would detect a small effect

((Indriidae, 0.25)-(*Propithecus*, 0.35)) with ≈80% power.

<u>Climate and body size effects on pelage density</u>

We conducted two principal component analyses (PCA) of hair and follicle density

values: one across all Indriidae taxa and a second within the *Propithecus* genus. We removed any

variables whose vectors strongly correlated in each of the PCAs. We used the resulting PC

values with eigenvalues over one as dependent variables in our PGLMM models, following the

methodology outlined above (see: climate and body size effects on pelage brightness and hue).

Using G*Power analysis (Faul et al., 2009), we estimated the Indriidae PGLMM would detect a

small effect (0.35), and the *Propithecus* PGLMM would detect a moderate effect (0.50) with

≈80% power. We also conducted an Analysis of Variance (ANOVA) to test for differences

between hair or follicle density across distinct body parts.

<u>Polymorphic trichromacy and pelage hue variation</u>

We ran one PCA on hue variables among a subset of individuals from which we had

paired opsin genotypes and pelage measurements from the same population. We used any PCs

with eigenvalues over one as the dependent variable in our linear models. To estimate the

relationship between pelage hue and color vision, we conducted both (1) a simple linear

regression and (2) PGLMM. Due to the small sample size (n = 10 localities/populations) (SI 6),

we assumed that the "true" result likely fell between a model where lambda = 0 (simple linear

model) and one where lambda = 1 (PGLMM). We followed the methods described previously,

except we used 'spectral sensitivity range' (i.e., difference in opsin spectral sensitivity) and the

'total number of opsin alleles' as our independent variables. Despite the low sample size, using

G*Power analysis (Faul et al., 2009), we estimated the simple linear model would detect a

moderate effect (0.65) with ≈80% power.

Results

Climate and body size effects on pelage brightness and hue

<u>Across Indriidae</u>

Across Indriidae, the first PC for the brightness PCA explained ≈61% of the variation (SI

7). The highest loadings on PC1 were hindlimb, tail, and cheek brightness (SI 8). We found a

statistically significant and positive association between hair brightness and precipitation

seasonality (Table 1). Thus, darker pelage across Indriidae is associated with forests of lower

precipitation seasonality (i.e., higher rainfall, more humid) (Figure 2A). Brightness across

Indriidae exhibited a moderate phylogenetic signal (0.68) (Table 1). On the hue PCA, the first

three PCs explained ≈68% of the variation (SI 9, SI 10). The highest loadings were forelimb,

hindlimb and ventral torso hue (PC1), cheek and ventral torso (PC2), as well as crown hue (PC3)

(SI 10). We found a statistically significant relationship between PC1 and PC2 with precipitation

seasonality (Table 2). Indriidae with redder pelage hues live in regions that experience less

seasonality (i.e., higher rainfall) (Figure 3, SI 24). Hue values across Indriidae exhibited low to

moderate phylogenetic signal (0.25-0.48) (Table 2).

<u>Within *Propithecus*</u>

For the brightness PCA, the first PC explained ≈70% of the variation within *Propithecus*

(SI 11). The highest loadings on PC1 were hindlimb, dorsal torso, and forelimb brightness (SI

12). We found a statistically significant and positive association between hair brightness and

mean annual temperature (Table 1). Thus, darker pelage within *Propithecus* is associated with

colder forests (Figure 2B). Brightness across *Propithecus* exhibited a high phylogenetic signal

(0.90) (Table 1). For the hue PCA, the first three PCs explained ≈80% of the variation within

Propithecus (SI 13, SI 14). The highest loadings were forelimb, dorsal torso, and hindlimb hue

(PC1), cheek, ventral torso, and hindlimb hue (PC2), and crown hue (PC3) (SI 14). We found a

statistically significant relationship between PC2 and mean annual temperature, precipitation

seasonality, precipitation of the wettest month, and body size (Table 3). Thus, redder hues within

Propithecus are associated with lower precipitation seasonality (and higher rainfall). We found

no statistically significant relationships between either PC1 or PC3 with the tested variables.

There was low to moderate phylogenetic signal for hair hue within *Propithecus* (0.22-0.63)

(Table 3).

Climate and body size effects on pelage density

<u>Across Indriidae</u>

Across Indriidae, the first three PCs explained ≈70% of the variation (SI 15, SI 16). The

highest loadings were forelimb, hindlimb, and crown follicle density (PC1), ventral torso hair

density and crown and hindlimb follicle density (PC2), and dorsal torso hair density (PC3) (SI

16). We found a statistically significant relationship between PC1 and precipitation seasonality

as well as between PC3 and body size (Table 4). Specifically, the results suggest that in forests

with high precipitation seasonality (dryer, open forests exposed to the sun) individuals exhibit an

increase of follicle density on the limbs and crown (Figure 4). Also, smaller taxa have the highest

hair density; and, larger individuals exhibit lower hair density on average (Figure 5). Pelage

density across Indriidae exhibited low phylogenetic signal (0.21-0.33) (Table 4). We also found a

statistically significant difference between hair density across body regions sampled (Table 5).

Specifically, we found the crown and forelimbs exhibit the greatest hair density, and the ventral

torso exhibits the least hair density (Figure 6A). However, we found no significant differences in

follicle density of regions sampled (Table 5, Figure 6B).

352
<u>Within *Propithecus*</u>

Within *Propithecus*, the first three PCs explained ≈70% of the variation (SI 17, SI 18).

The highest loadings were forelimb follicle density, crown hair density, and hindlimb follicle

density (PC1), crown and hindlimb follicle density as well as hindlimb hair density (PC2), and

ventral torso hair density (PC3) (SI 18). We found a statistically significant relationship between

PC1 and precipitation seasonality, similar to Indriidae (Table 6). However, unlike the Indriidae

results, none of the PCs were significant with body size. Though, we detected significant

differences between hair density and distinct body regions (SI 19), and no differences were

detected among follicle density within *Propithecus* (SI 19, SI 20).

Polymorphic trichromacy and pelage hue variation

For the hue PCA using only a subset of the data from *Indri* and *Propithecus*, the first two

PCs explained ≈66% of the variation (SI 21). The highest loadings to PC1 were forelimb, dorsal

torso, and hindlimb hue, and the highest loadings to PC2 were cheek, ventral torso, and hindlimb

hue (SI 22). From the simple linear model (lambda = 0), we found a statistically significant

relationship between PC1 with the difference in opsin spectral sensitivity (p = 0.03; R^2 = .51),

but not with the overall number of alleles in the population (Table 7). Specifically, we found a

significant negative linear relationship between spectral sensitivity and hue (Figure 7). Despite

the smaller sample size of the PGLMM (lambda = 1), we detected a similar association that

approaches statistical significance (pMCMC = 0.09) and only moderate phylogenetic signal

(0.41) (SI 23).

Discussion

Our overarching goals were to examine 1) if Indriidae pelage traits varied across distinct

phylogenetic scales and 2) if multiple non-exclusive selective pressures would impact the same

trait. We found support for both; distinct aspects of pelage color and density in Indriidae and

sifakas (genus *Propithecus*) vary with climate, body size, and/or color vision.

Climate and body size effects on pelage brightness and hue

Our first sub-aim sought to uncover how climate and body size influenced pelage

brightness and hue across two distinct scales (across a family [Indriidae] and within a genus

[*Propithecus*]). Our results show distinct trends across scales regarding dark pelage (i.e.,

brightness). Specifically, across Indriidae, greater brightness is associated with greater

precipitation seasonality (Table 1). Greater hue, similarly, is associated with less precipitation

seasonality across Indriidae and *Propithecus* (SI 24)) (Table 2). Thus, more pigmented coats

(i.e., individuals with blacker and redder hair) occur in regions of lower precipitation seasonality

(Figure 2A, Figure 3). Generally, in Madagascar, low precipitation seasonality is strongly and

negatively associated with high precipitation forests of the East (SI 24). The result for brightness

follows Gloger's rule, predicting an increase in eumelanin (i.e., black pigments) (Delhey, 2017).

Yet, within *Propithecus*, results indicate a relationship between brightness and mean annual

temperature (Table 1). Thus, dark-haired sifakas are generally associated with colder regions,

which serves as evidence of Bogert's rule (or the thermal melanism hypothesis) (Figure 2B)

(Bogert, 1949).

Recent work highlights that discovering evidence for Bogert's or Gloger's rule may

merely reflect differences in geographic space, scale, and climate gradients (Delhey et al., 2019).

We show in this analysis that different effects can be detected across the primate coat, depending

on the sampling scale. Additionally, the strength of selective pressures acting on pelage may also

play a role in detecting either rule. For example, if the need to camouflage or avoid parasites is

high, rainfall (which predicts Gloger's rule) may strongly affect pelage color (da Silva et al.,

2016). This is because higher rainfall may also be associated with higher loads of parasites

(Shearer & Ezenwa, 2020) or may indicate low light forested environments that enhance crypsis

for dark animals (Caro, 2005; Delhey, 2018). In contrast, if the need for thermoregulation is

high, temperature (which predicts Bogert's rule) may have a more substantial effect on pelage

color than rainfall (Galván et al., 2018). It is hypothesized that black colors aid with

thermoregulation in colder climates since they likely increase absorption of ultraviolet rays.

While this has been documented widely in ectoderms and recently in birds (Bishop et al., 2016;

Bogert, 1949; Clusella Trullas et al., 2007; Delhey et al., 2019; Xing et al., 2016), to date, our

results serve as the first empirical evidence for Bogert's rule in mammals. Notably, the range of

temperatures occupied by *Propithecus* species (≈10-27ºC) corresponds exceptionally well to the

range at which Bogert's rule is expected to influence plumage color (≈10-25ºC) in passerines

(Delhey et al., 2019). At this temperature range there is much nuance in the climatic space

between temperature and precipitation, which is in turn is hypothesized to influence the rate of

diversification and maintenance of species barriers (Delhey et al., 2019; Martin et al., 2010).

Insights from other primate species may further elucidate the importance of Bogert's and

Gloger's rules to broader primate evolution.

Both body size and phylogeny seem to have minimal impact on pelage color. First, body

size only influenced hair hue across the *Propithecus* clade and otherwise did not affect our

coloration models across either scale (Table 2, Table 3). At first glance, smaller individuals seem

to exhibit redder hues; however, this trend is driven mostly by *Propithecus* species on the

western side of the island. This may indicate subtler variation between species that our analysis

does not capture. However, these results may also reflect broader climatic trends, variations in

color vision (addressed below), or is perhaps the product of our body size sampling

methodology. Second, most of the coloration traits we sampled exhibited low to moderate

phylogenetic signals. Our brightness models, on the other hand, had a high phylogenetic signal

(≈0.80). Thus, brightness may implicate genetic drift (Losos, 2008; Revell et al., 2008).

However, it is well established that primate coat color exhibits little to no phylogenetic signal

and is a poor marker of phylogeny (Bell et al., 2021). An alternative explanation for such a high

signal is that it is an "artificial" result from taxonomic inflation that merely reflects patterns of

geographic variation (Kamilar & Muldoon, 2010).

Climate and body size effects on pelage density

Our second sub-aim sought to uncover how climate and body size influenced pelage hair

and follicle density across two distinct scales (across a family [Indriidae] and within a genus

[*Propithecus*]). We predicted that body size would affect density because these variables are

associated across primates. Body size may also have been one of the factors influencing the

reduction of the hairy coat in hominins (Sandel, 2013; Schwartz & Rosenblum, 1981). We found

support for this prediction across Indriidae (Table 4). Specifically, higher hair density is

associated with smaller body sizes (*Avahi*), and lower hair density is found in larger animals

across Indriidae (Figure 4). One explanation for this result may be that interfollicular distance

increases as surface area increases, thus, decreasing hair density in larger animals (Sandel, 2013).

In addition, this relationship may provide thermoregulatory benefits since larger animals may

struggle more to dissipate heat (Schwartz & Rosenblum, 1981). Our models show that climate

also influences pelage variation across Indriidae. Limb and crown follicle density increases as

precipitation seasonality increases (i.e., shifting to dry open habitats) (Table 4; Figure 4).

Interestingly, the crown and forelimbs are also the regions that exhibit the highest degree of hair

density (Figure 6A)—but not follicle density (Figure 6B). Thus, there are likely two mechanisms

to increase density: (1) increase hair follicle density or (2) increase the number of hairs emerging

from each follicle. Since follicle density is determined in early embryogenesis, follicle density

may be more strongly genetically determined than hair density (which may be more plastic to

rapid environmental changes, such as seasonality) (Tapanes et al., 2021). Alternatively, follicle

density may be easier to measure without error and could be a more reliable estimate for

'hairiness.'

Nevertheless, the shift towards drier habitats exposed to more sun (such as Western spiny

thickets) leads to an increase in follicle density in the limbs and crown (Figure 4), where hair

density is already the highest of all regions (Figure 6A). Since all Indriidae are vertical clingers

and leapers, the crown and upper limbs are the most directly impacted by the sun's rays—as they

are known to hold up their arms while jumping bipedally. Thus, increased hair density in these

regions may aid individuals in this clade with protection from ultraviolet rays (and/or protect the

underlying skin from abrasion in spiny thickets). We consider this density pattern may help

prevent photolysis of folate, an essential metabolite in the formation of the embryonic neural

tube. Protection against folate leaching was likely a selection target in the evolution of high

melanization within the skin of humans living in high UV areas (Jablonski & Chaplin, 2000). In

humans, lower levels of folate (or folate deficiency) are also often significantly associated with

hair loss (Almohanna et al., 2019; Ertug, 2018), and folate may play a significant role within the

hair follicle (Durusoy et al., 2009). Since the skin of most Indriidae is already highly melanized,

it is not surprising that populations exposed to hotter environments and more direct sun often

have lighter coats (Figure 2B, Figure 3)—potentially to prevent overheating due to lack of sweat

glands. We note that although *Avahi* is nocturnal relative to *Propithecus* and *Indri*, recent work

explains their activity patterns may be more accurately described as "cathermality sensu *lato*."

Mainly this is because certain species are active across daylight hours, depending on the lunar

phase (Campera et al., 2019). Therefore, implications of the potential function of the pelage

pattern detected across the family is not solely relegated to *Propithecus* and *Indri*. However, we

note that other aspects of hair morphology that we did not measure (length, curvature) may

influence the strength of this selective pressure differently in distinct genera (Lasisi, 2021).

Future studies should consider smaller phylogenetic scales, and other aspects of hair morphology

(e.g., length) and climate (e.g., wind speed). Nonetheless, the pattern that emerges from this data

may signal that variation in hair density (i.e., higher hair density in the scalp) across the human

body may have occurred early on as an early mechanism to prevent folate leaching from key

body regions exposed to direct UV rays. This body hair pattern potentially preceded the

evolution of "naked" skin and sweat glands.

Our data is also rare evidence supporting the body cooling hypothesis—from the

perspective of hair density in non-human primates. The body cooling hypothesis (Wheeler, 1985;

1992) argued that bipedal hominins evolved naked skin to allow for evaporative cooling.

However, Wheeler (1985, 1992) also argued as part of this hypothesis that bipedal hominins

likely retained hair on the crown and shoulders to protect those areas from UV exposure.

Indriidae, presenting similar postural locomotion to humans, exhibit follicle density increases in

their body regions most exposed to potential UV damage. Indriidae present a unique opportunity

to comparatively study potential tradeoffs between folate levels, hair density, hair color, and

thermoregulation across and within populations to clarify potential mechanisms driving this

pattern of hair variation in upright primates.

Our results indicate that body size is also likely contributing to hair density variation

(Figure 5). However, the smallest primates within the Indriidae clade are also nocturnal; larger-

bodied Indriidae are diurnal. Thus, "body size" is likely confounded by activity cycle. Our

results show that within *Propithecus* alone, there was no variation in hair density detected with

body size—but there was with climate (Table 6). This may be due to a smaller sample size

available for *Propithecus* density which we estimated would not pick up small effects (G* power

analyses detailed in methods). Alternatively, at smaller phylogenetic scales, other aspects of hair

morphology that we did not measure (e.g., overall shape, length, thickness) may directly relate to

body size best at a smaller phylogenetic scale. For example, there are no differences in hair

density between chimpanzees and humans, but there are significant differences in gland density,

sweat production, and overall hair morphology (Kamberov et al., 2018). Hair and follicle density

was likely one of a few aspects of hair evolution that were potentially malleable and allowed

rapid adaptation to changing climatic environments for hominins. More research is needed across

non-human and human primate hair to understand why and how humans retained hair in the

places they did (e.g., scalp, pubic)—a critical part of the story in the evolution of our apparent

"nakedness."

Polymorphic trichromacy and pelage hue variation

Our third sub-aim sought to understand the relationship between color vision and pelage

hue in diurnal genera. We found support for our prediction that populations with enhanced

capacity for trichromatic color vision would also contain individuals with redder pelage (*as*

hypothesized in (Sumner & Mollon, 2003)). Many primates are unique among placental

mammals in having trichromatic color vision, a feature that is thought to be advantageous for

foraging on "red" foods (i.e., ripe fruit, young leaves), detecting predators, and/or tracking

conspecifics (Kawamura et al., 2012; Surridge et al., 2003). Indeed, reddish skin coloration

observed in many primate species is thought to play a salient role in conspecific signaling for

primates (Moreira et al., 2019). However, previous support for the coevolution of red hair and

trichromacy in primates is mixed (Fernandez & Morris, 2007; Kamilar et al., 2013). Our results

indicate a significant positive relationship between hue ('redness') and measures of enhanced

trichromacy in Indriidae (Table 7). The similarity in results between both models (lambda=0 vs.

lambda=1) indicates a likely significant relationship between hue and opsin genotype variation in

diurnal Indriidae. Specifically, red hues are more likely to occur when there is a larger difference

in spectral sensitivity between opsin alleles (Figure 7), suggesting increased discriminability of

such hues. Results across Indriidae, which exhibits more subtle inter- and intra-species variation

in trichromacy than has been previously considered, suggests that there may be a more complex

and nuanced coevolutionary relationship between red hair coloration in primates and

trichromatic color vision that deserves further attention (Table 7).

Conclusions

Climate, body size, and color vision all impact hair variation across Indriidae, but the

extent that each exerts its influence on the phenotype varies by sampled phylogenetic scale.

These results have considerable implications for our understanding of primate (including human)

hair evolution because it suggests that larger macro-scale trends detected across primates may

not sufficiently explain variation within a genus, species, or population. Understanding wild non-

human primate hair can also potentially help shed light on the evolution of hominin hair. Further

work on primate hair should sample across smaller geographic or phylogenetic scales and from

diverse non-human and human populations.

Acknowledgments

We thank the American Museum of Natural History, the Smithsonian National Museum of

Natural History, and Harvard University's Museum of Comparative Zoology. Special thanks to

museum staff that helped with *Propithecus* collections, including Ian Tattersall, Darrin Lunde,

Mark Omura, Eleanor Hoeger, and Marisa Surovy. We also thank the Government of Madagascar

(Ministry of the Environment and Sustainable Development), SADABE, Madagascar Biodiversity

Partnership (MBP), Omaha Zoo, the University of Antananarivo, and Anthropobiologie et

Développement for facilitation and partnership. We thank members of GW PGL and Utah PEGL

labs for relevant discussions and feedback on ideas presented here. Illustrations copyright 2020

Stephen D. Nash/IUCN SSC Primate Specialist Group, used and modified with permission. The

findings and conclusions in this article are those of the author(s) and do not necessarily represent

the views of the U.S. Fish and Wildlife Service. The work was supported by the Leakey

Foundation, the Wenner Gren Foundation, the National Science Foundation (BCS #1546730, BCS

#1606360), the George Washington University, Northern Illinois University, National Geographic

CRE, Eppley Fund for Research, Primate Conservation Inc., and IdeaWild. Field work to collect

biological samples was supported by Conservation International, Margot Marsh Biodiversity

Foundation, and The Ahmanson Foundation. The funders had no role in study design, data collection

and analysis, decision to publish, or preparation of the manuscript.

References

Allen, W. L., & Higham, J. P. (2013). Analyzing visual signals as visual scenes. *American Journal of Primatology*, *75*(7), 664–682.

Almohanna, H. M., Ahmed, A. A., Tsatalis, J. P., & Tosti, A. (2019). The role of vitamins and minerals in hair loss: A review. *Dermatology and Therapy*, *9*(1), 51–70.

Avram, M. R., & Rogers, N. E. (2009). The use of low-level light for hair growth: Part i. *Journal of Cosmetic and Laser Therapy*, *11*(2), 110–117.

Barrett, R. D. H., Laurent, S., Mallarino, R., Pfeifer, S. P., Xu, C. C. Y., Foll, M., … Hoekstra, H. E. (2019). Linking a mutation to survival in wild mice. *Science*, *363*(6426), 499–504.

Bell, R. B., Bradley, B. J., & Kamilar, J. M. (2021). The evolutionary ecology of primate hair coloration: A phylogenetic approach. *Journal of Mammalian Evolution, 28,* 911-927.

Bergman, T. J., & Beehner, J. C. (2008). A simple method for measuring colour in wild animals: validation and use on chest patch colour in geladas (*Theropithecus gelada*). *Biological Journal of the Linnean Society*, *94*(2), 231–240.

Best, A., Lieberman, D., & Kamilar, J. (2019). Diversity and evolution of human eccrine sweat gland density. *Journal of Thermal Biology*, *84*, 331–338.

Bishop, T. R., Robertson, M. P., Gibb, H., van Rensburg, B. J., Braschler, B., Chown, S. L., … Parr, C. L. (2016). Ant assemblages have darker and larger members in cold environments. *Global Ecology and Biogeography*, *25*(12), 1489–1499.

Bogert, C. M. (1949). Thermoregulation in reptiles, a factor in evolution. *Evolution*, *3*, 195–211.

Bradley, B. J., Walsh, S., Nishimura, A., Karlsson, J., & Kamilar, J. M. (2014). The evolution of primate pelage: Morphological analysis of museum research skins. *Program of the 83rd Annual Meeting of the American Association of Physical Anthropologists*, *153*(S58).

591 Campera, M., Balestri, M., Chimienti, M., Nijman, V., Nekaris, K.A.I., & Donati, G. (2019).

592   Temporal niche seperation between the two ecologically similar nocturnal primates *Avahi*

593   *meridionalis* and *Lepilemur fleuretae*. *Behavioral Ecology and Sociobiology*, *73*, 55.

594 Caro, T. (2005). The adaptive significance of coloration in mammals. *BioScience*, *55*(2), 125-

595   136.

596 Chaplin, George, Jablonski, Nina G., & Cable, T. N. (1994). Physiology, thermoregulation and

597   bipedalism. *Journal of Human Evolution*, *27*(6), 497–510.

598 Clusella Trullas, S., van Wyk, J. H., & Spotila, J. R. (2007). Thermal melanism in ectotherms.

599   *Journal of Thermal Biology*, *32*(5), 235–245.

600 Cuthill, I. C., Allen, W. L., Arbuckle, K., Caspers, B., Chaplin, G., Hauber, M. E., … Caro, T.

601   (2017). The biology of color. *Science*, *357*(6350), aan0221.

602 da Silva, L. G., de Oliveira, T. G., Kasper, C. B., Cherem, J. J., Moraes, E. A., Paviolo, A., &

603   Eizirik, E. (2016). Biogeography of polymorphic phenotypes: Mapping and ecological

604   modelling of coat colour variants in an elusive Neotropical cat, the jaguarundi (*Puma*

605   *yagouaroundi*). *Journal of Zoology*, *299*(4), 295–303.

606 Delhey, K. (2017). Gloger's rule. *Current Biology*, *27*(14), R689–R691.

607 Delhey, K. (2018). Darker where cold and wet: Australian birds follow their own version of

608   Gloger's rule. *Ecography*, *41*(4), 673–683.

609 Delhey, K., Dale, J., Valcu, M., & Kempenaers, B. (2019). Reconciling ecogeographical rules:

610   rainfall and temperature predict global colour variation in the largest bird radiation. *Ecology*

611   *Letters*, *22*(4), 726–736.

612 Durusoy, C., Ozenli, Y., Adiguzel, A., Budakoglu, I. Y., Tugal, O., Arikan, S., & Uslu, A.

613   (2009). The role of psychological factors and serum zinc, folate and vitamin B 12 levels in

the aetiology of trichodyni: A case–control stud*y*. *Clinical and Experimental Dermatology,*

34(7), 789–792.

Dutton, A., Christopher, J., & Randall, E. (2003). Biomedical evaluation of free-ranging ring-

tailed lemurs (*Lemur catta*) in Tsimanampetsotsa Strict Nature Reserve, Madagascar.

Journal of Zoo and Wildlife Medicine, *34*(1), 16–24.

Ertug, E. Y. (2018). Reduced ferritin, folate, and vitamin B12 levels in female patients diagnosed

with telogen effluvium. *International Journal of Medical Biochemistry, 1*(3), 0–2.

Faul, F., Erdfelder, E., Buchner, A., & Lang, A. G. (2009). Statistical power analyses using

G*Power 3.1: Tests for correlation and regression analyses. *Behavior Research Methods*,

41(4), 1149–1160.

Fernandez, A. A., & Morris, M. R. (2007). Sexual selection and trichromatic color vision in

primates: Statistical support for the preexisting-bias hypothesis. *American Naturalist*,

170(1), 10–20.

Fratto, M. A., & Davis, A. K. (2011). Do black-furred animals compensate for high solar

absorption with smaller hairs? A test with a polymorphic squirrel species. *Current Zoology*,

57(6), 731–736.

Galván, I., Rodríguez-Martínez, S., & Carrascal, L. M. (2018). Dark pigmentation limits thermal

niche position in birds. *Functional Ecology*, *32*(6), 1531–1540.

Glogger, C.W.L. (1833). Das Abändern der Vögel durch Einfluss des Kilmas. Breslau: August

Schulz. & Co.

Godfrey, L.R., Samonds, K.E., Jungers, W.L., & Sutherland, M.R. (2001). Teeth, brains, and

primate life histories. *American Journal of Physical Anthropology*, *114*(3), 192-214.

Guevara, E.E. (2018). Genomics of primate longevity (Publication No. 13141656). [Doctoral

dissertation, Yale University]. ProQuest Dissertations and Theses.

Hadfield, J. D. (2010). MCMC methods for multi-response generalized linear mixed models: The

MCMCglmm R package. *Journal of Statistical Software*, *33*(2), 1–22.

Hansen, T. F., & Orzack, S. H. (2005). Assessing current adaptation and phylogenetic inertia as

explanations of trait evolution: The need for controlled comparisons. *Evolution*, *59*(10),

2063-2072.

Herrera, J. P., & Dávalos, L. M. (2016). Phylogeny and divergence times of lemurs inferred with

recent and ancient fossils in the tree. *Systematic Biology*, *65*(5), 772–791.

Hijmans, R. J., & Etten, J. V. (2012). Raster: Geographic analysis and modeling with raster data.

R package version 2.0-12.

Hoekstra, H. E., Krenz, J. G., & Nachman, M. W. (2005). Local adaptation in the rock pocket

mouse (*Chaetodipus intermedius*): Natural selection and phylogenetic history of

populations. *Heredity*, *94*(2), 217–228.

Irwin, M. T., Junge, R. E., Raharison, J. L., & Samonds, K. E. (2010). Variation in physiological

health of diademed sifakas across intact and fragmented forest at Tsinjoarivo, eastern

Madagascar. *American Journal of Primatology*, *72*(11), 1013–1025.

Jablonski, N. G., & Chaplin, G. (2000). The evolution of human skin coloration. *Journal of*

Human Evolution, *39*(1), 57–106.

Jacobs, R. L., MacFie, T. S., Spriggs, A. N., Baden, A. L., Morelli, T. L., Irwin, M. T., …

Bradley, B. J. (2017). Novel opsin gene variation in largebodied, diurnal lemurs. *Biology*

Letters, *13*(3), 1–6.

Jacobs, R. L., Spriggs, A. N., MacFie, T. S., Baden, A. L., Irwin, M. T., Wright, P. C., …

Bradley, B. J. (2016). Primate genotyping via high resolution melt analysis: rapid and

reliable identification of color vision status in wild lemurs. *Primates*, *57*(4), 541–547.

Jacobs, R. L., Veilleux, C. C., Louis, E. E., Herrera, J. P., Hiramatsu, C., Frankel, D. C., …

Bradley, B. J. (2019). Less is more: lemurs (*Eulemur* spp.) may benefit from loss of

trichromatic vision. *Behavioral Ecology and Sociobiology*, *73*, 22.

Kamberov, Y. G., Guhan, S. M., DeMarchis, A., Jiang, J., Wright, S. S., Morgan, B. A., …

Lieberman, D. E. (2018). Comparative evidence for the independent evolution of hair and

sweat gland traits in primates. *Journal of Human Evolution*, *125*, 99–105.

Kamilar, J. M., & Bradley, B. J. (2011). Interspecific variation in primate coat colour supports

Gloger's rule. *Journal of Biogeography*, *38*(12), 2270–2277.

Kamilar, J. M., & Cooper, N. (2013). Phylogenetic signal in primate behaviour, ecology and life

history. *Philosophical Transactions of the Royal Society of London. Series B, Biological*

Sciences, *368*, 20120341.

Kamilar, J. M., Heesy, C. P., & Bradley, B. J. (2013). Did trichromatic color vision and red hair

color coevolve in primates? *American Journal of Primatology, 75*(7), 740-751.

Kamilar, J. M., & Muldoon, K. M. (2010). The climatic niche diversity of Malagasy primates: A

phylogenetic perspective. *PLoS ONE*, *5*(6), e11073.

Kawamura, S., Hiramatsu, C., Melin, A. D., Schaffner, C. M., Aureli, F., & Fedigan, L. M.

(2012). Polymorphic color vision in primates: Evolutionary considerations. In H. Hirai, H.

Imai, & Y. Go (Eds.), *Post-Genome Biology of Primates* (pp. 93-120). Tokyo: Springer.

Kendal, D., Hauser, C. E., Garrard, G. E., Jellinek, S., Giljohann, K. M., & Moore, J. L. (2013).

Quantifying plant colour and colour difference as perceived by humans using digital

images. *PLoS ONE*, *8*(8), e72296.

King, T., Rakotonirina, L. H. F., Rakotoarisoa, A. H., Razafindramanana, J., & Ratsimbazafy, J.

(2014). Distributional limits and melanism in the south-west of the range of the crowned

sifaka (*Propithecus coronatus*), Madagascar. *Primate Conservation*, *28*(1), 55–64.

Lasisi, T., Ito, S., Wakamatsu, K., & Shaw, C. N. (2016). Quantifying variation in human scalp

hair fiber shape and pigmentation. *American Journal of Physical Anthropology*, *160*(2),

341–352.

Lasisi, T. (2021). The genetic architecture and evolutionary function of human scalp hair

morphology (Publication No. 28841761). [Doctoral dissertation, Penn State University].

ProQuest Dissertations and Theses.

Lawler, R.R., Caswell, H., Richard, A.F., Ratsirarson, J., Dewar, R.E., & Schwartz, M. 2009.

Demography of Verreauix's sifaka in a stochastic rainfall environment. *Oecologia*, 161,

491-504.

Lehman, S. M. (2007). Ecological and phylogenetic correlates to body size in the Indriidae.

International Journal of Primatology, *28*(1), 183–210.

Lewis, R. J., & Van Schaik, C. P. (2007). Bimorphism in male verreaux's sifaka in the Kirindy

Forest of Madagascar. *International Journal of Primatology*, *28*(1), 159–182.

Losos, J. B. (2008). Phylogenetic niche conservatism, phylogenetic signal and the relationship

between phylogenetic relatedness and ecological similarity among species. *Ecology Letters*,

11(10), 995–1003.

Lynch, M. (1991). Methods for the analysis of comparative data in evolutionary biology.

Evolution, *45*(5), 1065–1080.

Martin, P.R., Montgomerie, R., Lougheed, S.C. (2010). Rapid sympatry explains greater color

pattern divergence in high latitude birds. *Evolution, 64*(2), 336-347.

McLean, M. J., Bishop, P. J., & Nakagawa, S. (2013). Assessing the patterns of evolution in

anuran vocal sexual signals. *Evolutionary Biology*, *40*(1), 141–149.

Mittermeier, Russell A., Louis, Jr., Edward E., Richardson, M., Schwitzer, C., Langrand, O.,

Rylands, Anthony B., … MacKinnon, K. C. (Eds.) (2010). *Lemurs of Madagascar* (3rd ed.).

New York: Conservation International.

Moreira, L. A. A., Duytschaever, G., Higham, J. P., & Melin, A. D. (2019). Platyrrhine color

signals: New horizons to pursue. *Evolutionary Anthropology*, *28*(5), 236–248.

Norton, H. L., Edwards, M., Krithika, S., Johnson, M., Werren, E. A., & Parra, E. J. (2016).

Quantitative assessment of skin, hair, and iris variation in a diverse sample of individuals

and associated genetic variation. *American Journal of Physical Anthropology*, *160*(4), 570–

581.

Osorio, D., Smith, A. C., Vorobyev, M., & Buchanan-Smith, H. M. (2004). Detection of fruit

and the selection of primate visual pigments for color vision. *American Naturalist*, *164*(6),

696–708.

Paus, R., & Foitzik, K. (2004). In search of the "hair cycle clock": A guided tour. *Differentiation*,

72(9–10), 489–511.

Pebesma, E. J., & Bavand, R. S. (n.d.). Classes and methods for spatial data in R. *R News*, *5*(2),

9–13.

Petter, J. J., & Peyrieras, A. (1972). Melanization in the genus *Propithecus* Malagasey Lemur.

Journal of Human Evolution, *1*(4), 379–388.

Post, P. W., Szabó, G., & Keeling, M. E. (1975). A quantitative and morphological study of the

pigmentary system of the chimpanzee with the light and electron microscope. *American*

Journal of Physical Anthropology, *43*(3), 435–443.

Rakotonirina, H., Kappeler, P. M., & Fichtel, C. (2017). Evolution of facial color pattern

complexity in lemurs. *Scientific Reports*, *7*(1), 1–11.

Rakotonirina, L. H. F., Randriantsara, F., Rakotoarisoa, A. H., Rakotondrabe, R.,

Razafindramanana, J., Ratsimbazafy, J., & King, T. (2014). A preliminary assessment of

sifaka (*Propithecus*) distribution, chromatic variation and conservation in Western Central

Madagascar. *Primate Conservation*, *28*(1), 43–53.

Ramírez-Delgado, V. H., Sanabria-Urbán, S., Serrano-Meneses, M. A., & Cueva del Castillo, R.

(2016). The converse to Bergmann's rule in bumblebees, a phylogenetic approach. *Ecology*

and Evolution, *6*(17), 6160–6169.

Revell, L. J., Harmon, L. J., & Collar, D. C. (2008). Phylogenetic signal, evolutionary process,

and rate. *Systematic Biology*, *57*(4), 591–601.

Rowe, N. (1996). *The pictoral guide to the living primates*. East Hampton: Pagonias Press.

Sambrook, J., Fritch, E., & Maniatus, T. (1989). *Molecular cloning: a laboratory manual.* (2nd

ed.). New York: Cold Spring Harbor Press.

Sandel, A. A. (2013). Brief communication: Hair density and body mass in mammals and the

evolution of human hairlessness. *American Journal of Physical Anthropology*, *152*(1), 145–

150.

Santana, S. E., Alfaro, J. L., Noonan, A., & Alfaro, M. E. (2013). Adaptive response to sociality

and ecology drives the diversification of facial colour patterns in catarrhines. *Nature*

Communications, *4*, 1–7.

Santana, S. E., Lynch Alfaro, J., & Alfaro, M. E. (2012). Adaptive evolution of facial colour

patterns in Neotropical primates. *Proceedings. Biological Sciences / The Royal Society*,

279(1736), 2204–2211.

Schmidt, M. (2011). Locomotion and postural behaviour. *Advances in Science and Research*,

5(1), 23–39.

Schwartz, G. G., & Rosenblum, L. A. (1981). Allometry of primate hair density and the

evolution of human hairlessness. *American Journal of Physical Anthropology*, *55*(1), 9–12.

Seibert, H. C., & Steggerda, M. (1999). The size and shape of human head hair: Along its shaft.

The Journal of Heredity, *33*(8), 302–304.

Shearer, C. L., & Ezenwa, V. O. (2020). Rainfall as a driver of seasonality in parasitism.

International Journal for Parasitology: Parasites and Wildlife, *12*, 8–12.

Spriggs, A. N. (2017). Evolution of lemur pelage color variation (Publication No. 10682576).

[Doctoral dissertation, University at Albany, State University of New York]. ProQuest

Dissertations and Theses..

Steggerda, M., & Seibert, H. C. (1941). Size and shape of head hair from six racial groups.

Journal of Heredity, *32*(9), 315–318.

Stevens, M., Párraga, C. A., Cuthill, I. C., Partridge, J. C., & Troscianko, T. S. (2007). Using

digital photography to study animal coloration. *Biological Journal of the Linnean Society*,

90(2), 211–237.

Stevens, M., Stoddard, M. C., & Higham, J. P. (2009). Studying primate color: towards visual

system-dependent methods. *International Journal of Primatology*, *30*(6), 893–917.

Sumner, P., & Mollon, J. D. (2003). Colors of primate pelage and skin: Objective assessment of

conspicuousness. *American Journal of Primatology*, *59*(2), 67–91.

Surridge, A. K., Osorio, D., & Mundy, N. I. (2003). Evolution and selection of trichromatic

vision in primates. *Trends in Ecology and Evolution*, *18*(4), 198–205.

Tapanes, Elizabeth, Kamilar, Jason M, & Bradley, B. J. (2021). Molecular and cellular processes

of pelage growth and pigmentation in primate evolution. In K.M. Pitirri, J.T. & Richtsmeier

(Eds.). *Evolutionary Cell Processes in Primates* (pp. 1–33). Boca Raton: CRC Press.

Tapanes, E., Irwin, M., Spriggs, Amanda Nicole, Kamilar, Jason M., & Bradley, B. J. (2018).

Subtle sexual dichromatism and dimorphism detected in wild *Propithecus diadema*.

Program of the 87th Annual Meeting of the American Association of Physical

Anthropologists, *165*(S66).

Tattersall, I. (1986). Notes on the distribution and taxonomic status of some species of

Propithecus in Madagascar. *Folia Primatologica*, *46*, 51–63.

Villemereuil, P. de, & Nakagawa, S. (2014). General quantitative genetic methods for

comparative biology. In L.Z. Garamszegi (Ed.). *Modern Phylogenetic Comparative*

Methods and their Application in Evolutionary Biology (pp. 287–304). Berlin: Springer.

Wheeler, P. E. (1985). The loss of functional body hair in man: the influence of thermal

environment, body form and bipedality. *Journal of Human Evolution*, *14*(1), 23–28.

Wheeler, P. E. (1992). The influence of the loss of functional body hair on the water budgets of

early hominids. *Journal of Human Evolution*, *23*(5), 379–388.

Winters, S., Allen, W., & Higham, J. (2020). The structure of species discrimination signals

across a primate radiation. eLife, 9:e47428.

Xing, S., Bonebrake, T. C., Tang, C. C., Pickett, E. J., Cheng, W., Greenspan, S. E., …

Scheffers, B. R. (2016). Cool habitats support darker and bigger butterflies in Australian

tropical forests. *Ecology and Evolution*, *6*(22), 8062–8074.

Figure Legends

Figure 1 Indriidae species ranges and sampling localities for (A) *Avahi*, (B) *Propithecus*, and (C)

Indri; along with a (D) cladogram, and (E) Koppen-climate zones that span the genera's

distribution.

Key: *Avahi* spp. = (1a) Betampona, (1b) Sihanaka forest, (1c) Andasibe, (2) Forest west of Fort Dauphin, (3)

Vondrozo, (4) Ankarafatsika; *Propithecus* spp. = (5a) Betampona, (5b) Zahamahena, (5c) Didy, (5d) Sihanaka

forest, (5e) forest patch near Andasibe, (5f) Tsinjoarivo (eastern and western), (6a) Nandihizana, (6b) East of

Farohny River, (7a) Mandrare Valley, (7b) Bevala East, (7c) Tsinanampetsotsa, (7d) Lower Ifaty, (7e) Ifaty, (7f)

Lake Ihortry, (7g) Ankazoabo West, (7h) Upper Tsirbihina River, (8) Ambararatabe, (9a) Tsingy de Bemaraha, (9b)

Tsingy de Namoraka, (9c) Baie de Baly Nat Park, (10) Reserve of Analamerana, (11) Forest fragment near Daraina,

(12) Anjanaharibe-Sud Reserve; *Indri* indri = (13a) Anjanaharibe-Sud Reserve, (13b) 40 km West of Maroantsetra,

(13c) Sihanaka forest

Figure 2 Plot of brightness across (A) the entire Indriidae family, in association with

precipitation Seasonality (CV) (showing adherence to Gloger's Rule), and (B) only within the

Propithecus genus, in association with Annual Mean Temperature (°C) (showing adherence to

Bogert's Rule).

Figure 3 Plot of pelage hue trends across (A) the entire Indriidae family, in association with

precipitation seasonality (mm) (showing adherence to Gloger's Rule). Results are similar within

Propithecus.

Figure 4 Plot of hair density trends. The relationship between PC1 and precipitation seasonality

(CV) is significantly associated with hair density on the limbs and crown across the Indriidae

family. Lower PC1 values are associated with lower hair density on the limbs and crown.

Figure 5 Plot of hair density trends. The relationship between PC3 and body length (capturing

length, in mm) is significantly associated with hair density on the dorsal torso across Indriidae.

Lower PC3 values are associated with lower hair density across the dorsal torso.

Figure 6 Boxplots illustrating variation in (A) hair density and (B) follicle density across body

regions, for Indriidae. Black lines represent the average, and black dots represent outliers.

Figure 7 Plot of pelage hue trends across diurnal *Propithecus* and *Indri* genera, in association

with the range (i.e., difference) in opsin peak spectral sensitivity.

Table 1 Results of PGLMMs predicting hair brightness (RGB averages) across all Indriidae taxa

and only within *Propithecus*

Variable	Mean Est.	Lower CI	Upper CI	ESS	P MCMC
Across Indriidae (PC1; $H^2 = 0.666$)					
Intercept	-5.901	-15.911	4.432	9,000	0.242
BIO01	-0.540	-1.510	0.401	9,000	0.256
BIO04	0.901	-0.492	2.290	9,000	0.216
BIO07	-0.393	-2.264	1.396	9,000	0.662
BIO13	0.293	-0.171	0.762	9,000	0.218
BIO15	0.728	-0.016	1.431	9,000	**0.048**
Body size	0.010	-0.727	0.696	9,000	0.972
Within *Propithecus* (PC1; $H^2 = 0.902$)					
Intercept	0.710	-8.640	10.541	9,563	0.900
BIO01	-1.291	-2.064	-0.571	9,000	**0.002**
BIO04	-0.501	-1.920	0.863	9,410	0.471
BIO07	0.613	-0.921	2.176	9,000	0.430
BIO13	0.001	-0.398	0.436	9,000	0.999
BIO15	0.347	-0.289	0.988	9,000	0.284
Body size	0.527	-0.051	1.103	9,000	0.075[!]

Bold values indicate $p < 0.05$, [!] indicates $p < 0.10$, H^2 is akin to phylogenetic signal

859 **Table 2** Results of PGLMMs predicting hair hue (RG ratio values) across all Indriidae taxa

Variable	Mean Est.	Lower CI	Upper CI	ESS	P MCMC
PC1; $H^2 = 0.475$					
Intercept	-0.993	-9.445	7.609	9,000	0.817
BIO01	-0.221	-1.170	0.693	10,089	0.629
BIO04	0.126	-0.904	1.258	9,000	0.815
BIO07	-0.329	-1.858	1.265	9,000	0.688
BIO13	-0.047	-0.442	0.325	9,000	0.803
BIO15	0.860	0.198	1.515	9,000	**0.010**
Body size	-0.030	-0.707	0.658	9,000	0.921
PC2; $H^2 = 0.371$					
Intercept	3.750	-6.223	13.216	10,53	0.447
BIO01	0.778	-0.299	1.863	9,310	0.162
BIO04	-1.134	-2.341	0.094	9,441	0.061[!]
BIO07	0.959	-0.802	2.644	9,000	0.278
BIO13	-0.327	-0.740	0.123	9,955	0.134
BIO15	-0.751	-1.500	-0.040	9,000	**0.048**
Body size	0.523	-0.226	1.276	9,000	0.170
PC3; $H^2 = 0.255$					
Intercept	2.830	-6.910	13.510	9,000	0.581
BIO01	-0.127	-1.341	0.970	9,000	0.831
BIO04	-0.303	-1.471	0.916	9,000	0.615
BIO07	0.455	-1.272	2.176	9,372	0.592
BIO13	-0.072	-0.508	0.374	9,289	0.737
BIO15	-0.176	-0.940	0.553	8,955	0.637
Body size	0.052	-0.751	0.858	9,000	0.894

860 Bold values indicate significance below 0.05, [!] indicates p < 0.10, H^2 is akin to phylogenetic

861 signal.

866 **Table 3** Results of PGLMMs predicting hair hue (RG ratio values) within the *Propithecus* genus

Variable	Mean Est.	Lower CI	Upper CI	ESS	P MCMC
PC1; $H^2 = 0.632$					
Intercept	7.664	-5.813	21.243	8,635	0.268
BIO01	-0.341	-1.656	0.811	9,000	0.584
BIO04	-0.567	-2.530	1.366	8,581	0.565
BIO07	0.751	-1.599	3.290	8,631	0.548
BIO13	0.004	-0.612	0.607	8,699	0.984
BIO15	-0.969	-2.021	0.057	8,524	0.069[!]
Body size	-0.083	-1.063	0.831	9,375	0.857
PC2; $H^2 = 0.513$					
Intercept	1.773	-8.206	11.490	8,358	0.728
BIO01	1.434	0.448	2.381	9,322	**0.006**
BIO04	-1.220	-2.587	0.153	8,042	0.076[!]
BIO07	1.154	-0.598	2.923	9,000	0.187
BIO13	-0.450	-0.907	-0.004	9,000	**0.045**
BIO15	-1.089	-1.862	-0.314	9,000	**0.006**
Body size	0.806	0.047	1.531	8,829	**0.037**
PC3; $H^2 = 0.227$					
Intercept	3.296	-0.121	1.789	9,395	0.639
BIO01	-0.429	-2.065	1.199	9,000	0.596
BIO04	-0.155	-1.744	2.085	9,492	0.903
BIO07	-0.753	-3.283	1.704	9,765	0.545
BIO13	-0.008	-0.643	-0.637	9,000	0.975
BIO15	0.001	-1.241	1.213	9,325	1.000
Body size	-0.148	-1.166	1.471	9,000	0.812

867 Bold values indicate significance below 0.05, [!] indicates p < 0.10, H^2 is akin to phylogenetic

868 signal.

875
876
877

878   **Table 4** Results of PGLMMs predicting hair and follicle density across all Indriidae taxa

Variable	Mean Est.	Lower CI	Upper CI	ESS	P MCMC
PC1; H^2 = 0.222					
Intercept	-6.478	-22.987	18.456	9,456	0.401
BIO01	-0.036	-1.368	1.366	9,000	0.961
BIO04	0.770	-0.673	2.228	9,000	0.278
BIO07	-1.093	-3.262	1.047	9,434	0.314
BIO13	0.523	-0.085	1.190	9,000	0.101
BIO15	1.232	0.185	2.255	9,290	**0.020**
Body size	-0.275	-1.326	0.770	9,339	0.597
PC2; H^2 = 0.217					
Intercept	0.256	-15.042	16.829	9,000	0.981
BIO01	0.175	-1.238	1.587	9,000	0.812
BIO04	0.150	-1.334	1.643	9,000	0.835
BIO07	-0.990	-3.299	1.243	9,000	0.375
BIO13	0.110	-0.523	0.806	9,000	0.736
BIO15	0.518	-0.550	1.585	9,765	0.331
Body size	-0.161	-1.266	0.922	8,309	0.760
PC3; H^2 = 0.330					
Intercept	10.259	-0.399	20.576	9,000	0.057[!]
BIO01	-0.259	-1.175	0.700	9,000	0.602
BIO04	0.071	-0.953	1.056	9,307	0.876
BIO07	-0.129	-1.643	1.413	9,000	0.853
BIO13	-0.337	-0.772	0.081	9,000	0.123
BIO15	0.438	-0.276	1.164	9,908	0.228
Body size	-1.265	-2.012	-0.514	9,000	**0.002**

879   Bold values indicate significance below 0.05, [!] indicates p < 0.10, H^2 is akin to phylogenetic signal.

882   **Table 5** Summary of ANOVA examining hair and follicle density variation across distinct body

883   regions for Indriidae

	DF	Sum Sq.	Mean Sq.	F value	P value
Hair density					
Body Region	4	1458536	364634	5.808	**< 0.0005**
Residuals	60	3766879	62781		
Follicle density					
Body Region	4	108592	27148	1.524	0.207
Residuals	60	1068659	17811		

884
885

Table 6 Results of PGLMMs predicting hair and follicle density within the *Propithecus* genus

Variable	Mean Est.	Lower CI	Upper CI	ESS	P MCMC
PC1; $H^2 = 0.256$					
Intercept	16.640	-6.426	39.799	9,000	0.142
BIO01	-1.256	-3.056	0.696	8,178	0.180
BIO04	-1.333	-3.506	0.890	8,746	0.234
BIO07	2.437	-1.147	5.812	9,286	0.176
BIO13	-0.485	-1.359	0.364	9,000	0.259
BIO15	-1.673	-3.105	-0.200	9,423	**0.028**
Body size	-0.220	-2.195	1.495	9,000	0.802
PC2; $H^2 = 0.190$					
Intercept	9.252	-17.259	37.827	9,000	0.494
BIO01	1.437	-0.749	3.688	9,000	0.192
BIO04	0.351	-2.135	2.597	9,000	0.760
BIO07	-0.042	-3.693	3.627	9,571	0.988
BIO13	-0.090	-1.141	0.898	9,356	0.862
BIO15	-0.508	-2.082	1.157	9,000	0.530
Body size	-1.632	-3.695	0.666	9,000	0.134
PC3; $H^2 = 0.279$					
Intercept	-12.391	-31.523	5.691	9,000	0.186
BIO01	0.766	-0.815	2.345	9,000	0.330
BIO04	1.090	-0.802	2.988	9,000	0.224
BIO07	-1.264	-4.442	1.564	9,000	0.382
BIO13	0.437	-0.282	1.117	9,000	0.219
BIO15	0.309	-0.864	1.559	9,000	0.606
Body size	0.738	-0.698	2.194	9,000	0.310

Bold values indicate significance below 0.05, $^!$ indicates $p < 0.10$, H^2 is akin to phylogenetic signal.

919　**Table 7** Summary of linear models examining pelage coloration against difference in opsin peak

920　spectral sensitivity, using hue (PC1) scores of individuals across a subset of 10 sampling

921　locations

	Estimate	Std. Error	T value	P value
PC1				
[Intercept]	-1.529	1.611	-0.949	0.374
Difference in Spectral Sensitivity	-0.234	0.089	-2.630	**0.034**
Number of Opsin Alleles	2.635	1.501	1.746	0.124

922　SE: 1.05 on 7 degrees of freedom; R^2 = 0.516, F = 5.801 on 2 and 7 degrees of freedom; p = 0.033. Bold values

923　indicate significance below 0.05.

924
925