1

Prefrontal cortex supports speech perception in listeners with cochlear implants

Arefeh Sherafati[1], Noel Dwyer[2], Aahana Bajracharya[2], Mahlega S. Hassanpour[3],
Adam T. Eggebrecht[1, 4, 5, 6], Jill B. Firszt[2], Joseph P. Culver[1, 5, 6, 7], Jonathan E. Peelle[2*]

[1] Department of Radiology, Washington University in St. Louis, St. Louis, MO, USA

[2] Department of Otolaryngology, Washington University in St. Louis, St. Louis, MO, USA

[3] Moran Eye Center, University of Utah, Salt Lake City, UT, USA

[4] Department of Electrical & Systems Engineering, Washington University in St. Louis, St. Louis, MO, USA

[5] Department of Biomedical Engineering, Washington University in St. Louis, St. Louis, MO, USA

[6] Division of Biology and Biomedical Sciences, Washington University in St. Louis, St. Louis, MO, USA

[7] Department of Physics, Washington University in St. Louis, St. Louis, MO, USA

[*]jpeelle@wustl.edu

24   **Abstract**

Cochlear implants are neuroprosthetic devices that can restore hearing in people with severe to
profound hearing loss by electrically stimulating the auditory nerve. Because of physical
limitations on the precision of this stimulation, the acoustic information delivered by a cochlear
implant does not convey the same level of acoustic detail as that conveyed by normal hearing.
As a result, speech understanding in listeners with cochlear implants is typically poorer and more
effortful than in listeners with normal hearing. The brain networks supporting speech
understanding in listeners with cochlear implants are not well understood, partly due to difficulties
obtaining functional neuroimaging data in this population. In the current study, we assessed the
brain regions supporting spoken word understanding in adult listeners with right unilateral
cochlear implants (n=20) and matched controls (n=18) using high-density diffuse optical
tomography (HD-DOT), a quiet and non-invasive imaging modality with spatial resolution
comparable to that of functional MRI. We found that while listening to spoken words in quiet,
listeners with cochlear implants showed greater activity in left prefrontal cortex than listeners with
normal hearing, specifically in a region engaged in a separate spatial working memory task. These
results suggest that listeners with cochlear implants require greater cognitive processing during
speech understanding than listeners with normal hearing, supported by compensatory
recruitment of left prefrontal cortex.

Introduction

Cochlear implants (CIs) are neuroprosthetic devices that can restore hearing in people with severe to profound hearing loss by electrically stimulating the auditory nerve. Because of physical limitations on the precision of this stimulation—including, for example, the spatial spread of electrical current (Garcia et al., 2021)—the auditory stimulation delivered by a CI does not convey the same level of acoustic detail as normal hearing. As a result, speech understanding in listeners with CIs is poorer than in listeners with normal hearing (Firszt et al., 2004). Notably, even in quiet, listeners with CIs report increased effort during listening (Dwyer et al., 2014). In spite of these challenges, many listeners with CIs attain significant success in understanding auditory speech. This remarkable success raises the question of how listeners with CIs make sense of a degraded acoustic signal.

One area of key importance is understanding the degree to which listeners with CIs rely on nonlinguistic cognitive mechanisms to compensate for a degraded acoustic signal. In listeners with normal hearing, cognitive demands are increased when speech is acoustically challenging (Peelle, 2018). For example, even when speech is completely intelligible, acoustically-degraded speech can reduce later memory for what has been heard (Cousins et al., 2014; Koeritzer et al., 2018; Rabbitt, 1968; Ward et al., 2016). These findings suggest that to understand acoustically challenging speech, listeners need to engage domain-general cognitive resources during perception. In a limited-capacity cognitive system (Wingfield, 2016), such recruitment necessarily reduces the resources available for other tasks, including memory encoding. Importantly, even speech presented in quiet (that is, without background noise) is degraded by the time it reaches the auditory system of a listener with a CI.

Cognitive demands during speech understanding are supported by several brain networks that supplement classic frontotemporal language regions. The cingulo-opercular network, for example, is engaged during particularly challenging speech (Eckert et al., 2009; Vaden et al., 2017) and supports successful comprehension during difficult listening (Vaden et al., 2013). Prefrontal cortex (PFC) complements the cingulo-opercular network and varies parametrically with speech intelligibility (Davis & Johnsrude, 2003). Activity in PFC, particularly dorsolateral regions, is associated with cognitive demands in a wide range of tasks (Duncan, 2010), consistent with domain-general cognitive control (Braver, 2012). We thus hypothesized that listeners with CIs would rely more on PFC during listening than listeners with normal hearing, and in particular regions of PFC associated with non-linguistic tasks. However, the functional anatomy of PFC is also complex (Noyce et al., 2017), and dissociating nearby language and domain-general processing regions is challenging (Fedorenko et al., 2012).

A central question concerns the degree to which listeners with CIs rely on cognitive processing outside core speech processing regions, such as dorsolateral PFC. Obtaining precise spatially-localized images of regional brain activity has been difficult in listeners with CIs, given that functional MRI is not possible (or subject to artifact) due to the CI hardware. Thus, optical brain imaging (Peelle, 2017) has become a method of choice for studying functional activity in CI listeners (Anderson et al., 2017; Lawler et al., 2015; Lawrence et al., 2018; Olds et al., 2016; Zhou et al., 2018). In the current study, we use high-density diffuse optical tomography (HD-DOT) (Eggebrecht et al., 2014; Zeff et al., 2007), previously validated in speech studies in listeners with normal hearing (Hassanpour et al., 2015; Hassanpour et al., 2017; Schroeder et al., 2020). HD-DOT provides high spatial resolution and homogenous sensitivity over the field of view that captures known speech-related brain regions (White & Culver, 2010). We examine the brain regions supporting single word processing in listeners with a right unilateral CI relative to that in a group of matched, normal-hearing controls. We hypothesized that listeners with CIs would exhibit greater recruitment of PFC compared to normal-hearing controls.

Methods

Data and code availability
Stimuli, summary data, and analysis scripts are available from https://osf.io/nkb5v/.

Subjects
We recruited 21 adult CI patients aged 26–79 years (17 right-handed, 2 left-handed, 2 not available), and 19 age- and sex-matched controls (18 right-handed, 1 left-handed) (demographic information in Table 1). We excluded one CI user due to poor signal quality (evaluated as mean band limited SNR of all source-detectors) and one control due to excessive motion (see Appendix 1 for details). All CI patients had a unilateral right CI (manufacturer: 11 Cochlear, 6 Advanced Bionics, 3 Med El). All subjects were native speakers of English with no self-reported history of neurological or psychiatric disorders. All aspects of these studies were approved by the Human Research Protection Office of the Washington University School of Medicine. Subjects were recruited from the Washington University campus and the surrounding community (IRB 201101896, IRB 201709126). All subjects gave informed consent and were compensated for their participation in accordance with institutional and national guidelines.

Table 1: Demographic information.

Measure	Control	CI users
Number of subjects (# of females)	18 (11)	20 (11)
Mean age at test in years (std)	57.57 (12.74)	56.80 (14.09)
Mean years of CI use (std)	–	8.10 (6.51)
Mean speech perception score (AzBio sentences in quiet) (std)	0.99 (0.01)	0.88 (0.09)
Mean right ear 4fPTA (std)	16.02 (6.74)	21.85 (5.30) with CI on
Mean left ear 4fPTA (std)	16.61 (7.67)	91.25 (26.77) unaided
Mean left ear 4fPTA at test[1] (std)	–	73.28 (37.72)
Mean duration of deafness right ear	–	12.58 (11.74)

4fPTA, average pure tone threshold at 4 frequencies (500, 1000, 2000, 4000 Hz).
If no response at a given frequency, a value of 120 dB HL was assigned.
[1] With hearing aid, if the subject used amplification. Eight out of 20 CI users used hearing aids.

HD-DOT system
Data were collected using a continuous-wave HD-DOT system comprised of 96 sources (LEDs, at both 750 and 850 nm) and 92 detectors (coupled to avalanche photodiodes, APDs, Hamamatsu C5460-01) to enable oxy and deoxyhemoglobin spectroscopy (Figure 1) (Eggebrecht et al., 2014). The design of this HD-DOT system provides more than 1200 usable source-detector measurements per wavelength at a 10 Hz full-field frame rate. This system has been validated for successfully mapping cortical responses to language and naturalistic stimuli with comparable sensitivity and specificity to fMRI (Eggebrecht et al., 2014; Fishell et al., 2019; Hassanpour et al., 2015).

Figure 1 | HD-DOT system and the experimental design. (A) Schematic of a subject wearing the HD-DOT cap along with an illustration of the task design. **(B)** Simplified illustration of the HD-DOT system (far left), regional distribution of source-detector light levels (middle), and source-detector pair measurements (~1200 pairs) as gray solid lines illustrated in a flat view of the HD-DOT cap (far right). **(C)** An example point-spread-function (PSF) and the HD-DOT sensitivity profile illustrated on the volume and spatially registered on the cortical view of a standard atlas in lateral and posterior views.

Experimental design

Subjects were seated on a comfortable chair in an acoustically isolated room facing an LCD screen located 76 cm from them, at approximately eye level. The auditory stimuli were presented through two speakers located approximately 150 cm away at about +/- 21º from the subjects' ears at a sound level of approximately 65 dBA. Subjects were instructed to fixate on a white crosshair against a gray background while listening to the auditory stimuli, holding a keyboard on their lap for the stimuli that required their response (Figure 1A, left panel). The HD-DOT cap was fitted to the subject's head to maximize optode-scalp coupling, assessed via real-time coupling coefficient readouts using in-house software. The stimuli were presented using Psychophysics Toolbox 3 (Brainard, 1997) (RRID:SCR_002881) in MATLAB 2010b.

The spoken word recognition paradigm consisted of six blocks of spoken words per run. Each block contained 15 seconds of spoken words (one word per second), followed by 15 seconds of silence. Two runs were performed in each study session with a total of 180 words in about 6 minutes (Figure 1A, middle panel). This task was first introduced by Petersen et al. (1988) and subsequently replicated in other studies with the same HD-DOT system used in this study (Eggebrecht et al., 2014; Fishell et al., 2019; Sherafati et al., 2020). All sound files were generated from a list of short simple nouns (1 to 4 syllables) pronounced with a female digital-voice audio using AT&T Natural Voices® text-to-Speech generator with the 'Lauren' digital voice.

To better understand the left PFC activity we observed in our first few CI users, we adopted
a spatial working memory task introduced in previous studies (Fedorenko et al., 2011; Fedorenko
et al., 2013) in the remaining subjects to aid in functionally localizing domain-general regions of
prefrontal cortex. In this spatial working memory task, subjects were asked to remember four
locations (easy condition) or eight locations (hard condition) in a 3 × 4 grid, appearing one at a
time. Following each trial, subjects had to choose the pattern they saw among two choices, one
with correct and one with incorrect locations. This task requires keeping sequences of elements
in memory for a brief period and has been shown to activate PFC. Each run for the spatial working
memory task was about 8 minutes, with a total of 48 trials in the run.

Data processing
HD-DOT data were pre-processed using the NeuroDOT toolbox (Eggebrecht & Culver, 2019).
Source-detector (SD) pair light level measurements were converted to log-ratio by calculating the
temporal mean of a given SD-pair measurement as the baseline for that measurement. Noisy
measurements were empirically defined as those that had greater than 7.5% temporal standard
deviation in the least noisy (lowest mean motion) 60 seconds of each run (Eggebrecht et al., 2014;
Sherafati et al., 2020). The data were next high pass filtered at 0.02 Hz. The global superficial
signal was estimated as the average across the first nearest neighbor measurements (13 mm
SD-pair separation) and regressed from all measurement channels (Gregg et al., 2010). The
optical density time-courses were then low pass filtered with a cutoff frequency of 0.5 Hz to the
physiological brain signal band and temporally downsampled from 10 Hz to 1 Hz. A wavelength-
dependent forward model of light propagation was computed using an anatomical atlas including
the non-uniform tissue structures: scalp, skull, CSF, gray matter, and white matter (Ferradal et
al., 2014) (Figure 1C). The sensitivity matrix was inverted to calculate relative changes in
absorption at the two wavelengths via reconstruction using Tikhonov regularization and spatially
variant regularization (Eggebrecht et al., 2014). Relative changes in the concentrations of
oxygenated, deoxygenated, and total hemoglobin (ΔHbO, HbR, ΔHbT) were then obtained from
the absorption coefficient changes by the spectral decomposition of the extinction coefficients of
oxygenated and deoxygenated hemoglobin at the two wavelengths (750 nm and 850 nm). After
post-processing, we resampled all data to a $3 × 3 × 3 \text{ mm}^3$ standard atlas using a linear affine
transformation for group analysis. In addition to the standard HD-DOT pre-processing steps used
in the NeuroDOT toolbox, we used a comprehensive data quality assessment pipeline (see
Appendix 1) to exclude the data runs with low pulse SNR or high motion levels.
After pre-processing, the response for the speech task was estimated using a standard
general linear model (GLM) framework. The design matrix was constructed using onsets and
durations of each stimulus presentation convolved with a canonical hemodynamic response
function (HRF). This HRF was created using a two-gamma function (2 s delay time, 7 s time to
peak, and 17 s undershoot) fitted to the HD-DOT data described in a previous study (Hassanpour
et al., 2015). We included both runs for each subject in one design matrix using custom MATLAB
scripts (Appendix1—figure 2).
For modeling the spatial working memory task, we used a design matrix with two columns
representing easy and hard conditions. The duration of each easy or hard trial was modeled as
the total time of stimulus presentation and evaluation. Events were convolved with the same
canonical HRF described in the spoken word perception task to model hemodynamic responses
(Hassanpour et al., 2014). We used the easy + hard response maps as a reference for defining
the PFC ROI, which was more robust than the hard > easy contrast previously used for younger
populations.
After estimating the response (β map) for each subject for each task, we performed a
second-level analysis in SPM12 (Wellcome Trust Centre for Neuroimaging) version 7487
(RRID:SCR_007037). Extracted time traces for each subject were then calculated using a finite
impulse response model.

We only present the ΔHbO results in the main figures as we have found that the ΔHbO
signal exhibits a higher contrast-to-noise ratio compared to ΔHbR or ΔHbT (Eggebrecht et al.,
2014; Hassanpour et al., 2014).
192

Functionally-defined regions of interest
To perform a more focused comparison between controls and CI users, we defined three regions
of interest (ROIs), independent from our spoken word recognition dataset, for statistical analysis.
To accurately localize the elevated PFC activation in the CI group, we collected HD-DOT
data from 9 subjects (4 controls and 5 CI users in 13 sessions) using a spatial working memory
task (Fedorenko et al., 2012). The visual spatial working memory task robustly activates PFC due
to its working memory demands (and visual cortex because of its visual aspect). We chose this
task to localize the PFC ROI for performing an ROI-based statistical analysis between controls
and CI users. Our logic is that in prior work this task shows activity that dissociates from nearby
language-related activity (Fedorenko et al., 2012), and thus the region of PFC localized by this
task is more likely to reflect domain-general processing (as opposed to language-specific
processing). Our results show strong bilateral visual and PFC activations in response to this task
(Figure 2A left). We then defined the left PFC ROI as the cluster of activation in the left PFC
(Figure 2A right). This region was centered in inferior frontal sulcus, extending into both dorsal
IFG and inferior MFG. The location is broadly consistent with domain-general ("multiple demand")
activation (e.g., meta-analysis in Duncan, 2010).
To define the left and right auditory ROIs, we used a previously published fMRI resting
state dataset (Sherafati et al., 2020) that was masked by the field of view of our HD-DOT system.
We defined the left and right auditory ROIs by selecting a 5 mm radius seed in the contralateral
hemisphere and finding the Pearson correlation between the time-series of the seed region with
all other voxels in the field of view. Correlation maps in individuals were Fisher z-transformed and
averaged across subjects (Figure 2B-C, left). Right/left auditory ROIs were defined by masking
the correlation map to include only the right/left hemisphere (Figure 2B-C right). These ROIs
extend well beyond primary auditory cortex, which is important to capture responses to words
often observed in lateral regions of STG and MTG.
218

Figure 2 | Defining functional ROIs. (A) Spatial working memory activation for 5 CI users and 4 controls over 13 sessions. The PFC ROI was defined as the cluster of activation in the PFC region, after p < 0.05 (uncorrected) voxelwise thresholding. **(B)** Seed-based correlation map for a seed located in the left auditory cortex (left map). Right auditory ROI defined by masking the correlation map to include only the right hemisphere (right map). **(C)** Seed-based correlation map for a seed located in the right auditory cortex (left map). Left auditory ROI defined by masking the correlation map to include only the left hemisphere (right map).

Speech perception score
To measure auditory-only speech perception accuracy, we presented each participant with two
lists of AzBio sentences in quiet (Spahr et al., 2012) (except one participant who only heard one
set). We calculated speech perception accuracy as the proportion of correctly repeated words
across all sentences.

Hearing threshold
We summarized hearing thresholds using a four-frequency pure-tone average (4fPTA), averaging
thresholds at 500, 1000, 2000, and 4000 Hz. For unaided testing, a value of 120 dB was assigned
if there was no response at a given frequency; for aided or CI testing, a value of 75 dB was used.
For CI users, hearing was tested with the right ear CI alone, left ear alone unaided, and left ear
with a hearing aid, if worn at the time of testing. Figure Supplement 5 shows the audiograms for
both controls and CI users.

Quantification and statistical analysis

Based on prior optical neuroimaging studies with similar speech related tasks (Defenderfer et al., 2021; Eggebrecht et al., 2014; Hassanpour et al., 2015; Pollonini et al., 2014; Zhou et al., 2018), we anticipated 15-20 subjects per group would be sufficient to detect a moderate effect size while also within our resource limitations. We therefore aimed to have at least 15 subjects in each group. We performed a one-tailed two-sample t-test for our main hypothesis (increased recruitment of PFC in CI users), for which we had a strong directional hypothesis (greater activity in listeners with CIs than controls), and a two-tailed t-test for the left and right auditory cortex changes for which we did not have directional hypotheses. We adjusted for unequal variances for left PFC and right auditory ROIs based on the significance of Levene's test.

Results

Multi-session single subject results

Due to the variability across CI users and difficulties in defining single-subject ROIs, we performed a small multi-session study from one CI subject for 6 sessions (Figure 3). We collected 2 runs of spoken word perception per session (for 6 sessions) and 1 run of spatial working memory task per session (for 4 sessions). This multi-session analysis enabled localizing the left and right PFC based on the non-verbal spatial working memory task for this subject (Figure 3C). It also revealed the engagement of regions beyond the auditory cortex, including PFC, during the word perception task (Figure 3B). Time-traces of oxyhemoglobin concentration change show a clear event-related response for four selected regions in the word perception results.

Figure 3 | Single subject data from one CI user across multiple sessions. (A) A CI user wearing the HD-DOT cap. **(B)** Response to the spoken words across 6 sessions (72 trials, each 15 sec stimuli, 15 sec rest). Hemodynamic response time-traces are plotted for peak activation values across 6 sessions for 4 key brain regions. The seed colors match the plot boundaries with error bars indicating the standard error of the mean. **(C)** Response to the spatial working memory task for the same CI user across 4 sessions (40 min of data).

Mapping the brain response to spoken words

We first investigated the degree of auditory activation in both control and CI groups by assessing the activity in a block-design single word presentation task. We found strong bilateral superior temporal gyrus (STG) activations in controls similar to our previous studies using the same paradigm (Eggebrecht et al., 2014; Sherafati et al., 2020), as well as a strong left STG and a reduced right STG activation for the CI users (Figure 4A-B). In addition, we observed strong left-lateralized activations in regions beyond the auditory cortex, including parts of the prefrontal cortex, in the CI user group (Figure 4B).

Figure 4—figure supplement 1 | Spoken word recognition group maps. Response to the spoken words in **(A)** 18 controls and **(B)** 20 right-ear CI users. **(C)** Differential activation in response to the spoken words task in CIs > controls highlights the group differences. The first column shows unthresholded β maps and the second column shows t-statistic maps thresholded at voxelwise p < 0.05 (uncorrected) for each group.

Figure Supplement 1 provides the β maps of oxyhemoglobin (HbO), deoxyhemoglobin (HbR), and total hemoglobin (HbT) for controls (panel A), CI users (panel B), CI users greater than controls (panel C), and controls greater than CI users (panel D).

For our statistical analysis, we focused on our predefined regions of interest, shown in Figure 2. Figure 5A–C show unthresholded t maps, masked by our ROIs. We averaged beta values within each ROI, and statistically tested for group differences correcting for multiple comparisons across the three ROIs (Figure 5D). The temporal profile of the hemodynamic response in three selected ROIs also suggests the increased activity in the left PFC in the CI users relative to controls, and a decrease in both left and right auditory regions. Two sample t-statistics for the mean β values in each ROI support a statistically significant difference between the control and CI groups in left PFC, t(27) = -2.3, p = 0.015 (one-tailed) and right auditory cortex,

278 t(23) = 3.54, p = 0.0017 (two-tailed) (Figure 5D). The observed change in the left auditory cortex
279 was not statistically significant, t(36) = 1.46, p = 0.15 (two-tailed). The threshold for statistical
280 significance, Bonferroni corrected for multiple comparisons across three ROI analyses, was
281 0.016.

Figure 5—figure supplements 2, 3, 4 | ROI-based statistical analysis for spoken word recognition task. Unthresholded t-maps in response to the spoken words spatially masked in the three ROIs defined in **Figure 3** in **(A)** 18 controls, **(B)** 20 right-ear CI users, and **(C)** CIs > Controls, highlight the group differences in certain brain areas. **(D)** Temporal profile of the hemodynamic response in three selected ROIs (left PFC, left auditory, and right auditory cortices). Two-sample t-tests for mean β value in each ROI have been calculated between controls and the CI-user group, confirming a significant increase in left PFC (p = 0.015) and significant decrease in the right auditory cortex (p = 0.0017) in CI users (indicated by an asterisk above their corresponding box plots, corrected for multiple comparisons). The observed change in the left auditory cortex was not significant (p = 0.15).

282
To further investigate whether the significant decrease in right auditory cortex might be
due to lower light level values around the CI transducer, we first performed a simulation of the
HD-DOT sensitivity profile blocking the optodes around the CI transducer and found minimal
overlap between the CI-related signal loss and the right auditory ROI (Appendix 2). We also
performed a further simulation analysis by first calculating the mean value of optical light power
for each measurement. Then, for each source-detector location, we defined the approximate
coupling coefficient as the mean over the first nearest neighbor measurements that have that
source or detector in common. Figure Supplements 2 and 3 show the coupling coefficients of
each source and detector for all CI users and controls in the flat view, corrected by the average
sensitivity of the avalanche photo-diode detectors (APDs) and plotted on a log base 10 scale to
create linearly-distributed color scale. We defined noisy optodes in the CI cohort as ones that had
a coupling coefficient of 30% of maximum or lower based on the values for each participant across
their two speech runs. If no optode was identified using this threshold, we lowered the threshold
by 10% increments until at least one source or detector was identified (under the logic that the CI
transducer must block some light). Table Supplement 1 shows the source-detector numbers and

the threshold used for identifying them for each CI user. For controls, no source or detector had
a coupling coefficient of less than 30% of the maximum.
After identifying the noisy source-detector numbers for all 20 CI users, we preprocessed
the data 100 times. For each of these analyses, we blocked the identified sources and detectors
in a selected CI user for one control, assigned by a random permutation, effectively simulating
the CI-based dropout in our control participants. Then, we replicated the ROI analysis for the right
auditory cortex in Figure 5, far right panel, and found that in all 100 shuffles, the right auditory
cortex in control subjects still had a significantly larger activation compared to the CI users (Figure
Supplement 4).

Behavioral measures
An important consideration in studying CI users is the variability in their speech perception
abilities, hearing thresholds, and the relationship with brain activity. Figure 6 shows exploratory
analyses between the magnitude of the activation in the left PFC ROI for the CI cohort with respect
to the speech perception score, left ear hearing threshold un-aided, left ear hearing threshold at
test (aided if the subject used a hearing aid), and right ear hearing threshold (CI-aided). These
scores were unavailable for one control participant, who was thus not included in these analyses.
Using $p < 0.05$ (uncorrected) as a statistical threshold, left PFC activation positively
correlated with left ear unaided thresholds ($p = 0.01$, Pearson $r = 0.55$) and negatively correlated
with right ear CI-aided thresholds ($p = 0.02$, Pearson $r = -0.49$). Left PFC activation did not
correlate with speech perception score ($p = 0.4$, Pearson $r = 0.17$) and aided hearing threshold
for the left ear ($p = 0.1$, Pearson $r = 0.36$).
319

Figure 6—figure supplement 5 | Relationship between the magnitude of activation in left PFC and behavioral scores in CI users. Plots of the Pearson correlation r between the magnitude of the mean β value in the left PFC ROI are shown with respect to speech perception score, left ear hearing threshold unaided, left ear hearing threshold (aided if the subject used a hearing aid), and right ear CI-aided hearing threshold. Hearing threshold was defined as 4fPTA, pure tone average at 4 frequencies, 500, 1000, 2000, 4000 Hz.

Discussion

321 Using high-density optical brain imaging, we examined the brain networks supporting spoken
322 word recognition in listeners with CIs relative to a matched group of controls with bilateral normal
323 hearing. We found that relative to controls, when listening to words in quiet, listeners with CIs
324 showed reduced activity in the right auditory cortex and—critically—increased activity in left PFC.
325 We review these two findings in turn below.

Increased prefrontal cortex activity in CI users

When listening to spoken words in quiet, listeners with normal hearing typically engage the left and right superior temporal cortex, including primary and secondary auditory regions (Binder et al., 2000; Price et al., 1992; Rogers et al., 2020; Wiggins et al., 2016). Our current results for controls show this same pattern. However, when listeners with CIs performed the same task, we found that they also engaged left PFC significantly more than the controls.

Although we only tested a single level of speech difficulty (that is, speech in quiet), prior studies have parametrically varied speech intelligibility and found intelligibility-dependent responses in the prefrontal cortex. Use of several types of signal degradation (Davis & Johnsrude, 2003), revealed a classic "inverted-U" shape response in the prefrontal cortex as a function of speech intelligibility, with activity increasing until the speech became very challenging and then tapering off. A similar pattern was reported in fNIRS (Lawrence et al., 2018).

A pervasive challenge for understanding the role of PFC in speech understanding is the close anatomical relationship of core language processing regions and domain-general regions of prefrontal cortex (Fedorenko et al. 2012). We attempted to add some degree of functional specificity to our interpretation by including a spatial working memory task presumed to strongly engage domain-general regions with minimal reliance on language processing (Duncan, 2010; Woolgar et al., 2015). Ideally, we would have used functional ROIs individually created for each subject. However, we were not convinced that our data for this task were sufficiently reliable at the single-subject level, as we only had two runs. Furthermore, we did not have spatial working memory task data for all subjects. Thus, our functional localization relies on group-average spatial working memory responses. The region we identified—centered in left inferior frontal sulcus— corresponds well to other investigations of non-language tasks (e.g., Duncan, 2010), and supports our preferred interpretation of engagement of domain-general regions of dorsolateral PFC during listening. However, the ROI also extends into the dorsal portion of IFG, and we cannot rule out the possibility that this frontal activity relates to increased language (as opposed to domain-general) processing.

Reduced auditory cortical activity in CI users

We found reduced activity in the right auditory cortex in CI users relative to controls, which we attribute to differences in auditory stimulation. We limited our sample to CI listeners with unilateral right-sided implants but did not restrict left ear hearing. Most of our subjects with CIs had poor hearing in their left ears, which would result in reduced auditory information being passed to the contralateral (right) auditory cortex. This was as opposed to controls who had bilateral hearing. Prior fNIRS studies have also shown that activity in the superior temporal cortex corresponds with stimulation and comprehension (Olds et al., 2016; Zhou et al., 2018). We also leave open the possibility that the CI hardware in the right hemisphere interfered with the signal strength, although our simulation studies suggest this cannot fully explain the group differences (Appendix 2 and Figure Supplements 3, 4, and 5).

What is potentially more interesting is the hint at a lower level of activity in the left auditory cortex of the CI users compared to controls, even though all CI listeners were receiving adequate stimulation of their right auditory nerve with a right CI. There are several possible explanations for this finding. First, activity in superior temporal cortex does not reflect only "basic" auditory stimulation, but processing related to speech sounds, word meaning, and other levels of linguistic analysis. Thus, although subjects with CIs were certainly receiving stimulation and speech intelligibility scores were generally good, some variability was still present (mean speech perception score in quiet = 0.88, SD = 0.09). The overall level of speech processing was significantly ($p = 0.00005$) lower for CI users than controls (mean speech perception score = 0.99, SD = 0.01), resulting in decreased activity (indeed, because the depth of HD-DOT includes only about 1 cm of the brain, much of primary auditory cortex is not present in our field of view, and the observed group differences were localized in non-primary regions of STG and MTG).

Perhaps the most provocative explanation is that a reduction in top-down modulatory
processes (Davis & Johnsrude, 2007) plays out as reduced activity in the temporal cortex. That
is, given that effortful listening depends on attention (Wild et al., 2012), it might be that processes
related to top-down prediction (Cope et al., 2017; Sohoglu et al., 2012, 2014) are muted when too
much cognitive control is required for perceptual analysis. Reconciling this interpretation with
predictive coding accounts of speech perception (Blank & Davis, 2016; Sohoglu & Davis, 2020)
will require additional work. We emphasize that the group differences in left auditory regions were
not significant, and thus our interpretation is speculative.

Individual differences in prefrontal cortex activation during spoken word recognition
Because of the variability of outcomes in CI users (Firszt et al., 2004; Holden et al., 2013), one
promising thought is that individual differences in brain activation may help explain variability in
speech perception ability. Although our study was not powered for individual difference analysis
(Yarkoni & Braver, 2010), we conducted exploratory correlations to investigate this avenue of
inquiry. Interestingly, we saw a trend such that poorer hearing in the left (non-CI) ear was
correlated with increased activity in PFC. Our subjects with CIs had significant variability in left
ear hearing. Because the speech task was conducted using loudspeakers, we would expect both
ears to contribute to accurate perception. Thus, poorer hearing in the left ear would create a
greater acoustic challenge, with a correspondingly greater drain on cognitive resources. This
interpretation will need additional data to be properly tested.
395

Comparison with previous fNIRS studies in CI users
Due to limitations of using fMRI and EEG in studying CI users, fNIRS-based neuroimaging is an
attractive method for studying the neural correlates of speech perception in this population
(Hassanpour et al., 2015; Lawler et al., 2015; Saliba et al., 2016; Sevy et al., 2010; Zhou et al.,
2018). Some prior studies have looked at relationships of behavioral performance and brain
activity in CI users. For example, Olds and colleagues (2016) studied temporal lobe activity and
showed that both normal hearing and CI users with good speech perception, exhibited greater
cortical activations to natural speech than to unintelligible speech. In contrast, CI users with poor
speech perception had large, indistinguishable cortical activations to all stimuli. Zhou et al. (2018)
found that regions of temporal and frontal cortex had significantly different mean activation levels
in response to auditory speech in CI listeners compared with normal-hearing listeners, and these
activation levels were negatively correlated with CI users' auditory speech understanding.
Our current study differs in that we were able to simultaneously measure responses in
temporal and frontal lobes with a high-density array. Our finding of increased recruitment of left
PFC in CI listeners is broadly consistent with the recruitment of prefrontal cortex in fNIRS studies
in normal hearing participants listening to simulated distorted speech (Defenderfer et al., 2021;
Lawrence et al., 2018). Potential differences in anatomical localization could reflect the difference
between the type of the distortion created by an actual CI compared to simulated degraded
speech listening scenarios in the previous studies, or in source localization accuracy across
different imaging hardware. To our knowledge, our study is the first fNIRS-based study to use a
nonverbal spatial working memory task for localizing the PFC ROI in the same population, and it
may be that increased use of cross-domain functional localizers will prove to be a useful approach
in future work.

Caveats, considerations, and future directions
As with all fNIRS-based functional brain imaging, our ability to image neural activity is limited by
the placement of sources and detectors, and in depth to approximately 1 cm of the cortex. Our
HD-DOT cap covers the regions of superficial cortex commonly identified in fMRI studies of
speech processing, including bilateral superior and middle temporal gyri, and pars triangularis of

the left inferior frontal gyrus (Davis & Johnsrude, 2003; Rodd et al., 2005; Rogers et al., 2020;
Wild et al., 2012). We also have good coverage of occipital cortex and middle frontal gyrus.
However, in the present study we do not image pars opercularis of the inferior frontal gyrus, the
cerebellum, or any subcortical structures. Thus, it is certainly possible that additional regions not
identified here play different roles in understanding speech for CI listeners, a possibility that will
require fNIRS setups with improved cortical coverage and converging evidence from other
methods to explore.
We also note that some care should be taken in the degree to which we have identified
PFC. As we have pointed out, the functional organization of the frontal cortex is complex, and
differs from person to person; in particular, fMRI studies have demonstrated that regions
responding to language and non-language tasks lie nearby each other (Fedorenko et al., 2012).
Although we attempted to improve the interpretation of our results by using a functional localizer,
we were not able to define subject-specific ROIs, which would be preferable. Future work with
single-subject localizers and functional ROIs is needed to more clearly resolve this issue.
Cognitive demand during speech understanding frequently comes up in discussions of
effortful listening (Pichora-Fuller et al., 2016). Although understandable, it is important to keep in
mind that the construct of listening effort is not clearly defined (Strand et al., 2020) and different
measures of "effort" do not always agree with each other (Strand et al., 2018). Here, we interpret
increased activity in PFC as reflecting greater cognitive demand. Although we did not include an
independent measure of cognitive challenge, we note that simulated CI speech (i.e., noise-
vocoded speech) is associated with delayed word recognition (McMurray et al., 2017) and poorer
memory for what has been heard (Ward et al., 2016), consistent with increased dependence on
shared cognitive processes (Piquado et al., 2010). Relating activity in PFC to cognitive demand
is also consistent with *decreased* activity in PFC when speech becomes unintelligible (Davis &
Johnsrude, 2003).
Finally, we emphasize that in the current study we only measured responses to speech in
quiet. Everyday communication frequently occurs in the presence of background noise, which can
be particularly challenging for many listeners with cochlear implants (Firszt et al., 2004). Exploring
how activity in PFC and other regions fluctuates in response to speech at various levels of
background noise would be a highly interesting future extension of this work. Based on parametric
modulations of acoustic challenge in listeners with normal hearing, we might expect increasing
activity in PFC as speech gets more challenging (Davis & Johnsrude, 2003), followed by cingulo-
opercular activity once intelligibility is significantly hampered (Vaden et al., 2013).
457

Conclusions

459 Using high-density optical neuroimaging, we found increased activity in prefrontal cortex in
460 listeners with cochlear implants compared to listeners with normal hearing while listening to words
461 in quiet. Our findings are consistent with a greater reliance on domain-general cognitive
462 processing and provide a potential framework for the cognitive effort that many CI users need to
463 expend during speech perception.

Acknowledgements

465 AS would like to thank Abraham Z. Snyder, Andrew K. Fishell, Kalyan Tripathy, Karla M. Bergonzi,
466 Zachary E. Markow, Tracy M. Burns-Yocum, Mariel M. Schroeder, Monalisa Munsi, Emily Miller,
467 Timothy Holden, and Sarah McConkey for helpful discussions. We also want to thank our
468 participants for their time and interest in our study.

Competing interests

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

Anderson, C. A., Wiggins, I. M., Kitterick, P. T., & Hartley, D. E. H. (2017). Adaptive benefit of cross-modal plasticity following cochlear implantation in deaf adults. *Proceedings of the National Academy of Sciences of the United States of America*, *114*(38), 10256-10261. https://doi.org/10.1073/pnas.1704785114

Binder, J. R., Frost, J. A., Hammeke, T. A., Bellgowan, P. S. F., Springer, J. A., Kaufman, J. N., & Possing, E. T. (2000). Human temporal lobe activation by speech and nonspeech sounds. *Cerebral Cortex*, *10*(5), 512-528. https://doi.org/DOI 10.1093/cercor/10.5.512

Blank, H., & Davis, M. H. (2016). Prediction errors but not sharpened signals simulate multivoxel fMRI patterns during speech perception. *PLoS biology, 14*(11), e1002577.

Brainard, D. H. (1997). The psychophysics toolbox. *Spatial Vision*, *10*(4), 433-436. https://doi.org/Doi 10.1163/156856897x00357

Braver, T. S. (2012). The variable nature of cognitive control: a dual mechanisms framework. *Trends in Cognitive Sciences*, *16*(2), 106-113. https://doi.org/10.1016/j.tics.2011.12.010

Cope, T. E., Sohoglu, E., Sedley, W., Patterson, K., Jones, P. S., Wiggins, J., . . . Rowe, J. B. (2017). Evidence for causal top-down frontal contributions to predictive processes in speech perception. *Nature Communications*, *8*. https://doi.org/ARTN 2154 10.1038/s41467-017-01958-7

Cousins, K. A. Q., Dar, H., Wingfield, A., & Miller, P. (2014). Acoustic masking disrupts time-dependent mechanisms of memory encoding in word-list recall. *Memory & Cognition*, *42*(4), 622-638. https://doi.org/10.3758/s13421-013-0377-7

Davis, M. H., & Johnsrude, I. S. (2003). Hierarchical processing in spoken language comprehension. *Journal of Neuroscience*, *23*(8), 3423-3431. <Go to ISI>://WOS:000182475200037

Davis, M. H., & Johnsrude, I. S. (2007). Hearing speech sounds: Top-down influences on the interface between audition and speech perception. *Hearing Research*, *229*(1-2), 132-147. https://doi.org/10.1016/j.heares.2007.01.014

Defenderfer, J., Forbes, S., Wijeakumar, S., Hedrick, M., Plyler, P., & Buss, A. T. J. N. (2021). Frontotemporal activation differs between perception of simulated cochlear implant speech and speech in background noise: An image-based fNIRS study. *240*, 118385.

Duncan, J. (2010). The multiple-demand (MD) system of the primate brain: mental programs for intelligent behaviour. *Trends in Cognitive Sciences*, *14*(4), 172-179. https://doi.org/10.1016/j.tics.2010.01.004

Dwyer, N. Y., Firszt, J. B., Reeder, R. M. J. E., & hearing. (2014). Effects of unilateral input and mode of hearing in the better ear: self-reported performance using the speech, spatial and qualities of hearing scale. *35*(1).

Eckert, M. A., Menon, V., Walczak, A., Ahlstrom, J., Denslow, S., Horwitz, A., & Dubno, J. R. (2009). At the Heart of the Ventral Attention System: The Right Anterior Insula. *Human Brain Mapping*, *30*(8), 2530-2541. https://doi.org/10.1002/hbm.20688

Eggebrecht, A. T., & Culver, J. P. (2019). https://github.com/WUSTL-ORL/NeuroDOT_Beta

Eggebrecht, A. T., Ferradal, S. L., Robichaux-Viehoever, A., Hassanpour, M. S., Dehghani, H., Snyder, A. Z., . . . Culver, J. P. (2014). Mapping distributed brain function and networks with diffuse optical tomography. *Nature Photonics*, *8*(6), 448-454. https://doi.org/10.1038/nphoton.2014.107

Fedorenko, E., Behr, M. K., & Kanwisher, N. (2011). Functional specificity for high-level linguistic processing in the human brain. *Proceedings of the National Academy of Sciences of the United States of America*, *108*(39), 16428-16433. https://doi.org/10.1073/pnas.1112937108

Fedorenko, E., Duncan, J., & Kanwisher, N. (2012). Language-Selective and Domain-General Regions Lie Side by Side within Broca's Area. *Current Biology*, *22*(21), 2059-2062. https://doi.org/10.1016/j.cub.2012.09.011

Fedorenko, E., Duncan, J., & Kanwisher, N. (2013). Broad domain generality in focal regions of frontal and parietal cortex. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(41), 16616-16621. https://doi.org/10.1073/pnas.1315235110

Ferradal, S. L., Eggebrecht, A. T., Hassanpour, M., Snyder, A. Z., & Culver, J. P. (2014). Atlas-based head modeling and spatial normalization for high-density diffuse optical tomography: In vivo validation against fMRI. *Neuroimage*, *85*, 117-126. https://doi.org/10.1016/j.neuroimage.2013.03.069

Firszt, J. B., Holden, L. K., Skinner, M. W., Tobey, E. A., Peterson, A., Gaggl, W., . . . Wackym, P. A. (2004). Recognition of speech presented at soft to loud levels by adult cochlear implant recipients of three cochlear implant systems. *Ear and Hearing, 25*(4), 375-387. https://doi.org/10.1097/01.Aud.0000134552.22205.Ee

Fishell, A. K., Burns-Yocum, T. M., Bergonzi, K. M., Eggebrecht, A. T., & Culver, J. P. (2019). Mapping brain function during naturalistic viewing using high-density diffuse optical tomography. *Scientific Reports*, *9*. https://doi.org/ARTN 11115
10.1038/s41598-019-45555-8

Garcia, C., Goehring, T., Cosentino, S., Turner, R. E., Deeks, J. M., Brochier, T., . . . Carlyon, R. P. (2021). The Panoramic ECAP Method: Estimating Patient-Specific Patterns of Current Spread and Neural Health in Cochlear Implant Users. *Jaro-Journal of the Association for Research in Otolaryngology*. https://doi.org/10.1007/s10162-021-00795-2

Gregg, N. M., White, B. R., Zeff, B. W., Berger, A. J., & Culver, J. P. (2010). Brain specificity of diffuse optical imaging: improvements from superficial signal regression and tomography. *Frontiers in neuroenergetics*, *2*, 14.

Hassanpour, M. S., Eggebrecht, A. T., Culver, J. P., & Peelle, J. E. (2015). Mapping cortical responses to speech using high-density diffuse optical tomography. *Neuroimage*, *117*, 319-326. https://doi.org/10.1016/j.neuroimage.2015.05.058

Hassanpour, M. S., Eggebrecht, A. T., Peelle, J. E., & Culver, J. P. (2017). Mapping effective connectivity within cortical networks with diffuse optical tomography. *Neurophotonics*, *4*(4). https://doi.org/ARTN 041402
10.1117/1.NPh.4.4.041402

Hassanpour, M. S., White, B. R., Eggebrecht, A. T., Ferradal, S. L., Snyder, A. Z., & Culver, J. P. (2014). Statistical analysis of high density diffuse optical tomography. *Neuroimage*, *85*, 104-116. https://doi.org/10.1016/j.neuroimage.2013.05.105

Holden, L. K., Finley, C. C., Firszt, J. B., Holden, T. A., Brenner, C., Potts, L. G., . . . hearing. (2013). Factors affecting open-set word recognition in adults with cochlear implants. *34*(3), 342.

Koeritzer, M. A., Rogers, C. S., Van Engen, K. J., & Peelle, J. E. (2018). The Impact of Age, Background Noise, Semantic Ambiguity, and Hearing Loss on Recognition Memory for Spoken Sentences. *Journal of Speech Language and Hearing Research*, *61*(3), 740-751. https://doi.org/10.1044/2017_Jslhr-H-17-0077

Lawler, C. A., Wiggins, I. M., Dewey, R. S., & Hartley, D. E. (2015). The use of functional near-infrared spectroscopy for measuring cortical reorganisation in cochlear implant users: A possible predictor of variable speech outcomes? *Cochlear implants international*, *16*(sup1), S30-S32.

Lawrence, R. J., Wiggins, I. M., Anderson, C. A., Davies-Thompson, J., & Hartley, D. E. H. (2018). Cortical correlates of speech intelligibility measured using functional near-infrared spectroscopy (fNIRS). *Hearing Research*, *370*, 53-64. https://doi.org/10.1016/j.heares.2018.09.005

McMurray, B., Farris-Trimble, A., & Rigler, H. (2017). Waiting for lexical access: Cochlear implants or severely degraded input lead listeners to process speech less incrementally. *Cognition*, *169*, 147-164. https://doi.org/10.1016/j.cognition.2017.08.013

Noyce, A. L., Cestero, N., Michalka, S. W., Shinn-Cunningham, B. G., & Somers, D. C. (2017). Sensory-Biased and Multiple-Demand Processing in Human Lateral Frontal Cortex. *Journal of Neuroscience*, *37*(36), 8755-8766. https://doi.org/10.1523/Jneurosci.0660-17.2017

Olds, C., Pollonini, L., Abaya, H., Larky, J., Loy, M., Bortfeld, H., . . . Oghalai, J. S. (2016). Cortical Activation Patterns Correlate with Speech Understanding After Cochlear Implantation. *Ear and Hearing*, *37*(3), E160-E172. https://doi.org/10.1097/Aud.0000000000000258

Peelle, J. E. (2017). Optical neuroimaging of spoken language. *Language Cognition and Neuroscience*, *32*(7), 847-854. https://doi.org/10.1080/23273798.2017.1290810

Peelle, J. E. (2018). Listening Effort: How the Cognitive Consequences of Acoustic Challenge Are Reflected in Brain and Behavior. *Ear and Hearing*, *39*(2), 204-214. https://doi.org/10.1097/Aud.0000000000000494

Petersen, S. E., Fox, P. T., Posner, M. I., Mintun, M., & Raichle, M. E. (1988). Positron Emission Tomographic Studies of the Cortical Anatomy of Single-Word Processing. *Nature*, *331*(6157), 585-589. https://doi.org/DOI 10.1038/331585a0

Pichora-Fuller, M. K., Kramer, S. E., Eckert, M. A., Edwards, B., Hornsby, B. W. Y., Humes, L. E., . . . Wingfield, A. (2016). Hearing Impairment and Cognitive Energy: The Framework for Understanding Effortful Listening (FUEL). *Ear and Hearing*, *37*, 5s-27s. https://doi.org/10.1097/Aud.0000000000000312

Piquado, T., Cousins, K. A. Q., Wingfield, A., & Miller, P. (2010). Effects of degraded sensory input on memory for speech: Behavioral data and a test of biologically constrained computational models. *Brain Research*, *1365*, 48-65. https://doi.org/10.1016/j.brainres.2010.09.070

Pollonini, L., Olds, C., Abaya, H., Bortfeld, H., Beauchamp, M. S., & Oghalai, J. S. J. H. r. (2014). Auditory cortex activation to natural speech and simulated cochlear implant speech measured with functional near-infrared spectroscopy. *309*, 84-93.

Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2012). Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion (vol 59, pg 2142, 2012). *Neuroimage*, *63*(2), 999-999. https://doi.org/10.1016/j.neuroimage.2012.01.069

Price, C., Wise, R., Ramsay, S., Friston, K., Howard, D., Patterson, K., & Frackowiak, R. (1992). Regional Response Differences within the Human Auditory-Cortex When Listening to Words. *Neuroscience Letters*, *146*(2), 179-182. https://doi.org/Doi 10.1016/0304-3940(92)90072-F

Rabbitt, P. M. A. (1968). Channel-Capacity Intelligibility and Immediate Memory. *Quarterly Journal of Experimental Psychology*, *20*, 241-&. https://doi.org/Doi 10.1080/14640746808400158

Rodd, J. M., Davis, M. H., & Johnsrude, I. S. (2005). The neural mechanisms of speech comprehension: fMRI studies of semantic ambiguity. *Cerebral Cortex*, *15*(8), 1261-1269. https://doi.org/10.1093/cercor/bhi009

Rogers, C. S., Jones, M. S., McConkey, S., Spehar, B., Van Engen, K. J., Sommers, M. S., & Peelle, J. E. J. N. o. L. (2020). Age-related differences in auditory cortex activity during spoken word recognition. *1*(4), 452-473.

Saliba, J., Bortfeld, H., Levitin, D. J., & Oghalai, J. S. J. H. r. (2016). Functional near-infrared
spectroscopy for neuroimaging in cochlear implant recipients. *338*, 64-75.
Schroeder, M. L., Sherafati, A., Ulbrich, R. L., Fishell, A. K., Svoboda, A. M., Culver, J. P., &
Eggebrecht, A. T. (2020). Mapping Cortical Activations Underlying Naturalistic Language
Generation Without Motion Censoring Using HD-DOT. Optical Tomography and
Spectroscopy,
Sevy, A. B., Bortfeld, H., Huppert, T. J., Beauchamp, M. S., Tonini, R. E., & Oghalai, J. S. J. H.
r. (2010). Neuroimaging with near-infrared spectroscopy demonstrates speech-evoked
activity in the auditory cortex of deaf children following cochlear implantation. *270*(1-2),
39-47.
Sherafati, A. (2020). *Separating Signal from Noise in High-Density Diffuse Optical Tomography*
Washington University in St. Louis].
Sherafati, A., Eggebrecht, A. T., Bergonzi, K. M., Burns-Yocum, T. M., & Culver, J. P. (2018).
Improvements in functional diffuse optical tomography maps by global motion censoring
techniques. Optics and the brain,
Sherafati, A., Eggebrecht, A. T., Burns-Yocum, T. M., & Culver, J. P. (2017). A global metric to
detect motion artifacts in optical neuroimaging data (Conference Presentation). Neural
Imaging and Sensing,
Sherafati, A., Snyder, A. Z., Eggebrecht, A. T., Bergonzi, K. M., Burns-Yocum, T. M., Lugar, H.
M., . . . Culver, J. P. (2020). Global motion detection and censoring in high-density
diffuse optical tomography. *Human Brain Mapping*, *41*(14), 4093-4112.
https://doi.org/10.1002/hbm.25111
Smyser, C. D., Snyder, A. Z., & Neil, J. J. (2011). Functional connectivity MRI in infants:
Exploration of the functional organization of the developing brain. *Neuroimage*, *56*(3),
1437-1452. https://doi.org/10.1016/j.neuroimage.2011.02.073
Sohoglu, E., & Davis, M. H. (2020). Rapid computations of spectrotemporal prediction error
support perception of degraded speech. *Elife*, *9*. https://doi.org/ARTN e58077
10.7554/eLife.58077
Sohoglu, E., Peelle, J. E., Carlyon, R. P., & Davis, M. H. (2012). Predictive Top-Down
Integration of Prior Knowledge during Speech Perception. *Journal of Neuroscience*,
32(25), 8443-8453. https://doi.org/10.1523/Jneurosci.5069-11.2012
Sohoglu, E., Peelle, J. E., Carlyon, R. P., & Davis, M. H. (2014). Top-Down Influences of Written
Text on Perceived Clarity of Degraded Speech. *Journal of Experimental Psychology-*
Human Perception and Performance, *40*(1), 186-199. https://doi.org/10.1037/a0033206
Spahr, A. J., Dorman, M. F., Litvak, L. M., Van Wie, S., Gifford, R. H., Loizou, P. C., . . . hearing.
(2012). Development and validation of the AzBio sentence lists. *33*(1), 112.
Strand, J. F., Brown, V. A., Merchant, M. B., Brown, H. E., & Smith, J. (2018). Measuring
Listening Effort: Convergent Validity, Sensitivity, and Links With Cognitive and
Personality Measures. *Journal of Speech Language and Hearing Research*, *61*(6), 1463-
1486. https://doi.org/10.1044/2018_Jslhr-H-17-0257
Strand, J. F., Ray, L., Dillman-Hasso, N. H., Villanueva, J., Brown, V. A. J. A. p., & cognition.
(2020). Understanding speech amid the jingle and jangle: recommendations for
improving measurement practices in listening effort research. *3*(4), 169-188.
Vaden, K. I., Kuchinsky, S. E., Cute, S. L., Ahlstrom, J. B., Dubno, J. R., & Eckert, M. A. (2013).
The Cingulo-Opercular Network Provides Word-Recognition Benefit. *Journal of*
Neuroscience, *33*(48), 18979-18986. https://doi.org/10.1523/Jneurosci.1417-13.2013
Vaden, K. I., Teubner-Rhodes, S., Ahlstrom, J. B., Dubno, J. R., & Eckert, M. A. (2017).
Cingulo-opercular activity affects incidental memory encoding for speech in noise.
Neuroimage, *157*, 381-387. https://doi.org/10.1016/j.neuroimage.2017.06.028

Ward, C. M., Rogers, C. S., Van Engen, K. J., & Peelle, J. E. (2016). Effects of Age, Acoustic
Challenge, and Verbal Working Memory on Recall of Narrative Speech. *Experimental*
Aging Research, *42*(1), 126-144. https://doi.org/10.1080/0361073x.2016.1108785
White, B. R., & Culver, J. P. (2010). Quantitative evaluation of high-density diffuse optical
tomography: in vivo resolution and mapping performance. *Journal of Biomedical Optics*,
15(2). https://doi.org/Artn 026006
10.1117/1.3368999
Wiggins, I. M., Anderson, C. A., Kitterick, P. T., & Hartley, D. E. (2016). Speech-evoked
activation in adult temporal cortex measured using functional near-infrared spectroscopy
(fNIRS): Are the measurements reliable? *Hear Res*, *339*, 142-154.
https://doi.org/10.1016/j.heares.2016.07.007
Wild, C. J., Yusuf, A., Wilson, D. E., Peelle, J. E., Davis, M. H., & Johnsrude, I. S. (2012).
Effortful Listening: The Processing of Degraded Speech Depends Critically on Attention.
Journal of Neuroscience, *32*(40), 14010-14021. https://doi.org/10.1523/Jneurosci.1528-
12.2012
Wingfield, A. (2016). Evolution of Models of Working Memory and Cognitive Resources. *Ear and*
Hearing, *37*, 35s-43s. https://doi.org/10.1097/Aud.0000000000000310
Woolgar, A., Jackson, J., & Duncan, J. (2015). How domain general is information coding in the
brain? A meta-analysis of 93 multi-voxel pattern analysis studies. *Frontiers in Human*
Neuroscience, *9*.
Yarkoni, T., & Braver, T. S. (2010). Cognitive neuroscience approaches to individual differences
in working memory and executive control: conceptual and methodological issues. In
Handbook of individual differences in cognition (pp. 87-107). Springer.
Zeff, B. W., White, B. R., Dehghani, H., Schlaggar, B. L., & Culver, J. P. J. P. o. t. N. A. o. S.
(2007). Retinotopic mapping of adult human visual cortex with high-density diffuse
optical tomography. *104*(29), 12169-12174.
Zhou, X., Seghouane, A., Shah, A., Innes-Brown, H., Cross, W., Litovsky, R., & McKay, C. M.
(2018). Cortical Speech Processing in Postlingually Deaf Adult Cochlear Implant Users,
as Revealed by Functional Near-Infrared Spectroscopy. *Trends in Hearing*, *22*.
https://doi.org/10.1177/2331216518786850

718 **Table Supplement 1:** Noisy source-detectors in CI users
719
720

Subject ID	Noisy Source #	Noisy Detector #	Threshold (% max)[1]	
DOTCI01	69, 74	33, 65, 66, 70	30	722
DOTCI02	-	66	30	
DOTCI03	69, 70	70	50	723
DOTCI04	-	70	30	724
DOTCI05	12, 69, 77	70, 74	30	
DOTCI06	66	61, 62	30	725
DOTCI07	78	-	30	
DOTCI08	69	67, 70	30	726
DOTCI10	70	66	30	727
DOTCI11	8, 70, 74	71	30	
DOTCI12	79	77, 80	30	728
DOTCI13	-	69	30	729
DOTCI14	74	-	60	
DOTCI15	-	33	50	730
DOTCI16	65	68	50	
DOTCI17	73	70	40	731
DOTCI18	-	66, 73	50	732
DOTCI19	82	78, 79	70	
DOTCI20	-	66, 70	50	733
DOTCI21	65	61	80	
				734

Figure Supplement 1: Group results for different hemoglobin contrasts. β maps of Oxy (column 1, HbO), deoxy (column 2, HbR), and total (column 3, HbT) hemoglobin for the spoken word recognition task for **(A)** all controls, **(B)** all CI users, and **(C)** CIs > Controls. **(D)** Controls > CIs. Note that the first three rows in column 1 (HbO) are identical to the results shown in **Figure 4** column 1. The redundancy is to guide the reader for side-by-side comparisons with HbR and HbT results.

Figure Supplement 2: Coupling coefficients of each source and detector is shown for all CI users in a flat view. The dark red areas show the location of the signal drop for each CI user. The maximum coupling coefficient for each subject is shown above each plot.

Figure Supplement 3: Coupling coefficients of each source and detector is shown for all controls in a flat view. The maximum coupling coefficient for each subject is shown above each plot.

Figure Supplement 4: Effect of the simulated CI transducer in controls in the right auditory ROI analysis.
(A) Replicating the ROI-analysis for the right auditory cortex using 100 different shuffles of preprocessing for 18 controls by simulating the effect of the CI transducer. (B) Histogram of the p-values shows statistical significance (p<0.016 corrected) for all 100 shuffles across CI users and controls as found in the original preprocessing in Figure 5.

Figure Supplement 5: Audiograms for left and right ears. Audiograms for left and right ears. Individual subject (thin lines) and group mean (bold lines) hearing at 250, 500, 1000, 2000, 3000, 4000 and 6000 Hz for **(A)** left ear unaided, **(B)** left ear aided, and **(C)** right ear (unaided for controls and with a CI for CI users).

749 **Appendix 1**
750
751 **Data quality assessment steps:**
752

753 **Motion artifact detection**
754 Motion artifacts were detected in a time-point by time-point manner on the 10 Hz bandpass filtered
755 first nearest neighbor (13mm) measurements, using a previously described motion detection
756 method in HD-DOT, the global variance of the temporal derivatives (GVTD) (Sherafati et al., 2018;
757 Sherafati et al., 2017; Sherafati et al., 2020).
758
759 GVTD, similar to DVARS (derivative of variance) in fMRI (Power et al., 2012; Smyser et al., 2011),
760 is a vector g that is defined as the RMS of the temporal derivatives $(y_{ji} - y_{ji-1})$ across a set of
761 optical density measurements.
762

763
$$g = \begin{bmatrix} g_1 \\ \vdots \\ g_M \end{bmatrix}, \qquad g_i = \sqrt{\frac{1}{N} \sum_{j=1}^{N} (y_{ji} - y_{ji-1})^2}, \qquad \qquad (1)$$

764
765 N is the number of first nearest neighbor measurements, and M is the number of time-points.
766 GVTD indexes the global instantaneous change in the time-courses. Higher GVTD values indicate
767 high motion levels. We defined the motion criterion (g_{thresh}) (red line in Appendix 1—figure 1 and
768 5) based on the GVTD distribution mode ($\tilde{\kappa}$) plus a constant ($c = 10$) times the standard deviation
769 computed on the left (low) side of the mode (σ_L) .
770

771
$$g_{thresh} = \tilde{\kappa} + c\sigma_L,$$

772
773 Appendix 1—figure 1 shows two example GVTD time-traces for a low motion and a high motion
774 run. The GVTD threshold was calculated as $g_{thresh} = \tilde{\kappa} + 10\sigma_L$ and is shown as a red line.

Appendix 1—figure 1: Examples of GVTD time-traces for a low-motion and a high-motion spoken word recognition HD-DOT data. The red lines indicate the GVTD threshold of $\tilde{\kappa} + 10\sigma_L$ of each run.

Including motion regressors in the design matrix
After determining the time-points that passed the GVTD threshold, we then included a motion
regressor as a column for each time-point that passed the motion threshold with one for the noisy
time-point, and zeros for every other point in the design matrix, a method commonly known as
one hot encoding. An example design matrix including 2 runs in a session of spoken word
recognition task along with their motion regressors are shown in Appendix 1—figure 2.
781
782

783
784

Appendix 1—figure 2: Including motion regressors in the design matrix. (A) An example design matrix for the GLM model for the spoken word recognition task, including a constant column (cst), the task times for run1, one hot encoding columns for time-points passing the GVTD threshold for run 1, task times for run2, and one hot encoding columns for time-points passing the GVTD threshold for run 2. **(B)** The GVTD time-traces and the temporal masks for excluding the time-points passing the GVTD threshold are shown in **(B)** for run 1 and **(C)** for run 2.

Heartbeat (pulse) signal-to-noise-ratio (SNR) across HD-DOT field of view
One indicator of detecting physiological signal in optical neuroimaging is the detection of heartbeat frequency (pulse) (Sherafati, 2020). If the data has a very low heartbeat SNR it indicates that the lower frequencies emergent from the hemodynamic fluctuations might also have a low SNR. Therefore, we excluded the subjects that showed a very low mean pulse SNR across the field of view, as it indicates that their brain signal is also less reliable.

The underlying reason behind low pulse SNR is either a poor optode-scalp coupling or high levels of motion. If the optode-scalp coupling is good, normally pulse SNR is only low during high-motion (identified as high GVTD) epochs of data. However, if the optode-scalp coupling is poor, we will see low pulse SNR throughout the run. Note that there are not well-studied definitions and cutoffs for the calculation of pulse SNR in fNIRS based methods. Therefore, since we already regressed time-points contaminated with motion using the GVTD index, here we only exclude the subjects that had a very low mean pulse SNR across the cap compared to other subjects.

We calculated the pulse SNR using the NeuroDOT_Beta function PlotPhysiologyPower.m (https://github.com/WUSTL-ORL/NeuroDOT_Beta) following these steps: This function first calculates the Fast Fourier Transform (FFT) of the optical density signal (logmean ratio). Then, it calculates the pulse SNR based on the ratio of the band-limited pulse power ~1 Hz (P) and the noise floor (N).

The band-limited pulse power (P) is defined as the sum of the squares of the FFT magnitudes of a small frequency band around the peak of FFT magnitudes in the 0.5-2 Hz frequency bandwidth. The noise floor (N) is then defined as the median of the squares of the FFT

magnitudes of the FFT indices in the same 0.5-1 Hz frequency band, excluding the indices defined
as the peak frequency window (used for the heartbeat pulse power). Finally, the pulse SNR was
defined as $10 \times log_{10}(P/N)$.
For pulse SNR calculation, we only used the second nearest neighbor measurements
which penetrate through the cortex (30 mm source-detector separation), second wavelength (850
nm), and good measurements (ones with <7.5% std). Appendix 1—figure 3 shows an example
for high SNR values across the field of view (top) and its signature in a subset of the
measurements for the same run (bottom) (Appendix 1—figure 3A) and an example for low SNR
values across the field of view (top) and a subset of measurements for that run (bottom) (Appendix
1—figure 3B).

Appendix 1—figure 3: Examples of a good and a bad pulse SNR in HD-DOT data. Example pulse
SNR plot and a selection of the measurements from the HD-DOT array for **(A)** a high-quality pulse
SNR, and **(B)** a low-quality pulse SNR. Note the heartbeat frequency (~1 Hz) that appears as around
10 peaks in 10s.

Exclusion of subjects based on low pulse and high motion
818
819 We rank ordered all subjects based on their mean band limited SNR across the field of view and
820 their mean GVTD values (averaged over two spoken word recognition runs for each subject).

821 We then excluded one of the CI users due to a very low band limited SNR (Appendix 1—figure
822 4A) and one control due to a very high motion level (Appendix 1—figure 4B).

Appendix 1—figure 4: Exclusion of subjects with very low band limited SNR and very high motion levels. Sorting all subjects based on **(A)** mean band limited SNR value across the HD-DOT field of view, and **(B)** mean GVTD values across spoken word recognition runs. The red boxes indicate the subjects excluded based on each quality score.

Appendix 2

Simulating the effect of the CI transducer blocking the optodes

We identified the approximate area of the HD-DOT field of view that fell under the transducer across people. In most subjects, the transducer only blocked 3 to 5 optodes. Due to the high density of the optodes in HD-DOT systems (13mm nearest SD distance), even when an optode is blocked, the light still penetrates through that region from the neighboring sources. Therefore, in most cases, light level plots reveal sensitivity reduction in only 2 or 3 optodes. Appendix 2—figure 1A, shows the location of the CI transducer in one CI user and the optode numbers. Appendix 2—figure 1B shows the sensitivity of the HD-DOT cap if all the optodes are fully coupled with the scalp. Appendix 2—figures 1C and 1D demonstrate that the sensitivity drop due to the CI transducer is mainly affecting the inferior and middle temporal gyri when 4 and 6 source

837 detector pairs are simulated to be completely blocked. Note that the right auditory ROI (Figure
838 3B) mainly includes the superior temporal gyrus.
839

Appendix 2—figure 1: Effects of the CI transducer on HD-DOT sensitivity. (A) A right ear CI user wearing the HD-DOT cap. The yellow contours around the CI transducer with respect to the ear fiducials illustrate the source numbers (encircled in red) and detector numbers (encircled in blue) around the transducer. **(B)** The left panel shows the HD-DOT source-detector grid, overlaid on the MNI mesh used in the study for image reconstruction. The right panel shows the sensitivity of the HD-DOT cap around the MNI cortex including all optodes. **(C)** The left panel shows 4 sources (in red) and 4 detectors (in blue) excluded from the sensitivity calculation. The right panel shows the sensitivity of the HD-DOT cap excluding those sources and detectors. **(D)** The left panel shows 6 sources (in red) and 6 detectors (in blue) excluded from the sensitivity calculation. The right panel shows the sensitivity of the HD-DOT cap excluding those sources and detectors. Note that the exclusion of sources and detectors in (C) and (D) only resulted in sensitivity drop off in the inferior and middle temporal gyri.