1 **Title:** Synaptotagmin-7 enhances phasic dopamine release
2

3 **Authors:** Sarah A. Kissiwaa[1,*], Joseph J. Lebowitz[1,*], Kim A. Engeln[1], Anna M.
4 Bowman[1], John T. Williams[1], Skyler L. Jackman[1]
5
6 1Vollum Institute, Oregon Health & Science University, Portland, OR, United States
7 *These authors contributed equally

8
Abstract

Dopamine released from substantia nigra pars compacta (SNc) neurons modulates
movement, motivation, and reward. In addition to their tonic firing pattern, dopamine
neurons also fire high-frequency bursts that cause superlinear increases in dopamine
release. To study this poorly understood form of short-term plasticity, we used the
fluorescent dopamine sensor dLight1.3b to examine the role of the calcium-binding
protein synaptotagmin-7 (SYT7)**.** We report that SYT7 mediates a hidden component of
facilitation, which was unmasked by lowering initial release probability, or by low-
frequency stimulation of nerve terminals. In *Syt7* KO neurons, there was profound
synaptic depression that significantly reduced release during stimulations that mimic *in
vivo* firing patterns of SNc neurons. D2-mediated inhibitory postsynaptic currents in the
SNc revealed a similar role for SYT7 in somatodendritic release. Our results indicate
that SYT7 drives short-term facilitation of release from dopamine neurons, which likely
underlies frequency-dependence of dopamine signaling *in vivo.*

23
Introduction
Midbrain dopamine neurons modulate activity in disparate brain regions to shape affect,
attention, and action (Beeler & Dreyer, 2019). Dopamine neurons fire tonically at low
frequencies (~1-8 Hz), but also fire high-frequency (>20 Hz) bursts in response to
salient stimuli (Fiorillo et al., 2008; Fiorillo et al., 2003; Otomo et al., 2020). Phasic firing
releases dopamine more effectively than tonic firing. For example, appetitive rewards
elicit phasic firing in the midbrain (Schultz et al., 1997), and increase dopamine release
in forebrain terminal regions (Day et al., 2007). Stimulating the midbrain with high-
frequency bursts increases striatal dopamine concentrations more than the same
number of stimuli at low frequency (Gonon, 1988; Tsai et al., 2009). These observations
suggest that activity-dependent presynaptic plasticity increases dopamine release
during phasic firing (Cragg, 2003). Presynaptic plasticity is generally driven by
accumulation of calcium in axon terminals during high-frequency activity (Jackman &
Regehr, 2017). However, the presynaptic calcium sensor that confers frequency-
dependent enhancement of dopamine release has not been identified.
39
The vesicular Ca^{2+} sensor synaptotagmin-1 (SYT1) is required for synchronous
dopamine release in the striatum (Banerjee et al., 2020). However, when *Syt1* is deleted
from SNc neurons, a slow component of asynchronous striatal dopamine release
remains. Such asynchronous release has been theorized to be under control of
Synaptotagmin-7 (SYT7), a high-affinity Ca^{2+} sensor that mediates asynchronous
release and short-term synaptic facilitation in non-dopaminergic synapses (Chen et al.,
2017; Jackman et al., 2016; Luo & Südhof, 2017; Turecek & Regehr, 2018). *Syt7* RNA

is expressed throughout the brain, but is enriched in SNc neurons (Saunders et al.,
2018). SYT7 partially mediates release from dendrites within the SNc, and is present in
axon terminals of cultured dopamine neurons (Mendez et al., 2011), but whether SYT7
regulates release from dopamine axons remains unknown. Measurement of dopamine
release at sub-second intervals is required to test this hypothesis (Covey et al., 2013).
Here, we used high-speed photometry of a fluorescent dopamine sensor and whole-cell
electrophysiology in *Syt7* knockout mice to probe the role of SYT7 in dopamine release.
Release evoked by high-frequency stimulation was reduced in *Syt7* KO animals in
nerve terminals and in somatodendritic compartments. SYT7 enhanced dopamine
release during transitions from tonic to phasic firing. Our data establish an important
role for SYT7 in short-term plasticity and phasic dopamine release.
58
Results
We immunolabeled coronal brain sections with antibodies against SYT7 and tyrosine
hydroxylase (TH) (Figure 1A). In high-resolution images obtained using 3D structured
illumination microscopy, SYT7-positive puncta were abundant in the striatum, both
inside and outside the boundaries of TH-positive axons (Figure 1B). 3D reconstructions
of SYT7 immunoreactivity within TH positive regions revealed discrete puncta
throughout dopamine axons (Figure 1C-F), consistent with SYT7 localizing to the axon
terminals of dopamine neurons (Mendez et al., 2011). This localization is similar to non-
dopaminergic axons (Dean et al., 2012; Jackman et al., 2016; Turecek et al., 2017),
suggesting a conserved function for SYT7.
69

70
71
Figure 1. Immunohistochemistry suggests that SYT7 is present in dopaminergic axons.
(**A-B**) Coronal brain sections from a WT mouse labeled with antibodies against SYT7
(A) and tyrosine hydroxylase (TH) (B). (**C-F**) 0.5 µm-thick projections of a z-stack
acquired using 3D-structured illumination microscopy showing SYT7 (C) and TH
labeling (D), both channels overlaid (E), and a binary mask exclusively showing SYT7-
positive structures that lay within the boundaries of TH axons (F).
78

To probe the impact of SYT7 in striatal dopamine release, we monitored dopamine in
acute striatal slices from wildtype (WT) and *Syt7$^{-/-}$* (*Syt7* KO) mice (Chakrabarti et al.,
2003) using the genetically-encoded fluorescent dopamine sensor dLight1.3b (Patriarchi
et al., 2018). Adeno-associated viral vector was injected unilaterally into the dorsal
striatum to drive dLight expression in striatal neurons (Figure 2A-B). Sagittal brain slices
(270 µm) were subsequently prepared for imaging and electrophysiology. dLight
fluorescence was excited with dim light (mean intensity = 0.96 mW/mm^2) and recorded
at 10 kHz using a high-speed photodiode detector (Figure 2C). Dopamine axons were
activated by local extracellular stimulation, in artificial cerebrospinal fluid (ACSF)
containing 2 mM Ca^{2+}. To isolate action potential-driven dopamine release, recordings
were performed in the presence of antagonists for nicotinic, D2 and GABA receptors.
Electrical stimulation elicited rapid, low noise transients (Figure 2D). The use of dim
excitation light resulted in minimal photobleaching, permitting prolonged imaging over
the same region of the striatum (Figure 2E). Bath application of the D1 antagonist SKF-
83566 (10 µM) reduced the responses by 95 ± 3% (n = 4), confirming that fluorescence
was driven by binding of dopamine to dLight (Figure 2F). Lowering external Ca^{2+} to 0.5
mM reduced dLight signals by 81 ± 3% (n = 8) consistent with the expected calcium
dependence of dopamine release (Ford et al., 2010) (Figure 2G-H).
97

Figure 2. Optical monitoring of dopamine release in the striatum.
(**A**) Schematic illustrating method of expressing dLight1.3b by stereotaxic injection of
adeno-associated virus (AAV) into the dorsal striatum. (**B**) Representative image of a
sagittal brain slice showing dLight fluorescence. (**C**) Schematic illustrating optical
recording method. Fluorescence was captured using an amplified photodiode detector
mounted on an upright microscope. Dopamine axons were activated by local electrical
stimulation. (**D**) Representative fluorescence change evoked by electrical stimulation.
Average of 10 trials. (**E**) Averaged normalized response amplitudes during repeated
stimulation at 0.05 Hz (N = 7). (**F**) Average normalized responses during bath
application of the selective D1 receptor antagonist SKF-83566 (N = 4). (**G**)
Representative responses evoked in an acute brain slice in the presence of 2 and 0.5
mM external Ca^{2+} (Ca$_e$). (**H**) ΔF/F amplitudes in high and low Ca^{2+} ACSF (N = 8). Data
in all figures represents mean ± S.E.M.

112
dLight binds to dopamine with low affinity, and rapid on- and off-kinetics, providing an
approximate measure of the time-course of extracellular dopamine. Dopamine rises and
falls within ~100 ms of a release event (Condon et al., 2021), much faster than the
postsynaptic responses activated by G-protein coupled receptors. Because SYT7-
mediated asynchronous release can prolong the time-course of release at other
synapses (Bacaj et al., 2013; Turecek & Regehr, 2018), we evaluated whether the loss
of SYT7 affected the dopamine release kinetics as measured by dLight. Neither the
rise-time (20-80%), the time to the peak of the dLight transient, or the time for transients
to decay to 50% of the peak were altered in the absence of SYT7 (Figure 3A-B),
suggesting that SYT7 does not prolong the presence of dopamine in the extracellular
space by promoting asynchronous release.
124

125
Figure 3. Syt7 reduces short-term depression of striatal dopamine release.
(**A**) Representative dLight transients in striatal slices from WT and Syt7 KO animals. (**B**)
Averaged kinetics of dLight transients. (**C**) Representative responses to single stimuli
(light traces) and paired-pulse stimulation at 20 ms intervals (dark traces). (**D**) Averaged
paired-pulse ratios (A_2/A_1) at varying interstimulus intervals. Statistical significance in all
experiments was determined by unpaired Student's t-test after normal distribution and
homoscedasticity were confirmed. For comparisons of more than 2 samples, two-way
ANOVA tests were performed and Šidák corrected. Significance is shown as *: $p < 0.05$,
**: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$.
135
In addition to asynchronous release, SYT7 mediates short-term synaptic facilitation.
Such facilitation increases synchronous release for several hundred milliseconds after
an action potential, as assessed by the paired-pulse ratio (PPR) (Jackman & Regehr,
2017). Because dLight transients lasted for more than 100 ms, for short inter-stimulus

intervals (ISI) the amplitude of the second response was calculated by subtracting the
response driven by a single stimulus (Figure 3C). In WT animals the PPR was ~1 at
short intervals (0.87 ± 0.07 for ISI = 20 ms), followed by marked depression, and slow
recovery from depression (Figure 3D). In contrast, *Syt7* KOs displayed the hallmarks of
high release probability (*p*) synapses, depressing at short intervals (PPR = 0.34 ± 0.05
for ISI = 20 ms) and recovering slowly. This suggests that SYT7 serves to offset vesicle
depletion, resulting in sustained striatal dopamine release at short inter-stimulus
intervals.
148
Synapses that show short-term depression can still possess a hidden component of
facilitation that reduces depression at high stimulation frequencies (Turecek et al.,
2017). This hidden facilitation can be unmasked by decreasing external Ca^{2+} (Ca_e) to
reduce *p*. However, lowering Ca_e to 0.5 mM decreased dLight transients to the same
degree in WT and KO animals (Figure 4A-B), suggesting that the Ca^{2+}-dependence of
dopamine release is not affected by SYT7, as observed at other synapses (Jackman et
al., 2016; Liu et al., 2014). In WT synapses, dLight responses showed a ~2-fold
facilitation that decayed within 100 ms, whereas synapses in the *Syt7* KOs exhibited
mild depression (4C-D). Taken together, these experiments suggest that striatal
dopamine terminals have a high initial *p* and synaptic depression, but at high firing
frequencies SYT7 drives facilitation.
160

161
Figure 4. Syt7 mediates facilitation of dopamine release in low Ca_e.
(**A**) Representative dLight transients recorded in high and low Ca_e. (**B**) Average dLight
amplitudes in high and low Ca_e, normalized to amplitudes in 2 mM Ca_e. (**C**)
Representative responses evoked by single stimuli (light) and paired-pulse stimulation
at 20 ms intervals (dark) in 0.5 mM Ca_e. (**D**) Average paired-pulse ratios in 0.5 mM Ca_e
at varying interstimulus intervals in WT and *Syt7* KO.
168

Hidden facilitation can also be unmasked by depleting high-*p* vesicles with low
frequency stimulation (Doussau et al., 2017; Turecek et al., 2016). This possibility is
particularly relevant to dopamine release *in vivo*, as midbrain dopamine neurons switch
between low-frequency tonic firing, and high-frequency phasic bursts (Schultz et al.,
1997). Importantly, phasic firing is more effective at driving axonal dopamine release
(Tsai et al., 2009). To explore how SYT7 affects release during physiological patterns of
activity, we performed stimulations modeled after spike trains recorded *in vivo* from SNc
dopamine neurons in mice (Schiemann et al., 2012). The spike pattern contained
periods of tonic firing (instantaneous firing rates ranging from 1-6 Hz) as well as a tonic
burst (12-33 Hz). In 2 mM Ca$_e$, dopamine release was similar in WT and *Syt7* KO slices
during the first 3 tonic stimuli (Figure 5A). However, at the onset of a phasic burst,
dLight fluorescence increased rapidly in WT, often reaching the amplitude of the initial
pre-depressed response. In contrast, in KOs release after the phasic burst was
significantly decreased (Figure 5B-C). Because dopamine released during phasic bursts
may accumulate and act on longer time scales, we also calculated cumulative
dopamine release by integrating dLight fluorescence (Figure 5D). A significant
difference in cumulative fluorescence emerged between WT and *Syt7* KO synapses
after the onset of the phasic burst (Figure 5E-F). These data show that SYT7
contributes to the non-linear increase in striatal dopamine release during phasic firing of
SNc neurons.
189

190
191
192 **Figure 5**. Syt7 increases dopamine release during phasic firing.

(**A**) Stimulus train modeled after *in vivo* recordings from SNc dopamine neurons, and representative dLight responses from WT and *Syt7* KO animals. (**B**) Averaged responses from WT (N = 9) and KO (N = 6), normalized to the initial response amplitude. (**C**) Average normalized peak amplitudes for each stimulus, relative to the onset of the phasic burst. (**D**) Integrated dLight fluorescence (ΣF) calculated from the representative traces in (A), normalized to the end of the first response. (**E**) Averaged integrated fluorescence for WT and KO animals. (**F**) Difference in cumulative fluorescence between genotypes.

The data thus far support a role for SYT7 in driving short-term facilitation of dopamine release from striatal axon terminals during transitions from tonic to phasic firing. These transitions are in part regulated by D2 receptors expressed on dopamine cell dendrites which are activated by somatodendritic release (Lin et al., 2021). To determine the role of SYT7 in somatodendritic release, we performed whole-cell voltage-clamp recordings of D2-IPSCs. In *Syt7* KOs there was no change in the amplitudes of D2-IPSCs induced by single electrical stimuli (Figure 6A, WT: 28 pA ± 6 pA, KO: 25 ± 4 pA, p = 0.68), or a pair of stimuli at 40 Hz (Figure 6B). However, the amplitude of D2-IPSCs elicited by trains of 5 stimuli at 40Hz, normalized to the response to a single stimulus, was lower in KO animals (WT: 3.8 ± 0.4, Syt7 KO: 2.5 ± 0.2, p < 0.0001) (Figure 6C**)**. Paired-pulse ratios for stimuli presented at 1 Hz were similar in WT and KO animals **(**WT: 0.65 ± 0.02, KO: 0.64 ± 0.04, p = 0.90) (Figure 6F**)**, consistent with SYT7's minimal contribution to release at low frequencies in other synapses (Jackman et al., 2016). To examine whether SYT7-mediated facilitation could be revealed by pre-depressing synapses with low-frequency stimulation, we stimulated the SNc with 3 pre-pulses at 1 Hz, followed by a phasic train of 5 pulses at 100 Hz. Facilitation of release was measured as the ratio of the final pre-pulse IPSC (3rd) and the IPSC induced by the 100 Hz phasic stimulus train (4th) (Beckstead et al., 2007). In WT animals, the phasic IPSC was 4.6-fold larger than the pre-depressed IPSC, a greater ratio than that seen without a series of pre-pulses. However, in *Syt7* KO slices the ratio of pre-pulse to phasic IPSC was significantly lower than in WT slices, and did not differ from that seen without pre-depression in KO slices (WT: 4.6 ± 0.2, KO: 2.6 ± 0.2, p < 0.0001) (Figure 6E). Thus, Syt7 also mediates facilitation of dopamine release at somatodendritic sites.

Figure 6. Syt7 mediates facilitation of somatodendritic dopamine release.
(**A**) Representative D2-IPSCs induced by a single electrical stimulus, or by two or five
stimuli at 40Hz. (**B**) Amplitude of D2-IPSCs elicited by single stimuli in WT (N = 13 cells
from 4 animals) and *Syt7* KO animals (N = 19 cells from 4 animals). (**C**) Amplitude of
D2-IPSC induced by two or five stimuli at 40 Hz, normalized to the response elicited
with one stimulus. Same N as in (A) (**D**) Unmasking of facilitation protocol: three stimuli
were presented at 1 Hz, followed by a 5-stimuli train at 100 Hz. Facilitation was
quantified as the ratio of the 5-stimuli IPSC to the 3rd IPSC at 1 Hz. (**E**) Quantification of
facilitation in WT (N = 14 cells, 3 animals) and *Syt7* KOs (N = 15 cells, 3 animals) (**F**)
PPR measured for the first two 1 Hz stimuli.

Discussion

Here, we report that SYT7 drives short-term facilitation of dopamine release. SYT7
significantly increased release during phasic bursts, despite the fact that both axonal
and dendritic synapses exhibit a high initial release probability and overt synaptic
depression. A mixture of facilitation and depression was previously inferred from *in vivo*
electrochemical recordings of striatal dopamine concentrations during midbrain
stimulation (Montague et al., 2004). Similar "hidden" facilitation has been observed at
other depressing synapses, where SYT7 maintains release during sustained firing
(Chen et al., 2017; Turecek et al., 2017). Alternatively, tonic pacemaker activity *in vivo*
could deplete high release probability vesicles, accentuating release during phasic
bursts when SYT7 is activated by buildup of presynaptic calcium. Our experiments
highlight the utility of dLight1.3b for investigating rapid dopaminergic signaling, which
could be used to assess how presynaptic short-term plasticity impacts dopamine
release *in vivo*.

A potential limitation of this study is the use of global *Syt7* knockout animals, which
might lead to compensatory changes that confound the interpretation of our results.
However, in previous studies global *Syt7* deletion abolished facilitation selectively,
without changing other neuronal or synaptic properties (Bacaj et al., 2013; Jackman et
al., 2016). Moreover, facilitation is restored in global knockouts in a cell-autonomous
manner by presynaptic rescue of *Syt7* (Jackman et al., 2016; Turecek et al., 2017). In
biochemical assays SYT7 remains bound to calcium for hundreds of milliseconds (Hui
et al., 2005), similar to the time-course of facilitation we observed for dopamine release.
Hence, the most parsimonious interpretation for our results is that SYT7 binds to
presynaptic calcium following an action potential, and potentiates dopamine release for
several hundred milliseconds.
266
Despite progress in understanding the molecular basis of dopamine release, exactly
how neuronal activity and Ca^{2+} regulate release is unknown, and an understanding of
somatodendritic release mechanisms remains particularly incomplete. Dopaminergic
transmission relies on some of the same release machinery proteins that support fast,
calcium-dependent neurotransmitter release from non-dopamine synapses. Dopamine
release from both terminals and dendrites is dependent on RIM1, an active zone
scaffolding protein involved in vesicle priming, and Ca^{2+} channel tethering (Liu et al.,
2018; Robinson et al., 2019). However, dendrites and terminals may employ different
Ca^{2+} sensors for release, and exhibit different Ca^{2+}-dependence. The vesicular Ca^{2+}
sensor SYT1 is required for release from terminals (Banerjee et al., 2020), but not for
somatodendritic release (Delignat-Lavaud et al., 2021). Activity-evoked Ca^{2+} influx may
not be required for release from dendrites (Chen & Rice, 2001), although this is debated
(Ford et al., 2010; Yee et al., 2019). Further work is needed to clarify whether dendrites
and terminals employ distinct release mechanisms. The conserved role of SYT7 in
facilitating release from both compartments indicates that there are marked similarities
in the molecules that underlie dendritic and terminal dopamine release.
283
284
Methods
Animals
All mice were handled in accordance with NIH guidelines and protocols approved by
Oregon Health & Science University's Institutional Animal Care and Use Committee.
Syt7 knockout mice (Chakrabarti et al., 2003) and wild-type littermates of either sex
were used. Statistical tests were not used to predetermine sample size, but sample
sizes were similar to those in previous studies (Ford et al., 2010; Patriarchi et al., 2018).
All electrophysiological recordings and analysis were done blind to the genotype of the
animal, but for other experiments blinding and randomization were not performed.
294
Immunohistochemistry
Mice were anaesthetized with ketamine/xylazine (100/10 mg/kg) and transcardially
perfused with 4% paraformaldehyde (PFA) in PBS. Brains were removed and post-fixed
overnight at 4 °C. 50 µm slices were prepared using a Leica VT1000S vibratome, and
permeabilized for 1 h in vehicle (PBS, 10% normal goat serum, 0.3% Triton X-100),
then incubated overnight at 4 °C in vehicle with primary antibodies (mouse anti-Syt7,

1:200, Neuromab N275/14; rabbit anti-TH, 1:500, Millipore AB152). After 3 washes for 30 minutes in vehicle, slices were incubated with secondary antibodies in vehicle overnight at 4 °C (Alexa flor 488 goat anti mouse, Invitrogen A11001, 1:1000; Alexa flor 546 goat anti rabbit, Invitrogen A11035, 1:500). After 3 washes for 30 minutes in PBS, slices were mounted in Prolong Gold. Images were acquired using Zeiss LSM 980 or Zeiss Elyra 7 (airyscan).

Stereotaxic injections
Adult mice (P30-P90) were anaesthetized with ketamine/xylazine (100/10 mg/kg), placed in a stereotaxic apparatus (Kopf), and supplemented with 2% isoflurane. The head was shaved and disinfected with betadine and alcohol, an incision was made to expose the skull, and a small hole was drilled. 500 nl of AAV9.syn.dLight1.3b (4.66 10^{13} gc/ml) (Patriarchi et al., 2018) was injected unilaterally using glass pipettes (Drummond Scientific) at a rate of 100nl/minute (Nanoject 3000, Drummond). Injection coordinates from bregma were +1.4mm anterior, 1.4-1.5mm lateral and 3.5mm ventral. 10 minutes after injection the pipette was slowly retracted and scalp incision was closed with gluture. Post-injected analgesic (Carprofen, 5mg/kg) was administered subcutaneously for 48 hours.

dLight imaging
Acute brain slices were prepared from P60-120 mice 3-6 weeks after virus injection. Animals were anesthetized with isoflurane and euthanized. Brains were quickly removed into ice-cold cutting solution containing (in mM): 125 Choline-Cl, 25 NaHCO3, 10 glucose, 2.5 KCl, 1.25 NaH_2PO_4, 2 Na-pyruvate, 3 (3)-myo-inositol, 4.4 ascorbic acid, 7 MgCl2, 0.5 $CaCl_2$, bubbled continuously with 95% O_2/5% CO_2. 270 µm-thick sagittal slices were prepared using a Leica VT1200S vibratome, and transferred to a holding chamber with ACSF containing (in mM): 125 NaCl, 25 $NaHCO_3$, 10 glucose, 2.5 KCl, 1.25 NaH_2PO_4, 2 Na-pyruvate, 3 (3)-myo-inositol, 4.4 ascorbic acid, 1 $MgCl_2$, 2 $CaCl_2$, bubbled continuously with 95% O_2/5% CO_2. Slices were stored at room temperature (24±1 °C) prior to recording.

     Slices were transferred to a recording chamber and superfused with bubbled ACSF (~2 ml/min, 34±1 °C) containing 200 µM hexamethonium, 2 µM CGP, and 0.2 µM sulpiride. A bipolar stimulating electrode was placed in the dorsal striatum, and axons were excited using a constant-voltage stimulus isolator (Digitimer DS2A). dLight fluorescence was acquired through a 60X objective on an Olympus BX51WI microscope, equipped a 488 nm LED (Thorlabs) and a FITC XF100-2 filter set (Omega). Fluorescence was captured at 10 kHz using a custom-built amplified photodiode detector. Signals were digitized with an ITC-16 (Instrutech), and acquired in Igor Pro 8 (Wavemetrics) using custom-written routines (mafPC, courtesy of Matthew Xu-Friedman).

SNc whole-cell recordings
Acute SNc slices were prepared from P30-70 mice as previously described (Robinson et al., 2019, 2017). Animals were anesthetized with isoflurane before rapid decapitation into warmed (~35 °C) Krebs buffer containing (in mM): 126 NaCl, 1.2 $MgCl_2$, 2.4 $CaCl_2$, 1.4 NaH2PO4, 25 $NaHCO_3$, and 11 Dextrose. The brain was extracted and 222µm-thick

slices were cut in the horizontal plane at room temperature with continuous bubbling of 95% O_2/5% CO_2. Slices were allowed to recover for 30 min at 33 °C in bubbling Krebs. Typical preparations yielded 3 slices (ventral, medial, dorsal) which were hemisected for recording experiments. Brain extraction and slicing were all performed in the presence of 10 µM MK-801 to protect from NMDA-mediated excitotoxity. Recordings were obtained from hemisected slices continuously perfused with warm (~37 °C) Krebs containing 10 µM CNQX, 100 µM picrotoxin, and 300 nM CGP55845 to block AMPA-R, GABA-A, and GABA-B conductances respectively. Internal solution used for recording contained (in mM): 100 K-methanesulphonate, 20 NaCl, 1.5 MgCl2, 10 HEPES (K), 10 BAPTA, 2 ATP, 0.2 GTP, and 10 phosphocreatine. Giga-ohm seals were obtained using glass electrodes (1.1-1.4 MOhm resistance) and allowed to stabilize for 1-2 min before break-in. No overt difference in cell firing behavior was observed between genotypes in cell attached configuration (data not shown). Series resistance, capacitance, and membrane resistance were measured shortly after break-in using the average of 20 5 mV test pulses, and were monitored during and upon completion of all recordings. Cells were voltage-clamped at -58 mV using an AxoPatch1B amplifier and current was recorded at 1k Hz. In addition to spontaneous pacemaker activity in cell-attached configuration, DA cell identity was verified by the presence of an Ih current in response to a 50 mV hyperpolarizing step. D2-IPSCs were evoked using a platinum bipolar stimulator connected to a stimulus isolator (Warner Precision instruments) placed in the slice caudal to recorded cell bodies. All recordings are the average of 3-4 repetitions conducted once per minute that are baselined to the 500 ms immediately preceding stimulation before averaging. All recording and analysis of D2-IPSC experiments was conducted using AxoGraph software (Berkeley CA).

Acknowledgements

We thank Stefanie Kaech Petrie, Hannah Bronstein, and the Advanced Imaging Core at the Jungers Center for microscopy assistance, and Gary Westbrook and Dennis Weingarten for helpful comments on the manuscript. This work was supported by a Whitehall Foundation Grant (SLJ), the National Institutes of Health (RO1 DA004523 to JTW), the National Institute on Drug Abuse (T32DA007262 to JJL), and the National Institute of Neurological Disorders and Stroke (P30 NS061800 to the OHSU Imaging Center).

Competing Interests

The authors have no competing interests to declare.

References

Bacaj, T., Wu, D., Yang, X., Morishita, W., Zhou, P., Xu, W., Malenka, R. C., & Südhof, T. C. (2013). Synaptotagmin-1 and synaptotagmin-7 trigger synchronous and asynchronous phases of neurotransmitter release. *Neuron*, *80*(4), 947-959.

Banerjee, A., Lee, J., Nemcova, P., Liu, C., & Kaeser, P. S. (2020). Synaptotagmin-1 is the Ca2+ sensor for fast striatal dopamine release. *Elife*, *9*, e58359.

Beeler, J. A., & Dreyer, J. K. (2019). Synchronicity: the role of midbrain dopamine in whole-brain coordination. *eneuro*, *6*(2).

Chakrabarti, S., Kobayashi, K. S., Flavell, R. A., Marks, C. B., Miyake, K., Liston, D. R., Fowler, K. T., Gorelick, F. S., & Andrews, N. W. (2003). Impaired membrane resealing and autoimmune myositis in synaptotagmin VII–deficient mice. *The Journal of cell biology, 162*(4), 543-549.

Chen, B. T., & Rice, M. E. (2001). Novel Ca2+ dependence and time course of somatodendritic dopamine release: substantia nigra versus striatum. *Journal of Neuroscience, 21*(19), 7841-7847.

Chen, C., Satterfield, R., Young Jr, S. M., & Jonas, P. (2017). Triple function of synaptotagmin 7 ensures efficiency of high-frequency transmission at central GABAergic synapses. *Cell reports, 21*(8), 2082-2089.

Condon, A. F., Robinson, B. G., Asad, N., Dore, T. M., Tian, L., & Williams, J. T. (2021). The residence of synaptically released dopamine on D2 autoreceptors. *Cell reports, 36*(5), 109465.

Covey, D. P., Juliano, S. A., & Garris, P. A. (2013). Amphetamine elicits opposing actions on readily releasable and reserve pools for dopamine. *PLoS One, 8*(5), e60763.

Cragg, S. J. (2003). Variable dopamine release probability and short-term plasticity between functional domains of the primate striatum. *Journal of Neuroscience, 23*(10), 4378-4385.

Day, J. J., Roitman, M. F., Wightman, R. M., & Carelli, R. M. (2007). Associative learning mediates dynamic shifts in dopamine signaling in the nucleus accumbens. *Nature neuroscience, 10*(8), 1020-1028.

Dean, C., Dunning, F. M., Liu, H., Bomba-Warczak, E., Martens, H., Bharat, V., Ahmed, S., & Chapman, E. R. (2012). Axonal and dendritic synaptotagmin isoforms revealed by a pHluorin-syt functional screen. *Mol Biol Cell, 23*(9), 1715-1727. https://doi.org/10.1091/mbc.E11-08-0707

Delignat-Lavaud, B., Kano, J., Ducrot, C., Massé, I., Mukherjee, S., Giguère, N., Moquin, L., Lévesque, C., Nanni, S. B., & Bourque, M.-J. (2021). The calcium sensor synaptotagmin-1 is critical for phasic axonal dopamine release in the striatum and mesencephalon, but is dispensable for basic motor behaviors in mice. *bioRxiv*.

Doussau, F., Schmidt, H., Dorgans, K., Valera, A. M., Poulain, B., & Isope, P. (2017). Frequency-dependent mobilization of heterogeneous pools of synaptic vesicles shapes presynaptic plasticity. *Elife, 6*, e28935.

Fiorillo, C. D., Newsome, W. T., & Schultz, W. (2008). The temporal precision of reward prediction in dopamine neurons. *Nature neuroscience, 11*(8), 966-973.

Fiorillo, C. D., Tobler, P. N., & Schultz, W. (2003). Discrete coding of reward probability and uncertainty by dopamine neurons. *Science, 299*(5614), 1898-1902.

Ford, C. P., Gantz, S. C., Phillips, P. E., & Williams, J. T. (2010). Control of extracellular dopamine at dendrite and axon terminals. *Journal of Neuroscience, 30*(20), 6975-6983.

Gonon, F. (1988). Nonlinear relationship between impulse flow and dopamine released by rat midbrain dopaminergic neurons as studied by in vivo electrochemistry. *Neuroscience, 24*(1), 19-28.

Hui, E., Bai, J., Wang, P., Sugimori, M., Llinas, R. R., & Chapman, E. R. (2005). Three distinct kinetic groupings of the synaptotagmin family: candidate sensors for rapid

and delayed exocytosis. *Proceedings of the National Academy of Sciences*,
102(14), 5210-5214.
Jackman, S. L., & Regehr, W. G. (2017). The mechanisms and functions of synaptic
facilitation. *Neuron*, *94*(3), 447-464.
Jackman, S. L., Turecek, J., Belinsky, J. E., & Regehr, W. G. (2016). The calcium
sensor synaptotagmin 7 is required for synaptic facilitation. *Nature*, *529*(7584),
88-91. https://doi.org/10.1038/nature16507
Lin, M., Mackie, P. M., Shaerzadeh, F., Gamble-George, J., Miller, D. R., Martyniuk, C.
J., & Khoshbouei, H. (2021). In Parkinson's patient-derived dopamine neurons,
the triplication of α-synuclein locus induces distinctive firing pattern by impeding
D2 receptor autoinhibition. *Acta neuropathologica communications*, *9*(1), 1-25.
Liu, C., Kershberg, L., Wang, J., Schneeberger, S., & Kaeser, P. S. (2018). Dopamine
secretion is mediated by sparse active zone-like release sites. *Cell*, *172*(4), 706-
718. e715.
Liu, H., Bai, H., Hui, E., Yang, L., Evans, C. S., Wang, Z., Kwon, S. E., & Chapman, E.
R. (2014). Synaptotagmin 7 functions as a Ca2+-sensor for synaptic vesicle
replenishment. *Elife*, *3*, e01524.
Luo, F., & Südhof, T. C. (2017). Synaptotagmin-7-mediated asynchronous release
boosts high-fidelity synchronous transmission at a central synapse. *Neuron*,
94(4), 826-839. e823.
Mendez, J. A., Bourque, M.-J., Fasano, C., Kortleven, C., & Trudeau, L.-E. (2011).
Somatodendritic dopamine release requires synaptotagmin 4 and 7 and the
participation of voltage-gated calcium channels. *Journal of Biological Chemistry*,
286(27), 23928-23937.
Montague, P. R., McClure, S. M., Baldwin, P., Phillips, P. E., Budygin, E. A., Stuber, G.
D., Kilpatrick, M. R., & Wightman, R. M. (2004). Dynamic gain control of
dopamine delivery in freely moving animals. *Journal of Neuroscience*, *24*(7),
1754-1759.
Otomo, K., Perkins, J., Kulkarni, A., Stojanovic, S., Roeper, J., & Paladini, C. A. (2020).
In vivo patch-clamp recordings reveal distinct subthreshold signatures and
threshold dynamics of midbrain dopamine neurons. *Nature communications*,
11(1), 1-15.
Patriarchi, T., Cho, J. R., Merten, K., Howe, M. W., Marley, A., Xiong, W.-H., Folk, R.
W., Broussard, G. J., Liang, R., & Jang, M. J. (2018). Ultrafast neuronal imaging
of dopamine dynamics with designed genetically encoded sensors. *Science*,
360(6396).
Robinson, B. G., Cai, X., Wang, J., Bunzow, J. R., Williams, J. T., & Kaeser, P. S.
(2019). RIM is essential for stimulated but not spontaneous somatodendritic
dopamine release in the midbrain. *Elife*, *8*, e47972.
Saunders, A., Macosko, E. Z., Wysoker, A., Goldman, M., Krienen, F. M., de Rivera, H.,
Bien, E., Baum, M., Bortolin, L., & Wang, S. (2018). Molecular diversity and
specializations among the cells of the adult mouse brain. *Cell*, *174*(4), 1015-
1030. e1016.
Schiemann, J., Schlaudraff, F., Klose, V., Bingmer, M., Seino, S., Magill, P. J., Zaghloul,
K. A., Schneider, G., Liss, B., & Roeper, J. (2012). K-ATP channels in dopamine

substantia nigra neurons control bursting and novelty-induced exploration.
484   *Nature neuroscience*, *15*(9), 1272-1280.
485 Schultz, W., Dayan, P., & Montague, P. R. (1997). A neural substrate of prediction and
486   reward. *Science*, *275*(5306), 1593-1599.
487 Tsai, H.-C., Zhang, F., Adamantidis, A., Stuber, G. D., Bonci, A., De Lecea, L., &
488   Deisseroth, K. (2009). Phasic firing in dopaminergic neurons is sufficient for
489   behavioral conditioning. *Science*, *324*(5930), 1080-1084.
490 Turecek, J., Jackman, S. L., & Regehr, W. G. (2016). Synaptic specializations support
491   frequency-independent Purkinje cell output from the cerebellar cortex. *Cell*
492   *reports*, *17*(12), 3256-3268.
493 Turecek, J., Jackman, S. L., & Regehr, W. G. (2017). Synaptotagmin 7 confers
494   frequency invariance onto specialized depressing synapses. *Nature*, *551*(7681),
495   503-506. https://doi.org/10.1038/nature24474
496 Turecek, J., & Regehr, W. G. (2018). Synaptotagmin 7 mediates both facilitation and
497   asynchronous release at granule cell synapses. *Journal of Neuroscience*, *38*(13),
498   3240-3251.
499 Yee, A. G., Forbes, B., Cheung, P. Y., Martini, A., Burrell, M. H., Freestone, P. S., &
500   Lipski, J. (2019). Action potential and calcium dependence of tonic
501   somatodendritic dopamine release in the Substantia Nigra pars compacta.
502   *Journal of neurochemistry*, *148*(4), 462-479.
503