SIMBA: SIngle-cell eMBedding Along with features

Huidong Chen[1, 2, 3], Jayoung Ryu[1, 2, 4], Michael E. Vinyard[1, 2, 3, 5], Adam Lerer[6]+, Luca Pinello[1, 2, 3]+

1. Molecular Pathology Unit, Center for Cancer Research, Massachusetts General Hospital,
Charlestown, MA, USA
2. Department of Pathology, Harvard Medical School, Boston, MA, USA
3. Broad Institute of Harvard and MIT, Cambridge, MA, USA
4. Department of Biomedical Informatics, Harvard Medical School, Boston, MA, USA
5. Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA, USA
6. Facebook AI Research

+Corresponding author

Abstract

Recent advances in single-cell omics technologies enable the individual and joint profiling of
cellular measurements including gene expression, epigenetic features, chromatin structure and
DNA sequences. Currently, most single-cell analysis pipelines are cluster-centric, i.e., they first
cluster cells into non-overlapping cellular states and then extract their defining genomic
features. These approaches assume that discrete clusters correspond to biologically relevant
subpopulations and do not explicitly model the interactions between different feature types. In
addition, single-cell methods are generally designed for a particular task as distinct single-cell
problems are formulated differently. To address these current shortcomings, we present
SIMBA, a graph embedding method that jointly embeds single cells and their defining features,
such as genes, chromatin accessible regions, and transcription factor binding sequences into a
common latent space. By leveraging the co-embedding of cells and features, SIMBA allows for
the study of cellular heterogeneity, clustering-free marker discovery, gene regulation inference,
batch effect removal, and omics data integration. SIMBA has been extensively applied to
scRNA-seq, scATAC-seq, and dual-omics data. We show that SIMBA provides a single framework
that allows diverse single-cell analysis problems to be formulated in a unified way and thus
simplifies the development of new analyses and integration of other single-cell modalities.

Introduction

Recent progress in single-cell molecular profiling technologies have dramatically advanced our
ability to define cell types and states as well as discover key genes and regulatory regions in
development and disease. We can now measure multiple modalities of individual cells. At the
same time, the number of cells we are able to profile has increased rapidly. The emergence of
single-cell multi-omics technologies allows for the measurements of multiple cellular layers,
including genomics, epi-genomics, transcriptomics, and proteomics. Such assays have
pioneered an avenue to better understand the interplay between these 'omic' layers and cell

states based on diverse genomic and molecular features including genes, regulatory elements,
transcription factors. While single-cell multi-omic assays have quickly evolved towards the
incorporation of more modalities with increasing resolution, harnessing their full potential
poses significant computational challenges.
48
Many computational methods have been developed for single-cell single-modality analysis
(e.g., scRNA-seq or scATAC-seq analysis) [1-4]. Common to these methods is a workflow with
several standard steps, which typically include feature selection, dimension reduction,
clustering, and differential feature detection. These "cluster-centric" analysis methods rely on
accurately defined clustering solutions to discover meaningful and informative marker features.
Unfortunately, clustering solutions may range widely within the space of the user-defined
clustering resolution (number of clusters) and the chosen clustering algorithm. These
parameters may markedly influence the resulting cluster assignment and clusters may not
always correspond to the correct cell populations, thereby leading to inconsistent and
potentially misleading biological annotations[5]. Although initial efforts have been made recently
to develop clustering-free approaches to discover informative genes, they are specifically
designed for extracting gene signatures [6, 7] or identifying perturbations between experimental
conditions[8] from scRNA-seq data, and are therefore limited to single-modality and single-task
analysis.
63
In addition to single-batch/modality analysis, approaches have also been proposed for multi-
batch and cross-modality analysis, such as multimodal analysis (distinct cellular parameters are
measured in the same cell)[9], batch correction (the same cellular parameter is measure in
different batches) [10-12], and integration of multi-omics datasets (distinct cellular parameters are
measured in different cells)[11, 12]. These approaches play a critical role in removing batch effects
that confound true biological variation, improving the characterization of cell states by
leveraging the unique strengths of each assay, and providing insights into the complex
mechanisms of gene regulation. However, these tasks are formulated differently from those in
single-batch/modality settings and thus require development of new dedicated analysis
techniques. Also, while multiple types of cellular parameters (features) might be present, the
relation between features cannot be exploited directly by most current methods. Furthermore,
similar to single-batch/modality analysis methods, these methods identify marker features
based on groups of cells obtained by clustering and therefore are limited to clustering solutions.
Additionally, instead of directly identifying marker features in the integrated space, most batch
correction/multi-omics integration methods need to first detect marker features in the
uncorrected/unintegrated original space of each batch/modality independently, and then
combine them, thus resulting in potentially inconsistent interpretations between
batches/modalities.
82
To overcome the limitations in both single-batch/modality analysis and multi-batch/cross-
modality analysis, we propose SIMBA (**SIngle-cell eMBedding Along with features**), a versatile
single-cell embedding method that co-embeds cells and features into a shared latent space, in
which the relation between cells and features or between two types of features (e.g., genes,
peaks, or DNA sequences) can be assessed based on their proximity in the resulting embedding.

By formulating single-cell analyses as multi-entity graph embedding problems, SIMBA can be
used to solve popular single-cell tasks in a unified framework that would otherwise require the
development of distinct specialized approaches for each task. For each task, SIMBA constructs a
graph, wherein differing entities (i.e., cells and features) are represented as nodes and relations
between these entities are encoded as edges. Once the graph is constructed, SIMBA then
applies a multi-entity graph embedding algorithm derived from social networking technologies
as well as a Softmax-based transformation to embed the nodes/entities of the graph into a
common low-dimensional space wherein cells and features can be analyzed based on their
distance. We show that the SIMBA framework can perform many common and important
single-cell analysis tasks, including: 1) dimensionality reduction techniques for studying cellular
states; 2) clustering-free marker detection based on the similarity between single cells and
features; 3) Single-cell multimodal analysis and the study of gene regulation; 4) batch
correction and omics integration analysis as well as the simultaneous identification of marker
features. SIMBA is adapted to these diverse analysis tasks by simply modifying how the input
graph is constructed from the relevant single-cell data.
103
We extensively tested *SIMBA* in multiple scRNA-seq, scATAC-seq and dual-omics datasets
covering popular single-cell tasks including scRNA-seq analysis, scATAC-seq analysis, multimodal
analysis, batch correction, and multi-omics integration. We demonstrate that *SIMBA* jointly
learns low-dimensional representations of both cells and features and thus enables the ability
to simultaneously study cellular heterogeneity as well as proximity-based marker feature
detection or gene regulation inference in a clustering-free way. We also demonstrate that
SIMBA performs comparably to or better than current state-of-the-art methods specifically
developed for each task.
112
113 We believe that SIMBA, as a broadly applicable approach for single cell omics study, will
114 simplify the burden of developing methods for new single-cell tasks and measurements, while
115 increasing interpretability of cellular mechanisms and functions.
116

Results

118
Overview of SIMBA
119
SIMBA is a single-cell embedding method with support for single- or multi- modality analyses
that embeds cells and their associated genomic features into a shared latent space, generating
interpretable and comparable embeddings of cells and features. It leverages recent graph
embedding techniques that have been successful in modeling complex and hierarchical
information present in natural languages, social networks, and other domains, as "knowledge
graphs". In our case, the graph encodes different components of cellular regulatory circuits and
the relations between them.
127
SIMBA first encodes different types of entities such as cells, genes, open chromatin regions
(peaks or bins), transcription factor (TF) motifs, and *k*-mers (short sequences of a specific
length, *k*), into a single graph (**Fig. 1**, **Methods**) where each node represents an individual entity

and edges indicate relations between entities. For example, if a gene is expressed in a cell, an
edge is created between the gene and cell. The weight of this edge is determined by the gene
expression level. Similarly, an edge is added between a cell and a chromatin region if the region
is open in this cell, or between a chromatin region and a TF motif if the TF motif is found in the
region.

Figure1. SIMBA framework overview. SIMBA co-embeds cells and various features measured
during single-cell experiments into a shared latent space to accomplish both common tasks
involved in single-cell data analysis as well as tasks, which remain as open problems in single-cell
genomics. **(Left)** Examples of possible biological entities may be encoded by SIMBA including
cells, gene expression measurements, chromatin accessible regions, TF motifs, and k-mer
sequences found in reads. **(Middle)** SIMBA embedding plot with multiple types of entities into a
low-dimensional space. All entities represented as shapes (cell = circle, peak = triangle, gene =
square, TF motif = star, k-mer = hexagon) are colored by relevant cell type (green, orange, and
blue in this example). Non-informative features are colored dark grey. Within the graph, each
entity is a node, and an edge indicates a relation between entities (e.g., a gene is expressed in a
cell, a chromatin region is accessible in a cell, or a TF motif/k-mer is present within an open
chromatin region, etc.). Once connected in a graph, these entities may be embedded into a
shared low-dimensional space, with cell-type specific entities embedded in the same
neighborhood and non-informative features embedded elsewhere. **(Right)** Common single-cell
analysis tasks that may be accomplished using SIMBA.

Once the input graph is constructed, a low-dimensional representation of the graph nodes is
then computed using an unsupervised graph embedding method (**Methods**). This graph
embedding procedure leverages the PyTorch-BigGraph framework [13], which allows SIMBA to
scale to millions of cells (**Methods**). The resulting joint embedding of cells and features not only
reconstructs the heterogeneity of cells but also allows for the discovery of the defining features
for each single cell without relying on a clustering solution, separating cell-type specific features
from the non-informative features. In fact, the proximity between entity embeddings is
informative of the potential importance of a feature to a cell and the interplay between
features. When multiple types of features (e.g., transcriptomic and epigenetic features) are co-
embedded, SIMBA provides an intuitive way to study gene regulation and the regulatory
mechanisms underlying cellular differentiation and cell type formation.

Graph construction is inherently flexible, enabling SIMBA to be applied to a wide variety of
single-cell tasks. In the following sections, we demonstrate the application of SIMBA to several
popular single-cell tasks including scRNA-seq, scATAC-seq, multimodal analysis, batch
correction and multi-omics integration (**Fig. 1**).
166
167
SIMBA enables simultaneous learning of cellular heterogeneity and individual-cell-level
marker genes in scRNA-seq analysis

Single-cell RNA sequencing (scRNA-seq) is the most robust and widely used measurement to
profile single cells. **Figure 2a** provides an illustrative overview of the SIMBA graph construction
and the resulting low-dimensional embedding matrix of both cells and genes. Here we show
how SIMBA enables simultaneous dimensionality reduction and clustering-free marker gene
detection in scRNA-seq analysis. We applied SIMBA to a popular PBMCs dataset from 10x
Genomics (**Supplementary Table 1**), which is used in the tutorials of both Scanpy[2] and Seurat[1] to
illustrate their workflows. After the standard preprocessing steps including normalization and
log-transformation, SIMBA discretizes the gene expression matrix into multiple gene expression
levels (five levels, by default). The input graph is then constructed wherein two types of nodes –
cells and genes are connected by edges that embody the relation between them and are
weighted according to the corresponding multiple levels of gene expression. SIMBA then
generates embeddings of these nodes through a graph embedding procedure (**Fig. 2a;**
Methods).

We first visualized the SIMBA embeddings of cells using UMAP, which is a comparable output to
other current single-cell analysis methods. Each of the previously assigned eight cell types,
including B cells, megakaryocytes, CD14 monocytes, FCGR3A monocytes, dendritic cells, NK
cells, CD4 T, and CD8 T cells, was clearly separated (**Fig. 2b**). We next applied UMAP to visualize
the SIMBA embeddings of cells and genes together (**Fig. 2c**). The same set of marker genes used
to annotate these cells from Scanpy[2] was highlighted on the UMAP plot. In addition, as a
control, we also show the locations of two housekeeping genes *GAPDH* and *B2M*, which would
not be expected to associate with any particular cell type. From the UMAP plot, we can see that
SIMBA not only was able to embed major-cell-group specific genes to the correct locations

(e.g., *IL7R* was embedded into CD4T cells and *MS4A1* was embedded into B cells), but also was
robust to rare-cell-group specific genes (e.g., *PPBP* was embedded into megakaryocytes). On
the contrary, non-informative or non-cell-type specific genes such as the aforementioned
housekeeping genes were embedded in the middle of all cell groups.

Figure 2. Single-cell RNA-seq analysis of the 10x PBMCs dataset using SIMBA. **(a)** SIMBA graph
construction and embedding in scRNA-seq analysis. Biological entities including cells and genes
are represented as shapes and colored by relevant cell types (green and orange). Non-
informative genes are colored dark grey. Gene expression measurements for each cell are
organized into a cell-by-gene matrix. These normalized non-negative observed values undergo
discretization into five gene expression levels. Cells and genes are then assembled into a graph
with nodes representing cells and genes, and edges between them representing different gene
expression levels. This graph may then be embedded into a lower dimensional space resulting in
a #entities x #dimension (by default, 50) SIMBA embedding matrix. **(b)** UMAP visualization of
SIMBA embeddings of cells colored by cell type. **(c)** UMAP visualization of SIMBA embeddings of

cells and genes. Cells are colored according to cell type as defined in b. Genes are colored slate
blue. Cell-type-specific marker genes and housekeeping genes recovered by Scanpy are indicated
with text and arrows. Genes highlighted in red are shown in **d**, **e**, and **f**. **(d)** SIMBA barcode plots
of genes *CST3*, *MS4A1*, *NKG7*, and *GAPDH*. The x-axis indicates the ordering of a cell as ranked by
the probability for each cell to be associated with a given gene. The y-axis describes the
probability. The sum of probability over all cells is equal to 1. Each cell is one bar and colored
according to cell type as defined in b. **(e)** SIMBA ranking of genes based on the proposed metrics.
All the genes are plotted according to the Gini index against max score. The same set of genes as
in **c** are annotated. **(f)** UMAP visualization of SIMBA embeddings of cells colored by gene
expression of (left to right): *CST3*, *NKG7*, *MS4A1*, and *GAPDH*.
217

In addition to visualizing all the entities at once using UMAP, SIMBA also provides a feature-
specific 'barcode plot' to visualize the estimated probability of assigning a feature to a cell (**Fig.
2d, Methods**). The barcode plots in **Figure 2d** offer a rank-ordered probability of a given gene
being associated with each cell (colored by cell type) where the total probability over all cells
sums to one. An imbalance in probability indicates cell-type-specific association of a gene to a
sub-population of cells, whereas a uniform probability distribution indicates a non-cell-type-
specific gene. **Figure 2d** displays barcode plots for four genes, which are correspondingly
highlighted in red, in **Figure 2c**. Three of these genes are commonly used marker genes for
identifying subpopulations within PBMCs datasets (*CST3* for monocytes and dendritic cells,
MS4A1 for B cells, and *NGK7* for NK and CD8T cells). On the contrary, *GAPDH* is a housekeeping
gene expressed in all cell types. For marker genes, we observed a clear excess in the probability
of assigning each gene to their respective cell types. Conversely, for *GAPDH*, we observed a
more uniform distribution with much lower probability of associating that gene with the top-
ranked cells. Hence, SIMBA barcode plots serve as an informative way of visualizing gene
expression patterns by showing the cell assignment probability distribution.

In addition, SIMBA also provides several quantitative metrics, including max value, Gini index,
standard deviation, and entropy, to assess cell type specificity of various features (**Methods**). As
an example, by inspecting the gene metric plot of max value (a higher value indicates higher
cell-type specificity) vs Gini index (a higher value indicates higher cell-type specificity), we
observed that the marker genes (e.g., *CST3*, *NKG7*, *MS4A1*) fall in the upper right corner, as
opposed to housekeeping genes (e.g., *GAPDH*) in the lower left corner (**Fig. 2e**). Similar
separation is observed with other metrics (**Supplementary Fig. 1b**). The cell type specificity of
the selected marker genes was further confirmed by visualizing their expression pattern on
UMAP plots (**Fig. 2f and Supplementary Fig. 1c**), accompanied by SIMBA barcode plots
(**Supplementary Fig. 1d**).

To demonstrate that SIMBA provides a more robust means of detecting marker genes than the
statistical-tests-based methods implemented by tools such as Scanpy and Seurat, we compared
top 600 marker genes identified by SIMBA (based on max value and Gini index) with those
identified by Scanpy (based on the Wilcoxon rank-sum test, one of the statistical tests
recommended in Scanpy's tutorial) (**Supplementary Fig. 2a**). Upon comparison, we observed

that nearly half of the marker genes discovered by SIMBA overlap with the marker genes
identified by Scanpy (**Supplementary Fig. 2a**). However, on inspection of the top non-
overlapping marker genes, genes identified by SIMBA are found to be enriched only within
certain groups of cells (**Supplementary Figs. 2b and 2c**) while genes identified by Scanpy but
not by SIMBA include the housekeeping gene *B2M* and multiple ribosomal protein genes (e.g.,
RPS3 and *RPS6*) that are expressed ubiquitously in all cell types (**Supplementary Figs. 2b and**
2d). In addition, we noticed that some of the identified genes by SIMBA can be potentially
recovered using different statistical tests proposed in Scanpy. For example, IL7R was identified
only by using the t-test and FCER1A was identified only by using the Wilcoxon rank-sum test as
also noted in the Scanpy's tutorial. This example illustrates the limitations of a cluster-centric
approach combined with multiple statistical testing methods. In contrast, SIMBA successfully
identified both *IL7R* and *FCER1A* as informative genes with a single procedure and without
clustering the cells (**Fig. 2e and Supplementary Fig. 1b**).

Lastly, we showed that SIMBA does not require variable gene selection, which is an essential
step in standard scRNA-seq pipelines such as Seurat or Scanpy. SIMBA produces very similar
embeddings for cells and genes with and without variable gene selection (**Supplementary Fig.**
2e), though we observed that variable gene section does improve efficiency of the training
procedure.

SIMBA enables simultaneous characterization of cellular states and cis-regulatory elements
by jointly modeling accessible sites and DNA sequences in scATAC-seq analysis
268
As one of the most popular single-cell epigenomic techniques, single-cell assay for transposase-
accessible chromatin using sequencing (scATAC-seq) has been widely used to profile regions of
open chromatin and identify functional *cis*-regulatory elements such as enhancers and active
promoters. In scATAC-seq, cells are characterized by different types of features [14], such as regions
of accessible chromatin ("peaks" or "bins") and *cis*-regulatory elements (DNA sequences) within
these accessible regions including transcription factor (TF) motifs or *k*-mers. Unlike existing
methods that use only peaks/bins or the sequence content found within them, in SIMBA both
types of features can be encoded into a single graph. When only peaks/bins are used, SIMBA
constructs a graph with nodes representing cells and chromatin regions (peaks or bins) and edges
indicating the accessibility of the chromatin regions in cells (**Fig. 3a**). When the DNA sequences
for chromatin regions are available, SIMBA can also encode DNA sequences including TF motifs
and *k*-mers into the graph by adding edges between these entities as nodes and the existing
chromatin region nodes. The edges in this case indicate the presence of TF motifs/*k*-mers within
these chromatin-accessible regions. Through the embedding procedure, SIMBA generates
embeddings of cells along with peaks and DNA sequences (**Methods**). Therefore, SIMBA enables
dimensionality reduction by leveraging both chromatin accessible regions and cis-regulatory
sequences. Simultaneously, it highlights the cell-type-specific open chromatin regions and
regulatory DNA sequences in a clustering-free way.
287
To demonstrate the value of SIMBA embeddings for scATAC-seq analysis, we first applied
SIMBA to a scATAC-seq data of 2,034 human hematopoietic cells with FACS-characterized cell

types[15](**Supplementary Table 1**). For the embeddings of cells alone, as shown using UMAP (**Fig.**
3b), SIMBA accurately embeds cells such that cells belonging to distinct cell types are visually
separated. In addition to cells, SIMBA can also embed various types of features. The UMAP plot
in **Figure 3c** highlights how distinct features from both positional (peaks) as well as sequence-
content (TF motifs and k-mers) information are embedded together based on their biological
relations. Notably, these highlighted features that are embedded within each cell type all have
high cell-type specificity scores (shown in the upper right part of SIMBA metric plots in **Figure**
3d).

299
300

Figure 3. Single-cell ATAC-seq analysis of the human hematopoiesis dataset using SIMBA. **(a)**
SIMBA graph construction and embedding in scATAC-seq analysis. Biological entities including
cells, peaks/bins, TF motifs, k-mers are represented as shapes and colored by relevant cell types
(green and orange). Non-informative features are colored dark grey. Cells and chromatin
accessible features (peaks / bins) are organized into a cell x peaks / bins matrix. When sequence
information (TF motif or *k*-mer sequence) within these regions is available, they can be organized

into two sub-matrices to associate a TF motif or k-mer sequence with each peak/bin. These
constructed feature matrices are then binarized and assembled into a graph. When a single
feature (chromatin accessibility) is used, the graph encodes cells and peaks/bins as nodes. When
multiple features (both chromatin accessibility and DNA sequences) are used, this graph may
then be extended with the addition of TF motifs and k-mer sequences as nodes connected. Finally,
SIMBA embeddings of these entities are generated through a graph embedding procedure. **(b)**
UMAP visualization of SIMBA embeddings of cells colored by cell type. **(c)** UMAP visualization of
SIMBA embeddings of cells and features including TF motifs, k-mers, and peaks. Cells are colored
by cell type while motifs, k-mers, and peaks are colored green, blue, and pink, respectively. Cell
type specific features that are embedded near their corresponding cell types are indicated as the
text labels (colored according to feature type) with arrows. **(d)** SIMBA metric plots of TF motifs,
k-mers, and peaks. Cell-type specific features annotated in **(c)** are highlighted. **(e)** Genomic tracks
of aligned scATAC-seq fragments, separated and colored by cell type. Two marker peaks P1 and
P2 in red are shown beneath the alignment. Within the peak P1, k-mer GATAAG and its
resembling GATA1 motif logo are highlighted. **(f)** UMAP visualization of SIMBA embeddings of
cells colored by TF activity scores of the GATA1 motif and k-mer GATAAG enrichment. **(g)** SIMBA
barcode plots of the GATA1 motif, k-mer GATAAG, and the two peaks P1 and P2. Cells are colored
according to cell type labels described above. Dotted red line indicates the same cutoff used in
all four plots.
326
327 This analysis led to several key findings in human hematopoietic differentiation.
328
329 First, SIMBA analysis identified key master regulators of hematopoiesis. As highlighted in **Fig.**
330 **3c**, we observed that motifs of previously reported TFs were embedded near their respective
331 cell types in UMAP. For example, the GATA1 and GATA3 motifs are close to megakaryocyte-
332 erythroid progenitor (MEP) cells[16]; the PAX5 and EBF1 motifs are close to common lymphoid
333 progenitor (CLP) cells[17]; the CEBPB and CEBPD motifs are close to monocytes (mono)[18].
334
Second, SIMBA analysis identified an unbiased set of DNA sequences, i.e., *k*-mers, that
represent important TF binding motifs involved in hematopoiesis, enabling *de novo* motif
discovery. We observed that these *k*-mers were embedded near their resembling TF binding
motifs and relevant cell subpopulations (**Fig. 3c** and **3e**, **Supplementary Fig. 3b**). For example,
the DNA sequence CAGCTG is embedded in plasmacytoid dendritic cells (pDCs), and this
sequence matches the TCF12 binding motif, which controls dendritic cell lineage specification.
To further illustrate the interpretability of the SIMBA embeddings of TF motifs and k-mers, we
calculated per-cell TF activity scores[19] (high-variance TF motifs/*k*-mers) and visualized them on
SIMBA embeddings of cells . As shown in **Figure 3f**, the GATA1 TF motif and k-mer GATAAG that
were both embedded in MEP cells by SIMBA, also showed high-level activity in MEP cells. The
consistency between SIMBA embedding and TF activity was observed for most of other TF
motifs and *k*-mers as well (**Supplementary Fig. 3a, 3b**).
347
Third, SIMBA analysis identified differential accessible regions that may mediate cell-type
specific gene regulation. For example, the two peaks at chr19:12997999-12998154 (P1) and
chr19:12998329-12998592 (P2) that were embedded within MEP cells were almost exclusively

observed in MEP cells on KLF1 genome track (**Fig. 3e**). Interestingly, P1, upstream of *KLF1*,
contains the *k*-mer GATAAG that matches the GATA1 binding motif, while transcription factor
GATA1 is known to regulate the gene *KLF1* and plays a pivotal role in erythroid cell and
megakaryocyte development[20]. Therefore, by embedding these MEP-cell-related regulatory
elements into the neighborhood of MEP cells, SIMBA demonstrates a new analytic way for
studying cell-type-specific epigenetic landscape of the genome. To further validate the
differential accessible regions identified by SIMBA, we randomly picked 100 peaks within each
cell type in SIMBA co-embedding space. From the heatmap of chromatin accessibility, we
clearly see that the peaks embedded in the neighborhood of each cell type by SIMBA shows
strong cell-type specificity and this is robust to the number of cells within each cell type
(**Supplementary Fig. 3c**).
362
363 Although SIMBA strongly differs from existing computational methods for scATAC-seq analysis in
364 that SIMBA enables the co-embedding of cells and features, we still compared the resulting
365 SIMBA embeddings of cells with these specialized methods by their ability to distinguish cell types.
366 Our analyses show that SIMBA performs as well as or better than the methods evaluated, further
367 demonstrating the wide utility of SIMBA (**Supplementary Figs. 4 and 5; Supplementary Note 1**).
368
369
370 **SIMBA enables simultaneous learning of cellular heterogeneity and gene regulatory circuits**
371 **from integrated analysis of single-cell multimodal data**
372
scRNA-seq and scATAC-seq are two of the most widely adopted single-cell sequencing
technologies, but they are limited to measuring only a single aspect of cell state at a time. To
improve our ability to interrogate cellular states, several single-cell dual-omics technologies
have been recently developed [21-24] to jointly profile transcriptome and chromatin accessibility
within the same individual cells, therefore providing the potential to correlate gene expression
with accessible regulatory elements and further delineate the yet elusive principles of gene
regulation.
380
In this section, we demonstrate how SIMBA enables simultaneous learning of cellular
heterogeneity and gene regulatory circuit from single-cell multiome data. We applied SIMBA to
three recent single-cell dual-omics technologies: SHARE-seq[22], SNARE-seq[21], and a multiome
PBMCs dataset from 10x Genomics (**Supplementary Table 1**).
Figure 4a illustrates the procedure of graph construction and generation of the final SIMBA
embedding matrix. Briefly, for scRNA-seq, the gene expression matrix is discretized to generate
different levels of gene expression. For scATAC-seq, both the chromatin accessibility matrix and
motif/*k*-mer match matrix are binarized. In this graph, there are five entity (node) types,
including cells, genes, peaks, motifs, and *k*-mers. For scRNA-seq, an edge indicates whether a
gene is expressed in a cell and its weight indicates the gene expression level (five levels, by
default). For scATAC-seq, an edge indicates whether a peak is present in a cell or if a TF motif/*k*-
mer is present within a peak.

Figure 4. Multimodal analysis of the SHARE-seq hair follicle dataset using SIMBA. **(a)** SIMBA graph construction and embedding in multimodal analysis. Overview of SIMBA's approach to multimodal (scRNA-seq + scATAC-seq) data analysis. **(b)** SIMBA metric plots of genes, TF motifs, and peaks. All these features are plotted according to the Gini index against max score. Cell-type specific genes, TF motifs, and peaks are highlighted. **(c)** UMAP visualization of SIMBA embeddings of cells (Top-left), cells and genes (Top-right), and cells along with genes, TF motifs, and peaks (Bottom). **(d)** Ranked scatter plot of candidate master regulators as identified by SIMBA. **(e)** Schematic description of SIMBA's strategy for identifying target genes given a master regulator. **(f)** Top 30 target genes of transcription factors Lef1 and Hoxc13 as inferred by SIMBA.

To demonstrate the usefulness and versatility of the SIMBA embeddings, we analyzed the cell populations undergoing hair follicle differentiation from mouse skin profiled with SHARE-seq.

First, we used SIMBA to assess the cell-type specificity of different types of features, including genes, TF motifs, and peaks (**Fig. 4b, Methods**). As shown in **Figure 4b**, genes (e.g., *Lef1* and *Hoxc13*), which are associated with hair follicles each have relatively higher max values and Gini index scores. Similarly, TF motifs and peaks proximal to the genomic loci of these genes also

score in the upper right quadrant of the metric plots. SIMBA's cell-type specificity metrics
successfully revealed the key genes and regulatory factors important to the hair follicle
differentiation process.

Next, we visualized and interrogated the SIMBA embeddings of 1) cells; 2) cells and genes; and
3) cells together with genes, TF motifs, and peaks. **Figure 4c** shows the UMAP visualization of
the SIMBA embeddings of cells and informative features selected based on SIMBA metric plots.
The UMAP visualization of SIMBA embeddings of cells and the full set of features was also
performed (**Supplementary Fig. 6a**). However, we observed that with the SIMBA embeddings of
cells and the informative features selected by SIMBA, UMAP plots show more visible cell
subgroups without obstruction by entities that may dominate the space given their larger
number (e.g., peaks). SIMBA embeddings of cells were able to reveal the three fate decisions
from transit-amplifying cells (TACs), including inner root sheath (IRS), medulla, and
cuticle/cortex. SIMBA embeddings also uncovered important genes and regulatory factors
along the hair follicle differentiation trajectories. For example, the marker genes *Krt71*, *Krt31*,
and *Foxq1* were embedded into their corresponding cell types: IRS, cuticle/cortex, and medulla,
respectively. The Lef1 motif was embedded into the beginning of medulla and cuticle/cortex
lineages while the Hoxc13 motif was embedded into the late stage of cuticle/cortex
differentiation. Peaks near the *Lef1* and *Hoxc13* loci were also embedded into the nearby
regions of these genes and motifs, as expected. To show the robustness of SIMBA embedding,
we also performed the single-modality analyses within the SHARE-seq dataset, separating the
scRNA-seq and scATAC-seq components. With the consistent embedding results of cells and
features as in multimodal analysis, we further demonstrated that SIMBA embedding procedure
is robust to the type and number of features encoded in the input graph (**Supplementary Fig.**
6b,**6c**). Each marker gene was further validated using the UMAP plots with cells colored by gene
expression as well as using the SIMBA barcode plots. The two aforementioned TF motifs and
their respective peak sets were also validated visually using SIMBA barcode plots. As expected,
in SIMBA barcode plots these marker features clearly showed an imbalanced distribution with
much higher probabilities in the correct cell types (**Supplementary Fig. 7a-d**).

Further, we demonstrated that the SIMBA co-embedding space of cells and features provides
the potential to identify master regulators of differentiation and infer their target regulatory
genes. To define a master regulator *a priori*, we postulate that both its TF motif and TF gene
should be cell-type specific and given that active gene regulation involves the expression of TF
and accessibility of its binding sites, TF motif and TF gene should be embedded closely in the
shared latent space. Extending this logic to identify putative master regulators, we assessed the
cell-type-specificity of TF motifs and genes based on SIMBA metrics and ranked all potential
master regulators based on the distance between the TF motif and the respective TF gene in
the shared SIMBA embedding space (**Methods**). SIMBA successfully identified previously
described master regulators such as Lef1, Gata6, Nfatc1, and Hoxc13. as the top master
regulators related to lineage commitment in mouse skin (**Fig. 4d, Supplementary Table 2**). To
infer the target genes of a given master regulator, we postulate that in the shared SIMBA
embedding space, 1) the target gene is close to both the TF motif and the TF gene; 2) the
accessible regions (peaks) near the target gene loci must be close to both the TF motif and the

target TF gene. Resting on these assumptions of *cis*-regulatory dynamics, the inference of target
genes was performed by calculating the distance between target gene candidates and the
respective TF motif and gene. In addition, nearby peaks around the target gene's locus and the
presence of TF motif in these nearby peaks are also considered (**Fig. 4e, Methods**). The top 30
target genes of TF Lef1 and TF Hoxc13 inferred by SIMBA are shown respectively (**Fig. 4f,**
Supplementary Fig. 7e). The full list of ranked target genes is provided in **Supplementary Table**
3. Notably, our approach recovered targets genes that were also reported in the original
study[22]. For example, genes *Lef1*, *Jag1*, *Hoxc13*, *Gtf2ird1* are regulated by the TF Lef1, while
genes *Cybrd1*, *Hoxc13*, *St14* are regulated by the TF Hoxc13.

In addition to SHARE-seq, we also applied SIMBA to another two dual-omics datasets, the
mouse cerebral cortex dataset profiled by SNARE-seq[21] (**Supplementary Fig. 8**) and the
multiome PBMCs dataset from 10x Genomics (**Supplementary Fig. 9**). By validating the
embeddings of cells and features with given cell type labels (**Supplementary Fig. 8a and Fig.9a**),
marker genes from the original study (**Supplementary Fig. 8a,b,d and Fig.9a,b,d**), and
differentially accessible chromatin regions (**Supplementary Fig. 8c and Fig.9c**), we further
demonstrate the suitability of SIMBA for multimodal analysis.

468
469   **SIMBA enables simultaneous batch correction and clustering-free marker gene detection**
470
Efforts to collect data from single cells has grown to the level of consortia that span multiple
institutions with the hopes of finely mapping and characterizing specific tissues. This has
brought with it an increased demand for analysis methods that are capable of negating
technical covariates inherent to multi-batch data collection, including experimental replicate
identity, sample preparation, and sequencing platform. Batch correction that removes the
effects of technical covariation while preserving true biological signals is required prior to
downstream analysis [25, 26]. Unlike existing methods that require clustering and the merging of
marker genes detected in the original gene expression space of each batch, SIMBA enables
simultaneous batch effect removal and cell-type-specific marker gene detection in the same
integrated space without clustering.
481
We first demonstrate that SIMBA readily corrects batch effects and produces joint embeddings
of cells and genes across multiple datasets with different sequencing platforms and cell type
compositions. Thus, while previous methods primarily rely on specialized tools for batch
correction before downstream clustering and marker detection, SIMBA works as a stand-alone
package obviating the need for prior input data correction when applied to multi-batch scRNA-
seq dataset. In SIMBA, batch correction is accomplished by encoding multiple scRNA-seq
datasets into a single graph (**Fig. 5a**). Cells in different batches are linked to genes as in the
previously described scRNA-seq graph construction. Here, the gene nodes are shared between
the cell nodes of different batches. In addition to the experimentally measured edges, batch
correction is further enhanced through computationally inferred edges drawn between similar
cell nodes across datasets using a truncated randomized singular value decomposition (SVD)-
based procedure. SIMBA then generates the embeddings of all nodes including cells of each

batch and genes from the resulting graph, which may then be visualized using tools such as
UMAP as in the analyses presented in the sections above (**Methods**).
496

497
498

Figure 5. Batch correction analysis of scRNA-seq data using SIMBA. **(a)** SIMBA graph construction
and embedding in batch correction analysis. Overview of SIMBA's approach to batch correction
across scRNA-seq datasets. Distinct shapes indicate the type of entity (cell or gene). Colors
distinguish batches or cell types. **(b)** UMAP visualization of the scRNA-seq human pancreas
dataset with five batches of different studies before and after batch correction. Cells are colored
by scRNA-seq data source and cell type respectively. Top: UMAP visualization before batch
correction; Bottom: UMAP visualization after batch correction with SIMBA; **(c)** UMAP
visualization of SIMBA embeddings of cells and genes, with batch effect removed and known
marker genes highlighted.

508
SIMBA was applied to a mouse atlas dataset composed of two batches and a human pancreas
dataset that spans several batches used in a recent benchmark study[25] (**Supplementary Table**
1). The mouse atlas dataset contains two scRNA-seq datasets with shared cell types from
different sequencing platform. The human pancreas dataset contains five samples pooled from
five sources using four different sequencing techniques, in which not all cell types are shared
across each sample. For both datasets, SIMBA successfully corrected the batch effects where

the resulting embeddings of cells are clustered by the cell types while each batch is mixed
evenly, indicating the preservation of biological signal and the elimination of confounding
technical covariates (**Fig. 5b, Supplementary Fig. 12b**). It is important to note that the mouse
atlas dataset was collected from nine different organ systems, so there exists some expected
heterogeneity within the same cell type labels. Conversely, the human pancreas datasets are
curated from a single organ and SIMBA sufficiently separated cell types into transcriptionally-
distinct homogeneous cell clusters (**Fig. 5b**).

In addition to batch effect removal, SIMBA also simultaneously identifies cell-type-specific
marker genes (**Fig. 5c**). Having effectively eliminated differences between cells due to technical
covariates, marker genes are discoverable across multiple samples by querying according to
cells of each cell type within the batch-corrected SIMBA embedding. As shown in the SIMBA co-
embedding of cells and variable genes, known marker genes were correctly placed within each
cell type while non-marker genes were embedded away from any cell type (**Supplementary Fig.**
10, 11). The resulting marker genes recapitulated the clustering-based differential expression
(DE) analysis results for each dataset[27-32] (e.g. *Cdh5*, *Tie1*, *Myct1* for endothelial cell and *C1qc*,
Fcgr1 for macrophage, *S100a8*, *Trem3* for Neutrophil in the mouse atlas dataset and *KIF12* for
alpha cell and *KRT19* for ductal cell in the human pancreas dataset) and are shown to be
expressed specifically in the queried cell types (**Supplementary Fig. 10, 11**). This distinguishes
SIMBA from other batch correction methods, in which clustering is performed first in the batch-
corrected space and then marker genes of each batch are identified through DE analysis in the
original, uncorrected space of the corresponding batch.

While SIMBA is a generalizable graph embedding method capable of performing multiple tasks
outside of batch correction, we still evaluated its performance for this task with methods that
were specifically designed for batch correction. We considered three widely adopted batch
correction methods that demonstrated top-tier performance based on a recent benchmark
study[25]: Seurat3, LIGER and Harmony. Our results show that SIMBA achieved comparable batch
correction performance both qualitatively and quantitatively (**Supplementary Note 2,**
Supplementary Figure 12).

SIMBA enables simultaneous multi-omics integration and clustering-free multi-type marker
feature detection

Single-cell assays are now capable of measuring a broad range of cellular modalities including
mRNA, chromatin accessibility, DNA methylation and cell-surface proteins. Thus, data is being
generated that describes cells by varying features sets, which has motivated the need for
methods that leverage these features to perform multi-omics integration such that a more
comprehensive description of cell state may be learned. This is different from multi-modal
analysis because the correspondence between individual cells is unknown. We demonstrate
that SIMBA enables simultaneous multi-omics integration and clustering-free detection of
distinct marker features, specifically as it is applied to datasets comprised of scRNA-seq and
scATAC-seq.
558

SIMBA accomplishes this integration by first building one graph for scRNA-seq data and another
graph for scATAC-seq data independently as described in previous sections (**Fig. 6a**). To connect
these two graphs, SIMBA then calculates gene activity scores by summarizing accessible regions
from scATAC-seq data and then infers edges between cells of different assays based on their
shared gene expression modules as previously described in the batch correction section. Finally,
SIMBA embeds the graph of cells, genes, and peaks into a common, low-dimensional space. The
SIMBA embeddings of these multi-omics entities can be visualized using UMAP or similar
visualization tools (**Methods**).
567

568
Figure 6. Multi-omics integration of scRNA-seq + scATAC-seq data using SIMBA. **(a)** SIMBA graph
construction and embedding in multi-omics integration. Overview of SIMBA's approach to data
integration across scRNA-seq and scATAC-seq. Distinct shapes indicate the type of entity (cell,
gene, or peak). Colors distinguish batches or cell types. **(b)** UMAP visualization of the integrated
scRNA-seq and scATAC-seq data manually created from the 10x human PBMCs dataset before
and after data integration. Cells are colored by single-cell modality and cell type respectively. Top:
UMAP visualization before integration; Bottom: UMAP visualization after integration with SIMBA.
(c) UMAP visualization of SIMBA embeddings of cells, genes, and peaks with two cell modalities
integrated and known marker genes highlighted.
578

To facilitate the evaluation of data integration performance, we created datasets with ground-
truth labels by manually splitting the dual-omics datasets into two single-modality datasets (i.e.,
scRNA-seq and scATAC-seq), in which we know the true matching between cells across the two
modalities. We then applied SIMBA to the integration analysis of two case studies where
scRNA-seq and scATAC-seq datasets are generated from the SHARE-seq mouse skin dataset and
the 10x Genomics multiome human PBMCs dataset, respectively (**Supplementary Table 1**). We
observed that SIMBA was able to preserve cellular heterogeneity while evenly mixing the two
modalities (**Fig. 6b, Supplementary Fig. 15b**).

Importantly, SIMBA simultaneously identified marker genes and peaks at single-cell resolution.
In the co-embedding space, we observed that the neighbor genes of cells (highlighted in UMAP
plots), are each exclusively expressed in their corresponding cell types (**Supplementary Figs.
13a-e, 14a-c,e**). For example, in the SHARE-seq mouse skin dataset, *Foxq1* and *Shh* are located
within medulla and TAC-2, respectively; in the 10x PBMCs dataset, *PAPSS2* and *KCNMA1*, which
are the marker genes of blood monocytes, are embedded close to each other. Similarly, we
observed that the neighbor peaks of cells show a clear cell-type-specific accessibility pattern
that is robust to the cluster size of a given cell type (**Supplementary Figs. 13f and 14d).**

While SIMBA distinguishes itself from other multi-omics integration methods by generating
joint embeddings of cells and features and thus enabling simultaneous multi-omics integration
and marker feature detection, we still compared SIMBA with two leading and widely-adopted
single-cell multi-omics integration methods, Seurat3 and LIGER, based on their ability to
integrate single-cell modalities while persevering cellular heterogeneity (**Supplementary Note
3**). We observed that SIMBA achieved overall best performance on the mouse skin SHARE-seq
dataset and 10x PBMCs multiome dataset.

Discussion

Multimodal measurements of individual cells offer new and unexplored opportunities to
investigate complex interactions between components involved in establishing transcriptional
programs. Despite the exciting potential for discovery associated with these datasets, methods
to delineate cell states and cellular processes using multiple genomic features across modalities
remains under development.

As presented in this manuscript, SIMBA models cells and measured features as nodes encoded
in a graph and employs a scalable and efficient graph embedding procedure to embed nodes of
cells and features into a shared latent space. We demonstrate that direct graph representations
of single-cell data capture not only the relations between cells and the quantified features of
the experiment (e.g., gene expression or chromatin accessibility) but also the hierarchical
relations between features. An example of such a hierarchical relation is the coordinate-level
description of an ATAC-seq peak and the corresponding TF motifs and/or *k*-mer sequences
contained within that region. In the resulting joint embedding, proximity-based queries can be
performed to discover cell-type-specific regulatory mechanisms and the respective features
integral to such mechanisms. Therefore, SIMBA enables simultaneous learning of cellular

heterogeneity and cell-type-specific multimodal features, and complements the current gene
regulatory network analyses. SIMBA also circumvents the typical workflow led by cell clustering
and followed by differential feature detection, and thus avoids relying on user-defined
clustering resolution that may lead to artifactual discovery or false negative results.
627
SIMBA has been extensively benchmarked across single-cell modalities and tasks and obtained
comparable or better performance compared to current state-of-the-art methods developed
for the respective task. These results suggest a wide applicability of SIMBA's graph-based
framework and therefore obviates the need to optimally combine multiple analysis tools. This
contrasts with task-specific methods, which have been specially developed for a given data type
and modality.
634
A foreseeable extension of SIMBA would include encoding increasingly complex measurements
of single-cells such as spatial transcriptomics wherein transcriptomic and (real) proximity data
should be considered [33]. We also envision extending this framework to chromatin looping data
wherein the interaction between DNA segments can be encoded to represent how regulatory
regions are linked to genes. Another potential extension of SIMBA could consider single-cell
lineage-tracing datasets[34] wherein both cellular lineage information and gene expression
measurements are captured and can be potentially encoded into a SIMBA graph to represent
their longitudinal relations. In general, we are interested in the further incorporation of
external information and hierarchical relationships between features in the graph. While
extensions to different experimental designs is relatively straightforward, we note that the
interpretation of the output embedding in each case may vary as a function of the input graph
construction and training process thus requiring some level of domain-specific expertise.
647
Graph embedding methods hold significant promise for the analysis of biological data. Previous
applications of graph embedding include functional annotation of genes [35], transcription factor
binding to DNA motifs [36] and more recent single-cell RNA-seq analyses [37, 38]. The graph encoding
and embedding procedures we have outlined may be potentially improved and extended to
better represent biological entities and capture their respective relations. We envision
employing continuous edge weights as well as the consideration of both node attributes and
edge information to determine the proximity of node pairs [39]. Considering embedding distances
as a function of the node attribute may better equip our approach to extend to time series data
as well as to better align cells stratified by positions on a developmental axis. In addition, to
improve interpretability it may be possible to provide a metric of uncertainty alongside the
resulting embedding and generalization to the unseen nodes[40]. This could enable the mapping
of new cells in the same manifold and a straightforward and automatic cell type annotation.
660
The ability to generate high-quality, complex datasets that capture and quantify multiple
classes of molecules (omics) will undoubtedly continue to increase. Technological advances
have already outpaced our ability to gain integrative insights from such data, highlighting a
need for methods that break through previous limitations as well as extend easily to new tasks.
We believe SIMBA satisfies these conditions for such a comprehensive yet accessible method
for exploring cellular heterogeneity and investigating the regulatory mechanisms that drive

cellular diversity while laying a groundwork for the development of new non-cluster-centric
single cell omics computational methods.
669
670

Methods

Single-cell data preprocessing

a. Single-cell RNA-seq
Genes expressed in fewer than three cells were filtered.   Raw counts were library size-
normalized and subsequently log-transformed. Optionally, variable gene selection [12] (a
python version is implemented in SIMBA that is inspired by Scanpy[2]) may be performed
to remove non-informative genes and accelerate the training procedure. Notable
differences in the resulting cell embeddings were not observed upon limiting feature
input to those identified by variable gene selection but SIMBA embeddings of non-
variable genes will not be generated as they are not encoded in the graph.

b. Single-cell ATAC-seq
Peaks present in fewer than three cells were filtered. Optionally, we implemented a
scalable truncated-SVD-based procedure to select variable peaks as a preliminary step
to additionally filter non-informative peaks and accelerate the training procedure. First
the top k principal components (PCs) were selected, with k chosen based on the elbow
plot of variance ratio. Then for each of the top k PCs, peaks were automatically selected
based on the loadings using a knee point detection algorithm implemented by 'kneed'[41].
Finally, peaks selected for each PC were combined and denoted as "variable peaks".
Similar to the observation made with scRNA-seq data, the optional step of variable peak
selection has a negligible effect on the resulting cell embedding. Despite this minimal
impact on the resulting embedding, this feature selection step imparts a significant
practical advantage in reducing training procedure time.

k-mer and motif scanning was performed using packages 'Biostrings' and 'motifmatchr'
with JASPAR2020[42]. Included in the implementation of SIMBA is a convenient R
command line script "scan_for_kmers_motifs.R" , which will convert a list of peaks
(formatted in a bed file) to a sparse peaks-by-k-mers/motifs matrix, which is stored as
an hdf5-formated file.

Graph construction (five scenarios)

i.   Single-cell RNA-seq analysis
The distribution of non-zero values in the normalized gene expression matrix was first
approximated using a k-means clustering-based procedure. First, the continuous non-
zero values were binned into n intervals (by default n=5). Bin widths were defined using
1-dimensional k-means clustering wherein the values in each bin are assigned to the
same cluster center. The continuous matrix is then converted into a discrete matrix

wherein1, …, n are used to denote n levels of gene expression. Zero values are retained
in this matrix. Then the graph was constructed by encoding two types of entities, cells
and genes, as nodes and relations with n different weights between them, i.e., n levels
of gene expression, as edges. These n relation weights range from 1.0 to 5.0 with a step
size of $5/n$ denoting gene expression levels (lowest: 1.0, highest: 5.0), such that edges
corresponding to high expression levels affect embeddings more strongly than those
with intermediate or low expression levels. This discretization is implemented in the
SIMBA package using the function, "si.tl.discretize()".

ii.   Single-cell ATAC-seq analysis
Peak-by-cell matrices were binarized, with "1" indicating at least one read within a peak
and "0" otherwise. The graph was constructed by encoding two types of entities, cells
and peaks, as nodes and the relation between them, denoting the presence of a given
peak in a cell, as edges. The single relation type was assigned with a weight of 1.0. When
the DNA sequence features were available, they were encoded into the graph using k-
mer and motif sequence entities as nodes. This was performed by first binarizing the
peak-by-k-mer/motif matrix then constructing an extension to the original peak/cell
graph using the peaks, k-mers, and motifs as nodes and the presence of these entities
within peaks as edges between these additional nodes and the peak nodes. The relation
between k-mers and peaks was assigned a weight of 0.02 while the relation between TF
motifs was assigned a weight of 0.2. Of note, k-mers and motifs may be used
independently of each other as node inputs to the graph, depending on the specific
analysis task.

iii.   Multimodal analysis
Combination of the above outlined strategies for graph construction of scRNA-seq and
scATAC-seq data was used to construct a multi-omics graph.

iv.   Batch correction
A graph for each batch was constructed as described in i). Edges between cells of
different batches were inferred through a procedure based on truncated randomized
singular value decomposition (SVD) to link disjoint graphs of different batches. More
specifically, in the case of scRNA-seq data, consider two gene expression matrices
$X1_{n_1 \times m}$ and $X2_{n_2 \times m}$, where n_1, n_2 denotes the number of cells and m denotes the
number of the shared features, i.e., variable genes, between datasets. The matrix
$X_{n_1 \times n_2}$ was then computed by multiplying $X1$ and $X2$:

$$X = X1 \times X2^T$$

Truncated randomized SVD was subsequently performed on X:

$$X \approx U \times \Sigma \times V^T$$

where U is an $n_1 \times d$ matrix, Σ is an $d \times d$ matrix, and V is an $n_2 \times d$ matrix (by
default $d = 20$).

Both U and V were further $L2$ normalized. For each cell in U, we searched for k
nearest neighbors in V and vice versa (by default, $k = 20$). Eventually, only the mutual
nearest neighbors between U and V were retained as inferred edges between cells
(represented as dashed lines in **Fig. 5a**). The procedure of inferring edges between
cells of different batches is implemented in the function "si.tl.infer_edges()" in the
SIMBA package.

For multiple batches, SIMBA can flexibly infer edges between any pair of datasets. In
practice, however edges are inferred between the largest dataset(s) or the dataset(s)
containing the most complete set of expected cell types and other datasets.

v. Multi-omics integration
scRNA-seq and scATAC-seq graphs were constructed following steps i) and ii),
respectively. To infer the edges between cells of scRNA-seq and scATAC-seq, gene
activity scores were first calculated for scATAC-seq data[3]. More specifically, for each
gene, peaks within 100kb upstream and downstream of the TSS were considered.
Peaks overlapping gene body region or within 5kb upstream of gene bodies were
given the weight of 1.0. Otherwise, peaks were weighted based on their distances to
TSS using the exponential decay function: $e^{\frac{-distance}{5000}}$. Subsequently, the gene score of
each gene was computed as a weighted sum of the considered peaks. These gene
scores were then scaled to respective gene size. These steps are implemented by the
function "si.tl.gene_scores()" in SIMBA. For user convenience, the SIMBA package
curates the gene annotations of several commonly used reference genomes, including
hg19, hg38, mm9, and mm10. Once gene scores were obtained, the same procedure
described in iv) was performed to infer edges between cells profiled by scRNA-seq and
scATAC-seq using the function, "si.tl.infer_edges()" in SIMBA.

The procedure of generating constructed graphs is implemented in the function,
"si.tl.gen_graph()" in the SIMBA package.

Graph Embeddings with Type Constraints

Following the construction of a multi-relational graph between biological entities, we
adapted graph embedding techniques from the knowledge graph and recommendation
systems literature to construct unsupervised representations for these entities.

We provide as input a directed graph $G = (V, E)$, where V is a set of entities (vertices)
and E is a set of edges, with a generic edge $e = (u, v)$ between a source entity u and
destination entity v. We further assume that each entity has a distinct known type (e.g.,
cell, peak, etc.).

Graph embedding methods learn a D-dimensional embedding vector for each $v \in V$ by optimizing a link prediction objective via stochastic gradient descent, with D=50 used for our experiments. We will denote the full embedding matrix as $\theta \in R^{|V| \times D}$ and the embedding for an entity v as θ_v.

For an edge $e = (u, v)$, we denote $s_e = \theta_u * \theta_v$ as the score for e, and optimize a multi-class log loss

$$\mathcal{L} = -log \frac{\exp(s_e)}{\sum_{e' \in \mathcal{N}} \exp(s'_e)}$$

Where $\mathcal{N}$ is a set of "negative sampled" candidate edges generated by corrupting e [43]. This log loss objective attempts to maximize the score for all $(u, v) \in E$ and minimize it for $(u, v) \notin E$.

Negative samples are constructed by replacing either the source or target entity in the target edge $e = (u, v)$ with a randomly sampled entity. However, in graphs like ours where only edges between certain entity types are possible, previous work has shown that it is beneficial to optimize the loss only over candidate edges that satisfy the type constraints[44]. Thus, for e.g., a cell-peak edge we only sample negative candidates between cell and peak entities. This modification is crucial in our setting since most randomly selected edges will be of invalid type (e.g., peak-peak), forcing the embeddings to primarily be optimized for irrelevant tasks (e.g., having low dot product between every pair of peaks).

Furthermore, it has been frequently observed that in graphs with wide distribution of node degrees, it is advantageous to sample negatives proportional to some function of the node degree to produce more informative embeddings that don't merely capture the degree distribution [13, 45]. For each graph edge in the dataset encountered in a training batch, we produce 100 negatives by corrupting the edge with a source or destination sampled uniformly from the nodes with the correct types for this relation and 100 by corrupting the edge with a source or destination node sampled with probability proportional to its degree[13].

As with many ML methods, graph embeddings are prone to overfitting in a low-data regime (i.e., low ratio of edges to parameters). We observed overfitting measurable as a gap between training and validation loss on the link prediction task, which we addressed with $L2$ regularization on the embeddings θ,

$$\mathcal{L}_{reg} = \mathcal{L} + \lambda \sum_{u \in N} \sum_{d=1}^{D} \theta_{ud}^2.$$

with $\lambda = wd * wd\_interval$. For weight decay parameter (wd), by default it is calculated
automatically as $\frac{C}{N_e}$, where N_e is the training sample size (i.e., the total number of edges)
and C is a constant. For weight decay interval ($wd\_interval$), we set it to 50 for all
experiments.

We use the PyTorch-BigGraph framework, which provides efficient computation of
multi-relation graph embeddings over multiple entity types and can scale to graphs with
millions or billions of entities[13].

The resulting graph embeddings have two desirable properties that we will take
advantage of:
1.   First-order similarity: for two entity types T_1, T_2 with a relation between them,
edges with high likelihood should have higher dot product; specifically, for any $u \in T_1$,
the predicted probability distribution over edges to T_2 originating from u is
approximated as $\frac{e^{x_u * x_v}}{\sum_{v\prime \in T_2} e^{x_u * x_{v\prime}}}$.
2.   Second-order similarity: within a single entity type, entities that have 'similar
contexts', i.e., a similar distribution of edge probabilities, should have similar
embeddings. Thus, the embeddings of each entity type provide a low-rank latent space
that encodes the similarity of those entities' edge distributions.

Evaluation of the model during training

During the PyTorch-BigGraph training procedure, a small percent of edges is held out
(by default, the evaluation fraction is set to 5%) to monitor overfitting and evaluate the
final model. Five metrics are computed on the reserved set of edges, including mean
reciprocal rank (MRR, the average of the reciprocal of the ranks of all positives), R1 (the
fraction of positives that rank better than all their negatives, i.e., have a rank of 1), R10
(the fraction of positives that rank in the top 10 among their negatives), R50 (the
fraction of positives that rank in the top 50 among their negatives), and AUC (Area
Under the Curve). By default, we show MRR along with training loss and validation loss
while other metrics are also available in SIMBA package (**Supplementary Fig. 1a**). The
learning curves for validation loss and these metrics can be used to determine when
training has completed. The relative values of training and validation loss along with
these evaluation metrics can be used to identify issues with training (underfitting vs
overfitting) and tune the hyperparameters weight decay, embedding dimension, and
number of training epochs appropriately. For example, in **Supplementary Figure 1a**
training can be stopped once the validation loss plateaus. However, for most datasets
we find that the default parameters do not need tuning.

Softmax transformation

PyTorch-BigGraph training provides initial embeddings of all entities (nodes). However,
entities of different types (e.g., cells vs peaks, cells of different batches or modalities)

have different edge distributions and thus may lie on different manifolds of the latent
space. To make the embeddings of entities of different types comparable, we transform
the embeddings of features with the Softmax function by utilizing the first-order
similarity between cells (reference) and features (query). In the case of batch correction
or multi-omics integration, the Softmax transformation is also performed based on the
first-order similarity between cells of different batches or modalities.
886
887   Given the initial embeddings of cells (reference) $(v_{c_1}, \dots, v_{c_n})$ and features $(v_{f_1}, \dots, v_{d_m})$,
888   the model-estimated probability of an edge (c_i, f_j) obeys
889

$$P\left(v_{c_i,f_j}\right) \propto \exp\left(v_{c_i} \cdot v_{f_j}\right)$$

892   Therefore, if a random edge was sampled from feature f_j to a cell, the model would
893   estimate the distribution over such edges as
894

$$p_{c_i,f_j} = \frac{\exp\left(v_{c_i} \cdot v_{f_j}\right)}{\sum_{k=1}^{n} \exp\left(v_{c_k} \cdot v_{f_j}\right)}$$

897   i.e., the Softmax weights between all cells $\{c_i\}$ and the feature f_j. We can then compute
898   new embeddings for features as a linear combination of the cell embeddings weighted
899   by the edge probabilities raised to some power.
900

$$\hat{v}_{f_j} = \frac{\sum_{i=1}^{n} p_{c_i,f_j}^{T^{-1}} v_{c_i}}{\sum_{i=1}^{n} p_{c_i,f_j}^{T^{-1}}}$$

903   T is a temperature hyperparameter that controls the sharpness of the weighting over
904   cells. At $T = 1$, the cell embeddings are weighted by their estimated edge probabilities;
905   at $T \rightarrow 0$, each feature embedding is assigned the cell embedding of its nearest
906   neighbor; at $T \rightarrow \infty$, it becomes a discrete uniform distribution, and each query
907   becomes the average of reference embeddings. We set $T = 0.5$ for all the analyses.
908
909   These steps are implemented in the function "si.tl.embed()" in the SIMBA package.
910
911 **Metrics to assess cell-type specificity**
912
Four metrics are proposed to assess the cell type specificity of each feature from
different aspects, including max value (a higher value indicates higher cell-type
specificity), Gini index (a higher value indicates higher cell-type specificity), standard
deviation (a higher value indicates higher cell-type specificity), and entropy (a lower
value indicates higher cell-type specificity). We observe these four metrics generally give
consistent results. For SIMBA metric plot, by default, Gini index is plotted against max
value. For feature f_j :

The max value is defined as the average normalized similarity of top k cells (by default, k=50). The similarity normalization function is defined as:

$$norm(x_i) = x_i - log\frac{\sum_{j=1}^{n} \exp(x_j)}{n}$$

Where $i = 1, \ldots, n$. n is the number of cells and x_i represents the dot product of $\hat{v}_{f_j}$ and the embedding of cell i.

The max value is computed as:

$$\max(f_j) = \frac{\sum_{i=1}^{k} norm(x_i)}{k}$$

The Gini index is computed as:

$$\text{gini}(f_j) = \frac{\sum_{i=1}^{n}(2i - n - 1) * p_{c_i, f_j}}{n \sum_{i=1}^{n} p_{c_i, f_j}}$$

The standard deviation is computed as:

$$\text{std}(f_j) = \sqrt{\frac{1}{n-1}\sum_{i=1}^{n}(p_{c_i, f_j} - \mu)^2}$$

Where $\mu = \frac{1}{n}\sum_{i=1}^{n} p_{c_i, f_j}$.

Entropy is computed as:

$$\text{entropy}(f_j) = -\sum_{i=1}^{n} p_{c_i, f_j} log(p_{c_i, f_j})$$

Identification of master regulators

To identify master regulators, we take into consideration both the cell type specificity of each pair of TF motif and TF gene and the distance between them. More specifically, for each TF motif, first its distances (Euclidean distance by default) to all the genes are calculated in the SIMBA embedding space. Then the rank of this TF gene among all these genes is computed. In addition, we also assess the cell type specificity of this pair of TF motif and TF gene based on SIMBA metrics (by default, max value and Gini index are used). The same procedure is performed for all TFs. Finally, we identify master regulators by filtering out TFs with low cell-type specificity and scoring them based on

955   TF gene rank. This procedure is implemented in the function
956   "st.tl.find_master_regulators()" in SIMBA package.
957
958
Identification of TF target genes

Given a master regulator, its target genes are identified by comparing the locations of
the TF gene, TF motif, and the peaks near the genomic loci of candidate target genes in
the SIMBA co-embedding space (**Fig. 4e**). More specifically we first search for k nearest
neighbor genes around the motif (TF motif) and the gene (TF gene) of this master
regulator, respectively (k = 200 by default). The union of these neighbor genes is the
initial set of candidate target genes. These genes are then filtered based on the criterion
that open regions (peaks) within 100kb upstream and downstream of the TSS of a
putative target gene must contain the TF motif.

Next, for each candidate target gene, we compute four types of distances in SIMBA
embedding space: distances between the embeddings of 1) the candidate target gene
and TF gene; 2) the candidate target gene and TF motif; 3) peaks near the genomic locus
of the candidate target gene and TF motif; 4) peaks near the genomic locus of the
candidate target gene and the candidate gene. All the distances (Euclidean distances by
default) are converted to ranks out of all genes or all peaks to make the distances
comparable across different master regulators.

The final list of target genes is decided using the calculated ranks based on two criteria:
1) at least one of the nearest peaks to TF gene or TF motif is within a predetermined
range (top 1,000 by default); 2) the average rank of the candidate target gene is within a
predetermined range (top 5,000 by default). This procedure is implemented in the
function "st.tl. find_target_genes ()" in SIMBA.

Benchmarking scATAC-seq computational methods
981
To compare SIMBA to other scATAC-seq computational methods including SnapATAC [4],
Cusanovich2018[46], and cisTopic[47], we employed the previously developed benchmarking
framework from Chen et al[14](**Supplementary Table 1**). This framework evaluates
different methods based on their ability to distinguish cell types. We applied three
clustering algorithms: k-means clustering, hierarchical clustering, and Louvain on the
feature matrix derived from each method.
988
For datasets with ground-truth (FACS-sorted labels or known tissue labels), including
simulated bone marrow data, Buenrostro 2018, and sci-ATAC-seq subset, three metrics
including Adjusted Rand Index (ARI), Adjusted Mutual Information (AMI), and
Homogeneity are applied to evaluate the performance. ARI measures the similarity
between two clusters, comparing all pairs of samples assigned to matching or different
clusters in the predicted clustering solution vs the true cluster/cell type label. AMI

describes an observed frequency of co-occurrence compared to an expected frequency of co-occurrence between two variables, informing the mutual dependence or strength of association of these two variables. Homogeneity measures whether a clustering algorithm preserves cluster assignments towards samples that belong to a single class. A higher metric value indicates a better clustering solution.

For 10x PBMCs dataset with no ground truth, the Residual Average Gini Index (RAGI) proposed in the benchmarking study[14] is used as the clustering evaluation metric. RAGI measures the relative exclusivity of marker genes to their corresponding clusters in comparison to housekeeping genes, which should demonstrate low specificity to any given cluster. In brief, the mean Gini Index is computed for both marker genes and housekeeping genes. The difference between the means is computed to obtain the average residual specificity (i.e., RAGI) of a clustering solution with respect to marker genes. A higher RAGI indicates a better separation of biologically distinct clusters.

Benchmarking single-cell batch correction methods

The batch correction performance of SIMBA was compared to Seurat3[12], LIGER[11] and Harmony[10] in two benchmark datasets: the mouse atlas dataset and the human pancreas dataset (**Supplementary Table 1**). For Seurat3, LIGER and Harmony, the batch correction was done with the same parameters used in a previous benchmark study[25].

To evaluate the batch integration performance, average Silhouette width (ASW), adjusted Rand index (ARI), and local inverse Simpson's index (LISI)[10] were calculated for the batches and cell types using the Euclidean distance as described in a previous benchmark[25]. To make a fair evaluation, only the cell types that are present in all batches were considered. We used the same number of dimensions (50) for these methods and all other parameters were set as in the benchmark.

Average Silhouette width (ASW)

Average Silhouette width is the mean value of Silhouette scores calculated from each cell. Silhouette width measures the relative closeness of cells with the same label compared to the cells with the different label and ranges from -1 to +1. Silhouette score for a data point with a label is calculated as

$$s(i) = \frac{b(i) - a(i)}{\max\{a(i), b(i)\}}$$

where $a(i)$ is the distance to the closest point with the same label, and $b(i)$ is the distance to the closest point with different labels. A high Silhouette score means the point is located more closely with the same label, where a low Silhouette score closer to -1 means the point is located closer with different labels than that of itself. The ideal batch correction result will give a low ASW score for batch labels as the point is well

mixed with other batches and a high ASW score for the cell type labels as the cells of the same cell type should cluster together after the batch correction. The final score is calculated as the median ASW scores from 20 subsets of randomly sampled 80% cells.

Average Rand Index (ARI)

To evaluate the cell type purity, the true cell type labels and the k-means clustering solution were used to calculate the cell type ARI. To evaluate the batch correction performance, the true batch labels and the k-means clustering solution were used to calculate the batch ARI. The final ARI was calculated as the median ARI scores of 20 subsets comprised of randomly sampled 80% cells for batches and cell types, respectively. A superior batch correction will have a high cell type ARI (high agreement between the clustering solution and the true cell type labels), and a low batch ARI (the clustering solution is not mainly driven by batches and clusters contain cells with well-mixed batch labels).

Local Inverse Simpson's Index (LISI)

Local Inverse Simpson's Index (LISI) [10] measures the local batch and cell type mixing. For each data point, it considers the Gaussian kernel weighted distribution of labels in its neighborhood with a perplexity argument. We set perplexity to 50 40 as in the previous benchmark study. Using the weighted neighborhood label distribution, the inverse Simpson's index is calculated as $\frac{1}{\sum_l p(l)}$ where l is the batch or cell type labels and $p(l)$ is the probability of each label in the local neighborhood obtained with the kernel. For each cell, the LISI is the expected number of cells to be sampled locally before a cell of the same label is sampled. A perfect batch correction will have a cell type LISI (cLISI) of 1 and a batch LISI (integration LISI, iLISI) close to the number of batches. The final LISI score was calculated as the average LISI scores of all cells.

Further details are described in **Supplementary Note 2**.

Benchmarking single cell multi-omics integration methods

Two pairs of scRNA-seq and scATAC-seq datasets manually split from the dual-omics SHARE-seq mouse skin dataset and 10X PBMCs dataset respectively were used for the modality integration task. For Seurat3 and LIGER, the parameters and preprocessing were done as described in their documentations. However for the LIGER analysis of the SHARE-seq mouse skin dataset the parameter 'lambda' was set to 30 and the 'ref_dataset' was set to scATAC-seq to get a better alignment. For the Raw results, the activity matrix of scATAC-seq was constructed using Seurat3 and the first 20 PCs of the scRNA-seq count matrix and the activity matrix were used for the comparison. The integration results generated by each method were evaluated with four metrics—

1081	Anchoring distance, anchoring distance rank, Silhouette index, and cluster agreement—
1082	as described below.
1083	
1084	**Anchoring distance**
1085	The Anchoring distance was proposed in Dou et al., 2020[48] and is the normalized
1086	distance between the matched cells of two modalities (e.g. RNA and ATAC). Here we
1087	considered the Euclidean distance and normalized the distance by the mean of the
1088	distances calculated between random pairs of cells. The number of pairs randomly
1089	sampled was set to 10% of the total number of cells.
1090	
1091	**Anchoring distance rank**
1092	Given that the anchoring distance does not account for the local density of cells, we
1093	propose a new metric entitled *anchoring distance rank* (ADR). The ADR is based on the
1094	normalized rank of the distance between the matched cells of two modalities. For each
1095	cell x_{ij} with cell identity i and modality j, the distance between the cell and all the other
1096	cells of the other modality j', $d(x_{ij}, x_{kj'}), k = 1, ..., N$ is calculated, where N is the total
1097	number of cells. Then the rank of $r_i = d(x_{ij}, x_{ij'})$ within the calculated distances is
1098	normalized by the number of pairs $N - 1$ to obtain the final anchoring rank $m_i = \frac{r_i - 1}{N - 1}$.
1099	For each cell, an anchoring rank of 0 indicates an ideal modality integration performance
1100	as the matched cells are closest to each other in the embedding.
1101	
1102	**Silhouette index**
1103	The silhouette index was calculated as described in 10) based on the cluster assignment
1104	wherein each cluster consists of two cells, one cell from a scRNA-seq dataset and one
1105	cell from a scATAC-seq dataset.
1106	
1107	**Fraction in the same cluster**
1108	Fraction in the same cluster was calculated as the fraction of the matched cells from two
1109	modalities in the same cluster. The clusters of cells were generated using Louvain
1110	algorithm and the number of clusters is equal to the number of cell types in the dataset.
1111	
1112	Further details are described in **Supplementary Note 3**.
1113	
1114	

1115 Data availability:

1116

1117 All the datasets used in this study (eight scRNA-seq datasets, four scATAC-seq datasets, and
1118 three dual-omics datasets) are summarized in **Supplementary Table 1**. All these datasets are
1119 curated in the SIMBA package, and they can be easily downloaded and imported directly to
1120 reproduce the analyses presented in this manuscript.

1121

1122 Code availability:

1123

We provide a comprehensive Python package 'simba' available at https://anaconda.org/bioconda/simba and https://github.com/pinellolab/simba. All the proposed procedures are implemented in the "simba" package. 'simba' can be easily installed with conda "*conda install simba*". We also built a website (https://simba-bio.readthedocs.io), providing a detailed introduction of the 'simba' software and several SIMBA tutorials for different types of single-cell analyses presented in this manuscript.

Acknowledgements

The authors thank Ledell (Yu) Wu, Facebook, for the helpful discussions about Starspace; Dr.Sai Ma and Dr. Jason Buenrostro, Broad Institute of MIT and Harvard, for sharing the SHARE-seq data and metadata. The authors also acknowledge members of the Pinello lab, Mass General Hospital/Harvard Medical School, for helpful comments and feedback. This project has been made possible in part by grant number 2019-202669 from the Chan Zuckerberg Foundation to LP. LP is also partially supported by the National Human Genome Research Institute (NHGRI) Genomic Innovator Award (R35HG010717). M.E.V. is supported by the National Institutes of Health (NIH) under the Ruth L. Kirschstein National Research Service Award (1F31CA257625-01) from the National Cancer Institute (NCI).

Author contributions

H.C. and L.P. conceived this project. H.C. designed SIMBA, wrote the SIMBA package, and performed SIMBA analysis on all datasets. A.L. contributed to the adaption of PyTorch-BigGraph to single cell analysis. J.R. and H.C. performed the comparison analysis on batch correction and data integration. M.E.V. and H.C. performed the comparison analysis on scATAC-seq data. L.P. and A.L. provided guidance and supervised this project. All the authors wrote and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

References:

1. Satija, R., Farrell, J.A., Gennert, D., Schier, A.F. & Regev, A. Spatial reconstruction of single-cell gene expression data. *Nat Biotechnol* **33**, 495-502 (2015).
2. Wolf, F.A., Angerer, P. & Theis, F.J. SCANPY: large-scale single-cell gene expression data analysis. *Genome Biol* **19**, 15 (2018).
3. Granja, J.M. et al. ArchR is a scalable software package for integrative single-cell chromatin accessibility analysis. *Nat Genet* **53**, 403-411 (2021).
4. Fang, R. et al. Comprehensive analysis of single cell ATAC-seq data with SnapATAC. *Nat Commun* **12**, 1337 (2021).
5. Kiselev, V.Y., Andrews, T.S. & Hemberg, M. Challenges in unsupervised clustering of single-cell RNA-seq data. *Nat Rev Genet* **20**, 273-282 (2019).

6.   Cortal, A., Martignetti, L., Six, E. & Rausell, A. Gene signature extraction and cell identity
recognition at the single-cell level with Cell-ID. *Nat Biotechnol* (2021).

7.   Vandenbon, A. & Diez, D. A clustering-independent method for finding differentially
expressed genes in single-cell transcriptome data. *Nat Commun* **11**, 4318 (2020).

8.   Dann, E., Henderson, N.C., Teichmann, S.A., Morgan, M.D. & Marioni, J.C. Differential
abundance testing on single-cell data using k-nearest neighbor graphs. *Nat Biotechnol*
(2021).

9.   Hao, Y. et al. Integrated analysis of multimodal single-cell data. *Cell* (2021).

10.   Korsunsky, I. et al. Fast, sensitive and accurate integration of single-cell data with
Harmony. *Nat Methods* **16**, 1289-1296 (2019).

11.   Welch, J.D. et al. Single-Cell Multi-omic Integration Compares and Contrasts Features of
Brain Cell Identity. *Cell* **177**, 1873-1887 e1817 (2019).

12.   Stuart, T. et al. Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902
e1821 (2019).

13.   Lerer, A. et al. Pytorch-biggraph: A large-scale graph embedding system. *arXiv preprint*
arXiv:1903.12287 (2019).

14.   Chen, H. et al. Assessment of computational methods for the analysis of single-cell
ATAC-seq data. *Genome Biology* **20**, 241 (2019).

15.   Buenrostro, J.D. et al. Integrated Single-Cell Analysis Maps the Continuous Regulatory
Landscape of Human Hematopoietic Differentiation. *Cell* **173**, 1535-1548 e1516 (2018).

16.   Ferreira, R., Ohneda, K., Yamamoto, M. & Philipsen, S. GATA1 function, a paradigm for
transcription factors in hematopoiesis. *Molecular and cellular biology* **25**, 1215-1227
(2005).

17.   Tijchon, E., Havinga, J., Van Leeuwen, F. & Scheijen, B. B-lineage transcription factors
and cooperating gene lesions required for leukemia development. *Leukemia* **27**, 541-
552 (2013).

18.   Friedman, A. Transcriptional control of granulocyte and monocyte development.
Oncogene **26**, 6816-6828 (2007).

19.   Schep, A.N., Wu, B., Buenrostro, J.D. & Greenleaf, W.J. chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data. *Nat*
Methods **14**, 975-978 (2017).

20.   Moriguchi, T. & Yamamoto, M. A regulatory network governing Gata1 and Gata2 gene
transcription orchestrates erythroid lineage differentiation. *International journal of*
hematology **100**, 417-424 (2014).

21.   Chen, S., Lake, B.B. & Zhang, K. High-throughput sequencing of the transcriptome and
chromatin accessibility in the same cell. *Nat Biotechnol* (2019).

22.   Ma, S. et al. Chromatin Potential Identified by Shared Single-Cell Profiling of RNA and
Chromatin. *Cell* (2020).

23.   Cao, J. et al. Joint profiling of chromatin accessibility and gene expression in thousands
of single cells. *Science* **361**, 1380-1385 (2018).

24.   Zhu, C. et al. An ultra high-throughput method for single-cell joint analysis of open
chromatin and transcriptome. *Nat Struct Mol Biol* (2019).

25.   Tran, H.T.N. et al. A benchmark of batch-effect correction methods for single-cell RNA
sequencing data. *Genome Biol* **21**, 12 (2020).

26. Leek, J.T. et al. Tackling the widespread and critical impact of batch effects in high-
throughput data. *Nature Reviews Genetics* **11**, 733-739 (2010).
27. Han, X. et al. Mapping the mouse cell atlas by microwell-seq. *Cell* **172**, 1091-1107. e1017
(2018).
28. Tabula Muris, C. et al. Single-cell transcriptomics of 20 mouse organs creates a Tabula
Muris. *Nature* **562**, 367-372 (2018).
29. Baron, M. et al. A single-cell transcriptomic map of the human and mouse pancreas
reveals inter-and intra-cell population structure. *Cell systems* **3**, 346-360. e344 (2016).
30. Muraro, M.J. et al. A single-cell transcriptome atlas of the human pancreas. *Cell systems*
3, 385-394. e383 (2016).
31. Wang, Y.J. et al. Single-cell transcriptomics of the human endocrine pancreas. *Diabetes*
65, 3028-3038 (2016).
32. Segerstolpe, Å. et al. Single-cell transcriptome profiling of human pancreatic islets in
health and type 2 diabetes. *Cell metabolism* **24**, 593-607 (2016).
33. Palla, G. et al. Squidpy: a scalable framework for spatial single cell analysis. *bioRxiv*
2021.02.19.431994v2 (2021).
34. VanHorn, S. & Morris, S.A. Next-Generation Lineage Tracing and Fate Mapping to
Interrogate Development. *Dev Cell* **56**, 7-21 (2021).
35. Ietswaart, R., Gyori, B.M., Bachman, J.A., Sorger, P.K. & Churchman, L.S. GeneWalk
identifies relevant gene functions for a biological context using network representation
learning. *Genome Biol* **22**, 55 (2021).
36. Yuan, H., Kshirsagar, M., Zamparo, L., Lu, Y. & Leslie, C.S. BindSpace decodes
transcription factor binding signals by large-scale sequence embedding. *Nat Methods*
(2019).
37. Li, H., Xiao, X., Wu, X., Ye, L. & Ji, G. scLINE: A multi-network integration framework
based on network embedding for representation of single-cell RNA-seq data. *J Biomed*
Inform **122**, 103899 (2021).
38. Buterez, D., Bica, I., Tariq, I., Andrés-Terré, H. & Liò, P. CELLVGAE: AN UNSUPERVISED
SCRNA-SEQ ANALYSIS WORKFLOW WITH GRAPH ATTENTION NETWORKS. *bioRxiv*
2020.12.20.423645v1 (2020).
39. Sheikh, N., Kefato, Z. & Montresor, A. gat2vec: representation learning for attributed
graphs. *Computing* **101**, 187-209 (2018).
40. Bojchevski, A. & Günnemann, S. Deep gaussian embedding of graphs: Unsupervised
inductive learning via ranking. *arXiv preprint arXiv:1707.03815* (2017).
41. Satopaa, V., Albrecht, J., Irwin, D. & Raghavan, B. in 2011 31st international conference
on distributed computing systems workshops 166-171 (IEEE, 2011).
42. Fornes, O. et al. JASPAR 2020: update of the open-access database of transcription
factor binding profiles. *Nucleic acids research* **48**, D87-D92 (2020).
43. Kadlec, R., Bajgar, O. & Kleindienst, J. Knowledge base completion: Baselines strike back.
arXiv preprint arXiv:1705.10744 (2017).
44. Krompaß, D., Baier, S. & Tresp, V. in International semantic web conference 640-655
(Springer, 2015).
45. Mikolov, T., Chen, K., Corrado, G. & Dean, J. Efficient estimation of word representations
in vector space. *arXiv preprint arXiv:1301.3781* (2013).

46.   Cusanovich, D.A. et al. The cis-regulatory dynamics of embryonic development at single-
1256   cell resolution. *Nature* **555**, 538-542 (2018).
1257   47.   Bravo Gonzalez-Blas, C. et al. cisTopic: cis-regulatory topic modeling on single-cell ATAC-
1258   seq data. *Nat Methods* **16**, 397-400 (2019).
1259   48.   Dou, J. et al. Unbiased integration of single cell multi-omics data. *bioRxiv*,
1260   2020.2012.2011.422014 (2020).
1261
1262

Supplementary Figures

 Supplementary Figure 1. SIMBA analysis of the scRNA-seq 10x PBMCs dataset.

   a. Three default metrics used to evaluate SIMBA training procedure, including training loss
   (top), validation loss (middle), mean reciprocal rank (MRR)
   b. SIMBA metric plots of genes. All the genes are plotted according to the Gini index
   against max score, standard deviation (std) against max score, and entropy against max
   score, respectively. The same set of genes as in Figure 2c are highlighted.
   c. UMAP visualization of SIMBA embeddings of cells colored by cell type or gene
   expression of those genes highlighted in (b).
   d. SIMBA barcode plots of the genes highlighted in (b).

1278
Supplementary Figure 2. Comparison of SIMBA with Scanpy on the scRNA-seq 10x PBMCs
dataset.
a. Venn diagram of top marker genes identified by SIMBA and Scanpy
b. Scanpy-derived UMAP visualization of cells colored by cell type
c. Top marker genes detected only by SIMBA. Colored by intensity of gene expression.
d. Top marker genes detected only by Scanpy. Colored by intensity of gene expression.
e. SIMBA embedding result after implementing variable gene selection. Left: variable gene
selection step implemented in SIMBA. Middle: UMAP visualization of SIMBA
embeddings of cells. Right: UMAP visualization of SIMBA embeddings of cells and
variable genes.

Supplementary Figure 3. SIMBA analysis of the *Buenrostro2018* dataset

    a. UMAP visualization of SIMBA embeddings of cells colored by cell type (top-left), and TF activity scores of TF motifs calculated with chromVAR, respectively. The SIMBA barcode plot of each TF motif is shown below the UMAP plot.

    b. Top: UMAP visualization of SIMBA embeddings of cells colored by TF activity scores of k-mers calculated with chromVAR. Middle: SIMBA barcode plots of the corresponding k-mers. Bottom: the matching known motif against the enriched k-mer sequences.

    c. Heatmap of cells against neighboring peaks of each cell type that are selected in the SIMBA co-embedding space. Chromatin accessibility is binary and colored accordingly.

1305
1306
1307

1308
1309
1310 **Supplementary Figure 4**. Comparison of SIMBA performance using scATAC-seq peaks and DNA
1311 sequence content vs only scATAC-seq peaks. **Top**: UMAP visualization of SIMBA embeddings of
1312 cells for each indicated dataset generated from only scATAC-seq peak information. **Bottom**:
1313 UMAP visualization of SIMBA embeddings of cells for each indicated dataset generated from
1314 scATAC-seq peak information and DNA sequence content information.
1315

1316
1317 **Supplementary Figure 5**. Benchmark of SIMBA against top-performing scATAC-seq analysis
1318 methods.
1319
1320 Top: Evaluation of SIMBA and other methods including cisTopic, SnapATAC, *Cusanovich2018* for
1321 scATAC-seq analysis using metrics 1) ARI, AMI, and Homogeneity for datasets with ground truth
1322 cell type labels and 2) Residual Average Gini Index (RAGI) for the 10x PBMCs dataset without
1323 ground truth labels.
1324
1325 Bottom: UMAP visualization of feature matrices produced by each method on each dataset
1326 colored by cell type annotation or cluster label.
1327

1328
1329
1330 **Supplementary Figure 6**. SIMBA multimodal analysis of the SHARE-seq hair follicle dataset.
1331
a. SIMBA embedding results when both gene expression and chromatin accessibility are
encoded in the graph. Left: UMAP visualization of SIMBA embeddings of cells and genes.
Middle: UMAP visualization of SIMBA embeddings of cells along with genes, TF motifs, and
k-mers. Right: UMAP visualization of SIMBA embeddings of cells along with genes, peaks, TF
motifs, and k-mers.
b. SIMBA embedding results when only gene expression is encoded in the graph. Left: UMAP
visualization of SIMBA embeddings of cells. Right: UMAP visualization of SIMBA embeddings
of cells and variable genes.
c. SIMBA embedding results when only chromatin accessibility is encoded in the graph. Left:
UMAP visualization of SIMBA embeddings of cells. Right: UMAP visualization of SIMBA
embeddings of cells and peaks.
1343

Supplementary Figure 7. Cell type specific marker genes and the target genes of master regulators identified by SIMBA in the SHARE-seq hair follicle subset dataset.

    a. UMAP visualization of SIMBA embeddings of cells colored by cell type and gene expression intensity.

    b. SIMBA barcode plots of each gene plotted above.

    c. SIMBA barcode plots of TF motifs *Lef1* and *Hoxc13.*

    d. SIMBA barcode plots of peaks near the loci of *Lef1* and *Hoxc13.*

    e. Top 30 target genes of the master regulators *Relb, Gata6,* and *Nfatc1* as inferred by SIMBA.

Supplementary Figure 8. SIMBA multimodal analysis of the SNARE-seq mouse cerebral cortex dataset.

a. From top to bottom: UMAP visualization of SIMBA embeddings of (i) cells (ii) genes alongside cells (iii) genes, motifs, and k-mers alongside cells (iv) genes, peaks, motifs, and k-mers alongside cells.

b. UMAP visualization of SIMBA embeddings of cells colored by indicated gene expression intensity.

c. Heatmap of cells against neighboring peaks of each cell type that are selected in the SIMBA co-embedding space. Chromatin accessibility is binary and colored accordingly.

d. SIMBA barcode plots of the genes highlighted in (a).

1371
1372
1373 **Supplementary Figure 9**. SIMBA multimodal analysis of the 10x multiome PBMCs dataset.
1374
a. From top to bottom: UMAP visualization of SIMBA embeddings of (i) cells (ii) genes
alongside cells (iii) genes, motifs, and k-mers alongside cells (iv) genes, peaks, motifs,
and k-mers alongside cells.
b. UMAP visualization of SIMBA embeddings of cells colored by indicated gene expression
intensity.
c. Heatmap of cells against neighboring peaks of each cluster that are selected in the
SIMBA co-embedding space. Chromatin accessibility is binary and colored accordingly.
d. SIMBA barcode plots of the genes highlighted in (a).
1383

1386   **Supplementary Figure 10**. SIMBA-inferred marker genes for the scRNA-seq mouse atlas dataset
1387   in batch correction analysis.
1388
a. UMAP visualization of SIMBA embeddings of cells colored by cell type.
b. UMAP visualization of SIMBA embeddings of cells and genes.
c. UMAP visualization of SIMBA embeddings of cells and genes. Biological "query" points
are highlighted with a red "+". Nearby informative genes are colored accordingly.
d. UMAP visualization of SIMBA embeddings of cells colored by indicated gene expression
intensity, separated by cell type.
1395

1396
1397

Supplementary Figure 11. SIMBA-inferred marker genes for the scRNA-seq human pancreas
dataset in batch correction analysis.

1400

a. UMAP visualization of SIMBA embeddings of cells colored by cell type.
b. UMAP visualization of SIMBA embeddings of cells and genes.
c. UMAP visualization of SIMBA embeddings of cells and genes. Biological "query" points
   are highlighted with a red "+". Nearby informative genes are colored accordingly.
d. UMAP visualization of SIMBA embeddings of cells colored by indicated gene expression
   intensity, separated by cell type.

1407

1408
Supplementary Figure 12. Comparison of SIMBA to other methods for batch correction of the
mouse atlas (a-b) and human pancreas scRNA-seq datasets (c-d).
a, c. Quantitative comparison of SIMBA with three other batch correction methods
including Seurat3, LIGER and Harmony, using, left-to-right: average silhouette width (ASW),
adjusted Rand index (ARI), and local inverse Simpson's index (LISI)
b, d. UMAP visualization of raw and preprocessed data alongside the batch corrected
results produced by Seurat3, LIGER, Harmony, and SIMBA. Colored by technology (top) and
cell type (bottom).
1417

1418
1419
Supplementary Figure 13. SIMBA-inferred marker features for the SHARE-seq mouse skin
dataset in multi-omics integration analysis.

1422

    a. UMAP visualization of SIMBA embeddings of cells with two cellular modalities
        integrated.
    b. UMAP visualization of SIMBA embeddings of cells, genes, and peaks with two cellular
        modalities integrated.

c.   UMAP visualization of SIMBA embeddings of cells, genes, and peaks with two cellular
modalities integrated. Biological "query" points are highlighted with a red "+". Nearby
informative genes and peaks are colored accordingly.
d.   UMAP visualization of SIMBA embeddings of cells, genes, and peaks with two cell
modalities integrated and known marker genes highlighted.
e.   UMAP visualization of SIMBA embeddings of cells colored by indicated gene expression
intensity, separated by cell type.
f.   Heatmap of cells against neighboring peaks of each cell type that are selected in the
SIMBA co-embedding space. Chromatin accessibility is binary and colored accordingly.
1436

1437

1438
1439
1440 **Supplementary Figure 14**. SIMBA-inferred marker features for the 10x human PBMCs dataset in
1441 multi-omics integration analysis.
1442
a. UMAP visualization of SIMBA embeddings of cells with two cellular modalities
integrated.
b. UMAP visualization of SIMBA embeddings of cells, genes, and peaks with two cellular
modalities integrated.
c. UMAP visualization of SIMBA embeddings of cells, genes, and peaks with two cellular
modalities integrated. Biological "query" points are highlighted with a red "+". Nearby
informative genes and peaks are colored accordingly.
d. Heatmap of cells against neighboring peaks of each cluster that are selected in the
SIMBA co-embedding space. Chromatin accessibility is binary and colored accordingly.
e. UMAP visualization of SIMBA embeddings of cells colored by indicated gene expression
intensity, separated by cell type.
1454

Supplementary Figure 15. Comparison of SIMBA to other methods for multi-omics integration

of the SHARE-seq mouse skin (a-b) and 10x multiome human PBMCs (c-d) datasets.

a, c. Quantitative comparison of SIMBA with two other methods including Seurat3,

LIGER for multi-omics integration, using, left-to-right: anchoring distance rank,

anchoring distance, silhouette index, and Fraction in the same cluster.

b, d. UMAP visualization of the raw scRNA-seq and scATAC-seq data from the 10x

multiome human PBMCs dataset alongside the integrated results produced by Seurat3,

LIGER, and SIMBA. Colored by data modality (top) and cluster assignment (bottom). The

red intervals of violin plot of Anchoring distance and Silhouette index shows the 95%

of the mean.

Supplementary Notes

Supplementary Note 1: Comparison with scATAC-seq methods

To assess SIMBA's ability to cluster cell types based on scATAC-seq profiles, we compared SIMBA
with specialized methods specifically designed for this task. We observed that SIMBA yields
consistent embeddings of cells when using either a single feature (peaks) or multiple features
(peaks and DNA sequences from within those peaks) as input to the graph. This comparison was
performed across four scATAC-seq datasets of varying profiling technologies and organisms
(**Supplementary Fig. 4**). Given these differences, to create a fair comparison we used the same
set of features (i.e., peaks) for SIMBA as other methods. SIMBA's performance was compared
against three of the top methods, including SnapATAC[4], *Cusanovich*2018[46], and cisTopic[47]
recommended by our recent benchmark study[14]. This comparison was first made qualitatively
based on UMAP visualization and then quantitatively based on clustering performance. SIMBA
performed as well as or better than each of the methods evaluated. These results comparing
SIMBA to scATAC-seq-specialized methods highlight SIMBA's wide utility for single-cell analyses
(**Supplementary Fig. 5**).

Supplementary Note 2: Comparison with batch correction methods

Multiple methods have now been developed to correct for the technical effects of sample
preparation and data collection in single cells. To assess SIMBA's performance in removing batch
effects, we compared it to Seurat3[12], LIGER[11] and Harmony[10], three top-performing batch
correction methods recommended in a recent benchmark study[25].

Two datasets, including a mouse atlas dataset and a human pancreas dataset (see
Supplementary Table 1), were used for the evaluation. The mouse atlas dataset is composed of
two scRNA-seq subsets with shared cell types from different sequencing platforms. The human
pancreas dataset is composed of five samples pooled from five distinct sources using four
different sequencing techniques wherein not all cell types are shared across each sample.

To qualitatively compare these methods, we visualized cells of each dataset before and after
batch-correction in UMAP plots (**Supplementary Fig. 12b,d**). To quantitatively evaluate the
performance of each method, using the benchmarking pipeline laid out in Tran *et al*[25], we
measured the conservation of biological information and batch effect removal based on three
different metrics: average silhouette width (ASW), adjusted Rand index (ARI), and local inverse
Simpson's index (LSI)[10] as in the previously-mentioned benchmark study[25] (**Supplementary Fig.
12a,c**; **Methods**). Each metric measures the relative mixing of class labels, where optimal
performance is associated with maximal mixing in the batch labels and minimal mixing in the
cell type labels.

The "Raw" batch correction results are the first 50 principal components of the horizontally
concatenated gene-by-cell expression count matrix using *stats::prcomp* in R package with

1509   centering and scaling. The "Raw, preprocessed" batch correction used the preprocessed data
1510   with log normalization with scaling factor 10^4 and selection of 3000 highly variable genes with
1511   Seurat v3 with no restriction on the minimum number of cells and genes.
1512
1513   For batch correction using Seurat v3, default options are used for pancreas dataset whereas for
1514   mouse atlas dataset no cutoff was used for the minimum number of cells and genes as in Tran
1515   *et al.*[25]. The dimension of the batch corrected embedding is set as 50 dimensions following the
1516   default option for *Seurat::RunPCA* and for the consistency with SIMBA.
1517
1518   For batch correction using LIGER, the same arguments are used (lambda = 5, nrep = 3) are used
1519   for *liger::optimizeALS* in Tran *et al*. other than the number of factors k was set as 50 for
1520   consistency with other methods for both datasets.
1521
1522   For batch correction using Harmony, the same arguments are used as in Tran *et al.*[25] other than
1523   the number of dimensions of the output embedding was set to 50 instead of 20. We note that
1524   the output embedding of 20 dimensions would result in the similar result as when used 50
1525   dimensions in these methods.
1526
1527   **Supplementary Note 3**: **Comparison with multi-omics integration methods**
1528
1529   Seurat3 and LIGER are two of the most widely-adopted methods for single-cell data integration.
1530   Here, we demonstrate that SIMBA outperforms these methods on two separate datasets, the
1531   recently published SHARE-seq mouse skin dataset and the similarly recent 10x PBMCs multiome
1532   dataset (**Supplementary Table 1**). We focus on Seurat3 and LIGER as they have explicit
1533   documentation for the task of integrating scRNA-seq and scATAC-seq data.
1534
We first qualitatively evaluated these methods by inspecting UMAP visualization plots. For the
SHARE-seq dataset, we observed that all three methods perform comparably well in mixing
cells of two modalities though LIGER generated particularly small and noisy clusters
(**Supplementary Fig. 15b)**. For the 10X PBMCs dataset, SIMBA resulted in the best mixing of
cells belonging to each modality whereas other methods clustered cells separately within the
originating modalities (**Supplementary Fig. 15d**). We next quantitatively assessed the
integration performance of each method using four metrics that measure the distances
between matched cells in the integrated space (**Methods**). In addition to the commonly-used
metrics including anchoring distance, Silhouette index, and Fraction in the same cluster, we
developed an additional metric, *anchoring distance rank* (ADR)*,* which represents the
normalized rank of the distance between matching cells. If two matching cells from scRNA-seq
and scATAC-seq are mutually closest to one another, their ADR will be close to 0 (**Methods**) and
thus a minimized ADR is ideal. Overall SIMBA showed the best performance according to ADR
as well as cluster agreement while showing comparable or better performance according to the
remaining metrics for both datasets (**Supplementary Fig. 15a,c**).
1550
1551

1552 The modality integration procedure for Seurat v3 and LIGER follows the tutorial provided by the
1553 authors (Seurat v3:
1554 https://satijalab.org/seurat/archive/v3.1/atacseq_integration_vignette.html; LIGER:
1555 http://htmlpreview.github.io/?https://github.com/welch-
1556 lab/liger/blob/master/vignettes/Integrating_scRNA_and_scATAC_data.html).
1557
Both Seurat v3 and LIGER formulate the modality integration task between scRNA-seq and
scATAC-seq data as a batch correction task between scRNA-seq and gene activity matrix
constructed from scATAC-seq. In Seurat v3, the gene activity score of a gene is calculated as the
sum of the read counts in the peaks that falls within from 2kb upstream of the TSS to the end of
the gene body. In LIGER, this score is calculated as the sum of all read counts that falls within
3kb upstream of the TSS to the end of the gene body.
1564
The "Raw" results start from a scRNA-seq count matrix and a gene activity matrix calculated by
Seurat v3. Filtering for the shared genes in both modalities resulted in 16738 genes for the
SHARE-seq mouse skin dataset and 11045 genes for the 10X PBMCs dataset. Gene-by-cell gene
expression matrix and gene activity matrix were horizontally concatenated along matching rows
(genes). The output embedding is the first 20 principal components calculated by the R function
stats::prcomp with centering and scaling.
1571
1572 For the modality integration using Seurat v3, the gene expression count was filtered using the
1573 default parameters *min.cells = 3* and *min.features = 200*. The co-embedding was created as
1574 described in the tutorial of the package using the scRNA-seq. The output embedding consists of
1575 the first 50 principal components, which is the default option of *Seurat::RunPCA*.
1576
For the modality integration using LIGER, the gene expression count and gene activity matrices
were normalized and filtered for the genes that are shared between both matrices. The values
were then scaled according to the tutorial. In applying LIGER to the SHARE-seq mouse skin
dataset, the function, *liger::optimizeALS* was used with the default parameters, k = 20 and
lambda = 5. The scRNA-seq dataset was indicated as the reference in the function,
liger::quantile_norm as described in the documentation. The scRNA-seq and scATAC-seq
modalities of the 10X PBMC multiome dataset were unable to be aligned using the default
parameters. Thus *lambda = 30* and *max.iters = 100* were used for the *liger::optimizeALS*
function and the scATAC-seq dataset was indicated as the reference using the
liger::quantile_norm function to ensure a better alignment.
1587
1588 References
1589

1.   Satija, R., Farrell, J.A., Gennert, D., Schier, A.F. & Regev, A. Spatial reconstruction of
single-cell gene expression data. *Nat Biotechnol* **33**, 495-502 (2015).
2.   Wolf, F.A., Angerer, P. & Theis, F.J. SCANPY: large-scale single-cell gene expression data
analysis. *Genome Biol* **19**, 15 (2018).
3.   Granja, J.M. et al. ArchR is a scalable software package for integrative single-cell
chromatin accessibility analysis. *Nat Genet* **53**, 403-411 (2021).

4.   Fang, R. et al. Comprehensive analysis of single cell ATAC-seq data with SnapATAC. *Nat*
Commun **12**, 1337 (2021).
5.   Kiselev, V.Y., Andrews, T.S. & Hemberg, M. Challenges in unsupervised clustering of
single-cell RNA-seq data. *Nat Rev Genet* **20**, 273-282 (2019).
6.   Cortal, A., Martignetti, L., Six, E. & Rausell, A. Gene signature extraction and cell identity
recognition at the single-cell level with Cell-ID. *Nat Biotechnol* (2021).
7.   Vandenbon, A. & Diez, D. A clustering-independent method for finding differentially
expressed genes in single-cell transcriptome data. *Nat Commun* **11**, 4318 (2020).
8.   Dann, E., Henderson, N.C., Teichmann, S.A., Morgan, M.D. & Marioni, J.C. Differential
abundance testing on single-cell data using k-nearest neighbor graphs. *Nat Biotechnol*
(2021).
9.   Hao, Y. et al. Integrated analysis of multimodal single-cell data. *Cell* (2021).
10.   Korsunsky, I. et al. Fast, sensitive and accurate integration of single-cell data with
Harmony. *Nat Methods* **16**, 1289-1296 (2019).
11.   Welch, J.D. et al. Single-Cell Multi-omic Integration Compares and Contrasts Features of
Brain Cell Identity. *Cell* **177**, 1873-1887 e1817 (2019).
12.   Stuart, T. et al. Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902
e1821 (2019).
13.   Lerer, A. et al. Pytorch-biggraph: A large-scale graph embedding system. *arXiv preprint*
arXiv:1903.12287 (2019).
14.   Chen, H. et al. Assessment of computational methods for the analysis of single-cell
ATAC-seq data. *Genome Biology* **20**, 241 (2019).
15.   Buenrostro, J.D. et al. Integrated Single-Cell Analysis Maps the Continuous Regulatory
Landscape of Human Hematopoietic Differentiation. *Cell* **173**, 1535-1548 e1516 (2018).
16.   Ferreira, R., Ohneda, K., Yamamoto, M. & Philipsen, S. GATA1 function, a paradigm for
transcription factors in hematopoiesis. *Molecular and cellular biology* **25**, 1215-1227
(2005).
17.   Tijchon, E., Havinga, J., Van Leeuwen, F. & Scheijen, B. B-lineage transcription factors
and cooperating gene lesions required for leukemia development. *Leukemia* **27**, 541-
552 (2013).
18.   Friedman, A. Transcriptional control of granulocyte and monocyte development.
Oncogene **26**, 6816-6828 (2007).
19.   Schep, A.N., Wu, B., Buenrostro, J.D. & Greenleaf, W.J. chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data. *Nat*
Methods **14**, 975-978 (2017).
20.   Moriguchi, T. & Yamamoto, M. A regulatory network governing Gata1 and Gata2 gene
transcription orchestrates erythroid lineage differentiation. *International journal of*
hematology **100**, 417-424 (2014).
21.   Chen, S., Lake, B.B. & Zhang, K. High-throughput sequencing of the transcriptome and
chromatin accessibility in the same cell. *Nat Biotechnol* (2019).
22.   Ma, S. et al. Chromatin Potential Identified by Shared Single-Cell Profiling of RNA and
Chromatin. *Cell* (2020).
23.   Cao, J. et al. Joint profiling of chromatin accessibility and gene expression in thousands
of single cells. *Science* **361**, 1380-1385 (2018).

24. Zhu, C. et al. An ultra high-throughput method for single-cell joint analysis of open chromatin and transcriptome. *Nat Struct Mol Biol* (2019).

25. Tran, H.T.N. et al. A benchmark of batch-effect correction methods for single-cell RNA sequencing data. *Genome Biol* **21**, 12 (2020).

26. Leek, J.T. et al. Tackling the widespread and critical impact of batch effects in high-throughput data. *Nature Reviews Genetics* **11**, 733-739 (2010).

27. Han, X. et al. Mapping the mouse cell atlas by microwell-seq. *Cell* **172**, 1091-1107. e1017 (2018).

28. Tabula Muris, C. et al. Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris. *Nature* **562**, 367-372 (2018).

29. Baron, M. et al. A single-cell transcriptomic map of the human and mouse pancreas reveals inter-and intra-cell population structure. *Cell systems* **3**, 346-360. e344 (2016).

30. Muraro, M.J. et al. A single-cell transcriptome atlas of the human pancreas. *Cell systems* **3**, 385-394. e383 (2016).

31. Wang, Y.J. et al. Single-cell transcriptomics of the human endocrine pancreas. *Diabetes* **65**, 3028-3038 (2016).

32. Segerstolpe, Å. et al. Single-cell transcriptome profiling of human pancreatic islets in health and type 2 diabetes. *Cell metabolism* **24**, 593-607 (2016).

33. Palla, G. et al. Squidpy: a scalable framework for spatial single cell analysis. *bioRxiv 2021.02.19.431994v2* (2021).

34. VanHorn, S. & Morris, S.A. Next-Generation Lineage Tracing and Fate Mapping to Interrogate Development. *Dev Cell* **56**, 7-21 (2021).

35. Ietswaart, R., Gyori, B.M., Bachman, J.A., Sorger, P.K. & Churchman, L.S. GeneWalk identifies relevant gene functions for a biological context using network representation learning. *Genome Biol* **22**, 55 (2021).

36. Yuan, H., Kshirsagar, M., Zamparo, L., Lu, Y. & Leslie, C.S. BindSpace decodes transcription factor binding signals by large-scale sequence embedding. *Nat Methods* (2019).

37. Li, H., Xiao, X., Wu, X., Ye, L. & Ji, G. scLINE: A multi-network integration framework based on network embedding for representation of single-cell RNA-seq data. *J Biomed Inform* **122**, 103899 (2021).

38. Buterez, D., Bica, I., Tariq, I., Andrés-Terré, H. & Liò, P. CELLVGAE: AN UNSUPERVISED SCRNA-SEQ ANALYSIS WORKFLOW WITH GRAPH ATTENTION NETWORKS. *bioRxiv 2020.12.20.423645v1* (2020).

39. Sheikh, N., Kefato, Z. & Montresor, A. gat2vec: representation learning for attributed graphs. *Computing* **101**, 187-209 (2018).

40. Bojchevski, A. & Günnemann, S. Deep gaussian embedding of graphs: Unsupervised inductive learning via ranking. *arXiv preprint arXiv:1707.03815* (2017).

41. Satopaa, V., Albrecht, J., Irwin, D. & Raghavan, B. in 2011 31st international conference on distributed computing systems workshops 166-171 (IEEE, 2011).

42. Fornes, O. et al. JASPAR 2020: update of the open-access database of transcription factor binding profiles. *Nucleic acids research* **48**, D87-D92 (2020).

43. Kadlec, R., Bajgar, O. & Kleindienst, J. Knowledge base completion: Baselines strike back. *arXiv preprint arXiv:1705.10744* (2017).

1684   44.   Krompaß, D., Baier, S. & Tresp, V. in International semantic web conference 640-655
1685   (Springer, 2015).
1686   45.   Mikolov, T., Chen, K., Corrado, G. & Dean, J. Efficient estimation of word representations
1687   in vector space. *arXiv preprint arXiv:1301.3781* (2013).
1688   46.   Cusanovich, D.A. et al. The cis-regulatory dynamics of embryonic development at single-
1689   cell resolution. *Nature* **555**, 538-542 (2018).
1690   47.   Bravo Gonzalez-Blas, C. et al. cisTopic: cis-regulatory topic modeling on single-cell ATAC-
1691   seq data. *Nat Methods* **16**, 397-400 (2019).
1692   48.   Dou, J. et al. Unbiased integration of single cell multi-omics data. *bioRxiv*,
1693   2020.2012.2011.422014 (2020).
1694
1695