SIMBA: SIngle-cell eMBedding Along with features

Huidong Chen[1, 2, 3], Jayoung Ryu[1, 2, 4], Michael E. Vinyard[1, 2, 3, 5], Adam Lerer[6]+, Luca Pinello[1, 2, 3]+

1. Molecular Pathology Unit, Center for Cancer Research, Massachusetts General Hospital,
Charlestown, MA, USA
2. Department of Pathology, Harvard Medical School, Boston, MA, USA
3. Broad Institute of Harvard and MIT, Cambridge, MA, USA
4. Department of Biomedical Informatics, Harvard Medical School, Boston, MA, USA
5. Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA, USA
6. Facebook AI Research

+Corresponding author

Abstract

Recent advances in single-cell omics technologies enable the individual and joint profiling of
cellular measurements. Currently, most single-cell analysis pipelines are cluster-centric and
cannot explicitly model the interactions between different feature types. In addition, single-cell
methods are generally designed for a particular task as distinct single-cell problems are
formulated differently. To address these current shortcomings, we present *SIMBA*, a graph
embedding method that jointly embeds single cells and their defining features, such as genes,
chromatin accessible regions, and transcription factor binding sequences into a common latent
space. By leveraging the co-embedding of cells and features, SIMBA allows for the study of
cellular heterogeneity, clustering-free marker discovery, gene regulation inference, batch effect
removal, and omics data integration. SIMBA has been extensively applied to scRNA-seq,
scATAC-seq, and dual-omics data. We show that SIMBA provides a single framework that allows
diverse single-cell analysis problems to be formulated in a unified way and thus simplifies the
development of new analyses and integration of other single-cell modalities. SIMBA is
implemented as an efficient, comprehensive, and extensible Python library (https://simba-
bio.readthedocs.io) for the analysis of single-cell omics data using graph embedding.

Introduction

Technology to profile single cells has advanced to several molecular modalities, dramatically
advancing our ability to characterize cell states as well as discover key molecular machinery
that underlies both development and disease. Individual cells are now measured using multiple
molecular modalities, simultaneously. At the same time, single-cell experiments have scaled
such that tens of thousands of cells can be routinely profiled. The emergence of single-cell
multi-omics technologies allows for the measurements of multiple cellular layers, including
genomics, epi-genomics, transcriptomics, and proteomics. Such assays have pioneered an
avenue toward a better understanding of the interplay between layers as they jointly define cell

states based on diverse genomic and molecular features including genes, regulatory elements,
and transcription factors. While single-cell multi-omic assays have quickly evolved towards the
incorporation of additional modalities with increasing resolution, harnessing their full potential
has posed several significant computational challenges.
48
Many single-cell computational methods have been developed for the analysis of one modality
(e.g., scRNA-seq or scATAC-seq analysis) [1-4]. Common to these methods is a workflow that
includes routine steps such as feature selection, dimension reduction, clustering, and
differential feature detection. These "cluster-centric" analysis methods rely on accurately
defined clustering solutions to discover meaningful and informative marker features.
Unfortunately, clustering solutions may range widely within the space of the user-defined
clustering resolution (number of clusters) and the chosen clustering algorithm. These
parameters may markedly influence the resulting cluster assignment and clusters may not
always correspond to the correct or intended cell populations, thereby leading to inconsistent
and potentially misleading biological annotations[5]. Although initial efforts have been made
recently to develop clustering-free approaches to discover informative genes, they are
specifically designed for extracting gene signatures [6, 7] or identifying perturbations between
experimental conditions[8] from scRNA-seq data, and are therefore limited to single-modality
and single-task analysis.
63
In addition to single-batch/modality analysis, approaches have also been proposed for multi-
batch and cross-modality analysis, such as multimodal analysis (distinct cellular parameters are
measured in the same cell)[9], batch correction (the same cellular parameter is measure in
different batches) [10-12], and integration of multi-omics datasets (distinct cellular parameters are
measured in different cells)[11, 12]. These approaches play a critical role in removing batch effects
that confound true biological variation, improving the characterization of cell states by
leveraging the unique strengths of each assay, and providing insights into the complex
mechanisms of gene regulation. However, these tasks are formulated differently from those in
single-batch/modality settings and thus require development of new dedicated analysis
techniques. Also, while multiple types of cellular features might be present, the relation
between features cannot be exploited directly by most current methods. Furthermore, similar
to single-batch/modality analysis methods, these methods identify marker features based on
groups of cells obtained by clustering and therefore are limited to clustering solutions.
Additionally, instead of directly identifying marker features in the integrated space, most batch
correction/multi-omics integration methods need to first detect marker features in the
uncorrected/unintegrated original space of each batch/modality independently, and then
combine them, thus resulting in potentially inconsistent interpretations between
batches/modalities.
82
To overcome the limitations in both single-batch/modality analysis and multi-batch/cross-
modality analysis, we propose SIMBA (**SIngle-cell eMBedding Along with features**), a versatile
single-cell embedding method that co-embeds cells and features into a shared latent space, in
which various types of tasks can be performed based on the proximity between entities
including cells and features such as genes, peaks, and DNA sequences. Unlike existing methods

that require featurization of cells, SIMBA directly encodes the cell-feature or feature-feature
relations into a large multi-entity graph. For each task, SIMBA constructs a graph, wherein
differing entities (i.e., cells and features) are represented as nodes and relations between these
entities are encoded as edges. Once the graph is constructed, SIMBA then applies a multi-entity
graph embedding algorithm derived from social networking technologies as well as a Softmax-
based transformation to embed the nodes/entities of the graph into a common low-
dimensional space wherein cells and features can be analyzed based on their distance. Hence
SIMBA provides an information-rich embedding space containing cells and all the features,
serving as an informative database of entities. Depending on the task, we can define biological
queries on the "SIMBA database" by considering neighboring entities of either a cell (or cells) or
a feature (or features) at the individual-cell and individual-feature level (**Methods**). For
example, the query for a cell's neighboring features can be used to identify marker features
(e.g., marker genes or peaks) or to study the interaction between features (e.g., peak-gene)
while the query for features' neighboring cells can be used to annotate cells.
102
By formulating single-cell analyses as multi-entity graph embedding problems, we show SIMBA
can be used to solve popular single-cell tasks in a unified framework that would otherwise
require the development of distinct specialized approaches for each task, including: 1)
dimensionality reduction techniques for studying cellular states; 2) clustering-free marker
detection based on the similarity between single cells and features; 3) Single-cell multimodal
analysis and the study of gene regulation; 4) batch correction and omics integration analysis as
well as the simultaneous identification of marker features. SIMBA is adapted to these diverse
analysis tasks by simply modifying how the input graph is constructed from the relevant single-
cell data.
112
We extensively tested SIMBA in multiple scRNA-seq, scATAC-seq and dual-omics datasets
covering popular single-cell tasks including scRNA-seq analysis, scATAC-seq analysis, multimodal
analysis, batch correction, and multi-omics integration. We demonstrate that SIMBA learns the
joint low-dimensional representations of both cells and features and thus enables the ability to
simultaneously study cellular heterogeneity as well as proximity-based marker feature
detection or gene regulation inference in a clustering-free way. We also demonstrate that
SIMBA performs better than or comparably to current state-of-the-art methods specifically
developed for each task.
121
Importantly, we developed a scalable and comprehensive Python package that enables
seamless interaction between graph construction, training with PyTorch for graph embedding,
and post-training analysis. The SIMBA package not only provides a self-contained framework
that covers preprocessing, graph embedding, and visualization, but also is compatible with
popular single cell analysis tool Scanpy[2]. SIMBA with detailed documentation and extensive
tutorials is available at https://simba-bio.readthedocs.io.
128
129   We believe that SIMBA, as a broadly applicable approach for single cell omics study, not only
130   outperforms current cluster-centric analysis, but also will simplify the burden of developing

131 methods for new single-cell tasks and measurements, while increasing interpretability of
132 cellular mechanisms and functions.
133
134 **Results**
135
Overview of SIMBA

SIMBA is a single-cell embedding method with support for single- or multi- modality analyses
that embeds cells and their associated genomic features into a shared latent space, generating
interpretable and comparable embeddings of cells and features. It leverages recent graph
embedding techniques that have been successful in modeling complex and hierarchical
information present in natural languages, social networks, and other domains, as "knowledge
graphs". In our case, the graph encodes cells, different components of cellular regulatory
circuits, and the relations between them.

Figure1. SIMBA framework overview. SIMBA co-embeds cells and various features
measured during single-cell experiments into a shared latent space to accomplish both
common tasks involved in single-cell data analysis as well as tasks, which remain as open
problems in single-cell genomics. **(Left)** Examples of possible biological entities may be
encoded by SIMBA including cells, gene expression measurements, chromatin accessible
regions, TF motifs, and k-mer sequences found in reads. **(Middle)** SIMBA embedding plot

with multiple types of entities into a low-dimensional space. All entities represented as
shapes (cell = circle, peak = triangle, gene = square, TF motif = star, k-mer = hexagon) are
colored by relevant cell type (green, orange, and blue in this example). Non-informative
features are colored dark grey. Within the graph, each entity is a node, and an edge
indicates a relation between entities (e.g., a gene is expressed in a cell, a chromatin region
is accessible in a cell, or a TF motif/k-mer is present within an open chromatin region,
etc.). Once connected in a graph, these entities may be embedded into a shared low-
dimensional space, with cell-type specific entities embedded in the same neighborhood
and non-informative features embedded elsewhere. **(Right)** Common single-cell analysis
tasks that may be accomplished using SIMBA.
161

SIMBA first encodes different types of entities such as cells, genes, open chromatin regions
(peaks or bins), transcription factor (TF) motifs, and k-mers (short sequences of a specific
length, k), into a single graph (**Fig. 1**, **Methods**) where each node represents an individual entity
and edges indicate relations between entities. For example, if a gene is expressed in a cell, an
edge is created between the gene and cell. The weight of this edge is determined by the gene
expression level. Similarly, an edge is added between a cell and a chromatin region if the region
is open in this cell, or between a chromatin region and a TF motif if the TF motif is found in the
region.

Once the input graph is constructed, a low-dimensional representation of the graph nodes is
then computed using an unsupervised graph embedding method. This graph embedding
procedure leverages the PyTorch-BigGraph framework [13], which allows SIMBA to scale to
millions of cells (**Methods**). The obtained SIMBA space provides an intuitive way to study gene
regulation and the regulatory mechanisms underlying cell differentiation and specification. The
resulting joint embedding of cells and features not only reconstructs the heterogeneity of cells
but also allows for the discovery of the defining features for each individual cell without relying
on a clustering solution, separating cell-type specific features from the non-informative
features. In fact, the relationship between cells and features can be explored directly through
their mutual proximity in the SIMBA embedding as the distance between embedded nodes
reflects their edge probability, which is informative of the potential importance of a feature to a
cell and the interplay between features (**Methods**).

Therefore, cell-type-specific features such as marker genes, cis-regulatory elements can be
discovered without clustering in two different ways. When the labels of cells are known, marker
features can be identified as the neighboring features of cells by performing biological queries
(**Methods**). When these labels are unknown, marker features can be identified through
calculating the imbalance of edge probabilities between a feature and all cells using metrics
such as the Gini index (**Methods**).

Importantly, graph construction is inherently flexible, enabling SIMBA to be applied to a wide
variety of single-cell tasks. In the following sections, we demonstrate the application of SIMBA
to several popular single-cell tasks including scRNA-seq, scATAC-seq, multimodal analysis, batch

correction and multi-omics integration (**Fig. 1**). Extensions to additional tasks will become
readily apparent to the reader and are later discussed.

193
194
SIMBA enables simultaneous learning of cellular heterogeneity and individual-cell-level
marker genes in scRNA-seq analysis

Single-cell RNA sequencing (scRNA-seq) is the most robust and widely used measurement to
profile single cells. **Figure 2a** provides an illustrative overview of the SIMBA graph construction
and the resulting low-dimensional embedding matrix of both cells and genes. Here we show
how SIMBA enables simultaneous dimensionality reduction and clustering-free marker gene
detection in scRNA-seq analysis. We applied SIMBA to a popular PBMCs dataset from 10x
Genomics (**Supplementary Table 1**) to illustrate its workflow. After the standard preprocessing
steps including normalization and log-transformation, SIMBA discretizes the gene expression
matrix into multiple gene expression levels (five levels, by default). The input graph is then
constructed wherein two types of nodes –cells and genes are connected by edges that embody
the relation between them and are weighted according to the corresponding multiple levels of
gene expression. SIMBA then generates embeddings of these nodes through a graph
embedding procedure (**Fig. 2a; Methods**). Depending on the task, we have the full flexibility to
visualize either the whole SIMBA embeddings (embeddings of cells and all genes in
Supplementary Fig. 1c) or the partial SIMBA embeddings (embeddings of cells in **Fig. 2b,** or
embeddings of cells and variable genes in **Fig. 2c**, or embeddings of any entities of interest)
using visualization tools such as UMAP.

When the SIMBA embeddings of cells were visualized, each of the eight cell types, including B
cells, megakaryocytes, CD14 monocytes, FCGR3A monocytes, dendritic cells, NK cells, CD4 T,
and CD8 T cells, was clearly separated (**Fig. 2b**). When the SIMBA embeddings of both cells and
genes were visualized, the co-embedding space showed that SIMBA not only recovered the
cellular heterogeneity, but also correctly embedded informative genes close to relevant cell
types (**Fig. 2c**). The same set of marker genes used to annotate these cells from Scanpy[2] was
highlighted on the UMAP plot. In addition, as a control, we also show the locations of two
housekeeping genes *GAPDH* and *B2M*, which would not be expected to associate with any
particular cell type. From the UMAP plot, we can see that SIMBA not only was able to embed
major-cell-group specific genes to the correct locations (e.g., *IL7R* was embedded into CD4T
cells and *MS4A1* was embedded into B cells), but also was robust to rare-cell-group specific
genes (e.g., *PPBP* was embedded into megakaryocytes). On the contrary, non-informative or
non-cell-type specific genes such as *GAPDH* and *B2M* were embedded in the middle of all cell
groups (**Fig. 2c and Supplementary Fig. 1c**).

These highlighted genes can be further confirmed with "barcode plot", which visualizes the
estimated probability of assigning a feature to a cell by SIMBA based on the recovered edge
confidence (**Fig. 2d, Supplementary Fig. 1e, Methods**). An imbalance in probability indicates
the association of a gene to a sub-population of cells (often corresponding to known cell-types),
whereas a uniform probability distribution indicates a non-cell-type-specific gene. For marker

genes (*CST3* for monocytes and dendritic cells, *MS4A1* for B cells, and *NGK7* for NK and CD8T cells), we observed a clear excess in the probability of assigning each gene to their respective cell types. Conversely, for the housekeeping gene *GAPDH*, we observed a more uniform distribution with much lower probability of associating that gene with the top-ranked cells.

Figure 2. Single-cell RNA-seq analysis of the 10x PBMCs dataset using SIMBA. **(a)** SIMBA graph construction and embedding in scRNA-seq analysis. Biological entities including cells and genes are represented as shapes and colored by relevant cell types (green and orange). Non-informative genes are colored dark grey. Gene expression measurements for each cell are organized into a cell-by-gene matrix. These normalized non-negative observed values undergo discretization into five gene expression levels. Cells and genes are then assembled into a graph with nodes representing cells and genes, and edges between them representing different gene expression levels. This graph may then be embedded into a lower dimensional space resulting in a #entities x #dimension (by default, 50) SIMBA embedding matrix. **(b)** UMAP visualization of SIMBA embeddings of cells

colored by cell type. **(c)** UMAP visualization of SIMBA embeddings of cells and variable
genes. Cells are colored according to cell type as defined in b. Genes are colored slate
blue. Cell-type-specific marker genes and housekeeping genes recovered by Scanpy are
indicated with text and arrows. Genes highlighted in red are shown in **d**, **e**, and **f. (d)**
SIMBA barcode plots of genes *CST3*, *MS4A1*, *NKG7*, and *GAPDH*. The x-axis indicates the
ordering of a cell as ranked by the probability for each cell to be associated with a given
gene. The y-axis describes the probability. The sum of probability over all cells is equal to
1. Each cell is one bar and colored according to cell type as defined in b. € SIMBA ranking
of genes based on the proposed metrics. All the genes are plotted according to the Gini
index against max score. The same set of genes as in **c** are annotated. **(f)** UMAP
visualization of SIMBA embeddings of cells colored by gene expression of (left to right):
CST3, *NKG7*, *MS4A1*, and *GAPDH*.
259

SIMBA also provides several quantitative metrics (termed "SIMBA metrics"), including max
value, Gini index, standard deviation, and entropy, to assess cell-type specificity of various
features without requiring the prior knowledge such as cluster labels, predefined cell types, or
known marker genes (**Methods**). As an example, by inspecting the gene metric plot of max
value (a measurement of maximum probability, a higher value indicates higher cell-type
specificity) vs Gini index (a measurement of imbalance, a higher value indicates higher cell-type
specificity), we observed that the marker genes (e.g., *CST3*, *NKG7*, *MS4A1*) fall in the upper right
corner, as opposed to housekeeping genes (e.g., *GAPDH*) in the lower left corner (**Fig. 2e**).
Similar separation is observed with other metrics (**Supplementary Fig. 1b**). The cell type
specificity of the selected marker genes was further confirmed by visualizing their expression
pattern on UMAP plots (**Fig. 2f and Supplementary Fig. 1d**), accompanied by SIMBA barcode
plots (**Supplementary Fig. 1d**). As a certain feature (e.g., genes) might notably outnumber cells
or other features (when multiple types of features are present), SIMBA metrics not only serve
as an efficient way of ranking features based on their cell type specificity, but also provides a
straightforward way to filter out non-informative (non-cell-type-specific) features so that only
the embeddings of cells and informative features will be visualized and the SIMBA space will
not be crowded with non-informative features (e.g., house-keeping genes).

We next compared the top 600 marker genes identified by SIMBA (based on max value and Gini
index) with those identified by the clustering-based statistical-tests method implemented in
Scanpy (based on z-score calculated from the two-sided Wilcoxon rank-sum test with a
Benjamini-Hochberg p-value correction, one of the statistical tests recommended in Scanpy's
tutorial) (**Supplementary Fig. 2a**). Upon comparison, we observed that nearly half of the marker
genes discovered by SIMBA overlap with the marker genes identified by Scanpy
(**Supplementary Fig. 2a**). However, on inspection of the top non-overlapping marker genes,
genes identified by SIMBA are found to be enriched only within certain groups of cells
(**Supplementary Figs. 2b and 2c**) while genes identified by Scanpy but not by SIMBA include the
housekeeping gene *B2M* and multiple ribosomal protein genes (e.g., *RPS3* and *RPS6*) that are
expressed ubiquitously in all cell types (**Supplementary Figs. 2b and 2d**). Furthermore, a
combination of different statistical tests proposed in Scanpy is required to recover the genes

identified only by SIMBA. For example, IL7R was identified only by using the t-test and FCER1A
was identified only by using the Wilcoxon rank-sum test, as also noted in the Scanpy's tutorial,
while SIMBA successfully identified both *IL7R* and *FCER1A* as informative genes with a single
procedure and without clustering the cells (**Fig. 2e and Supplementary Fig. 1b**). These examples
illustrated some limitations of the clustering-based statistical-tests methods.

Lastly, we showed that SIMBA does not require variable gene selection, which is an essential
step in standard scRNA-seq pipelines such as Seurat or Scanpy. SIMBA produces very similar
embeddings for cells with and without variable gene selection (**Fig. 2b** and **Supplementary Fig.
2e**), though we observed that variable gene section does improve efficiency of the training
procedure.

SIMBA enables simultaneous characterization of cell states and cis-regulatory elements by
jointly modeling accessible sites and DNA sequences in scATAC-seq analysis
301
As one of the most popular single-cell epigenomic techniques, single-cell assay for transposase-
accessible chromatin using sequencing (scATAC-seq) has been widely used to profile regions of
open chromatin and identify functional *cis*-regulatory elements such as enhancers and active
promoters. In scATAC-seq, cells are characterized by different types of features [14], such as regions
of accessible chromatin ("peaks" or "bins") and *cis*-regulatory elements (DNA sequences) within
these accessible regions including transcription factor (TF) motifs or *k*-mers.
308
Unlike existing methods that can only use peaks/bins or the DNA sequence within them, SIMBA
can leverage simultaneously both types of features to learn cell states due to its flexibility in
graph construction. Also, as SIMBA encodes cell-feature or feature-feature relations into the
graph based on the simple binary presence of a feature, SIMBA does not need additional
normalization steps such as term frequency-inverse document frequency (TF-IDF), which is
required by most scATAC-seq analyses. When only peaks/bins are used, SIMBA constructs a graph
with nodes representing cells and chromatin regions (peaks or bins) and edges indicating the
accessibility of the chromatin regions in cells (**Fig. 3a**). When the DNA sequences for chromatin
regions are available, SIMBA can also encode DNA sequences including TF motifs and *k*-mers into
the graph by adding edges between these entities as nodes and the existing chromatin region
nodes. The edges in this case indicate the presence of TF motifs/*k*-mers within these chromatin-
accessible regions. Through the embedding procedure, SIMBA generates embeddings of cells
along with peaks and DNA sequences (**Methods**). Finally, either the partial SIMBA embeddings
(embeddings of cells in **Fig.3b**) or the whole SIMBA embeddings (embeddings of cells and all the
features in **Fig.3c**) can be visualized. Therefore, SIMBA enables dimensionality reduction by
leveraging both chromatin accessible regions and cis-regulatory sequences. Simultaneously, it
highlights the cell-type-specific open chromatin regions and regulatory DNA sequences in a
clustering-free way.
327
To demonstrate the value of SIMBA embeddings for scATAC-seq analysis, we first applied
SIMBA to a scATAC-seq data of 2,034 human hematopoietic cells with FACS-characterized cell
types[15](**Supplementary Table 1**). For the SIMBA embeddings of cells alone, as shown in **Fig. 3b**,

SIMBA accurately separated cells such that cells belonging to distinct cell types are visually
separated. For the SIMBA embeddings of cells together with various types of features, as shown
in **Fig. 3c**, SIMBA successfully embedded distinct features from both positional (peaks/bins) as
well as sequence-content (TF motifs and k-mers) information together based on their biological
relations. Notably, based on SIMBA metrics, these highlighted features that are embedded
within each cell type all have high cell-type specificity scores (shown in the upper right part of
SIMBA metric plots in **Figure 3d**).

339
340
Figure 3. Single-cell ATAC-seq analysis of the human hematopoiesis dataset using SIMBA.
(a) SIMBA graph construction and embedding in scATAC-seq analysis. Biological entities
including cells, peaks/bins, TF motifs, k-mers are represented as shapes and colored by
relevant cell types (green and orange). Non-informative features are colored dark grey.
Cells and chromatin accessible features (peaks / bins) are organized into a cell x peaks /
bins matrix. When sequence information (TF motif or *k*-mer sequence) within these
regions is available, they can be organized into two sub-matrices to associate a TF motif

or k-mer sequence with each peak/bin. These constructed feature matrices are then
binarized and assembled into a graph. When a single feature (chromatin accessibility) is
used, the graph encodes cells and peaks/bins as nodes. When multiple features (both
chromatin accessibility and DNA sequences) are used, this graph may then be extended
with the addition of TF motifs and k-mer sequences as nodes connected. Finally, SIMBA
embeddings of these entities are generated through a graph embedding procedure. **(b)**
UMAP visualization of SIMBA embeddings of cells colored by cell type. **(c)** UMAP
visualization of SIMBA embeddings of cells and features including TF motifs, k-mers, and
peaks. Cells are colored by cell type while motifs, k-mers, and peaks are colored green,
blue, and pink, respectively. Cell type specific features that are embedded near their
corresponding cell types are indicated as the text labels (colored according to feature type)
with arrows. **(d)** SIMBA metric plots of TF motifs, k-mers, and peaks. Cell-type specific
features annotated in **(c)** are highlighted. **€** Genomic tracks of aligned scATAC-seq
fragments, separated and colored by cell type. Two marker peaks P1 and P2 in red are
shown beneath the alignment. Within the peak P1, k-mer GATAAG and its resembling
GATA1 motif logo are highlighted. **(f)** UMAP visualization of SIMBA embeddings of cells
colored by TF activity scores of the GATA1 motif and k-mer GATAAG enrichment. **(g)**
SIMBA barcode plots of the GATA1 motif, k-mer GATAAG, and the two peaks P1 and P2.
Cells are colored according to cell type labels described above. Dotted red line indicates
the same cutoff used in all four plots.

Our analysis using SIMBA led to several key findings in human hematopoietic differentiation.

First, SIMBA identified key master regulators of hematopoiesis. As highlighted in **Fig. 3c**, we
observed that motifs of previously reported TFs were embedded near their respective cell types
in the UMAP plot. For example, the GATA1 and GATA3 motifs are proximal to megakaryocyte-
erythroid progenitor (MEP) cells[16], the PAX5 and EBF1 motifs are near to common lymphoid
progenitor (CLP) cells[17], and the CEBPB and CEBPD motifs are proximal to monocyte (mono)
population[18].

Second, SIMBA revealed an unbiased set of DNA sequences, i.e., *k*-mers, that represent
important TF binding motifs involved in hematopoiesis. We observed that these *k*-mers were
embedded near their resembling TF binding motifs and relevant cell subpopulations (**Fig. 3c** and
3e, **Supplementary Fig. 3b**), indicating that this methodological framework is capable of *de
novo* motif discovery. For example, the DNA sequence, CAGCTG is embedded in plasmacytoid
dendritic cells (pDCs); this sequence matches the TCF12 binding motif, which controls dendritic
cell lineage specification. To further illustrate the interpretability of the SIMBA embeddings of
TF motifs and k-mers, we calculated per-cell TF activity scores[19] (high-variance TF motifs/*k*-
mers) and visualized them on SIMBA embeddings of cells. As shown in **Figure 3f**, the GATA1 TF
motif and k-mer GATAAG that were both embedded in MEP cells by SIMBA, also showed high-
level activity in MEP cells. The consistency between SIMBA embedding and TF activity was
observed for most of other TF motifs and *k*-mers as well (**Supplementary Fig. 3a, 3b**).

Third, SIMBA identified differentially accessible chromatin regions that may mediate cell-type
specific gene regulation. For example, the two peaks with coordinates chr19:12997999-
12998154 (P1) and chr19:12998329-12998592 (P2) that were embedded within MEP cells were
almost exclusively observed in MEP cells on KLF1 genome track (**Fig. 3e**). Interestingly, P1,
upstream of *KLF1*, contains the *k*-mer GATAAG that matches the GATA1 binding motif, while
transcription factor GATA1 is known to regulate the gene *KLF1* and plays a pivotal role in
erythroid cell and megakaryocyte development[20]. Therefore, by embedding these MEP-cell-
related regulatory elements into the neighborhood of MEP cells, SIMBA demonstrates a novel
means of studying the epigenetic landscape of cell differentiation. To further validate the
differentially accessible regions identified by SIMBA, we selected 100 peaks at random from
each annotated cell type in SIMBA co-embedding space. From the heatmap of chromatin
accessibility, we clearly see that the peaks embedded nearby respective types correlate with
strong cell-type specificity. This observation is robust to the number of cells within each cell
type (**Supplementary Fig. 3c**).
405
Available methods for scATAC-seq analysis visualize only cells. While SIMBA diverges from these
available workflows, enabling the co-embedding of cells and features, we still qualitatively and
quantitatively compared the SIMBA embeddings of cells to state-of-the-art scATAC-seq analysis
methods by their ability to distinguish cell types. Our analyses show that SIMBA overall
performs better than the methods evaluated, further demonstrating the wide utility of SIMBA
(**Supplementary Figs. 4 and 5; Supplementary Note 1**).
412
413
414   **SIMBA enables simultaneous learning of cellular heterogeneity and gene regulatory circuits**
415   **from integrated analysis of single-cell multimodal data**
416
scRNA-seq and scATAC-seq are two of the most widely adopted single-cell sequencing
technologies, but they are limited to measuring only a single aspect of cell state at a time. To
improve our ability to interrogate cell states, several single-cell dual-omics technologies have
been recently developed [21-24] to jointly profile transcriptome and chromatin accessibility within
the same individual cells, therefore providing the potential to correlate gene expression with
accessible regulatory elements and further delineate the yet elusive principles of gene
regulation. This section outlines the SIMBA's ability to simultaneously learn cell heterogeneity
as well as gene regulatory circuits from single-cell multiomic data. We applied SIMBA to three
recent single-cell dual-omics technologies: SHARE-seq[22], SNARE-seq[21], and a multiome PBMCs
dataset from 10x Genomics (**Supplementary Table 1**).
Figure 4a illustrates the procedure of graph construction and generation of the final SIMBA
embedding matrix. Briefly, for scRNA-seq, the gene expression matrix is discretized to generate
different levels of gene expression. For scATAC-seq, both the chromatin accessibility matrix and
motif/*k*-mer match matrix are binarized. In this graph, there are five entity (node) types,
including cells, genes, peaks, motifs, and *k*-mers. For scRNA-seq, an edge indicates whether a
gene is expressed in a cell and its weight indicates the gene expression level (five levels, by
default). For scATAC-seq, an edge indicates whether a peak is present in a cell or if a TF motif/*k*-

mer is present within a peak. Once the graph is constructed, the graph embedding procedure is
performed to generate SIMBA embeddings of cells and different types of features. scATAC-seq
peaks generally greatly outnumber cells and other features and many of these peaks are non-
informative, resulting in them dominating the space if the whole SIMBA embeddings are
visualized (**Supplementary Fig. 6a, c**). In such cases, we leverage the flexibility of SIMBA
embedding to only visualize the partial SIMBA embeddings to improve the visibility of cells and
cell-type-specific features.

Figure 4. Multimodal analysis of the SHARE-seq hair follicle dataset using SIMBA. **(a)**
SIMBA graph construction and embedding in multimodal analysis. Overview of SIMBA's
approach to multimodal (scRNA-seq + scATAC-seq) data analysis. **(b)** SIMBA metric plots
of genes, TF motifs, and peaks. All these features are plotted according to the Gini index
against max score. Cell-type specific genes, TF motifs, and peaks are highlighted. **(c)** UMAP
visualization of SIMBA embeddings of cells (Top-left), cells and genes (Top-right), and cells
along with genes, TF motifs, and peaks (Bottom). **(d)** Ranked scatter plot of candidate
master regulators as identified by SIMBA. **€** Schematic description of SIMBA's strategy for
identifying target genes given a master regulator. **(f)** Top 30 target genes of transcription
factors Lef1 and Hoxc13 as inferred by SIMBA.

To demonstrate the usefulness and versatility of the SIMBA embeddings, we analyzed
the cell populations undergoing the dynamic process of hair follicle differentiation from
mouse skin profiled with SHARE-seq.

First, we calculated SIMBA metrics (max values and Gini index scores) to assess the cell-type
specificity of different types of features, including genes, TF motifs, and peaks (**Fig. 4b,**
Methods). As shown in **Figure 4b**, based on these metrics, we successfully recovered genes
associated with hair follicles such as *Lef1* and *Hoxc13*. Similarly, TF motifs and peaks proximal to
the genomic loci of these genes also score in the upper right quadrant of the metric plots.
SIMBA's cell-type specificity metrics successfully revealed the key genes and regulatory factors
important to the hair follicle differentiation process.

Next, we visualized and interrogated the SIMBA embeddings of 1) cells; 2) cells and top-ranked
genes based on SIMBA metrics; and 3) cells, top-ranked genes and TF motifs based on SIMBA
metrics, and the neighboring peaks of these genes and TF motifs by querying the SIMBA space
(**Methods**). **Figure 4c** shows the UMAP visualization of the partial SIMBA embeddings of cells
and informative features. The UMAP visualization of SIMBA embeddings of cells and the full set
of features was also performed (**Supplementary Fig. 6a**).

The SIMBA embeddings of cells were able to reveal the three fate decisions from transit-
amplifying cells (TACs), including inner root sheath (IRS), medulla, and cuticle/cortex. The
SIMBA embeddings of cells and informative features uncovered important genes and regulatory
factors along the hair follicle differentiation trajectories. For example, the marker genes *Krt71*,
Krt31, and *Foxq1* were embedded into their corresponding cell types: IRS, cuticle/cortex, and
medulla, respectively. The Lef1 motif was embedded into the beginning of medulla and
cuticle/cortex lineages while the Hoxc13 motif was embedded into the late stage of
cuticle/cortex differentiation. Peaks near the *Lef1* and *Hoxc13* loci were also embedded into the
nearby regions of these genes and motifs, as expected.

To show the robustness of SIMBA, we separated the scRNA-seq and scATAC-seq components
within the SHARE-seq dataset and performed each respective single-modality analysis. With the
consistent embedding results of cells as in multimodal analysis, we further demonstrated that
SIMBA embedding procedure is robust to the type and the number of features encoded in the
input graph (**Supplementary Fig. 6b,6c**). Each reported marker gene was corroborated using
the UMAP plots with cells colored by gene expression as well as using the SIMBA barcode plots.
The two aforementioned TF motifs and their respective peak sets were supported by the
corresponding SIMBA barcode plots, wherein we observed an imbalanced distribution with high
probability towards the correct cell type labels (**Supplementary Fig. 7a-d**).

Further, we demonstrated that the SIMBA co-embedding space of cells and features provides
the potential to identify master regulators of differentiation and infer their target regulatory
genes. To define a master regulator *a priori*, we postulated that both its TF motif and TF gene

should be cell-type specific, given that active gene regulation involves both the expression of a
TF and accessibility of its binding sites. Thus, the TF motif and TF gene should be embedded
closely in the shared latent space. Extending this logic to identify putative master regulators, we
assessed the cell-type-specificity of TF motifs and genes based on SIMBA metrics and ranked all
potential master regulators based on the distance between the TF motif and the respective TF
gene in the shared SIMBA embedding space (**Methods**). SIMBA successfully identified
previously described master regulators such as Lef1, Gata6, Nfatc1, and Hoxc13 as the top
master regulators related to lineage commitment in mouse skin (**Fig. 4d, Supplementary Table
2**). To infer the target genes of a given master regulator, we postulate that in the shared SIMBA
embedding space, 1) the target gene is close to both the TF motif and the TF gene; 2) the
accessible regions (peaks) near the target gene loci must be close to both the TF motif and the
target TF gene. Resting on these assumptions of *cis*-regulatory dynamics, the inference of target
genes was performed by calculating the distance between target gene candidates and the
respective TF motif and gene. In addition, nearby peaks around the target gene's locus and the
presence of TF motif in these nearby peaks are also considered (**Fig. 4e, Methods**). The top 30
target genes of TF Lef1 and TF Hoxc13 inferred by SIMBA are shown respectively (**Fig. 4f,
Supplementary Fig. 7e**). The full list of ranked target genes is provided in **Supplementary Table
3**. Notably, our approach recovered targets genes that were also reported in the original
study[22]. For example, genes *Lef1*, *Jag1*, *Hoxc13*, *Gtf2ird1* are regulated by the TF Lef1, while
genes *Cybrd1*, *Hoxc13*, *St14* are regulated by the TF Hoxc13.

In addition to SHARE-seq, we also applied SIMBA to another two dual-omics datasets, the
mouse cerebral cortex dataset profiled by SNARE-seq[21] (**Supplementary Fig. 8**) and the
multiome PBMCs dataset from 10x Genomics (**Supplementary Fig. 9**). By validating the
embeddings of cells and features with given cell type labels (**Supplementary Fig. 8a and Fig.9a**),
marker genes from the original study (**Supplementary Fig. 8a,b,d and Fig. 9a,b,d**), and
differentially accessible chromatin regions (**Supplementary Fig. 8c and Fig. 9c**), we further
demonstrate the suitability of SIMBA for multimodal analysis.

517
518   **SIMBA enables simultaneous batch correction and clustering-free marker gene detection**
519
Efforts to collect data from single cells has grown to the level of consortia that span multiple
institutions with the hopes of finely mapping and characterizing specific tissues. This has
brought with it an increased demand for analysis methods that are capable of negating
technical covariates inherent to multi-batch data collection, including experimental replicate
identity, sample preparation, and sequencing platform. Batch correction that removes the
effects of technical covariation while preserving true biological signals is required prior to
downstream analysis [25, 26]. Existing methods follow a workflow with four primary steps. The first
step is the actual batch correction, which often generates a "batch corrected" latent space. The
second step clusters cells in this batch corrected space. Based on the clustering result the third
step detects marker genes in the original gene expression space of each batch because the low-
dimensional "batch corrected" space is no longer comprised of genes. The fourth step finally
combines the marker genes detected from each batch. However, these methods are clustering-

dependent and may result in the inconsistent explanation of marker genes as marker genes are
detected in each original batch as opposed to the batch-corrected space. Unlike current
methods, in addition to embeddings of cells, SIMBA generates comparable embeddings of
genes and therefore relieves marker gene discovery from a dependence on the original gene
expression space. Thus, SIMBA enables simultaneous batch effect removal and cell-type-
specific marker gene detection in the same integrated space without clustering.
538

539
540
Figure 5. Batch correction analysis of scRNA-seq data using SIMBA. **(a)** SIMBA graph
construction and embedding in batch correction analysis. Overview of SIMBA's approach
to batch correction across scRNA-seq datasets. Distinct shapes indicate the type of entity
(cell or gene). Colors distinguish batches or cell types. **(b)** UMAP visualization of the
scRNA-seq human pancreas dataset with five batches of different studies before and after
batch correction. Cells are colored by scRNA-seq data source and cell type respectively.
Top: UMAP visualization before batch correction; Bottom: UMAP visualization after batch
correction with SIMBA; **(c)** UMAP visualization of SIMBA embeddings of cells and genes,
with batch effect removed and known marker genes highlighted.

550
We first demonstrate that SIMBA readily corrects batch effects and produces joint embeddings
of cells and genes across multiple scRNA-seq datasets generated from varying sequencing

platforms and cell type compositions. While existing methods for scRNA-seq analysis rely on
specialized tools for batch correction, SIMBA works as a stand-alone method obviating the need
for prior input data correction when applied to multi-batch scRNA-seq dataset. SIMBA
accomplishes batch correction by encoding multiple scRNA-seq datasets into a single graph (**Fig.**
5a). Cell nodes across batches are connected to gene nodes through experimentally measured
edges as in the previously described scRNA-seq graph construction. Here, the gene nodes are
shared between the cell nodes of different batches. In addition to the experimentally measured
edges, batch correction is further enhanced through computationally inferred edges drawn
between similar cell nodes across datasets using a truncated randomized singular value
decomposition (SVD)-based procedure. SIMBA then generates the embeddings of all nodes
including cells of each batch and genes from the resulting graph (**Methods**). The SIMBA
embeddings of cells naturally represent the batch-corrected space. In addition, the whole
SIMBA embeddings of all entities provide the batch-corrected space, in which cells and genes
co-exist, and therefore allow for individual-cell-level marker detection by performing biological
queries of cells in the SIMBA space (**Methods**). We visualized both SIMBA embeddings of cells
(**Fig. 5b**), and the whole SIMBA embeddings of cells and genes (**Fig. 5c**) in UMAP.
569
We applied SIMBA to two multi-batch scRNA-seq datasets; a mouse atlas dataset composed of
two batches and a human pancreas dataset spanning five batches used in a recent benchmark
study[25] (**Supplementary Table 1**). The mouse atlas dataset contains two scRNA-seq datasets
with shared cell types from different sequencing platform. The human pancreas dataset
contains five samples pooled from five sources using four different sequencing techniques, in
which not all cell types are shared across each sample. For both datasets, SIMBA successfully
corrected batch effects, evenly mixing batches within annotated cell type clusters, while
maintaining the segregation of these clusters in the resulting embedding, indicating
preservation of biological signal and elimination of confounding technical covariates (**Fig. 5b,**
Supplementary Fig. 12b). It is important to note that the mouse atlas dataset was collected
from nine different organ systems, so there exists some expected heterogeneity within cell type
labels. Conversely, the human pancreas datasets are curated from a single organ and SIMBA
sufficiently separated cell types into transcriptionally distinct, homogeneous cell clusters (**Fig.**
5b).
584
Through removing batch effects during graph embedding, SIMBA simultaneously identifies cell-
type-specific marker genes (**Fig. 5c**). In the absence of the eliminated technical covariation,
marker genes are identifiable by performing biological queries for neighboring genes within cell
types in the SIMBA embeddings of cells and genes (**Methods**). In the case of unknown cell
labels, marker genes can be identified by calculating SIMBA metrics (**Methods**). SIMBA correctly
embeds known cell-type-specific marker genes proximal to the correct cell type labels, while
non-marker genes were non-proximal to specifically-labelled cells (**Supplementary Fig. 10, 11**).
The resulting marker genes recapitulated the clustering-based differential expression (DE)
analysis results for each dataset[27-32] (e.g. *Cdh5*, *Tie1*, *Myct1* for endothelial cell and *C1qc*, Fcgr1
for macrophage, *S100a8*, *Trem3* for Neutrophil in the mouse atlas dataset and *KIF12* for alpha
cell and *KRT19* for ductal cell in the human pancreas dataset) and are shown to be expressed
specifically in the queried cell types (**Supplementary Fig. 10, 11**). Taken together, these results

distinguish SIMBA from existing batch correction methods that rely on clustering in a batch-
corrected space, followed by differential gene expression analysis in the original, uncorrected
space of each batch.
600
While SIMBA is a versatile graph embedding method that can perform multiple tasks and
generate embeddings of both cells and genes, we evaluated the SIMBA embeddings of cells for
this task with methods that were specifically designed for batch correction. We considered
three widely adopted batch correction methods that demonstrated top-tier performance based
on a recent benchmark study[25]: Seurat3, LIGER and Harmony. Our results indicate that SIMBA
achieved comparable batch correction performance both qualitatively and quantitatively while
enabling simultaneous marker gene detection by providing the additional SIMBA embeddings
of genes. (**Supplementary Note 2, Supplementary Figure 12**).
609
610   **SIMBA enables simultaneous multi-omics integration and clustering-free multi-type marker**
611   **feature detection**
612
Single-cell assays are now capable of measuring a broad range of cellular modalities and data is
being generated that describes cells by varying features sets, which has motivated the need for
methods that leverage these features to perform multi-omics integration such that a more
comprehensive description of cell state may be learned. This is different from multi-modal
analysis because the correspondence between individual cells is unknown. Current multi-omics
integration methods follow a similar workflow as the previously described batch correction
methods, including: 1) generating a low-dimensional integrated space of cells; 2) clustering cells
in the integrated space; 3) detecting marker features in the original feature (e.g., genes, peaks)
space of each modality because the low-dimensional integrated space no longer consists of the
original features. Unlike existing multi-omics integration methods that cannot directly explore
multi-type features in the integrated space and require clustering for identifying marker
features, we demonstrate that SIMBA enables simultaneous multi-omics integration and
clustering-free detection of distinct marker features, specifically as it is applied to datasets
comprised of scRNA-seq and scATAC-seq.
627
SIMBA accomplishes this integration by first building one graph for scRNA-seq data and another
graph for scATAC-seq data independently as described in previous sections (**Fig. 6a**). To connect
these two graphs, SIMBA then calculates gene activity scores by summarizing accessible regions
from scATAC-seq data and then infers edges between cells of different assays based on their
shared gene expression modules as previously described in the batch correction section. Finally,
SIMBA embeds the graph of cells, genes, and peaks into a common, low-dimensional space. The
SIMBA embeddings of cells naturally represent the integrated space of multiple modalities.
Furthermore, the SIMBA embeddings of all entities provide the integrated space containing cell,
genes, and peaks, and therefore enable the individual-cell-level marker detection of multi-type
features by performing biological queries of cells in SIMBA space (**Methods**). The SIMBA
embeddings of these multi-omics entities can be visualized either partially or as a whole using
UMAP or similar visualization tools.
640

641
Figure 6. Multi-omics integration of scRNA-seq + scATAC-seq data using SIMBA. **(a)** SIMBA
graph construction and embedding in multi-omics integration. Overview of SIMBA's
approach to data integration across scRNA-seq and scATAC-seq. Distinct shapes indicate
the type of entity (cell, gene, or peak). Colors distinguish batches or cell types. **(b)** UMAP
visualization of the integrated scRNA-seq and scATAC-seq data manually created from the
10x human PBMCs dataset before and after data integration. Cells are colored by single-
cell modality and cell type respectively. Top: UMAP visualization before integration;
Bottom: UMAP visualization after integration with SIMBA. **(c)** UMAP visualization of
SIMBA embeddings of cells, genes, and peaks with two cell modalities integrated and
known marker genes highlighted.

652
To facilitate the evaluation of data integration performance, we created datasets with ground-
truth labels by manually splitting the dual-omics datasets into two single-modality datasets (i.e.,
scRNA-seq and scATAC-seq), in which we know the true matching between cells across the two
modalities. We then applied SIMBA to the integration analysis of two case studies where
scRNA-seq and scATAC-seq datasets are generated from the SHARE-seq mouse skin dataset and
the 10x Genomics multiome human PBMCs dataset, respectively (**Supplementary Table 1**).
659

We first visualized the SIMBA embeddings of cells and observed that SIMBA was able to
preserve cellular heterogeneity while evenly mixing the two modalities (**Fig. 6b, Supplementary**
Fig. 15b). We then visualized the SIMBA embeddings of cells, genes, and top-ranked peaks
based on SIMBA metrics and observed that in addition to learning cellular heterogeneity,
SIMBA simultaneously identified marker genes and peaks at single-cell resolution. In the co-
embedding space, we observed that the neighbor genes of cells (highlighted in UMAP plots),
are each exclusively expressed in their corresponding cell types (**Supplementary Figs. 13a-e,**
14a-c,e). For example, in the SHARE-seq mouse skin dataset, *Foxq1* and *Shh* are located within
medulla and TAC-2, respectively; in the 10x PBMCs dataset, *PAPSS2* and *KCNMA1*, which are
the marker genes of blood monocytes, are embedded close to each other. Similarly, we
observed that the neighbor peaks of cells show a clear cell-type-specific accessibility pattern
that is robust to the cluster size of a given cell type (**Supplementary Figs. 13f and 14d).**
672
The joint embedding of cells and features produced by SIMBA is fundamentally distinguished
from other multi-omics integration methods in that it simultaneously achieves integration as
well as marker feature discovery. However, we still sought to compare the SIMBA embeddings
of cells with two widely-adopted single-cell multi-omics integration methods, Seurat3 and
LIGER, based on their ability to integrate single-cell modalities while persevering cellular
heterogeneity (**Supplementary Note 3**). We observed that SIMBA achieved the overall best
performance on the mouse skin SHARE-seq dataset and 10x PBMCs multiome dataset.
680

Discussion

682
683   Multimodal measurements of individual cells offer new and unexplored opportunities to study
684   cell identity as a function of the complex interactions between omic layers. While these
685   datasets offer an exciting potential for discovery, computational analysis methods to fully
686   delineate the cell states and molecular processes across multiple genomic features remain
687   insufficient.
688
As presented in this manuscript, SIMBA models cells and measured features as nodes encoded
in a graph and employs a scalable and efficient graph embedding procedure to embed nodes of
cells and features into a shared latent space. We demonstrate that direct graph representations
of single-cell data capture not only the relations between cells and the quantified features of
the experiment (e.g., gene expression or chromatin accessibility) but also hierarchical relations
between features. An example of such a hierarchical relation is the coordinate-level description
of an ATAC-seq peak and the corresponding TF motifs and/or *k*-mer sequences contained
within that region. In the resulting joint embedding, proximity-based biological queries can be
performed to discover cell-type-specific co-regulatory machinery across modalities. Therefore,
SIMBA enables simultaneous learning of cellular heterogeneity and cell-type-specific
multimodal features and complements the current gene regulatory network analyses. SIMBA
also circumvents the ordinary reliance on cell clustering for cell sub-population feature
discovery. We thus avoid user-defined clustering solutions, which may lead to artifactual
discovery or false negative results.
703

704 SIMBA has been extensively benchmarked across single-cell modalities and tasks, obtaining
705 better or comparable performance metrics when compared to current state-of-the-art methods
706 developed for the respective task. In contrast to tools developed and optimized for a single,
707 specific task these results suggest a wide applicability of SIMBA's graph-based framework,
708 obviating the need to combine multiple analysis tools.
709
710 Graph embedding methods hold significant promise for the analysis of biological data. Previous
711 applications of graph embedding include functional annotation of genes [33], transcription factor
712 binding to DNA motifs [34] and more recent single-cell RNA-seq analyses [35, 36]. The graph encoding
713 and embedding procedures we have outlined may be potentially improved and extended to
714 better represent biological entities and capture their respective relations.
715
Foreseeable extensions of SIMBA may include the analyses of increasingly complex datasets.
For example, in the analysis of spatial transcriptomics wherein transcriptomic measurements
are mapped to the true cell coordinates within a tissue [37], we can encode the spatial proximity
into a SIMBA graph. We also envision extending this framework to data describing 3-D
chromatin conformation wherein the interaction between DNA segments can be encoded to
represent how regulatory regions are linked to genes[38]. Another potential extension of SIMBA
could consider single-cell lineage-tracing datasets[39] wherein both cellular lineage information
and gene expression measurements are captured and can be potentially encoded into a SIMBA
graph to represent their longitudinal relations. In general, we are interested in the further
incorporation of external information and hierarchical relations between features in the graph.
We anticipate our comprehensive and extensible SIMBA framework (https://simba-
bio.readthedocs.io/) will provide the possibility to leverage *a priori* knowledge for graph
embedding and the flexibility to extend to new experimental designs.
729
It is likely that multi-omics assays will continue to improve as well as expand in scope. Already,
innovation in these data-generating technologies have outpaced the development of
corresponding computational frameworks required to gain integrative insights from such rich
data. This disparity highlights a need for methods that break through previous limitations and
are easily extended to future cell measurements. We believe SIMBA satisfies these conditions
as a comprehensive and extensible method for exploring cellular heterogeneity and
investigating the regulatory mechanisms that drive cellular diversity while laying a groundwork
for the development of new non-cluster-centric analysis methods for single cell omics data.
738
739

Methods

Single-cell data preprocessing

a. Single-cell RNA-seq
Genes expressed in fewer than three cells were filtered.   Raw counts were library size-
normalized and subsequently log-transformed. Optionally, variable gene selection [12] (a
python version is implemented in SIMBA that is inspired by Scanpy[2]) may be performed

to remove non-informative genes and accelerate the training procedure. Notable
differences in the resulting cell embeddings were not observed upon limiting feature
input to those identified by variable gene selection but SIMBA embeddings of non-
variable genes will not be generated as they are not encoded in the graph.
752
b. Single-cell ATAC-seq
Peaks present in fewer than three cells were filtered. Optionally, we implemented a
scalable truncated-SVD-based procedure to select variable peaks as a preliminary step
to additionally filter non-informative peaks and accelerate the training procedure. First
the top k principal components (PCs) were selected, with k chosen based on the elbow
plot of variance ratio. Then for each of the top k PCs, peaks were automatically selected
based on the loadings using a knee point detection algorithm implemented by 'kneed'[40].
Finally, peaks selected for each PC were combined and denoted as "variable peaks".
Similar to the observation made with scRNA-seq data, the optional step of variable peak
selection has a negligible effect on the resulting cell embedding. Despite this minimal
impact on the resulting embedding, this feature selection step imparts a significant
practical advantage in reducing training procedure time.
765
766   k-mer and motif scanning was performed using packages 'Biostrings' and 'motifmatchr'
767   with JASPAR2020[41]. Included in the implementation of SIMBA is a convenient R
768   command line script "scan_for_kmers_motifs.R" , which will convert a list of peaks
769   (formatted in a bed file) to a sparse peaks-by-k-mers/motifs matrix, which is stored as
770   an hdf5-formated file.
771
Graph construction (five scenarios)
773
i. Single-cell RNA-seq analysis
The distribution of non-zero values in the normalized gene expression matrix was first
approximated using a k-means clustering-based procedure. First, the continuous non-
zero values were binned into n intervals (by default n=5). Bin widths were defined using
1-dimensional k-means clustering wherein the values in each bin are assigned to the
same cluster center. The continuous matrix is then converted into a discrete matrix
wherein1, ... , n are used to denote n levels of gene expression. Zero values are retained
in this matrix. Then the graph was constructed by encoding two types of entities, cells
and genes, as nodes and relations with n different weights between them, i.e., n levels
of gene expression, as edges. These n relation weights range from 1.0 to 5.0 with a step
size of $5/n$ denoting gene expression levels (lowest: 1.0, highest: 5.0), such that edges
corresponding to high expression levels affect embeddings more strongly than those
with intermediate or low expression levels. This discretization is implemented in the
SIMBA package using the function, "si.tl.discretize()".
788
789   ii. Single-cell ATAC-seq analysis
790   Peak-by-cell matrices were binarized, with "1" indicating at least one read within a peak
791   and "0" otherwise. The graph was constructed by encoding two types of entities, cells

792	and peaks, as nodes and the relation between them, denoting the presence of a given
793	peak in a cell, as edges. The single relation type was assigned with a weight of 1.0. When
794	the DNA sequence features were available, they were encoded into the graph using k-
795	mer and motif sequence entities as nodes. This was performed by first binarizing the
796	peak-by-k-mer/motif matrix then constructing an extension to the original peak/cell
797	graph using the peaks, k-mers, and motifs as nodes and the presence of these entities
798	within peaks as edges between these additional nodes and the peak nodes. The relation
799	between k-mers and peaks was assigned a weight of 0.02 while the relation between TF
800	motifs was assigned a weight of 0.2. Of note, k-mers and motifs may be used
801	independently of each other as node inputs to the graph, depending on the specific
802	analysis task.
803	
804	iii. Multimodal analysis
805	Combination of the above outlined strategies for graph construction of scRNA-seq and
806	scATAC-seq data was used to construct a multi-omics graph.
807	
808	iv. Batch correction
809	A graph for each batch was constructed as described in i). Edges between cells of
810	different batches were inferred through a procedure based on truncated randomized
811	singular value decomposition (SVD) to link disjoint graphs of different batches. More
812	specifically, in the case of scRNA-seq data, consider two gene expression matrices
813	$X1_{n_1 \times m}$ and $X2_{n_2 \times m}$, where n_1, n_2 denotes the number of cells and m denotes the
814	number of the shared features, i.e., variable genes, between datasets. The matrix
815	$X_{n_1 \times n_2}$ was then computed by multiplying $X1$ and $X2$:
816	
817	

$$X = X1 \times X2^T$$

818	
819	Truncated randomized SVD was subsequently performed on X:
820	
821	

$$X \approx U \times \Sigma \times V^T$$

822	
823	where U is an $n_1 \times d$ matrix, Σ is an $d \times d$ matrix, and V is an $n_2 \times d$ matrix (by
824	default $d = 20$).
825	
826	Both U and V were further $L2$ normalized. For each cell in U, we searched for k
827	nearest neighbors in V and vice versa (by default, $k = 20$). Eventually, only the mutual
828	nearest neighbors between U and V were retained as inferred edges between cells
829	(represented as dashed lines in **Fig. 5a**). The procedure of inferring edges between
830	cells of different batches is implemented in the function "si.tl.infer_edges()" in the
831	SIMBA package.
832	
833	For multiple batches, SIMBA can flexibly infer edges between any pair of datasets. In
834	practice, however edges are inferred between the largest dataset(s) or the dataset(s)
835	containing the most complete set of expected cell types and other datasets.

v. Multi-omics integration

scRNA-seq and scATAC-seq graphs were constructed following steps i) and ii), respectively. To infer the edges between cells of scRNA-seq and scATAC-seq, gene activity scores were first calculated for scATAC-seq data[3]. More specifically, for each gene, peaks within 100kb upstream and downstream of the TSS were considered. Peaks overlapping gene body region or within 5kb upstream of gene bodies were given the weight of 1.0. Otherwise, peaks were weighted based on their distances to TSS using the exponential decay function: $e^{\frac{-distance}{5000}}$. Subsequently, the gene score of each gene was computed as a weighted sum of the considered peaks. These gene scores were then scaled to respective gene size. These steps are implemented by the function "si.tl.gene_scores()" in SIMBA. For user convenience, the SIMBA package curates the gene annotations of several commonly used reference genomes, including hg19, hg38, mm9, and mm10. Once gene scores were obtained, the same procedure described in iv) was performed to infer edges between cells profiled by scRNA-seq and scATAC-seq using the function, "si.tl.infer_edges()" in SIMBA.

The procedure of generating constructed graphs is implemented in the function, "si.tl.gen_graph()" in the SIMBA package.

Graph Embeddings with Type Constraints

Following the construction of a multi-relational graph between biological entities, we adapted graph embedding techniques from the knowledge graph and recommendation systems literature to construct unsupervised representations for these entities.

We provide as input a directed graph $G = (V, E)$, where V is a set of entities (vertices) and E is a set of edges, with a generic edge $e = (u, v)$ between a source entity u and destination entity v. We further assume that each entity has a distinct known type (e.g., cell, peak, etc.).

Graph embedding methods learn a D-dimensional embedding vector for each $v \in V$ by optimizing a link prediction objective via stochastic gradient descent, with D=50 used for our experiments. We will denote the full embedding matrix as $\theta \in R^{|V| \times D}$ and the embedding for an entity v as θ_v.

For an edge $e = (u, v)$, we denote $s_e = \theta_u * \theta_v$ as the score for e, and optimize a multi-class log loss

$$\mathcal{L} = -log\frac{\exp{(s_e)}}{\sum_{e' \in \mathcal{N}} \exp{(s'_e)}}$$

876   Where $\mathcal{N}$ is a set of "negative sampled" candidate edges generated by corrupting e [42].
877   This log loss objective attempts to maximize the score for all $(u, v) \in E$ and minimize it
878   for $(u, v) \notin E$.
879

Negative samples are constructed by replacing either the source or target entity in the
target edge $e = (u, v)$ with a randomly sampled entity. However, in graphs like ours
where only edges between certain entity types are possible, previous work has shown
that it is beneficial to optimize the loss only over candidate edges that satisfy the type
constraints[43]. Thus, for e.g., a cell-peak edge we only sample negative candidates
between cell and peak entities. This modification is crucial in our setting since most
randomly selected edges will be of invalid type (e.g., peak-peak), forcing the
embeddings to primarily be optimized for irrelevant tasks (e.g., having low dot product
between every pair of peaks).
889

Furthermore, it has been frequently observed that in graphs with wide distribution of
node degrees, it is advantageous to sample negatives proportional to some function of
the node degree to produce more informative embeddings that don't merely capture
the degree distribution [13, 44]. For each graph edge in the dataset encountered in a
training batch, we produce 100 negatives by corrupting the edge with a source or
destination sampled uniformly from the nodes with the correct types for this relation
and 100 by corrupting the edge with a source or destination node sampled with
probability proportional to its degree[13].
898

899   As with many ML methods, graph embeddings are prone to overfitting in a low-data
900   regime (i.e., low ratio of edges to parameters). We observed overfitting measurable as a
901   gap between training and validation loss on the link prediction task, which we addressed
902   with $L2$ regularization on the embeddings θ,
903

904
$$\mathcal{L}_{reg} = \mathcal{L} + \lambda \sum_{u \in N} \sum_{d=1}^{D} \theta_{ud}^2.$$

905
906   with $\lambda = wd * wd\_interval$. For weight decay parameter (wd), by default it is calculated
907   automatically as $\frac{C}{N_e}$, where N_e is the training sample size (i.e., the total number of edges)
908   and C is a constant. For weight decay interval $(wd\_interval)$, we set it to 50 for all
909   experiments.
910

911   We use the PyTorch-BigGraph framework, which provides efficient computation of
912   multi-relation graph embeddings over multiple entity types and can scale to graphs with
913   millions or billions of entities[13]. For 1.3 million cells, the PyTorch-BigGraph training itself
914   takes only ~ 1.5 hours using 12 cores without the requirement of GPU (https://simba-
915   bio.readthedocs.io/en/latest/rna_10x_mouse_brain_1p3M.html).
916

The resulting graph embeddings have two desirable properties that we will take
advantage of:
1.   First-order similarity: for two entity types T_1, T_2 with a relation between them,
edges with high likelihood should have higher dot product; specifically, for any $u \in T_1$,
the predicted probability distribution over edges to T_2 originating from u is
approximated as $\frac{e^{x_u * x_v}}{\sum_{v\prime \in T_2} e^{x_u * x_{v\prime}}}$.
2.   Second-order similarity: within a single entity type, entities that have 'similar
contexts', i.e., a similar distribution of edge probabilities, should have similar
embeddings. Thus, the embeddings of each entity type provide a low-rank latent space
that encodes the similarity of those entities' edge distributions.
927

928 **Evaluation of the model during training**
929

During the PyTorch-BigGraph training procedure, a small percent of edges is held out
(by default, the evaluation fraction is set to 5%) to monitor overfitting and evaluate the
final model. Five metrics are computed on the reserved set of edges, including mean
reciprocal rank (MRR, the average of the reciprocal of the ranks of all positives), R1 (the
fraction of positives that rank better than all their negatives, i.e., have a rank of 1), R10
(the fraction of positives that rank in the top 10 among their negatives), R50 (the
fraction of positives that rank in the top 50 among their negatives), and AUC (Area
Under the Curve). By default, we show MRR along with training loss and validation loss
while other metrics are also available in SIMBA package (**Supplementary Fig. 1a**). The
learning curves for validation loss and these metrics can be used to determine when
training has completed. The relative values of training and validation loss along with
these evaluation metrics can be used to identify issues with training (underfitting vs
overfitting) and tune the hyperparameters weight decay, embedding dimension, and
number of training epochs appropriately. For example, in **Supplementary Figure 1a**
training can be stopped once the validation loss plateaus. However, for most datasets
we find that the default parameters do not need tuning.
946

947 **Softmax transformation**
948

PyTorch-BigGraph training provides initial embeddings of all entities (nodes). However,
entities of different types (e.g., cells vs peaks, cells of different batches or modalities)
have different edge distributions and thus may lie on different manifolds of the latent
space. To make the embeddings of entities of different types comparable, we transform
the embeddings of features with the Softmax function by utilizing the first-order
similarity between cells (reference) and features (query). In the case of batch correction
or multi-omics integration, the Softmax transformation is also performed based on the
first-order similarity between cells of different batches or modalities.
957

958   Given the initial embeddings of cells (reference) $(v_{c_1}, \dots, v_{c_n})$ and features $(v_{f_1}, \dots, v_{d_m})$,
959   the model-estimated probability of an edge (c_i, f_j) obeys

$$P\left(v_{c_i, f_j}\right) \propto \exp\left(v_{c_i} \cdot v_{f_j}\right)$$

Therefore, if a random edge was sampled from feature f_j to a cell, the model would estimate the distribution over such edges as

$$p_{c_i, f_j} = \frac{\exp\left(v_{c_i} \cdot v_{f_j}\right)}{\sum_{k=1}^{n} \exp\left(v_{c_k} \cdot v_{f_j}\right)}$$

i.e., the Softmax weights between all cells $\{c_i\}$ and the feature f_j. We can then compute new embeddings for features as a linear combination of the cell embeddings weighted by the edge probabilities raised to some power.

$$\hat{v}_{f_j} = \frac{\sum_{i=1}^{n} p_{c_i, f_j}^{T^{-1}} v_{c_i}}{\sum_{i=1}^{n} p_{c_i, f_j}^{T^{-1}}}$$

T is a temperature hyperparameter that controls the sharpness of the weighting over cells. At $T = 1$, the cell embeddings are weighted by their estimated edge probabilities; at $T \to 0$, each feature embedding is assigned the cell embedding of its nearest neighbor; at $T \to \infty$, it becomes a discrete uniform distribution, and each query becomes the average of reference embeddings. We set $T = 0.5$ for all the analyses.

These steps are implemented in the function "si.tl.embed()" in the SIMBA package.

Metrics to assess cell-type specificity

Four metrics are proposed to assess the cell type specificity of each feature from different aspects, including max value (a higher value indicates higher cell-type specificity), Gini index (a higher value indicates higher cell-type specificity), standard deviation (a higher value indicates higher cell-type specificity), and entropy (a lower value indicates higher cell-type specificity). We observe these four metrics generally give consistent results. For SIMBA metric plot, by default, Gini index is plotted against max value. For feature f_j :

The max value is defined as the average normalized similarity of top k cells (by default, k=50). The similarity normalization function is defined as:

$$norm(x_i) = x_i - log\frac{\sum_{j=1}^{n} \exp\left(x_j\right)}{n}$$

Where $i = 1, \dots, n$. n is the number of cells and x_i represents the dot product of $\hat{v}_{f_j}$ and the embedding of cell i.

1000   The max value is computed as:

1001

$$\max(f_j) = \frac{\sum_{i=1}^{k} norm(x_i)}{k}$$

1002

1003

1004   The Gini index is computed as:

1005

$$\mathrm{gini}(f_j) = \frac{\sum_{i=1}^{n}(2i - n - 1) * p_{c_i, f_j}}{n \sum_{i=1}^{n} p_{c_i, f_j}}$$

1006

1007

1008   The standard deviation is computed as:

1009

$$\mathrm{std}(f_j) = \sqrt{\frac{1}{n-1} \sum_{i=1}^{n} (p_{c_i, f_j} - \mu)^2}$$

1010

1011   Where $\mu = \frac{1}{n} \sum_{i=1}^{n} p_{c_i, f_j}$.

1012

1013   Entropy is computed as:

$$\mathrm{entropy}(f_j) = - \sum_{i=1}^{n} p_{c_i, f_j} \log(p_{c_i, f_j})$$

1014

1015

1016 **Queries of entities in SIMBA space**

1017

The informative SIMBA embedding space serves as a database of entities including cells
and features. To query the "SIMBA database" for the neighboring entities of a given cell
or feature, we first build a k-d tree of all entities based on their SIMBA embeddings. We
then search for the nearest neighbors in the tree using Euclidean distance. To do so,
SIMBA query can perform either K-nearest neighbors (KNN) or nearest neighbor search
within a specified radius. SIMBA also provides the option to limit the search to entities
of certain types, which is useful when a certain type of entity significantly outnumbered
others. For example, the K nearest features of a given cell may be all peaks while genes
are the features of interest. In this case, SIMBA allows users to add "filters" to ensure
that nearest neighbor search is performed within the specified types of entities. This
procedure is implemented in the function "st.tl.query()" and its visualization is
implemented in the function "st.pl.query()" in the SIMBA package.

1030

1031 **Identification of master regulators**

1032

To identify master regulators, we take into consideration both the cell type specificity of
each pair of TF motif and TF gene and the distance between them. More specifically, for

each TF motif, first its distances (Euclidean distance by default) to all the genes are
calculated in the SIMBA embedding space. Then the rank of this TF gene among all these
genes is computed. In addition, we also assess the cell type specificity of this pair of TF
motif and TF gene based on SIMBA metrics (by default, max value and Gini index are
used). The same procedure is performed for all TFs. Finally, we identify master
regulators by filtering out TFs with low cell-type specificity and scoring them based on
TF gene rank. This procedure is implemented in the function
"st.tl.find_master_regulators()" in SIMBA package.
1043
1044

Identification of TF target genes

Given a master regulator, its target genes are identified by comparing the locations of
the TF gene, TF motif, and the peaks near the genomic loci of candidate target genes in
the SIMBA co-embedding space (**Fig. 4e**). More specifically we first search for k nearest
neighbor genes around the motif (TF motif) and the gene (TF gene) of this master
regulator, respectively (k = 200 by default). The union of these neighbor genes is the
initial set of candidate target genes. These genes are then filtered based on the criterion
that open regions (peaks) within 100kb upstream and downstream of the TSS of a
putative target gene must contain the TF motif.

Next, for each candidate target gene, we compute four types of distances in SIMBA
embedding space: distances between the embeddings of 1) the candidate target gene
and TF gene; 2) the candidate target gene and TF motif; 3) peaks near the genomic locus
of the candidate target gene and TF motif; 4) peaks near the genomic locus of the
candidate target gene and the candidate gene. All the distances (Euclidean distances by
default) are converted to ranks out of all genes or all peaks to make the distances
comparable across different master regulators.

The final list of target genes is decided using the calculated ranks based on two criteria:
1) at least one of the nearest peaks to TF gene or TF motif is within a predetermined
range (top 1,000 by default); 2) the average rank of the candidate target gene is within a
predetermined range (top 5,000 by default). This procedure is implemented in the
function "st.tl. find_target_genes ()" in SIMBA.

Benchmarking scATAC-seq computational methods
1067
To compare SIMBA to other scATAC-seq computational methods including SnapATAC [4],
Cusanovich2018[45], and cisTopic[46], we employed the previously developed benchmarking
framework from Chen et al[14](**Supplementary Table 1**). This framework evaluates
different methods based on their ability to distinguish cell types. We applied three
clustering algorithms: k-means clustering, hierarchical clustering, and Louvain on the
feature matrix derived from each method.
1074

1075	For datasets with ground-truth (FACS-sorted labels or known tissue labels), including
1076	simulated bone marrow data, Buenrostro 2018, and sci-ATAC-seq subset, three metrics
1077	including Adjusted Rand Index (ARI), Adjusted Mutual Information (AMI), and
1078	Homogeneity are applied to evaluate the performance. ARI measures the similarity
1079	between two clusters, comparing all pairs of samples assigned to matching or different
1080	clusters in the predicted clustering solution vs the true cluster/cell type label. AMI
1081	describes an observed frequency of co-occurrence compared to an expected frequency
1082	of co-occurrence between two variables, informing the mutual dependence or strength
1083	of association of these two variables. Homogeneity measures whether a clustering
1084	algorithm preserves cluster assignments towards samples that belong to a single class. A
1085	higher metric value indicates a better clustering solution.
1086	
1087	For 10x PBMCs dataset with no ground truth, the Residual Average Gini Index (RAGI)
1088	proposed in the benchmarking study[14] is used as the clustering evaluation metric. RAGI
1089	measures the relative exclusivity of marker genes to their corresponding clusters in
1090	comparison to housekeeping genes, which should demonstrate low specificity to any
1091	given cluster. In brief, the mean Gini Index is computed for both marker genes and
1092	housekeeping genes. The difference between the means is computed to obtain the
1093	average residual specificity (i.e., RAGI) of a clustering solution with respect to marker
1094	genes. A higher RAGI indicates a better separation of biologically distinct clusters.
1095	

Benchmarking single-cell batch correction methods

1097	
1098	The batch correction performance of SIMBA was compared to Seurat3[12], LIGER[11] and
1099	Harmony[10] in two benchmark datasets: the mouse atlas dataset and the human
1100	pancreas dataset (**Supplementary Table 1**). For Seurat3, LIGER and Harmony, the batch
1101	correction was done with the same parameters used in a previous benchmark study[25].
1102	
1103	To evaluate the batch integration performance, average Silhouette width (ASW),
1104	adjusted Rand index (ARI), and local inverse Simpson's index (LISI)[10] were calculated for
1105	the batches and cell types using the Euclidean distance as described in a previous
1106	benchmark[25]. To make a fair evaluation, only the cell types that are present in all
1107	batches were considered. We used the same number of dimensions (50) for these
1108	methods and all other parameters were set as in the benchmark.
1109	

Average Silhouette width (ASW)

1111	
1112	Average Silhouette width is the mean value of Silhouette scores calculated from each
1113	cell. Silhouette width measures the relative closeness of cells with the same label
1114	compared to the cells with the different label and ranges from -1 to +1. Silhouette score
1115	for a data point with a label is calculated as
1116	

$$s(i) = \frac{b(i) - a(i)}{\max\{a(i), b(i)\}}$$

where $a(i)$ is the distance to the closest point with the same label, and $b(i)$ is the distance to the closest point with different labels. A high Silhouette score means the point is located more closely with the same label, where a low Silhouette score closer to -1 means the point is located closer with different labels than that of itself. The ideal batch correction result will give a low ASW score for batch labels as the point is well mixed with other batches and a high ASW score for the cell type labels as the cells of the same cell type should cluster together after the batch correction. The final score is calculated as the median ASW scores from 20 subsets of randomly sampled 80% cells.

Average Rand Index (ARI)

To evaluate the cell type purity, the true cell type labels and the k-means clustering solution were used to calculate the cell type ARI. To evaluate the batch correction performance, the true batch labels and the k-means clustering solution were used to calculate the batch ARI. The final ARI was calculated as the median ARI scores of 20 subsets comprised of randomly sampled 80% cells for batches and cell types, respectively. A superior batch correction will have a high cell type ARI (high agreement between the clustering solution and the true cell type labels), and a low batch ARI (the clustering solution is not mainly driven by batches and clusters contain cells with well-mixed batch labels).

Local Inverse Simpson's Index (LISI)

Local Inverse Simpson's Index (LISI) [10] measures the local batch and cell type mixing. For each data point, it considers the Gaussian kernel weighted distribution of labels in its neighborhood with a perplexity argument. We set perplexity to ~~50~~ 40 as in the previous benchmark study. Using the weighted neighborhood label distribution, the inverse Simpson's index is calculated as $\frac{1}{\sum_l p(l)}$ where l is the batch or cell type labels and $p(l)$ is the probability of each label in the local neighborhood obtained with the kernel. For each cell, the LISI is the expected number of cells to be sampled locally before a cell of the same label is sampled. A perfect batch correction will have a cell type LISI (cLISI) of 1 and a batch LISI (integration LISI, iLISI) close to the number of batches. The final LISI score was calculated as the average LISI scores of all cells.

Further details are described in **Supplementary Note 2**.

Benchmarking single cell multi-omics integration methods

Two pairs of scRNA-seq and scATAC-seq datasets manually split from the dual-omics SHARE-seq mouse skin dataset and 10X PBMCs dataset respectively were used for the modality integration task. For Seurat3 and LIGER, the parameters and preprocessing

1161	were done as described in their documentations. However, for the LIGER analysis of the
1162	SHARE-seq mouse skin dataset the parameter 'lambda' was set to 30 and the
1163	'ref_dataset' was set to scATAC-seq to get a better alignment. For the Raw results, the
1164	activity matrix of scATAC-seq was constructed using Seurat3 and the first 20 PCs of the
1165	scRNA-seq count matrix and the activity matrix were used for the comparison. The
1166	integration results generated by each method were evaluated with four metrics—
1167	Anchoring distance, anchoring distance rank, Silhouette index, and cluster agreement—
1168	as described below.
1169	
1170	**Anchoring distance**
1171	The Anchoring distance was proposed in Dou et al., 2020[47] and is the normalized
1172	distance between the matched cells of two modalities (e.g. RNA and ATAC). Here we
1173	considered the Euclidean distance and normalized the distance by the mean of the
1174	distances calculated between random pairs of cells. The number of pairs randomly
1175	sampled was set to 10% of the total number of cells.
1176	
1177	**Anchoring distance rank**
1178	Given that the anchoring distance does not account for the local density of cells, we
1179	propose a new metric entitled *anchoring distance rank* (ADR). The ADR is based on the
1180	normalized rank of the distance between the matched cells of two modalities. For each
1181	cell x_{ij} with cell identity i and modality j, the distance between the cell and all the other
1182	cells of the other modality j', $d(x_{ij}, x_{kj'}), k = 1, ..., N$ is calculated, where N is the total
1183	number of cells. Then the rank of $r_i = d(x_{ij}, x_{ij'})$ within the calculated distances is
1184	normalized by the number of pairs $N - 1$ to obtain the final anchoring rank $m_i = \frac{r_i - 1}{N - 1}$.
1185	For each cell, an anchoring rank of 0 indicates an ideal modality integration performance
1186	as the matched cells are closest to each other in the embedding.
1187	
1188	**Silhouette index**
1189	The silhouette index was calculated as described in 10) based on the cluster assignment
1190	wherein each cluster consists of two cells, one cell from a scRNA-seq dataset and one
1191	cell from a scATAC-seq dataset.
1192	
1193	**Fraction in the same cluster**
1194	Fraction in the same cluster was calculated as the fraction of the matched cells from two
1195	modalities in the same cluster. The clusters of cells were generated using Louvain
1196	algorithm and the number of clusters is equal to the number of cell types in the dataset.
1197	
1198	Further details are described in **Supplementary Note 3**.
1199	
1200	
1201	**Data availability:**
1202	

All the datasets used in this study (eight scRNA-seq datasets, four scATAC-seq datasets, and three dual-omics datasets) are summarized in **Supplementary Table 1**. All these datasets are curated in the SIMBA package, and they can be easily downloaded and imported directly to reproduce the analyses presented in this manuscript.

Code availability:

We provide a comprehensive Python package 'simba' available at https://anaconda.org/bioconda/simba and https://github.com/pinellolab/simba. All the proposed procedures are implemented in the "simba" package. 'simba' can be easily installed with conda "*conda install simba*". We also built a website (https://simba-bio.readthedocs.io), providing a detailed introduction of the 'simba' software and several SIMBA tutorials for different types of single-cell analyses presented in this manuscript.

Acknowledgements

The authors thank Ledell (Yu) Wu, Facebook, for the helpful discussions about Starspace; Dr.Sai Ma and Dr. Jason Buenrostro, Broad Institute of MIT and Harvard, for sharing the SHARE-seq data and metadata. The authors also acknowledge members of the Pinello lab, Mass General Hospital/Harvard Medical School, for helpful comments and feedback. This project has been made possible in part by grant number 2019-202669 from the Chan Zuckerberg Foundation to LP. LP is also partially supported by the National Human Genome Research Institute (NHGRI) Genomic Innovator Award (R35HG010717). M.E.V. is supported by the National Institutes of Health (NIH) under the Ruth L. Kirschstein National Research Service Award (1F31CA257625-01) from the National Cancer Institute (NCI).

Author contributions

H.C. and L.P. conceived this project. H.C. developed SIMBA, wrote the SIMBA package, and performed SIMBA analysis on all datasets. A.L. contributed to the adaption of PyTorch-BigGraph to single cell analysis. J.R. and H.C. performed the comparison analysis on batch correction and data integration. M.E.V. and H.C. performed the comparison analysis on scATAC-seq data. L.P. and A.L. provided guidance and supervised this project. All the authors wrote and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

References:

1. Satija, R., Farrell, J.A., Gennert, D., Schier, A.F. & Regev, A. Spatial reconstruction of single-cell gene expression data. *Nat Biotechnol* **33**, 495-502 (2015).
2. Wolf, F.A., Angerer, P. & Theis, F.J. SCANPY: large-scale single-cell gene expression data analysis. *Genome Biol* **19**, 15 (2018).
3. Granja, J.M. et al. ArchR is a scalable software package for integrative single-cell chromatin accessibility analysis. *Nat Genet* **53**, 403-411 (2021).
4. Fang, R. et al. Comprehensive analysis of single cell ATAC-seq data with SnapATAC. *Nat Commun* **12**, 1337 (2021).
5. Kiselev, V.Y., Andrews, T.S. & Hemberg, M. Challenges in unsupervised clustering of single-cell RNA-seq data. *Nat Rev Genet* **20**, 273-282 (2019).
6. Cortal, A., Martignetti, L., Six, E. & Rausell, A. Gene signature extraction and cell identity recognition at the single-cell level with Cell-ID. *Nat Biotechnol* (2021).
7. Vandenbon, A. & Diez, D. A clustering-independent method for finding differentially expressed genes in single-cell transcriptome data. *Nat Commun* **11**, 4318 (2020).
8. Dann, E., Henderson, N.C., Teichmann, S.A., Morgan, M.D. & Marioni, J.C. Differential abundance testing on single-cell data using k-nearest neighbor graphs. *Nat Biotechnol* (2021).
9. Hao, Y. et al. Integrated analysis of multimodal single-cell data. *Cell* (2021).
10. Korsunsky, I. et al. Fast, sensitive and accurate integration of single-cell data with Harmony. *Nat Methods* **16**, 1289-1296 (2019).
11. Welch, J.D. et al. Single-Cell Multi-omic Integration Compares and Contrasts Features of Brain Cell Identity. *Cell* **177**, 1873-1887 e1817 (2019).
12. Stuart, T. et al. Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902 e1821 (2019).
13. Lerer, A. et al. Pytorch-biggraph: A large-scale graph embedding system. *arXiv preprint arXiv:1903.12287* (2019).
14. Chen, H. et al. Assessment of computational methods for the analysis of single-cell ATAC-seq data. *Genome Biology* **20**, 241 (2019).
15. Buenrostro, J.D. et al. Integrated Single-Cell Analysis Maps the Continuous Regulatory Landscape of Human Hematopoietic Differentiation. *Cell* **173**, 1535-1548 e1516 (2018).
16. Ferreira, R., Ohneda, K., Yamamoto, M. & Philipsen, S. GATA1 function, a paradigm for transcription factors in hematopoiesis. *Molecular and cellular biology* **25**, 1215-1227 (2005).
17. Tijchon, E., Havinga, J., Van Leeuwen, F. & Scheijen, B. B-lineage transcription factors and cooperating gene lesions required for leukemia development. *Leukemia* **27**, 541-552 (2013).
18. Friedman, A. Transcriptional control of granulocyte and monocyte development. *Oncogene* **26**, 6816-6828 (2007).
19. Schep, A.N., Wu, B., Buenrostro, J.D. & Greenleaf, W.J. chromVAR: inferring transcription-factor-associated accessibility from single-cell epigenomic data. *Nat Methods* **14**, 975-978 (2017).

20. Moriguchi, T. & Yamamoto, M. A regulatory network governing Gata1 and Gata2 gene
transcription orchestrates erythroid lineage differentiation. *International journal of*
hematology **100**, 417-424 (2014).
21. Chen, S., Lake, B.B. & Zhang, K. High-throughput sequencing of the transcriptome and
chromatin accessibility in the same cell. *Nat Biotechnol* (2019).
22. Ma, S. et al. Chromatin Potential Identified by Shared Single-Cell Profiling of RNA and
Chromatin. *Cell* (2020).
23. Cao, J. et al. Joint profiling of chromatin accessibility and gene expression in thousands
of single cells. *Science* **361**, 1380-1385 (2018).
24. Zhu, C. et al. An ultra high-throughput method for single-cell joint analysis of open
chromatin and transcriptome. *Nat Struct Mol Biol* (2019).
25. Tran, H.T.N. et al. A benchmark of batch-effect correction methods for single-cell RNA
sequencing data. *Genome Biol* **21**, 12 (2020).
26. Leek, J.T. et al. Tackling the widespread and critical impact of batch effects in high-
throughput data. *Nature Reviews Genetics* **11**, 733-739 (2010).
27. Han, X. et al. Mapping the mouse cell atlas by microwell-seq. *Cell* **172**, 1091-1107. e1017
(2018).
28. Tabula Muris, C. et al. Single-cell transcriptomics of 20 mouse organs creates a Tabula
Muris. *Nature* **562**, 367-372 (2018).
29. Baron, M. et al. A single-cell transcriptomic map of the human and mouse pancreas
reveals inter-and intra-cell population structure. *Cell systems* **3**, 346-360. e344 (2016).
30. Muraro, M.J. et al. A single-cell transcriptome atlas of the human pancreas. *Cell systems*
3, 385-394. e383 (2016).
31. Wang, Y.J. et al. Single-cell transcriptomics of the human endocrine pancreas. *Diabetes*
65, 3028-3038 (2016).
32. Segerstolpe, Å. et al. Single-cell transcriptome profiling of human pancreatic islets in
health and type 2 diabetes. *Cell metabolism* **24**, 593-607 (2016).
33. Ietswaart, R., Gyori, B.M., Bachman, J.A., Sorger, P.K. & Churchman, L.S. GeneWalk
identifies relevant gene functions for a biological context using network representation
learning. *Genome Biol* **22**, 55 (2021).
34. Yuan, H., Kshirsagar, M., Zamparo, L., Lu, Y. & Leslie, C.S. BindSpace decodes
transcription factor binding signals by large-scale sequence embedding. *Nat Methods*
(2019).
35. Li, H., Xiao, X., Wu, X., Ye, L. & Ji, G. scLINE: A multi-network integration framework
based on network embedding for representation of single-cell RNA-seq data. *J Biomed*
Inform **122**, 103899 (2021).
36. Buterez, D., Bica, I., Tariq, I., Andrés-Terré, H. & Liò, P. CELLVGAE: AN UNSUPERVISED
SCRNA-SEQ ANALYSIS WORKFLOW WITH GRAPH ATTENTION NETWORKS. *bioRxiv*
2020.12.20.423645v1 (2020).
37. Longo, S.K., Guo, M.G., Ji, A.L. & Khavari, P.A. Integrating single-cell and spatial
transcriptomics to elucidate intercellular tissue dynamics. *Nat Rev Genet* **22**, 627-644
(2021).
38. Kempfer, R. & Pombo, A. Methods for mapping 3D chromosome architecture. *Nat Rev*
Genet **21**, 207-226 (2020).

39.   VanHorn, S. & Morris, S.A. Next-Generation Lineage Tracing and Fate Mapping to
1333   Interrogate Development. *Dev Cell* **56**, 7-21 (2021).
1334   40.   Satopaa, V., Albrecht, J., Irwin, D. & Raghavan, B. in 2011 31st international conference
1335   on distributed computing systems workshops 166-171 (IEEE, 2011).
1336   41.   Fornes, O. et al. JASPAR 2020: update of the open-access database of transcription
1337   factor binding profiles. *Nucleic acids research* **48**, D87-D92 (2020).
1338   42.   Kadlec, R., Bajgar, O. & Kleindienst, J. Knowledge base completion: Baselines strike back.
1339   *arXiv preprint arXiv:1705.10744* (2017).
1340   43.   Krompaß, D., Baier, S. & Tresp, V. in International semantic web conference 640-655
1341   (Springer, 2015).
1342   44.   Mikolov, T., Chen, K., Corrado, G. & Dean, J. Efficient estimation of word representations
1343   in vector space. *arXiv preprint arXiv:1301.3781* (2013).
1344   45.   Cusanovich, D.A. et al. The cis-regulatory dynamics of embryonic development at single-
1345   cell resolution. *Nature* **555**, 538-542 (2018).
1346   46.   Bravo Gonzalez-Blas, C. et al. cisTopic: cis-regulatory topic modeling on single-cell ATAC-
1347   seq data. *Nat Methods* **16**, 397-400 (2019).
1348   47.   Dou, J. et al. Unbiased integration of single cell multi-omics data. *bioRxiv*,
1349   2020.2012.2011.422014 (2020).
1350
1351

Supplementary Figures

 Supplementary Figure 1. SIMBA analysis of the scRNA-seq 10x PBMCs dataset.

  a. Three default metrics used to evaluate SIMBA training procedure, including training loss
   (top), validation loss (middle), mean reciprocal rank (MRR)
  b. SIMBA metric plots of genes. All the genes are plotted according to the Gini index
   against max score, standard deviation (std) against max score, and entropy against max
   score, respectively. The same set of genes as in Figure 2c are highlighted.
  c. UMAP visualization of the SIMBA embeddings of cells and the SIMBA embeddings of
   cells and all genes. Genes highlighted in (b) are also highlighted in the UMAP plot.
  d. UMAP visualization of the SIMBA embeddings of cells, colored by gene expression of the
   genes highlighted in (b).
  e. SIMBA barcode plots of the genes highlighted in (b).

1367
1368

1369
Supplementary Figure 2. Comparison of SIMBA with Scanpy on the scRNA-seq 10x PBMCs
dataset.
a. Venn diagram of top marker genes identified by SIMBA and Scanpy
b. Scanpy-derived UMAP visualization of cells colored by cell type
c. Top marker genes detected only by SIMBA. Colored by intensity of gene expression.
d. Top marker genes detected only by Scanpy. Colored by intensity of gene expression.
e. SIMBA embedding result after implementing variable gene selection. Left: variable gene
selection step implemented in SIMBA. Middle: UMAP visualization of SIMBA

embeddings of cells. Right: UMAP visualization of SIMBA embeddings of cells and
variable genes.
1380

1381
1382
1383   **Supplementary Figure 3**. SIMBA analysis of the *Buenrostro2018* dataset
1384
a. UMAP visualization of SIMBA embeddings of cells colored by cell type (top-left), and TF
activity scores of TF motifs calculated with chromVAR, respectively. The SIMBA barcode
plot of each TF motif is shown below the UMAP plot.
b. Top: UMAP visualization of SIMBA embeddings of cells colored by TF activity scores of k-
mers calculated with chromVAR. Middle: SIMBA barcode plots of the corresponding k-
mers. Bottom: the matching known motif against the enriched k-mer sequences.
c. Heatmap of cells against neighboring peaks of each cell type that are selected in the
SIMBA co-embedding space. Chromatin accessibility is binary and colored accordingly.
1393

1399
1400
1401 **Supplementary Figure 4**. Comparison of SIMBA performance using scATAC-seq peaks and DNA
1402 sequence content vs only scATAC-seq peaks. **Top**: UMAP visualization of SIMBA embeddings of
1403 cells for each indicated dataset generated from only scATAC-seq peak information. **Bottom**:
1404 UMAP visualization of SIMBA embeddings of cells for each indicated dataset generated from
1405 scATAC-seq peak information and DNA sequence content information.
1406

1407
1408 **Supplementary Figure 5**. Benchmark of SIMBA against top-performing scATAC-seq analysis
1409 methods.
1410
1411 Top: Evaluation of SIMBA and other methods including cisTopic, SnapATAC, *Cusanovich2018* for
1412 scATAC-seq analysis using metrics 1) ARI, AMI, and Homogeneity for datasets with ground truth
1413 cell type labels and 2) Residual Average Gini Index (RAGI) for the 10x PBMCs dataset without
1414 ground truth labels.
1415
1416 Bottom: UMAP visualization of feature matrices produced by each method on each dataset
1417 colored by cell type annotation or cluster label.
1418

1419
1420

1421 **Supplementary Figure 6**. SIMBA multimodal analysis of the SHARE-seq hair follicle dataset.

1422

a. SIMBA embedding results when both gene expression and chromatin accessibility are
encoded in the graph. Left: UMAP visualization of SIMBA embeddings of cells and genes.
Middle: UMAP visualization of SIMBA embeddings of cells along with genes, TF motifs, and
k-mers. Right: UMAP visualization of SIMBA embeddings of cells along with genes, peaks, TF
motifs, and k-mers.
b. SIMBA embedding results when only gene expression is encoded in the graph. Left: UMAP
visualization of SIMBA embeddings of cells. Right: UMAP visualization of SIMBA embeddings
of cells and variable genes.
c. SIMBA embedding results when only chromatin accessibility is encoded in the graph. Left:
UMAP visualization of SIMBA embeddings of cells. Right: UMAP visualization of SIMBA
embeddings of cells and peaks.

1434
1435

1436
1437 **Supplementary Figure 7**. Cell type specific marker genes and the target genes of master
1438 regulators identified by SIMBA in the SHARE-seq hair follicle subset dataset.
1439
a. UMAP visualization of SIMBA embeddings of cells colored by cell type and gene
expression intensity.
b. SIMBA barcode plots of each gene plotted above.
c. SIMBA barcode plots of TF motifs *Lef1* and *Hoxc13.*
d. SIMBA barcode plots of peaks near the loci of *Lef1* and *Hoxc13.*
e. Top 30 target genes of the master regulators *Relb, Gata6,* and *Nfatc1* as inferred by
SIMBA.
1447

1448
1449

1450   **Supplementary Figure 8**. SIMBA multimodal analysis of the SNARE-seq mouse cerebral cortex
1451   dataset.

1452

a. From top to bottom: UMAP visualization of SIMBA embeddings of (i) cells (ii) genes
alongside cells (iii) genes, motifs, and k-mers alongside cells (iv) genes, peaks, motifs,
and k-mers alongside cells.
b. UMAP visualization of SIMBA embeddings of cells colored by indicated gene expression
intensity.
c. Heatmap of cells against neighboring peaks of each cell type that are selected in the
SIMBA co-embedding space. Chromatin accessibility is binary and colored accordingly.
d. SIMBA barcode plots of the genes highlighted in (a).

1461

1462
1463

1464 **Supplementary Figure 9**. SIMBA multimodal analysis of the 10x multiome PBMCs dataset.

1465

a. From top to bottom: UMAP visualization of SIMBA embeddings of (i) cells (ii) genes
alongside cells (iii) genes, motifs, and k-mers alongside cells (iv) genes, peaks, motifs,
and k-mers alongside cells.
b. UMAP visualization of SIMBA embeddings of cells colored by indicated gene expression
intensity.
c. Heatmap of cells against neighboring peaks of each cluster that are selected in the
SIMBA co-embedding space. Chromatin accessibility is binary and colored accordingly.
d. SIMBA barcode plots of the genes highlighted in (a).

1474

1475
1476
1477 **Supplementary Figure 10**. SIMBA-inferred marker genes for the scRNA-seq mouse atlas dataset
1478 in batch correction analysis.
1479
a. UMAP visualization of SIMBA embeddings of cells colored by cell type.
b. UMAP visualization of SIMBA embeddings of cells and genes.
c. UMAP visualization of SIMBA embeddings of cells and genes. Biological "query" points
are highlighted with a red "+". Nearby informative genes are colored accordingly.
d. UMAP visualization of SIMBA embeddings of cells colored by indicated gene expression
intensity, separated by cell type.
1486

1487
1488
1489 **Supplementary Figure 11**. SIMBA-inferred marker genes for the scRNA-seq human pancreas
1490 dataset in batch correction analysis.
1491
a. UMAP visualization of SIMBA embeddings of cells colored by cell type.
b. UMAP visualization of SIMBA embeddings of cells and genes.
c. UMAP visualization of SIMBA embeddings of cells and genes. Biological "query" points
are highlighted with a red "+". Nearby informative genes are colored accordingly.
d. UMAP visualization of SIMBA embeddings of cells colored by indicated gene expression
intensity, separated by cell type.
1498

Mouse atlas

Pancreas

1499
Supplementary Figure 12. Comparison of SIMBA to other methods for batch correction of the
mouse atlas (a-b) and human pancreas scRNA-seq datasets (c-d).
a, c. Quantitative comparison of SIMBA with three other batch correction methods
including Seurat3, LIGER and Harmony, using, left-to-right: average silhouette width (ASW),
adjusted Rand index (ARI), and local inverse Simpson's index (LISI)
b, d. UMAP visualization of raw and preprocessed data alongside the batch corrected
results produced by Seurat3, LIGER, Harmony, and SIMBA. Colored by technology (top) and
cell type (bottom).
1508

1509
1510

Supplementary Figure 13. SIMBA-inferred marker features for the SHARE-seq mouse skin
dataset in multi-omics integration analysis.

a. UMAP visualization of SIMBA embeddings of cells with two cellular modalities
   integrated.
b. UMAP visualization of SIMBA embeddings of cells, genes, and peaks with two cellular
   modalities integrated.
c. UMAP visualization of SIMBA embeddings of cells, genes, and peaks with two cellular
   modalities integrated. Biological "query" points are highlighted with a red "+". Nearby
   informative genes and peaks are colored accordingly.
d. UMAP visualization of SIMBA embeddings of cells, genes, and peaks with two cell
   modalities integrated and known marker genes highlighted.
e. UMAP visualization of SIMBA embeddings of cells colored by indicated gene expression
   intensity, separated by cell type.
f. Heatmap of cells against neighboring peaks of each cell type that are selected in the
   SIMBA co-embedding space. Chromatin accessibility is binary and colored accordingly.

1528

1529
1530
1531 **Supplementary Figure 14**. SIMBA-inferred marker features for the 10x human PBMCs dataset in
1532 multi-omics integration analysis.
1533
a. UMAP visualization of SIMBA embeddings of cells with two cellular modalities
integrated.
b. UMAP visualization of SIMBA embeddings of cells, genes, and peaks with two cellular
modalities integrated.
c. UMAP visualization of SIMBA embeddings of cells, genes, and peaks with two cellular
modalities integrated. Biological "query" points are highlighted with a red "+". Nearby
informative genes and peaks are colored accordingly.
d. Heatmap of cells against neighboring peaks of each cluster that are selected in the
SIMBA co-embedding space. Chromatin accessibility is binary and colored accordingly.
e. UMAP visualization of SIMBA embeddings of cells colored by indicated gene expression
intensity, separated by cell type.
1545

1546

Supplementary Figure 15. Comparison of SIMBA to other methods for multi-omics integration
of the SHARE-seq mouse skin (a-b) and 10x multiome human PBMCs (c-d) datasets.
a, c. Quantitative comparison of SIMBA with two other methods including Seurat3,
LIGER for multi-omics integration, using, left-to-right: anchoring distance rank,
anchoring distance, silhouette index, and Fraction in the same cluster.
b, d. UMAP visualization of the raw scRNA-seq and scATAC-seq data from the 10x
multiome human PBMCs dataset alongside the integrated results produced by Seurat3,
LIGER, and SIMBA. Colored by data modality (top) and cluster assignment (bottom). The
red intervals of violin plot of Anchoring distance and Silhouette index shows the 95%
of the mean.

1557

Supplementary Notes

1559

Supplementary Note 1: Comparison with scATAC-seq methods

1561

To assess SIMBA's ability to cluster cell types based on scATAC-seq profiles, we compared SIMBA with specialized methods specifically designed for this task. We observed that SIMBA yields consistent embeddings of cells when using either a single feature (peaks) or multiple features (peaks and DNA sequences from within those peaks) as input to the graph. This comparison was performed across four scATAC-seq datasets of varying profiling technologies and organisms (**Supplementary Fig. 4**). Given these differences, to create a fair comparison we used the same set of features (i.e., peaks) for SIMBA as other methods. SIMBA's performance was compared against three of the top methods, including SnapATAC[1], *Cusanovich*2018[2], and cisTopic[3] recommended by our recent benchmark study[4]. This comparison was first made qualitatively based on UMAP visualization and then quantitatively based on clustering performance. SIMBA performed as well as or better than each of the methods evaluated. These results comparing SIMBA to scATAC-seq-specialized methods highlight SIMBA's wide utility for single-cell analyses (**Supplementary Fig. 5**).

1575

Supplementary Note 2: Comparison with batch correction methods

1577

Multiple methods have now been developed to correct for the technical effects of sample preparation and data collection in single cells. To assess SIMBA's performance in removing batch effects, we compared it to Seurat3[5], LIGER[6] and Harmony[7], three top-performing batch correction methods recommended in a recent benchmark study[8].

1582

Two datasets, including a mouse atlas dataset and a human pancreas dataset (see **Supplementary Table 1**), were used for the evaluation. The mouse atlas dataset is composed of two scRNA-seq subsets with shared cell types from different sequencing platforms. The human pancreas dataset is composed of five samples pooled from five distinct sources using four different sequencing techniques wherein not all cell types are shared across each sample.

1588

To qualitatively compare these methods, we visualized cells of each dataset before and after batch-correction in UMAP plots (**Supplementary Fig. 12b,d**). To quantitatively evaluate the performance of each method, using the benchmarking pipeline laid out in Tran *et al*[8], we measured the conservation of biological information and batch effect removal based on three different metrics: average silhouette width (ASW), adjusted Rand index (ARI), and local inverse Simpson's index (LSI)[7] as in the previously-mentioned benchmark study[8] (**Supplementary Fig. 12a,c**; **Methods**). Each metric measures the relative mixing of class labels, where optimal performance is associated with maximal mixing in the batch labels and minimal mixing in the cell type labels.

1598

1599 The "Raw" batch correction results are the first 50 principal components of the horizontally
1600 concatenated gene-by-cell expression count matrix using *stats::prcomp* in R package with
1601 centering and scaling. The "Raw, preprocessed" batch correction used the preprocessed data
1602 with log normalization with scaling factor 10^4 and selection of 3000 highly variable genes with
1603 Seurat v3 with no restriction on the minimum number of cells and genes.
1604
1605 For batch correction using Seurat v3, default options are used for pancreas dataset whereas for
1606 mouse atlas dataset no cutoff was used for the minimum number of cells and genes as in Tran
1607 *et al.*[8]. The dimension of the batch corrected embedding is set as 50 dimensions following the
1608 default option for *Seurat::RunPCA* and for the consistency with SIMBA.
1609
1610 For batch correction using LIGER, the same arguments are used (lambda = 5, nrep = 3) are used
1611 for *liger::optimizeALS* in Tran *et al.* other than the number of factors k was set as 50 for
1612 consistency with other methods for both datasets.
1613
1614 For batch correction using Harmony, the same arguments are used as in Tran *et al.*[8] other than
1615 the number of dimensions of the output embedding was set to 50 instead of 20. We note that
1616 the output embedding of 20 dimensions would result in the similar result as when used 50
1617 dimensions in these methods.
1618
1619 **Supplementary Note 3**: **Comparison with multi-omics integration methods**
1620
1621 Seurat3 and LIGER are two of the most widely-adopted methods for single-cell data integration.
1622 Here, we demonstrate that SIMBA outperforms these methods on two separate datasets, the
1623 recently published SHARE-seq mouse skin dataset and the similarly recent 10x PBMCs multiome
1624 dataset (**Supplementary Table 1**). We focus on Seurat3 and LIGER as they have explicit
1625 documentation for the task of integrating scRNA-seq and scATAC-seq data.
1626
We first qualitatively evaluated these methods by inspecting UMAP visualization plots. For the
SHARE-seq dataset, we observed that all three methods perform comparably well in mixing
cells of two modalities though LIGER generated particularly small and noisy clusters
(**Supplementary Fig. 15b)**. For the 10X PBMCs dataset, SIMBA resulted in the best mixing of
cells belonging to each modality whereas other methods clustered cells separately within the
originating modalities (**Supplementary Fig. 15d**). We next quantitatively assessed the
integration performance of each method using four metrics that measure the distances
between matched cells in the integrated space (**Methods**). In addition to the commonly-used
metrics including anchoring distance, Silhouette index, and Fraction in the same cluster, we
developed an additional metric, *anchoring distance rank* (ADR)*,* which represents the
normalized rank of the distance between matching cells. If two matching cells from scRNA-seq
and scATAC-seq are mutually closest to one another, their ADR will be close to 0 (**Methods**) and
thus a minimized ADR is ideal. Overall SIMBA showed the best performance according to ADR
as well as cluster agreement while showing comparable or better performance according to the
remaining metrics for both datasets (**Supplementary Fig. 15a,c**).
1642

1643

1644 The modality integration procedure for Seurat v3 and LIGER follows the tutorial provided by the
1645 authors (Seurat v3:
1646 https://satijalab.org/seurat/archive/v3.1/atacseq_integration_vignette.html; LIGER:
1647 http://htmlpreview.github.io/?https://github.com/welch-
1648 lab/liger/blob/master/vignettes/Integrating_scRNA_and_scATAC_data.html).

1649

Both Seurat v3 and LIGER formulate the modality integration task between scRNA-seq and
scATAC-seq data as a batch correction task between scRNA-seq and gene activity matrix
constructed from scATAC-seq. In Seurat v3, the gene activity score of a gene is calculated as the
sum of the read counts in the peaks that falls within from 2kb upstream of the TSS to the end of
the gene body. In LIGER, this score is calculated as the sum of all read counts that falls within
3kb upstream of the TSS to the end of the gene body.

1656

The "Raw" results start from a scRNA-seq count matrix and a gene activity matrix calculated by
Seurat v3. Filtering for the shared genes in both modalities resulted in 16738 genes for the
SHARE-seq mouse skin dataset and 11045 genes for the 10X PBMCs dataset. Gene-by-cell gene
expression matrix and gene activity matrix were horizontally concatenated along matching rows
(genes). The output embedding is the first 20 principal components calculated by the R function
stats::prcomp with centering and scaling.

1663

1664 For the modality integration using Seurat v3, the gene expression count was filtered using the
1665 default parameters *min.cells = 3* and *min.features = 200*. The co-embedding was created as
1666 described in the tutorial of the package using the scRNA-seq. The output embedding consists of
1667 the first 50 principal components, which is the default option of *Seurat::RunPCA*.

1668

For the modality integration using LIGER, the gene expression count and gene activity matrices
were normalized and filtered for the genes that are shared between both matrices. The values
were then scaled according to the tutorial. In applying LIGER to the SHARE-seq mouse skin
dataset, the function, *liger::optimizeALS* was used with the default parameters, k = 20 and
lambda = 5. The scRNA-seq dataset was indicated as the reference in the function,
liger::quantile_norm as described in the documentation. The scRNA-seq and scATAC-seq
modalities of the 10X PBMC multiome dataset were unable to be aligned using the default
parameters. Thus *lambda = 30* and *max.iters = 100* were used for the *liger::optimizeALS*
function and the scATAC-seq dataset was indicated as the reference using the
liger::quantile_norm function to ensure a better alignment.

1679

1680 References

1681

1682

1.   Fang, R. et al. Comprehensive analysis of single cell ATAC-seq data with SnapATAC. *Nat*
Commun **12**, 1337 (2021).
2.   Cusanovich, D.A. et al. The cis-regulatory dynamics of embryonic development at single-
cell resolution. *Nature* **555**, 538-542 (2018).

3.   Bravo Gonzalez-Blas, C. et al. cisTopic: cis-regulatory topic modeling on single-cell ATAC-
seq data. *Nat Methods* **16**, 397-400 (2019).
4.   Chen, H. et al. Assessment of computational methods for the analysis of single-cell
ATAC-seq data. *Genome Biology* **20**, 241 (2019).
5.   Stuart, T. et al. Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902
e1821 (2019).
6.   Welch, J.D. et al. Single-Cell Multi-omic Integration Compares and Contrasts Features of
Brain Cell Identity. *Cell* **177**, 1873-1887 e1817 (2019).
7.   Korsunsky, I. et al. Fast, sensitive and accurate integration of single-cell data with
Harmony. *Nat Methods* **16**, 1289-1296 (2019).
8.   Tran, H.T.N. et al. A benchmark of batch-effect correction methods for single-cell RNA
sequencing data. *Genome Biol* **21**, 12 (2020).
1699