Strip1 regulates retinal ganglion cell survival by suppressing Jun-mediated apoptosis to promote retinal neural circuit formation

Mai Ahmed, Yutaka Kojima, and Ichiro Masai*

Developmental Neurobiology Unit, Okinawa Institute of Science and Technology Graduate University, Tancha, Okinawa, Japan.

* Correspondence should be addressed: masai@oist.jp

Key words: Zebrafish, Strip1, retinal ganglion cells, Jun, lamination, apoptosis, migration, Striatin 3.

Abstract

In the vertebrate retina, an interplay between retinal ganglion cells (RGCs), amacrine and bipolar cells establishes a synaptic layer called the inner plexiform layer (IPL). This circuit conveys signals from photoreceptors to visual centers in the brain. However, the molecular mechanisms involved in its development remain poorly understood. Striatin-interacting protein 1 (Strip1) is a core component of the STRIPAK complex, and it has shown emerging roles in embryonic morphogenesis. Here, we uncover the importance of Strip1 in inner retina development. Using zebrafish, we show that loss of Strip1 causes defects in IPL formation. In *strip1* mutants, RGCs undergo dramatic cell death shortly after birth. Amacrine and bipolar cells subsequently invade the degenerating RGC layer, leading to a disorganized IPL. Thus, Strip1 promotes IPL formation through RGC maintenance. Mechanistically, zebrafish Strip1 interacts with its STRIPAK partner, Striatin3, to promote RGC survival by suppressing Jun-mediated apoptosis. In addition to its function in RGC maintenance, Strip1 is required for RGC dendritic patterning, which likely contributes to proper IPL formation. Taken together, we propose that a series of Strip1-mediated regulatory events coordinates inner retinal circuit formation by maintaining RGCs during development, which ensures proper positioning and neurite patterning of inner retinal neurons.

Introduction

The retina has a highly organized neural circuit that consists of six major classes of neurons, which are assembled into three cellular layers with two synaptic or plexiform layers in between. This beautiful layered architecture is termed "retinal lamination" (Avanesov and Malicki, 2010, D'Orazi et al., 2014, Dowling, 1987). Lamination is conserved among vertebrates and critical for processing of visual information (Baden et al., 2020, Nassi and Callaway, 2009). During development, neurogenesis, cell migration and neurite patterning are spatially and temporally coordinated to form retinal lamination. Any defect in these events can disrupt retinal wiring and consequently compromise visual function (Amini et al., 2018). However, molecular mechanisms that govern retinal neural circuit formation are not fully understood.

The retinal neural circuit processes visual signals through two synaptic neuropils (Figure 1A). At the apical side, the outer plexiform layer (OPL) harbors synapses that transmit input from photoreceptors (PRs) in the outer nuclear layer (ONL) to bipolar (BP) and horizontal cells (HCs) in the inner nuclear layer (INL). At the basal side of the retina, the inner plexiform layer (IPL) is densely packed with synaptic connections formed between BPs and amacrine cells (ACs) in the INL and retinal ganglion cells (RGCs) in the ganglion cell layer (GCL). The retina contains one type of glial cells called Müller glia (MGs) which span the apico-basal axis of the retina (Hoon et al., 2014, Huberman et al., 2010).

RGCs are the first-born retinal neurons, which extend their axons to exit the eye cup and innervate visual centers in the brain (D'Souza and Lang, 2020, Robles et al., 2014). In mouse and zebrafish models, when RGCs are absent or exhibit defects in axon projections, vision is compromised (Kay et al., 2001, Moshiri et al., 2008, Rick et al., 2000). So, RGCs are indispensable for vision. RGC degeneration is often a secondary defect in optic neuropathies and the leading cause of blindness. Thus, tremendous efforts are being dedicated to decipher signaling pathways involved in RGC death (Maes et al., 2017, Munemasa and Kitaoka, 2012).

Strip1 is a recently identified protein with emerging functions in neuronal development. It was first described as one of the core components of the striatin-interacting phosphatases and kinases (STRIPAK) complex (Goudreault et al., 2009).

STRIPAK complex is an evolutionarily conserved supramolecular complex with diverse functions in cell proliferation, migration, vesicular transport, cardiac development, and cancer progression (He et al., 2010, Hwang and Pallas, 2014, La Marca et al., 2019, Madsen et al., 2015, Neisch et al., 2017, Shi et al., 2016). In addition, several STRIPAK components participate in dendritic development, axonal transport, and synapse assembly (Chen et al., 2012, Li et al., 2018, Schulte et al., 2010). In *Drosophila*, Strip (a homolog of mammalian Strip1/2) is essential for axon elongation by regulating early endosome trafficking and microtubule stabilization (Sakuma et al., 2014, Sakuma et al., 2015). In addition, Strip, together with other STRIPAK members, modulates synaptic bouton development and prevents ectopic retina formation (Neal et al., 2020, Sakuma et al., 2016). On the other hand, loss of mouse Strip1 causes early mesoderm migration defects leading to embryonic lethality (Bazzi et al., 2017, Zhang et al., 2021). Thus, the role of Strip1 in the vertebrate nervous system is largely unknown.

Here, we report an essential role for Strip1 in neural circuit formation of zebrafish retina. In zebrafish *strip1* mutants, retinal lamination, especially IPL formation, is disrupted. Loss of Strip1 causes RGC death shortly after birth. Cells in the INL subsequently infiltrate the degenerating GCL, leading to a disorganized IPL. Strip1 cell-autonomously promotes RGC survival; however, it is not required in INL cells for IPL formation. Therefore, Strip1-mediated RGC maintenance is required to establish the IPL. Mechanistically, Strip1 interacts with its STRIPAK partner, Striatin3, to promote RGC survival through the suppression of the Jun-mediated apoptotic pathway. We also found that Strip1 is necessary for RGC dendritic patterning, which likely promotes interaction between RGCs and ACs for IPL formation. Collectively, we demonstrate that Strip1 is crucial for RGC survival during development and thereby coordinates proper wiring of the inner retina.

Results

Strip1 is essential for inner retinal neural circuit development

To understand mechanisms of retinal neural circuit formation, we screened zebrafish retinal lamination-defective mutants (Masai et al., 2003) and identified the *rw147* mutant. At 4 days post-fertilization (dpf), *rw147* mutant embryos have small eyes, lower jaw atrophy and cardiac edema (Figure 1B). *rw147* mutants also show defects in retinal lamination, in which retinal layers, especially in the inner retina, fluctuate in a wave-like pattern (Figure 1C). The *rw147* mutation is lethal by 6 dpf due to cardiac edema. Mapping of the *rw147* mutation revealed a missense mutation in exon 7 of *striatin interacting protein 1* (*strip1*) gene of the *rw147* mutant genome (Figure 1D).

Next, we performed Crispr/cas9-mediated mutagenesis to generate a 10-base deletion mutant, *strip1$^{crispr\Delta10}$* (Figure 1–figure supplement 1A). *strip1$^{crispr\Delta10}$* mutants show similar morphology and retinal lamination defects to those of *strip1^{rw147}* mutants (Figure 1–figure supplement 1B-1D). Likewise, knockdown of Strip1 using translation-blocking morpholinos (MO-Strip1) phenocopied *strip1^{rw147}* mutants (Figure 1–figure supplement 1E and 1F). We verified the specificity of MO-Strip1 using a custom-made zebrafish Strip1 antibody that fails to detect a 93-kDa protein band corresponding to zebrafish Strip1 in the morphants (Figure 1–figure supplement 1G). Furthermore, we generated transgenic lines that express wild-type and *rw147* mutant forms of zebrafish Strip1 protein under the control of the heat shock promotor, *Tg[hsp:WT.Strip1-GFP]* and *Tg[hsp:Mut.Strip1-GFP],* respectively. Wild-type Strip1, but not the mutant form, rescued the retinal defects of *strip1^{rw147}* (Figure 1–figure supplement 1H and 1I). Taken together, the *strip1* mutation is the cause of retinal lamination defects.

Next, we examined Strip1 expression in wild-type retinas by labeling with the zebrafish Strip1 antibody. Strip1 was expressed in RGCs and ACs at 2 dpf (Figure 1E). *In situ* hybridization shows that *strip1* mRNA is maternally and zygotically expressed and by 2 dpf, expression becomes restricted to the eyes, optic tectum, and heart (Figure 1–figure supplement 2A and 2B). Like Strip1 protein, *strip1* mRNA was expressed in RGCs and ACs (Figure 1–figure supplement 2C). To visualize retinal neuropils, we performed whole-mount staining of the retina with anti-acetylated α-tubulin antibody. In wild-type retinas, IPL and OPL are evident at 3 dpf. In contrast, IPL shows abnormal

morphology, whereas OPL is relatively normal in *strip1^{rw147}* mutants (Figure 1F). We

tracked IPL development using Bodipy TR stain. In wild-type retinas, a rudimentary IPL

was formed as early as 52 hours post-fertilization (hpf); however, it was less defined in

strip1^{rw147} mutants (Figure 1–figure supplement 2D). At 62 hpf, mutants exhibited a

wave-like IPL. This temporal profile coincides with development of RGCs and ACs.

Next, we visualized neurite morphology of RGCs, ACs, and BPs by transiently

expressing fluorescent proteins under control of *ath5* (Masai et al., 2003), *ptf1a* (Jusuf

and Harris, 2009), and *nyx* promoters (Schroeter et al., 2006) respectively. In wild-type

siblings, RGCs and ACs normally extend their dendrites toward the IPL; however,

strip1^{rw147} mutants show randomly directed dendritic patterns of RGCs and ACs (Figures

1G and 1H). In wild-type siblings, BPs normally extend their axons and dendrites toward

IPL and OPL, respectively; however, BPs of *strip1^{rw147}* mutants show mis-routed axons

and abnormal dendritic branching (Figure 1I). Thus, Strip1 is required for IPL formation

and correct neurite patterning of RGCs, ACs, and BPs.

RGCs are reduced and INL cells infiltrate the GCL in *strip1* mutants

To examine how IPL is disrupted in *strip1* mutants, we combined *strip1^{rw147}* mutants with

two transgenic lines, *Tg[ath5:GFP; ptf1a:mCherry-CAAX]*, to visualize RGCs and ACs.

In *Tg[ath5:GFP]*, GFP is expressed strongly in RGCs and weakly in ACs and

photoreceptors under control of the *ath5* enhancer (Masai et al., 2003). In

Tg[ptf1a:mCherry-CAAX], membrane-targeted mCherry is expressed in ACs and HCs

under control of *ptf1a* promoter (Jusuf and Harris, 2009). Live imaging of 3-dpf retinas

revealed that RGCs are severely reduced in *strip1^{rw147}* mutants (Figure 2A and 2B). On

the other hand, there was no significant change in the number of ptf1a+ ACs between

strip1^{rw147} mutants and wild-type siblings (Figure 2A and 2C). However, ptf1a+ ACs

abnormally extended their dendrites to form an irregular pattern of IPL (Figure 2A). In

wild-type retinas, the majority of ACs reside in the INL, except displaced ACs (Jusuf and

Harris, 2009). However, in *strip1^{rw147}* mutants, a significant fraction of ptf1a+ ACs were

abnormally located in the GCL (Figure 2A, bottom panels and 2D). Such abnormal

positioning of ACs is correlated with the severity of IPL defects (Figure 2–figure

supplement 1A). This phenotype is reminiscent of *ath5* mutants, *lakritz*, in which RGCs

fail to undergo neurogenesis, leading to infiltration of ACs into the GCL and transient IPL formation defects (Kay et al., 2001, Kay et al., 2004). We confirmed similar IPL defects in *ath5* morphant retinas at 3 dpf (Figure 2–figure supplement 1B), albeit weaker than those of *strip1^{rw147}* mutants.

Next, we visualized ACs using anti-Pax6 antibody, which strongly labels ACs and weakly labels RGCs (Macdonald and Wilson, 1997). In wild-type siblings, most strong Pax6+ cells were in the INL, and only 9.84±4.13% were in the GCL (Figure 2E, 2G). However, in *strip1^{rw147}* mutants, a significant percentage of Pax6+ cells (44.26±17.8%) was in the GCL (Figure 2E, 2G). The total number of Pax6+ cells did not differ between wild-type siblings and *strip1^{rw147}* mutants (Figure 2F). Next, we visualized BPs using anti-Prox1 antibody, which labels BPs and HCs (Jusuf and Harris, 2009). In wild-type, 100% of Prox1+ cells were in the INL (Figure 2H and 2J). However, 10.6±6.26% of Prox1+ cells were abnormally located in the GCL in *strip1^{rw147}* mutants (Figures 2H and 2J). The total number of Prox1+ cells did not differ between wild-type siblings and *strip1^{rw147}*mutants (Figure 2I). We performed labeling of double-cone and rod photoreceptors using zpr1 and zpr3 antibodies, respectively (Nishiwaki et al., 2008). Apart from occasional mildly disrupted areas, the photoreceptor cell layer was grossly intact, with no positioning defects (Figure 2–figure supplement 1C and 1D). Müller glia and proliferating cells at the ciliary marginal zone were visualized using anti-glutamine synthetase (GS) (Peterson et al., 2001) and anti-PCNA antibodies (Raymond et al., 2006), respectively. Both cell types showed grossly normal positioning in *strip1^{rw147}* mutants (Figure 2–figure supplement 1E and 1F). Thus, in the absence of Strip1, INL cells abnormally infiltrate the GCL and seem to replace the reduced RGCs.

Strip1 cell-autonomously promotes RGC survival

In zebrafish, RGC genesis starts in the ventronasal retina at 25 hpf, spreads into the entire retina by 36 hpf and is completed by 48 hpf (Avanesov and Malicki, 2010, Hu and Easter, 1999). The reduction of RGCs in *strip1* mutants could be due to compromised RGC genesis or RGC death after birth. To clarify which, we examined RGC genesis by monitoring ath5:GFP expression, and apoptosis by TUNEL. In *strip1^{rw147}* mutants, RGCs are normally produced at 36 hpf; however, apoptosis occurred in the GCL at 48 hpf

(Figure 3A). The number of apoptotic cells in GCL reached its highest level at 60 hpf, and apoptotic cells were eliminated by 96 hpf (Figure 3A and 3B). Accordingly, RGCs were significantly lower in *strip1^{rw147}* mutants than in wild-type siblings at 60 hpf and progressively reduced by 96 hpf (Figure 3C). In contrast, other retinal layers of *strip1^{rw147}* mutants showed slightly, but not significantly increased apoptosis at 72 hpf (Figure 3–figure supplement 1A), suggesting a specific function of Strip1 in RGC survival. In addition, despite the reduction in ath5:GFP+ area, the total presumptive GCL area, which was defined by retinal area between the lens and the IPL, was unchanged in *strip1^{rw147}* mutants throughout the stages (Figure 3–figure supplement 1B), suggesting that infiltrating INL cells replace the lost RGCs. We confirmed RGC death in *strip1^{crisprΔ10}* mutants (Figure 3–figure supplement 1C and 1D). Interestingly, we observed apoptosis in the optic tectum of *strip1^{rw147}* mutants (Figure 3–figure supplement 1E and 1F), suggesting a common Strip1-dependent survival mechanism in the optic tectum. RGCs are the only retinal neurons whose axons project to the optic tectum. In *strip1^{rw147}* mutants, RGC axons appeared to exit from the eye cup and formed an optic chiasm at 3 dpf (Figure 3–figure supplement 1G). However, consistent with the reduction of RGCs, the optic nerve was thinner in *strip1^{rw147}* mutants than in wild-type siblings and showed elongation defects towards the optic tectum.

To determine whether Strip1 cell-autonomously promotes RGC survival, we conducted cell transplantation from *strip1^{rw147}* mutant donor cells into wild-type host embryos at the blastula stage. TUNEL of transplanted retinas at 60 hpf revealed that *strip1^{rw147}* mutant donor RGCs underwent apoptosis in wild-type host retinas (Figures 3D-3F). Thus, Strip1 is cell-autonomously required for survival of RGCs.

RGC death triggers abnormal positioning of ACs, leading to IPL disruption

ACs are proposed to be the main cell type responsible for IPL formation (Godinho et al., 2005, Huberman et al., 2010). To clarify how RGC death influences infiltration of ACs into GCL and IPL disruption, we performed time-lapse imaging of wild-type and *strip1^{rw147}* mutant retinas combined with the transgenic line *Tg[ath5:GFP; ptf1a:mCherry-CAAX]*. At 48 hpf, there were no apparent differences in position or morphology of RGCs and ACs between wild-type siblings and *strip1^{rw147}* mutants

(Figure 4A and Videos 1 and 2). In *strip1^rw147* mutants at 52 hpf, RGCs started to
disappear, creating an empty spot in the GCL (Figure 4A, asterisks). However, ACs
were still located in the INL. At 55 hpf, a rudimentary IPL was observed in the central
retina of both wild-type siblings and *strip1^rw147* mutants. At 59 hpf, ACs started to invade
the empty spaces in the GCL (Figure 4A, arrowheads). Infiltration of ACs into the GCL
was more prominent at 62 hpf, resulting in a fluctuating IPL. Thus, loss of RGCs triggers
infiltration of ACs into the GCL in *strip1^rw147* mutants.

To examine whether Strip1 is required in ACs for IPL formation, we performed
cell transplantation using donor embryos carrying the transgene *Tg[ptf1a:mCherry-*
CAAX] (Figure 4B). When mutant ACs were transplanted into wild-type host retinas,
most donor ACs were normally positioned in the INL and extended dendrites toward the
IPL, as in the case of wild-type donor ACs transplanted into a wild-type host retina
(Figure 4C, 4D). Occasionally, three ACs extended 2 dendritic trees instead of 1 among
73 transplanted ACs; however, such dendritic mis-projection did not perturb IPL
formation (Figure 4–figure supplement 1A). On the other hand, as with mutant donor
ACs transplanted into mutant host retinas, when wild-type donor ACs were transplanted
to mutant host retinas, they showed irregular neurite projection with many somas
abnormally located toward the basal side, resulting in IPL formation defects (Figure 4C,
4D and Figure 4–figure supplement 1A). These data suggest a non-cell autonomous
function of Strip in ACs for IPL formation.

Similarly, we conducted cell transplantation to assess the role of Strip1 in BPs.
Mutant donor BPs labeled with the transgene *Tg[xfz43]* (Zhao et al., 2009) normally
projected axons toward the IPL in wild-type host retinas, in the same fashion as wild-
type donor BPs (Figure 4–figure supplement 1B-1D). Few transplanted columns of
mutant donors showed extra lateral branching and excessive elongation of BP arbors
(Figure 4–figure supplement 1D, arrows). However, such arbor defects did not disrupt
the IPL. On the other hand, when wild-type donor BPs labeled with the transgene,
Tg[xfz3], were transplanted into a mutant host retina, wild-type donor bipolar axons
failed to project toward the mutant host IPL, but rather seemed to be guided toward the
wild-type donor IPL (Figure 4–figure supplement 1E-1G). Thus, Strip1 is not required in
ACs or BPs for neurite projection to the IPL, although we do not exclude the possibility

that Strip1 is cell-autonomously required in a small subset of ACs and BPs to regulate
dendritic branching and neurite extension. Taken together, it is likely that Strip1-
mediated RGC maintenance is essential for proper neurite patterning of ACs and BPs
and subsequent IPL formation.

Strip1 interacts with Striatin3 to regulate RGC survival

To identify which molecules interact with Strip1 to regulate RGC survival, we conducted
a co-immunoprecipitation experiment coupled with mass spectrometry (Co-IP/MS).
Head lysates of wild-type embryos combined with the transgenic line *Tg[hsp:WT.Strip1-*
GFP] were used to pull-down wild-type Strip1 and its interacting partners. As a negative
control, we used lysates from 2 other lines; *Tg[hsp:Mut.Strip1-GFP]* and
Tg[hsp:Gal4;UAS:GFP], to rule out proteins enriched by the mutant form of Strip1 or by
GFP alone (Figure 5A and 5B). Six proteins were enriched only by the wild-type form of
Strip1, 5 of which are components of the STRIPAK complex (Figures 5C, 5D, and
Figure 5–figure supplement 1A and 1B). Since none of these components has been
studied in zebrafish, we analyzed previously published single-cell RNA sequencing data
on transcriptomes from zebrafish embryonic retinas at 2 dpf (Xu et al., 2020), and found
that only *strip1* and *striatin3 (strn3)* mRNA are abundantly expressed in retinal cells
(Figure 5–figure supplement 1C-1I). Next, we knocked down Strn3 using morpholino
and observed a significant increase in apoptotic cells in the GCL at 60 hpf (Figure 5E
and 5F). This leads to a significant reduction in RGCs at 60 and 76 hpf, as assessed by
the *ath5:GFP* signal (Figures 5G-5I). Although *strn3* morphants showed a similar RGC
loss to *strip1* mutants, it was weaker. IPL defects were also milder in *strn3* morphants
than in *strip1^{rw147}* mutants at 76 hpf (Figure 5H). Taken together, Strip1 and Strn3
cooperatively regulate RGC survival.

Jun is a key mediator of RGC death in the absence of Strip1

To determine what kinds of molecules mediate RGC apoptosis in *strip1* mutants, we
performed RNA sequencing on transcriptomes from 62-hpf eye cups of *strip1^{rw147}*
mutants. Compared to wild-type siblings, *strip1* mutants had 131 significantly
upregulated genes and 75 downregulated genes (Figure 6A, log$_2$FC>|1|, and

FDR<0.05). Most downregulated genes were markers of RGCs, like *isl2b*, *pou4f3* (also

known as *brn3c*) and *tbr1b,* which reflects the reduction in RGCs. Genes related to

synaptic development and transmission were also downregulated (Figures 6A, 6B and

Figure 6–figure supplement 1A). On the other hand, many of the significantly

upregulated genes were related to apoptosis, oxidative phosphorylation, cellular

response to stress, and the MAP kinase (MAPK) signaling pathway (Figure 6A, 6B and

Figure 6–figure supplement 1B). Most reports of RGCs undergoing stress are in

glaucoma and optic nerve injury models, where adult RGCs undergo cell death in

response to injury (Bahr, 2000). So, we compared the transcriptomic profiles of *strip1*

mutant eyes to those of adult zebrafish RGCs following optic nerve injury (Veldman et

al., 2007) or adult eyes after optic nerve crush (McCurley and Callard, 2010). Indeed,

there were several genes commonly upregulated in all three models, namely, *jun, atf3,*

gap43, stmn4, sox11b, and *adcyap1b* (Figure 6–figure supplement 1C). These findings

suggest that Strip1-deficient zebrafish RGCs share a similar stress response with adult

RGCs following optic nerve injury.

In *strip1^{rw147}* mutants, *jun* was among the top upregulated stress response

markers. Jun (the zebrafish homolog of mammalian c-Jun) is the canonical target of the

Jun N-terminal kinase (JNK) pathway, which belongs to the MAPK super family. JNK/c-

Jun signaling is a key regulator of stress-induced apoptosis (Dhanasekaran and Reddy,

2008, Ham et al., 2000). Activation of the JNK pathway involves phosphorylation events

that end with c-Jun phosphorylation and transactivation, which in turn activates *c-jun*

gene expression (Eilers et al., 1998). We stained *strip1^{rw147}* mutant retinas with anti-

phosphorylated c-Jun (p-Jun) antibody. At 54 hpf, *strip1^{rw147}* mutants showed

significantly elevated levels of p-Jun compared to wild-type siblings. This elevation is

specifically localized in RGCs visualized with zn5 antibody (Figures 6C and 6D).

Likewise, p-Jun was significantly elevated in RGCs of *strn3* morphants at 49 hpf

compared to control injected embryos (Figures 6E and 6F). Next, we knocked down Jun

using morpholino (MO-jun). At 60 hpf, apoptosis was significantly inhibited in the GCL of

strip1^{rw147} mutants injected with MO-jun, compared to *strip1^{rw147}* mutants injected with a

standard control morpholino (Figures 6G and 6H). Accordingly, at 76 hpf, RGCs were

partially but significantly recovered in *strip1^{rw147}* mutants injected with MO-jun,

compared to *strip1^{rw147}* mutants injected with a standard control morpholino (Figures 6I

and 6J). Taken together, Strip1 interacts with Strn3 to suppress Jun-mediated apoptotic

signaling in RGCs.

Bcl2 rescues RGC survival in *strip1* mutants, but surviving RGCs do not project

their dendrites to the IPL

The anti-apoptotic B-cell lymphoma 2 (Bcl2) is a key regulator of mitochondria-

dependent apoptosis in neurons, including RGCs, both during survival and in response

to injury (Anilkumar and Prehn, 2014, Bahr, 2000, Bonfanti et al., 1996, Maes et al.,

2017). In addition, JNK/c-Jun activation induces neuronal apoptosis by modulating

BCL2 family proteins (Guan et al., 2006, Hollville et al., 2019, Whitfield et al., 2001).

Thus, we combined *strip1^{rw147}* mutants with the transgenic line *Tg[hsp:mCherry-Bcl2]*,

which overexpresses mCherry-tagged Bcl2 protein under control of a heat shock

promoter (Nishiwaki and Masai, 2020). Bcl2 overexpression significantly inhibited RGC

apoptosis in *strip1^{rw147}* mutants (Figures 7A and 7B). Accordingly, at 78 hpf, RGCs were

partially but significantly recovered in *strip1^{rw147}* mutants overexpressing Bcl2, compared

to non-transgenic mutants (Figures 7C and 7D). Thus, loss of RGCs in *strip1^{rw147}*

mutants depends on the mitochondria-mediated apoptotic pathway.

Surprisingly, *strip1^{rw147}* mutants overexpressing Bcl2 still displayed IPL defects.

In this case, the IPL was not formed at the interface between surviving RGCs and ACs,

but instead, thin IPL-like neuropil was ectopically formed in the intermediate position of

presumptive ACs (Figure 7C). Thus, a fraction of presumptive ACs were abnormally

located between surviving RGCs and IPL-like neuropil, although this AC fraction does

not intermingle with surviving RGCs (Figure 7C, bottom panels, asterisks). In addition,

surviving RGCs in *strip1^{rw147}* mutants apparently fail to project their dendrites to the IPL-

like thin neuropil. Upon closer examination, few surviving RGCs successfully innervate

the IPL, and such areas show less infiltration of ACs (Figure 7C, bottom panels, arrow

heads). These data suggest an additional role of Strip1 in dendritic patterning of RGCs,

which is likely to prevent ectopic IPL-like neuropil formation in the AC layer.

Discussion

Over the past decade, Strip1/Strip has emerged as an essential protein in embryonic development (Bazzi et al., 2017, La Marca et al., 2019, Neal et al., 2020, Sakuma et al., 2014, Sakuma et al., 2015). Using zebrafish, we demonstrate that Strip1 performs multiple functions in development of inner retinal neural circuit. First, we discovered a novel neuroprotective mechanism governed by Strip1, probably through the STRIPAK complex, to suppress Jun-mediated pro-apoptotic signaling in RGCs during development (Figure 8A). In addition, we demonstrate that Strip1-mediated RGC maintenance is essential for laminar positioning of other retinal neurons and structural integrity of the IPL.

RGCs are the most susceptible retinal neurons to cell death, both during development and in response to injury. Unlike zebrafish retina, in which only 1.06 % of RGCs die during development (Biehlmaier et al., 2001), around 50% of mammalian RGCs undergo apoptotic cell death (Bahr, 2000, Fawcett et al., 1984). Similarly, the survival rate of mouse RGCs following optic nerve injury (ONI) is only ~8% compared to a survival rate of ~75% of zebrafish RGCs (Li et al., 2020, Zou et al., 2013). This suggests that RGC survival signals are more active in zebrafish retina. Recently, many studies are seeking to identify such zebrafish-specific survival mechanisms, with the aim to develop therapy that can prevent death of mammalian RGCs (Chen et al., 2021). Interestingly, loss of zebrafish Strip1 causes an elevated apoptotic stress response profile in embryonic retinas with a degree of overlap with adult RGCs post-ONI. Indeed, 5 out of the 6 overlapping upregulated markers (*jun, atf3, stmn4, sox11b* and *adcyap1b)* are commonly upregulated in retinal transcriptomic studies of mammalian ONI models (Wang et al., 2021). Jun is the canonical target of JNK signaling and JNK/Jun activation is a major cause of axonal injury- or glaucoma-induced RGC death (Fernandes et al., 2012, Fernandes et al., 2013, Syc-Mazurek et al., 2017). Thus, Jun signaling appears to be a common mediator of RGC death among vertebrates. Our findings will open exciting new research avenues to determine whether Strip1-mediated Jun suppression can modulate proapoptotic signaling in adult RGCs of both zebrafish and higher vertebrates.

What is the molecular link between Strip1 and Jun suppression in RGCs? Our
proteomic assays revealed that recruitment of many STRIPAK components is
compromised in *strip1* mutants. Also, we demonstrate a functional interaction between
Strip1 and Strn3 to regulate RGC survival. Recently, several studies on the human
STRIPAK complex found that STRIP1 and STRN3 are organizing centers for the
STRIPAK complex and that their mutant forms compromise complex assembly and
function (Jeong et al., 2021, Tang et al., 2019). Modulation of JNK/Jun signaling by the
STRIPAK complex is supported by several studies. MAP4 kinases activate the JNK
signaling pathway and they are among kinase family members that are recruited and
dephosphorylated by the STRIPAK complex (Fuller et al., 2021, Hwang and Pallas,
2014, Kim et al., 2020, Seo et al., 2020). Moreover, JNK signaling is activated in
STRIP1/2 knockout human cell lines (Chen et al., 2019). Similarly, the interaction
between Strip and CKa (*Drosophila* homolog of Striatins) suppresses JNK signaling in
Drosophila testis (La Marca et al., 2019). Thus, it is likely that Strip1 and Strn3 function
in the context of the STRIPAK complex to modulate JNK/Jun activity, thereby promoting
RGC survival. To our knowledge, this study is the first *in vivo* evidence for a functional
interaction between STRIPAK components and Jun signaling in vertebrates.

There are still many gaps in our knowledge of mechanisms underlying IPL
development. Which molecular cues dictate the laminar positioning of inner retinal
neurons? Which cell types are essential for IPL formation? It has been proposed that
IPL development is a robust process. Upon genetic elimination of different inner retinal
cells, the remaining cells manage to form an IPL-like neuropil (Randlett et al., 2013).
However, it is widely agreed that ACs play the dominant role in IPL initiation. Elegant
time-lapse experiments show that ACs project their neurites to form a proto-IPL (Chow
et al., 2015, Godinho et al., 2005). This presumed IPL guides RGCs to extend dendritic
arbors and stratify, although they are born earlier than ACs (Mumm et al., 2006). Other
studies propose an active role for RGCs in shaping the developing IPL (Kay et al.,
2004). We found that RGC death in *strip1* mutants is strongly linked to abnormal
infiltration of ACs and BPs in the GCL and the perturbed IPL. This coincides with the
phenotype of the *lakritz* mutant, in which similar defects occur when RGC genesis is
inhibited (Kay et al., 2001, Kay et al., 2004). Knockout mice in which atypical Cadherin

Fat3 is absent in both RGCs and ACs, ACs invade the GCL abnormally. However, AC-specific knock out mice do not exhibit such positioning defects (Deans et al., 2011). Therefore, we re-introduce a model proposed by (Kay et al., 2004), in which both RGCs and ACs play distinct roles in shaping the developing IPL. In this model, RGCs provide positional cues for migrating ACs to initiate proper IPL program, whereas ACs subsequently project their dendritic plexus and establish the foundation for a proto-IPL.

So far, the RGC-dependent mechanisms that instruct the laminar positioning of ACs and IPL development remain unknown. However, Bcl2-rescued *strip1* mutants provide valuable new insights on such mechanisms. Bcl2-rescued mutants show defects in dendritic patterns of surviving RGCs, which are associated with an ectopic IPL formed within presumptive ACs. Thus, we propose that RGCs serve dual functions in IPL development (summarized in Figure 8B); (1) RGCs act as a physical barrier that prevents abnormal infiltration of ACs into the GCL and (2) RGCs show dendritic guidance cues that establish interactions between RGCs and ACs for a proper IPL program. It is unclear what guidance cues participate in this process. Possible candidates are N-Cadherin and Semaphorin-3 receptors, Neuropillin-1 (Nrp1) and PlexinA1. A hypomorphic allele of zebrafish *Ncad* mutants compromises IPL formation with abnormal neurite patterning of INL cells (Masai et al., 2003). In *Xenopus*, Nrp1 and PlexinA1 inhibition induced randomly oriented dendritic patterning of RGCs, similar to zebrafish *strip1* mutants (Kita et al., 2013). Future experiments will clarify whether these molecules participate in communication among RGCs and ACs to establish a proper IPL. Lastly, we show that Strip1 may function in neurite patterning of RGCs, and small subsets of ACs and BPs. This is supported by established roles of *Drosophila* Strip in dendritic branching and axon elongation (Sakuma et al., 2014). Future studies on cell-specific Strip1 knockout models will help understand Strip1 function in retinal neurite morphogenesis.

In summary, we demonstrate that a series of Strip1-mediated regulatory mechanisms constructs retinal neural circuit through RGC survival and neurite patterning of retinal neurons. Our findings provide valuable insights to mechanistic understanding of JNK/Jun-mediated apoptotic pathway and correct assembly of synaptic neural circuits in the brain. For medical perspective, our findings on a similar

432   stress response between zebrafish *strip1* mutants and mammalian ONI models pave

433   the way for future research on potential Strip1-mediated therapeutic targets that could

434   help mitigate RGC degeneration in glaucoma and optic neuropathies.

Author contributions

Conceptualization: MA, IM; Methodology: MA, IM; Software: MA; Validation: MA; Formal analysis: MA; Investigation: MA, YK, IM; Resources: MA, IM; Data curation: MA, IM; Writing - original draft: MA; Writing - review & editing: MA, IM; Visualization: MA, IM; Supervision: IM; Project administration: MA, IM; Funding acquisition: IM.

Acknowledgments

We are grateful to Rachel Wong for providing the pZNYX-GalVP16 construct and *Tg[UAS:MYFP]* fishline and Francesco Argenton for the pG1[ptf1a:GFP] construct. We thank lab members for supporting experiments, especially Yuki Takeuchi, Tetsuya Harakuni and Yuko Nishiwaki. We thank Fumiyasu Imai, Asuka Yoshizawa, and Sachihiro Suzuki for supporting cloning experiments and fishline generation. We are grateful to Alejandro Villar Briones in the Instrumental analysis section and the sequencing section of the Research Support Division of OIST for supporting mass spectrometry and RNA sequencing, respectively. This work was funded by a grant from the Okinawa Institute of Science and Technology Graduate University to IM.

Declaration of interests

The authors declare no competing interests.

457 **Materials and methods**

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
genetic reagent (*Danio rerio*)	*Tg[hsp:gal4]^{kca4}*	(Scheer et al., 2002)	ZDB-ALT-020918-6	
genetic reagent (*Danio rerio*)	*Tg[UAS:EGFP]*	(Koster and Fraser, 2001)	N/A	
genetic reagent (*Danio rerio*)	*Tg[UAS:MYFP]*	(Schroeter et al., 2006)	N/A	
genetic reagent (*Danio rerio*)	*Tg[ath5:GFP] ^{rw021}*	(Masai et al., 2003)	N/A	
genetic reagent (*Danio rerio*)	*Tg[ptf1a:mCherry-CAAX]^{oki067}*	This paper	N/A	
genetic reagent (*Danio rerio*)	*Tg[Gal4-VP16,UAS:EGFP]xf z43* or *xfz43*	(Zhao et al., 2009)	ZDB-ALT-100201-1	
genetic reagent (*Danio rerio*)	*Tg[Gal4-VP16,UAS:EGFP]xf z3* or *xfz3*	(Zhao et al., 2009)	ZDB-ALT-100201-2	
genetic reagent (*Danio rerio*)	*Tg[hs:mCherry-tagged Bcl2]^{oki029}*	(Nishiwaki and Masai, 2020)	ZDB-ALT-210524-5	
genetic reagent (*Danio rerio*)	*Tg[hsp:WT.Strip1-GFP]^{oki068}*	This paper	N/A	
genetic reagent (*Danio rerio*)	*Tg[hsp:Mut.Strip1:G FP]^{oki069}*	This paper	N/A	
genetic reagent (*Danio rerio*)	*strip1^{rw147}*	This paper	N/A	
genetic reagent (*Danio rerio*)	*strip1$^{crispr\Delta10}$* or *strip1^{oki8}*	This paper	N/A	
genetic reagent (*Danio rerio*)	*roy*	(D'Agati et al., 2017)	ZDB-GENE-040426-1168	
antibody	Mouse anti-acetylated α-tubulin	Sigma-Alrich	T6793	1:1000

antibody	Rabbit anti-Pax6	BioLegned	901301	1:500
antibody	Rabbit anti-Prox1	Genetex	GTX128354	1:500
antibody	Mouse anti-PCNA	Sigma-Aldrich	P8825	1:200
antibody	Mouse Zpr1	ZIRC	ZDB-ATB-081002-43	1:100
antibody	Mouse Zpr3	ZIRC	ZDB-ATB-081002-45	1:100
antibody	Mouse anti-glutamine synthetase	Sigma-Alrich	MAB302	1:150
antibody	Mouse Zn5	ZIRC	ZDB-ATB-081002-19	1:50
antibody	Rabbit anti-p-Jun	Cell Signal	9164S	1:100
antibody	Rabbit anti-Strip1	This paper	N/A	IF: 1:1000 WB: 1:500
antibody	Goat anti-rabbit Alexa 488 secondary antibody	Life Technologies	A11034	1:500
antibody	Goat anti-mouse Alexa 488 secondary antibody	Life Technologies	A11029	1:500
antibody	Goat anti-mouse Alexa 546 secondary antibody	Life Technologies	A11030	1:500
antibody	Goat anti-mouse Alexa 647 secondary antibody	Life Technologies	A21236	1:500
antibody	Anti-rabbit IgG, HRP-linked Antibody	Cell Signalling	7074	1:5000
antibody	Rabbit anti-GFP	Thermo Fisher Scientific	A11122	1:500
antibody	Rabbit anti β-actin	Abcam	AB8227	1:5000
recombinant DNA reagent	pTol2[ptf1a:mCherry-CAAX]	This paper	N/A	

recombinant DNA reagent	pG1[ptf1a:GFP]	Francesco Argenton Laboratory (Jusuf et al., 2012)	N/A	
recombinant DNA reagent	pTol2[hsp:WT.Strip1-GFP]	This paper	N/A	
recombinant DNA reagent	pTol2[hsp:Mut.Strip1:GFP]	This paper	N/A	
recombinant DNA reagent	pBluescript SK (+)	Stratagene	N/A	
recombinant DNA reagent	pT2AL200R150G	(Urasaki et al., 2006)	N/A	
recombinant DNA reagent	pB[ath5:Gal4-VP16]	This paper	N/A	
recombinant DNA reagent	pZNYX-GalVP16	Rachel Wong Laboratory (Schroeter et al., 2006)	N/A	
sequence-based reagent	Standard control MO (STD-MO)	GeneTools	N/A	5'-CCTCTTACCTCAGTTACAATTTATA-3'
sequence-based reagent	Strip1 morpholino (MO-strip1)	GeneTools	N/A	5'-TAGCACATAAACCGACACCGTCCAT-3'
sequence-based reagent	Ath5 morpholino (MO-ath5)	GeneTools	ZDB-MRPHLNO-100405-2	5'-TTCATGGCTCTTCAAAAAAGTCTCC-3'
sequence-based reagent	Striatin3 morpholino (MO-strn3)	GeneTools	N/A	5'-CCTGCTAGAAGTCGCCGATTGTTAC -3'
sequence-based reagent	Jun morpholino (MO-jun)	GeneTools	ZDB-MRPHLNO-080908-1	5' -CTTGGTAGACATAGAAGGCAAAGCG - 3'
peptide, recombinant protein	Cas9 protein	FASMAC	GE-006-S	

commercial assay or kit	JB-4® Embedding Kit	Polysciences	00226-1	
commercial assay or kit	In Situ Cell Death Detection Kit, TMR Red	Roche	12156792910	
commercial assay or kit	In Situ Cell Death Detection Kit, Fluorescein	Roche	11684795910	
commercial assay or kit	DIG RNA Labeling Kit	Roche	11277073910	
commercial assay or kit	GFP Trap Agarose	Chromotek	gta-20	
commercial assay or kit	Arcturus PicoPure RNA Isolation Kit	Thermo Fisher Scientific	KIT0204	
commercial assay or kit	NEB Next® Ultra™ II Directional RNA Library Prep Kit	New England BioLabs	E7760L	
chemical compound, drug	Acridine Orange (AO)	Nacalai tesque	1B-307	
chemical compound, drug	CellTrace™ BODIPY™ TR Methyl Ester	Thermo Fisher Scientific	C34556	
chemical compound, drug	Ethyl-3-aminobenzoate de methanesulfonate (Tricaine, MS-222)	Nacalai tesque	14805-82	
chemical compound, drug	PTU (N-Phenylthiourea)	Nacalai tesque	27429-22	
chemical compound, drug	Fast DiO solid	Thermo Fisher Scientific	D3898	
chemical compound, drug	Fast DiI solid	Thermo Fisher Scientific	D7756	
chemical compound, drug	TO-PRO™-3 Iodide (642/661)	Thermo Fisher Scientific	T3605	
chemical compound, drug	Hoechst 33342	Wako	346-07951	
chemical compound, drug	Toluidine Blue	Nacalai tesque	1B-481	

chemical compound, drug	Dextran, Tetramethylrhodamine	Thermo Fisher Scientific	D1817 LTJ	
chemical compound, drug	Dextran, Alexa Flour-488	Thermo Fisher Scientific	D22910	
chemical compound, drug	Dextran, Alexa Flour-647	Thermo Fisher Scientific	D22914	
chemical compound, drug	Dextran, Cascade Blue	Thermo Fisher Scientific	D1976	
software, algorithm	chopchop	chopchop	https://chopchop.cbu.uib.no	
software, algorithm	ImageJ (Fiji)	(Schneider et al., 2012)	https://imagej.nih.gov/ij/; RRID: SCR_003070	
software, algorithm	Imaris	Bitplane	http://www.bitplane.com/imaris; RRID: SCR_007370	
software, algorithm	Proteome Discoverer	Thermo	https://www.thermofisher.com/store/products/OPTON-30945#/OPTON-30945	
software, algorithm	STRING	(Szklarczyk et al., 2017)	https://string-db.org	
software, algorithm	Metascape	(Zhou et al., 2019)	https://metascape.org	
software, algorithm	R v3.4.2 , RStudio interface	(Team, 2016)	https://string-db.org	
software, algorithm	BioVenn	(Hulsen et al., 2008)	http://www.biovenn.nl/	
software, algorithm	Graphpad Prism v9.1.0.	Graphpad Prism	https://www.graphpad.com/scientific-software/prism/	

 Ethics statement

Zebrafish (*Danio rerio*) were maintained on a 14:10 hour light: dark cycle at 28°C.

Collected embryos were cultured in E3 embryo medium (5mM NaCl, 0.17mM KCl,

0.33mM CaCl$_2$, 0.33mM MgSO$_4$) containing 0.003% 1-phenyl-2-thiouera (PTU) to

prevent pigmentation and 0.01% methylene blue to prevent fungal growth. All

experiments were performed on zebrafish embryos between 36 hpf and 4 dpf prior to

sexual differentiation. Therefore, sexes of the embryos could not be determined.

All zebrafish experiments were performed in accordance with the Animal Care and Use

Program of Okinawa Institute of Science and Technology Graduate School (OIST),

Japan, which is based on the Guide for the Care and Use of Laboratory Animals by the

National Research Council of the National Academies. The OIST animal care facility

has been accredited by the Association for Assessment and Accreditation of Laboratory

Animal Care (AAALAC International). All experimental protocols were approved by the

OIST Institutional Animal Care and Use Committee.

Transgenic fish lines

To visualize RGCs, the transgenic line *Tg[ath5:GFP]^{rw021} (Masai et al., 2003)* was used.

In this line, GFP is expressed under control of the *ath5* (also referred to as *atoh7*)

promoter. The transgenic lines *Tg(Gal4-VP16,UAS:EGFP)xfz43* or *xfz43* and *Tg(Gal4-*

VP16,UAS:EGFP)xfz3 or *xfz3* (Zhao et al., 2009) are enhancer trap lines and were both

used to label distinct populations of BPs. To visualize ACs, the transgenic line

Tg[Ptf1a:mCherry-CAAX]^{oki067} was generated by injecting the plasmid

pTol2[ptf1a:mCherry-CAAX] into one-cell-stage fertilized eggs, together with Tol2

transposase mRNA.

For Bcl2 overexpression experiments, the line *Tg[hs:mCherry-tagged Bcl2]^{oki029}*

(Nishiwaki and Masai, 2020) was employed, in which mCherry-tagged Bcl2 at the N-

terminus is overexpressed under control of the heat shock promoter. For over-

expression of wild-type and *rw147* mutant forms of zebrafish Strip1, the lines

Tg[hsp:WT.Strip1-GFP]^{oki068} and *Tg[hsp:Mut.Strip1:GFP]^{oki069}* were generated,

respectively, to express GFP-tagged Strip1 at the C-terminus under control of the heat

shock promoter. For line generation, the DNA constructs *pTol2[hsp:WT.Strip1-GFP]* and

pTol2[hsp:Mut.Strip1-GFP] were injected into one-cell-stage fertilized eggs together with

Tol2 transposase mRNA. These injected F0 embryos were bred up to the adult stage

and used to identify founder fish that produce F1 generation embryos showing stable

GFP expression. Transgenic lines were established in F2 generation. The transgenic

line *Tg[hsp:Gal4;UAS:EGFP]* was generated by combining *Tg[hsp:gal4]^{kca4}* (Scheer et

al., 2002) with *Tg[UAS:EGFP] (Koster and Fraser, 2001)* to express EGFP under

control of the heat shock promoter, while *Tg[UAS:MYFP]* expresses EYFP fused to the

membrane targeting palmitoylation signal of gap43 under the control of 14XUAS E1b

promoter (Schroeter et al., 2006). Some of the mentioned transgenic lines were

combined with the mutant line *strip1^{rw147}*. The steps of plasmid construction are

mentioned below in details.

Mutant line generation, mutant Identification, and genotyping

The *strip1^{rw147}* mutant line was generated from a mutagenesis screen (Masai et al.,

2003) that used RIKEN Wako (RW) as a wild-type strain. Mutation mapping and

subsequent experiments were carried out in the genetic background of WIK and

Okinawa wildtype (oki), respectively. The *rw147* mutation was mapped on a genomic

region in chromosome 22 flanked by two self-designed polymorphic markers;

AL928817-12:   (5'-TTCAACATCTGCTTTTCCTCCT-3'   and

5'TCATGTCCCAGAAATCACACAT3')   and   zk253D23-4   (5'-

CATTCTTCATTAAAGAGATCAGTGTGA-3' and 5'-AGTGATCACACACCCCCACT-3').

In addition, location of the *rw147* mutation was further restricted using another self-

designed polymorphic marker Zk286J17-3 (5'- TTCACATTTACATTTTTCTGAACATTT-

3' and 5'-CACACAGCCTTCTCTTGCAC -3') as no recombination was detected.

From 3 dpf, *strip1^{rw147}* homozygous mutants (*strip1$^{-/-}$*) are distinguished from wild-type

siblings (*strip1$^{+/+}$* or *strip1$^{+/-}$*) by external morphology since they exhibit cardiac edema,

an abnormal lower jaw, and smaller eyes. From 54 to 72 hpf, *strip1^{rw147}* homozygous

mutants are screened by Acridine Orange (AO) live staining to detect apoptotic cells

(Casano et al., 2016) or Bodipy TR live staining to visualize lamination defects (Choi et

al., 2010). Prior to 54 hpf, genotyping of *strip1^{rw147}* mutant embryos was performed by

sequencing.   The   primer   set,   5'-CGTGTGTTTTCAGGGTGTT-3'   and   5'-

TCACCATCCCAAACAGCATA-3', was used. The 257-bp PCR amplicons were
amplified using Phusion Hot Start II (Thermo Fisher Scientific) and sequenced for
genotyping.

The *strip1^crisprΔ10^* (officially referred to as *strip1^oki8^*) mutant line was generated using
Crispr/Cas9 gene editing technology. The gRNA sequence, 5'-
CCCGCGTCCGCCTCTGACCTCAT-3', was designed using chopchop
(https://chopchop.cbu.uib.no) and it targets exon 9 of *strip1* gene. One-cell-stage
embryos were injected with 200ng/µL gRNA and 500ng/µL Cas9 protein (FASMAC). F1
mutant founders were identified by sequencing. A 10-bp deletion was introduced at
nucleotide 932 of the Strip1 coding sequence, resulting in a frameshift at amino acid
313 and a premature stop at amino acid 330. The primer set 5'-
CGTTCCAAATCATTGAAACAGA-3' and 5'-TGTTTGTGATGTGTTGACCTTG-3' was
used for genotyping. PCR amplicons were run on 15% polyacrylamide gel for
identification of wild-type siblings and mutants.

All generated transgenic and mutant lines were combined with the zebrafish
pigmentation mutant, *roy orbison (roy) (D'Agati et al., 2017)* to remove iridophores and
enhance live imaging.

Molecular cloning

To generate pTol2[hsp:WT.Strip1-GFP] and pTol2[hsp:Mut.Strip1:GFP], a PCR strategy
was used to amplify ~2.5 kb Strip1 cDNA from total cDNA of 4 dpf wild-type and
*strip1^rw147^* zebrafish embryos using the primers 5'-
AGACTTGTGTCAGCGTGACGCGAG-3' and 5'-
ACTCTAGCAAGTGTAGTGTTGTTGATG-3'. Then, using a Gibson Assembly Cloning
Kit, a *strip1* cDNA fragment was cloned into a Tol2 transposon vector pT2AL200R150G
(Urasaki et al., 2006) at the *XhoI* and *ClaI* sites with a heat-shock inducible promoter
(hsp) at the N-terminus (Halloran et al., 2000) and a GFP tag at the C-terminus
(separated by a linker sequence, CTCGAGGGAGGTGGAGGT). For
pTol2[ptf1a:mCherry-CAAX] construction, pG1[ptf1a:GFP] was used as donor plasmid,
which was kindly provided by the Francesco Argenton lab. A 5.5-kb fragment of the
ptf1a promoter sequence was retrieved at *HindIII* and *SmaI* sites and inserted into a

pBluescript SK (+) (Stratagene) shuttle vector upstream of the membrane-targeting
mCherry-CAAX sequence. Then, the ptf1a:mCherry-CAAX sequence was inserted into
the *XhoI and BglII* sites of pT2AL200R150G. The pB[ath5:Gal4-VP16] plasmid was
constructed by inserting a 6.6-kb fragment of the ath5 5'-enhancer/promoter region
(including 5' UTR) into the *BamHI* site of the pB[Gal4-VP16] plasmid provided by Dr. R.
Köster (Koster and Fraser, 2001).

In vivo **cell labeling**
Single-cell mosaic labeling to visualize the morphology of RGCs was done by injecting
20 ng/μL of pB[ath5:Gal4] into 1-cell-stage embryos from intercrosses of *strip1^{rw147}*
heterozygous fish combined with *Tg[UAS:MYFP]* (Schroeter et al., 2006). Likewise,
pZNYX-GalVP16, a kind gift from the Rachel Wong Laboratory, was injected to visualize
ON-BPs (Schroeter et al., 2006). Single AC labeling was performed by injecting the
DNA construct pG1[ptf1a:GFP] into 1-cell-stage embryos from intercrosses of *strip1^{rw147}*
heterozygous fish at concentration 20 ng/μL (Jusuf et al., 2012).

Morpholino knockdown assay
Embryos produced by intercrosses of wild-type or *strip1^{rw147}* heterozygous fish were
injected with antisense morpholino oligonucleotides at 1-cell stage. MO-strip1, MO-
strn3, and MO-ath5 (Ranawat and Masai, 2021) were injected at a concentration of 250
μM, whereas MO-jun (Han et al., 2016) was injected at a concentration of 125 μM. For
each morpholino experiment, the same concentration of the standard control
morpholino (STD-MO) was used as a negative control. Detailed morpholino sequences
are listed in the Key resources table.

DiI/DiO Injections
To trace the RGC axon projections into the optic tectum, 3-dpf embryos were fixed in
4% paraformaldehyde (PFA) and after washing with PBS several times, were injected
with 2 mg/mL of the lipophilic dyes, DiI and DiO, in the area between the lens and
retina. Large injection volumes were applied to label all RGCs. Embryos were incubated
overnight at 4°C, and then mounted in 75% glycerol for confocal imaging.

Histological methods

Zebrafish embryos were embedded for JB4 plastic sectioning and toluidine blue counterstaining, as previously described (Sullivan-Brown et al., 2011). Immunolabeling of cryosections and paraffin sections were carried out according to standard protocols (Imai et al., 2010, Masai et al., 2003). An antigen retrieval step was performed on paraffin sections by heating in 10 mM citrate buffer pH 6.0 for 5 min at 121°C. TUNEL (terminal deoxynucleotidyl transferase dUTP nick end labeling) was performed using an In Situ Cell Death Detection Kit (Roche) according to the manufacturer's protocol. Nuclear staining was carried out using 1nM TOPRO3 or 1ng/mL Hoechst 33342.

Antibodies used in this study and their dilutions are as follows: anti-acetylated α-tubulin (1:1000), anti-Pax6 and anti-Prox1 (1:500), anti-PCNA (1:200), anti-glutamine synthetase (1:150), antibodies against zpr1 and zpr3 (1:100), zn5 antibody (1:50) and anti-p-Jun (1:100). Antibody against the peptide sequence of zebrafish Strip1 (amino acids 344-362: EKDPYKADDSHEDEEENDD) was generated using a synthetic peptide and used for immunostaining at 1:1,000. For adsorption control, purified antibody was preincubated with 3.6 μg/mL of corresponding blocking peptide for 1 h at room temperature. Secondary antibodies employed in this study were: Alexa488, 546 and 647 fluorophore-conjugated secondary antibodies used at a concentration of 1:500.

For whole-mount immunostaining against acetylated α-tubulin, 3-dpf embryos were fixed at room temperature for 3 h in 2% trichloroacetic acid (TCA). Then, embryos were washed in PBTr (Westerfield, 1995) (PO$_4$ buffer (0.1M, pH7.3)+0.1% Triton X-100) followed by permeabilization in 0.2% trypsin for 4 min at 4^{o}C. After washing, a post-fixation step in 4% PFA for 5 min at 4^{o}C was applied. Next, blocking was done in 10% goat serum in PBTr for 1 h at room temperature followed by incubation in mouse anti-acetylated α-tubulin in 1% goat serum/PBTr overnight at 4^{o}C. After washing, embryos were incubated in secondary antibody diluted in 1% goat serum in PBTr overnight at 4^{o}C. Whole-mount immunostaining against p-Jun was performed following standard protocols (Ungos et al., 2003). After staining, embryos were mounted in 75% glycerol for confocal imaging.

Whole-mount *in situ* hybridization was performed on wild-type zebrafish embryos at specific developmental stages as previously described (Xu et al., 1994). Hybridization was performed overnight at 65^{o}C using *strip1* RNA probe at the concentration 2.5 ng/µL in hybridization buffer. *strip1* probe synthesis was performed according to standard protocols (Thisse and Thisse, 2008). Template regions were amplified from *strip1* cDNA using the primers, 5'-AATGCTGCCGAATAAAATGCGAG-3' and 5'-CCCAGAGTGAACAGGATGCTCT-3'. Antisense and sense probes were synthesized by *in vitro* transcription using a DIG RNA Labeling Kit (Roche). Following labeling, whole embryos were mounted in 75% glycerol for imaging. To visualize expression patterns in the retina, cryosections were prepared from whole mount embryos post-hybridization.

Live staining

To visualize lamination patterns, live staining of retinal landmarks was performed by incubating live zebrafish embryos in 100 nM solution of Bodipy TR methyl ester (Thermo Fisher Scientific) in E3 embryo rearing media for 1 h at room temperature following the manufacturer's protocol. To examine DNA condensation of apoptotic cells in the GCL, zebrafish embryos were incubated for 30 min in 5 µg/mL of AO stain dissolved in egg water. Following staining, embryos were extensively washed with egg water and observed using epifluorescence or imaged using confocal microscopy.

Overexpression experiments

For rescue experiments, the wild-type form of Strip1, the *rw147* mutant form of Strip1 or Bcl2 was overexpressed in *strip1^{rw14}* mutants by heat shock treatment using the transgenic lines *Tg[hsp:WT.Strip1-GFP]*, *Tg[hsp:Mut.Strip1:GFP]* and *Tg[hs:mCherry-tagged Bcl2]*, respectively. To perform heat shock, embryos from heterozygous intercrosses were incubated for 1 h at 39^{o}C starting from 27-30 hpf and applied every 12 hours until the designated timepoints. For Co-IP/MS, heat shock was applied to embryos from intercrosses of wild-type zebrafish combined with *Tg[hsp:WT.Strip1-GFP]*, *Tg[hsp:Mut.Strip1:GFP] or Tg[hsp:Gal4;UAS:EGFP].* After screening for transgenic embryos, embryos were either fixed in 4% PFA for histological assays or processed for protein extraction.

Cell transplantation assays

Single-cell transplantations were performed at blastula stage, as previously described (Kemp et al., 2009). Genotypes of donor and host embryos were determined at 3-4 dpf based on morphological phenotype or they were genotyped at earlier time points by sequencing or AO live-staining of apoptotic cells. To trace transplanted donor cells in host retinas, 2-5% lysine-fixable dextran rhodamine, Alexa-488 dextran, Alexa-647 dextran or cascade blue dextran were injected in 1-2-cell-stage donor embryos, depending on study design. To assess the cell autonomy of Strip1 in RGC death, donor embryos from intercrosses of $strip1^{rw147}$ heterozygous fish were transplanted into wild-type host embryos. Host embryos with successful retinal transplants were fixed in 4% PFA at 60 hpf and processed for TUNEL labeling. To assess the cell autonomy of Strip1 in AC or BP development, donor embryos from intercrosses of $strip1^{rw147}$ heterozygous fish combined with *Tg[ptf1a:mCherry-CAAX]* or *Tg[Gal4-VP16,UAS:EGFP]xfz3/xfz43* were transplanted into embryos from intercrosses of $strip1^{rw147}$ heterozygous fish, live imaging of host retinas with successful transplants was done at 3-4 dpf to assess the morphology of donor ACs labeled with mCherry or donor BPs labeled with EGFP. To visualize the retinal lamination phenotype, some hosts were stained with Bopidy TR live stain prior to imaging.

Microscopy

Imaging of toluidine blue-stained sections and retinal sections following *in situ* hybridization was performed using a Zeiss upright Axioplan2 equipped with an AxioCam HRC camera, while imaging of whole-mount *in situ* hybridization embryos was done using a Keyence BZ-X700. An inverted Zeiss LSM 780 was used to scan immunostained retinal sections with a 40x / 1.40 Plan-Apochromat Oil objective and whole-mount immunostained embryos using a 40x / 1.0 Plan-Apochromat water objective. Glycerol-mounted embryos were placed on glass-bottom depression slides for scanning. For live imaging, zebrafish embryos were anesthetized using 0.02% tricaine (3-amino benzoic acid ethyl ester) dissolved in E3 embryonic medium and mounted laterally in 1% low-melting agarose. Image acquisition of embryo retina was

carried out using an upright Zeiss LSM 710 with a 40x / 1.0 W Plan-Apochromat
objective or an upright Fluoview FV3000 (Olympus) confocal microscope with a 40x /
0.8 water immersion objective.

To perform time-lapse imaging of retinal development, several embryos from
intercrosses of $strip1^{rw147}$ heterozygous fish carrying the transgenes
Tg[ath5:GFP;ptf1a:mCherry-CAAX] were mounted simultaneously in a culture dish
covered with E3 embryonic medium containing 0.003% PTU and 0.02% tricaine and
overlayed with a thin layer of mineral oil to prevent evaporation of E3 medium to
minimize embryotoxicity. Retinal z-stacks were acquired consecutively in 1-µm steps
every ~2 hrs, starting at 48 hpf with undetermined genotypes. Scanning was done using
the Multi Area Time Lapse (MATL) Software module of the FV3000 (Olympus) confocal
microscope and a motorized XYZ-rotation stage.

All Images were processed using ImageJ (NIH, v2.1.0/1.53C), Imaris (Bitplane, v9.1.2)
and Adobe Illustrator software. 3D rendering and analysis of time-lapse movies were
performed on Imaris software. Whenever necessary, brightness and contrast display
levels for the whole image were adjusted to aid visualization or decrease background
noise.

Western Blotting and Co-immunoprecipitation (CO-IP)

Heads of non-injected or MO-strip1-injected wild-type embryos were dissected at 2 dpf
in Leibovitz's L-15 ice cold medium and homogenized in lysis buffer (125mM NaCl, 50
mM Tris (pH 7.5), 0.5 mM EDTA (pH 8), 1% Triton X-100 and 1× cocktail protease
inhibitors). Lysates were clarified by centrifugation at 10,000 g for 10 min at 4 °C. Equal
amounts of denatured clarified lysates were run on 10% Mini-PROTEAN TGX gels for
SDS-PAGE and transferred to PVDF membranes using Trans-Blot Turbo PVDF
Transfer system. After blocking with 5% skim milk in 0.1% tween-20 in TBS,
immunoblotting was performed using anti-Strip1 (1:500). HRP-linked rabbit IgG was
used as a secondary antibody. Chemiluminescence signals were detected using a FUJI
Las 4000 luminescence image analyzer.

For co-immunoprecipitation, wild-type embryos carrying the transgenes
Tg[hsp:WT.Strip1-GFP], Tg[hsp:Mut.Strip1-GFP] or Tg[hsp:Gal4;UAS:GFP] were

exposed to 12-hr interval heat shocks starting at 27 hpf. At 2 dpf, lysates for each

biological replicate were prepared from a pool of around 150 embryo heads in NP-40-

based lysis buffer (150mM NaCl, 10 mM Tris (pH 7.5), 0.5 mM EDTA (pH 8), 0.5% NP-

40 and 1× cocktail protease inhibitors), as described above. Immunoprecipitation was

performed on clarified lysates using anti-GFP (GFP-Trap agarose beads, Chromotek)

according to the manufacturer's protocol. Briefly, clarified lysates were diluted in wash

buffer (150mM NaCl, 10 mM Tris (pH 7.5), 0.5 mM EDTA (pH 8) and 1× cocktail

protease inhibitors) to reach 0.1% NP-40. Then, lysates were incubated with pre-

equilibrated GFP-Trap beads for 1 h at 4°C. Afterwards, beads were collected by

centrifugation and washed in wash buffer 5 times. To confirm that GFP-fused proteins

were successfully pulled down, proteins were eluted from beads by boiling in 1x sample

buffer for 5 mins. Then, 5% of pre-pulldown lysate (input) and 10% of the pulled-down

proteins were run for western blotting as described above using anti-GFP (1:500).

Mass spectrometry (MS) and data analysis

To prepare protein samples for MS analysis, immunoprecipitated protein complexes

were eluted from GFP-Trap beads using an on-bead trypsin-based digestion protocol

according to manufacturer's protocol. Digestion was performed overnight at 32°C and

under rotation at 400 rpm. Afterwards, digested peptides were cleaned and desalted

using C_{18} stage tips, as previously described (Rappsilber et al., 2007). Eluted peptides

were vacuum-dried and re-constituted in 1% acetic acid, 0.5% formic acid for MS

analysis using an Orbitrap-Fusion Lumos mass spectrometer coupled to a Waters

nanoACQUITY Liquid Chromatography System. Samples were trapped on a

nanoACQUITY UPLC 2G-V/M Trap 5 μm Symmetry C_{18}, 180 μm × 20 mm column

and analytical separation was performed on a nanoACQUITY UPLC HSS T3 1.8 μm,

75 μm × 150 mm column. Peptides were fractionated over a 60-min gradient from

1-32% acetonitrile with 0.1% formic acid. Solvent flow rate was 500 nL/min and column

temperature was 40°C.

LC-MS raw data files were analyzed using Proteome Discoverer (PD, v.2.2, Thermo

Fisher Scientific). The SEQUEST algorithm was used to match MS data to the *Danio*

rerio (zebrafish) database downloaded from UniProt (July, 2021) and the common

Repository of Adventitious Protein (cRAP, https://www.thegpm.org/crap). Database
search parameters included carbamidomethylation of cysteine as fixed modification
and oxidation of methionine, deamidation of glutamine and asparagine as dynamic
modifications. Trypsin was specified as a cleavage enzyme with up to 2 missed
cleavages. Normalization was performed based on specific protein amount (Trypsin)
and proteins were filtered based on a false discovery rate of q < 0.05. Abundance ratios
were generated for wild-type compared to mutant and wild-type compared to GFP
control. Enriched proteins with an abundance ratio ≥ 2 and adjusted p-value <0.05 were
considered statistically significant. Search Tool for the Retrieval of Interacting
Genes/Proteins (STRING, v11.0) was used to visualize the Strip1-interaction network
with the enriched proteins and calculate protein-protein interaction (PPI) value
(Szklarczyk et al., 2017).
748
RNA sequencing and analysis
Total RNA was isolated from 4 independent biological replicates of 60-64 hpf wild-type
siblings and $strip^{rw147}$ mutant eye cups using a PicoPure RNA Isolation Kit (Thermo
Fisher Scientific) according to the manufacturer's instructions. Each biological replicate
represented a pool of 20-30 eyes. All samples had RNA integrity number (RIN) values
greater than 8.5. The purified RNA was used for Poly(A)-selected mRNA library
preparation with a NEBNext® Ultra™ II Directional RNA Library Prep Kit for Illumina and
sequenced on a NovaSeq6000 SP to generate 150 bp paired-end reads. Sequencing
reads were quality checked using FastQC (Andrews, 2010) and trimmed with FastP
(Chen et al., 2018). The resulting reads were mapped using hisat v2.1.0 (Kim et al.,
2015) to the zebrafish reference genome (GRCz11). Mapped reads were counted using
featureCounts, v1.6.2 (Liao et al., 2014) and differential gene expression analysis
(Mutant vrs. Wildtype) was carried out on the counts files using the EdgeR package,
v3.32.1 (Robinson et al., 2010) in RStudio, v1.4.1106 (Team, 2016). Genes with FDR
<0.05 and $Log_2FC > |1|$ were considered statistically significant. EnhancedVolcano,
v1.8.0 (Blighe et al., 2019) and pheatmap, v 1.0.12 (Kolde, 2012) packages were used
to generate the volcano plot and heatmap, respectively. Gene Ontology analyses were
performed using Metascape with *D. Rerio* as the input species and *M. Musculus* as the

analysis species (Zhou et al., 2019). To analyze published scRNA sequencing data of

zebrafish retina at 48 hpf, raw count matrices were analyzed with the Seurat package,

v4.0.1 (Satija et al., 2015), as previously described (Xu et al., 2020). Clustering results

were visualized using Uniform Manifold Approximation and Projection (UMAP). The

BioVenn web application was used to generate Venn diagrams to compare Upregulated

DEGs in this study with published upregulated DEGs in optic nerve injury models

(Hulsen et al., 2008).

Quantification and statistical analysis

To quantify RGC area, masks were generated for areas with strong ath5+ signals in the

retina and quantified using the Color Threshold tool in ImageJ (Schneider et al., 2012).

Afterwards, the retinal outline was defined using the lasso tool and retinal area was

calculated. Data were represented as the percentage of ath5+ area to total retinal area.

To calculate the area of p-Jun signal in RGCs, an ROI containing the GCL was defined

based on zn5 antibody signal (antigen: alcama, previously referred to as DM-GRASP)

to exclude noise at the retina boundary. Areas of p-Jun and zn5 signals were calculated

as described above and data were represented as the percentage of p-Jun+ area to

zn5+ area.

To quantify apoptotic cells, the number of TUNEL+ cells in GCL or retina was calculated

manually within a single retinal section. To quantify the number of ptf1a+ cells that

contribute to IPL formation (ACs), cells were manually counted in a unified area (8,500

μm^2) across all samples. Ptf1a+ cells that contributed to the OPL (presumably HCs)

were excluded from quantification. To quantify the migration patterns of ptf1a+ cells that

contribute to the IPL, cells located at the apical side relative to the IPL were assigned

INL+, while cells located at the basal side of the IPL or near the lens were assigned

GCL+. To determine the distribution of ptf1a+ cells within transplanted columns, ptf1a+

cells that contributed to the IPL were manually calculated in a z-stack and the

distribution pattern was represented as the percentage of basally or apically located

ptf1a+ cells to the total number of ptf1a+ cells. Numbers of strong Pax6+ and Prox1+

cells were calculated using the analyze particles tool in ImageJ and the distribution of

cells (INL+ or GCL+) was assigned based on their location relative to the IPL, according

to the nuclear stain pattern. Distribution data were represented as the percentage of

INL+ or GCL+ to the total number of Prox1+ or Pax6+ cells.

Statistical analysis was conducted using Graphpad Prism 9.1.0. Data are represented

as means ±SD. Comparisons between two samples were done using the Mann Whitney

U test or Student's t test with Welch's correction for normally distributed data. For

multiple comparisons, 2-way ANOVA with the Tukey post hoc test was used. Details of

statistical tests and number of samples used are in figures and figure legends.

Significance level is indicated as $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$,

n.s. indicates not significant.

Data availability

Raw RNA-seq data files have been deposited in DDBJ under accession number

DRA012640.

Mass spectrometry raw data and result files have been deposited in the

ProteomeXchange Consortium (http://proteomecentral.proteomexchange.org) via the

jPOST partner repository (https://jpostdb.org) (Okuda et al., 2017) under accession

number PXD028131.

References

AMINI, R., ROCHA-MARTINS, M. & NORDEN, C. 2018. Neuronal Migration and Lamination in the Vertebrate Retina. *Frontiers in Neuroscience,* 11.

ANDREWS, S. 2010. FastQC: A Quality Control Tool for High Throughput Sequence Data.

ANILKUMAR, U. & PREHN, J. H. 2014. Anti-apoptotic BCL-2 family proteins in acute neural injury. *Front Cell Neurosci,* 8**,** 281.

AVANESOV, A. & MALICKI, J. 2010. Analysis of the retina in the zebrafish model. *Methods Cell Biol,* 100**,** 153-204.

BADEN, T., EULER, T. & BERENS, P. 2020. Understanding the retinal basis of vision across species. *Nat Rev Neurosci,* 21**,** 5-20.

BAHR, M. 2000. Live or let die - retinal ganglion cell death and survival during development and in the lesioned adult CNS. *Trends Neurosci,* 23**,** 483-90.

BAZZI, H., SOROKA, E., ALCORN, H. L. & ANDERSON, K. V. 2017. STRIP1, a core component of STRIPAK complexes, is essential for normal mesoderm migration in the mouse embryo. *Proc Natl Acad Sci U S A,* 114**,** E10928-E10936.

BIEHLMAIER, O., NEUHAUSS, S. C. & KOHLER, K. 2001. Onset and time course of apoptosis in the developing zebrafish retina. *Cell Tissue Res,* 306**,** 199-207.

BLIGHE, K., RANA, S. & LEWIS, M. 2019. EnhancedVolcano: Publication-ready volcano plots with enhanced colouring and labeling. *R package version,* 1.

BONFANTI, L., STRETTOI, E., CHIERZI, S., CENNI, M. C., LIU, X. H., MARTINOU, J. C., MAFFEI, L. & RABACCHI, S. A. 1996. Protection of retinal ganglion cells from natural and axotomy-induced cell death in neonatal transgenic mice overexpressing bcl-2. *J Neurosci,* 16**,** 4186-94.

CASANO, A. M., ALBERT, M. & PERI, F. 2016. Developmental Apoptosis Mediates Entry and Positioning of Microglia in the Zebrafish Brain. *Cell Rep,* 16**,** 897-906.

CHEN, R., XIE, R., MENG, Z., MA, S. & GUAN, K. L. 2019. STRIPAK integrates upstream signals to initiate the Hippo kinase cascade. *Nat Cell Biol,* 21**,** 1565-1577.

CHEN, S., LATHROP, K. L., KUWAJIMA, T. & GROSS, J. M. 2021. Retinal ganglion cell survival after severe optic nerve injury is modulated by crosstalk between JAK/STAT signaling and innate immune responses in the zebrafish retina. *bioRxiv***,** 2021.04.08.439090.

CHEN, S., ZHOU, Y., CHEN, Y. & GU, J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics,* 34**,** i884-i890.

CHEN, Y.-K., CHEN, C.-Y., HU, H.-T. & HSUEH, Y.-P. 2012. CTTNBP2, but not CTTNBP2NL, regulates dendritic spinogenesis and synaptic distribution of the striatin–PP2A complex. *Molecular biology of the cell,* 23**,** 4383-4392.

CHOI, J.-H., LAW, M.-Y., CHIEN, C.-B., LINK, B. A. & WONG, R. O. 2010. In vivo development of dendritic orientation in wild-type and mislocalized retinal ganglion cells. *Neural development,* 5**,** 29.

CHOW, R. W., ALMEIDA, A. D., RANDLETT, O., NORDEN, C. & HARRIS, W. A. 2015. Inhibitory neuron migration and IPL formation in the developing zebrafish retina. *Development,* 142**,** 2665-77.

D'AGATI, G., BELTRE, R., SESSA, A., BURGER, A., ZHOU, Y., MOSIMANN, C. & WHITE, R. M. 2017. A defect in the mitochondrial protein Mpv17 underlies the transparent casper zebrafish. *Dev Biol,* 430**,** 11-17.

D'SOUZA, S. & LANG, R. A. 2020. Retinal ganglion cell interactions shape the developing mammalian visual system. *Development,* 147.

D'ORAZI, F. D., SUZUKI, S. C. & WONG, R. O. 2014. Neuronal remodeling in retinal circuit assembly, disassembly, and reassembly. *Trends in neurosciences,* 37**,** 594-603.

DEANS, M. R., KROL, A., ABRAIRA, V. E., COPLEY, C. O., TUCKER, A. F. & GOODRICH, L. V. 2011. Control of neuronal morphology by the atypical cadherin Fat3. *Neuron,* 71**,** 820-32.

DHANASEKARAN, D. N. & REDDY, E. P. 2008. JNK signaling in apoptosis. *Oncogene,* 27**,** 6245-51.

DOWLING, J. E. 1987. *The retina: an approachable part of the brain*, Harvard University Press.

EILERS, A., WHITFIELD, J., BABIJ, C., RUBIN, L. L. & HAM, J. 1998. Role of the Jun kinase pathway in the regulation of c-Jun expression and apoptosis in sympathetic neurons. *J Neurosci,* 18**,** 1713-24.

FAWCETT, J. W., O'LEARY, D. D. & COWAN, W. M. 1984. Activity and the control of ganglion cell death in the rat retina. *Proc Natl Acad Sci U S A,* 81**,** 5589-93.

FERNANDES, K. A., HARDER, J. M., FORNAROLA, L. B., FREEMAN, R. S., CLARK, A. F., PANG, I. H., JOHN, S. W. & LIBBY, R. T. 2012. JNK2 and JNK3 are major regulators of axonal injury-induced retinal ganglion cell death. *Neurobiol Dis,* 46**,** 393-401.

FERNANDES, K. A., HARDER, J. M., KIM, J. & LIBBY, R. T. 2013. JUN regulates early transcriptional responses to axonal injury in retinal ganglion cells. *Exp Eye Res,* 112**,** 106-17.

FULLER, S. J., EDMUNDS, N. S., MCGUFFIN, L. J., HARDYMAN, M. A., CULL, J. J., ALHARBI, H. O., MEIJLES, D. N., SUGDEN, P. H. & CLERK, A. 2021. MAP4K4 expression in cardiomyocytes: multiple isoforms, multiple phosphorylations and interactions with striatins. *Biochem J,* 478**,** 2121-2143.

GODINHO, L., MUMM, J. S., WILLIAMS, P. R., SCHROETER, E. H., KOERBER, A., PARK, S. W., LEACH, S. D. & WONG, R. O. 2005. Targeting of amacrine cell neurites to appropriate synaptic laminae in the developing zebrafish retina. *Development,* 132**,** 5069-79.

GOUDREAULT, M., D'AMBROSIO, L. M., KEAN, M. J., MULLIN, M. J., LARSEN, B. G., SANCHEZ, A., CHAUDHRY, S., CHEN, G. I., SICHERI, F. & NESVIZHSKII, A. I. 2009. A PP2A phosphatase high density interaction network identifies a novel striatin-interacting phosphatase and kinase complex linked to the cerebral cavernous malformation 3 (CCM3) protein. *Molecular & Cellular Proteomics,* 8**,** 157-171.

GUAN, Q. H., PEI, D. S., ZONG, Y. Y., XU, T. L. & ZHANG, G. Y. 2006. Neuroprotection against ischemic brain injury by a small peptide inhibitor of c-Jun N-terminal kinase (JNK) via nuclear and non-nuclear pathways. *Neuroscience,* 139**,** 609-27.

HALLORAN, M. C., SATO-MAEDA, M., WARREN, J. T., SU, F., LELE, Z., KRONE, P. H., KUWADA, J. Y. & SHOJI, W. 2000. Laser-induced gene expression in specific cells of transgenic zebrafish. *Development,* 127**,** 1953-60.

HAM, J., EILERS, A., WHITFIELD, J., NEAME, S. J. & SHAH, B. 2000. c-Jun and the transcriptional control of neuronal apoptosis. *Biochem Pharmacol,* 60**,** 1015-21.

HAN, R., WANG, R., ZHAO, Q., HAN, Y., ZONG, S., MIAO, S., SONG, W. & WANG, L. 2016. Trim69 regulates zebrafish brain development by ap-1 pathway. *Sci Rep,* 6**,** 24034.

HE, Y., ZHANG, H., YU, L., GUNEL, M., BOGGON, T. J., CHEN, H. & MIN, W. 2010. Stabilization of VEGFR2 signaling by cerebral cavernous malformation 3 is critical for vascular development. *Sci Signal,* 3**,** ra26.

HOLLVILLE, E., ROMERO, S. E. & DESHMUKH, M. 2019. Apoptotic cell death regulation in neurons. *FEBS J,* 286**,** 3276-3298.

HOON, M., OKAWA, H., DELLA SANTINA, L. & WONG, R. O. 2014. Functional architecture of the retina: development and disease. *Progress in retinal and eye research,* 42**,** 44-84.

HU, M. & EASTER, S. S. 1999. Retinal neurogenesis: the formation of the initial central patch of postmitotic cells. *Dev Biol,* 207**,** 309-21.

HUBERMAN, A. D., CLANDININ, T. R. & BAIER, H. 2010. Molecular and cellular mechanisms of lamina-specific axon targeting. *Cold Spring Harb Perspect Biol,* 2**,** a001743.

HULSEN, T., DE VLIEG, J. & ALKEMA, W. 2008. BioVenn–a web application for the comparison and visualization of biological lists using area-proportional Venn diagrams. *BMC genomics,* 9**,** 1-6.

HWANG, J. & PALLAS, D. C. 2014. STRIPAK complexes: structure, biological function, and involvement in human diseases. *The international journal of biochemistry & cell biology,* 47**,** 118-148.

IMAI, F., YOSHIZAWA, A., FUJIMORI-TONOU, N., KAWAKAMI, K. & MASAI, I. 2010. The ubiquitin proteasome system is required for cell proliferation of the lens epithelium and for differentiation of lens fiber cells in zebrafish. *Development,* 137**,** 3257-68.

JEONG, B. C., BAE, S. J., NI, L., ZHANG, X., BAI, X. C. & LUO, X. 2021. Cryo-EM structure of the Hippo signaling integrator human STRIPAK. *Nat Struct Mol Biol,* 28**,** 290-299.

JUSUF, P. R., ALBADRI, S., PAOLINI, A., CURRIE, P. D., ARGENTON, F., HIGASHIJIMA, S., HARRIS, W. A. & POGGI, L. 2012. Biasing amacrine subtypes in the Atoh7 lineage through expression of Barhl2. *J Neurosci,* 32**,** 13929-44.

JUSUF, P. R. & HARRIS, W. A. 2009. Ptf1a is expressed transiently in all types of amacrine cells in the embryonic zebrafish retina. *Neural Dev,* 4**,** 34.

KAY, J. N., FINGER-BAIER, K. C., ROESER, T., STAUB, W. & BAIER, H. 2001. Retinal ganglion cell genesis requires lakritz, a Zebrafish atonal Homolog. *Neuron,* 30**,** 725-36.

KAY, J. N., ROESER, T., MUMM, J. S., GODINHO, L., MREJERU, A., WONG, R. O. & BAIER, H. 2004. Transient requirement for ganglion cells during assembly of retinal synaptic layers. *Development,* 131**,** 1331-42.

KEMP, H. A., CARMANY-RAMPEY, A. & MOENS, C. 2009. Generating chimeric zebrafish embryos by transplantation. *J Vis Exp.*

KIM, D., LANGMEAD, B. & SALZBERG, S. L. 2015. HISAT: a fast spliced aligner with low memory requirements. *Nat Methods,* 12**,** 357-60.

KIM, J. W., BERRIOS, C., KIM, M., SCHADE, A. E., ADELMANT, G., YEERNA, H., DAMATO, E., INIGUEZ, A. B., FLORENS, L., WASHBURN, M. P., STEGMAIER, K., GRAY, N. S., TAMAYO, P., GJOERUP, O., MARTO, J. A., DECAPRIO, J. & HAHN, W. C. 2020. STRIPAK directs PP2A activity toward MAP4K4 to promote oncogenic transformation of human cells. *Elife,* 9.

KITA, E. M., BERTOLESI, G. E., HEHR, C. L., JOHNSTON, J. & MCFARLANE, S. 2013. Neuropilin-1 biases dendrite polarization in the retina. *Development,* 140**,** 2933-41.

KOLDE, R. 2012. Pheatmap: pretty heatmaps. *R package version,* 1**,** 726.

KOSTER, R. W. & FRASER, S. E. 2001. Tracing transgene expression in living zebrafish embryos. *Dev Biol,* 233**,** 329-46.

LA MARCA, J. E., DIEPSTRATEN, S. T., HODGE, A. L., WANG, H., HART, A. H., RICHARDSON, H. E. & SOMERS, W. G. 2019. Strip and Cka negatively regulate JNK signalling during Drosophila spermatogenesis. *Development,* 146.

LI, D., MUSANTE, V., ZHOU, W.-L., PICCIOTTO, M. R. & NAIRN, A. C. 2018. Striatin-1 is a B subunit of protein phosphatase PP2A that regulates dendritic arborization and spine development in striatal neurons. *Journal of Biological Chemistry***,** jbc. RA117. 001519.

LI, L., HUANG, H., FANG, F., LIU, L., SUN, Y. & HU, Y. 2020. Longitudinal Morphological and Functional Assessment of RGC Neurodegeneration After Optic Nerve Crush in Mouse. *Front Cell Neurosci,* 14**,** 109.

LIAO, Y., SMYTH, G. K. & SHI, W. 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics,* 30**,** 923-30.

MACDONALD, R. & WILSON, S. W. 1997. Distribution of Pax6 protein during eye development suggests discrete roles in proliferative and differentiated visual cells. *Dev Genes Evol,* 206**,** 363-369.

MADSEN, C. D., HOOPER, S., TOZLUOGLU, M., BRUCKBAUER, A., FLETCHER, G., ERLER, J. T., BATES, P. A., THOMPSON, B. & SAHAI, E. 2015. STRIPAK components determine mode of cancer cell migration and metastasis. *Nat Cell Biol,* 17**,** 68-80.

MAES, M. E., SCHLAMP, C. L. & NICKELLS, R. W. 2017. BAX to basics: How the BCL2 gene family controls the death of retinal ganglion cells. *Prog Retin Eye Res,* 57**,** 1-25.

MASAI, I., LELE, Z., YAMAGUCHI, M., KOMORI, A., NAKATA, A., NISHIWAKI, Y., WADA, H., TANAKA, H., NOJIMA, Y. & HAMMERSCHMIDT, M. 2003. N-cadherin mediates retinal lamination, maintenance of forebrain compartments and patterning of retinal neurites. *Development,* 130**,** 2479-2494.

MCCURLEY, A. T. & CALLARD, G. V. 2010. Time Course Analysis of Gene Expression Patterns in Zebrafish Eye During Optic Nerve Regeneration. *J Exp Neurosci,* 2010**,** 17-33.

MOSHIRI, A., GONZALEZ, E., TAGAWA, K., MAEDA, H., WANG, M., FRISHMAN, L. J. & WANG, S. W. 2008. Near complete loss of retinal ganglion cells in the math5/brn3b double knockout elicits severe reductions of other cell types during retinal development. *Dev Biol,* 316**,** 214-27.

MUMM, J. S., WILLIAMS, P. R., GODINHO, L., KOERBER, A., PITTMAN, A. J., ROESER, T., CHIEN, C.-B., BAIER, H. & WONG, R. O. 2006. In vivo imaging reveals dendritic targeting of laminated afferents by zebrafish retinal ganglion cells. *Neuron,* 52**,** 609-621.

MUNEMASA, Y. & KITAOKA, Y. 2012. Molecular mechanisms of retinal ganglion cell degeneration in glaucoma and future prospects for cell body and axonal protection. *Front Cell Neurosci,* 6**,** 60.

NASSI, J. J. & CALLAWAY, E. M. 2009. Parallel processing strategies of the primate visual system. *Nat Rev Neurosci,* 10**,** 360-72.

NEAL, S. J., ZHOU, Q. & PIGNONI, F. 2020. STRIPAK-PP2A regulates Hippo-Yorkie signaling to suppress retinal fate in the Drosophila eye disc peripodial epithelium. *J Cell Sci,* 133.

NEISCH, A. L., NEUFELD, T. P. & HAYS, T. S. 2017. A STRIPAK complex mediates axonal
transport of autophagosomes and dense core vesicles through PP2A regulation. *J Cell*
Biol, 216**,** 441-461.
NISHIWAKI, Y., KOMORI, A., SAGARA, H., SUZUKI, E., MANABE, T., HOSOYA, T.,
NOJIMA, Y., WADA, H., TANAKA, H., OKAMOTO, H. & MASAI, I. 2008. Mutation
of cGMP phosphodiesterase 6alpha'-subunit gene causes progressive degeneration of
cone photoreceptors in zebrafish. *Mech Dev,* 125**,** 932-46.
NISHIWAKI, Y. & MASAI, I. 2020. beta-SNAP activity in the outer segment growth period is
critical for preventing BNip1-dependent apoptosis in zebrafish photoreceptors. *Sci Rep,*
10**,** 17379.
OKUDA, S., WATANABE, Y., MORIYA, Y., KAWANO, S., YAMAMOTO, T.,
MATSUMOTO, M., TAKAMI, T., KOBAYASHI, D., ARAKI, N., YOSHIZAWA, A.
C., TABATA, T., SUGIYAMA, N., GOTO, S. & ISHIHAMA, Y. 2017. jPOSTrepo: an
international standard data repository for proteomes. *Nucleic Acids Res,* 45**,** D1107-
d1111.
PETERSON, R. E., FADOOL, J. M., MCCLINTOCK, J. & LINSER, P. J. 2001. Muller cell
differentiation in the zebrafish neural retina: evidence of distinct early and late stages in
cell maturation. *J Comp Neurol,* 429**,** 530-40.
RANAWAT, N. & MASAI, I. 2021. Mechanisms underlying microglial colonization of
developing neural retina in zebrafish. *bioRxiv***,** 2021.05.20.444912.
RANDLETT, O., MACDONALD, R. B., YOSHIMATSU, T., ALMEIDA, A. D., SUZUKI, S.
C., WONG, R. O. & HARRIS, W. A. 2013. Cellular requirements for building a retinal
neuropil. *Cell Rep,* 3**,** 282-90.
RAPPSILBER, J., MANN, M. & ISHIHAMA, Y. 2007. Protocol for micro-purification,
enrichment, pre-fractionation and storage of peptides for proteomics using StageTips. *Nat*
Protoc, 2**,** 1896-906.
RAYMOND, P. A., BARTHEL, L. K., BERNARDOS, R. L. & PERKOWSKI, J. J. 2006.
Molecular characterization of retinal stem cells and their niches in adult zebrafish. *BMC*
Dev Biol, 6**,** 36.
RICK, J. M., HORSCHKE, I. & NEUHAUSS, S. C. 2000. Optokinetic behavior is reversed in
achiasmatic mutant zebrafish larvae. *Curr Biol,* 10**,** 595-8.
ROBINSON, M. D., MCCARTHY, D. J. & SMYTH, G. K. 2010. edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics, 26**,** 139-40.
ROBLES, E., LAURELL, E. & BAIER, H. 2014. The retinal projectome reveals brain-area-
specific visual representations generated by ganglion cell diversity. *Curr Biol,* 24**,** 2085-
2096.
SAKUMA, C., KAWAUCHI, T., HARAGUCHI, S., SHIKANAI, M., YAMAGUCHI, Y.,
GELFAND, V. I., LUO, L., MIURA, M. & CHIHARA, T. 2014. Drosophila Strip serves
as a platform for early endosome organization during axon elongation. *Nat Commun,* 5**,**
5180.
SAKUMA, C., OKUMURA, M., UMEHARA, T., MIURA, M. & CHIHARA, T. 2015. A
STRIPAK component Strip regulates neuronal morphogenesis by affecting microtubule
stability. *Sci Rep,* 5**,** 17769.
SAKUMA, C., SAITO, Y., UMEHARA, T., KAMIMURA, K., MAEDA, N., MOSCA, T. J.,
MIURA, M. & CHIHARA, T. 2016. The Strip-Hippo Pathway Regulates Synaptic

Terminal Formation by Modulating Actin Organization at the Drosophila Neuromuscular Synapses. *Cell Rep,* 16**,** 2289-97.

SATIJA, R., FARRELL, J. A., GENNERT, D., SCHIER, A. F. & REGEV, A. 2015. Spatial reconstruction of single-cell gene expression data. *Nature biotechnology,* 33**,** 495-502.

SCHEER, N., RIEDL, I., WARREN, J. T., KUWADA, J. Y. & CAMPOS-ORTEGA, J. A. 2002. A quantitative analysis of the kinetics of Gal4 activator and effector gene expression in the zebrafish. *Mech Dev,* 112**,** 9-14.

SCHNEIDER, C. A., RASBAND, W. S. & ELICEIRI, K. W. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods,* 9**,** 671-5.

SCHROETER, E. H., WONG, R. O. & GREGG, R. G. 2006. In vivo development of retinal ON-bipolar cell axonal terminals visualized in nyx:: MYFP transgenic zebrafish. *Visual neuroscience,* 23**,** 833-843.

SCHULTE, J., SEPP, K. J., JORQUERA, R. A., WU, C., SONG, Y., HONG, P. & LITTLETON, J. T. 2010. DMob4/Phocein regulates synapse formation, axonal transport, and microtubule organization. *Journal of Neuroscience,* 30**,** 5189-5203.

SEO, G., HAN, H., VARGAS, R. E., YANG, B., LI, X. & WANG, W. 2020. MAP4K Interactome Reveals STRN4 as a Key STRIPAK Complex Component in Hippo Pathway Regulation. *Cell Rep,* 32**,** 107860.

SHI, Z., JIAO, S. & ZHOU, Z. 2016. STRIPAK complexes in cell signaling and cancer. *Oncogene,* 35**,** 4549.

SULLIVAN-BROWN, J., BISHER, M. E. & BURDINE, R. D. 2011. Embedding, serial sectioning and staining of zebrafish embryos using JB-4 resin. *Nat Protoc,* 6**,** 46-55.

SYC-MAZUREK, S. B., FERNANDES, K. A. & LIBBY, R. T. 2017. JUN is important for ocular hypertension-induced retinal ganglion cell degeneration. *Cell Death Dis,* 8**,** e2945.

SZKLARCZYK, D., MORRIS, J. H., COOK, H., KUHN, M., WYDER, S., SIMONOVIC, M., SANTOS, A., DONCHEVA, N. T., ROTH, A., BORK, P., JENSEN, L. J. & VON MERING, C. 2017. The STRING database in 2017: quality-controlled protein-protein association networks, made broadly accessible. *Nucleic Acids Res,* 45**,** D362-D368.

TANG, Y., CHEN, M., ZHOU, L., MA, J., LI, Y., ZHANG, H., SHI, Z., XU, Q., ZHANG, X., GAO, Z., ZHAO, Y., CHENG, Y., JIAO, S. & ZHOU, Z. 2019. Architecture, substructures, and dynamic assembly of STRIPAK complexes in Hippo signaling. *Cell Discov,* 5**,** 3.

TEAM, R. 2016. RStudio: Integrated development environment for R. Boston, MA.

THISSE, C. & THISSE, B. 2008. High-resolution in situ hybridization to whole-mount zebrafish embryos. *Nat Protoc,* 3**,** 59-69.

UNGOS, J. M., KARLSTROM, R. O. & RAIBLE, D. W. 2003. Hedgehog signaling is directly required for the development of zebrafish dorsal root ganglia neurons. *Development,* 130**,** 5351-62.

URASAKI, A., MORVAN, G. & KAWAKAMI, K. 2006. Functional dissection of the Tol2 transposable element identified the minimal cis-sequence and a highly repetitive sequence in the subterminal region essential for transposition. *Genetics,* 174**,** 639-49.

VELDMAN, M. B., BEMBEN, M. A., THOMPSON, R. C. & GOLDMAN, D. 2007. Gene expression analysis of zebrafish retinal ganglion cells during optic nerve regeneration identifies KLF6a and KLF7a as important regulators of axon regeneration. *Dev Biol,* 312**,** 596-612.

WANG, J., STRUEBING, F. L. & GEISERT, E. E. 2021. Commonalities of optic nerve injury
and glaucoma-induced neurodegeneration: Insights from transcriptome-wide studies. *Exp*
Eye Res, 207**,** 108571.
WESTERFIELD, M. 1995. *The zebrafish book: a guide for the laboratory use of zebrafish*
(Brachydanio rerio), University of Oregon press.
WHITFIELD, J., NEAME, S. J., PAQUET, L., BERNARD, O. & HAM, J. 2001. Dominant-
negative c-Jun promotes neuronal survival by reducing BIM expression and inhibiting
mitochondrial cytochrome c release. *Neuron,* 29**,** 629-43.
XU, B., TANG, X., JIN, M., ZHANG, H., DU, L., YU, S. & HE, J. 2020. Unifying
developmental programs for embryonic and postembryonic neurogenesis in the zebrafish
retina. *Development,* 147.
XU, Q., HOLDER, N., PATIENT, R. & WILSON, S. W. 1994. Spatially regulated expression of
three receptor tyrosine kinase genes during gastrulation in the zebrafish. *Development,*
120**,** 287-99.
ZHANG, S., DONG, Y., QIANG, R., ZHANG, Y., ZHANG, X., CHEN, Y., JIANG, P., MA, X.,
WU, L., AI, J., GAO, X., WANG, P., CHEN, J. & CHAI, R. 2021. Characterization of
Strip1 Expression in Mouse Cochlear Hair Cells. *Front Genet,* 12**,** 625867.
ZHAO, X.-F., ELLINGSEN, S. & FJOSE, A. 2009. Labelling and targeted ablation of specific
bipolar cell types in the zebrafish retina. *BMC neuroscience,* 10**,** 107.
ZHOU, Y., ZHOU, B., PACHE, L., CHANG, M., KHODABAKHSHI, A. H., TANASEICHUK,
O., BENNER, C. & CHANDA, S. K. 2019. Metascape provides a biologist-oriented
resource for the analysis of systems-level datasets. *Nat Commun,* 10**,** 1523.
ZOU, S., TIAN, C., GE, S. & HU, B. 2013. Neurogenesis of retinal ganglion cells is not essential
to visual functional recovery after optic nerve injury in adult zebrafish. *PLoS One,* 8**,**
e57280.
1112
1113

 Figure legends

Figure 1. Strip1 is essential for inner retinal neural circuit development.

(A) Zebrafish retinal neural circuit showing retinal neurons and synaptic layers. GCL, ganglion cell layer; IPL, inner plexiform layer; INL, inner nuclear layer; OPL, outer plexiform layer; ONL, outer nuclear layer; RGCs, retinal ganglion cells; ACs, amacrine cells; BPs, bipolar cells; HCs, horizontal cells; PRs, photoreceptors; MGs, Müller glia.

(B) Morphology of wild-type and *rw147* embryos at 4 dpf. Dotted lines demarcate the eye. An arrowhead indicates abnormal lower jaw. An asterisk indicates heart edema.

(C) Wild-type and *rw147* mutant retinas at 4 dpf. Red and yellow arrowheads indicate the IPL and OPL, respectively.

(D) A missense mutation occurs in *strip1* gene of *rw147* mutants leading to replacement of Leu195 with arginine.

(E) Wild-type retinas labeled with anti-Strip1 antibody (upper panels) and anti-Strip1 plus Strip1 blocking peptide as a negative control (lower panels). Nuclei are stained with Hoechst. Scale, 50 µm. Right panels show higher magnification of outlined areas. Scale, 10 µm.

(F) Whole-mount labeling of 3-dpf wild-type and *strip1^{rw147}* mutant retinas with anti-acetylated α-tubulin antibody. Bottom panels show higher magnification of outlined areas. Scale, 50 µm.

(G) Projection images of single RGCs at 2 dpf expressing *ath5:Gal4VP16; UAS:MYFP* in wild type and *strip1^{rw147}* mutants. Scale, 10 µm.

(H) Projection images of single ACs at 3 dpf expressing *ptf1a:GFP* in wild type and *strip1^{rw147}* mutants. Scale, 10 µm.

(I) Projection images of single BPs at 3 dpf expressing *nyx:Gal4VP16; UAS:MYFP* in wild type and *strip1^{rw147}* mutants. Scale, 10 µm.

Figure 2. RGCs are reduced and INL cells infiltrate the GCL in *strip1* mutants.

(A) Confocal sections of wild-type and *strip1^{rw14}* mutant retinas combined with the
transgenic line *Tg[ath5:GFP; ptf1a:mCherry-CAAX]* to label RGCs and ACs.
Middle panels represent higher magnification. Lower panels show the magenta
channel. INL, inner nuclear layer; GCL, retinal ganglion cell layer. Scale, 50 μm
(upper panels) and 10 μm (middle and lower panels).
(B) Percentage of ath5+ area relative to total retinal area. Student's t test with Welch's
correction, n≥4.
(C) AC numbers per unified retinal area (8,500 μm^2). Student's t test with Welch's
correction, n≥3.
(D) Distribution of ACs (GCL or INL) per unified retinal area (8,500 μm^2). Two-way
ANOVA with the Tukey multiple comparison test, n≥3.
(E) Wild-type and *strip1^{rw14}* mutant retinas at 3 dpf labeled with anti-Pax6 which
strongly labels ACs. Arrows indicate strong Pax6+ cells that infiltrate the GCL.
Nuclei are stained with TOPRO. Scale, 50 μm.
(F) The number of strong Pax6+ cells per retina. Student's t test with Welch's
correction, n=5.
(G) Percentage of strong Pax6+ cells (GCL+ or INL+) to the total number of strong
Pax6+ cells. Two-way ANOVA with the Tukey multiple comparison test, n=5.
(H) Wild-type and *strip1^{rw14}* mutant retinas at 3 dpf labeled with anti-Prox1. Arrows
indicate Prox1+ cells that infiltrate the GCL. Nuclei are stained with TOPRO.
Scale, 50 μm.
(I) The number of Prox1+ cells per retina. Student's t test with Welch's correction,
n=5.
(J) Percentage of Prox1+ cells (GCL+ or INL+) to the total number of Prox1+. Two-
way ANOVA with the Tukey multiple comparison test, n=5.
For all graphs, data are represented as means□±□SD. ns: not significant, *p□<□0.05,
**p□<□0.01, and ****p□<□0.0001.
1171
Figure 3. Strip1 cell-autonomously promotes RGC survival.
(A) TUNEL of wild-type and *strip1^{rw147}* mutant retinas carrying the transgene
Tg[ath5:GFP] to label RGCs. Nuclei are stained with Hoechst. Scale, 50 μm.

(B) The number of TUNEL+ cells in GCL. Two-way ANOVA with the Tukey multiple
comparison test, n≥3.
(C) Percentage of ath5+ area relative to total retinal area. Two-way ANOVA with the
Tukey multiple comparison test, n≥3.
(D) Cell transplantation design to evaluate the cell autonomy of Strip1 in RGC survival.
Donor embryos from a *strip1*[rw147] mutant background are labeled with dextran
rhodamine and transplanted into host wild-type embryos. Hosts that show
transplanted retinal columns at 60 hpf were subjected to TUNEL.
(E) 60-hpf host retinas stained with TUNEL FL to visualize apoptotic cells in wild type
to wild type (upper panel) or *strip1*[rw147] mutant to wild type (lower panel). Arrows
indicate the presence of apoptotic donor cells. Scale, 10 μm.
(F) Percentage of TUNEL+ donor RGCs relative to total donor RGCs. Mann-Whitney
U test, n=4.
For all graphs, data are represented as means ± SD. *p < 0.05, **p < 0.01, and
****p < 0.0001.
1190
Figure 4. RGC death triggers abnormal positioning of ACs leading to IPL
disruption.
(A) Time-lapse imaging of wild-type and *strip1*[rw147] mutant retinas combined with the
transgenic line *Tg[ath5:GFP; ptf1a:mCherry-CAAX]* to track ACs and RGCs during
IPL formation. Asterisks denote empty areas in the GCL. Arrowheads represent
infiltration of ACs into empty spaces in the GCL. Panels on the right show higher
magnification of outlined areas. Scale, 50 μm.
(B) Cell transplantation design to evaluate the cell autonomy of Strip1 in AC-mediated
IPL formation. Donor embryos are from intercross of *strip1*[rw147] heterozygous fish
combined with *Tg[ptf1a:mCherry-CAAX]* to label ACs. Host embryos are
generated by non-transgenic intercross of *strip1*[rw147] heterozygous fish. Donor
cells are labeled with dextran Alexa-488 and transplanted into host embryos to
make chimeric host retinas with donor-derived retinal columns.
(C) Confocal images of four combinations of transplantation outcomes: wild type to
wild type, wild type to mutant, mutant to wild type and mutant to mutant.

Arrowheads represent abnormal positioning of ACs in basal side of IPL. Scale, 20
μm.
(D) Percentage of ACs (either at the apical or the basal side of the IPL) relative to total
number of ACs within a transplanted column. Two-way ANOVA with the Tukey
multiple comparison test, n≥4. Data are represented as means ± SD.
**p < 0.01, and ****p < 0.0001.
1212

Figure 5. Strip1 interacts with Strn3 to regulate RGC survival.
(A) Design of co-immunoprecipitation coupled with mass spectrometry (Co-IP/MS) to
identify zebrafish Strip1 interacting partners. Embryos carrying the transgenes
Tg[hsp:WT.Strip1-GFP], Tg[hsp:Mut.Strip1-GFP] or Tg[hsp:Gal4;UAS:GFP] were
used to pull down wild-type GFP-tagged Strip1, mutant GFP-tagged Strip1 or only
GFP, respectively. Head lysates from 2-dpf zebrafish embryos were subjected to
immunoprecipitation using GFP-Trap beads. Immunoprecipitates were digested
and analyzed by mass spectrometry.
(B) Western blotting of whole head lysates (input) and immunoprecipitates (IP) using
anti-GFP antibody. Red and black arrowheads indicate the expected band sizes
for Strip1-GFP (120 kDa) and GFP (26 kDa), respectively.
(C) Venn Diagram comparing proteins significantly enriched in WT.Strip1-GFP relative
to Control GFP (blue) and WT.Strip1-GFP relative to Mut.Strip1.GFP (magenta).
Six proteins are commonly enriched in both groups, FC>2, p < 0.05. n=3 for
WT.Strip1-GFP and Mut. Strip1-GFP and n=2 for GFP-control.
(D) Five components of the STRIPAK complex found from six proteins commonly
enriched in (C).
(E) TUNEL of 60-hpf retinas of *Tg[ath5:GFP]* transgenic embryos injected with
standard MO and MO-Strn3. RGCs and apoptotic cells are labeled with *ath5:GFP*
and TUNEL, respectively. Nuclei are stained with Hoechst (blue).
(F) The number of TUNEL+ cells in GCL. Mann-Whitney U test, n≥6.
(G) Percentage of ath5+ area relative to total retinal area. Student's t test with Welch's
correction, n≥3.

(H)   Confocal images of retinas of 76-hpf *Tg[ath5:GFP; ptf1a:mCherry-CAAX]*
transgenic embryos injected with standard MO and MO-Strn3. *ath5:GFP* and
ptf1a:mCherry-CAAX] label RGCs and ACs, respectively.

(I)   Percentage of ath5+ area relative to total retinal area. Student's t test with
Welch's correction, n=8.

Scale, 50 µm (E, H).

For all graphs, data are represented as means $\pm$ SD. **$p < 0.01$, ***$p < 0.001$ and
****$p < 0.0001$.

1244

Figure 6. Jun is a key mediator of RGC death in the absence of Strip1.

(A)   Volcano plot showing differentially expressed genes (DEGs) in *strip1^{rw147}* mutants
compared to wild-type siblings. Colored points represent genes that are
significantly upregulated (131 genes, red) or downregulated (75 genes, blue). Data
are obtained from four independent collections of 62-hpf embryo eye cups.
FDR<0.05, Log$_2$FC > |1|.

(B)   Heatmap of expression values (z-score) representing selected DEGs in *strip1^{rw147}*
mutants compared to wild-type siblings.

(C)   Whole-mount labeling of 54-hpf wild-type and *strip1^{rw147}* mutant retinas with anti-
phospho-Jun antibody and zn5 antibody, which label active Jun and RGCs,
respectively.

(D)   Percentage of phospho-Jun area relative to zn5 area at 54-58 hpf. Student's t test
with Welch's correction, n=6.

(E)   Whole-mount labeling of 49-hpf wild-type embryos injected with standard MO or
MO-strn3 with anti-phospho-Jun antibody and zn5 antibody, respectively.

(F)   Percentage of phospho-Jun area relative to zn5 area at 49 hpf. Student's t test
with Welch's correction, n=5.

(G)   TUNEL and zn5 antibody labeling of 60-hpf wild-type and *strip1^{rw147}* mutant retinas
injected with standard MO and MO-Jun. Nuclei are stained with Hoechst.

(H)   The number of TUNEL+ cells in GCL per retina. Two-way ANOVA with the Tukey
multiple comparison test, n=6.

(I) Confocal images of 76-hpf wild-type and *strip1^{rw14}* mutant retinas injected with
standard-MO and MO-Jun. Embryos carry the transgene *Tg[ath5:GFP]* to label
RGCs and are stained with bodipy TR methyl ester to visualize retinal layers.
(J) Percentage of ath5+ area relative to total retinal area, Two-way ANOVA with the
Tukey multiple comparison test, n≥3.
Scale, 50 µm (C, E, G, I).
For all graphs, data are represented as means ± SD. ns: not significant, $*p < 0.05$,
$**p < 0.01$ and $****p < 0.0001$.
1274

Figure 7. Bcl2 rescues RGC death in *strip1* mutants, but surviving RGCs do not
project their dendrites to the IPL.
(A) 60-hpf wild-type and *strip1^{rw147}* mutants combined with the transgenic line
Tg[hsp:mCherry-Bcl2]. Non-transgenic embryos (Bcl2-, top panels) are compared
to transgenic embryos (Bcl2+, bottom panels) after heat-shock treatment.
Apoptotic cells are visualized by TUNEL FL and fluorescent signals from mCherry-
Bcl2 are shown. Nuclei are stained with Hoechst.
(B) The number of TUNEL+ cells in GCL. Two-way ANOVA with the Tukey multiple
comparison test, n=3.
(C) 78-hpf wild-type and *strip1^{rw147}* mutant retinas combined with the transgenic lines,
Tg[ath5:GFP] and *Tg[hsp:mCherry-Bcl2]*. Non-transgenic embryos (Bcl2-, top
panels) are compared to transgenic embryos (Bcl2+, bottom panels) after heat-
shock treatment. RGCs are labeled with *ath5:GFP* and fluorescent signals from
mCherry-Bcl2 are shown. Anti-acetylated α-tubulin labels the IPL. Nuclei are
stained with Hoechst.
(D) Percentage of ath5+ area relative to retinal area. Two-way ANOVA with the Tukey
multiple comparison test, n≥3.
Scale: 50 µm (A, C).
For all graphs, data are represented as means ± SD. ns: not significant, $*p < 0.05$,
$**p < 0.01$, $***p < 0.001$ and $****p < 0.0001$.
1295

Figure 8. Summary of developmental and molecular events that underlie Strip1
function in inner retinal circuit formation.

(A) In wild-type retina, Strip1 suppresses Jun-mediated pro-apoptotic signals, probably

through the STIPAK complex, to maintain RGCs during development. In absence

of Strip1, Jun is activated within RGCs leading to severe degeneration of RGCs as

early as 2 dpf. Subsequently, cells in the INL abnormally infiltrate the GCL leading

to a disrupted IPL.

(B) Proposed model for Strip1 role within RGCs to regulate AC positioning and IPL

formation. In wild type, Strip1 regulates (1) RGC survival to prevent AC infiltration,

and (2) RGC dendritic patterning to promote RGC-AC interactions. In *strip1^{rw147}*

mutants, both mechanisms are perturbed, leading to AC infiltration, increased AC-

AC interactions, and IPL defects. In Bcl2-rescued *strip1^{rw147}* mutants, survived

RGCs prevent AC infiltration. However, RGC dendritic defects lead to increased

AC-AC interactions and ectopic IPL formation.

1310

Supplementary materials legends

Figure 1–figure supplement 1. Retinal lamination defects in *strip1^{rw147}* mutants are

phenocopied by either CRISPR- or MO-mediated knockdown of *strip1* and

rescued by overexpression of Strip1.

(A) CRISPR-mediated, targeted mutagenesis of zebrafish *strip1* gene. A 10-bp

deletion was introduced into exon 9, leading to a premature stop codon in *strip1*

gene.

(B) Morphology of wild-type and *strip1$^{crispr\Delta10}$* embryos at 4 dpf. Dotted lines demarcate

the eye outline. The arrowhead indicates an abnormal lower jaw. An asterisk

indicates heart edema.

(C) Plastic sections of wild-type and *strip1$^{crispr\Delta10}$* mutant retinas stained with toluidine

blue at 4 dpf.

(D) A polyacrylamide gel image showing PCR amplicons using primers that flank the

10-base deletion of *strip1$^{crispr\Delta10}$*. Wild-type and *strip1$^{crispr\Delta10}$* mutant bands are

expected to appear at 279 and 269 bps, respectively. Different band sizes were

consistently obtained from progeny produced by pair-wise cross of *strip1$^{crispr\Delta10}$*

heterozygous mutant fish.

(E) Morphology of non-injected and MO-strip1 injected wild-type embryos at 4 dpf.
Dotted lines outline the eye. An arrowhead indicates the abnormal lower jaw. An
asterisk indicates heart edema.
(F) Retinal plastic sections of non-injected and MO-strip1 injected wild-type embryos
at 4 dpf.
(G) Western blotting of 2-dpf head lysates from non-injected and MO-strip1 injected
wild-type embryos. Red and black arrows show expected band sizes for zebrafish
Strip1 (93 kDa) and the loading control b-actin (42 kDa), respectively.
(H) Plastic sections of 4 dpf wild-type and *strip1^{rw147}* mutant retinas without transgene
(top panels), and with the transgenes *Tg[hsp:WT.Strip1-GFP]* (middle panels) or
Tg[hsp:Mut.Strip1-GFP] (bottom panels) post heat shock treatment. Over
expression of GFP-tagged wild-type Strip1 rescues retinal defects in *strip1^{rw147}*
mutants, while the GFP-tagged mutant form fails to rescue.
(I) A staggered bar graph, which represents the percentage of *strip1^{rw147}* mutants with
a defective retina vs. mutants with a rescued retina in *strip1^{rw147}* mutants without
transgenes, and with *Tg[hsp:WT.Strip1-GFP]* or *Tg [hsp:Mut.Strip1-GFP].*
1344
Figure 1–figure supplement 2. *strip1* mRNA is expressed in developing retinas
and is required for IPL formation.
(A) Expression patterns of *strip1* mRNA during embryonic development. *In situ*
hybridization using a *strip1* antisense probe is shown at different developmental
stages: 4-cell, blastula, 75% epiboly, 1 dpf, 2 dpf and 3 dpf. *In situ* hybridization
using the sense probe at 1 dpf is shown as a negative control. *strip1* mRNA is
ubiquitously expressed until gastrula stage. At 1-2 dpf, expression becomes
restricted in the heart, optic tectum, and eyes. Ventral view of 2-dpf retina shows a
stronger signal in GCL (left panels). OT, optic tectum; GCL, ganglion cell layer.
Scale bar: 100 μm.
(B) Frontal cryosections of 2-dpf heads following *in situ* hybridization show that *strip1*
mRNA is predominantly expressed in the inner retina and the optic tectum.

(C) Retinal cryosections show higher expression in RGCs and ACs. Right panels show
higher magnification of outlined areas. RGCs, retinal ganglion cells; ACs, amacrine
cells; HCs, horizontal cells; PRs, photoreceptors.

(D) Live confocal snapshots of the same wild-type and *strip1^{rw147}* mutant retinas at
different developmental stages; 40, 52, 62 and 76 hpf stained with Bodipy TR. A
primitive IPL is first observed in wild-type retinas as early as 52 hpf. However, it is
not as prominent in *strip1^{rw147}* mutant retinas. By 62 hpf, *strip1^{rw147}* mutants display
severe defects in IPL formation. Arrowheads indicate rudimentary IPL. Asterisks
show IPL defects in *strip1^{rw147}* mutants. Scale bar: 50 μm.

1366

Figure 2–figure supplement 1. ACs are abnormally positioned at the basal side of
the IPL in *strip1* mutants and *ath5* morphants.

(A) Live confocal images of wild-type and *strip1^{rw147}* mutant retinas at 3 dpf combined
with the transgenic line *Tg[ptf1a:mCherry-CAAX]*, which labels ACs. Middle and
lower panels show higher magnification of outlined areas. Abnormal positioning of
ACs in the GCL is more prominent in areas of strong IPL defects than in areas of
weak lamination defects. Scale bar, 50 μm.

(B) Live confocal images of retinas of STD-MO and MO-ath5 injected wild-type
embryos combined with the transgenic line *Tg[ath5:GFP; ptf1a:mCherry-CAAX]*,
which labels RGCs and ACs. Middle and bottom panels show higher magnification
images to show abnormal localization of ACs in GCL (arrowheads) and disrupted
IPL. Scale bar, 50 μm (upper panels) and 20 μm (middle and lower panels).

(C-F) Labelling of 4-dpf wild-type and *strip1^{rw147}* mutant retinas with zpr3 (C), zpr1 (D),
anti-glutamine synthetase (GS) (E) and anti-PCNA (F) antibodies, which visualize
rod photoreceptors, double cone photoreceptors, Müller glia and proliferative cells
in the ciliary marginal zone (CMZ), respectively. Both rods and cones are in ONL
and are mostly intact in *strip1^{rw147}* mutants. Müller glia genesis and retinal stem
cells and progenitor cells in the CMZ appear to be unaffected in *strip1^{rw147}* mutants.
Scale bar, 50 μm.

1386

Figure 3–figure supplement 1. *strip1* **mutants show apoptosis in RGCs and optic tectum and elongation defects in retinal axons.**

(A)   The number of TUNEL+ in the GCL compared to the rest of the retina at 36, 48, 60, 72 and 96 hpf in s*trip1^{rw147}* mutant retinas, for experiments shown in Figure 3A. Apoptosis gradually increases to reach its peak at 72 hpf in other retinal areas, but its number at 72 hpf is significantly lower than that of the GCL at 72 hpf. In addition, there is no significant difference in apoptotic cell number of other retinal areas between 72 hpf and other time points (36, 60, 96 hpf). Two-way ANOVA with the Tukey multiple comparison test, n≥3.

(B)   Percentage of GCL area (area between the lens and IPL) relative to the total retinal area in wild-type and s*trip1^{rw147}* mutant retinas at 36, 48, 60, 72 and 96 hpf, for experiments shown in Figure 3A. There is no significant difference between wild-type siblings and s*trip1^{rw147}* mutants at any stage. Two-way ANOVA with the Tukey multiple comparison test, n≥3.

(C)   Live confocal images of wild-type and *strip1$^{crispr\Delta10}$* mutant retinas stained with acridine orange (AO) to label apoptotic cells at 60 hpf. Scale bar, 50 µm.

(D)   Percentage of AO area relative to total retina area in 60 hpf wild-type and *strip1$^{crispr\Delta10}$* mutant retinas. Mann Whitney U test, n=5.

(E)   TUNEL of 60 hpf wild-type and *strip1^{rw147}* mutant heads combined with *Tg[ath5:GFP]* to label RGCs. All nuclei are counterstained with Hoechst. Apoptosis occurs markedly in RGCs and optic tectum in *strip1^{rw147}* mutants. Scale bar, 100 µm.

(F)   The number of TUNEL+ in the optic tectum of wild-type and s*trip1^{rw147}* mutants at 60 hpf. The number of TUNEL+ cells is significantly higher in *strip1^{rw147}* mutants than in wild-type siblings. Student's t test with Welch's correction, n=3.

(G)   Dorsal view of the optic tectum of 3-dpf wild-type and s*trip1^{rw147}* mutants. RGC axons are labeled using intraretinal injections of DiO or DiI. Axons of s*trip1^{rw147}* appear to exit the optic disc normally and the optic chiasm is formed. However, the optic nerve is thinner in s*trip1^{rw147}* mutants than in wild-type siblings. In addition, most axons fail to elongate properly and do not arborize within the optic tectum

1417   (arrowheads) or very few axons elongate poorly to reach more posterior

1418   arborization fields within the optic tectum (asterisk). Scale bar: 50 μm.

1419 For all graphs, data are represented as means □±□SD. ns: not significant, *p□<□0.05,
1420 and ***p□<□0.001.

1421

Figure 4–figure supplement 1. Strip1 is not required in ACs and BPs for their
1422

neurite projections to the IPL.

(A)   3D confocal live images showing neurite projection of ACs to the IPL in

transplantation assays outlined in Figure 4B-C. Panels represent three

combinations of transplantation experiments: from wild-type donor to wild-type

host (left columns), from $strip1^{rw147}$ mutant donor to wild-type host (middle

columns), and from wild-type donor to $strip1^{rw147}$ mutant host (right columns).

Donor cells are labeled with dextran-Alexa-488 in green and ACs are labeled with

ptf1a:mCherry-CAAX in magenta. Mutant donor ACs show a normal dendritic

pattern and normal projection to the IPL in wild-type host retina, although on rare

occasion we observed that mutant ACs display two dendritic trees (arrow in the

bottom middle panel), but still contribute to a normal IPL. On the other hand, wild-

type donor ACs transplanted into mutant host retinas show abnormal projection

patterns and infiltrate the GCL (arrowheads in the right panels). Scale bar, 20 μm.

(B)   Confocal image of 3-dpf live wild-type and $strip1^{rw14}$ mutant retinas combined with

the transgenic line, *xfz43*, to label subsets of BPs. Scale bar, 10 μm.

(C)   Schematic drawings of cell transplantation design to evaluate cell autonomy of

Strip1's role in BP development. $strip1^{rw147}$ mutant donor cells carrying the

transgene *xfz43* (green) are labeled with dextran-cascade blue or dextran-Alexa-

flour 647 (blue) and transplanted into wild-type host embryos at blastula stage,

leading to chimeric wild-type retina with $strip1^{rw147}$ mutant retinal columns. Retinal

lamination of the host retina is visualized with Bodipy TR (magenta).

(D)   3D confocal live images of four wild-type host retinas with $strip1^{rw14}$ mutant donor

retinal columns outlined in (C). $strip1^{rw14}$ mutant donor BPs show normal neurite

projection patterns to the IPL in wild-type host retina. Occasionally, bipolar axon

terminals are abnormally extended laterally (arrow in the bottom panel); however,

such defects are confined to a seemingly normal IPL. Scale bar, 10 μm.

(E)   Confocal image of 3-dpf live wild-type and *strip1^{rw14}* mutant retinas combined with

the transgenic line, *xfz3*, to label subsets of BPs. Scale bar, 10 µm.

(F)   Schematic drawings of cell transplantation design to evaluate the cell autonomy of

Strip1 role in BP development. Wild-type donor cells carrying the transgene *xfz3*

(green) are labeled with dextran-rhodamine (magenta) and transplanted into

strip1^{rw14} mutant host embryos at blastula stage, leading to chimeric *strip1^{rw14}*

mutant retinas with wild-type retinal columns.

(G)   3D confocal live images of two different *strip1^{rw147}* mutant host retinas with wild-

type donor retinal columns outlined in (F). Wild-type donor BPs show misguided

neurite projections and excessive branching. Scale bar, 10 µm.

1459

Figure 5–figure supplement 1. Components of the STRIPAK complex are highly

enriched in the interactome of zebrafish Strip1 and their retinal expression,

according to published scRNA-seq data.

(A)   Schematic diagram showing the core components of the mammalian STRIPAK

complex and some of its accessory molecules. Core STRIPAK components

include serine/threonine-protein phosphatase catalytic subunit known as PP2Ac

and the scaffolding subunit PP2Aa, together with striatins (STRN, STRN3 and

STRN4), Mob3, STRIP1 or STRIP2, and a germinal center kinase GCKIII kinase

bound via cerebral cavernous malformation 3 (Ccm3). Some molecules bind in a

mutually exclusive pattern to the core components. Arrows show that sarcolemmal

membrane-associated protein (SLMAP) and the suppressor of IKKe (SIKE) are not

detected in STRIPAK complexes containing cortactin binding proteins

(CTTNBP2/NL) and vice versa. (Figure adapted from Hwang J. and Pallas D.C.,

2014).

(B)   STRING protein-protein interaction (PPI) network analysis of six zebrafish Strip1

interacting partners significantly enriched from IP-MS analysis highlighted in Figure

5C. Network edges represent known and/or predicted functional interactions in the

STRING database. Edge thickness reflects the combined STRING evidence score

for each binary relationship. Thicker edges represent increased interaction

evidence. PPI enrichment p-value <1.0e-16.

(C)　UMAP plots showing different clusters of retinal cells at 48 hpf analyzed with
Seuret R pipeline using data published by Xu B. et al, 2020. Twelve clusters are
identified and categorized to different retinal cell types or sub-types. RPCs, retinal
progenitor cells; RGCs, retinal ganglion cells; ACs, amacrine cells; BPs, bipolar
cells; HCs, horizontal cells; PRs, photoreceptors; MGs, Müller glia.
(D-I) UMAP plots showing retinal mRNA expression patterns of *strip1* (D) and its
interacting STRIPAK components, identified from Co-IP/MS; *cttnbp2* (E),
cttnbp2nlb (F), *strn* (G), *strn3* (H) and *strn4* (I), Only *strip1* and *strn3* show high
retinal expression levels at 48 hpf compared to other partners.
1489

Figure 6–figure supplement 1. Gene ontology (GO) enrichment analysis of
differentially expressed genes (DEGs) from RNA-seq of *strip1* mutants and
comparison between upregulated DEGs in *strip1* mutant transcriptomes vs.
transcriptomes of zebrafish RGCs under stress.
(A-B) List of enriched GO terms in DEGs downregulated (A) and upregulated (B) in
strip1^{rw147} mutant relative to wild-type siblings as analyzed with Metascape,
FDR<0.05 and Log$_2$FC > |1|. Downregulated genes show many GO terms related
to the nervous system, RGCs, embryo morphogenesis, synaptic development, and
transmission. Upregulated genes show terms related to stress response, cell death
and MAPK signaling.
(C)　Venn diagram showing overlap of upregulated DEGs in RNA sequencing datasets
of *strip1^{rw147}* mutant relative to wild-type (FDR<0.05 and FC >1.5) in comparison
with published upregulated DEGs in microarray data of adult zebrafish RGCs at 3
days post-optic nerve injury (ONI) (Veldman et al., 2007) and adult zebrafish eyes
at 4 days-post optic nerve crush (ONX) (McCurley and Callard, 2010).
1505

1506　**Legends of video files**
1507

Figure 4–video 1.
Development of ACs and IPL formation in wild-type sibling retina.

Time-lapse movie to track the development of ACs and RGCs in wild-type sibling

combined with the transgenic line *Tg[ath5:GFP; ptf1a:mCherry-CAAX]* to label RGCs in

green and ACs in magenta. Acquisition started ~48hpf. Movie on the right represents

outlined areas for *ptf1a:mCherry-CAAX* channel.

1514

Figure 4–video 2

Development of ACs and IPL formation in *strip1^{rw147}* mutant retina.

Time-lapse movie to track the development of ACs and RGCs in a *strip1^{rw147}* mutant

combined with the transgenic line *Tg[ath5:GFP; ptf1a:mCherry-CAAX]* to label RGCs in

green and ACs in magenta. Acquisition started ~48 hpf. The movie on the right

represents outlined areas for the *ptf1a:mCherry-CAAX* channel. Asterisks depict the

onset of severe RGC loss and arrows show areas of ectopic migration of ACs into the

GCL.

1523

1524

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Figure 8.