Good moms: dependent-young and their mothers cope better than others with longer dry season in plains zebras

Lucie Thel[1], Christophe Bonenfant[1, 4] & Simon Chamaillé-Jammes[2, 3, 4]

Author details[1]

[1] Laboratoire de Biométrie et Biologie Évolutive, Unité Mixte de Recherche 5558, Bâtiment 711, Université Lyon I, 43 Boulevard du 11 novembre 1918, F-69622 Villeurbanne Cedex, France.

[2] CEFE, Université Montpellier, CNRS, EPHE, IRD, Montpellier, France.

[3] Mammal Research Institute, Department of Zoology & Entomology, University of Pretoria, Pretoria, South Africa.

[4] LTSER France, Zone Atelier "Hwange", Hwange National Park, Bag 62, Dete, Zimbabwe-CNRS HERD (Hwange Environmental Research Development) program.

Corresponding author: Thel Lucie, lucie.mc.thel@gmail.com

[1] **Author contributions**: LT, CB and SCJ were involved in the Conceptualization, Investigation, Methodology, Writing, Funding acquisition. CB and SCJ were involved in the Supervision and Validation. LT and CB were involved in the Formal analysis. LT and SCJ were involved in the Data curation. LT was involved in the Visualization. SCJ was involved in the Data collection, Project administration.

Abstract

In large herbivores, the timing of births often matches the seasonal peak of availability of food resources, likely to improve juvenile survival and reduce reproduction costs. Some species however breed year round, even in seasonal environments. To what extent being born during the lean season reduces survival of juveniles and reproductive females remains mostly unknown. This holds particularly true in tropical ecosystems where the demographic processes underlying population dynamics are often poorly studied. We investigated survival rates in plains zebras (*Equus quagga*) in Hwange National Park (Zimbabwe), a highly seasonal savanna ecosystem. We used capture-recapture models to analyse long-term demographic data (2008-2019). We investigated the effect of seasonality as a categorical (wet versus dry season) and continuous (duration of the dry season) variable on survival. We found little variability in the early juvenile survival (< 6 m.o.), whereas late juvenile and yearling survival rates, which were higher on average, decreased with increasing length of the dry season. Female survival was high but decreased with increasing exposure to the dry season in non-reproductive females only. The probability of females becoming reproductive the next year was not affected by the length of the dry season. Together, our results highlight the importance of individual quality in reproductive performance, as reproductive females seem to buffer the effect of environmental variability on their own survival and that of their foal. Timing of birth in this population might be the result of a trade-off between minimising mother's reproductive costs and maximising juvenile survival.

Keywords: capture-mark-recapture, *Equus quagga*, Hwange National Park, reproduction, tropical ecosystem.

Introduction

In long-lived species, the recruitment of juveniles in the population is often the most variable demographic rate and the major determinant of the variation in population growth rate (Gaillard et al. 1998, Gaillardet al. 2000, Kiffner and Lee 2019). Specifically, in large mammalian herbivores, recruitment depends mostly on juvenile survival (Gaillard et al. 2003), which is affected primarily by mothers during the last months of gestation and by newborns during their first year of life (Lindström 1999, Beckerman et al. 2003, Kiffner and Lee 2019). For instance, exposure to harsh winters reduce fawn survival in red deer (*Cervus elaphus*, Loison and Langvaten 1998). In reindeer (*Rangifer tarandus*), high rain-on-snow during the gestation period increases the negative effect of a low body mass of the mother on her probability to have a calf at heel during the next summer (Pigeon et al. 2022). For capital breeders, female usually enter a reproductive phase if they reach a threshold body mass (Jönsson 1997). The energy reserves built prior to reproduction critically contribute to the female's ability to feed the fetus during pregnancy and the newborn during lactation. The level of reserves thus affects the weight of the newborn at birth (Cameron et al. 1993) and its survival to weaning (Jackson et al. 2021). In seasonal environments where food resource availability and quality vary strongly within a year, the timing of developmental stages and of the reproductive cycle is therefore critical for juvenile survival and the mother's reproductive success.

In temperate and boreal ecosystems, the timing of birth contributes to generate silver-spoon effects on life history traits (sensu Grafen 1988). Timing of birth is associated with early-life survival in the roe deer (*Capreolus capreolus*, Plard et al. 2015), and individual fitness in the collared flycatcher (*Ficedula albicollis*, Spagopoulou et al. 2020). By determining the early environmental conditions experienced by newborns during the first months of life, timing of birth contributes directly (*e.g.* through thermal stress; Grovenburg et

al. 2012) and indirectly (*e.g.* through potentially lower food availability reducing growth rate;

Feder et al. 2008) to the annual variation of juvenile survival. The season of parturition is

critical to reproductive females themselves, as they must cater to their offspring's needs in

addition to their own. Lactation costs are particularly high in mammals (Sadleir 1982,

Clutton-Brock et al. 1989), and could make reproductive females more susceptible to harsh

environmental conditions. These selection pressures acting on newborns and mothers are

thought to underlie the commonly observed strong seasonality in timing of birth across

ungulates in temperate ecosystems (Bronson 1989). Additionally, the timing of parturition

can generate reproductive costs for females by lowering the chances of successfully

reproducing the next year (Green and Rothstein 1993). In red deer, the later in the year a hind

gives birth, the less likely she is to conceive again during the following breeding season

(Clutton-Brock et al. 1983). In short, in temperate and boreal systems, the timing of birth

influences juvenile survival and reproductive success of the mother of large herbivores,

explaining why births are synchronised with the best period in terms of food resource

availability and quality (Post 2003).

Even though less predictable than temperate and boreal ecosystems, arid and semi-

arid ecosystems also show a pronounced seasonality, i.e. wet and dry seasons mainly driven

by the distribution of rains within the year (Owen-Smith and Ogutu 2013). Although some

large herbivore species give birth within a relatively short time window in such environments

(e.g. 80% of birth occur within 3 weeks in the blue wildebeest (*Connochaetes taurinus*);

Sinclair et al. 2000), others have more extended breeding periods (Owen-Smith and Ogutu

2013), or even breed year-round (e.g. the plains zebras (*Equus quagga*); Sinclair et al. 2000).

In such species, and even if a peak of birth is often observed during the wet season, some

females therefore give birth in less favorable conditions than others (Boutton et al. 1988a,

Boutton et al. 1988b, Feng et al. 2013), potentially exposing themselves and their offspring to

a higher risk of mortality (Valeix et al. 2009) and creating cohort effects (i.e. individuals

presenting similar life histories because they experience the same environmental conditions,

Lindström 1999). An interesting and yet unanswered question is therefore: for such species,

to what extent does giving birth in the dry unfavourable season reduce the probability of

survival of both the newborn and its mother?

In order to grasp a better understanding of the phenology of reproduction and the

factors that can influence it, more empirical studies exploring juvenile survival and

reproductive costs both in species reproducing year-round and in tropical large herbivores are

needed (Clutton-Brock and Sheldon 2010, Burthe et al. 2011, Festa-Bianchet et al. 2017, but

see Lee et al. 2017, MacKay et al. 2018). In this study, we investigated the effect of the

season of birth on the annual survival of young and mares in wild plains zebras. We explored

the impact of the duration of the dry season, which varies according to the timing of rainfall,

on the cohort-specific annual survival of foals (i.e. individuals of the same age), yearlings and

adult females. We took advantage of a long-term individual-based monitoring program of

plains zebras conducted in Hwange National Park (HNP) in Zimbabwe. Although their

environment is seasonal (Chamaille-Jammes et al. 2006), plains zebras breed year round in

this ecosystem (Fig. 1), and are exposed to a high inter-annual variability in the duration of

the dry season. This constitutes an adequate framework to study the impact of the timing of

reproduction on the survival of young and females in seasonal tropical environments.

We hypothesised that the dry season, and in particular long dry seasons which lead to

harsh conditions for herbivores, should decrease the survival of plains zebra foal until

weaning (0 to 6 months old) because this period corresponds to the most vulnerable stage of

life (Caughley 1966, Festa-Bianchet 1988). Similarly, harsh environmental conditions could

reduce the survival of foals when they become independent from their mother (between 6 and

12 months old). Older foals are arguably less vulnerable, but are still growing and cannot rely

on their mother anymore for food provisioning and protection (Smuts 1976, Côté and Festa-

Bianchet 2001). We expected a comparatively smaller effect of exposure to harsh conditions

on yearling (1 to 2 years old) survival than on foal survival, because older individuals should

have more resources to survive through periods of low food availability (Gaillard et al. 2000).

Regarding mares, we expected that costs of reproduction would be reflected in a lower

survival of reproductive females compared to non-reproductive females. Indeed, reproductive

females spend more energy than non-reproductive females because they have to raise a foal

(energy allocation hypothesis; Cody 1966, Williams 1966). Alternatively, one could expect

the opposite, with reproductive females coping better than non-reproductive females. The

latter could be of lower phenotypic quality, explaining that they do not even breed (Reznick

et al. 2000, Hamel et al. 2009), and overall would be more vulnerable to environmental

harshness (Weladji et al. 2008).

Materials and methods

1. Context of the study

1.1. Study site

HNP (19°00' S, 26°30' E), Zimbabwe, covers 14,651 km^2 of bushlands (*Combretum spp.,*

Terminalia sericea, Colophospermum mopane, Acacia spp.), woodlands (mostly

Colophospermum mopane and *Baikiaea plurijuga*, Rogers et al. 1993) and scrublands

interspersed with grasslands (Arraut et al. 2018). Dominant grass species in HNP are *Aristida*

sp., Bothriochloa radicans, Cynodon dactylon, Digitaria eriantha, Eragrostis sp.,

Heteropogon contortus, Panicum maximum, Pogonarthria squarrosa and Schimidtia

pappophoroides (unpub. data). Altitude varies between 900 and 1,100m a.s.l.. Average

annual rainfall is approximately 600mm, with marked inter-annual variations (Chamaille-

Jammes et al. 2006). On average, 98% of the rainfall events occur during the wet season,

generally from October to April (Chamaille-Jammes et al. 2006, Chamaille-Jammes et al.

2007). The start of the wet season is characterised by a high inter-annual variability, leading

to a variable duration of the dry season. The study took place in the north-east of the park,

where artificial waterholes retain water year round and where areas > 8km from a waterhole

are rare, even in the dry season (Chamaille-Jammes et al. 2007). The densities of lion

(*Panthera leo*) and spotted hyena (*Crocuta crocuta*), the two main predators of plains zebra,

are relatively high (Loveridge et al. 2007, Drouet-Hoguet 2007). There is no hunting in the

park.

1.2. Study species

Plains zebras live in harems composed of a stallion, several mares and their offspring under

two years old (Klingel 1969). Females can give birth to a singleton at any time of the year in

most of their geographic range, including Zimbabwe (Dasmann and Mossman 1962, Barnier

et al. 2012), even if a birth peak occurs from January to March in this area (Fig. 1). Gestation

length ranges between 340 and 390 days and foals are weaned between 6 and 12 months of

age (Smuts 1976, Owen-Smith and Mason 2005, Pluháček et al. 2007). Plains zebra foals are

considered as "followers" on the hider-follower gradient (Lent 1974), as they stand and

follow their mother a couple of hours after birth (Sinclair2000). Plains zebras are grazers

(*sensu* Hofman 1989), feeding virtually only on grasses, hence the availability of their food

resource is directly linked to rainfall amount and distribution within the year, as well as with

the general availability of habitats having a grass layer (Du Plessis 2001). Similarly to the

other large herbivores occupying African savannas, the plains zebra is considered to be a

capital breeder (Jönsson 1997, Stephens et al. 2009), although there is currently little

empirical assessment of its energy allocation strategy (but see Ogutu et al. 2014). In HNP, the

population of plains zebras is mostly resident thanks to the artificial waterholes providing

water year round inside the park, but some seasonal shifts in home-ranges have been

observed for some harems (*unpublished GPS data*).

1.3. Seasonality and plains zebra life histories

Following the protocol presented in Grange et al. (2015), we recorded the presence of

individually identified plains zebras between 2008 and 2019 using visual identification of

their unique stripe pattern. Observations were conducted twice a year, around March and

around August, during field sessions (hereafter called "sessions", *n* = 24 sessions, *mean*

session duration ± SD = 45 ± 25 days, *range* = 13 - 88 days). Individuals of more than 24

months were identified as adults while the age of the young individuals was determined more

accurately based on their relative size compared to adults, using the criteria of Smuts (1975)

and Penzhorn (1982), and photographs of individuals of known age from HNP. We also

attributed an accuracy to this age estimation, based on our experience from the field. The age

of very young foals was considered accurate to a few days, whereas young seen at an older

stage for the first time could be attributed an age ± 2 months, for instance.

The main goal of this study was to explore the effect of seasonality as a categorical

(i.e. wet season versus dry season) or continuous (i.e. duration of the dry season) variable on

the survival of young and mares, accounting for the reproductive state of the mare and the

age-class of the juvenile. We also assessed the probability of transition between reproductive

and non-reproductive states in mares from one session to the next. Explanatory variables are

described in the following sections and summarised in Table 1.

2. Explanatory variables

For each year, we identified the transition date between wet and dry season using 500m

spatial resolution, bi-monthly Normalised Difference Vegetation Index (NDVI) records from

the NASA website (MOD13A1 product, computed from atmospherically corrected bi-

directional surface reflectances masked for water, clouds, heavy aerosols, and cloud shadows;

temporal resolution: 15 days; https://modis.gsfc.nasa.gov) and daily rainfall records from the

Climate Hazards Center website (Rainfall Estimates from Rain Gauge and Satellite

Observations, https://www.chc.ucsb.edu). During the study period and according to our

estimations, the wet season in HNP started between the 1st of November and the 19th of

December, while the dry season started between the 9th of May and the 29th of July (Fig. 1,

Online Resource 1). We attributed a season to each session according to the date of the

session (variable *season*, Table 1). We also estimated the proportion of days of dry season

between the first day of the session *s* and the first day of the following session *s+1* (variable

pds, namely "proportion of dry season", Table 1). We defined three age-classes among the

young: "younger foals" of less than six months old, "older foals" aged between six and 12

months old, and "yearlings" between one and two years old (Gaillard et al. 2000, Table 1).

3. Statistical analysis

3.1. Capture-Mark-Recapture framework

We used a Capture-Mark-Recapture (CMR, Lebreton et al. 1992) framework to account for

the variability of the probability of recapture (i.e. re-observation) of the individuals of this

population (Grange et al. 2015). We considered each session as a discrete event, summarised

by its starting date. As the time interval between two successive sessions varied, we used the

time.intervals argument of the *process.data* function to correct survival estimates for the

duration of exposure to mortality risk (*RMark* package, Laake 2013). For this, we calculated

the proportion of time elapsed between two successive sessions in relation to a year of 365

days: $\Delta t_{session2-session1} = (Starting\ date_{session2} - Starting\ date_{session1}) / 365$. We performed all CMR

analyses using the program MARK (www.phidot.org/software/mark) and the R statistical

software (version 4.3.1, www.r-project.org), thanks to the package *RMark* (Laake 2013). The

goodness-of-fit tests (GOF tests: see below) were conducted with the R package *R2ucare*

(Gimenez et al. 2017).

3.2. Survival of the young

Similarly to previous demographic analyses performed on these data (Grange et al. 2015,

Vitet et al. 2020, Vitet et al. 2021), we built capture histories for each individual as follows: 0

corresponding to "no sighting of the individual during the session", and 1 corresponding to

"at least one sighting of the individual during the session". We ran the analyses on

individuals whose date of birth was known with an accuracy ranging from 0 to ± 90 days,

based either on direct observation in the field or on hormonal assay of the feces of the female

(n = 310). Indeed, the dataset contained individuals observed at least once in the field (n =

290) and individuals never observed in the field but whose mother was detected to be

pregnant (n = 20). For the latter, the date of parturition was estimated based on opportunistic

faecal sampling from the mare and subsequent 20-oxopregnanes and oestrogens enzyme

immunoassay (Online Resource 2, Ncube et al. 2011). We recorded these individuals as being

identified at birth only and never seen again, assuming no foetal loss or abortion as usually

reported in large mammals. We assigned all the individuals born close to the same session to

the same cohort and assumed that they experienced the same environmental conditions during

the last months of gestation and beginning of lactation (i.e. *season* and *proportion of dry*

season variables). Based on the rich literature dealing with northern ungulates which breed

synchronously during a restricted period, a cohort is generally defined as the ensemble of

individuals born during the same year (Gaillard et al. 1993, Gaillard et al. 2003). However,

working on annual cohorts is less meaningful in the case of asynchronous breeders, for which

births are more spread out and can happen during different seasons. In our situation, it made

more sense to define a cohort at the seasonal scale, such as Lee et al. (2017) did to study

juvenile survival of giraffes *Girafa camelopardalis*. We kept sub-adults and adults (> 2 years

old) in the model to get better estimations of juvenile and yearling survival. In other words,

individuals re-observed only at an age of 2 years or more but missed during the previous

sessions still contributed to the estimation of survival of the young, even if adult survival *per*

se was the focus of another analysis. Indeed, because detection of individuals is imperfect, a

zebra can be unseen for several years in a raw, even if still alive. If a foal is only seen again at

the adult stage but adult stage is not included in the juvenile survival analyses, these

individuals would be considered as dead and juvenile survival would be overall biased low.

In a preliminary step, we performed goodness-of-fit tests to check if the dataset of

young individuals did fit the CMR model assumptions (Lebreton et al. 1992). The GOF tests

of the fully time-dependent model (Gimenez et al. 2018) highlighted possible overdispersion

(Test 2.CL: $\chi^2 = 40.79$, $df = 16$, $P < 0.01$; Test 3.Sm: $\chi^2 = 11.05$, $df = 19$, $P = 0.92$), owing to

trap-dependence ($\chi^2 = 139.35$, $df = 17$, $P < 0.01$) and transience issues ($\chi^2 = 70.47$, $df = 22$, P

< 0.01). After examination of Test 2.CL, we noticed that overdispersion was mainly caused

by three sessions over the 24 available in the dataset, so we considered this lack of fit as

marginal and ignored overdispersion in the analyses.

We accounted for trap-dependence by adding a trap-dependence effect in the

recapture model (*td* variable, Table 1). As plains zebras' distribution within the park could

depend on seasonal dynamics, we tested the effect of the *season* or *proportion of dry season*

in the recapture model. Transience was likely due to the age structure as young individuals

often have lower survival in large herbivores (Gaillard et al. 2000) and are thus less

frequently recaptured than adults, leading to a lower recapture probability. Thus, we fitted the

survival model with an *age-class* effect (Table 1). We also explored the effect of several

poolings of those age-classes (e.g. older foals and yearlings pooled together as one single

age-class versus younger foals). We tested the effect of the *season* or *proportion of dry*

season in the young survival models (Online Resource 3). For comparison purposes, we also

conducted an analysis of the survival of the juveniles using a Generalised Linear Model

(GLM) approach instead of a CMR approach (Online Resource 4).

3.3. Mare survival

To study the probability of survival and transition between reproductive states in adult

females (n = 322), we used a multi-state capture-recapture model (Lebreton et al. 2009). We

built capture histories for each individual as follows: 0 corresponding to "no sighting of the

mare during the session", 1 corresponding to "all sightings of the mare without any

dependent juvenile during the session" and 2 corresponding to "at least one sighting of the

mare with a dependent juvenile during the session". We considered a dependent juvenile

(corresponding to our "younger foal" age-class) as an individual that was still suckling and

entirely relying on its mother's milk to feed, so under 6 months of age.

As there were too few observations to conduct the GOF tests of the multi-states

model, we conducted the GOF tests on the single-state model instead (i.e. without taking into

account reproductive states). The tests revealed overdispersion (Test 2.CL: χ^2 = 49.02, df =

19, P < 0.01; Test 3.Sm: χ^2 = 72.10, df = 20, P < 0.01), trap-dependence (χ^2 = 112.13, df = 21,

P < 0.01) and transience (χ^2 = 62.23, df = 21, P < 0.01). After examination of Test 2.CL and

3.Sm, we noticed that overdispersion was mainly caused by five and three sessions over the

24 available in the dataset, respectively. Again, we considered it as marginal and ignored

overdispersion in the analyses. We accounted for trap-dependence by adding a trap-

dependence effect (*td* variable, Table 1) in the recapture model. To take into account

transience, we added a categorical variable *sight* in the survival model to differentiate

between mares captured for the first time during the survey and mares already captured at

least once during the survey, following the method described by Pradel et al. (1997; *sight*

variable, Table 1). We evaluated the effect of the *reproductive state* (Table 1), and *season* or

proportion of dry season on recapture, survival and transition probabilities. Age being not

known for most adult females, we were unable to include this variable in the model.

3.4. Model selection

For both juveniles and mares, we conducted model selection by comparing models ranging

from the most complex full time-dependent model (*time* variable, i.e. chronological number

identifying the session, Table 1) to the reference model (i.e. depending exclusively on the

covariables related to GOF corrections, Online Resource 3). Because of the large number of

possible combinations between recapture and survival (and transition for the mares dataset)

models, we conducted a sequential model selection. We started by modelling recapture, then

survival and finally transition probabilities (Lebreton et al. 1992). We conducted a first model

selection to identify the best recapture model, while survival model was set to depend

exclusively on the covariables related to GOF corrections (i.e. *age-class* for young

individuals and *sight* for mares) and transition probability was set constant for mares. We

then modeled survival probability using the best recapture model and constant transition

probability for mares. We eventually selected the best transition probability model with the

best recapture and survival models for mares.

We used the Akaike Information Criterion adjusted for small sample sizes (AICc) and

we selected the model with the lowest number of parameters for models having a ΔAICc < 2

(i.e. models receiving substantial empirical support, Burnham and Anderson 2002, Arnold

2010) when compared to the best model (parsimony principle, Burnham and Anderson 2002).

In order to estimate the overall explanatory power of the variables *time in dry season* versus

season, we calculated the cumulated weight of all the models including the effect of the

proportion of dry season or the *season* on the probability of survival. The weight of the

model *i* was calculated as follows:

$$weight_i = e^{-\Delta AIC_i/2} / \Sigma(e^{-\Delta AIC_i/2}),$$

where ΔAIC_i is the difference between the AIC of the best model and the model *i*. We also

report the number of models within a ΔAIC < 7 (i.e. models receiving substantial or some

form of empirical support, Burnham and Anderson 2002, Arnold 2010) from the best model

which included the effect of the *proportion of dry season* or the *season* on the probability of

survival. Following Arnold (2010), we reported the 85% confidence intervals of estimated

parameters, in accordance with our AICc model selection procedure. We used the scaled

value of *proportion of dry season* in the models to ease model convergence, but for

convenience, we back-transformed the coefficients to their initial scale in the Results section.

Results

1. Survival of the young

Results on recapture probabilities of juveniles are presented in Online Resource 5. In

summary, recapture probability was best described by an effect of *trap dependence* and

proportion of dry season. Recapture probability ranged between 0.293 ± 0.039 SE, 85% CI

[0.240; 0.353] and 0.483 ± 0.030 SE, 85% CI [0.440; 0.526] over the study period. According

to the most parsimonious model, survival varied between age-classes (see Online Resource 3

for the detailed model selection). The early juvenile survival was lower ($\varphi = 0.458 \pm 0.044$

SE, 85% CI [0.395; 0.522]) than the late juvenile and yearling survival, which decreased with

the *proportion of dry season* ($\beta = -0.026 \pm 0.016$ SE, 85% CI [-0.049; -0.002]). The predicted

probability of survival of older foals and yearlings (pooled together) varied from $0.850 \pm$

0.095 SE, 85% CI [0.661; 0.943] when the *proportion of dry season* was the shortest (i.e. 9 %

of the time) to 0.480 ± 0.120 SE, 85% CI [0.316; 0.648] when the *proportion of dry season*

was the longest (i.e. 80 % of the time, Fig. 2). The *season* variable was not retained in the

best models of either the CMR or the GLM approaches (Online Resource 4). Even though

non-significant, a quick exploration of late juvenile survival and yearling survival according

to the season and time showed that it was likely lower and more variable during the dry

(0.536 ± 0.221 SD) than during the wet season (0.707 ± 0.172 SD). Overall, the cumulated

weight of the models including the effect of the *proportion of dry season* on the probability of

survival was 0.738 ($n = 15$ models in a $\Delta AIC < 7$), whereas the cumulated weight of the

models including the effect of the *season* was only 0.214 ($n = 9$ models in a ΔAIC < 7).

2. Mare survival

The recapture probability was higher for mares than for young individuals, within the range

0.449 ± 0.033 SE, 85% CI [0.402; 0.496] - 0.849 ± 0.083 SE, 85% CI [0.690; 0.934] (Online

Resource 5). The best model for survival probability included an additive effect of *sight*,

proportion of dry season and *reproductive state*. The best model for transition probability

included an additive effect of *time* and *reproductive state* (see Online Resource 3 for the

detailed model selection). As the *sight* variable had a significant effect on the probability of

survival, we present results for mares in their second and following observations only.

Results relying on mares observed multiple times are a better representation of the average

behaviour of the population than those from mares observed for the first time or only once

because the latter could be transient young adult females. The *proportion of dry season* had a

negative effect on survival ($\beta = -0.035 \pm 0.011$ SE, 85% CI [-0.050; -0.019], Fig. 3a). The

probability of survival of reproductive females varied slightly from 0.995 ± 0.006 SE, 85%

CI [0.970; 0.999] when the *proportion of dry season* was the shortest to 0.945 ± 0.059 SE,

85% CI [0.768; 0.989] when the *proportion of dry season* was the longest. Although the

interaction between the *proportion of dry season* and the reproductive state of the females

was not retained in the best model, we noted that the probability of survival of non-

reproductive females appeared to vary more largely, from 0.966 ± 0.019 SE, 85% CI [0.924;

0.985] to 0.703 ± 0.057 SE, 85% CI [0.615; 0.779] as the *proportion of dry season* increased.

Overall, the cumulated weight of the models including the effect of the *proportion of dry*

season on the probability of survival was 0.964 ($n = 3$ models in a ΔAIC < 7), whereas the

cumulated weight of the models including the effect of the *season* was only 0.024 ($n = 0$

models in a ΔAIC < 7).

The probability of transition between reproductive states was not dependent on either

the *season* or the *proportion of dry season* as none of these variables were selected in the best

model, but was highly variable according to *time*. The probability for a mare to switch from

the reproductive to the non-reproductive state between two consecutive sessions was higher

(from 0.582 ± 0.094 SE, 85% CI [0.443; 0.709] to 0.961 ± 0.015 SE, 85% CI [0.933; 0.978])

than the probability to transition from the non-reproductive to the reproductive state (from

0.049 ± 0.018 SE, 85% CI [0.028; 0.083] to 0.475 ± 0.093 SE, 85% CI [0.346; 0.608], Fig.

3b). Based on the model including the season and reproductive state as explanatory variables

for the probability of transition, the average probability to be reproductive one year later was

not different between reproductive females (0.423) and non-reproductive females (0.420).

Discussion

Our study highlighted an influence of duration of dry season on the survival of some juvenile

age-classes. We found the year-round breeding strategy of female plains zebras to have little

consequences on early survival of foals, which could be related to the long gestation length

and the capital breeding strategy of energy allocation to reproduction. We documented,

however, a within-year and between years variability in young survival rates with potential

consequences for population dynamics.

Although we did not find major evidence for an effect of the season as a categorical

variable, possibly because our field sessions sometimes happened at different stages of a

given season due to variability in their timing, we identified a marked effect of the duration

of the dry season, a more accurate variable to estimate the possible influence of harsh

conditions on survival. In contrast with early juvenile survival, late juvenile survival and

yearling survival were lower during long dry seasons. Georgiadis et al. (2003) suggest in their

theoretical approach that plains zebra foals under one year of age are more sensitive to

fluctuations in rainfall than adults. Sensitivity to environmental conditions of young

individuals is a commonly observed pattern, even in tropical herbivores: for instance,

juveniles and yearlings' survival is highly correlated with the preceding annual rainfall in the

greater kudu (*Tragelaphus strepsiceros*, Owen-Smith 1990) and calf survival decreases

during the dry season in the wildebeest (*Connochaetes taurinus*, Mduma et al. 1999). Young

plains zebras only reach their adult size around three years of age and adult body mass around

five years of age (Smuts 1975). During this critical period, older foals and yearlings showed a

decreasing survival with increasing time spent in dry season. This decrease in survival could

result from energy allocation to body growth that may induce a higher vulnerability to poor

environmental conditions or predation (Grange et al. 2004).

Reproductive females usually time their reproduction to match the most costly part of

the reproductive cycle, corresponding to the late gestation and early lactation in large

herbivores (Clutton-Brock et al. 1989), with the period of best food quality or availability

(Bronson 1989). In mammals, this matching between food resources and costly reproductive

events should minimise subsequent fitness costs such as a lower survival, litter size or

probability of reproducing again at the following opportunity. Accordingly, most of the plains

zebra mares in HNP gave birth during the wet season (Fig. 1). Our results suggest that when

environmental conditions are harsher (longer dry seasons), mothers appear to buffer the costs

for the dependent foal (i.e. younger foal). This supports the idea that plains zebras might be

capital breeders, with females engaging in reproduction conditional to a sufficient acquisition

of energy prior the reproductive period. This would increase their ability to cope with their

offspring's requirements in case of an unexpectedly long dry season. However, dry season

births do not mean death sentence for either the foal or the mare. Plains zebras are able to

survive on rather low quality grasses, even if these may not allow them to not loose condition

in the dry season (Duncan et al. 2024). Therefore, reproductive timing leading to dry season

births may not have been selected against so strongly. Additionally, the presence of

permanent waterholes in the study area greatly removes the effect that drought usually have

on herbivores via reduced water availability.

Given our results and somewhat counterintuitively, the optimal timing of birth in

order to minimise mortality risks of young plains zebras should be the beginning of the dry

season. Juveniles could thus benefit from their mother's protection during the harsher season

and enter the age of independence during the most favourable one. Similarly, Lee et al.

(2017) showed that giraffes born during the dry season experienced a higher survival

probability in Tanzania. Even though most of the plains zebras' births in our population

happened during the wet season, ensuring sufficient food resources for the mother at a critical

stage of their reproductive cycle, the birth peak also extended towards the beginning of the

dry season (Fig. 1), which might be more favourable to the offspring. Such a phenomenon

could illustrate a potential trade-off between minimising mother's reproductive costs and

maximising juveniles' survival, as it has been described in primates (Dezeure et al. 2021).

However, the strength of this trade-off might not be so large. The length of the dry

season only had a small impact on the survival of reproductive females, whose survival

always remained very high (~ 0.95 or above). Our results also suggest that reproductive

females are not more sensitive, and possibly are less sensitive, than non-reproductive females

to environmental conditions. Females engaging in reproduction thus appear to not only be

able to buffer the effect of the environment on their young foal, but also to bear the costs of

reproduction, even under harsh conditions, with minimal impact. As shown by the similar

probability of transition to the reproductive state both from the non-reproductive and the

reproductive state the previous year (0.420 and 0.423 respectively), reproductive females are

as able as non-reproductive females to engage in future reproduction. All these results point

toward the fact that reproductive females might be those of higher reproductive potential, or

experience. If these effects have been well described and studied in temperate or boreal

ungulates (e.g. Hamel et al. 2009), our results generalise these observations to tropical

species. As both reproductive and non-reproductive females can be found in the same

harems, with thus similar landscape use, we can rule out that differences in habitat

availability or habitat selection strategies explain the observed contrasts between females of

different reproductive states. Given the importance of age and dominance status (which are

often related in equids) on reproduction previously demonstrated in horses and zebra species

(Lloyd and Rasa 1989, Pluháček et al. 2007, Rubenstein and Nuñez 2009), we suspect that

they are key traits underlying the observed difference. Unfortunately, age, which was

sometimes available (dominance status was not) could not be included in our analysis without

reducing its power.

Overall, we documented marked variations in demographic rates (juvenile survival

and females' reproductive rates) of plains zebras in HNP, both between seasons and across

the years, partly accounted for by the duration of the dry season. Such variations can create

cohort effects, which have been documented in many long-lived species in temperate areas

(see Sæther 1997, Gaillard et al. 2003 for reviews in large herbivores, Kiffner and Lee 2019)

but so far have not been extensively studied in tropical ecosystems (e.g. in the greater kudu,

Owen-Smith 1990). The seasonal variation of juvenile survival has the potential to generate

short-term and long-term cohort effects on demographic rates (Gaillard et al. 2003).

Compared to strictly seasonal breeders, plains zebras have more variable cohort-specific

recruitment rates. A potential large juvenile survival variability might limit the long-term

stochastic population growth rate because of its sensitivity to the temporal variance of

demographic rates (Tuljapurkar 1989).

Acknowledgments

This long-term demographic survey was initiated by Patrick Duncan, who also took part in many of the field sessions. The authors are therefore indebted to him, and also acknowledge all the other people involved in data collection over the years, in particular S. Grange, F. Barnier, H. Ncube, C. Vitet and A. Duncan. J. Dezeure assisted with NDVI and rainfall data retrieval.

Declarations

Funding: This work was supported by a PhD grant from the "Ministère Français de l'Enseignement Supérieur, de la Recherche et de l'Innovation" through the "Ecole Doctorale E2M2" of the "Université Claude Bernard Lyon 1" attributed to LT. This long-term study benefited from some financial support from the OSU-OREME (CNRS UAR 3282).

Conflict of Interest: The authors declare that they have no conflict of interest.

Ethical approval: Ethics approval was not required for this study according to local legislation [Parks and Wildlife Act].

Consent to participate: Not applicable.

Consent for publication: Not applicable.

Availability of data and material: Data available from the corresponding author on reasonable request.

Code availability: Not applicable.

Authors' contributions: LT, CB and SCJ were involved in the Conceptualization, Investigation, Methodology, Writing, Funding acquisition. CB and SCJ were involved in the Supervision and Validation. LT and CB were involved in the Formal analysis. LT and SCJ were involved in the Data curation. LT was involved in the Visualization. SCJ was involved in the Data collection, Project administration.

References

Apollonio M, Merli E, Chirichella R, Pokorny B, Alagić A, Flajšman K, Stephens PA (2020) Capital-income breeding in male ungulates: Causes and consequences of strategy differences among species. Front ecol evol 8:521767. https://doi.org/10.3389/fevo.2020.521767

Arnold TW (2010) Uninformative parameters and model selection using Akaike's Information Criterion. J Wildl Manage 74:1175–1178. https://doi.org/10.1111/j.1937-2817.2010.tb01236.x

Arraut EM, Loveridge AJ, Valls-Fox H, Macdonald DW, Chamaillé-Jammes S (2018) The 2013-2014 vegetation structure map of Hwange National Park, Zimbabwe, produced using free satellite images and software. Koedoe African Prot Area Conserv Sci 60:1–10. https://doi.org/10.4102/koedoe.v60i1.1497

Barnier F, Grange S, Ganswindt A, Ncube H, Duncan P (2012) Inter-birth interval in zebras is longer following the birth of male foals than after female foals. Acta Oecologica 42, 11–1. https://doi.org/10.1016/j.actao.2011.11.007

Beckerman AP, Benton TG, Lapsley CT, Koesters N (2003) Talkin' 'bout my generation: environmental variability and cohort effects. Am Nat 162, 754–767. https://doi.org/10.1086/381056

Boutton TW, Tieszen LL, Imbamba SK (1988a) Biomass dynamics of grassland vegetation in Kenya. Afr J Ecol 26, 89–101. https://doi.org/10.1111/j.1365-2028.1988.tb00960.x

Boutton TW, Tieszen LL, Imbamba SK (1988b) Seasonal changes in the nutrient content of East African grassland vegetation. Afr J Ecol 26, 103–115. https://doi.org/10.1111/j.1365-2028.1988.tb00961.x

Bronson FH (1989) Mammalian reproductive biology. University of Chicago Press, Chicago,

USA

Burnham K, Anderson D (2002) Model Selection and Multimodel Inference: A Practical

Information-Theoretic Approach. Springer, New York, USA

Burthe S, Butler A, Searle KR, Hall SJ, Thackeray SJ, Wanless S (2011) Demographic

consequences of increased winter births in a large aseasonally breeding mammal (*Bos*

taurus) in response to climate change. J Anim Ecol 80, 1134–1144. https://doi.org/10.11

11/j.1365-2656.2011.01865.x

Cameron RD, Smith WT, Fancy SG, Gerhart KL, White RG (1993) Calving success of

female caribou in relation to body weight. Can J Zool 71, 480–486. https://doi.org/10.11

39/z93-069

Caughley G (1966) Mortality patterns in mammals. Ecol 47, 906–918. https://doi.org/10.2307

/1935638

Chamaillé-Jammes S, Fritz H, Murindagomo F (2007) Detecting climate changes of concern

in highly variable environments: Quantile regressions reveal that droughts worsen in

Hwange National Park, Zimbabwe. J Arid Environ 71, 321–326. https://doi.org/10.1016/

j.jaridenv.2007.05.005

Chamaillé-Jammes S, Fritz H, Murindagomo F (2006) Spatial patterns of the NDVI-rainfall

relationship at the seasonal and interannual time scales in an African savanna. Int J

Remote Sens 27, 5185–5200. https://doi.org/10.1080/01431160600702392

Clutton-Brock T, Sheldon BC (2010) Individuals and populations: the role of long-term,

individual-based studies of animals in ecology and evolutionary biology. Trends Ecol &

Evol 25, 562–573. https://doi.org/10.1016/j.tree.2010.08.002

Clutton-Brock TH, Albon SD, Guinness FE (1989) Fitness costs of gestation and lactation in

wild mammals. Nature 337, 260–262. https://doi.org/10.1038/337260a0

Clutton-Brock TH, Guinness FE, Albon SD (1983) The costs of reproduction to red deer

hinds. J Anim Ecol 52, 367–383. https://doi.org/10.2307/4560

Cody ML (1966) A general theory of clutch size. Evolution 20, 174–184. https://doi.org/10.2

307/2406571

Côté SD, Festa-Bianchet M (2001) Birthdate, mass and survival in mountain goat kids:

effects of maternal characteristics and forage quality. Oecologia 127, 230–238. https://d

oi.org/10.1007/s004420000584

Dasmann RF, Mossman AS (1962) Reproduction in some ungulates in Southern Rhodesia. J

Mammal 43, 533–537. https://doi.org/10.2307/1376916

Dezeure J, Baniel A, Carter A, Cowlishaw G, Godelle B, Huchard E (2021) Birth timing

generates reproductive trade-offs in a non-seasonal breeding primate. Proc R Soc B 288,

20210286. https://doi.org/10.1098/rspb.2021.0286

Drouet-Hoguet N (2007) Influence des activités anthropogéniques sur la régime alimentaire

et la réponse numérique de la hyène tachetée en savane arborée dystrophique dominée

par l'éléphant. PhD Thesis Lyon 1 University, France.

Duncan P, Valeix M, Loveridge AJ, Grange S, Barnier F, Ncube H, Périquet S, Vitet C,

Davidson Z, Mbizah M, Madhlamoto D, Fritz H, Chamaillé-Jammes S (2024)

Population limitation of a non-ruminant in a nutrient-poor ecosystem—predation rather

than food. African Journal of Ecology, 62(3), e13294. https://doi.org/10.1111/aje.13294

Du Plessis WP (2001) Effective rainfall defined using measurements of grass growth in the

Etosha National Park, Namibia. J Arid Environ 48, 397–417. https://doi.org/10.1006/jare.

2000.0752

Feder C, Martin JGA, Festa-Bianchet M, Bérubé C, Jorgenson J (2008) Never too late?

Consequences of late birthdate for mass and survival of bighorn lambs. Oecologia 156,

773–781. https://doi.org/10.1007/s00442-008-1035-9

Feng X, Porporato A, Rodriguez-Iturbe I (2013) Changes in rainfall seasonality in the tropics.

Nat Clim Chang 3, 811–815. https://doi.org/10.1038/nclimate1907

Festa-Bianchet M (1988) Birthdate and survival in bighorn lambs (*Ovis canadensis*). J Zool

214, 653–661. https://doi.org/10.1111/j.1469-7998.1988.tb03764.x

Festa-Bianchet M, Douhard M, Gaillard J-M, Pelletier F (2017) Successes and challenges of

long-term field studies of marked ungulates. J Mammal 98, 612–620. https://doi.org/10.

1093/jmammal/gyw227

Gaillard J-M, Delorme D, Jullien J (1993) Effects of cohort, sex, and birth date on body

development of roe deer (*Capreolus capreolus*) fawns. Oecologia 94, 57–61.

https://doi.org/10.1007/bf00317301

Gaillard J-M, Festa-Bianchet M, Yoccoz NG, Loison A Toigo C (2000) Temporal variation

in fitness components and population dynamics of large herbivores. Annu Rev Ecol Syst

31, 367–393. https://doi.org/10.1146/annurev.ecolsys.31.1.367

Gaillard J-M, Festa-Bianchet M, Yoccoz NG (1998) Population dynamics of large

herbivores: variable recruitment with constant adult survival. Trends Ecol & Evol 13,

58–63. https://doi.org/10.1016/s0169-5347(97)01237-8

Gaillard J-M, Loison A, ToÏgo C, et al (2003) Cohort effects and deer population dynamics.

Ecoscience 10, 412–420. https://doi.org/10.1080/11956860.2003.11682789

Georgiadis N, Hack M, Turpin K (2003) The influence of rainfall on zebra population

dynamics: implications for management. J Appl Ecol 40, 125–136. https://doi.org/10.10

46/j.1365-2664.2003.00796.x

Gimenez O, Lebreton JD, Choquet R, Pradel R (2017) R2ucare: Goodness-of-Fit Tests for

Capture-Recapture Models

Gimenez O, Lebreton JD, Choquet R, Pradel R (2018) R2ucare: An R package to perform

goodness-of-fit tests for capture-recapture models. bioRxiv 192468. https://doi.org/10.11

01/192468

Grafen A (1988) On the uses of data on lifetime reproductive success. Pages 454–471 in T H

Clutton-Brock, editor. Reproductive success. University of Chicago Press, Chicago,

Illinois, USA.

Grange S, Barnier F, Duncan P, Gaillard J-M, Valeix M, Ncube H, Périquet S, Fritz H (2015)

Demography of plains zebras (*Equus quagga*) under heavy predation. Popul Ecol 57,

201–214. https://doi.org/10.1007/s10144-014-0469-7

Grange S, Duncan P, Gaillard J-M, Sinclair AR, Gogan PJ, Packer C, Hofer H, East M

(2004). What limits the Serengeti zebra population? Oecologia 140, 523–532.

https://doi.org/10.1007/s00442-004-1567-6

Green WCH, Rothstein A (1993) Persistent influences of birth date on dominance, growth

and reproductive success in bison. J Zool 230, 177–186. https://doi.org/10.1111/j.1469-

7998.1993.tb02680.x

Grovenburg TW, Klaver RW, Jenks JA (2012) Survival of white-tailed deer fawns in the

grasslands of the northern Great Plains. J Wildl Manage 76, 944–956. https://doi.org/10.

1002/jwmg.339

Hamel S, Côté SD, Gaillard J-M, Festa-Bianchet M (2009) Individual variation in

reproductive costs of reproduction: High-quality females always do better. J Anim Ecol

78, 143–151. https://doi.org/10.1111/j.1365-2656.2008.01459.x

Hofmann RR (1989) Evolutionary steps of ecophysiological adaptation and diversification of

ruminants: a comparative view of their digestive system. Oecologia 78, 443–457. https://

doi.org/10.1007/bf00378733

Jackson NJ, Stewart KM, Wisdom MJ, Clark DA, Rowland MM (2021) Demographic

performance of a large herbivore: effects of winter nutrition and weather. Ecosphere 12,

e03328. https://doi.org/10.1002/ecs2.3328

Jönsson KI (1997) Capital and income breeding as alternative tactics of resource use in

reproduction. Oikos 78, 57–66. https://doi.org/10.2307/3545800

Joubert E (1974) Composition and limiting factors of a Khomas Hochland population of

Hartmann zebra *equus zebra hartmannae*. Madoqua 49–53.

Kerby J, Post E (2013) Capital and income breeding traits differentiate trophic match-

mismatch dynamics in large herbivores. Philos Trans R Soc Lond B Biol Sci 368,

20120484. https://doi.org/10.1098/rstb.2012.0484

Kiffner C, Lee DE (2019) The ecology of browsing and grazing II. In: Population dynamics

of browsing and grazing ungulates in the Anthropocene, Springer Nature, Switzerland,

pp 155–179. https://doi.org/10.1007/978-3-030-25865-8_6

Klingel H (1969) The social organisation and population ecology of the plains zebra (*Equus*

quagga). African Zool 4, 249–263. https://doi.org/10.1080/00445096.1969.11447374

Laake JL (2013) RMark: An R Interface for Analysis of Capture-Recapture Data with

MARK

Lebreton JD, Burnham KP, Clobert J, Anderson DR (1992) Modeling survival and testing

biological hypotheses using marked animals: a unified approach with case studies. Ecol

Monogr 62, 67–118. https://doi.org/10.2307/2937171

Lebreton JD, Nichols JD, Barker RJ, Pradel R, Spendelow JA (2009) Modeling individual

animal histories with multistate capture-recapture models. Adv Ecol Res 41, 87–173.

Lee DE, Bond ML, Bolger DT (2017) Season of birth affects juvenile survival of giraffe.

Popul Ecol 59, 45-54. https://doi.org/10.1007/s10144-017-0571-8

Lent PC (1974) Mother-Infant relationships in ungulates. In: The Behaviour of Ungulates and

its relation to management. International Union for Conservation of Nature and Natural

Resources, Switzerland, pp 14–55.

Lindström J (1999) Early development and fitness in birds and mammals. TREE 14, 343–

348. https://doi.org/10.1016/s0169-5347(99)01639-0

Lindström J, Kokko H (2002) Cohort effects and population dynamics. Ecol Lett 5, 338–344.

https://doi.org/10.1046/j.1461-0248.2002.00317.x

Lloyd P, Rasa O (1989) Status, reproductive success and fitness in Cape mountain zebra

(*Equus zebra zebra*). Behav Ecol Sociobiol 25, 411–420. https://doi.org/10.1007/bf0030

0187

Loison A, Langvatn R (1998) Short-and long-term effects of winter and spring weather on

growth and survival of red deer in Norway. Oecologia 116, 489–500. https://doi.org/10.1

007/s004420050614

Loveridge AJ, Searle AW, Murindagomo F, Macdonald DW (2007) The impact of sport-

hunting on the population dynamics of an African lion population in a protected area.

Biol Conserv 134, 548–558. https://doi.org/10.1016/j.biocon.2006.09.010

MacKay AE, Forsyth DM, Coulson G, Festa-Bianchet M (2018) Maternal resource allocation

adjusts to timing of parturition in an asynchronous breeder. Behav Ecol Sociobiol 72, 1-

10. https://doi.org/10.1007/s00265-017-2419-9

Mduma SAR, Sinclair ARE, Hilborn R (1999) Food regulates the Serengeti wildebeest: a 40-

year record. J Anim Ecol 68, 1101–1122. https://doi.org/10.1046/j.1365-2656.1999.0035

2.x

Ncube H, Duncan P, Grange S, et al (2011) Pattern of faecal 20-oxopregnane and oestrogen

concentrations during pregnancy in wild plains zebra mares. Gen Comp Endocrinol 172,

358–362. https://doi.org/10.1016/j.ygcen.2011.03.027

Ogutu JO, Piepho HP, Dublin HT (2014) Responses of phenology, synchrony and fecundity

of breeding by African ungulates to interannual variation in rainfall. Wildl Res 40, 698–

717. https://doi.org/10.1071/wr13117

Owen-Smith N (1990) Demography of a large herbivore, the greater kudu *Tragelaphus*

strepsiceros, in relation to rainfall. J Anim Ecol 59, 893–913. https://doi.org/10.2307/50

21

Owen-Smith and Mason (2005) Comparative changes in adult vs. juvenile survival affecting

population trends of African ungulates. J Anim Ecol 74, 762-773. https://doi.org/10.111

1/j.1365-2656.2005.00973.x

Owen-Smith N, Ogutu JO (2013) Controls over reproductive phenology among ungulates:

allometry and tropical-temperate contrasts. Ecography 36, 256–263. https://doi.org/10.1

111/j.1600-0587.2012.00156.x

Penzhorn BL (1984) Age determination in cape mountain zebras *Equus zebra zebra* in the

Mountain Zebra National Park. Koedoe 25, 89-102.

Pigeon G, Albon S, Loe LE, Bischof R, Bonenfant C, Forchhammer M, Irvine RJ, Ropstad E,

Veiberg V, Stien A (2022) Context-dependent fitness costs of reproduction despite

stable body mass costs in an Arctic herbivore. J Anim Ecol 91, 61–73. https://doi.org/10.

1111/1365-2656.13593

Plard F, Gaillard J-M, Coulson T, Hewison AM, Douhard M, Klein F, Delorme D, Warnant

C, Bonenfant C (2015) The influence of birth date via body mass on individual fitness in

a long-lived mammal. Ecology 96, 1516–1528. https://doi.org/10.1890/14-0106.1

Pluháček J, Bartoš L, Doležalová M, Bartošová-Víchová J (2007) Sex of the foetus

determines the time of weaning of the previous offspring of captive plains zebra (*Equus*

burchelli). Appl Anim Behav Sci 105, 192–204. https://doi.org/10.1016/j.applanim.2006.

05.019

Post E (2003) Timing of reproduction in large mammals. In: Phenology: an integrative

environmental science. Springer, Dordrecht, pp 437–449

Pradel R, Hines JE, Lebreton JD, Nichols JD (1997) Capture-recapture survival models

taking account of transients. Biometrics 60–72. https://doi.org/10.2307/2533097

Reznick D, Nunney L, Tessier A (2000) Big houses, big cars, superfleas and the costs of

reproduction. TREE 15, 421–425. https://doi.org/10.1016/s0169-5347(00)01941-8

Rogers CM and others (1993) A woody vegetation survey of Hwange National Park. A

woody vegetation survey of Hwange National Park.. Department of National Parks and

Wildlife Management

Rubenstein DI, Nuñez CM (2009) Sociality and reproductive skew in horses and zebras. In:

Reproductive skew in vertebrates: proximate and ultimate causes. Cambridge University

Press Cambridge, pp 196–226

Sadleir R (1982) Energy consumption and subsequent partitioning in lactating black-tailed

deer. Can J Zool 60, 382–386. https://doi.org/10.1139/z82-051

Sæther BE (1997) Environmental stochasticity and population dynamics of large herbivores:

a search for mechanisms. TREE 12, 143–149. https://doi.org/10.1016/s0169-5347(96)10

068-9

Sinclair ARE, Mduma SAR, Arcese P (2000) What determines phenology and synchrony of

ungulate breeding in Serengeti? Ecology 81, 2100–2111. https://doi.org/10.1890/0012-9

658(2000)081[2100:wdpaso]2.0.co;2

Smuts GL (1975) Pre- and postnatal growth phenomena of Burchell's zebra *Equus burchelli*

antiquorum. Koedoe 18, 69–102.

Smuts GL (1976) Reproduction in the zebra mare *Equus burchelli antiquorum* from the

Kruger National Park. Koedoe 19, 89–132. https://doi.org/10.4102/koedoe.v19i1.1186

Spagopoulou F, Teplitsky C, Chantepie S, Lind MI, Gustafsson L, Maklakov AA (2020)

Silver-spoon upbringing improves early-life fitness but promotes reproductive ageing in

a wild bird. Ecol Lett 23, 994–1002. https://doi.org/10.1111/ele.13501

Stephens PA, Boyd IL, McNamara JM, Houston AI (2009) Capital breeding and income

breeding: their meaning, measurement, and worth. Ecology 90, 2057–2067. https://doi.o

rg/10.1890/08-1369.1

Tuljapurkar S (1989) An uncertain life: demography in random environments. Theor Popul

Biol 35, 227–294. https://doi.org/10.1016/0040-5809(89)90001-4

Valeix M, Loveridge AJ, Chamaillé-Jammes S, et al (2009) Behavioral adjustments of

African herbivores to predation risk by lions: spatiotemporal variations influence habitat

use. Ecology 90, 23–30. https://doi.org/10.1890/08-0606.1

Vitet C, Duncan P, Ganswindt A, Mabika C, Chamaillé-Jammes S (2021) Do infanticides

occur in harem-forming equids? A test with long-term sociodemographic data in wild

plains zebras. Anim Behav 177, 9–18. https://doi.org/10.1016/j.anbehav.2021.04.012

Vitet C, Duncan P, Gimenez O, Mabika C, Chamaillé-Jammes S (2020) Plains zebras bring

evidence that dilution and detection effects may not always matter behaviorally and

demographically. Ecosphere 11, e03288. https://doi.org/10.1002/ecs2.3288

Weladji RB, Loison A, Gaillard J-M, Holand Ø, Mysterud A, Yoccoz NG, Nieminen M,

Stenseth NC (2008) Heterogeneity in individual quality overrides costs of reproduction

in female reindeer. Oecologia 156, 237-247. https://doi.org/10.1007/s00442-008-0961-x

Williams GC (1966) Adaptation and natural selection. Princeton University Press, Princeton,

USA.

Tables

Table 1: Set of explanatory variables used in the models investigating survival in young

individuals and mares plains zebra (*Equus quagga*), and transition probability between

reproductive and non-reproductive states in mares plains zebra (Hwange National Park,

Zimbabwe, 2008-2019).

Variable	Type	Description	Used in the model(s)
Season	Categorical	"Wet" or "Dry" season	Young & Mares
Time in dry season	Continuous	Proportion of dry season between two consecutive sessions [range: 9-80 %]	Young & Mares
Time	Categorical	ID number of the field session [range: 1-24]	Young & Mares
Trap-dependence	Categorical	The individual has been seen during the previous field session (1) or not (0)	Young & Mares
Age-class	Categorical	"Younger foal" (less than six months old), "Older foal" (between six and 12 months old) or "Yearling" (between one and two years old)	Young
Reproductive state	Categorical	The mare is seen with a dependent juvenile (1) or not (0)	Mares
Sight	Categorical	The individual is seen for the first time (1) or not (0)	Mares

 Figure legends

Figure 1: Distribution of plains zebra (*Equus quagga*) births per month, estimated from the

individual-based monitoring conducted in Hwange National Park, Zimbabwe, between 2008

and 2019 (*n* = 310 foals). The black line represents the total number of births per month over

the complete study period (i.e. all years summed). The coloured areas indicate the seasons:

blue = main wet season (approx. mid-December to mid-May), orange = main dry season

(approx. August to October), mixed colour = inter-annual variability in the start of the

seasons (start of the wet season: November to mid-December; start of the dry season: mid-

May to July).

Figure 2: Probability of annual survival of plains zebra (*Equus quagga*) young individuals in

Hwange National Park, Zimbabwe (2008-2019), according to the proportion of dry season

between two consecutive sessions. Purple: younger foals (between birth and six months old),

brown: older foals (between six and 12 months old) combined with yearlings (between one

and two years old). Solid lines represent predicted values from the best model. Shaded areas

represent 85% confidence intervals of these predicted values. Dark coloured dots represent

the survival predicted by the time model, shaded areas around the dot are proportional to

1/standard error of these predicted values.

Figure 3: a) Probability of annual survival of plains zebra (*Equus quagga*) mares in Hwange

National Park, Zimbabwe (2008-2019), according to the proportion of dry season between

two consecutive sessions and the reproductive state (for mares in their second and subsequent

observations only, see text for details). Red: non-reproductive females, blue: reproductive

females. Solid lines represent predicted values from the best model. Shaded areas represent

85% confidence intervals of these predicted values. Dots represent females survival predicted

by the time model, vertical bars represent 85% confidence intervals of these predicted values.

b) Probability of transition between reproductive and non-reproductive states for plains zebra

mares according to the session. Red: from reproductive to non-reproductive state, blue: from

non-reproductive to reproductive state. Dark coloured dots represent the survival predicted by

the time model, shaded areas around the dot are proportional to 1/standard error of these

predicted values.

Figures

Figure 1:

Figure 2:

Figure 3: