# Good moms: dependent-young and their mothers cope better than others with longer dry season in plains zebras

Lucie Thel[1], Christophe Bonenfant[1, 4] & Simon Chamaillé-Jammes[2, 3, 4]

**Author details[1]**

[1] Laboratoire de Biométrie et Biologie Évolutive, Unité Mixte de Recherche 5558, Bâtiment 711, Université Lyon I, 43 Boulevard du 11 novembre 1918, F-69622 Villeurbanne Cedex, France.

[2] CEFE, Université Montpellier, CNRS, EPHE, IRD, Montpellier, France.

[3] Mammal Research Institute, Department of Zoology & Entomology, University of Pretoria, Pretoria, South Africa.

[4] LTSER France, Zone Atelier "Hwange", Hwange National Park, Bag 62, Dete, Zimbabwe-CNRS HERD (Hwange Environmental Research Development) program.

**Corresponding author**: Thel Lucie, lucie.mc.thel@gmail.com

[1] **Author contributions**: LT, CB and SCJ were involved in the Conceptualization, Investigation, Methodology, Writing, Funding acquisition. CB and SCJ were involved in the Supervision and Validation. LT and CB were involved in the Formal analysis. LT and SCJ were involved in the Data curation. LT was involved in the Visualization. SCJ was involved in the Data collection, Project administration.

## Abstract

In large herbivores, the timing of births often matches the seasonal peak of availability of food resources, likely to improve juvenile survival and reduce reproduction costs. Some species however breed year round, even in seasonal environments. To what extent being born during the lean season reduces juveniles and reproductive females survival remains mostly unknown. This holds particularly true in tropical ecosystems where the demographic processes underlying population dynamics are often poorly studied. We investigate the variability of survival for plains zebras (*Equus quagga*) in Hwange National Park (Zimbabwe), a highly seasonal savanna ecosystem. We used capture-recapture models to analyse long-term demographic data (2008-2019). We investigated the effect of seasonality as a categorical (wet versus dry season) and continuous (duration of the dry season) variable on survival. We found little variability in the survival of younger foals (< 6 m.o.), whereas older foals and yearlings' survival, which was higher on average, decreased with the length of the dry season. Female survival was high but decreased with increasing exposure to the dry season for the non-reproductive ones only. The probability of females becoming reproductive the next year was not affected by the length of the dry season. Together, our results highlight the importance of individual quality in reproductive performance, as reproductive females seem to buffer the effect of environmental variability on their own survival and that of their foal. Timing of birth in this population might be the result of a trade off between minimising mother's reproductive costs and maximising juvenile survival.

**Keywords**: capture-mark-recapture, *Equus quagga*, Hwange National Park, reproduction, tropical ecosystem.

## Introduction

In long-lived species, the recruitment of juveniles in the population is often the most variable demographic rate, and therefore the major determinant of the variation in population growth rate (Gaillard et al. 1998, Gaillardet al. 2000, Kiffner and Lee 2019). Specifically, in large mammalian herbivores, recruitment depends mostly on juvenile survival (Gaillard et al. 2003), which is affected primarily by environmental conditions experienced by mothers during the last months of gestation and the first year of life of newborns (Sadleir 1982, Lindström 1999, Beckerman et al. 2003, Kiffner and Lee 2019). For instance, a lower fawn survival is observed in red deer (*Cervus elaphus*) when they are exposed to harsher winters (Loison and Langvaten 1998). Environmental conditions also interact with the mothers' condition and their ability to feed their offspring (Pigeon et al. 2022). For capital breeders, female body condition during the previous year or even during the current pregnancy is conditioned by resource availability at this time and can in turn affect the weight of the newborn at birth (Cameron et al. 1993) and its survival to weaning (Jackson et al. 2021). In seasonal environments where food resource availability and quality vary strongly within a year, the timing of developmental stages and of the reproductive cycle is therefore critical for juvenile survival and the mother's reproductive success.

It is well known from studies in temperate and boreal ecosystems that the timing of birth is a major life history trait of organisms, accounting for early-life survival and individual fitness (Plard et al. 2015). By determining the early environmental conditions experienced by newborns during the first months of life, timing of birth contributes directly (*e.g.* through thermal stress which may cause hypothermia; Grovenburg et al. 2012) and also indirectly (*e.g.* through potentially lower food availability reducing growth rate; Feder et al. 2008) to the annual variation of juvenile survival. The season of parturition is critical to reproductive females themselves, as they endure their offspring's needs in addition to their

own. Lactation costs are particularly high in mammals (Sadleir 1982, Clutton-Brock et al.

1989), and could make reproductive females more susceptible to harsh environmental

conditions. These selection pressures acting on newborns and mothers are thought to underlie

the commonly observed strong seasonality in time-of-birth across ungulates in temperate

ecosystems (Bronson 1989). Additionally, the timing of parturition can generate reproductive

costs for females by lowering the chances of successfully reproducing the next year (Green

and Rothstein 1991). In red deer the later a hind gives birth at any given year, the less likely

she is to conceive again during the following breeding season (Clutton-Brock et al. 1983). In

short, in temperate and boreal systems, the timing of birth accounts for the variation in

juvenile survival and reproductive success of the mother of large herbivores, explaining why

births are synchronised with the best period in terms of food resource availability and quality

(Post 2003).

Even though less predictable than temperate and boreal ecosystems, arid and semi-

arid ecosystems also show a pronounced seasonality, i.e. wet and dry seasons mainly driven

by the distribution of rains within the year (Desmukh 1984, Owen-Smith and Ogutu 2013).

Although some large herbivores reproduce according to the seasons (e.g. the blue wildebeest

(*Connochaetes taurinus*); Sinclair et al. 2000), others have more extended breeding periods

(Owen-Smith and Ogutu 2013), or even breed year-round (e.g. the plains zebras (*Equus*

*quagga*); Sinclair et al. 2000). In such species, and even if a peak of birth is often observed

during the wet season, some females give birth during the dry season, at a time when food

and water are more scarce (Boutton et al. 1988a, Boutton et al. 1988b, Feng et al. 2013) and

susceptibility to predation is increased (Valeix et al. 2009). This means that a variable share

of the females of a population give birth in less favourable conditions than the others,

potentially exposing their offspring and themselves to a higher mortality risk. An interesting

and yet unanswered question is therefore: for such species, to what extent does giving birth in

the dry unfavourable season reduce the probability of survival of both the newborn and its

mother?

To date, we lack empirical studies exploring juvenile survival and reproductive costs

both in species reproducing year-round and in tropical large herbivores in general (Clutton-

Brock and Sheldon 2010, Burthe et al. 2011, Festa-Bianchet et al. 2017, but see Lee et al.

2017). In this study, we investigated the effect of the season of birth on the annual survival of

young and mares in relation to environmental conditions in wild plains zebras (*Equus*

*quagga*). We explored the impact of the duration of the dry season, which varies according to

the timing of rainfalls, on the cohort-specific annual survival of foals, yearlings and adult

females. We took advantage of a long-term individual-based monitoring program of plains

zebras conducted in Hwange National Park (HNP) in Zimbabwe. Even if their environment is

seasonal (Chamaille-Jammes et al. 2006), plains zebras breed year round in this ecosystem

(Fig. 1), and are exposed to a high inter-annual variability in the duration of the dry season.

This constitutes an adequate framework to study the impact of the timing of reproduction on

the survival of young and females in seasonal tropical environments.

Specifically, we hypothesised that the dry season, and in particular long dry seasons

which lead to harsh conditions for herbivores, should decrease plains zebra foal survival until

weaning (0 to 6 months old) because this period corresponds to the most vulnerable stage

(Caughley 1966, Festa-Bianchet 1988). Similarly, harsh environmental conditions during

early life should decrease foal survival between weaning and complete independence from

the mother (6 to 12 months old) because they are arguably less vulnerable, are still growing

and cannot rely on their mother anymore for food provisioning and protection (Smuts 1976,

Côté and Festa-Bianchet 2001). We expected a comparatively smaller effect of exposure to

harsh conditions on yearling (1 to 2 years old) survival than on foal, because older individuals

should have more resources to survive through periods of low food availability (Gaillard et

al. 2000). Regarding mares, we expected that costs of reproduction would be reflected in a lower survival of reproductive females compared to non-breeding ones. Indeed, a dependent foal imposes additional energy expenditure for reproduction whereas the amount of energy available to the mother is limited (energy allocation hypothesis; Cody 1966, Williams 1966). Alternatively, one could expect the opposite, with reproductive females coping better than breeding females as non-reproductive females could be those of lower phenotypic quality, explaining that they do not even breed (Reznick et al. 2000, Hamel et al. 2009), and overall would be more vulnerable to environmental harshness.

## Materials and methods

*1. Context of the study*

*1.1. Study site*

HNP (19°00' S, 26°30' E), Zimbabwe, covers 14,651 km$^2$ of bushlands, woodlands (mostly *Colophospermum mopane*, *Combretum spp.*, *Acacia spp.*, *Baikiaea plurijuga* and *Terminalia sericea*, Rogers et al. 1993) and scrublands interspersed with grassland (Arraut et al. 2018), and is located between 900 and 1,100m a.s.l.. Average annual rainfall is approximately 600mm, with marked inter-annual variations (Chamaille-Jammes et al. 2006). On average, 98% of the rainfall events occur during the wet season, generally from October to April (Chamaille-Jammes et al. 2006, Chamaille-Jammes et al. 2007). The start of the wet season is characterised by a high inter-annual variability, leading to variable duration of the dry season. The study took place in the north-east of the park, where artificial waterholes retain water year round and where areas > 8km from a waterhole are rare, even in the dry season (Chamaille-Jammes et al. 2007). The densities of lion (*Panthera leo*) and spotted hyena (*Crocuta crocuta*), the two main predators of plains zebra, are relatively high (Loveridge et al. 2007, Drouet-Hoguet 2007). There is no hunting in the park.

*1.2. Study species*

Plains zebras live in harems composed of a stallion, several mares and their offspring under two years old (Klingel 1969). Females give birth to a singleton year round in most of their geographic range, including Zimbabwe (Dasmann and Mossman 1962, Barnier et al. 2012), even if a birth peak can occur from January to March in this area (Fig. 1). Gestation length ranges between 340 and 390 days and foals are weaned around 11 months of age (Smuts 1976). Plains zebra foals are considered as "followers" on the hider-follower gradient (Lent 1974), as they stand and follow their mother a couple of hours after birth (Sinclair2000). Plains zebras are grazers (*sensu* Hofman 1989), feeding virtually only on grasses, hence the availability of their food resource is directly linked to rainfall amount and distribution within the year, as well as with the general availability of habitats having a grass layer (DuPlessis 2001). As the other large herbivores occupying African savannas, the plains zebra is generally considered to be located more towards the capital end of the capital-income breeder continuum (Jönsson 1997, Stephens et al. 2009), even if there is currently little empirical assessment of this tactic of energy allocation (but see Ogutu et al. 2014). In HNP, the population of plains zebras is mostly resident thanks to the artificial waterholes providing water year round inside the park, but some seasonal shifts in home-ranges have been observed for some harems (unpublished GPS data).

*1.3. Seasonality and plains zebra life histories*

Following the protocol presented in Grange et al. (2015), we recorded the presence of individually identified plains zebras between 2008 and 2019 using visual identification of their unique stripe pattern. Observations were conducted twice a year, around March and August, during field sessions (hereafter called "sessions") at the transition between wet and dry seasons ($n = 24$ sessions, *mean session duration* $\pm$ *SD* $= 45 \pm 25$ days, *range* $= 13 - 88$ days). When first sighted, individuals were classified according to three age-classes: foal

(from birth to 12 months old), yearling (12 to 24 months old) and adults (> 24 months old).

The age of the young individuals was then determined more accurately based on their relative

size compared to adults, using the criteria of Smuts (1975) and Penzhorn (1984), and

photographs of individuals of known age from HNP. We also arbitrarily attributed an

accuracy to this age estimation, based on our experience from the field. The age of very

young foals was considered accurate to a few days, whereas young seen at an older age for

the first time could be attributed an age $\pm$ 2 months, for instance.

The main goal of this study was to explore the effect of seasonality as a categorical

(i.e. wet season versus dry season) or continuous (i.e. duration of the dry season) variable on

the survival of young and mares, accounting for the reproductive state of the mare and the

age-class of the young. We also assessed the probability of transition between reproductive

and non-reproductive states in mares from one session to the next. Explanatory variables are

described in the following sections and summarised in Table 1.

*2. Explanatory variables*

For each year, we identified the transition date between wet and dry season using 500m

spatial resolution, bi-monthly Normalised Difference Vegetation Index (NDVI) records from

the NASA website (MOD13A1 product, computed from atmospherically corrected bi-

directional surface reflectances masked for water, clouds, heavy aerosols, and cloud shadows;

temporal resolution: 15 days; https://modis.gsfc.nasa.gov) and daily rainfall records from the

Climate Hazards Center website (Rainfall Estimates from Rain Gauge and Satellite

Observations, https://www.chc.ucsb.edu). During the study period and according to our

estimations, the wet season in HNP started between the 1st of November and the 19th of

December, while the dry season started between the 9th of May and the 29th of July (Fig. 1,

Online Resource 1). We attributed a season to each session according to the date of the

session (variable *season*, Table 1). We also estimated the proportion of days of dry season

between the first day of the session *s* and the first day of the following session *s+1* (variable

*pds*, namely "proportion of dry season", Table 1). We defined three age-classes among the

young: "younger foals" of less than six months old, "older foals" aged between six and 12

months old, and "yearlings" between one and two years old (Gaillard et al. 2000, Table 1).

*3. Statistical analysis*

*3.1. Capture-Mark-Recapture framework*

We used a Capture-Mark-Recapture (CMR, Lebreton et al. 1992) framework to account for

the variability of the probability of recapture (i.e. re-observation) of the individuals of this

population (Grange et al. 2015). We considered each session as a discrete event, summarised

by its starting date. As the time interval between two successive sessions varied, we used the

*time.intervals* argument of the *process.data* function to correct survival estimates for the

duration of exposure to mortality risk (*RMark* package, Laake 2013). For this, we calculated

the proportion of time elapsed between two successive sessions in relation to a year of 365

days: $\Delta t_{session2-session1} = (Starting\ date_{session2} - Starting\ date_{session1}) / 365$. We performed all

CMR analyses using the program MARK (www.phidot.org/software/mark) and the R

statistical software (version 4.3.1, www.r-project.org), thanks to the package *RMark* (Laake

2013). The goodness-of-fit tests (GOF tests: see below) were conducted with the R package

*R2ucare* (Gimenez et al. 2017).

*3.2. Survival of the young*

Similarly to previous demographic analyses performed on these data (Grange et al. 2015,

Vitet et al. 2020, Vitet et al. 2021), we built capture histories for each individual as follows: 0

corresponded to "no sighting of the individual during the session", and 1 corresponded to "at

least one sighting of the individual during the session". We ran the analyses on individuals of

known date of birth with an accuracy ranging from 0 to $\pm$ 90 days (*n* = 310). The dataset

contained individuals observed at least once in the field (*n* = 290) and individuals never

observed in the field but whose mother was detected to be pregnant ($n = 20$). For the latter,

the date of parturition was estimated based on opportunistic faecal sampling from the mare

and subsequent 20-oxopregnanes and oestrogens enzyme immunoassay (Online Resource 2,

Ncube et al. 2011). We recorded these individuals as being identified at birth only and never

seen again, assuming no foetal loss or abortion as usually reported in large mammals. All the

individuals born in the same cohort (i.e. which had the same session as the one closest to their

date of birth) were considered to have experienced the same environmental conditions (i.e.

*season* and *proportion of dry season* variables). Because of the imperfect detection, we kept

sub-adults and adults (> 2 years old) in the model to get better estimations of yearling

survival. In other words, individuals re-observed only at an age of 2 years or more but missed

during the previous sessions still contributed to the estimation of survival of the young, even

if adult survival *per se* was not presented in this analysis.

In a preliminary step, we performed goodness-of-fit tests to check if the dataset of

young individuals did fit the CMR model assumptions (Lebreton et al. 1992). The GOF tests

of the fully time-dependent model (Gimenez et al. 2018) highlighted possible overdispersion

(Test 2.CL: $\chi^2 = 40.79$, $df = 16$, $P < 0.01$; Test 3.Sm: $\chi^2 = 11.05$, $df = 19$, $P = 0.92$), owing to

trap-dependence ($\chi^2 = 139.35$, $df = 17$, $P < 0.01$) and transience issues ($\chi^2 = 70.47$, $df = 22$, $P$

$< 0.01$). After examination of Test 2.CL, we noticed that overdispersion was mainly caused

by three sessions over the 24 available in the dataset, so we considered this lack of fit as

marginal and ignored overdispersion in the analyses.

We accounted for trap-dependence by adding a trap-dependence effect in the

recapture model (*td* variable, Table 1). As plains zebras' distribution within the park could

depend on the seasonal dynamics, we tested the effect of the *season* or *proportion of dry*

*season* in the recapture model. Transience was likely due to the age structure as young

individuals often have lower survival in large herbivores (Gaillard et al. 2000) and are thus

less frequently recaptured than adults, leading to a lower recapture probability. Thus, we

fitted the survival model with an *age-class* effect (Table 1). We also explored the effect of

several poolings of those age-classes (e.g. older foals and yearlings pooled together as one

single age-class versus younger foals). We tested the effect of the *season* or *proportion of dry*

*season* in the young survival models (Online Resource 3). For comparison purposes, we also

conducted an analysis of the survival of the juveniles using a Generalised Linear Model

(GLM) approach instead of a CMR approach (Online Resource 4).

*3.3. Mare survival*

To study the probability of survival and transition between reproductive states in adult

females ($n = 322$), we used a multi-states capture-recapture model (Lebreton et al. 2009). We

built capture histories for each individual as follows: 0 corresponds to "no sighting of the

mare during the session", 1 corresponding to "at least one sighting of the mare without her

dependent juvenile during the session" and 2 corresponding to "at least one sighting of the

mare with her dependent juvenile during the session". We considered a dependent juvenile as

an individual that was still suckling, so under 6 months of age. Even if weaning generally

occurs around 11 months of age in zebras, juveniles can survive without their mother from 9

months old onwards (Smuts 1976). We chose 6 months to exclude such possibility and match

our sessions frequency.

As there were too few observations to conduct the GOF tests of the multi-states

model, we conducted the GOF tests on the one-state model instead (i.e. without taking into

account reproductive states). The tests revealed overdispersion (Test 2.CL: $\chi^2 = 49.02$, $df =$

19, $P < 0.01$; Test 3.Sm: $\chi^2 = 72.10$, $df = 20$, $P < 0.01$), trap-dependence ($\chi^2 = 112.13$, $df = 21$,

$P < 0.01$) and transience ($\chi^2 = 62.23$, $df = 21$, $P < 0.01$). After examination of Test 2.CL and

3.Sm, we noticed that overdispersion was mainly caused by five and three sessions over the

24 available in the dataset, respectively. Again, we considered it as marginal and ignored

overdispersion in the analyses. We accounted for trap-dependence by adding a trap-

dependence effect (*td* variable, Table 1) in the recapture model. To take into account

transience, we added a categorical variable *sight* in the survival model to differentiate

between mares captured for the first time during the survey and mares already captured at

least once during the survey, following the method described by Pradel et al. (1997; *sight*

variable, Table 1). We evaluated the effect of the *reproductive state* (Table 1), and *season* or

*proportion of dry season* on recapture, survival and transition probabilities. Age being not

known for most adult females, we were unable to include this variable in the model.

*3.4. Model selection*

For both young and mares, we conducted model selection by comparing models ranging from

the most complex full time-dependent model (*time* variable, i.e. chronological number

identifying the session, Table 1) to the reference model (i.e. depending exclusively on the

covariables related to GOF corrections, Online Resource 3). Because of the huge number of

possible combinations between recapture and survival (and transition for the mares dataset)

models, we conducted a sequential model selection starting by modelling recapture, survival

and then transition probabilities (Lebreton et al. 1992). We conducted a first model selection

to identify the best recapture model, while survival model was set to depend exclusively on

the covariables related to GOF corrections (i.e. *age-class* for young individuals and *sight* for

mares) and transition probability was set constant for mares. We then modelled survival

probability using the best recapture model and constant transition probability for mares. We

eventually selected the best transition probability model with the best recapture and survival

models for mares.

We used the Akaike Information Criterion adjusted for small sample sizes (AICc) and

we selected the model with the lowest number of parameters for models having a $\Delta$AICc < 2

when compared to the best model (parsimony principle, Burnham and Anderson 2002).

Following Arnold (2010), we reported the 85% confidence intervals of estimated parameters,

in accordance with our AICc model selection procedure. We used the scaled value of

*proportion of dry season* in the models to ease model convergence, but for convenience, we

back-transformed the coefficients to their initial scale in the Results section.

**Results**

*1. Survival of the young*

Results on recapture probabilities of juveniles are presented in Online Resource 5. In

summary, recapture probability was best described by an effect of *trap dependence* and

*proportion of dry season*, and overall was within the 0.3 - 0.5 range. According to the most

parsimonious model, survival varied between age-classes (see Online Resource 3 for the

detailed model selection). The average probability of annual survival was lower for younger

foals ($\varphi = 0.458 \pm 0.044$ SE, 85% CI [0.395; 0.522]) than for the two other age-classes, which

had a similar survival that decreased with the *proportion of dry season* ($\beta = -0.026 \pm 0.016$

SE, 85% CI [-0.049; -0.002]). The predicted probability of survival of older foals and

yearlings varied from $0.850 \pm 0.095$ SE, 85% CI [0.661; 0.943] when the *proportion of dry*

*season* was the shortest (i.e. 9 % of the time) to $0.480 \pm 0.120$ SE, 85% CI [0.316; 0.648]

when the *proportion of dry season* was the longest (i.e. 80 % of the time, Fig. 2). The *season*

variable was not retained in the best models of both the CMR or the GLM approaches. Even

though non-significant, a quick exploration of survival probabilities according to the season

and time in older foals and yearling survival showed that it was likely lower and more

variable during the dry ($0.536 \pm 0.221$ SD) than during the wet season ($0.707 \pm 0.172$ SD).

Note that an alternative, simplest, analysis based on a GLM approach provided similar results

(Online Resource 4).

*2. Mare survival*

Overall, the recapture probability was higher for mares than for young individuals, generally

around 0.7 (Online Resource 5). The best model for survival probability included an additive

effect of *sight*, *proportion of dry season* and *reproductive state*, and an additive effect of *time*

and *reproductive state* for transition probability (see Online Resource 3 for the detailed

model selection). As the *sight* variable had a significant effect on the probability of survival,

we present results for mares in their second and following observations only, because results

relying on the first observation of a mare are less robust and informative. The *proportion of*

*dry season* had a negative effect on survival ($\beta = -0.035 \pm 0.011$ SE, 85% CI [-0.050; -0.019],

Fig. 3a). The probability of survival of reproductive females varied slightly from $0.995 \pm$

$0.006$ SE, 85% CI [0.970; 0.999] when the *proportion of dry season* was the shortest to 0.945

$\pm 0.059$ SE, 85% CI [0.768; 0.989] when the *proportion of dry season* was the longest.

Although the interaction between the *proportion of dry season* and the reproductive state of

the females was not retained in the best model, we noted that the probability of survival of

non-reproductive females appeared to vary more largely, from $0.966 \pm 0.019$ SE, 85% CI

[0.924; 0.985] to $0.703 \pm 0.057$ SE, 85% CI [0.615; 0.779] as the *proportion of dry season*

increased.

The probability of transition between reproductive states was not dependent on either

the *season* or the *proportion of dry season* as neither of these variables were selected in the

best model, but was highly variable according to *time*. As could be expected from the

phenotypic quality hypothesis, the probability for a mare to switch from the reproductive to

the non-reproductive state between two consecutive sessions was higher (from $0.582 \pm 0.094$

SE, 85% CI [0.443; 0.709] to $0.961 \pm 0.015$ SE, 85% CI [0.933; 0.978]) than the probability

to transition from the non-reproductive to the reproductive state (from $0.049 \pm 0.018$ SE, 85%

CI [0.028; 0.083] to $0.475 \pm 0.093$ SE, 85% CI [0.346; 0.608], Fig. 3b). Based on the model

including the season and reproductive state as explanatory variables for the probability of

transition, the average probability to be reproductive one year later was not different between

reproductive females (0.423) and non-reproductive females (0.420).

**Discussion**

Our study shows that even if the general survival pattern of mammalian populations

(Caughley 1966, Gaillard et al. 2000) also fits plains zebra in a tropical context, the expected

negative effects of the duration of dry season on juvenile survival was not consistently

observed across all age classes. We found the year-round breeding strategy of female plains

zebras to have little consequences for early survival of foals, and we relate it to the long

gestation length and the possible strategy of energy allocation of mares to their offspring. We

document, however, a within-year and between years variability of young survival rates with

potential consequences for population dynamics.

In contrast with the younger foals, older foals and yearlings had a lower survival

when experiencing long dry seasons. To our knowledge, studies so far mostly addressed the

difference between juvenile and adult survival, but comparisons of survival between different

"young" age classes remains rare. Georgiadis et al. (2003) modelling work suggested a higher

sensitivity of the survival of plains zebra foals under one year of age to fluctuations in

rainfalls than adults. Sensitivity to environmental conditions of young individuals is a

commonly observed pattern, even in tropical herbivores: for instance, juveniles and

yearlings' survival is highly correlated with the preceding annual rainfalls in the greater kudu

(*Tragelaphus strepsiceros*, Owen-Smith 1990) and calf survival decreases during the dry

season in the wildebeest (*Connochaetes taurinus*, Mduma et al. 1999). Young plains zebras

are weaned around 11 months (Smuts 1976), but they only reach their adult size around three

years of age (Smuts 1975). During this critical period, older foals and yearlings, which are

still growing, remain more vulnerable than adults in the face of poor environmental

conditions or predation, explaining their lower survival.

Breeding females usually time their reproduction to match the most costly part of the

reproductive cycle, corresponding to the late gestation and early lactation in large herbivores

(Sadleir 1982, Clutton-Brock et al. 1989), with the period of best food quality or availability

(Bronson 1989). This matching between food resources and costly reproductive events should

minimise subsequent fitness costs like a lower survival, litter size or probability to reproduce

after a successful reproduction. Accordingly, most of the plains zebra mares in HNP gave

birth during the wet season (Fig. 1). The clear and striking result of our study is that, when

environmental conditions are harsher (longer dry seasons), mothers appear to buffer the costs

for the dependent young (i.e. younger foal). This suggests that plains zebras might be capital

breeders, with females engaging in reproduction conditional to a sufficient acquisition of

energy during the prior non-reproductive period. This would guarantee their ability to cope

with their offspring requirements in case of an unexpectedly long dry season. Only in the case

of extreme drought reproductive mares and their dependent foals have been found to

experience higher mortality than usual (Joubert 1974). In our system, the impact of droughts

on plains zebras, which are able to feed on rather low quality grasses, is likely buffered by the

presence of permanent waterholes. Combining these demographic data with explorations of

the existence of accumulation and storage of energy by the females during favourable periods

would help in clarifying the plains zebra's reproductive strategy. Also, it would shed light on

the validity of the capital-income breeder concept in tropical herbivores. Indeed, such a

classification has so far rarely been used in these systems because of the lack of adequate

demographic data, contrary to temperate ecosystems where it is a common conceptual

framework (e.g. Kerby and Post 2013, Apollonio et al. 2020).

Given our results and somewhat counterintuitively, the optimal timing of birth in

order to minimise mortality risks of young plains zebras should be the beginning of the dry

season. Juveniles could thus benefit from their mother's protection during the harsher season and enter the age of independence during the most favourable one. Similarly, Lee et al. (2017) showed that giraffes (*Giraffa camelopardalis*) born during the dry season experienced a higher survival probability in Tanzania. Even though most of the plains zebras' births in our population happened during the wet season, ensuring sufficient food resources for the mother at a critical stage of their reproductive cycle, we note that the birth peak also extended towards the beginning of the dry season (Fig. 1), which is more favourable to the offspring. Such a phenomenon could illustrate a potential trade-off between minimising mother's reproductive costs and maximising juveniles' survival, as it has been described in primates (Dezeure et al. 2021).

The strength of this trade-off might not however be so large. Indeed, the length of the dry season had only a small impact on the survival of reproductive females, whose survival always remained very high (~ 0.95 or above). Data also suggested that reproductive females are not more sensitive, and possibly are less sensitive, than non-reproductive females to environmental conditions. Females engaging in reproduction thus appear to not only be able to buffer the effect of the environment on their young foal, but also to bear the costs of reproduction, even under harsh conditions, with minimal impact. They are also able to engage in future reproduction as quickly as non-reproductive females. All these results point toward the fact that females that reproduce might be those of high intrinsic phenotypic quality, or experience. Again, if these effects have been well described and studied in temperate or boreal ungulates (e.g. Hamel et al. 2009), our results generalise these observations to tropical species. Although this was not the goal of this research, getting a more mechanistic understanding of what makes some females more prone to reproduce would be interesting. As both reproductive and non-reproductive females can be found in the same harems, with thus similar landscape use, we can rule out that differences in habitat availability or habitat

selection strategies explain the observed contrasts between females of different reproductive

states. Given the importance of age and dominance status (which are often related in equids)

on reproduction previously demonstrated in horses and zebra species (Lloyd and Rasa 1989,

Pluháček et al. 2007, Rubenstein and Nuñez 2009), we suspect that they are key traits

underlying the observed difference. Unfortunately, age, which was sometimes available

(dominance status was not) could however not be included in our analysis without reducing

its power.

Overall, we document marked variations in demographic rates (juvenile survival and

females' reproductive rates) of plains zebras in HNP, both between seasons and across the

years, partly accounted for by the duration of the dry season. This creates cohort effects,

which have been documented in many long-lived species in temperate areas (see Sæther

1997, Gaillard et al. 2003 for reviews in large herbivores, Kiffner and Lee 2019) but so far

haven't been much studied in tropical ecosystems (e.g. in the greater kudu, Owen-Smith

1990). Here, our results indicate that juvenile survival of plains zebras is structured according

to two levels of temporal variation, the seasonal and yearly levels. These two sources of

juvenile survival variation have the potential to generate more cohorts of individuals with

contrasting life histories in the population than for seasonal breeding species. This particular

pattern of juvenile survival variability might destabilise plains zebra population dynamics

because of cohort effects (Lindström and Kokko 2002) and reduce the long-term stochastic

population growth rate because of its sensitivity to the temporal variance of demographic

rates (Tuljapurkar 1989).

## Acknowledgments

This long-term demographic survey was initiated by Patrick Duncan, who also took part in many of the field sessions. The authors are therefore indebted to him, and also acknowledge all the other people involved in data collection over the years, in particular S. Grange, F. Barnier, H. Ncube, C. Vitet and A. Duncan. J. Dezeure assisted with NDVI and rainfall data retrieval.

## Declarations

**Funding**: This work was supported by a PhD grant from the "Ministère Français de l'Enseignement Supérieur, de la Recherche et de l'Innovation" through the "Ecole Doctorale E2M2" of the "Université Claude Bernard Lyon 1" attributed to LT.

**Conflict of Interest**: The authors declare that they have no conflict of interest.

**Ethical approval**: Ethics approval was not required for this study according to local legislation [Parks and Wildlife Act].

**Consent to participate**: Not applicable.

**Consent for publication**: Not applicable.

**Availability of data and material**: Data available from the corresponding author on reasonable request.

**Code availability**: Not applicable.

**Authors' contributions**: LT, CB and SCJ were involved in the Conceptualization, Investigation, Methodology, Writing, Funding acquisition. CB and SCJ were involved in the Supervision and Validation. LT and CB were involved in the Formal analysis. LT and SCJ were involved in the Data curation. LT was involved in the Visualization. SCJ was involved in the Data collection, Project administration.

## References

**References**

Apollonio M, Merli E, Chirichella R, Pokorny B, Alagić A, Flajšman K, Stephens PA (2020)

Capital-income breeding in male ungulates: Causes and consequences of strategy

differences among species. Front ecol evol 8:521767.

https://doi.org/10.3389/fevo.2020.521767

Arnold TW (2010) Uninformative parameters and model selection using Akaike's

Information Criterion. J Wildl Manage 74:1175–1178. https://doi.org/10.1111/j.1937-

2817.2010.tb01236.x

Arraut EM, Loveridge AJ, Valls-Fox H, Macdonald DW, Chamaillé-Jammes S (2018) The

2013-2014 vegetation structure map of Hwange National Park, Zimbabwe, produced

using free satellite images and software. Koedoe African Prot Area Conserv Sci 60:1–

10. https://doi.org/10.4102/koedoe.v60i1.1497

Barnier F, Grange S, Ganswindt A, Ncube H, Duncan P (2012) Inter-birth interval in zebras

is longer following the birth of male foals than after female foals. Acta Oecologica 42,

11–1. https://doi.org/10.1016/j.actao.2011.11.007

Beckerman AP, Benton TG, Lapsley CT, Koesters N (2003) Talkin' 'bout my generation:

environmental variability and cohort effects. Am Nat 162, 754–767.

https://doi.org/10.1086/381056

Boutton TW, Tieszen LL, Imbamba SK (1988a) Biomass dynamics of grassland vegetation in

Kenya. Afr J Ecol 26, 89–101. https://doi.org/10.1111/j.1365-2028.1988.tb00960.x

Boutton TW, Tieszen LL, Imbamba SK (1988b) Seasonal changes in the nutrient content of

East African grassland vegetation. Afr J Ecol 26, 103–115.

https://doi.org/10.1111/j.1365-2028.1988.tb00961.x

Bronson FH (1989) Mammalian reproductive biology. University of Chicago Press, Chicago,

USA

Burnham K, Anderson D (2002) Model Selection and Multimodel Inference: A Practical

Information-Theoretic Approach. Springer, New York, USA

Burthe S, Butler A, Searle KR, Hall SJ, Thackeray SJ, Wanless S (2011) Demographic

consequences of increased winter births in a large aseasonally breeding mammal (*Bos*

*taurus*) in response to climate change. J Anim Ecol 80, 1134–1144.

https://doi.org/10.1111/j.1365-2656.2011.01865.x

Cameron RD, Smith WT, Fancy SG, Gerhart KL, White RG (1993) Calving success of

female caribou in relation to body weight. Can J Zool 71, 480–486.

https://doi.org/10.1139/z93-069

Caughley G (1966) Mortality patterns in mammals. Ecol 47, 906–918.

https://doi.org/10.2307/1935638

Chamaillé-Jammes S, Fritz H, Murindagomo F (2007) Detecting climate changes of concern

in highly variable environments: Quantile regressions reveal that droughts worsen in

Hwange National Park, Zimbabwe. J Arid Environ 71, 321–326.

https://doi.org/10.1016/j.jaridenv.2007.05.005

Chamaillé-Jammes S, Fritz H, Murindagomo F (2006) Spatial patterns of the NDVI-rainfall

relationship at the seasonal and interannual time scales in an African savanna. Int J

Remote Sens 27, 5185–5200. https://doi.org/10.1080/01431160600702392

Clutton-Brock T, Sheldon BC (2010) Individuals and populations: the role of long-term,

individual-based studies of animals in ecology and evolutionary biology. Trends Ecol &

Evol 25, 562–573. https://doi.org/10.1016/j.tree.2010.08.002

Clutton-Brock TH, Albon SD, Guinness FE (1989) Fitness costs of gestation and lactation in

wild mammals. Nature 337, 260–262. https://doi.org/10.1038/337260a0

Clutton-Brock TH, Guinness FE, Albon SD (1983) The costs of reproduction to red deer

hinds. J Anim Ecol 52, 367–383. https://doi.org/10.2307/4560

Cody ML (1966) A general theory of clutch size. Evolution 20, 174–184.

https://doi.org/10.2307/2406571

Côté SD, Festa-Bianchet M (2001) Birthdate, mass and survival in mountain goat kids:

effects of maternal characteristics and forage quality. Oecologia 127, 230–238.

https://doi.org/10.1007/s004420000584

Dasmann RF, Mossman AS (1962) Reproduction in some ungulates in Southern Rhodesia. J

Mammal 43, 533–537. https://doi.org/10.2307/1376916

Deshmukh I (1984) A common relationship between precipitation and grassland peak

biomass for east and southern Africa African. J Ecol 22, 181–186.

https://doi.org/10.1111/j.1365-2028.1984.tb00693.x

Dezeure J, Baniel A, Carter A, Cowlishaw G, Godelle B, Huchard E (2021) Birth timing

generates reproductive trade-offs in a non-seasonal breeding primate. Proc R Soc B 288,

20210286. https://doi.org/10.1098/rspb.2021.0286

Drouet-Hoguet N (2007) Influence des activités anthropogéniques sur la régime alimentaire

et la réponse numérique de la hyène tachetée en savane arborée dystrophique dominée

par l'éléphant. PhD Thesis Lyon 1 University, France.

Du Plessis WP (2001) Effective rainfall defined using measurements of grass growth in the

Etosha National Park, Namibia. J Arid Environ 48, 397–417.

https://doi.org/10.1006/jare.2000.0752

Feder C, Martin JGA, Festa-Bianchet M, Bérubé C, Jorgenson J (2008) Never too late?

Consequences of late birthdate for mass and survival of bighorn lambs. Oecologia 156,

773–781. https://doi.org/10.1007/s00442-008-1035-9

Feng X, Porporato A, Rodriguez-Iturbe I (2013) Changes in rainfall seasonality in the tropics.

Nat Clim Chang 3, 811–815. https://doi.org/10.1038/nclimate1907

Festa-Bianchet M (1988) Birthdate and survival in bighorn lambs (*Ovis canadensis*). J Zool

214, 653–661. https://doi.org/10.1111/j.1469-7998.1988.tb03764.x

Festa-Bianchet M, Douhard M, Gaillard J-M, Pelletier F (2017) Successes and challenges of

long-term field studies of marked ungulates. J Mammal 98, 612–620.

https://doi.org/10.1093/jmammal/gyw227

Gaillard JM, Festa-Bianchet M, Yoccoz NG, Loison A Toigo C (2000) Temporal variation in

fitness components and population dynamics of large herbivores. Annu Rev Ecol Syst

31, 367–393. https://doi.org/10.1146/annurev.ecolsys.31.1.367

Gaillard JM, Festa-Bianchet M, Yoccoz NG (1998) Population dynamics of large herbivores:

variable recruitment with constant adult survival. Trends Ecol & Evol 13, 58–63.

https://doi.org/10.1016/s0169-5347(97)01237-8

Gaillard JM, Loison A, ToÏgo C, et al (2003) Cohort effects and deer population dynamics.

Ecoscience 10, 412–420. https://doi.org/10.1080/11956860.2003.11682789

Georgiadis N, Hack M, Turpin K (2003) The influence of rainfall on zebra population

dynamics: implications for management. J Appl Ecol 40, 125–136.

https://doi.org/10.1046/j.1365-2664.2003.00796.x

Gimenez O, Lebreton JD, Choquet R, Pradel R (2017) R2ucare: Goodness-of-Fit Tests for

Capture-Recapture Models

Gimenez O, Lebreton JD, Choquet R, Pradel R (2018) R2ucare: An R package to perform

goodness-of-fit tests for capture-recapture models. bioRxiv 192468.

https://doi.org/10.1101/192468

Grange S, Barnier F, Duncan P, Gaillard JM, Valeix M, Ncube H, Périquet S, Fritz H (2015)

Demography of plains zebras (*Equus quagga*) under heavy predation. Popul Ecol 57,

201–214. https://doi.org/10.1007/s10144-014-0469-7

Green WCH, Rothstein A (1991) Trade-offs between growth and reproduction in female

bison. Oecologia 86, 521–527. https://doi.org/10.1007/bf00318318

Grovenburg TW, Klaver RW, Jenks JA (2012) Survival of white-tailed deer fawns in the

grasslands of the northern Great Plains. J Wildl Manage 76, 944–956.

https://doi.org/10.1002/jwmg.339

Hamel S, Côté SD, Gaillard J-M, Festa-Bianchet M (2009) Individual variation in

reproductive costs of reproduction: High-quality females always do better. J Anim Ecol

78, 143–151. https://doi.org/10.1111/j.1365-2656.2008.01459.x

Hofmann RR (1989) Evolutionary steps of ecophysiological adaptation and diversification of

ruminants: a comparative view of their digestive system. Oecologia 78, 443–457.

https://doi.org/10.1007/bf00378733

Jackson NJ, Stewart KM, Wisdom MJ, Clark DA, Rowland MM (2021) Demographic

performance of a large herbivore: effects of winter nutrition and weather. Ecosphere 12,

e03328. https://doi.org/10.1002/ecs2.3328

Jönsson KI (1997) Capital and income breeding as alternative tactics of resource use in

reproduction. Oikos 78, 57–66. https://doi.org/10.2307/3545800

Joubert E (1974) Composition and limiting factors of a Khomas Hochland population of

Hartmann zebra *equus zebra hartmannae*. Madoqua 49–53.

Kerby J, Post E (2013) Capital and income breeding traits differentiate trophic match-

mismatch dynamics in large herbivores. Philos Trans R Soc Lond B Biol Sci 368,

20120484. https://doi.org/10.1098/rstb.2012.0484

Kiffner C, Lee DE (2019) The ecology of browsing and grazing II. In: Population dynamics

of browsing and grazing ungulates in the Anthropocene, Springer Nature, Switzerland,

pp 155–179. https://doi.org/10.1007/978-3-030-25865-8_6

Klingel H (1969) The social organisation and population ecology of the plains zebra (*Equus*

*quagga*). African Zool 4, 249–263. https://doi.org/10.1080/00445096.1969.11447374

Laake JL (2013) RMark: An R Interface for Analysis of Capture-Recapture Data with

MARK

Lebreton JD, Burnham KP, Clobert J, Anderson DR (1992) Modeling survival and testing

biological hypotheses using marked animals: a unified approach with case studies. Ecol

Monogr 62, 67–118. https://doi.org/10.2307/2937171

Lebreton JD, Nichols JD, Barker RJ, Pradel R, Spendelow JA (2009) Modeling individual

animal histories with multistate capture-recapture models. Adv Ecol Res 41, 87–173.

Lee DE, Bond ML, Bolger DT (2017) Season of birth affects juvenile survival of giraffe.

Popul Ecol 59, 45-54. https://doi.org/10.1007/s10144-017-0571-8

Lent PC (1974) Mother-Infant relationships in ungulates. In: The Behaviour of Ungulates and

its relation to management. International Union for Conservation of Nature and Natural

Resources, Switzerland, pp 14–55.

Lindström J ( 1999) Early development and fitness in birds and mammals. TREE 14, 343–

348. https://doi.org/10.1016/s0169-5347(99)01639-0

Lindström J, Kokko H (2002) Cohort effects and population dynamics. Ecol Lett 5, 338–344.

https://doi.org/10.1046/j.1461-0248.2002.00317.x

Lloyd P, Rasa O (1989) Status, reproductive success and fitness in Cape mountain zebra

(*Equus zebra zebra*). Behav Ecol Sociobiol 25, 411–420.

https://doi.org/10.1007/bf00300187

Loison A, Langvatn R (1998) Short-and long-term effects of winter and spring weather on

growth and survival of red deer in Norway. Oecologia 116, 489–500.

https://doi.org/10.1007/s004420050614

Loveridge AJ, Searle AW, Murindagomo F, Macdonald DW (2007) The impact of sport-

hunting on the population dynamics of an African lion population in a protected area.

Biol Conserv 134, 548–558. https://doi.org/10.1016/j.biocon.2006.09.010

Mduma SAR, Sinclair ARE, Hilborn R (1999) Food regulates the Serengeti wildebeest: a 40-

year record. J Anim Ecol 68, 1101–1122. https://doi.org/10.1046/j.1365-

2656.1999.00352.x

Ncube H, Duncan P, Grange S, et al (2011) Pattern of faecal 20-oxopregnane and oestrogen

concentrations during pregnancy in wild plains zebra mares. Gen Comp Endocrinol 172,

358–362. https://doi.org/10.1016/j.ygcen.2011.03.027

Ogutu JO, Piepho HP, Dublin HT (2014) Responses of phenology, synchrony and fecundity

of breeding by African ungulates to interannual variation in rainfall. Wildl Res 40, 698–

717. https://doi.org/10.1071/wr13117

Owen-Smith N (1990) Demography of a large herbivore, the greater kudu *Tragelaphus*

*strepsiceros*, in relation to rainfall. J Anim Ecol 59, 893–913.

https://doi.org/10.2307/5021

Owen-Smith N, Ogutu JO (2013) Controls over reproductive phenology among ungulates:

allometry and tropical-temperate contrasts. Ecography 36, 256–263.

https://doi.org/10.1111/j.1600-0587.2012.00156.x

Penzhorn BL (1984) A long-term study of social organisation and behaviour of Cape

mountain zebras *Equus zebra zebra*. Z Tierpsychol 64, 97–146.

Pigeon G, Albon S, Loe LE, Bischof R, Bonenfant C, Forchhammer M, Irvine RJ, Ropstad E,

Veiberg V, Stien A (2022) Context-dependent fitness costs of reproduction despite

stable body mass costs in an Arctic herbivore. J Anim Ecol 91, 61–73.

https://doi.org/10.1111/1365-2656.13593

Plard F, Gaillard JM, Coulson T, Hewison AM, Douhard M, Klein F, Delorme D, Warnant C,

Bonenfant C (2015) The influence of birth date via body mass on individual fitness in a

long-lived mammal. Ecology 96, 1516–1528. https://doi.org/10.1890/14-0106.1

Pluháček J, Bartoš L, Doležalová M, Bartošová-Víchová J (2007) Sex of the foetus

determines the time of weaning of the previous offspring of captive plains zebra (*Equus*

*burchelli*). Appl Anim Behav Sci 105, 192–204.

https://doi.org/10.1016/j.applanim.2006.05.019

Post E (2003) Timing of reproduction in large mammals. In: Phenology: an integrative

environmental science. Springer, Dordrecht, pp 437–449

Pradel R, Hines JE, Lebreton JD, Nichols JD (1997) Capture-recapture survival models

taking account of transients. Biometrics 60–72. https://doi.org/10.2307/2533097

Reznick D, Nunney L, Tessier A (2000) Big houses, big cars, superfleas and the costs of

reproduction. TREE 15, 421–425. https://doi.org/10.1016/s0169-5347(00)01941-8

Rogers CM and others (1993) A woody vegetation survey of Hwange National Park. A

woody vegetation survey of Hwange National Park.. Department of National Parks and

Wildlife Management

Rubenstein DI, Nuñez CM (2009) Sociality and reproductive skew in horses and zebras. In:

Reproductive skew in vertebrates: proximate and ultimate causes. Cambridge University

Press Cambridge, pp 196–226

Sadleir R (1982) Energy consumption and subsequent partitioning in lactating black-tailed

deer. Can J Zool 60, 382–386. https://doi.org/10.1139/z82-051

Sæther BE (1997) Environmental stochasticity and population dynamics of large herbivores:

a search for mechanisms. TREE 12, 143–149. https://doi.org/10.1016/s0169-

5347(96)10068-9

Sinclair ARE, Mduma SAR, Arcese P (2000) What determines phenology and synchrony of

ungulate breeding in Serengeti? Ecology 81, 2100–2111. https://doi.org/10.1890/0012-

9658(2000)081[2100:wdpaso]2.0.co;2

Smuts GL (1976) Reproduction in the zebra mare *Equus burchelli antiquorum* from the

Kruger National Park. Koedoe 19, 89–132. https://doi.org/10.4102/koedoe.v19i1.1186

Smuts GL (1975) Pre- and postnatal growth phenomena of Burchell's zebra *Equus burchelli*

*antiquorum*. Koedoe 18, 69–102.

Stephens PA, Boyd IL, McNamara JM, Houston AI (2009) Capital breeding and income

breeding: their meaning, measurement, and worth. Ecology 90, 2057–2067.

https://doi.org/10.1890/08-1369.1

Tuljapurkar S (1989) An uncertain life: demography in random environments. Theor Popul

Biol 35, 227–294. https://doi.org/10.1016/0040-5809(89)90001-4

Valeix M, Loveridge AJ, Chamaillé-Jammes S, et al (2009) Behavioral adjustments of

African herbivores to predation risk by lions: spatiotemporal variations influence habitat

use. Ecology 90, 23–30. https://doi.org/10.1890/08-0606.1

Vitet C, Duncan P, Ganswindt A, Mabika C, Chamaillé-Jammes S (2021) Do infanticides

occur in harem-forming equids? A test with long-term sociodemographic data in wild

plains zebras. Anim Behav 177, 9–18. https://doi.org/10.1016/j.anbehav.2021.04.012

Vitet C, Duncan P, Gimenez O, Mabika C, Chamaillé-Jammes S (2020) Plains zebras bring

evidence that dilution and detection effects may not always matter behaviorally and

demographically. Ecosphere 11, e03288. https://doi.org/10.1002/ecs2.3288

Williams GC (1966) Adaptation and natural selection. Princeton University Press, Princeton,

USA.

# Tables

Table 1: Set of explanatory variables used in the models investigating survival in young individuals and mares plains zebra (*Equus quagga*), and transition probability between reproductive and non-reproductive states in mares plains zebra (Hwange National Park, Zimbabwe, 2008-2019).

| Variable | Type | Description | Used in the model(s) |
|---|---|---|---|
| *Season* | Categorical | "Wet" or "Dry" season | Young & Mares |
| *Time in dry season* | Continuous | Proportion of dry season between two consecutive sessions [range: 9-80 %] | Young & Mares |
| *Time* | Categorical | ID number of the field session [range: 1-24] | Young & Mares |
| *Trap-dependence* | Categorical | The individual has been seen during the previous field session (1) or not (0) | Young & Mares |
| *Age-class* | Categorical | "Younger foal" (less than six months old), "Older foal" (between six and 12 months old) or "Yearling" (between one and two years old) | Young |
| *Reproductive state* | Categorical | The mare is seen with a dependent juvenile (1) or not (0) | Mares |
| *Sight* | Categorical | The individual is seen for the first time (1) or not (0) | Mares |

**Figure legends**

Figure 1: Distribution of plains zebra (*Equus quagga*) births per month, estimated from the individual-based monitoring conducted in Hwange National Park, Zimbabwe, between 2008 and 2019 (*n* = 310 foals). The black line represents the total number of births per month over the complete study period (i.e. all years summed). The coloured areas indicate the seasons: blue = main wet season (approx. mid-December to mid-May), orange = main dry season (approx. August to October), mixed colour = inter-annual variability in the start of the seasons (start of the wet season: November to mid-December; start of the dry season: mid-May to July).

Figure 2: Probability of annual survival of plains zebra (*Equus quagga*) young individuals in Hwange National Park, Zimbabwe (2008-2019), according to the proportion of dry season between two consecutive sessions. Black: younger foals (between birth and six months old), blue: older foals (between six and 12 months old) combined with yearlings (between one and two years old). Solid lines represent predicted values from the best model. Shaded areas represent 85% confidence intervals of these predicted values. Dark coloured dots represent the survival predicted by the time model, shaded areas around the dot are proportional to 1/standard error of these predicted values.

Figure 3: a) Probability of annual survival of plains zebra (*Equus quagga*) mares in Hwange National Park, Zimbabwe (2008-2019), according to the proportion of dry season between two consecutive sessions and the reproductive state (for mares in their second and subsequent observations only, see text for details). Red: non-reproductive females, green: reproductive females. Solid lines represent predicted values from the best model. Shaded areas represent 85% confidence intervals of these predicted values. Dots represent females survival predicted

by the time model, vertical bars represent 85% confidence intervals of these predicted values.

b) Probability of transition between reproductive and non-reproductive states for plains zebra mares according to the session. Red: from reproductive to non-reproductive state, green: from non-reproductive to reproductive state. Dark coloured dots represent the survival predicted by the time model, shaded areas around the dot are proportional to 1/standard error of these predicted values.

# Figures

Figure 1:

Figure 2:

Figure 3: