Genetic and genomic architecture of species-specific cuticular hydrocarbon variation in parasitoid wasps

Jan Buellesbach[*], Henrietta Holze[*], Lukas Schrader[*], Jürgen Liebig[†], Thomas Schmitt[‡], Juergen Gadau[*] & Oliver Niehuis[§]

[*] Molecular Evolution and Sociobiology Group, Institute for Evolution & Biodiversity, University of Münster, Hüfferstr. 1, DE-48149 Münster, Germany

[†] School of Life Sciences, Arizona State University, P.O. Box 874701, Tempe, Arizona 85287-4501, USA

[‡] Department of Animal Ecology and Tropical Biology, Biocenter, University of Würzburg, Am Hubland, D-97074 Würzburg, Germany

[§] Department of Evolutionary Biology and Ecology, Institute of Biology I (Zoology), Albert Ludwig University of Freiburg, Hauptstr. 1, 79104 Freiburg, Germany

Running title: Wasp cuticular hydrocarbon genetics

Keywords: comparative genomics, gene mapping, ortholog inference, quantitative trait loci,

28 chemical ecology, entomology, *Drosophila*, *Nasonia*, Hymenoptera, Pteromalidae

30 **Corresponding author**: Jan Buellesbach, Institute for Evolution & Biodiversity (Room 221),

Hüfferstr. 1, DE-48149 Münster, Germany, Phone: +49 (0) 251-83-21637, Email:

32 buellesb@uni-muenster.de

ABSTRACT

46   Cuticular hydrocarbons (CHCs) serve two fundamental functions in insects: protection against

desiccation and chemical signaling. CHC profiles can consist of dozens of different compounds and

48   are considered a prime example for a complex trait. How the interaction of genes shapes CHC

profiles, which are essential for insect survival, adaptation, and reproductive success, is still poorly

50   understood. Here we investigate the genetic and genomic basis of CHC biosynthesis and

variation in parasitoid wasps of the genus *Nasonia*. Taking advantage of the wasps' haplo-

52   diploid sex determination and cross-species fertility, we mapped 91 quantitative trait loci

(QTL) explaining variation of a total of 43 CHCs in F_2 hybrid males from interspecific crosses

54   between three *Nasonia* species. To identify candidate genes, we localized orthologs of CHC

biosynthesis-related genes in the *Nasonia* genomes. By doing so, we discovered multiple

56   genomic regions where the location of QTL coincides with the location of CHC biosynthesis-

related candidate genes. Most conspicuously, on a region on chromosome 1 close to the

58   centromere, multiple CHC biosynthesis-related candidate genes co-localize with several QTL

explaining variation in methyl-branched alkanes. The genetic underpinnings behind this

60   compound class are not well understood so far, despite their high potential for encoding

chemical information as well as their prevalence in both *Nasonia* CHC profiles and many other

62   Hymenoptera. Our study considerably extends our knowledge on the so far little-known

genetic and genomic architecture governing biosynthesis and variation of this fundamental

64   compound class, establishing a model for methyl-branched alkane genetics in the

Hymenoptera in general.

INTRODUCTION

Understanding the genetic basis of quantitative phenotypic traits remains one of the central challenges in population genetics and evolutionary biology despite several advances in this field during the last decades (Yang et al. 2007; Mackay et al. 2009; Hartl 2020). Genes and regulatory nucleotide sequences governing natural variation in a set of phenotypic traits can be mapped (and potentially identified) in a genome by searching for phenotypic covariation with polymorphic genetic markers. This technique is commonly referred to as quantitative trait locus (QTL) mapping (Lander and Botstein 1989; Haley and Knott 1992; Mackay 2001a). Evolutionary genetic studies addressing variation and divergence of traits between closely related species with varying degrees of phylogenetic distance between them particularly benefit from QTL mapping techniques (Orr 2001; Slate 2005; Hartl 2020). Two components are of paramount importance for these kinds of studies: first, the availability of genetically tractable and crossable species with comparably short generation times and, ideally, established phylogenetic relationships (Mackay 2001b; Orr 2001; Mackay et al. 2009). Second, polygenic traits of interest that differ between the crossed species (Erickson et al. 2004; Slate 2005). Insects have proven to be particularly useful for QTL studies, since they combine various beneficial attributes: a multitude of quantifiable phenotypes, short generation times, large offspring numbers and, most commonly, relatively small genomes (Hoy 2005; Hunter and Kole 2008; Thomas et al. 2020). Cuticular hydrocarbons (CHCs), lipids constituting a major part of the waxy layer covering the insects' epicuticle, are particular promising traits to dissect genetically via QTL mapping, as they are comparatively easily quantifiable, polygenic, and phenotypically as well as genotypically traceable (Chung and Carroll 2015; Blomquist and Ginzel 2021; Holze et al. 2021).

CHCs are long-chained molecules whose primary function is to protect terrestrial

94 insects from desiccation (Ramsay 1935; Lockey 1980; Howard and Blomquist 1982, Fig. 1). As

versatile semiochemicals, CHCs have also been demonstrated to function in a wide array of

96 chemical communication systems integral to insect survival, reproduction, and adaptation

(Carlson et al. 1971; Blomquist and Bagnères 2010; Blomquist and Ginzel 2021). Differences

98 between CHC profiles arise from presence, absence, and abundance of individual CHCs, which

can differ in their chain length as well as in the presence and position of double bonds and

100 methyl groups (Lockey 1988; Howard and Blomquist 2005). The most commonly occurring

compound classes are saturated straight-chain alkanes (*n*-alkanes), unsaturated straight-chain

102 alkenes (*n*-alkenes) and alkadienes, and methyl-branched alkanes (Blomquist et al. 1987;

Blomquist and Bagnères 2010). Despite considerable diversity of CHC profiles across insects,

104 to the current stage of knowledge, the pathway of CHC biosynthesis appears to be mostly

conserved (Blomquist and Ginzel 2021; Holze et al. 2021). Briefly, the biosynthetic pathway

106 consists of the elongation of fatty-acyl-Coenzyme A units to produce very long-chain fatty

acids (VLCFs) that are subsequently converted to hydrocarbons by subducting the carboxyl

108 group (Nelson and Blomquist 1995; Howard and Blomquist 2005; Blomquist and Bagnères

2010, Fig. 1).

Most of what we currently know about the genetics governing CHC biosynthesis and

112 CHC profile diversity is based on studies on species of the Dipteran genus *Drosophila* (e.g.

Jallon 1984; Coyne and Oyama 1995; Ferveur 2005). This taxonomic bias has so far impeded a

114 more holistic view on the genetic underpinnings of this fundamental process across insect

orders (Blomquist and Bagnères 2010; Holze et al. 2021). Particularly little is known about the

Figure 1: Simplified overview of CHC biosynthesis, with the main enzymes catalyzing the intermediate reactions highlighted (compare to Fig. 3-5 for the localization of the corresponding genes). The pathway branches at different stages, eventually synthesizing the different CHC compound classes that are then transported to the cuticle. The main CHC compound classes are methyl-branched alkanes (mono-, di-, tri-, and tetra-), straight-chain alkanes, and unsaturated alkanes. Acetyl-CoA as the initial reactant of CHC biosynthesis is mainly provided by the citric acid cycle. Respective percentages of the individual compound classes found on the cuticle of *Nasonia* males (averaged over our three study species *N. vitripennis*, *N. longicornis* and *N. giraulti*) are indicated as well. Schematic insect drawing depicting an *N. vitripennis* male adapted from Ward's science, Rochester, NY, USA, 2002. Abbreviations: CoA: Coenzyme A, ACC: Acetyl-CoA carboxylase, FAS: Fatty acid synthase (m: microsomal, c: cytosolic, f: fat body associated). Adapted from Holze et al (2021).

116   genetic factors governing the differentiation and variation of methyl-branched alkanes, a

dominant compound class in many insect taxa with high potential for encoding chemical

118   information (Niehuis et al. 2011; Chung and Carroll 2015; Holze et al. 2021). In contrast to

Drosophila, where methyl-branched alkanes only comprise a small fraction of their CHC

120   profiles, this particular compound class dominates the CHC profiles in most species of the

insect order Hymenoptera (Martin and Drijfhout 2009; Kather and Martin 2015; Sprenger and

122   Menzel 2020). However, no single gene with a demonstrated impact on methyl-branched CHC

biosynthesis and variation has been unambiguously identified in Hymenoptera so far. To

124   address this knowledge gap, suitable model organisms are required to provide a basis for

studying the genetic background of CHC production, variation, and divergence.

    The parasitoid jewel wasp genus *Nasonia* (Hymenoptera: Pteromalidae) has emerged

128   as a model system well-suited to study quantitative as well as polygenic phenotypic traits in

the Hymenoptera. It combines ease of maintenance and handling, the possibility to generate

130   hybrids between its four described species, haplo-diploid genetics, and a growing genetic tool

kit (Gadau et al. 2008; Werren and Loehlin 2009; Werren et al. 2010). The *Nasonia* species

132   complex consists of four species, *N. vitripennis*, *N. longicornis*, *N. giraulti*, and *N. oneida*

(Darling and Werren 1990; Raychoudhury et al. 2010). Natural populations of *Nasonia* are all

134   infected with different strains of intracellular *Wolbachia* bacteria, leading to bidirectional

incompatibility in most interspecific crosses (Breeuwer and Werren 1990; Bordenstein and

136   Werren 1998; Bordenstein et al. 2001). However, the *Wolbachia* infections can be cured

(Richardson et al. 1987), rendering hybrids between all four species readily obtainable

138   (Breeuwer and Werren 1990; Bordenstein and Werren 1998; Raychoudhury et al. 2010). This

has proven to be particularly useful for QTL mapping of interspecific differences in a variety of

140 phenotypic traits (Gadau et al. 2002; Gadau et al. 2008; Niehuis et al. 2008). An exemplary

study comparing CHC variation through QTL mapping between two *Nasonia* species,

142 *N. vitripennis* and *N. giraulti*, shed some first light on the genetic basis of *Nasonia* CHC profile

differences (Niehuis et al. 2011). A study on CHC variation between all so far described

144 *Nasonia* species established clearly distinguishable species- and sex-specific CHC blends

(Buellesbach et al. 2013).

Here we dissect the genomic and genetic architecture of CHC variation in hybrid cross-

148 comparisons between three *Nasonia* species. We present results from conducting QTL

analyses on CHC variation in recombinant F_2 hybrid males obtained by crossing *N. vitripennis*,

150 *N. longicornis*, and *N. giraulti,* and from mapping CHC biosynthesis candidate genes. We

additionally screened for selection signatures considering candidate gene orthologs from two

152 more distantly related parasitoid wasp species within the family Pteromalidae: *Muscidifurax*

raptorellus and *Trichomalopsis sarcophagae* (Fig. 2). Finally, we discuss how CHC profile

154 divergence between the three *Nasonia* species can be explained phenotypically when

considering the newly gained knowledge on the genetic architecture of CHC profile differences

156 between them.

Figure 2: Phylogenetic relationship of the *Nasonia* species complex rooted by representatives of its two most closely related genera, *Trichomalopsis* and *Muscidifurax*, based on the D2 expansion region of their 28S rDNA sequences (Campbell et al. 2000) and mitochondrial DNA sequences (Werren et al. 2010). The three *Nasonia* species used in the current study are highlighted in grey.

MATERIALS AND METHODS

Nasonia strains and crossing experiments

166 We used the standard laboratory strains AsymCX, RV2x(U), and IV7(U) of the species *Nasonia*

vitripennis, *N. giraulti*, and *N. longicornis*, respectively, to conduct the cross experiments

168 (Breeuwer and Werren 1995; Werren et al. 2010). These strains are derived from wild-type

strains and have been antibiotically cured from their *Wolbachia* infection, enabling them to

170 hybridize under laboratory conditions. All strains were maintained in an incubator at 25 °C

under permanent light and were provided pupae of the flesh fly *Sarcophaga bullata*. Cross

172 experiments were conducted as described by Niehuis et al. (2008). We generated two types

of F1 hybrid females: 1) by crossing virgin *N. vitripennis* females with *N. longicornis* males and

174 2) by crossing virgin *N. giraulti* females with *N. longicornis* males. All F1 hybrid females were

kept unmated and thus generated recombinant (i.e., each genome constitutes a unique

176 combinations of the parental genomes) haploid F2 males. These F2 recombinant hybrid males

were collected 36 h after they had eclosed and were then freeze-killed and stored at -80 °C

178 before characterizing their genotypes and CHC phenotypes.

CHC profile analyses

182 We analyzed the CHC profiles of 100 *N. vitripennis* x *N. longicornis* (LV) and 101 *N. longicornis*

x *N. giraulti* (LG) recombinant hybrid F2 males. We additionally characterized the CHC profiles

184 of twelve *N. giraulti*, eleven *N. longicornis*, and 32 *N. vitripennis* males of the investigated

strains for comparison. CHCs were extracted by submersing individual wasps for 10 min in 10

186   µl n-hexane (≥ 99.0%; Sigma-Aldrich, St. Louis, MO, USA) using 1-ml glass vials equipped with

a 0.1-ml micro insert (Alltech, Deerfield, IL, USA). The CHC extracts were subsequently

188   transferred to new vials and concentrated under a stream of nitrogen to approximately 1 µl

total volume. The CHC extracts were then injected into a gas chromatograph coupled with a

190   mass spectrometer (GC: 6890N; MS: 5975C; Agilent, Santa Clara, CA, USA) for analysis. The

GC-MS was operated with in the splitless mode with an injector temperature of 250 °C.

192   Separation of compounds was performed on a J&W DB-5MS fused silica capillary column (30

m × 0.25 mm ID; df = 0.25 µm; Agilent, Santa Clara, CA, USA) and applying the following

194   temperature program: 60 °C start temperature, temperature increase by 40 °C per min to 200

°C, followed by an increase of 5 °C per min to 320 °C. Helium with a constant flow of 1 ml per

196   min was used as carrier gas. The MS-quad temperature was 150 °C, the MS-source

temperature was 230 °C, and the MSD-transfer line temperature was 315 °C. The obtained

198   chromatograms were analyzed with the software Enhanced Chemstation E.01.00.237 (Agilent,

Santa Clara, CA, USA). Peak identification was carried out automatically with the initial area

200   reject parameter set to 0, the initial peak width set to 0.043, and the initial threshold set to

14. Shoulder detection was turned off. CHC compounds were identified according to their

202   retention indices, diagnostic ions, and mass spectra (Carlson et al. 1998; Niehuis et al. 2011;

Buellesbach et al. 2013).

DNA extraction and QTL analysis

We extracted genomic DNA of all wasps after extraction of their CHCs. DNA extraction was

208   done as outlined by Niehuis et al. (2007). Genotype data were collected for 29 length-

polymorphic markers with known positions in the *Nasonia* nuclear genome (Tab. S1, Niehuis

et al. 2011). The species-specific length of the markers was assessed by separating PCR-generated amplicons of them on a denaturing polyacrylamide gel following the procedure given by Niehuis et al. (2008). We used the following PCR temperature program for obtaining the amplicon: 5 min. at 95 °C, followed by 30 cycles of 1 min. at 95 °C, 1 min. at 55 °C, and 1 min. at 72 °C, and followed by 10 min. at 72 °C. PCR oligonucleotide primer sequences with their respective amplicon lengths are listed in Tab. S1.

Quantitative trait loci (QTL) analyses were conducted with the R/qtl package version 1.14-2 (Broman et al. 2003; Broman and Sen 2009) on a 64-bit-build of R version 2.10.1 (R Development Core Team 2018). Significant phenotypic differences in the relative abundances of CHC compounds were assessed with Bonferroni-corrected Wilcoxon rank-sum tests with continuity correction (Hollander and Wolfe 1999; Zar 1999). Each CHC compound was subjected to a one-dimensional one-QTL and to a two-dimensional two-QTL scan using Haley-Knott regression (Haley and Knott 1992) with an assumed genotypic error probability of 0.001 and a step width of 1 cM. We additionally applied a multiple QTL model to each CHC trait with forward/backward model selection as implemented in R/qtl. The maximum number of QTL for each trait was set to 10, with QTL positions refined after each step of the forward and backward selection. Penalties for model selection were set based on the permutation from the two-dimensional two-QTL genome scan of the respective CHC trait with a significance level of 0.01 (Manichaikul et al. 2009). Significance thresholds for QTL presence were estimated from 10,000 permutations of each respective phenotypic CHC abundance. Since our QTL mapping was based on the *Nasonia* linkage map inferred by (Niehuis et al. 2010) and on genome assembly 1.0 (Nvit 1.0, Werren et al. 2010), we transferred marker positions and QTL to the most recent high-resolution linkage map of *Nasonia* (Nvit 2.1, Desjardins et al. 2013) where we performed the mapping of candidate CHC biosynthesis genes (Tab. S1, see below).

234 Though the position of several markers had slightly shifted on the new linkage map, their

respective initial linkage groups remained the same, enabling the re-assessment of the QTL

236 positions by their relative position between the two closest markers. A fixed confidence

interval of 20 cM was chosen for each QTL, since this distance was equivalent to the average

238 distance between markers.

Search for candidate CHC biosynthesis genes

242 Thirty-eight well-characterized candidate genes with a demonstrated impact on CHC variation

via targeted knockdown studies were selected from *Drosophila melanogaster* (Tab. S2, see

244 also Holze et al. 2021) via FlyBase, Version FB2019_02 (http://flybase.org/, Gramates et al.

2017). To screen for orthologs of these candidate genes in the *Nasonia* Nvit 2.0 reference

246 genome, three complementary methods were used: first, the amino-acid sequences, including

all possible isoforms encoded by *D. melanogaster* CHC biosynthesis genes of interest (GOI),

248 were searched against the *N. vitripennis* proteome available on NCBI (*N. vitripennis*,

https://www.ncbi.nlm.nih.gov/) using the BLAST+ software suite (version 2.8.1, Camacho et

250 al. 2009). Amino-acid sequences with an e-value smaller than 10^{-10} were then again searched

reciprocally against the *D. melanogaster* proteome. Reciprocal best hits were considered as

252 orthologs. Second, the program OrthoFinder (version 2.3.1, Emms and Kelly 2015) was used

to identify ortholog groups at the hierarchical level of the last common ancestor of the

254 investigated species. OrthoFinder generated a) a list of ortholog groups that contained one

gene from each of the tested species only (single-copy genes) and b) rooted phylogenetic trees

256 of genes that belong to ortholog groups with paralogous gene copies. The software DIAMOND

(Buchfink et al. 2014) was used to align the amino-acid sequences of a given ortholog group.

Third, we queried the WaspAtlas database (http://cyverse.warwick.ac.uk:3000/, Davies and Tauber 2015) for ortholog groups (Flicek et al. 2014).

We were unable to identify unambiguous orthologs of some of the *D. melanogaster* CHC biosynthesis candidate genes in the *N. vitripennis* genome with the above-outlined methods

despite partially high sequence similarities. Therefore, BLAST+-inferred hits in the *N. vitripennis* proteome with amino-acid sequence similarity thresholds of >-30-% identity, <-10-

^{10} e-value, and >-85-% query coverage to *D. melanogaster* were additionally considered, although these may not necessarily constitute direct orthologs. The identification of

orthologous candidate proteins of two protein families posed particular problems: fatty acid elongases (ELOs) and fatty-acyl-CoA reductases (FARs). Candidate ELOs and FARs in the *N.*

vitripennis proteome with the highest amino-acid sequence similarity (lowest e-value and highest bit score) to the *D. melanogaster* ELOs and FARs with demonstrated impact on CHC

variation (Tab. S2) were used as queries to search for homologs in the *Nasonia* proteome using blastp of the BLAST+ software suite. Hits above a similarity threshold of e-value < 10^{-20} were

considered to belong to the same protein/gene family, and domain structure of the respective hits was taken into account to ensure that the derived gene sequences encode functional

proteins. This was achieved with hmmscan of the HMMER software (http://hmmer.org/, version 3.1b2, February 2015) and use of the Pfam-A database (version 32, El-Gebali et al.

2019).

    For visualization of the QTL regions and candidate gene positions on the *N. vitripennis* linkage

278   map Nvit 2.1, the R package LinkageMapView was used (Ouellette et al. 2017).

CHC profile discriminant analysis

Discriminant analysis (DA) was performed with the R package "MASS" (Venables and Ripley 2002) to test whether CHC profiles statistically differ between the six investigated groups of male wasps (i.e., those of the parental strains and those of the F_2 hybrids). To standardize the peak area values for DA, the normalization method of the function "decostand" of the community ecology R package "vegan" was used based on the following formula (Dixon 2003):

$$T_{x,y} = P_{x,y} / \sqrt{\Sigma P_y^2}$$

$T_{x,y}$ is the transformed peak area x of individual y, $P_{x,y}$ is the absolute peak area x of individual y and ΣP_y^2 to the squared sums of all absolute peak areas of individual y. This widely applied method for normalizing phenotypic data was chosen to make the peak areas comparable between our groups and to highlight the relative peak area differences. To visualize the data by plotting the first three discriminant functions simultaneously, the R package "scatterplot3d" was used (Ligges and Maechler 2003). Wilk's λ was calculated to measure the quality of the DA.

Gene orthology and screens for signatures of positive selection

To test for signatures of positive selection in our candidate genes, we extended our ortholog inference analysis to the fourth *Nasonia* species, *N. oneida*, as well as two closely related parasitoid wasp genera, *Trichomalopsis sarcophagae* and *Muscidifurax raptorellus*. The nucleotide sequences of the candidate genes from the *N. vitripennis* genome were extracted with SAMtools (Li et al. 2009) and aligned with the software BLAT (v35, Kent 2002; Bhagwat

306   et al. 2012) to the nucleotide sequences of the latest genome assemblies of *N. giraulti*, *N.*

longicornis, *N. oneida*, and *T. sarcophagae* kindly provided by Xiaozhu Wang (unpublished

308   data). Aligned nucleotide sequences that shared at least 50 identical nucleotide positions and

were aligned along at least 80 % query nucleotide sequence length were considered as

310   homologous. To ensure these were also orthologous to the candidate genes of the

N. vitripennis genome, their amino-acid sequences were reciprocally searched against the

312   *N. vitripennis* proteome with BLAST+ (version 2.8.1, Camacho et al. 2009). Only genes that

fulfilled the best reciprocal hit criterion were considered as orthologous. To annotate genes

314   in the draft genomes of *N. giraulti*, *N. oneida*, *N. longicornis*, and *T. sarcophagae*, we used the

program GeMoMa (version 1.6, Keilwagen et al. 2016). The reliability of the annotation by the

316   program can be improved by providing it the coding sequence of an orthologous gene if a

phylogenetically conserved gene structure can be assumed. We applied this approach for

318   structurally annotating candidate genes in the *N. giraulti*, *N. oneida*, *N. longicornis*, and

T. sarcophagae genomes, using nucleotide sequences of the orthologous *N. vitripennis* genes

320   as references. Since BLAT was not sensitive enough to identify orthologs in *M. raptorellus*

(comparatively distantly related to *N. vitripennis*, see Fig. 2), we *de novo*-inferred gene models

322   in a draft genome of this species, kindly provided by Eva Jongepier (unpublished data), with

the software GeMoMa. To obtain the respective coding nucleotide sequences from the

324   GeMoMa-predicted annotations, the program BEDtools (Version 2.28.0, Quinlan and Hall

2010) with either the nucleotide sequences of the gene regions or the whole genome

326   assembly.

Candidate genes were screened for signatures of positive selection by analyzing the ratio of

328   nonsynonymous to synonymous substitutions (ω = dN/dS). The nucleotide sequences of each

candidate gene ortholog group containing the orthologs of all six wasp species were aligned

330 with the codon-sensitive alignment tool PRANK (version 170427, Löytynoja and Goldman

2008; Löytynoja 2014). The alignment process was supported with a phylogenetic tree (Fig. 2)

332 of the study species whose evolutionary distances are based on the D2 expansion region of

their 28S rDNA sequences (Campbell et al. 2000) and mitochondrial DNA sequences (Werren

334 et al. 2010). Poorly aligned regions were trimmed with Gblocks (version 0.91b, Castresana

2000). Each sequence alignment was screened for signatures of positive selection on at least

336 one site on at least one branch of the given phylogeny with the BUSTED (**B**ranch-**S**ite

Unrestricted **S**tatistical **T**est for **E**pisodic **D**iversification) algorithm (Murrell et al. 2015),

338 implemented in HyPhy (version 2.5, http://hyphy.org). Resulting p values were Benjamini-

Hochberg-corrected for multiple testing (Benjamini and Hochberg 1995).

RESULTS

QTL explaining CHC variation in Nasonia *F2 hybrid males*

346 We detected a total of 91 QTL explaining CHC variation in recombinant F_2 hybrid males. Of

those, 80 were found in the recombinant F2 hybrid males obtained from crossing

348 *N. longicornis* (♂) and *N. vitripennis* (♀) and eleven were found in the recombinant F2 hybrid

males obtained from crossing *N. longicornis* (♂) and *N. giraulti* (♀) (Fig. 3 and 4, Tab. 1). The

350 detected QTL explain variation in 43 (of a total of 53 analyzed) CHCs. The QTL of seven CHCs

(two mono-methyl-branched, three di-methyl-branched, and two tetra-methyl-branched

352 alkanes) were found in hybrids from both crosses, but corresponding QTL are located on

Figure 3: Linkage map based on F2 hybrid males from a cross between *Nasonia longicornis* (♂) and *Nasonia vitripennis* (♀) depicting positions of 80 quantitative trait loci (QTL) for 42 individual cuticular hydrocarbons (CHCs). Additionally, the positions of CHC biosynthesis-related candidate genes are shown. CHCs are indicated by symbols corresponding to their respective compound class (alkanes, alkenes, mono-, di-, tri-, and tetra-methyl-alkanes), their retention indices (RI) and numbers inside the symbols reflecting their carbon chain length. The five Nasonia chromosomes are labeled 1–5. Black horizontal lines on the chromosomes depict the respective positions of the 29 genotyped molecular markers used for determining the QTL positions. Orthologs to genes with a clear impact on CHC production derived from *Drosophila melanogaster* are color coded according to their putative position in the CHC biosynthesis pathway where possible (see Fig. 1). Regions of low recombination (> 0.8 Mb / cM) and centromeric regions are derived from Desjardins et al. (2013) and marked in light and dark grey, respectively.

Figure 4: Linkage map based on F2 hybrid males from a cross between *Nasonia longicornis* (♂) and *Nasonia giraulti* (♀) depicting positions of 11 quantitative trait loci (QTL) for 9 individual cuticular hydrocarbons (CHCs). Additionally, the positions of CHC biosynthesis-related candidate genes are shown. QTL could only be detected for the variation of methyl-branched alkanes (mono-, di-, and tetra-methyl-alkanes). Labels of chromosomes, candidate genes, molecular markers and regions of low recombination are indicated as in Fig. 3

different chromosomes or on different chromosomal regions (Tab. 1). The QTL of one single

CHC compound, 15,17-DiMeC29 (RI: 2982), was detected exclusively in hybrids that originated

from crossing *N. longicornis* and *N. giraulti* (Fig. 4 and Tab. 1). Sixty-nine QTL, explaining

variation of 35 CHCs, were exclusively detected in hybrids that originated from crossing

N. longicornis and *N. vitripennis* (Fig. 3 and Tab. 1). The QTL detected in F2 hybrid males of the

latter cross are spread across all five chromosomes and encompass all six compound classes

identified in *Nasonia* CHC profiles (i.e., alkanes, alkenes, mono-, di-, tri-, and tetra-methyl-

branched alkanes). Several QTL explaining quantitative variation of structurally similar CHCs

(mostly methyl-branched alkanes) clustered together, with two clusters each on

chromosomes 1 and 2, and one cluster each on chromosomes 3, 4, and 5 (Fig. 3 and Table 1).

Eleven QTL were found in recombinant F2 hybrid males obtained from crossing *N. longicornis*

and *N. giraulti* (Fig. 2). All of them explain variation of methyl-branched alkanes (mono-, di-,

and tetra-methy-alkanes). These QTL are located on chromosomes 2, 4, and 5, with clusters

of them detected on chromosomes 4 and 5 (Fig. 4). Although these QTL clusters are in spatial

proximity to the ones detected on the same two chromosomes in the F2 male hybrids

obtained from crossing *N. longicornis* and *N. vitripennis*, they explain variation of different

CHCs (compare Fig. 3 and 4).

CHC biosynthesis-related candidate genes

Of the candidate genes from *Drosophila melanogaster* with a demonstrated impact on CHC

variation through targeted knockdown studies (Table S2), we found 15 to be represented by

exactly one ortholog in the *Nasonia* reference genome, irrespective of the applied ortholog

inference method (i.e., reciprocal BLAST search, OrthoFinder, and WaspAtlas) (Table 2).

Ortholog inference of the remaining 15 *D. melanogaster* candidate genes generated

inconsistent results when comparing the three ortholog inference methods. For instance,

reciprocal BLAST search (RBS) of three fatty acid synthase (FAS) genes with a demonstrated

impact on *Drosophila* CHC biosynthesis identified nine putative *fas* orthologs in the *Nasonia*

reference genome, whereas OrthoFinder found five. However, upon closer inspection, several

of the putative *fas* orthologs identified by RBS are actually comprised of very short nucleotide

sequences that unlikely encode fully functional proteins. These potential pseudogenes were

excluded as CHC biosynthesis candidate genes in *Nasonia*. Furthermore, of all three ortholog

inference methods, WaspAtlas was the only tool that led to the identification of an *N.*

vitripennis ortholog to the *D. melanogaster* gene *cyp4g1*. However, due to a complimentary

identification of an orthologous member of the *Cyp4g* gene family in the *N. vitripennis* genome

with the highest nucleotide sequence similarity to *cyp4g1* by Qiu et al (2012), this ortholog

was still included in the list of potential CHC biosynthesis candidate genes.

Two further gene families emerged in our analysis with high nucleotide sequence similarity to

Drosophila CHC biosynthesis candidate genes despite no evidence for direct orthology:

elongases (*elos*) and *fatty-acyl-CoA reductases* (*fars*). Of the four *elos* and the two *fars* in the

D. melanogaster genome with a clear impact on CHC profile composition, we were able to

identify twelve and 22 homologous genes with high nucleotide sequence similarity in the

Nasonia reference genome (Tab. 3 and Fig. 5). Mapping of those genes on the *Nasonia* linkage

map revealed a region on chromosome 1 around the centromere where eight *elos* and

thirteen *fars* cluster together in close spatial vicinity (Fig. 5). Two additional candidate genes,

orthologous to the *Drosophila* CHC biosynthesis genes *cyp4g1* and *CG14688*, also map in this

chromosomal region.

Figure 5: *Nasonia* linkage map of chromosome 1 with an enlarged region densely packed with CHC biosynthesis related candidate genes. Eight elongase (elo) and thirteen fatty-acyl-reductase (far) cluster together in close spatial vicinity in addition to two additional genes genes cyp4g1 and CG14688 (see Tab. 2 and 3) in this chromosomal region. Note that this region maps around the centromere and is characterized by a particularly low recombination rate (> 0.8 Mb / cM). Labels of chromosomes and molecular markers are indicated as in Fig. 3 and 4.

Correlation in the genomic localization of candidate genes and QTL

QTL from the second cluster on chromosome 1 explaining variation of CHCs in recombinant F_2 male hybrids of *N. longicornis* and *N. vitripennis* co-localize with CHC biosynthesis-related candidate genes in close spatial vicinity to the centromere (Fig. 3). Intriguingly, this also encompasses the conspicuous chromosomal region with a high density of *elo* and *far* genes (Fig. 5). Of the two detected QTL clusters on chromosome 2 from the cross mentioned above, the first one co-localizes with an ortholog of the candidate gene *spidey* that codes for a component of the elongase enzyme complex (Wicker-Thomas et al. 2015; Holze et al. 2021, see Fig. 1 and 3). Some QTL from the second cluster co-localize with several *far* genes (compare Fig. 3 and Fig. 5). On chromosome 3, the detected QTL cluster maps closely to the position of a gene orthologous to *CG16979*, a candidate gene with an impact on CHC variation in *D. melanogaster* despite an unclear role in the CHC biosynthesis pathway (Dembeck et al. 2015; Holze et al. 2021). On chromosome 4, several QTL map closely to a *hacd1*-ortholog, another gene that codes for an elongase enzyme complex component (Wicker-Thomas et al. 2015; Holze et al. 2021). On chromosome 5, a particularly large cluster, consisting almost entirely of QTL explaining variation in methyl-branched alkanes, co-localizes with a *fas* gene ortholog (*fas5*). Another QTL cluster identified in male F_2 hybrids from the cross between *N. longicornis* and *N. giraulti* on chromosome 4 maps close to the *hacd1* ortholog, in close proximity to the above-mentioned QTL cluster from the cross between *N. vitripennis* and *N. longicornis* on the same chromosome (compare Fig. 3 and 4).

CHC divergence between F_0 parental and F_2 hybrid males

426   To assess whether CHC profiles statistically differ between males of the parental strains and

the recombinant F_2 hybrid males, we conducted a discriminant analysis (DA) on the CHC profile

428   data of 256 male wasps belonging to five groups: *N. giraulti* (NG)*, N. longicornis* (NL),

N. vitripennis (NV), *N. longicornis* x *N. vitripennis* F2 hybrids (LV), and *N. longicornis* x *N. giraulti*

430   F2 hybrids (LG) (Fig. 6). All groups differed in their CHC profiles significantly from each other

(Wilk's $\lambda < 0.001$, $\chi^2 = 23.22$, $p < 0.001$), and the recombinant F_2 males consistently clustered

432   in between their respective F_0 parental males.

Orthology and signatures of positive selection for CHC candidate genes in other

436   *parasitoid wasps*

We screened all 26 CHC biosynthesis-related candidate genes identified in the *N. vitripennis*

438   reference genome (see Tab. 2) for orthologs in the respective genomes of the other three

Nasonia species and in the genomes of two additional pteromalid wasp species: *Muscidifurax*

440   *raptorellus* and *Trichomalopsis sarcophagae* (Fig. 2). In the genome of the latter, we were

unable to identify orthologs of seven of the 26 *N. vitripennis* CHC biosynthesis candidate genes

442   (*CG14688, fas2, fas4, fas5, hacd2, nrt*). Of these seven, two (*fas5* and *hacd2*) remained

unidentified in *T. sarcophagae* and three remained unidentified in *N. giraulti* (*fatp1, nrt,*

444   *hacd2*). Interestingly, *hacd2* was the only *N. vitripennis*-exclusive CHC biosynthesis-related

candidate gene with no orthologs in any of the other investigated parasitoid wasp species

446   (Tab. 4). Orthologs of the remaining 19 CHC biosynthesis-related candidate genes were found

in all investigated species and were screened for signatures of positive selection based on the

◯	F$_0$ *N. giraulti* males (NG)
⊗	F$_2$ *N. longicornis* / *N. giraulti* hybrids (LG)
✕	F$_0$ *N. longicornis* males (NL)
⊠	F$_2$ *N. longicornis* / *N. vitripennis* hybrids (LV)
☐	F$_0$ *N. vitripennis* males (NV)

Figure 6: Plot of the first three discriminant functions showing the divergence between CHC profiles of F0 parental and F2 hybrid males between the three investigated *Nasonia* species in three dimensions. F0 NG: *N. giraulti* males (n = 12), F0 NL: *N. longicornis* males (n = 11), F0 NV: *N. vitripennis* males (n = 32), F2 LV: *N. longicornis* / *N. vitripennis* hybrid males (n = 101), F2 LG: *N. longicornis* / *N. giraulti* hybrid males (n = 100), total n = 256. All groups were significantly differentiated from each other (Wilk's λ < 0.001, χ2 =23.22, p < 0.001), the variation each function explains is indicated in percentages.

ratios of non-synonymous (dN) to synonymous (dS) nucleotide substitutions. After Benjamini-Hochberg correction for multiple testing, the signature for positive selection in one gene, *fatty acid synthase 1* (*fas1*), was considered statistically significant.

DISCUSSION

Genomic regions harboring candidate genes and QTL explaining CHC variation

By analyzing recombinant F2 hybrid males from crosses between *N. longicornis* and *N. vitripennis* (LV) and between *N. longicornis* x *N. giraulti* (LG), we identified several QTL clusters mapping to genomic regions that harbor CHC biosynthesis candidate genes. Four of these regions additionally coincide with the location of QTL explaining CHC variation in recombinant F2 hybrid males of *N. vitripennis* and *N. giraulti* (VG, Niehuis et al. 2011, Fig. S1 and Tab. S3). The coinciding QTL of the latter study additionally explain variation in six CHCs (i.e., 7-MeC30, 5-MeC30, 9-C31ene, 3,7-DiMeC31, 3,25-DiMeC31, 3-MeC32, 15-; 17-; 19-MeC37) that we found no QTL of in the present investigation. Interestingly, QTL explaining variation of identical compounds in different hybrid crosses map only in very few instances to the same genomic regions (e.g., QTL for 11,15-; 11,23- 11,25-DiMeC33 on Chr. 5 in LV and in VG crosses, QTL for 7,19-; 7,21-; 7,23-DiMeC37 on Chr. 2 in LV and in LG crosses). Conversely, in the majority of cases, QTL explaining variation of the same CHC compounds map to different chromosomal regions, or to different chromosomes entirely (Fig. 3, 4 and S1). This illustrates

the genetic complexity and potential pleiotropy of the governance of most investigated CHC

472   compounds. However, specifically comparing our cross between *N. vitripennis* and

N. longicornis with the previously analyzed cross between *N. vitripennis* and *N. giraulti*, we

474   identified overlapping QTL clusters governing the variation of structurally similar CHC

compounds. The three most obvious of these clusters map close to the centromeric regions

476   of chromosomes 1, 2, and 5 (compare Fig. 3 and S1, Tab. S3). The respective CHC compounds

are all methyl-branched alkanes, partially sharing methyl-branching patterns and chain

478   lengths (Tab. 1). Intriguingly, all of the above-mentioned QTL clusters co-localize with

orthologs of candidate genes implicated with CHC biosynthesis. Particularly, the shared cluster

480   on chromosome 1 mapped to the identified chromosomal region with the highest density of

orthologs of CHC biosynthesis candidate genes (compare Fig. 3 and Fig. 5). As the

482   recombination rate around the centromeres is very low, the close proximity of these gene

orthologs is concordant with a higher likelihood of passing the whole unaltered gene cluster

484   to the next generation, rendering gene shuffling in this region rare (Niehuis et al. 2010). The

other two clusters co-localize with the positions of the gene orthologs *spidey* and *fas5* on

486   chromosome 2 and 5, respectively. *Spidey* encodes a 3-hydroxy-acyl-CoA-dehydratase,

constituting a part of the elongase enzyme complex (Wicker-Thomas et al. 2015), whereas

488   *fas5* encodes a fatty acid synthase, instrumental for the formation of long-chain fatty acids as

precursors for the elongation process (Juarez et al. 1992; Gu et al. 1997, see Fig. 2). These

490   findings render these three distinct chromosomal regions particularly promising targets for

further investigating the impact of the candidate genes mapped in these regions on CHC

492   biosynthesis.

CHC compound class variation and gene orthology between Nasonia and

496 *Drosophila*

The vast majority of QTL that we detected in the hybrids from both crosses explained variation

498 of methyl-branched alkanes (79 of 91 in total). This compound class dominates in CHC profiles

of *Nasonia* wasp males and accounts for 87 % of the total amount of CHCs when averaged

500 across all three investigated species (Fig. 1 and Tab. S4). This result is in accordance with those

of most studies on CHC profiles in Hymenoptera, consistently reporting methyl-branched

502 alkanes as the dominant CHC class (e.g. Martin and Drijfhout 2009; Kather and Martin 2015;

Sprenger and Menzel 2020). In species of other insect orders, however, methyl-branched

504 alkanes seem to be less dominant, which is particularly true for *Drosophila melanogaster*:

analyzing CHC profiles in lines of the *D. melanogaster* Reference Panel (DGRP), Dembeck et al.

506 (2015) detected only mono-methyl-branched alkanes in *D. melanogaster* CHC profiles, and

these account for 16.2 % and 24.1 % of the total CHCs in CHC profiles of males and females,

508 respectively (Tab. S5). This might be one of the reasons why our understanding of the genetics

governing the variation in methyl-branched alkanes is comparatively limited.

510 The split between Hymenoptera and other holometabolous insects, including Diptera, is

estimated to have occurred around 327 Mya (Misof et al. 2014). Thus, it is not surprising that

512 were we unable to reliably identify orthologs of several genes with an impact on CHC profile

composition in *Drosophila* in the genome of *Nasonia* wasps. For instance, the potentially best

514 studied CHC biosynthesis-related genes in *Drosophila* are desaturases, catalyzing the

introduction of double bounds in hydrocarbon chains (e.g. Wicker-Thomas et al. 1997; Coyne

516 et al. 1999; Dallerac et al. 2000, Fig. 1). Although several *desaturase* genes have been

identified in the *Nasonia* genome (Niehuis et al. 2011), we were not able to clearly confirm

518 their orthology to *Drosophila* desaturases in the present study. However, the proportions of

unsaturated CHCs vastly differ between the CHC profiles of *Drosophila* flies and those of

520 *Nasonia* wasps (on average 59.2 % vs. 1.4 % in males, Tab. S4 and S5). Thus, since the

proportion of unsaturated compounds is considerably less in *Nasonia* CHC profiles (compared

522 to CHC profiles of *Drosophila* flies), the main genes governing their biosynthesis and variation

might have deviated considerably after more than 300 Mya of estimated evolutionary

524 divergence between *Nasonia* and *Drosophila* (Misof et al. 2014).

Similar in their unresolved orthology to *Drosophila*, we found numerous *elo* and *far* genes

526 across the *Nasonia* genome (Tab. 3). In *D. melanogaster*, 19 *elo* genes have been identified in

total (Chung and Carroll 2015; Zuo et al. 2018), but only four have been clearly associated with

528 CHC biosynthesis and variation (Chertemps et al. 2007; Dembeck et al. 2015). Concerning *far*

genes, from 17 identified in the *D. melanogster* genome (Finet et al. 2019), two have a

530 demonstrated impact on CHC biosynthesis (Dembeck et al. 2015). This already indicates the

difficulty in associating members in these two large and diverse gene families with CHC

532 biosynthesis and variation, as they can also be involved in vastly different processes, including

tracheal gas exchange, reproductive tissue growth, and larval development (Jaspers et al.

534 2014; Cinnamon et al. 2016). However, there are two further properties of *Nasonia* CHC

profiles that greatly differentiate them from their *Drosophila* counterparts: first, in *Drosophila*,

536 CHC chain lengths apparently do not exceed beyond C31 (Dembeck et al. 2015), whereas in

Nasonia, CHC compounds with chain lengths of up to C37 have been identified, and additional

538 CHCs with chain length of up to C52 have recently been described (Bien et al. 2019). Since

elongases contribute to the elongation of long chain fatty acids (see Fig. 1), the longer and

540 more diverse chain lengths in the *Nasonia* profiles could mean that a larger and more diverse

set of *elongase* genes is functionally involved in CHC biosynthesis in *Nasonia* wasps. Second,

542 *Nasonia* wasp CHC profiles appear to be more complex, generally containing more compound

classes, including di-, tri-, tetra-methyl-branched alkanes, in contrast to CHC profiles of

544 *D. melanogaster* (compare Tab. S4 and S5). It has been demonstrated that the *far* gene family

shows particularly quick evolutionary turnover rates, which allows, in turn, for rapid

546 diversification of CHC profiles between *Nasonia* species (Finet et al. 2019). This also argues for

the involvement of a larger set of *far* genes in CHC biosynthesis in *Nasonia* than in *Drosophila*

548 due to the more diverse and complex nature of *Nasonia* CHC profiles. Hence, the region on

chromosome 1 containing eight *elo* and eleven *far* genes while harboring QTL mostly

550 explaining methyl-branched alkane variation in two of our analyzed hybrid crosses constitutes

a promising target-rich region for future functional genetic studies. *Nasonia* might prove to be

552 a particularly well-suited model system for investigating the genetic underpinnings of methyl-

branched alkane biosynthesis and diversification, with our current study providing a

554 comprehensive foundation.

Phylogenetic divergence in relation to CHC differentiation and selection

558 ***signatures***

It has been estimated that *N. vitripennis* has diverged around 1 Mya from the lineage from

560 which the remaining *Nasonia* species evolved (Campbell et al. 1993). The divergences of the

remaining *Nasonia* lineages occurred approximately between 500 000 and 400 000 years ago

562 (Werren et al. 2010, Fig. 2). We discovered the smallest number of QTL governing CHC

variation between the more recently diverged species pair studied by us: *N. longicornis* and

564 *N. giraulti*. This finding matches well with the considerably shorter evolutionary time frame to

accumulate genetic differences compared to the longer divergence time between each of the

566 two species and *N. vitripennis*. Therefore, it is all the more surprising that phenotypically, the

overall CHC profile divergence sufficiently discriminates all species with F_2 hybrid males

568   clustering as intermediate phenotypes between the respective F_0 males (Fig. 6). As

N. longicornis and *N. giraulti* males also differ considerable in their CHC composition from

570   each other, this obvious phenotypic divergence is not reflected in the comparably low

genomic divergence between these two species hinted at by our QTL comparison. A potential

572   explanation for this might be other mechanisms, *e.g.*, phenotypic plasticity (Otte et al. 2018)

or epigenetic regulation (Morandin et al. 2019), that factor into CHC profile divergence

574   between these phylogenetically less distant species.

Concerning CHC biosynthesis candidate gene orthology, we extended our analysis beyond the

576   genus *Nasonia* to two more distantly related parasitoid wasp species in the family

Pteromalidae, *Trichomalopsis sarcophagae and Muscidifurax raptorellus* (Fig. 2).

578   Unsurprisingly, in the genome of the latter, most distantly related species, we detected the

fewest orthologs in comparison to the *N. vitripennis* reference genome (19 out of 26, Tab. 4).

580   Furthermore, taking into account all shared 19 candidate genes in our six investigated wasp

genomes, we detected signatures of positive selection in only one of them, *fas1*. The gene

582   *fas1*, an ortholog to *FASN1* and *FASN2* in *Drosophila*, plays an important role in the early steps

of CHC biosynthesis and impacts the abundance of methyl-branched alkanes (Chung et al.

584   2014; Wicker-Thomas et al. 2015). Additionally, *fas1* maps in close proximity to the

conspicuous cluster of CHC biosynthesis gene orthologs (Fig. 5) that also co-localizes with QTL

586   explaining variation of mostly methyl-branched alkanes between *N. vitripennis* and *N. giraulti*

and between *N. longicornis* and *N. giraulti* (compare Fig. 3 and S1). This renders *fas1* a

588   particularly promising candidate gene for further functional genetic studies investigating its

impact on CHC profiles not only in *Nasonia*, but also in other Hymenoptera. Generally,

590   establishing a functional link between the identified candidate genes and CHC variation will

be instrumental in further elucidating the genomic and genetic architecture underlying

species-specific CHC diversification in parasitoid wasps in particular and other hymenopteran

species in general.

Conclusions and Outlook

With our study on the genetic and genomic architecture of CHC variation between parasitoid

wasps of the genus *Nasonia*, we uncovered several genomic regions where QTL explaining

variation in similar CHC compounds and CHC biosynthesis-related candidate genes clustered.

Especially one region on chromosome 1 close to the centromere is conspicuous in this regard,

as it harbors many CHC biosynthesis-related candidate genes and QTL explaining variation of

methyl-branched alkanes, the prevalent CHC compound class in the Hymenoptera in general

and in *Nasonia* in particular. This underlines the considerable potential of *Nasonia* as a model

system to further investigate the little-know genetic and genomic architecture of methyl-

branched alkane biosynthesis and variation.

ACKNOWLEDGEMENTS

Thanks to Andrea K. Judson, Nadine Brehm and Sylvia Geeritsma for their help in acquiring the

chemical and QTL data, Joshua D. Gibson for his valuable input in conceptualizing the

manuscript in its current form, and Valerio Vitali for helpful suggestions on the first draft of

the manuscript.

FUNDING

This research was partially funded by the Deutsche Forschungsgemeinschaft (DFG, German
Research Foundation) – 427879779; 403980864.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

All data underlying the presented study will be made available at the dryad data repository

REFERENCES

Benjamini, Y. and Y. Hochberg. 1995. Controlling the false discovery rate: A practical and
powerful approach to multiple testing. J. Roy. Stat. Soc. Ser. B. (Stat. Method.). 57(1):
289-300.

Bhagwat, M., L. Young and R. R. Robison. 2012. Using BLAT to find sequence similarity in
closely related genomes. Current protocols in bioinformatics. 37: 10.18.11-10.18.24.

638   Bien, T., J. Gadau, A. Schnapp, J. Y. Yew, C. Sievert and K. Dreisewerd. 2019. Detection of

very long-chain hydrocarbons by laser mass spectrometry reveals novel species-, sex-

640   , and age-dependent differences in the cuticular profiles of three *Nasonia* species.

Anal. Bioanal. Chem. 411(13): 2981-2993.

Blomquist, G. J. and A. G. Bagnères. 2010. Insect hydrocarbons: Biology, biochemistry, and

644   chemical ecology. Cambridge University Press, Cambridge, Massachusetts.

646   Blomquist, G. J. and M. D. Ginzel. 2021. Chemical Ecology, Biochemistry, and Molecular

Biology of Insect Hydrocarbons. Annu. Rev. Entomol. 66(1): 45-60.

Blomquist, G. J., D. R. Nelson and M. Derenobales. 1987. Chemistry, biochemistry, and

650   physiology of insect cuticular lipids. Arch. Insect Biochem. Physiol. 6(4): 227-265.

652   Bordenstein, S. R., F. P. O'Hara and J. H. Werren. 2001. *Wolbachia*-induced incompatibility

precedes other hybrid incompatibilities in *Nasonia*. Nature. 409(6821): 707.

Bordenstein, S. R. and J. H. Werren. 1998. Effects of A and B *Wolbachia* and host genotype

656   on interspecies cytoplasmic incompatibility in *Nasonia*. Genetics. 148(4): 1833-1844.

658   Breeuwer, J. A. J. and J. H. Werren. 1990. Microorganisms associated with chromosome

destruction and reproductive isolation between two insect species. Nature. 346: 558-

660   560.

662   Breeuwer, J. A. J. and J. H. Werren. 1995. Hybrid breakdown between two haplodiploid

        species: The role of nuclear and cytoplasmic genes. Evolution. 49: 705-717.

        Broman, K. W. and S. Sen. 2009. A Guide to QTL Mapping with R/QTL. Springer, Dordrecht.

        Broman, K. W., H. Wu, S. Sen and G. A. Churchill. 2003. R/qtl: QTL mapping in experimental

668   crosses. Bioinformatics. 19(7): 889–890.

670   Buchfink, B., C. Xie and D. H. Huson. 2014. Fast and sensitive protein alignment using

        DIAMOND. Nat. Methods. 12: 59–60.

        Buellesbach, J., J. Gadau, L. W. Beukeboom, F. Echinger, R. Raychoudhury, J. H. Werren and

674   T. Schmitt. 2013. Cuticular hydrocarbon divergence in the jewel wasp *Nasonia*:

        Evolutionary shifts in chemical communication channels? J. Evol. Biol. 26(11): 2467-

676   2478.

678   Camacho, C., G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K. Bealer and T. L. Madden.

        2009. BLAST+: architecture and applications. BMC Bioinformatics. 10: 421.

        Campbell, B., J. Heraty, J.-Y. Rasplus, K. Chan, J. Steffen-Campbell and C. Babock. 2000.

682   Molecular Systematics of the Chalcidoidea using 28S-D2 rDNA, in Hymenoptera:

        Evolution, Biodiversity and Biological Control edited by A. D. Austin and M. Dowton.

684   CISRO Publishing, Collingwood.

686   Campbell, B. C., J. D. Steffen-Campbell and J. H. Werren. 1993. Phylogeny of the Nasonia

            species complex (Hymenoptera: Pteromalidae) inferred from an internal transcribed

688   spacer (ITS2) and 28S rDNA sequences. Insect Biochem. Mol. Biol. 2(4): 225-237.

690   Carlson, D. A., U. R. Bernier and B. D. Sutton. 1998. Elution patterns from capillary GC for

            methyl-branched alkanes. J. Chem. Ecol. 24(11): 1845-1865.

        Carlson, D. A., M. S. Mayer, D. L. Silhacek, J. D. James, M. Beroza and B. A. Bierl. 1971. Sex

694   attractant pheromone of the housefly: Isolation, identification, and synthesis.

            Science. 174: 76-77.

        Castresana, J. 2000. Selection of conserved blocks from multiple alignments for their use in

698   phylogenetic analysis. Mol. Biol. Evol. 17(4): 540–552.

700   Chertemps, T., L. Duportets, C. Labeur, R. Ueda, K. Takahashi, K. Saigo and C. Wicker-

            Thomas. 2007. A female-biased expressed elongase involved in long-chain

702   hydrocarbon biosynthesis and courtship behavior in *Drosophila melanogaster*. Proc.

            Natl. Acad. Sci. USA. 104(11): 4273-4278.

        Chung, H. and S. B. Carroll. 2015. Wax, sex and the origin of species: Dual roles of insect

706   cuticular hydrocarbons in adaptation and mating. Bioessays. 37(7): 822-830.

708   Chung, H., D. W. Loehlin, H. D. Dufour, K. Vaccarro, J. G. Millar and S. B. Carroll. 2014. A

single gene affects both ecological divergence and mate choice in *Drosophila*.

710   Science. 343(6175): 1148-1151.

712   Cinnamon, E., R. Makki, A. Sawala, L. P. Wickenberg, G. J. Blomquist, C. Tittiger, Z. e. Paroush

and A. P. Gould. 2016. *Drosophila* Spidey/Kar regulates oenocyte growth via PI3-

714   kinase signaling. PLoS Genet. 12(8): e1006154.

716   Coyne, J. A. and R. Oyama. 1995. Localization of pheromonal sexual dimorphism in

Drosophila melanogaster and its effect on sexual isolation. Proc. Natl. Acad. Sci. USA.

718   92(21): 9505-9509.

720   Coyne, J. A., C. Wicker-Thomas and J. M. Jallon. 1999. A gene responsible for a cuticular

hydrocarbon polymorphism in *Drosophila melanogaster*. Genet. Res. 73(3): 189-203.

Dallerac, R., C. Labeur, J. M. Jallon, D. C. Knippie, W. L. Roelofs and C. Wicker-Thomas. 2000.

724   A Δ9 desaturase gene with a different substrate specificity is responsible for the

cuticular diene hydrocarbon polymorphism in *Drosophila melanogaster*. Proc. Natl.

726   Acad. Sci. USA. 97(17): 9449-9454.

728   Darling, D. C. and J. H. Werren. 1990. Biosystematics of *Nasonia* (Hymenoptera:

Pteromalidae): Two new species reared from birds' nests in North America. Annals of

730   the Entomological Society of America. 83: 352-370.

732   Davies, N. J. and E. Tauber. 2015. WaspAtlas: a *Nasonia vitripennis* gene database and

analysis platform. Database. bav103: 1-7.

Dembeck, L. M., K. Böröczky, W. Huang, C. Schal, R. R. H. Anholt, T. F. C. Mackay and D. J.

736   Kliebenstein. 2015. Genetic architecture of natural variation in cuticular hydrocarbon

composition in *Drosophila melanogaster*. eLife. 4: e09861.

Desjardins, C. A., J. Gadau, J. A. Lopez, O. Niehuis, A. R. Avery, D. W. Loehlin, S. Richards, J. K.

740   Colbourne and J. H. Werren. 2013. Fine-scale mapping of the Nasonia genome to

chromosomes using a high-density genotyping microarray. G3: Genes, Genomes,

742   Genetics. 3(2): 205-215.

744   Dixon, P. 2003. VEGAN, a package of R functions for community ecology. J. Veg. Sci. 14(6):

927-930.

El-Gebali, S., J. Mistry, A. Bateman, S. R. Eddy, A. Luciani, S. C. Potter, M. Qureshi, L. J.

748   Richardson, G. A. Salazar, A. Smart et al. 2019. The Pfam protein families database in

2019. Nucleic Acids Research. 47(D1): D427-D432.

Emms, D. M. and S. Kelly. 2015. OrthoFinder: solving fundamental biases in whole genome

752   comparisons dramatically improves orthogroup inference accuracy. Genome Biology.

16(1): 157.

Erickson, D. L., C. B. Fenster, H. K. Stenøien and D. Price. 2004. Quantitative trait locus

756   analyses and the study of evolutionary process. Mol. Ecol. 13(9): 2505-2522.

758   Ferveur, J.-F. 2005. Cuticular hydrocarbons: Their evolution and roles in *Drosophila*

       pheromonal communication. Behav. Genet. 35(3): 279-295.

       Finet, C., K. Slavik, J. Pu, S. B. Carroll and H. Chung. 2019. Birth-and-death evolution of the

762   fatty acyl-CoA reductase (FAR) gene family and diversification of cuticular

       hydrocarbon synthesis in *Drosophila*. Genome Biol. Evol. 11(6): 1541–1551.

       Flicek, P., M. R. Amode, D. Barrell, K. Beal, K. Billis, S. Brent, D. Carvalho-Silva, P. Clapham, G.

766   Coates, S. Fitzgerald et al. 2014. Ensembl 2014. Nucleic acids research. 42: D749-

       D755.

       Gadau, J., O. Niehuis, A. Peire, J. H. Werren, E. Baudry and L. W. Beukeboom. 2008. The

770   jewel wasp, *Nasonia* spp., p. 27-41 in Genome Mapping and Genomics in Animals

       edited by W. Hunter and C. Kole. Springer, Berlin Heidelberg.

       Gadau, J., R. E. Page and J. H. Werren. 2002. The genetic basis of the interspecific differences

774   in wing size in *Nasonia* (Hymenoptera: Pteromalidae): Major quantitative trait loci

       and epistasis. Genetics. 161: 673-684.

Gramates, L. S., S. J. Marygold, G. d. Santos, J.-M. Urbano, G. Antonazzo, B. B. Matthews, A.

778   J. Rey, C. J. Tabone, M. A. Crosby, D. B. Emmert et al. 2017. FlyBase at 25: looking to

the future. Nucleic Acids Research. 45(D1): D663-D671.

Gu, P., W. H. Welch, L. Guo, K. M. Schegg and G. J. Blomquist. 1997. Characterization of a

782   novel microsomal fatty acid synthetase (FAS) compared to a cytosolic FAS in the

housefly, *Musca domestica*. Comp. Biochem. Physiol. B-Biochem. Mol. Biol. 118(2):

784   447–456.

786   Haley, C. S. and S. A. Knott. 1992. A simple regression method for mapping quantitative trait

loci in line crosses using flanking markers. Heredity. 69: 315-324.

Hartl, D. 2020. Population genetics of complex traits, p. 225-261 in A primer of population

790   genetics and genomics edited by. 4 ed. Oxford University Press, Oxford.

792   Hollander, M. and D. A. Wolfe. 1999. Nonparametric Statistical Methods. Wiley, New York.

794   Holze, H., L. Schrader and J. Buellesbach. 2021. Advances in deciphering the genetic basis of

insect cuticular hydrocarbon biosynthesis and variation. Heredity. 126(2): 219-234.

Howard, R. W. and C. Blomquist. 1982. Chemical ecology and biochemistry of insect

798   hydrocarbons. Annu. Rev. Entomol. 27: 149-172.

800   Howard, R. W. and G. J. Blomquist. 2005. Ecological, behavioral, and biochemical aspects of

         insect hydrocarbons. Annu. Rev. Entomol. 50: 371-393.

       Hoy, M. A. 2005. Behavior of Insects: Genetic Analysis by Crossing and Selection, p. 278-281

804   in Encyclopedia of Entomology edited by J. L. Capinera. Springer, Dordrecht.

806   Hunter, W. and C. Kole. 2008. Genome Mapping and Genomics in Arthropods. Springer,

         Berlin Heidelberg.

       Jallon, J. M. 1984. A few chemical words exchanged by *Drosophila* during courtship and

810   mating. Behav. Genet. 14(5): 441-478.

812   Jaspers, M. H. J., R. Pflanz, D. Riedel, S. Kawelke, I. Feussner and R. Schuh. 2014. The fatty

         acyl-CoA reductase Waterproof mediates airway clearance in *Drosophila*. Dev. Biol.

814   385(1): 23–31.

816   Juarez, P., J. Chase and G. J. Blomquist. 1992. A microsomal fatty acid synthetase from the

         integument of *Blattella germanica* synthesizes methyl-branched fatty acids,

818   precursors to hydrocarbon and contact sex pheromone. Arch. Biochem. Biophys.

         293(2): 333–341.

       Kather, R. and S. J. Martin. 2015. Evolution of cuticular hydrocarbons in the Hymenoptera: A

822   meta-analysis. J. Chem. Ecol. 41(10): 871-883.

824 Keilwagen, J., M. Wenk, J. L. Erickson, M. H. Schattat, J. Grau and F. Hartung. 2016. Using

intron position conservation for homology-based gene prediction. Nucleic Acids

826 Research. 44(9): e89.

828 Kent, W. J. 2002. BLAT--the BLAST-like alignment tool. Genome Res. 12(4): 656–664.

830 Lander, E. S. and D. Botstein. 1989. Mapping mendelian factors underlying quantitative traits

using RFLP linkage maps. Genetics. 121(1): 185-199.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis and R.

834 Durbin. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics.

25(16): 2078–2079.

Ligges, U. and M. Maechler. 2003. Scatterplot3d – An R package for visualizing multivariate

838 data. J. Stat. Soft. 8(11): 1-20.

840 Lockey, K. H. 1980. Insect cuticular hydrocarbons. Comp. Biochem. Physiol. B: Biochem. Mol.

Biol. 65(3): 457-462.

Lockey, K. H. 1988. Lipids of the insect cuticle: origin, composition and function. Comp.

844 Biochem. Physiol. B: Biochem. Mol. Biol. 89(4): 595-645.

846 Löytynoja, A. 2014. Phylogeny-aware alignment with PRANK, p. 155-170 in Multiple

Sequence Alignment Methods edited by D. J. Russell. Humana Press, Totowa, NJ.

Löytynoja, A. and N. Goldman. 2008. Phylogeny-aware gap placement prevents errors in

850   sequence alignment and evolutionary analysis. Science. 320(5883): 1632–1635.

852   Mackay, T. F. C. 2001a. The Genetic Architecture of Quantitative Traits. Annu. Rev. Genet.

35(1): 303-339.

Mackay, T. F. C. 2001b. Quantitative trait loci in *Drosophila*. Nature Reviews Genetics. 2(1):

856   11-20.

858   Mackay, T. F. C., E. A. Stone and J. F. Ayroles. 2009. The genetics of quantitative traits:

challenges and prospects. Nature Reviews Genetics. 10(8): 565-577.

Manichaikul, A., J. Y. Moon, S. Sen, B. S. Yandell and K. W. Broman. 2009. A model selection

862   approach for the identification of quantitative trait loci in experimental crosses,

allowing epistasis. Genetics. 181(3): 1077-1086.

Martin, S. and F. Drijfhout. 2009. A review of ant cuticular hydrocarbons. J. Chem. Ecol.

866   35(10): 1151-1161.

868   Misof, B., S. Liu, K. Meusemann, R. S. Peters, A. Donath, C. Mayer, P. B. Frandsen, J. Ware, T.

Flouri, R. G. Beutel et al. 2014. Phylogenomics resolves the timing and pattern of

870   insect evolution. Science. 346(6210): 763-767.

872 Morandin, C., V. P. Brendel, L. Sundstrom, H. Helantera and A. S. Mikheyev. 2019. Changes in

gene DNA methylation and expression networks accompany caste specialization and

874 age-related physiological changes in a social insect. Mol. Ecol. 28(8): 1975-1993.

876 Murrell, B., S. Weaver, M. D. Smith, J. O. Wertheim, S. Murrell, A. Aylward, K. Eren, T.

Pollner, D. P. Martin, D. M. Smith et al. 2015. Gene-wide identification of episodic

878 selection. Mol. Biol. Evol. 32(5): 1365–1371.

880 Nelson, D. R. and G. J. Blomquist. 1995. Insect waxes, p. 1-90 in Waxes: Chemistry, molecular

biology and functions edited by R. J. Hamilton. Oily Press, Dundee, Scotland.

Niehuis, O., J. Buellesbach, A. K. Judson, T. Schmitt and J. Gadau. 2011. Genetics of cuticular

884 hydrocarbon differences between males of the parasitoid wasps *Nasonia giraulti* and

Nasonia vitripennis. Heredity. 107(1): 61-70.

Niehuis, O., J. D. Gibson, M. S. Rosenberg, B. A. Pannebakker, T. Koevoets, A. K. Judson, C. A.

888 Desjardins, K. Kennedy, D. Duggan, L. W. Beukeboom et al. 2010. Recombination and

its impact on the genome of the haplodiploid parasitoid wasp *Nasonia*. PloS one.

890 5(1): e8597.

892 Niehuis, O., A. K. Judson and G. J. 2008. Cytonuclear genic incompatibilities cause increased

mortality in male F2 hybrids of *Nasonia giraulti* and *N. vitripennis*. Genetics. 178(1):

894 413-426.

896 Niehuis, O., A. K. Judson, J. H. Werren, W. B. Hunter, P. M. Dang, S. E. Dowd, B. Grillenberger,

L. W. Beukeboom and J. Gadau. 2007. Species-diagnostic single-nucleotide

898 polymorphism and sequence-tagged site markers for the parasitic wasp genus

Nasonia (Hymenoptera: Pteromalidae). J. Econ. Entomol. 100(4): 1033-1036.

Orr, A. H. 2001. The genetics of species differences. Trends Ecol. Evol. 16(7): 343-350.

Otte, T., M. Hilker and S. Geiselhardt. 2018. Phenotypic plasticity of cuticular hydrocarbon

904 profiles in insects. J. Chem. Ecol. 44(3): 235-247.

906 Ouellette, L. A., R. W. Reid, S. G. Blanchard and C. R. Brouwer. 2017. LinkageMapView -

Rendering High Resolution Linkage and QTL Maps. Bioinformatics. 34(2): 306-307.

Quinlan, A. R. and I. M. Hall. 2010. BEDTools: a flexible suite of utilities for comparing

910 genomic features. Bioinformatics. 26(6): 841–842.

912 R Development Core Team. 2018. R: A language and environment for statistical computing.

Vienna, Austria, R foundation for statistical computing, http://www.R-project.org.

Ramsay, J. A. 1935. The evaporation of water from the cockroach. J. Exp. Biol. 12: 373-383.

Raychoudhury, R., C. A. Desjardins, J. Buellesbach, D. W. Loehlin, B. K. Grillenberger, L. W.

918 Beukeboom, T. Schmitt and J. H. Werren. 2010. Behavioral and genetic characteristics

of a new species of *Nasonia*. Heredity. 104(3): 278-288.

Richardson, P. M., W. P. Holmes and G. B. Saul. 1987. The effect of tetracycline on
nonreciprocal cross incompatibility in *Mormoniella* [= *Nasonia*] vitripennis. J.
Invertebr. Pathol. 50(3): 176-183.

Slate, J. 2005. Quantitative trait locus mapping in natural populations: progress, caveats and
future directions. Mol. Ecol. 14(2): 363-379.

928   Sprenger, P. P. and F. Menzel. 2020. Cuticular hydrocarbons in ants (Hymenoptera:
Formicidae) and other insects: How and why they differ among individuals, colonies,
930   and species. Myrmecol. News. 30: 1-26.

932   Thomas, G. W. C., E. Dohmen, D. S. T. Hughes, S. C. Murali, M. Poelchau, K. Glastad, C. A.
Anstead, N. A. Ayoub, P. Batterham, M. Bellair et al. 2020. Gene content evolution in
934   the arthropods. Genome Biology. 21(1): 15.

936   Venables, W. N. and B. D. Ripley. 2002. Modern Applied Statistics with S. Springer, New York.

938   Werren, J. H. and D. W. Loehlin. 2009. The parasitoid wasp *Nasonia*: An emerging model
system with haploid male genetics. Cold Spring Harbor Protocols. 10.

Werren, J. H., S. Richards, C. A. Desjardins, O. Niehuis, J. Gadau, J. K. Colbourne and The
942   Nasonia Genome Working Group. 2010. Functional and evolutionary insights from
the genomes of three parasitoid *Nasonia* species. Science. 327(5963): 343-348.

Wicker-Thomas, C., D. Garrido, G. Bontonou, L. Napal, N. Mazuras, B. Denis, T. Rubin, J.-P.

946   Parvy and J. Montagne. 2015. Flexible origin of hydrocarbon/pheromone precursors

in *Drosophila melanogaster*. J. Lipid Res. 56(11): 2094-2101.

Wicker-Thomas, C., C. Henriet and R. Dallerac. 1997. Partial characterization of a fatty acid

950   desaturase gene in *Drosophila melanogaster*. Insect Biochem. Mol. Biol. 27(11): 963-

972.

Yang, J., J. Zhu and R. W. Williams. 2007. Mapping the genetic architecture of complex traits

954   in experimental populations. Bioinformatics. 23(12): 1527-1536.

956   Zar. 1999. Biostatistical Analysis. Prentice-Hall, Englewood Cliffs, NJ.

958   Zuo, W., C. Li, Y. Luan, H. Zhang, X. Tong, M. Han, R. Gao, H. Hu, J. Song, F. Dai et al. 2018.

Genome-wide identification and analysis of elongase of very long chain fatty acid

960   genes in the silkworm, Bombyx mori. Genome. 61(3): 167–176.

TABLES

Tab. 1: Cuticular hydrocarbons (CHCs) identified in male *Nasonia* profiles and the corresponding QTL from crosses between *Nasonia longicornis* and *Nasonia vitripennis* as well as between *N. longicornis* and *N. girauli*. Indicated are the retention indices (RI), compound identifications or possible configurations in case of ambiguities, chromosome number (Chr.), QTL position (pos.), confidence interval for the respective QTL position, hybrid cross (LV or LG), ratios of the CHC compounds in parental males of the two species for each respective cross (percentages with standard deviations) and significance assessments comparing the two ratios (Benjamini Hochberg-corrected Mann-Whitney U tests).

RI	Compound identification / possible configurations	Chr.	QTL pos.	Confidence interval	Cross	NL ♂ ratio	NV ♂ ratio	NG ♂ ratio	Significance
2900	C29	4	47.6	37.6 - 57.6	LV	1.11 ± 0.44	0.98 ± 0.5		n.s.
2934	9-; 11-; 13-; 15-MeC29	4	56.8	46.8 - 66.8	LV	0.13 ± 0.21	0.08 ± 0.12		n.s.
		5	64.2	54.2 - 73	LG	0.13 ± 0.21		0.01 ± 0.05	0.02346
2942	7-MeC29	4	60	50 - 70	LV	0.78 ± 0.69	0.6 ± 0.31		n.s.
2951	5-MeC29	4	57.9	47.9 - 67.9	LV	0.36 ± 0.28	0.37 ± 0.16		n.s.
2973	3-MeC29	2	49.6	39.6 - 59.6	LV	0.15 ± 0.19	0.11 ± 0.06		n.s.
		4	57.9	47.9 - 67.9	LV				
		5	63.3	53.3 - 73	LG	0.15 ± 0.20		0.04 ± 0.07	n.s.
2982	15,17-DiMeC29	2	71.5	61.5 - 81.5	LG	0.09 ± 0.1		0.01 ± 0.03	n.s.
3100	C31	1	0	0 - 10	LV				
		1	48.9	38.9 - 58.9	LV				
		2	49.6	39.6 - 59.6	LV				
		3	77.9	67.9 - 87.9	LV				
3131	9-; 11-; 13-; 15-MeC31	2	61.2	51.2 - 71.2	LV	2.13 ± 0.89	1.99 ± 0.95		n.s.
3141	7-MeC31	1	13.3	3.3 - 23.3	LV	22.71 ± 4.49	11.43 ± 1.11		< 0.0001
		2	0.7	0 - 10.7	LV				
		2	54.3	44.3 - 64.3	LV				
		3	60.2	50.2 - 70.2	LV				
3149	5-MeC31	2	53.3	43.3 - 63.3	LV	7.17 ± 1.8	4.42 ± 0.65		0.00021
		5	42	32 - 52	LV				
3158	11,15-; 11,17-; 11,21-DiMeC31	2	69.4	59.4 - 79.4	LV	0.37 ± 0.12	0.46 ± 0.13		n.s.
		3	50.3	40.3 - 60.3	LV				
3171	7,11-DiMeC31	2	86.9	76.9 - 86.9	LV	4.84 ± 1.06	3.73 ± 0.35		0.01025

RI	Compound identification / possible configurations	Chr.	QTL pos.	Confidence interval	Cross	NL ♂ ratio	NV ♂ ratio	NG ♂ ratio	Significance
3177	3-MeC31 (+7,23-DiMeC31)	2	46	36 - 56	LV	3.42 ± 0.73	2.55 ± 0.55		n.s.
		5	49.8	39.8 - 59.8	LV				
3171	7,11-DiMeC31	2	86.9	76.9 - 86.9	LV	4.84 ± 1.06	3.73 ± 0.35		0.01025
3187	7,25-DiMeC31	2	44.9	34.9 - 54.9	LV	0.93 ± 0.2	0.61 ± 0.18		0.00655
		3	0	0 - 10	LV				
		5	45.9	35.9 - 55.9	LV				
3226	3,7,9-; 3,7,11-; 3,7,15-TriMeC31	1	57.7	47.7 - 67.7	LV	0.23 ± 0.09	0.16 ± 0.08		n.s.
		2	71.5	61.5 - 81.5	LV				
3234	6-MeC32	1	63	53 - 73	LV	0.47 ± 0.14	0.39 ± 0.05		n.s.
3261	3,7,11,15-TetraMeC31	1	60.8	50.8 - 70.8	LV	0.53 ± 0.35	0.33 ± 0.1		n.s.
		5	59.4	49.4 - 69.4	LG	0.53 ± 0.35		0.67 ± 0.18	n.s.
3273	9-C33ene	2	46	36 - 56	LV	0.31 ± 0.18	1.45 ± 0.27		< 0.0001
		2	82.9	72.9 - 86.9	LV				
		3	29.2	19.2 - 39.2	LV				
3281	7-C33ene	1	44.5	34.5 - 54.5	LV	0.17 ± 0.09	0.96 ± 0.2		< 0.0001
		3	39.5	29.5 - 49.5	LV				
3300	C33	1	48	38 - 58	LV	0.41 ± 0.2	0.99 ± 0.25		< 0.0001
		5	49.8	39.8 - 59.8	LV				
3330	9-; 11-; 13-; 15-MeC33	2	72.5	62.5 - 82.5	LV	4.98 ± 1.11	3.97 ± 1.26		n.s.
		3	61.1	51.1 - 71.1	LV				
		3	33.2	23.2 - 43.2	LG	4.98 ± 1.11		8.02 ± 0.92	0.00044
		4	55.5	45.5 - 65.5	LG				
		5	52.7	42.7 - 62.7	LG				
3339	7-MeC33	5	48.8	38.8 - 58.8	LV	3.07 ± 1.03	2.93 ± 0.68		n.s.
3355	11,15-; 11,23- 11,25-DiMeC33	2	72.5	62.5 - 82.5	LV	1.97 ± 0.87	0.66 ± 0.42		0.00059
		5	45.9	35.9 - 55.9	LV				
		4	55.5	45.5 - 65.5	LG				
3360	7,19-; 7,21-DiMeC33	1	59.8	49.8 - 69.8	LV	1.23 ± 0.36	1.68 ± 0.56		n.s.
		5	43.8	33.8 - 53.8	LV				
		4	55.5	45.5 - 65.5	LG	1.23 ± 0.36		1.24 ± 0.21	n.s.
3370	7,23-DiMeC33	2	56.3	46.3 - 66.3	LV	7.52 ± 1.63	7.71 ± 1.21		n.s.
		2	86.9	76.9 - 86.9	LV				
		5	45	35 - 55	LV				

RI	Compound identification / possible configurations	Chr.	QTL pos.	Confidence interval	Cross	NL ♂ ratio	NV ♂ ratio	NG ♂ ratio	Significance
3376	5,9-; 5,15-; 5,17-; 5,19-DiMeC33 (+11,15,23-TriMeC33)	2	54.3	44.3 - 64.3	LV	4.4 ± 1.22	5.85 ± 0.81		0.01573
		3	63.8	53.8 - 73.8	LV				
3396	3,9-; 3,15-; 3,17-DiMeC33	2	49.3	39.3 - 59.3	LV	0.78 ± 0.64	1.09 ± 0.23		n.s.
3405	5,9,11-; 5,9,15-TriMeC33	1	2	0 - 12	LV	0.04 ± 0.1	2.04 ± 0.68		< 0.0001
		2	74.6	64.6 - 84.6	LV				
		3	69.4	59.4 - 79.4	LV				
		5	50.7	40.7 - 60.7	LV				
3430	8,10-; 8,12-; 8,14-; 8,16-; 8,18-DiMeC34	1	5.1	0 - 15.1	LV	0.32 ± 0.14	0.92 ± 0.28		< 0.0001
		2	72.5	62.5 - 82.5	LV				
		5	43.8	33.8 - 53.8	LV				
3453	3,7,11,15-TetraMeC33	2	75.7	65.7 - 85.7	LV	0.89 ± 0.4	1.57 ± 0.35		0.00211
		5	58.4	48.4 - 68.4	LG	0.89 ± 0.4		4 ± 0.71	0.00011
3477	7,11-; 7,13-; 7,15-DiMeC34	2	56.3	46.3 - 66.3	LV	0.32 ± 0.19	0.41 ± 0.17		n.s.
3524	13-; 15-;17-MeC35	2	81.9	71.9 - 86.9	LV	1.65 ± 0.51	2.58 ± 0.44		0.00086
		5	0	0 - 10	LV				
3534	7-MeC35	2	81.9	71.9 - 86.9	LV	0.36 ± 0.1	0.77 ± 0.18		< 0.0001
		5	49.8	39.8 - 59.8	LV				
3549	11,17-; 11,19-; 11,21-; 11,23-; 13,17-; 13,19-; 13,21-; 13,17-; 13,19-; 13,21-;13,23-; 15,17-; 15,19-; 15,21-; 15,23-DiMeC35	1	60.8	50.8 - 70.8	LV	4.35 ± 0.85	3.08 ± 0.92		0.02070
		5	43.8	33.8 - 53.8	LV				
3563	7,19-; 7,21-; 7,23-DiMeC35	1	51.5	41.5 - 61.5	LV	2.33 ± 0.68	7.07 ± 1.47		< 0.0001
		2	77.7	67.7 - 86.9	LV				
		5	43.8	33.8 - 53.8	LV				
3572	5,15-; 5,17-; 5,19-; 5,21-; 5,23-DiMeC35	2	4.7	0 - 14.7	LV	2.84 ± 0.58	5.21 ± 0.73		< 0.0001
		2	58.2	48.2 - 68.2	LV				
3585	7,15,21-; 7,15,23-; 7,17,21-; 7,17,23-; 7,19,21-; 7,19,23-TriMeC35	2	71.5	61.5 - 81.5	LV	0.71 ± 0.34	1.04 ± 0.29		n.s.
		3	54.8	44.8 - 64.8	LV				
3603	5,9,13-; 5,9,15-; 5,9,17-; 5,9,19-; 5,9,21-; 5,11,13-; 5,11,15-; 5,11,17-; 5,11,19-; 5,11,21-TriMeC35	2	58.2	48.2 - 68.2	LV	0.42 ± 0.3	1.65 ± 0.4		< 0.0001
3721	13-; 15-; 17-; 19-MeC37	2	79.8	69.8 - 86.9	LV				
3745	11,19-; 11,21-; 11,23-; 11,25-; 13,19-; 13,21-; 13,23-; 15,19-; 15,21-; 15,23-DiMeC37	2	86.9	76.9 - 86.9	LV	1.48 ± 0.4	1.38 ± 0.46		n.s.
		5	43.8	33.8 - 53.8	LV				

RI	Compound identification / possible configurations	Chr.	QTL pos.	Confidence interval	Cross	NL ♂ ratio	NV ♂ ratio	NG ♂ ratio	Significance
3764	7,19-; 7,21-; 7,23-DiMeC37	1	60.8	50.8 - 70.8	LV	0.58 ± 0.27	2.39 ± 0.69		< 0.0001
		2	79.8	69.8 - 86.9	LV				
		5	43.8	33.8 - 53.8	LV				
		2	82.9	72.9 - 86.9	LG	0.58 ± 0.27		0.61 ± 0.16	n.s.
3768	5,15-; 5,17-; 5,23-DiMeC37	2	69.4	59.4 - 79.4	LV	1.03 ± 0.46	1.42 ± 0.35		n.s.
3779	7,15,17-; 7,15,19-; 7,15,23-; 7,19,17-; 7,19,19-; 7,19,23-TriMeC37	2	76.7	66.7 - 86.7	LV	0.12 ± 0.14	0.53 ± 0.36		0.00336

Tab. 2: Candidate genes for CHC biosynthesis and variation in *Nasonia vitripennis* based on orthology towards *Drosophila melanogaster,* where the impact on CHC profiles of the respective genes has been demonstrated through targeted knockdown studies. Indicated are the gene annotations, gene IDs obtained from RefSeq, descriptions of the gene function obtained from NCBI, linkage map positions (centimorgan (cM) on the respective chromosome (c)), recombination rates (Mb/cM), and evidence of orthology based on the three methods reciprocal blast search (RB), OrthoFinder (OF) and WaspAtlas (WA). Genes for which exactly one ortholog could be unambiguously identified are indicated with a '+', in other cases the method(s) that have generated the most consistently reliable results in each case are shown. Genes that can clearly be assigned to a particular step in CHC biosynthesis (compare to Fig. 1) are indicated in bold.

Gene annotation	RefSeq ID	Description NCBI	Linkage map position	Recomb. rate (Mb/cM)	Evidence of orthology
acc	**100123347**	acetyl-CoA carboxylase	c1: 75.9 cM	0.1114	+
app	100121627	palmitoyltransferase ZDHHC18	c2: 38 cM	13.3806	+
CG14688	100120956	phytanoyl-CoA dioxygenase	c1: 48.2 cM	20.6198	+
CG16979	100119053	ufm1-specific protease 2-like	c3: 64.2 cM	0.1015	+
CG5599	100122501	lipoamide acyltransferase component ofbranched-chain alpha-keto acid dehydrogenase complex, mitochondrial	c4: 6.6 cM	0.2967	+
CG7724	100120136	3 beta-hydroxysteroid dehydrogenase/Delta 5-->4-isomerase type 4	c2: 1.5 cM	0.5931	+
Ndufs6	100119717	NADH dehydrogenase (ubiquinone) Fe-Sprotein 6, 13kDa (NADH-coenzyme Q reductase)	c2: 43.8 cM	1.1653	+
cyp4g44	**100113546**	cytochrome P450 4G44	c1: 47.5 cM	2.6119	WA
desi	100115157	uncharacterized LOC100115157	c4: 35.8 cM	0.1267	+

Gene annotation	RefSeq ID	Description NCBI	Linkage map position	Recomb. rate (Mb/cM)	Evidence of orthology
fas1	**100121447**	fatty acid synthase	c1: 60.6 cM	0.3184	OF
fas2	**100122099**	fatty acid synthase-like	c3: 5.8 cM	0.556	OF
fas3	**100122083**	fatty acid synthase-like	c3: 5.8 cM	0.556	OF
fas4	**100119597**	fatty acid synthase-like	c1: 49.6 cM	20.78	OF
fas5	**100678737**	fatty acid synthase-like	c5: 45.3 cM	0.7646	OF
fatp1	**100120211**	long-chain fatty acid transport protein 4-like	c2: 20.4 cM	0.8866	RB
hacd1	**100119292**	very-long-chain (3R)-3-hydroxyacyl-CoAdehydratase hpo-8	c4: 50.4 cM	4.7456	+
hacd2	**100115280**	very-long-chain (3R)-3-hydroxyacyl-CoAdehydratase	c3: 38 cM	11.8001	+
nrt	100121165	neurotactin	c3: 73 cM	0.1823	+
phgpx_1	100677983	probable phospholipid hydroperoxideglutathione peroxidase	c3: 38 cM	11.8001	RB,WA
phgpx_2	100114603	probable phospholipid hydroperoxideglutathione peroxidase	c4: 4.4 cM	0.2203	RB,WA
phgpx_3	100123141	phospholipid hydroperoxide glutathioneperoxidase, mitochondrial-like	c4: 4.4 cM	0.2203	RB,WA
prx6005	100117827	peroxiredoxin-6	c2: 38 cM	13.3806	
pxd	100119885	peroxidase	c3: 68.6 cM	0.3195	OF,WA
pxn	100116337	peroxidasin	c1: 93.4 cM	0.1259	+
sc2	**107980451**	very-long-chain enoyl-CoA reductase	c4: 32.1 cM	0.1186	+
spidey	**100121752**	very-long-chain 3-oxoacyl-CoA reductase	c2: 52.6 cM	0.3081	+

Tab. 3: *Elongase* (*ELO*) and *fatty acyl-CoA reductase* (*FAR*) genes in the *Nasonia* reference genome homologous to *D. melanogaster ELOs* and *FARs* with a demonstrated impact on CHC profiles (*ELO*s: *CG18609*, *CG30008, CG9458, eloF*; *FAR*s: *CG10097, CG13091;* see Tab. S2). Indicated are the gene annotations, gene and respective protein IDs obtained from RefSeq, linkage map positions and recombination rates (Kb/cM).

Gene annotation	RefSeq Gene ID	RefSeq Protein IDs	Linkage map position	Recombination rate (Kb/cM)
elo1	100114999	XP_016845887.1 XP_008214189.1 XP_008214188.1 XP_001599838.1	c1: 46.7 cM	3.6211
elo2	100115035	XP_001599867.1	c1: 46.7 cM	3.6211
elo3	100115068	XP_003428006.1	c1: 46.7 cM	3.6211
elo4	100115103	XP_008214193.1 XP_001599914.2 XP_016845633.1	c1: 46.7 cM	3.6211
elo5	100115131	XP_001599942.1	c1: 46.7 cM	3.6211
elo6	100115199	XP_001599996.2 XP_016838633.1	c1: 46.7 cM	3.6211
elo7	100115239	XP_016838697.1 XP_001600017.1	c1: 46.7 cM	3.6211
elo8	100115278	XP_001600048.1	c1: 46.7 cM	3.6211
elo9	100116202	XP_008205141.1	c3: 42.3 cM	0.3664
elo10	100118619	XP_008205429.1	c4: 47.5 cM	0.8598
elo11	100120094	XP_001603768.2	c3: 38 cM	11.8001
elo12	100123468	XP_001607111.2	c5: 16.1 cM	1.8931
far1	100115559	XP_008213286.1	c1: 48.9 cM	22.4845
far2	100115595	XP_008213287.2	c1: 48.9 cM	22.4845
far3	100115642	XP_001600309.1	c1: 48.9 cM	22.4845
far4	100115711	XP_001601168.1	c1: 49.6 cM	20.7846
far5	100117683	XP_016845554.1 XP_016845553.1 XP_016845552.1 XP_016845551.1 XP_008205703.1	c1: 49.6 cM	20.7846
far6	100117801	XP_008205709.1 XP_001601942.1	c1: 49.6 cM	20.7846
far7	100117837	XP_001601970.1	c1: 49.6 cM	20.7846
far8	100118857	XP_008212002.1 XP_008212003.1	c1: 51.8 cM	2.0759
far9	100118893	XP_001602762.2	c1: 51.8 cM	2.0759
far10	100119003	XP_001602857.1	c1: 51.8 cM	2.0759
far11	100121470	XP_016841641.1 XP_016841638.1 XP_016841633.1	c1: 48.2 cM	20.6198

Gene annotation	RefSeq Gene ID	RefSeq Protein IDs	Linkage map position	Recombination rate (Kb/cM)
far12	100121766	XP_008217013.1	c5: 19 cM	0.2367
far13	100123790	XP_001607507.3	c1: 83.2 cM	0.5953
far14	103317077	XP_016836513.1	c1: 51.8 cM	2.0759
far15	100117218	XP_001601521.2	c2: 85.4 cM	0.1672
far16	100117254	XP_001601550.4	c2: 85.4 cM	0.1672
far17	100117104	XP_001601438.1	c2: 85.4 cM	0.1672
far18	100117143	XP_001601466.1	c2: 85.4 cM	0.1672
far19	100117180	XP_001601494.1	c2: 85.4 cM	0.1672
far20	100117760	XP_016836801.1 XP_016836800.1	c1: 49.6 cM	20.7846
far21	100679544	XP_003426741.2	c2: 86.9 cM	NA
far22	100117878	XP_016843183.1	c3: 38 cM	0.1544

Tab. 4.: Orthologous candidate genes for CHC biosynthesis and variation in the *Nasonia* species complex as well as two representative species from the two most closely related genera, *Trichomalopsis sarcophagae* and *Muscidifurax raptorellus* (see Fig. 2 for phylogeny). Genes with identified orthologs in all analyzed species were subjected to a d_N/d_S ratio analysis to screen for signatures of positive selection with BUSTED. Indicated are gene annotations and gene IDs obtained from the *N. vitripennis* reference genome, species IDs where no orthologous gene could be unambiguously identified, and Benjamini-Hochberg-corrected p-values from the BUSTED analysis for signatures of positive selection. Species abbreviations: MR: *M. raptorellus*, TS: *T. sarcophagae*, NG: *N. giraulti*, NO: *N. oneida*, NL: *N. longicornis*.

Nv gene annotation	Nv gene ID	Positive selection (BUSTED, *p*-value)	Species with no identifiable ortholog
acc	100123347	0.081	
app	100121627	0.5	
CG14688	100120956		MR
CG16979	100119053	0.283	
CG5599	100122501	0.073	
CG7724	100120136	0.5	
Ndufs6	100119717	0.5	
cyp4g44	100113546	0.5	
desi	100115157	0.5	
fas1	**100121447**	**0.013**	
fas2	100122099		MR

Nv gene annotation	Nv gene ID	Positive selection (BUSTED, *p*-value)	Species with no identifiable ortholog
fas3	100122083	0.5	
fas4	100119597		MR
fas5	100678737		TS, MR
fatp1	100120211		NG, MR
hacd1	100119292	0.5	
hacd2	100115280		NO, NG, NL, TS, MR
nrt	100121165		NG, MR
phgpx_1	100677983	0.5	
phgpx_2	100114603	0.5	
phgpx_3	100123141	0.5	
prx6005	100117827	0.363	
pxd	100119885	0.5	
pxn	100116337	0.071	
sc2	107980451	0.363	
spidey	100121752	0.5	