1

An improved vector for baculovirus-mediated protein production in mammalian cells

Wenjun Guo[1,4], Mengmeng Wang[1,2,3,4], and Lei Chen[1,2,3]*

[1] State Key Laboratory of Membrane Biology, College of Future Technology, Institute of Molecular Medicine, Peking University, Beijing Key Laboratory of Cardiometabolic Molecular Medicine, Beijing 100871, China

[2] Peking-Tsinghua Center for Life Sciences, Peking University, Beijing 100871, China

[3] Academy for Advanced Interdisciplinary Studies, Peking University, Beijing 100871, China

[4] These authors contribute equally

*Correspondence: Lei Chen, chenlei2016@pku.edu.cn

Key words: BacMam; baculovirus; membrane protein; TRPC; K_{ATP}.

Abstract

BacMam system utilizes baculovirus to deliver exogenous genes into mammalian cells and is

extensively used for recombinant production of eukaryotic proteins. Here, we described an

improved BacMam vector (pBMCL1, Addgene#178203) which allows convenient tracing of

virus production, provides higher infection efficiency towards mammalian cells, minimizes

unwanted transcription of toxic genes in insect cells, and provides the capability for co-

expression of multiple proteins via a single virus. We demonstrate the successful application of

the pBMCL1 vector for the expression of not only the human TRPC3 channel but also the

heteromeric KATP channel.

1. Introduction

Baculoviruses are double-stranded DNA viruses that can infect and replicate in insect

populations. *Autographa californica* multiple nuclear polyhedrosis virus (AcMNPV) is one of

the most common isolates of baculovirus. A modified version of AcMNPV, of which the

polyhedrin gene is replaced by a target gene, is widely used for over-expression of recombinant

protein in insect cells. Since it was found that the modified AcMNPV could mediate gene-

transfer in mammalian cells [1-3], baculovirus is more and more extensively used as a vehicle for

recombinant protein production in mammalian cells and this procedure is called BacMam system.

BacMam system has several advantages over other eukaryotic expression systems, including

insect cells or yeast expression systems. BacMam system allows exogenous gene transcription

from several available strong promoters such as CMV in mammalian cells. Moreover, the

BacMam system provides genuine mammalian cell environment for the posttranslational

modifications of target proteins, and the glycosylation pattern of membrane protein or secreted

protein can be modified using available cell line such as GnT1^{-}[4]. Furthermore, BacMam system

often provides high level of membrane protein production compared to other expression systems,

allowing the access to milligram-level purified sample of previously intractable membrane

protein targeted for structural or biochemical studies [5].

On the membrane envelope of baculovirus, there are several copies of GP64 surface protein

which might bind to the heparin sulfate proteoglycans on the plasma membrane of mammalian

cells [6]. The virus then enters the cell through clathrin-mediated endocytosis process [7]. Many

exogenous membrane proteins can be functionally displayed on the baculovirus envelope during

virus assembly and budding. It is reported that the vesicular stomatitis virus glycoprotein G

(VSV-G) protein displayed on baculovirus envelope can enhance the infectivity of baculovirus

towards mammalian cells [8] and a short fragment of VSV-G (VSV-GED) is sufficient to facilitate

baculovirus-mediated gene delivery in vertebrate cells [9,10]. Moreover, the large physical size of

baculovirus provides high capacity for packing exogenous DNA, allowing delivering multiple

genes in one single virus. However, the current available vectors did not fully exploit these

advantages of BacMam system. Here, we describe the design and application of an improved

vector for BacMam system.

2. Materials and methods

2.1. Cell lines and reagents

FreeStyle 293F cells (Thermo Fisher Scientific, R79007), Sf9 cells in Sf-900 III SFM (Thermo

Fisher Scientific, 12659017), FreeStyle 293 Expression Medium (Thermo Fisher Scientific,

12338018), Sf-900 III SFM medium (Thermo Fisher Scientific, 12658019), SIM SF Expression

Medium, (Sino Biological Inc, MSF1), SMM 293-TI (Sino Biological Inc., M293TI), Fetal

bovine serum (Vistech), PEI (Polyscience, 687600), Lauryl Maltose Neopentyl Glycol, LMNG

(Antrance, NG310), Cholesteryl Hemisuccinate Tris Salt, CHS (Antrance, CH210), digitonin

(Biosynth)

2.2. Construction of pBMCL1 vector

We synthesized two DNA sequences that contain the open reading frame of (GP64 signal

peptide)-(VSV-GED)-(CreiLOV) fusion protein and (LINK1)-(CMV promoter)-(multiple

cloning site, MCS)-(Woodchuck Hepatitis Virus Posttranscriptional Regulatory Element,

WPRE)-(SV40 polyA signal)-(LINK2) expression cassette. We inserted (GP64 signal peptide)-

(VSV-GED)-(CreiLOV) into pFastBac1 vector and replaced its SV40 polyA signal with

HSVtkPolyA signal to generate a vector that displays VSV-GED protein on insect cell

membrane and baculovirus envelope. Then we insert the expression cassette of (LINK1)-(CMV

promoter)-(MCS)-(WPRE)-(SV40 polyA signal)-(LINK2) into the vector to the opposite

direction of the pH promoter to minimize mammalian gene expression in insect cells. We name

this vector pBMCL1 (Figure 1).

2.3. Generation of expression constructs

The cDNA of human TRPC3 (hTRPC3) with N-terminal GFP was cloned into the MCS of

pBMCL1 vector for protein expression using BacMam system. The cDNA of SUR1 of

Mesocricetus auratus (XP_005084737) [11] and the cDNA of Kir6.2 with C-terminal GFP tag

from *Mus musculus* [12] were cloned into the MCS of pBMCL1 BacMam expression vector to

generate pBMCL1-Kir6.2-CGFP and pBMCL1-SUR1 vector. Then these two vectors were

merged into one vector by the LINK sequences using ligation-independent cloning method [13] to

generate the pBMCL1-K_{ATP} vector.

2.4. Generation of baculoviruses and infection of mammalian cells

pBMCL1 vectors containing target proteins were transformed into DH10Bac *Escherichia coli* cells. Recombinant baculoviruses were generated using the Bac-to-Bac Baculovirus Expression system (Invitrogen). Sf9 cells were cultured in Sf-900 III SFM medium (Gibco) at 27°C and seeded in 12-well plate. After transfection for 7 days, the fluorescence of CreiLOV was detected and the supernatant was collected. Higher titer P2 virus was amplified using sf9 cells cultured in SIM SF medium. At the time of cell mortality over 20%, the supernatant that containing baculovirus was collected by centrifugation and used to infect FreeStyle 293F cells.

2.5. Fluorescence-detection size-exclusion (FSEC)

Cells that expressed hTRPC3 channel were solubilized in TBS (20 mM Tris pH 8.0 at 4°C, 150 mM NaCl) with 10 mM MNG, 0.1% CHS and protease inhibitors including 1 μg/ml aprotinin, 1 μg/ml leupeptin, and 1 μg/ml pepstatin on ice for 30 min. Cell lysates were removed by centrifugation at 40,000 rpm for 30 min and supernatants were loaded onto Superose 6 Increase (GE Healthcare) for FSEC analysis.

Cells that expressed K_{ATP} channels were harvested and solubilized in TBS with 1% digitonin and protease inhibitors including 1 μg/ml aprotinin, 1 μg/ml leupeptin, and 1 μg/ml pepstatin on ice for 30 min. Cell lysates were removed by centrifugation at 40,000 rpm for 30 min and supernatant was loaded onto Superose 6 Increase for FSEC analysis.

2.6. Microscopy imaging

Imaging experiment was performed on sf9 cells 7 days post transfection of bacmid using EVOS FL imaging system. FreeStyle 293F cells (cultured in Freestyle 293 medium) were firstly plated

in 35 mm dish without shaking for 30 min, then baculoviruses were added into the medium. 48 h

post infection, GFP signal was detected. Green fluorescence images were obtained using Axio

Image M2 microscope.

2.7. Electrophysiology

For patch clamp recording, pBMCL1-K_{ATP} plasmids were transfected into FreeStyle 293F cells

(cultured in FreeStyle 293 medium) at density of 1×10^6 cells/ml using PEI and cells were

cultured for 32 h before recording. Patch electrodes were pulled from a horizontal microelectrode

puller (P-1000, Sutter Instrument Co, USA) to a tip resistance of 2-3 MΩ. Currents were

recorded using inside-out mode at +60 mV using an Axopatch 200B amplifier (Axon

Instruments, USA). We used KINT solution which contained (mM):140 KCl, 1EGTA, 10

HEPES (pH 7.4, KOH) in both pipette and bath. Signals were acquired at 20 kHz and low-pass

filtered at 5 kHz. Data were further analyzed with pClampFit 9.0 software and filtered at 1 kHz.

3. Results and discussion

3.1 Design of the pBMCL1 vector

We adopted the popular Bac-to-Bac system which utilizes the site-specific Tn7-mediated

transposition for bacmid generation. We modified the established BacMam expression cassette [5]

to minimize linker lengths between each elements and to remove common restriction enzyme

cutting sites out of the MCS. To confer the capability to express multiple genes simultaneously,

we appended LINK1 and LINK2 sequences at 5' and 3' of the expression cassette respectively.

LINK sequences allow the unidirectional fusion of unlimited number of expression cassettes

using a ligation-independent and PCR-independent cloning method [13]. LINK sequences are

found to be functional in BacMam system based on our previous successful application on the

expression of DUOX1-DUOXA1 complex [14]. The resulting expression cassette contains a 5'

LINK1 sequence, a CMV promoter, an MCS, a WPRE, a SV40 polyA signal and a 3' LINK2

sequence.

Figure 1. The design of pBMCL1 vector. **a**, Map of pBMCL1 vector. Elements that are essential
for function are shown on the vector. **b**, The sequence of MCS and restriction enzyme
recognition sites. **c**, LINK1 and LINK2 sequences and their restriction enzyme recognition sites
by SwaI and PacI. These sequences are used for merging multiple expression cassettes into one
vector by ligation-independent cloning.

The successful application of BacMam system highly relies on the titer of baculovirus produced

from insect cells. The strong expression of some proteins, such as certain ion channels, in insect

cells might be toxic and lead to low virus-titer. Therefore, we intentionally insert the

aforementioned expression cassette in reverse direction to the polyhedrin promoter of pFastbac1

vector to prevent the high level transcription of target gene in insect cells. In order to trace the

successful transfection of bacmid and production of baculovirus, we utilize the polyhedrin

promoter to drive the expression of a fluorescent reporter, VSV-GED-LOV fusion protein in

insect cells. A GP64 signal peptide directs the fusion protein to the plasma membrane of insect

cell and virus envelope. VSV-GED fragment is used to enhance the infectivity of baculovirus [9,10].

CreiLOV is a small, thermostable, photostable, and fast-maturing monomeric flavin-based green

fluorescent protein [15]. The commonly available green channel on fluorescent microscope for

GFP detection can be used to visualize the bright CreiLOV signals. If the insect cells are

successfully transfected with bacmids and are producing baculovirus, they would show green

signals generated from CreiLOV which is under the control of polyhedrin promoter (Figure 2).

Figure 2. Successful baculovirus generation indicated by green fluorescence signal in sf9 cells. **a**,
CreiLOV signal in sf9 cells after transfection for 7 days detected by a fluorescent microscope
using green channel. **b**, Bright filed image of transfected sf9 cells. **c**, The merged image of green
signal and bright field signal.

3.2 Using pBMCL1 vector to express human TRPC3 channel in mammalian cells

To test the function of pBMCL1 vector, we used it to express human TRPC3 (hTRPC3) channel

in FreeStyle 293F cells. Full length hTRPC3 is a canonical TRP channel with molecular weight

around 90 kDa. Functional hTRPC3 is a homo-tetrameric channel. We cloned the full length

hTRPC3 with N-terminal GFP tag into the pBMCL1 vector. After transfection of the bacmid into

sf9 insect cells for 7 days, we observed green fluorescence signal using GFP channel (Figure 3a)

and P1 viruses were harvested and further amplified to obtain the P2 virus. P2 viruses were used

for overexpression of hTRPC3 in FreeStyle 293F cells. 60 h post infection, we observed GFP

signal from the majority of FreeStyle 293F cells (Figure 3b). We also found the band of NGFP-

hTRPC3 protein at correct molecular weight on SDS-PAGE using in-gel fluorescence. (Figure

3c).

To optimize the protein expression condition, we screened different virus-cell ratio, expression

duration and expression medium. We found that at the volume ratio of 1:10 (virus: cell), the

expression level was highest (Figure 3d). Moreover, the expression level of hTRPC3 peaked at

48 h post-infection but decreased thereafter (Figure 3e). Furthermore, we found 293-TI medium

supported higher level of hTRPC3 expression than the FreeStyle 293 medium under conditions

tested (Figure 3f).

Figure 3. Expression of hTRPC3 channel via BacMam system using pBMCL1 vector. **a**,
CreiLOV signal of sf9 cell 7 days post transfection of bacmid. **b**, GFP signal of FreeStyle 293F
cells 48 h post infection by BacMam virus with virus-cell ratio 1:10 (v/v). **c**, In-gel fluorescence
of NGFP-hTRPC3 protein. Band of NGFP-hTRPC3 protein is boxed in red. **d**, FSEC profiles of
NGFP-hTRPC3 tetramer produced from FressStyle 293F cells infected with different virus to

cell ratio (v/v). **e**, FSEC profiles of NGFP-hTRPC3 tetramer produced from FressStyle 293F
cells that were cultured in different medium. **f**, FSEC profiles of NGFP-hTRPC3 tetramer
produced from FressStyle 293F cells expressed for various time.

3.3 Using pBMCL1 vector to express pancreatic K$_{ATP}$ channel in mammalian cells

To explore the capability of co-expressing more than one protein using a single pBMCL1 vector,

we expressed the hetero-octameric pancreatic K$_{ATP}$ channel in FreeStyle 293F cells. K$_{ATP}$

channel is a potassium channel, the activity of which is sensitive to intracellular ATP/ADP ratio.

Pancreatic K$_{ATP}$ channel in β cells couples blood glucose level to insulin secretion and is a drug

target for diabetes. Pancreatic K$_{ATP}$ channel is assembled from four Kir6.2 subunits and four

SUR1 subunits. Kir6.2 subunit is the ion channel pore with molecular weight 43.5 kDa. SUR1

subunit is the regulatory subunit with molecular weight 177 kDa. To trace the expression of K$_{ATP}$

channel, we fused mouse Kir6.2 with a C-terminal GFP tag and cloned it into pBMCL1 vector to

generate pBMCL1-Kir6.2-CGFP. We cloned non-tagged SUR1 subunit into pBMCL1 to

generate pBMCL1-SUR1 vector. The expression cassette of pBMCL1-SUR1 was inserted into

pBMCL1-Kir6.2-CGFP to obtain the pBMCL1-K$_{ATP}$ vector. Transient transfection of pBMCL1-

K$_{ATP}$ into FreeStyle 293F cells allowed the recording of robust ATP-sensitive potassium currents

in inside-out mode (Figure 4a). Moreover, the currents were sensitive to the insulin secretagogue

repaglinide, suggesting functional K$_{ATP}$ channels were correctly assembled and trafficked onto

the plasma membrane. When solubilized, the K$_{ATP}$ complex ran faster than Kir6.2 channel

expressed alone on FSEC, indicating the hetero-octamer assembly (Figure 4b). We generated

BacMam virus using pBMCL1-K$_{ATP}$ vector for infection of FreeStyle 293F cells. We found K$_{ATP}$

complex had robust expression 48 h post-infection (Figure 4c). We also showed the expression

of both Kir6.2 and SUR1 subunits by western blot (Figure 4d). These data collectively

demonstrate pBMCL1 vector is capable for the expression of eukaryotic membrane protein

complex.

Figure 4. Expression of K_{ATP} channel via BacMam system using pBMCL1 vector. **a**, Recording
of robust ATP-sensitive potassium currents in inside-out mode in FreeStyle 293F cells which
were transfected by pBMCL1-K_{ATP} plasmid, indicating functional co-expression of both Kir6.2
and SUR1 subunits. **b**, The FSEC profile of K_{ATP} complex (using pBMCL1-K_{ATP} plasmid) and
Kir6.2 channel in transfected cells. The earlier elution of K_{ATP} complex on size-exclusion
chromatography suggests the successful hetero-complex formation between Kir6.2 and SUR1
subunits. **c**, The FSEC profile of K_{ATP} complex in FreeStyle 293F cells infected by pBMCL1-
K_{ATP} BacMam virus. **d**, In virus-infected cells, the expression of Kir6.2 was detected by in-gel
GFP fluorescence and the expression of SUR1 subunit was detected by western blot using anti-
SUR1 antibody.

4. Conclusion

In this paper, we have described the design of pBMCL1 vector with several new features. First, this vector facilitates the tracing of successful virus generation by green fluorescence signal from LOV. Second, this vector might enhance the infectivity of BacMam virus towards mammalian cells by displaying VSV-GED fragment on baculovirus surface. Third, the expression of target protein in insect cells is minimized by placing CMV promoter in reverse direction to the polyhedrin promoter. Fourth, the incorporation of LINK sequences permits co-expression of multiple genes using BacMam system at the same time. Using this vector, we successfully expressed homo-tetrameric ion channel hTRPC3. Furthermore, we demonstrated the functional expression of hetero-octameric pancreatic K_{ATP} channel using this vector. We believe the aforementioned new features of this vector would help researchers to fully unleash the capability of BacMam system for recombinant protein expression.

Funding sources

This work is supported by grants National Natural Science Foundation of China (91957201, 31870833 and 31821091 to L.C.)

Credit authorship contribution statement

Wenjun Guo: Methodology, Validation, Formal analysis, Investigation, Writing - review & editing; **Mengmeng Wang:** Methodology, Validation, Formal analysis, Investigation, Writing - review & editing; **Lei Chen:** Conceptualization, Methodology, Validation, Formal analysis, Investigation, Writing - original draft, Writing - review & editing, Visualization, Supervision, Project administration, Funding acquisition.

Competing interests

The authors declare no competing interests

Acknowledgements

We thank Kaili Xue from Prof. Yifu Qiu lab and Peixue Xia from Prof. Ying Liu lab for kindly

help of microscopy imaging. We thank Chen lab members for kindly help.

References

Hofmann, C. *et al.* Efficient gene transfer into human hepatocytes by baculovirus vectors. *Proc. Natl. Acad. Sci. U. S. A.* **92**, 10099-10103, doi:10.1073/pnas.92.22.10099 (1995).

Boyce, F. M. & Bucher, N. L. Baculovirus-mediated gene transfer into mammalian cells. *Proc. Natl. Acad. Sci. U. S. A.* **93**, 2348-2352, doi:10.1073/pnas.93.6.2348 (1996).

Shoji, I. *et al.* Efficient gene transfer into various mammalian cells, including non-hepatic cells, by baculovirus vectors. *J. Gen. Virol.* **78 (Pt 10)**, 2657-2664, doi:10.1099/0022-1317-78-10-2657 (1997).

Dukkipati, A., Park, H. H., Waghray, D., Fischer, S. & Garcia, K. C. BacMam system for high-level expression of recombinant soluble and membrane glycoproteins for structural studies. *Protein Expr. Purif.* **62**, 160-170, doi:10.1016/j.pep.2008.08.004 (2008).

Goehring, A. *et al.* Screening and large-scale expression of membrane proteins in mammalian cells for structural studies. *Nat. Protoc.* **9**, 2574-2585, doi:10.1038/nprot.2014.173 (2014).

Wu, C. & Wang, S. A pH-sensitive heparin-binding sequence from Baculovirus gp64 protein is important for binding to mammalian cells but not to Sf9 insect cells. *J. Virol.* **86**, 484-491, doi:10.1128/JVI.06357-11 (2012).

Long, G., Pan, X., Kormelink, R. & Vlak, J. M. Functional entry of baculovirus into insect and mammalian cells is dependent on clathrin-mediated endocytosis. *J. Virol.* **80**, 8830-8833, doi:10.1128/JVI.00880-06 (2006).

Barsoum, J., Brown, R., McKee, M. & Boyce, F. M. Efficient transduction of mammalian cells by a recombinant baculovirus having the vesicular stomatitis virus G glycoprotein. *Hum. Gene Ther.* **8**, 2011-2018, doi:10.1089/hum.1997.8.17-2011 (1997).

Kaikkonen, M. U. *et al.* Truncated vesicular stomatitis virus G protein improves baculovirus transduction efficiency in vitro and in vivo. *Gene Ther.* **13**, 304-312, doi:10.1038/sj.gt.3302657 (2006).

Graves, L. P. *et al.* Improved Baculovirus Vectors for Transduction and Gene Expression in Human Pancreatic Islet Cells. *Viruses* **10**, doi:10.3390/v10100574 (2018).

Aguilar-Bryan, L. *et al.* Cloning of the beta cell high-affinity sulfonylurea receptor: a regulator of insulin secretion. *Science* **268**, 423-426 (1995).

12 Woo, S. K., Kwon, M. S., Ivanov, A., Gerzanich, V. & Simard, J. M. The sulfonylurea
receptor 1 (Sur1)-transient receptor potential melastatin 4 (Trpm4) channel. *J. Biol. Chem.*
288, 3655-3667, doi:10.1074/jbc.M112.428219 (2013).
13 Scheich, C., Kummel, D., Soumailakakis, D., Heinemann, U. & Bussow, K. Vectors for
co-expression of an unrestricted number of proteins. *Nucleic Acids Res.* **35**, e43,
doi:10.1093/nar/gkm067 (2007).
14 Wu, J. X., Liu, R., Song, K. & Chen, L. Structures of human dual oxidase 1 complex in
low-calcium and high-calcium states. *Nat Commun* **12**, 155, doi:10.1038/s41467-020-
20466-9 (2021).
15 Mukherjee, A. *et al.* Engineering and characterization of new LOV-based fluorescent
proteins from Chlamydomonas reinhardtii and Vaucheria frigida. *ACS Synth Biol* **4**, 371-
377, doi:10.1021/sb500237x (2015).

290